Differential Effects of Value and Salience on Saccade Latencies

Mohammad Shams-Ahmar & Peter Thier

Department of Cognitive Neurology, Hertie Institute for Clinical Brain Research, University of Tübingen, 72076 Tübingen, Germany.

Address for correspondence:

Mohammad Shams-Ahmar
Department of Cognitive Neurology
Hertie Institute for Clinical Brain Research
Otfried-Müller-Str. 27
72076 Tübingen, Germany
m.shamsahmar@gmail.com

or

Prof. Dr. Peter Thier
Department of Cognitive Neurology
Hertie Institute for Clinical Brain Research
Hoppe-Seyler-Str. 3
72076 Tübingen, Germany
thier@uni-tuebingen.de

ABSTRACT

Express saccades, a mode of visually guided saccades that are distinguished from regular saccades by extremely short reaction times, are triggered by inserting a temporal gap between the fixation dot and the saccade target. It is usually assumed that they are produced by a specific pathway in which the superior colliculus plays a key role. Whether and how this pathway deals with information on the subjective value of a saccade target is unknown. We therefore studied the influence of varying reward expectancies and compared it with the impact of the presence and absence of a temporal gap between the disappearance of the fixation dot and the appearance of the target on the visually guided saccades of two rhesus macaques (*Macaca mulatta*). We observed that the introduction of a gap shifted the entire saccadic reaction time distribution to shorter latencies while increasing the probability of express saccades. On the other hand, promoting the monkey's reward expectancy shortened reaction times and increased peak velocities of regular saccades, and increased the probability of express saccades. Importantly, we observed that the reaction time and peak velocity of express saccades were not sensitive to the value of the saccade target, suggesting that the express pathway does not have access to information on value. We propose a new model on express saccades that treats the salience of visual objects in the scene differently from the subjective value assigned to them.

KEY WORDS

Express saccade; value; salience; reaction time distribution; superior colliculus;

INTRODUCTION

In a visually guided saccade task, inserting a temporal gap between the fixation dot offset and the saccade target onset increases the probability of express saccades at the expense of regular saccades, highlighting two distinct modes of saccadic reaction times (Dorris et al. 1997; Fischer et al. 1984; Fischer and Boch 1983; Fischer and Ramsperger 1984; Mayfrank et al. 1986; Pare and Munoz 1996; Schiller et al. 2004). It is commonly assumed that these two modes arise from two—at least partially—distinct pathways, with one, responsible for express saccades, being significantly faster (Chen et al. 2013; Isa and Hall 2009; Schiller and Tehovnik 2005). The existence of these two pathways might explain the bimodality of the saccadic reaction time distribution. However, the question why such parallel pathways exist in the first place, and under which circumstances the one or the other is given preference, remains unanswered.

It is well established that both saccade reaction time and saccade peak velocity of regular saccades depend on the expected value of the saccade target (Lauwereyns et al. 2002; Milstein and Dorris 2007; Takikawa et al. 2002). Targets with large expected value are associated with shorter reaction times and higher peak velocities, features that increase the probability of getting hold of a potentially fleeing valuable target, yet, at the expense of significant investments of costly resources. On the other hand, reacting less vigorously in cases of targets of low expected value, whose loss one may bear more easily, has the advantage that these investments are lower. In any case, assessing the expected value will inevitably take time with detrimental consequences for targets that may be expected to be of existential importance. Hence, in this case, skipping the time-consuming assessment of value to react as quickly as possible – for instance to something potentially dangerous – may be the better strategy. Could it be that express saccades are a manifestation of such a strategy? Of course, to take advantage of such a selection, reliable prior assumptions on the relevance of stimuli would be needed. However, independent of the question of what their basis might be, one would expect to see no correlation between the metrics of express saccades and the expected value of the saccade target.

To test our hypothesis of a differential influence of value for express and regular saccades, we carried out a study of visually guided saccades made by well-trained rhesus monkeys. In our experiments, we manipulated the expected value and the temporal context of the presentation of the saccade target: the earlier was realized by providing a cue during the fixation period for the amount of upcoming reward, and the latter by randomly inserting a temporal gap between the fixation dot offset and the onset of the saccade target. Our results support the concept of two—partially separated—pathways, fed by a common input that is responsible for the influence of physical properties of the stimuli on both. The target value, along with the target's physical properties, influences the routing between the two pathways and as hypothesized, its influence is largely confined to the regular saccade pathway.

METHODS

Subjects and setup. Two male rhesus monkeys (8-13 kg) were trained on a visually guided saccade task with two distinct features. First, on randomly chosen trials a temporal gap separated fixation dot offset and target onset, whereas on the other trials the two coincided. Second, monkeys received prior information on the varying amount of reward to be expected in the case of a successful trial. Eye movements were recorded with an ISCAN ETL-200 video tracker, resampled at 1 kHz. During experiments, the monkeys sat in a primate chair with their heads immobilized, 60 cm away from an LED monitor in a dim room. The surgical techniques used for the implantation of the head posts, a scleral search coil and a chamber for later recordings from the cerebellum, followed protocols described in detail before (Arnstein et al. 2015). The procedure was approved by the local animal care committee under European and German law and the National Institutes of Health's Guide for the Care and Use of Laboratory Animals.

Behavioral procedure. Each trial began after a variable inter-trial interval of 200-500 ms, followed by a fixation period of 500 ms duration. The fixation dot ($d = 0.25^\circ$) appeared in the center of the screen and the monkey was required to keep his gaze within a window of 1.5° centering on the fixation dot. The initial fixation dot was replaced by a 1° diameter symbol for 300 ms, still requiring fixation, besides informing the monkey on the amount of reward to be expected. Two reward cues (Fig. 1A) were chosen at random. In both monkeys, the mapping between the two cues and the two levels of reward was reversed after about half of the experimental sessions. After the disappearance of the reward cue, the fixation dot reappeared and lasted for another 300-600 ms. On almost half of the trials, chosen at random, the fixation dot offset was followed by the immediate appearance of a target dot ($d = 0.25^\circ$) at 8° eccentricity on the horizontal axis, at random left or right. On the other half of trials, the onset of the peripheral target followed the offset of the fixation dot after a temporal gap of 200 ms during which the monkey had to keep fixating the position of the fixation dot. The monkeys had to make a saccade to the saccade target within 500 ms following its onset and hold their gaze within a fixation control window for 500 ms to receive a reward (Fig. 1A). All cues used were monochromatic, had a luminance of 53 cd/m^2 , and appeared on a black background of 0.2 cd/m^2 .

To gauge the subjective value of the two reward levels possible, the standard trials were occasionally replaced by free choice trials, interspersed at random with a probability of 5%. In these free-choice trials, the high-reward and the low-reward cue appeared simultaneously on the left and right side of the fixation dot at 8° , serving as optional targets. Their onset took place simultaneously with the offset of the central fixation dot after 500-800 ms. The monkeys could freely choose between the two target options and received the expected reward. Once a new association of reward cue and reward level was chosen, it took the monkeys 2 to 3 sessions to prefer the high-reward target in more than 90% of free-

choice trials. Data from the main trials were used for the analysis of the influence of the gap and expected reward only after this choice probability had been reached.

Eye data analysis. The eye position was recorded using the ISCAN digital eye tracker at a sampling rate of 120 Hz. To obtain a better estimation of the saccade times and positions, the eye signals were smoothed by fitting a second-degree polynomial model, using a local regression method of size 50 ms around each sample. For saccade detection, the local extrema in the velocity profile which had absolute values between 100 to 1000 deg/s were identified as potential saccades. Then the onset and offset of each potential saccade were defined as the first data points that fell below 25 deg/s around the extremum. Saccades that landed within one degree of the targets were labeled as valid. Saccades that took less than 60 ms were considered as anticipatory saccades and were not included in further analyses.

Distinguishing express saccades and regular saccades. To this end, we subjected saccadic reaction times to a cluster analysis using the k -means algorithm, with the number of clusters set to two. Reaction times in the range of 60 ms to 150 ms were fed into the clustering algorithm for potential outliers not to bias the decision of the algorithm. Clustering was run 10 times, each time with a new set of initial cluster centroids. The iteration resulting in the least sum of all the points' distances to their corresponding centroid was taken as the final output.

Saccade amplitude matching of peak velocities. First, the distributions of saccade amplitudes in the low-reward condition (LR_{cond}) and high-reward condition (HR_{cond}) were characterized by histograms with an equal bin width (0.2° or 0.3°). Then, from each amplitude bin, we drew N saccades from each of the two distributions, with N representing the smaller of the two counts of saccades obtained for that bin for the two conditions. The corresponding peak velocities of these N trials across all amplitude bins were extracted to form the amplitude-matched distribution of peak velocities. For calculating the average peak velocity of the session, an average of 100 iterations of the same procedure was taken as the final output.

Unless otherwise stated, all statistical comparisons were based on Wilcoxon signed-rank tests ($p < 0.001$ by ***, $p < 0.01$ by **, and $p < 0.05$ indicated by *).

RESULTS

Differential effects of reward expectancy and the gap on the express saccade boundary. We collected a total number of 35 experimental sessions ($n_{\text{monkey1}} = 21$, $n_{\text{monkey2}} = 14$). The resulting saccadic reaction time distributions of both monkeys showed distinct express saccade modes, easily discernible by eye, independent of whether a gap was present or not. Nonetheless, to avoid any subjective bias in determining the reaction time boundary between regular saccades and express saccades (X_{bound}), we deployed the k -means clustering algorithm (see Methods). Fig. 1B illustrates the distribution of saccadic reaction times in an exemplary session with a gap between the disappearance of the fixation dot and the appearance of the target for low reward trials. In this particular example, the k -means algorithm suggested an X_{bound} separating express and regular saccades at 105 ms. The k -means clustering analysis was carried out separately for gap and no-gap condition trials, furthermore separating low- and high-reward trials. Figs. 1C-E summarize the effects of the two variables reward expectancy and the presence or absence of a gap (gap_{cond} vs. no-gap_{cond}) on the X_{bound} . The reward level did not influence the X_{bound} in the no-gap_{cond}, and reduced it only slightly, yet significantly, in the gap_{cond} (median reduction = 1 ms; Fig. 1C). On the other hand, the introduction of a gap caused a shift of the X_{bound} towards lower saccadic reaction times in both reward conditions. A “gap effect” i.e. the mean saccadic reaction times difference between gap_{cond} and no-gap_{cond} (Jin and Reeves 2009; Kingstone and Klein 1993; Rolfs and Vitu 2007), pooled over both saccade modes of about 12 ms, was observed in both reward conditions. Thus, the introduction of a gap strongly modulated the position of the X_{bound} , whereas the effect of reward expectancy was negligible.

Differential effects of reward expectancy and the gap on the express saccade probability. We next analyzed how the reward expectancy influenced the probability of express saccades generated in the gap_{cond} vs. no-gap_{cond}. In the no-gap_{cond}, both monkeys showed a significantly higher probability of generating express saccades in the HR_{cond} compared to the LR_{cond}. In the gap_{cond}, the same effect was obtained for one of the two monkeys ($p = 0.0004$) but neither seen in the other monkey ($p = 0.27$) nor in data pooled from both (Fig. 2A). One may safely assume that the subjective value of both the low- and high-level rewards attained by the monkey will steadily decrease in a session until a point of complete satiety is reached, stopping the session. Hence, the aforementioned promotion of express saccades by large reward expectancy might be particularly strong early in a session. In fact, this was the case. The probability of express saccades was highest in the first decade of a session and decreased steadily over sessions (Fig. 2B). In the presence of a gap, the probability of express saccades reached 75% in the first decade, no matter if the reward was large or small and decreased to about 50% in the last decade. In the no-gap_{cond}, in which the overall probability of express saccade was less, with significantly more express saccades for larger reward expectancies, the decrease with the trial number was much weaker, albeit still significant for the LR_{cond} and not significant for HR_{cond}.

Differential effects of reward expectancy and the gap on saccadic reaction times and peak velocities. Next, we asked if manipulating reward expectancies and introducing a temporal gap between the fixation dot disappearance and the target appearance influenced saccadic reaction times and velocities of saccades. As shown in Fig. 3A and C, the saccadic reaction times of express saccades as well as their peak velocity did not show a change due to reward expectancy. However, the variability of saccadic reaction times dropped strongly (median STD difference = 1.2, $p = 1.45 \times 10^{-7}$). The mean of the same two factors underwent a strong modulation in the case of regular saccades, whereas mean saccadic reaction times dropped and mean peak velocities increased robustly due to a higher expectancy of reward. Besides, we found that the peak velocity of express saccades was on average below that of regular saccades. For the LR_{cond} the difference was 17.48 deg/s ($p = 7.39 \times 10^{-12}$) and for the HR_{cond} it was 25.68 deg/s ($p = 7.49 \times 10^{-11}$).

Modulation across reaction time bins. The analyses presented in the preceding paragraphs were based on the assumption of two distinct saccade modes, an express saccade mode and a regular saccade mode, and the expectation of qualitatively distinct effects of the two reward levels and the presence or absence of a gap on saccades in the two modes. The results seem to meet this expectation. However, the a priori separation of saccades into two modes might conceal that the effects of the two experimental variables are simply graded with saccade latency, independent of a saccade interpreted as an express or regular saccade. To address this concern, we calculated the probability of saccades in individual latency bins for a pool of saccades characterized by the same reward level and the presence, or absence of a gap, bin-by-bin. Next, we subtracted the resulting probability distribution for the LR_{cond} from the one for the HR_{cond}. The resulting distributions are shown in Fig. 4A, separately for the gap and the no-gap conditions. In both gap conditions, we saw that reward increased the number of saccades in the earlier saccadic reaction time bins at the expense of the later ones. In contrast to what the assumption of a graded influence of saccade latency would predict, the resulting pattern is bimodal. In the no-gap_{cond}, the redistribution of later saccades to early saccades peaked at 140 ms and 100 ms, whereas in the gap_{cond} it peaked at 125 ms and 100 ms. This pattern suggests that a large reward expectancy speeded up all saccades within their respective modes while still obeying the X_{bound} .

To compare the total amount of modulation between the no-gap_{cond} and the gap_{cond}, we defined a modulation index for each session, given by the sum of the absolute probability differences across all saccadic reaction time bins. This comparison revealed that changes in reward expectancy had a stronger effect on saccade latencies in the no-gap_{cond} than in the gap_{cond}. Using a modulation index that captures both changes in express saccade probability and in mean saccadic reaction times, implies that express saccades in no-gap_{cond} are most susceptible to the subjective value assigned to the saccade target.

Finally, to fully describe the saccadic reaction time distributions for the four conditions distinguished by reward level and the presence or absence of a fixation gap with a high temporal resolution, we

quantified each resulting saccadic reaction time mode by four parameters: mean, standard deviation, skewness, and kurtosis and modeled the result by resorting to the Pearson system probability distributions. The distributions then allowed us to generate artificial trials to make a visual comparison between the differential effects of value and salience on the saccadic reaction time distribution in one frame (Fig. 4B).

DISCUSSION

In this study, we found different sensitivities of saccades in the express mode and the regular mode to the expected value of the saccade. Promoting the expected value increased the peak velocities of regular saccades but not of express saccades. Furthermore, it biased the reaction times of regular saccades toward a minimum, corresponding to the express saccade boundary. Finally, it also caused an increase in the express saccade probability. On the other hand, introducing a temporal gap shifted the express saccade boundary to shorter reaction times and increased the probability of express saccades.

By taking the findings of other studies into account that showed the effect of physical properties of visual stimuli on the probability and timing of express saccades (Boch et al. 1984; Heeman et al. 2017; Marino et al. 2015; Marino and Munoz 2009), we propose a conceptual model of express saccades that accommodates the aforementioned differential effects on regular and express saccades (Fig. 5). The first stage of the model processes the physical properties of all visual objects, namely the fixation dot, the eventual target, and the “distractors” in the visual field, independently of the subjective value assigned to them. The visual objects compete with each other to become the target of an upcoming saccade, shifting the fovea to them, based on a comparison of their “visual drive”. We assume that this comparison is based on a divisive normalization algorithm (Carandini and Heeger 2012; Louie et al. 2013; Louie and Glimcher 2019). The salience of individual visual objects is obtained by dividing the individual visual drive by the sum of the visual drives of all objects, resulting in values between zero and one. In a natural setting, the object associated with the largest salience will win and correspondingly bind the observer’s foveal attention. The larger the salience of the winning object is, the shorter the saccadic reaction time will be. In a scenario in which just two objects staggered in time are seen, the first one available will bind the fovea first, i.e. serve as fixation dot, while the latter will serve as the saccade target. The advanced removal of the first one, the fixation dot in our paradigm, will maximize the salience of the later object, the saccade target.

The salience signal feeds two parallel pathways, namely the regular and the express saccade pathway, thereby influencing all saccades, i.e. shaping the entire reaction time distribution. Information processing based on the regular pathway takes longer than the processing accommodated by the express pathway. The reason is that the former has to incorporate information on the value to adjust the timing and the peak velocity of regular saccades according to the subjective value assigned to the saccade target. The express pathway is faster because it skips this process, remaining indifferent to the value of the saccade target. Hence, value affects the regular saccade mode of the reaction time distribution, while not touching the express saccade mode. Whether information is transferred to the premotor circuits from the one or the other pathway is determined by a switch that is controlled by the joint influence of salience and value signals. As a result, both value and salience signals influence the probability of express saccades.

The importance of the superior colliculus in generating express saccades became clear after it was shown that the unidirectional ablation of SC eliminates express saccades in the contralateral direction (Schiller et al. 1987). Additionally, some pieces of evidence suggest that the intermediate layers of the superior colliculus (SCi) might be a good candidate to serve as the integration and decision module in the scheme (Fig. 5, the green box). First, they receive visual information from the retina and visual cortex via superficial layers of the superior colliculus (Coe and Munoz 2017). Hence, the salience signal might originate from these structures. Alternatively, it might be derived from the posterior parietal cortex, which has direct projections to SCi (White et al. 2017; Wolf et al. 2015). Second, the basal ganglia, a system that plays a central role in encoding value, communicate with the SCi via the substantia nigra pars reticulata (Ikeda and Hikosaka 2003; Munoz and Everling 2004). Other studies have shown that the amount of build-up activity of visuomotor neurons in the deeper layers of the superior colliculus during the gap is a good predictor of the occurrence of an express saccade (Dorris et al. 1997; Edelman and Keller 1996). This build-up activity is controlled by target predictability (Basso and Wurtz 1997, 1998), as well as by target brightness (Marino et al. 2015). Whether value signals are represented in the SCi or deeper layers during the gap, and the question of how the integration of salience and value takes place in those layers, will require further experimental and computational studies.

CONTRIBUTIONS: M.S. conceived and designed research, performed experiment, analyzed data, prepared figures, and drafted manuscript. M.S. and P.T. interpreted results and revised manuscript.

ACKNOWLEDGMENTS: The authors would like to thank Dr. F. Bunjes for implementing the behavioral paradigm, Dr. P. Dicke, for assisting in the surgical procedure, and A. Ridderbusch for proofreading the manuscript. This study was funded by Deutsche Forschungsgemeinschaft <http://www.dfg.de> (grant number FOR 1847-A3, TH 425/13-1) received by PT within the framework of the Research Unit “Primate Systems Neuroscience”. The funder had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

DISCLOSURE: The authors declare no conflict of interest

REFERENCES

- Arnstein D, Junker M, Smilgin A, Dicke PW, Thier P.** Microsaccade Control Signals in the Cerebellum. *J Neurosci* 35: 3403–3411, 2015.
- Basso MA, Wurtz RH.** Modulation of neuronal activity by target uncertainty. *Nature* 389: 66–69, 1997.
- Basso MA, Wurtz RH.** Modulation of Neuronal Activity in Superior Colliculus by Changes in Target Probability. *J Neurosci* 18: 7519–7534, 1998.
- Boch R, Fischer B, Ramsperger E.** Express-saccades of the monkey: Reaction times versus intensity, size, duration, and eccentricity of their targets. *Experimental Brain Research* 55, 1984.
- Carandini M, Heeger DJ.** Normalization as a canonical neural computation. *Nat Rev Neurosci* 13: 51–62, 2012.
- Chen M, Liu Y, Wei L, Zhang M.** Parietal Cortical Neuronal Activity Is Selective for Express Saccades. *Journal of Neuroscience* 33: 814–823, 2013.
- Coe BC, Munoz DP.** Mechanisms of saccade suppression revealed in the anti-saccade task. *Phil Trans R Soc B* 372: 20160192, 2017.
- Dorris MC, Paré M, Munoz DP.** Neuronal Activity in Monkey Superior Colliculus Related to the Initiation of Saccadic Eye Movements. *The Journal of Neuroscience* 17: 8566–8579, 1997.
- Edelman JA, Keller EL.** Activity of visuomotor burst neurons in the superior colliculus accompanying express saccades. *Journal of Neurophysiology* 76: 908–926, 1996.
- Fischer B, Boch R.** Saccadic eye movements after extremely short reaction times in the monkey. *Brain Research* 260: 21–26, 1983.
- Fischer B, Boch R, Ramsperger E.** Express-saccades of the monkey: Effect of daily training on probability of occurrence and reaction time. *Experimental Brain Research* 55, 1984.
- Fischer B, Ramsperger E.** Human express saccades: extremely short reaction times of goal directed eye movements. *Experimental Brain Research* 57, 1984.
- Heeman J, Van der Stigchel S, Theeuwes J.** The influence of distractors on express saccades. *Journal of Vision* 17: 35, 2017.
- Ikeda T, Hikosaka O.** Reward-Dependent Gain and Bias of Visual Responses in Primate Superior Colliculus. *Neuron* 39: 693–700, 2003.
- Isa T, Hall WC.** Exploring the Superior Colliculus In Vitro. *Journal of Neurophysiology* 102: 2581–2593, 2009.
- Jin Z, Reeves A.** Attentional release in the saccadic gap effect. *Vision Research* 49: 2045–2055, 2009.
- Kingstone A, Klein RM.** Visual offsets facilitate saccadic latency: does predisengagement of visuospatial attention mediate this gap effect? *J Exp Psychol Hum Percept Perform* 19: 1251–1265, 1993.
- Lauwereyns J, Watanabe K, Coe B, Hikosaka O.** A neural correlate of response bias in monkey caudate nucleus. *Nature* 418: 413–417, 2002.

Louie K, Glimcher PW. Normalization Principles in Computational Neuroscience. In: *Oxford Research Encyclopedia of Neuroscience*. Oxford University Press.

Louie K, Khaw MW, Glimcher PW. Normalization is a general neural mechanism for context-dependent decision making. *Proceedings of the National Academy of Sciences* 110: 6139–6144, 2013.

Marino RA, Levy R, Munoz DP. Linking express saccade occurrence to stimulus properties and sensorimotor integration in the superior colliculus. *Journal of Neurophysiology* 114: 879–892, 2015.

Marino RA, Munoz DP. The effects of bottom-up target luminance and top-down spatial target predictability on saccadic reaction times. *Experimental Brain Research* 197: 321–335, 2009.

Mayfrank L, Mobashery M, Kimmig H, Fischer B. The role of fixation and visual attention in the occurrence of express saccades in man. *Eur Arch Psychiatry Neurol Sci* 235: 269–275, 1986.

Milstein DM, Dorris MC. The Influence of Expected Value on Saccadic Preparation. *Journal of Neuroscience* 27: 4810–4818, 2007.

Munoz DP, Everling S. Look away: the anti-saccade task and the voluntary control of eye movement. *Nat Rev Neurosci* 5: 218–228, 2004.

Pare M, Munoz DP. Saccadic reaction time in the monkey: advanced preparation of oculomotor programs is primarily responsible for express saccade occurrence. *Journal of Neurophysiology* 76: 3666–3681, 1996.

Rolfs M, Vitu F. On the limited role of target onset in the gap task: support for the motor-preparation hypothesis. *J Vis* 7: 7.1-20, 2007.

Schiller PH, Haushofer J, Kendall G. How do target predictability and precueing affect the production of express saccades in monkeys? *European Journal of Neuroscience* 19: 1963–1968, 2004.

Schiller PH, Sandell JH, Maunsell JH. The effect of frontal eye field and superior colliculus lesions on saccadic latencies in the rhesus monkey. *Journal of Neurophysiology* 57: 1033–1049, 1987.

Schiller PH, Tehovnik EJ. Neural mechanisms underlying target selection with saccadic eye movements. In: *Progress in Brain Research*. Elsevier, p. 157–171.

Takikawa Y, Kawagoe R, Itoh H, Nakahara H, Hikosaka O. Modulation of saccadic eye movements by predicted reward outcome. *Experimental Brain Research* 142: 284–291, 2002.

White BJ, Berg DJ, Kan JY, Marino RA, Itti L, Munoz DP. Superior colliculus neurons encode a visual saliency map during free viewing of natural dynamic video. *Nat Commun* 8: 14263, 2017.

Wolf AB, Lintz MJ, Costabile JD, Thompson JA, Stubblefield EA, Felsen G. An integrative role for the superior colliculus in selecting targets for movements. *Journal of Neurophysiology* 114: 2118–2131, 2015.

Fig. 1. The behavioral task and the express boundary (X_{bound}). **A:** the stimulus sequence. The cues were selected from the cue set and swapped their meaning after almost half of the sessions for each monkey. Dashed circles show the potential target positions. **B:** illustration of an exemplary session. The graph includes trials in the gap_{cond} and the LR_{cond} . Each dot represents a trial and the vertical line shows the calculated X_{bound} . **C:** effect of reward expectancy on X_{bound} separated by gap condition ($\text{median}_{\text{gap}} = 1$ ms, $p = 0.5$; $\text{median}_{\text{no-gap}} = -1$ ms, $p = 0.0027$). **D:** effect of introduction of a temporal gap on X_{bound} separated by reward condition ($\text{median}_{\text{LR}} = -3$, $p = 0.0021$; $\text{median}_{\text{HR}} = -6$, $p = 10^{-6}$). **E:** The gap effect separated by the reward condition ($\text{median}_{\text{LR}} = 11.94$, $\text{median}_{\text{HR}} = 11.69$, $p = 0.90$). In **C-E**, each dot represents an experimental session ($n = 35$), and the horizontal black bars indicate the median across sessions.

Fig. 2. Effect of reward expectancy on the probability of express saccades. *A*: comparison between the probability of express saccades in LR_{cond} and HR_{cond} in both gap conditions ($\text{median}_{\text{gap}} = -0.03$, $p = 0.09$; $\text{median}_{\text{no-gap}} = -0.07$, $p = 2 \times 10^{-6}$). Each marker represents an experimental session. The triangles above the histogram indicate the median of the corresponding distribution. *B*: the probability of express saccades throughout a session. Except for the HR_{cond}–no-gap_{cond}, the other trials show a significant difference of express saccade probability across trial decades (Kruskal-Wallis one-way test, LR_{cond}, no-gap_{cond}: $p = 8.37 \times 10^{-7}$, HR_{cond}, no-gap_{cond}: $p = 0.21$, LR_{cond}, gap_{cond}: $p = 9.63 \times 10^{-13}$, HR_{cond}, gap_{cond}: $p = 2.03 \times 10^{-8}$). The relative probability of express saccades in HR_{cond} was higher across trial bins in gap_{cond} ($p = 0.0039$) and no-gap_{cond} ($p = 0.002$). Vertical lines indicate the standard error of the mean (SEM) across sessions. Asterisks indicate those trial bins with a significant difference ($p < 0.05$) between HR_{cond} and LR_{cond}, and triangles show trial bins lying significantly ($p < 0.05$) above the chance level, specified by the horizontal gray line (both corrected for the multiple comparison testing using the Benjamini-Hochberg method).

Fig. 3. Effect of reward on saccade latency and peak velocity. *A-B*: comparison between the saccadic reaction time means in the LR_{cond} and the HR_{cond}, for express saccades (*A*; median difference (HR - LR) = -1.17 ms, $p = 0.79$) and regular saccades (*B*; median difference (HR - LR) = -4.38 ms, $p = 3.69 \times 10^{-11}$). *C-D*: comparison between the average saccade peak velocity in two reward conditions for express saccades (*C*; median difference (HR - LR) = 2.77 deg/s, $p = .92$) and regular saccades (*D*; median difference (HR - LR) = 3.55 deg/s, $p = 5.06 \times 10^{-7}$). Each circle represents an experimental session. The triangles above the histograms show the median of each distribution.

Fig. 4. Effect of reward and gap across saccadic reaction time bins. *A*: for each gap condition, the probability of each bin in each LR_{cond} was subtracted from that of the HR_{cond}. Vertical lines indicate the SEM across sessions. Gray and black asterisks indicate significantly different reaction time bins ($p < 0.05$) from zero in the gap_{cond} and the no-gap_{cond} respectively (corrected for the multiple comparison testing using the Benjamini-Hochberg method). *B*: the result of a simulation, based on features extracted from each session, using the Pearson system of probability distributions.

Fig. 5. Partially separated pathways for express and regular saccades. The physical properties of the saccade target are assessed by a divisive normalization process (gray box) that results in a saliency signal of the saccade target (blue arrows). This signal reaches the express saccade pathway and as well the regular saccade pathway. The latter involves a receiver of value signals (red arrows). An integration and decision module (green box), made of a switch under the joint influence of saliency and value signals, decides which of the two pathways gets access to the premotor circuitry for saccades.