Species dynamics and interactions via metabolically informed consumer-resource models

Mario E. Muscarella · James P. O'Dwyer

the date of receipt and acceptance should be inserted later

Abstract Quantifying the strength, sign, and origin of species interactions, along with their dependence on environmental context, is at the heart of prediction and understanding in ecological communities. Pairwise interaction models like Lotka-Volterra provide an important and flexible foundation, but notably absent is an explicit mechanism mediating interactions. Consumer-resource models incorporate mechanism, but describing competitive and mutualistic interactions is more ambiguous. Here, we bridge this gap by modeling a coarse-grained version of a species' true, cellular metabolism to describe resource consumption via uptake and conversion into biomass, energy, and byproducts. This approach does not require detailed chemical reaction information, but it provides a more explicit description of underlying mechanisms than pairwise interaction or consumer-resource models. Using a model system, we find that when metabolic reactions require two distinct resources we recover Liebig's Law and multiplicative co-limitation in particular limits of the intracellular reaction rates. In between these limits, we derive a more general phenomenological form for consumer growth rate, and we find corresponding rates of secondary metabolite production, allowing us to model competitive and non-competitive interactions (e.g., facilitation). Using the more general form, we show how secondary metabolite production can support coexistence even when two species compete for a shared resource, and we show how differences in metabolic rates change species' abundances in equilibrium. Building on these findings, we make the case for incorporating coarse-grained metabolism to update the phenomenology we use to model species interactions.

Mario E. Muscarella

Department of Plant Biology, University of Illinois, ORCID ID: 0000-0001-6183-6037, E-mail: mmuscar@illinois.edu

Current Address: Département des Sciences Biologiques, Université du Québec à Montréal

James P. O'Dwyer

Department of Plant Biology & Carl R. Woese Institute for Genomic Biology, University of Illinois,

ORCID ID: 0000-0003-1180-8622, E-mail: jodwyer@illinois.edu

Keywords Species interactions · Consumer-resource models · Metabolic model · Efficiency · Mutualism

Introduction

A central goal in ecology is to understand and predict the dynamics in communities of interacting species (Holt, 1977; Loreau, 2010; Vellend, 2010, 2016). Mathematical models allow us to generate and test theoretical predictions, and the development of such models leads to a hierarchy of challenges. The first challenge is determining an appropriate functional form describing species dynamics. A range of functional forms with increasing complexity has been used and each has strengths and weaknesses which are often context dependent (Holling, 1959; Abrams, 1982; DeAngelis et al., 1989; Murdoch et al., 2003; Mougi and Kondoh, 2012). Second, we need to parameterize these equations, for example by quantifying the strength and sign of species interactions in a given environmental context. While some attempts have been made to parameterize natural systems, accurately fitting interaction strengths remains challenging in both empirical and theoretical work (Schoener, 1983; Tilman, 1987; Ives et al., 2003; Marino et al., 2014; Carrara et al., 2015; Terry et al., 2017; Barner et al., 2018). Finally, we may wish to determine how changes in the environmental context will modify species interactions, dynamics, and even coexistence. Integrating each of these goals will lead to the development of robust models which can predict the dynamics of complex communities, even when the environmental landscape changes within and across ecosystems.

There are two primary approaches to mathematically model the dynamics of interacting species. The Lotka-Volterra equations characterize species interactions in terms of the net, direct effect of one population on another's growth rate (Lotka, 1932; Volterra, 1926, see Supplemental). However, because these models lack an explicit description of the mechanisms mediating interactions, it is often difficult to translate the inferred interactions in different environmental contexts (Abrams, 1983; Grilli et al., 2017). For example, if two species compete, it is often because they consume common resources (Gause and Witt, 1935; MacArthur, 1970; Schoener, 1983). But, these models assume that resource dynamics can be safely ignored because resource dynamics are faster than consumer dynamics (MacArthur, 1970). This exposes an important context-dependence of Lotka-Volterra type equations: the strength and even the sign of a pairwise interaction may depend on what resources are present (Xiao et al., 2017). An alternate approach, consumer-resource equations, characterizes competitive interactions as the explicit result of shared resource consumption (Tilman et al., 1982; Grover, 1990; Litchman, 2003; Abrams, 2009, see Supplemental). These models produce species interactions as an emergent property dependent on shared resource consumption. Assuming we can infer or otherwise estimate consumer preferences, one critical aspect of the environmental context is now explicitly characterized, via resource

input rates. As such, species dynamics across resource landscapes can be understood better than in the case of Lotka-Volterra, where the effect of the environment is implicit (Tilman, 1977; Grover, 1990, 2011).

The more explicit mechanism in a consumer-resource model gives an advantage, but there is a cost. Lotka-Volterra equations are extremely flexible and can straightforwardly incorporate a mixture of antagonistic and mutualistic interactions simply by altering the signs of entries in the community matrix (Mougi and Kondoh, 2012; Allesina and Tang, 2012). But for consumer-resource models we have to be careful about how those mechanisms are formulated. Consumption can take a variety of forms depending, for example, on whether resources are substitutable or essential (Tilman, 1980), and mutualistic interactions can occur via resource production and exchange (i.e., cross-feeding) (Loreau, 2001; Freilich et al., 2011; Zeleznik et al., 2015; Sun et al., 2019). The latter in particular is underexplored relative to consumption (Butler and O’Dwyer, 2018), and an open question is to what extent the precise form of exchange might affect community dynamics and species coexistence.

So how do we retain the advantages of consumer-resource models, but also incorporate a mixture of species interactions? Current consumer-resource models are largely agnostic to what happens inside cells or organisms. For many systems, this approach may be valid especially when the resources (i.e., prey) belong to higher trophic levels, self-regulate, and/or closely match the stoichiometric requirements of the consumer (Stern and Elser, 2002; Cherif and Loreau, 2007; Hall, 2009). However, this may not be valid for microorganisms. First, most microorganisms consume abiotic resources which do not self-regulate. Second, any single resource generally does not satisfy the full stoichiometric requirements of metabolism. For example, heterotrophic microorganisms require organic carbon, but they still require nitrogen, phosphorus, and other resources to grow and reproduce. Because growth depends on multiple resources, population dynamics may depend on a single limiting resource (e.g., Liebig’s Law of the Minimum: von Liebig and Gregory, 1842; Odum, 1959) or an interaction between resources (e.g., multiplicative co-limitation: Harpole et al., 2011). Likewise, the consumption and transformation of resources depends on how cells produce biomass, energy, and metabolic byproducts, and species interactions may therefore depend on metabolic rates and the release of metabolic byproducts.

Here, we propose that going one level deeper into cellular metabolism will allow us to generalize consumer-resource models in a meaningful way and give them the same flexibility to describe multiple interaction types as models of direct species interactions. First, we introduce a general metabolically informed consumer-resource model for a single species. In this model, we include internal cellular dynamics via a simplified metabolic network, which require less knowledge of a particular species’ idiosyncrasies but still captures the major metabolic events transforming resources. Using this model, we explore population dynamics in variable resource environments. Next, we expand the approach to

incorporate a second species. Here, both species share a common resource, but the second species also uses a metabolic byproduct produced by the first species. As such, our model includes competition for the shared resource and facilitation via metabolic cross-feeding. We then explore how the resource landscape and internal metabolic rates interact to demonstrate how competition and facilitation mediate species dynamics and coexistence conditions under different resource use scenarios. In summary, our findings demonstrate how a metabolically informed consumer-resource model for a single species can produce both Liebig's Law and multiplicative co-limitation in particular limits of the metabolic rates. Furthermore, when a second species is included we expand the expectations of Liebig's Law and multiplicative co-limitation to include cross-feeding and demonstrate both species dynamics and equilibrium abundances across resource landscapes.

Methods and Results

The Metabolically Informed Consumer-Resource Model. We know a substantial amount about the internal physiology of microorganisms, and there have been large advancements in the development of flux balance models which use biochemistry and genomics to describe (to some level of approximation) every reaction that occurs within a cell (Kauffman et al., 2003; Orth et al., 2010). Likewise, models based on the dynamic energy budget theory provide a complete mass and energy balance approach for molecular processes, cellular physiology, and population growth (Kooijman, 2001; Nisbet et al., 2008). More recently, flux balance models have been applied in the context of entire microbial communities and their interactions (Embree et al., 2015; Zomorodi and Segrè, 2016; Pacheco et al., 2019). However, we propose that including a full description of a metabolic network may not be required to develop a useful ecological model. Simplified metabolic models have been used to understand metabolic partitioning (Kempes et al., 2012) and trophic tradeoffs (Chakraborty et al., 2017). Furthermore, models that add some intracellular dynamics, such as Droop's model which accounts for cell quotas and intracellular resource storage (Droop, 1974), have been used to better understand how consumers contend with nutrient limitations and how trait trade-offs underlie coexistence (Cherif and Loreau, 2010; Litchman et al., 2015). Here, we model the internal dynamics of a microorganism via simplified metabolic networks, which require less knowledge of the particular species' idiosyncrasies but still capture the major metabolic events transforming resources. Our prototypical metabolic model is based on a basic fermentation reaction, homolactic fermentation, which uses glucose and phosphate and produces lactate (Figs. 1 & S1). In this reaction, glucose and phosphate are used to form new biomass, and glucose is used to produce chemical energy via fermentation. The energetic component results in the

Fig. 1 Conceptual Model. Our metabolically informed consumer-resource model uses fermentation as a prototype. Fermentation is an anaerobic—usually sugar consuming—metabolic lifestyle, and is the primary anaerobic energy-producing reaction for many microorganisms (Gottschalk, 1986). A signature of fermentation is that it results in byproducts such as organic acids, alcohols, and/or gases, which are produced due to the incomplete resource oxidation during the energy producing reactions. When organic acids (e.g., lactate) are produced, they are often used as resources by other microorganisms. In our model, we simplify the true biochemistry (see Supplemental, Fig. S1) by assuming glucose (R_1) and phosphate (R_2) enter the cell at uptake rates k_i , biomass (N_1) is generated at rate $\beta M_1 M_2$, and the metabolic byproduct, lactate, is exported from the cell at rate v_3 . In addition, we assume that any unused glucose and phosphate are exported at rates v_1 and v_2 .

production of a byproduct, lactate, which is exported back into the environment (i.e., excretion); therefore, efficiency is emergent property determined by the balance between the biomass and energy production.

Here, we assume that cellular metabolism relies on the interaction of sugar and phosphate, producing new biomass and a byproduct (i.e., lactate).

R_1 here is the sugar, R_2 represents a source of phosphorus, and R_3 is the metabolic byproduct lactate. We model this intracellular reaction as occurring at rate β . Next, we assume that the organism grows in a chemostat-type environment, where resource inflow rates are held constant. In addition, we assume that metabolic rates are not limited by any other factors and that resources are not toxic at high concentrations and therefore do not inhibit growth. Given that we consider one phosphate molecule (i.e., we assume 1:1 stoichiometry), this already simplifies the resource requirements relative to the true process. But we will see general principles emerge.

We now assume that intracellular processes are always close to equilibrium, so that we can balance fluxes for the internal cellular densities of the two input resources (labeled M_1 and M_2). This leads to:

$$\begin{aligned} v_{bio} &= \beta M_1 M_2 \\ v_1 &= k_1 - \beta M_1 M_2 \\ v_2 &= k_2 - \beta M_1 M_2 \\ v_3 &= \beta M_1 M_2, \end{aligned} \tag{2}$$

for uptake rates k_1 and k_2 , which can depend in an arbitrary way on external resource concentrations R_1 , R_2 , and R_3 , outflow (i.e., export) rates v_1 , v_2 , and v_3 , and biomass production v_{bio} (Fig. 1). We make a natural assumption that export of each resource is determined by passive excretion: i.e. that $v_\alpha = \lambda_\alpha M_\alpha$ for each metabolite, where λ_α is a species- and resource-specific constant. This resource excretion, or leakiness, prevents unbounded internal resource accumulation, but it can also mediate species interactions (e.g., cross-feeding) and therefore structure communities (Pfeiffer and Bonhoeffer, 2004; Morris, 2015; Sun et al., 2019). As such, the outflow rates (v_α) represent *use it or lose it* internal resource dynamics and therefore intracellular resources that are not used are exported back into the external environment.

We can solve this system of polynomial equations (Eq. 2) for internal resource concentrations to obtain:

$$\begin{aligned} M_1 &= \frac{k_1 - k_2}{2\lambda_1} - \frac{\lambda_2}{2\beta} + \frac{1}{2\lambda_1} F(\beta, k_1, k_2) \\ M_2 &= \frac{k_2 - k_1}{2\lambda_2} - \frac{\lambda_1}{2\beta} + \frac{1}{2\lambda_2} F(\beta, k_1, k_2), \end{aligned} \tag{3}$$

while biomass production is given by:

$$v_{bio} = \frac{\lambda_1 \lambda_2}{2\beta} - \frac{F(\beta, k_1, k_2)}{2} + \frac{k_1 + k_2}{2}, \tag{4}$$

and where $F(\beta, k_1, k_2) = \sqrt{\left(k_1 + k_2 + \frac{\lambda_1 \lambda_2}{\beta}\right)^2 - 4k_1 k_2}$ depends on uptake, export, and reaction rates.

Using these intracellular concentrations, we can now develop a set of generalized consumer-resource equations using uptake rates and this simplified flux balance analysis (Eq. 4) as the only building blocks. First, we focus on one species and model its uptake rate k_α of resource α as $C_\alpha R_\alpha$, where R_α is the *external* (environmental) concentration of this

Table 1 Model Notation. Note: resource concentrations are given in cell equivalents – moles required to generated one cell.

Symbol	Meaning	Units
N_i	Number of cells in population i	cell
R_α	Concentration of resource α	cell equivalent
ρ_α	Inflow rate of resource α	cell equivalent hr^{-1}
η_α	Outflow rate of resource α	cell equivalent hr^{-1}
$C_{\alpha i}$	Uptake rate of resource α by population i	cell equivalent hr^{-1} cell $^{-1}$
v_i	Mortality rate of population i	cell hr^{-1}
λ_α	Leakage rate of resource α	cell equivalent hr^{-1}
β_i	Internal metabolic rate of population i	cell hr^{-1}

resource. Applying this assumption, we derive a general set of equations for the consumer and three resources:

$$\begin{aligned}\frac{dR_1}{dt} &= \rho_1 - \eta_1 R_1 - R_1 C_{11} N_1 + v_1 N_1 \\ &= \rho_1 - \eta_1 R_1 - N_1 \left(\frac{\lambda_1 \lambda_2}{2\beta} - \frac{F(\beta, C_{11} R_1, C_{21} R_2)}{2} + \frac{C_{11} R_1 + C_{21} R_2}{2} \right)\end{aligned}\quad (5a)$$

$$\begin{aligned}\frac{dR_2}{dt} &= \rho_2 - \eta_2 R_2 - R_2 C_{21} N_1 + v_2 N_1 \\ &= \rho_2 - \eta_2 R_2 - N_1 \left(\frac{\lambda_1 \lambda_2}{2\beta} - \frac{F(\beta, C_{11} R_1, C_{21} R_2)}{2} + \frac{C_{11} R_1 + C_{21} R_2}{2} \right)\end{aligned}\quad (5b)$$

$$\begin{aligned}\frac{dR_3}{dt} &= \rho_3 - \eta_3 R_3 + v_3 N_1 \\ &= \rho_3 - \eta_3 R_3 + N_1 \left(\frac{\lambda_1 \lambda_2}{2\beta} - \frac{F(\beta, C_{11} R_1, C_{21} R_2)}{2} + \frac{C_{11} R_1 + C_{21} R_2}{2} \right)\end{aligned}\quad (5c)$$

$$\begin{aligned}\frac{dN_1}{dt} &= v_{bio} N_1 - \mu_1 N_1 \\ &= N_1 \left(\frac{\lambda_1 \lambda_2}{2\beta} - \frac{F(\beta, C_{11} R_1, C_{21} R_2)}{2} + \frac{C_{11} R_1 + C_{21} R_2}{2} \right) - \mu_1 N_1,\end{aligned}\quad (5d)$$

where ρ_α and η_α are inflow and outflow rates for each of the three resources, and μ_1 is the mortality rate of the consumer (see Table 1). For the resources (Eqs. 5a–5c), the extracellular resource concentrations are determined by the inflow and outflow rates minus the consumer-density dependent components which include uptake, leakage, and conversion into biomass (Eq. 4). For the consumer (Eq. 5d), the production of new biomass is determined by the density dependent uptake, leakage, and conversion of resources into biomass (Eq. 4) minus the density dependent mortality. Here, we have assumed that uptake, k_α , increases linearly with resource concentration, but this could easily be modified to saturate using Monod dynamics by changing k_α to $(C_{\alpha 1} R_\alpha)/(K_{\alpha 1} + R_\alpha)$, where K , the half-saturation constant, is a species- and resource-specific constant.

We now note two limits of the simplified flux balance analysis (Eqs 3–4). The first limit is where β (i.e., the internal metabolic rate) is large relative to the other rates,

$$\begin{aligned} M_1 &\simeq \frac{k_1 - k_2 + |k_1 - k_2|}{2\lambda_1} + \frac{\lambda_2}{2\beta} \left(\frac{k_1 + k_2}{|k_1 - k_2|} - 1 \right) + O(1/\beta^2) \\ M_2 &\simeq \frac{k_2 - k_1 + |k_1 - k_2|}{2\lambda_2} + \frac{\lambda_1}{2\beta} \left(\frac{k_1 + k_2}{|k_1 - k_2|} - 1 \right) + O(1/\beta^2) \\ v_1 &= \lambda_1 M_1 \\ v_2 &= \lambda_2 M_2 \\ v_3 &= v_{bio} \simeq \min(k_1, k_2) + O(1/\beta). \end{aligned} \quad (6)$$

Where $O(1/\beta)$ represents the linear approximation of the higher order terms of the Taylor Series expansion. Note that we have to keep the $O(1/\beta)$ terms for M_1 and M_2 , because for at least one of the two (depending on whether $k_1 > k_2$ or $k_1 < k_2$) the $O(1/\beta)$ term vanishes in this limit of fast reaction rate β . Therefore, when β is large relative to the other rates the consumer-resource equations become:

$$\begin{aligned} \frac{dR_1}{dt} &= \rho_1 - \eta_1 R_1 - R_1 C_{11} N_1 \\ &\quad + \frac{N_1}{2} \left[R_1 C_{11} - R_2 C_{21} + |R_1 C_{11} - R_2 C_{21}| + \frac{\lambda_1 \lambda_2}{\beta} \left(\frac{R_1 C_{11} + R_2 C_{21}}{|R_1 C_{11} - R_2 C_{21}|} - 1 \right) \right] \\ \frac{dR_2}{dt} &= \rho_2 - \eta_2 R_2 - R_2 C_{21} N_1 \\ &\quad + \frac{N_1}{2} \left[R_2 C_{21} - R_1 C_{11} + |R_1 C_{11} - R_2 C_{21}| + \frac{\lambda_1 \lambda_2}{\beta} \left(\frac{R_1 C_{11} + R_2 C_{21}}{|R_1 C_{11} - R_2 C_{21}|} - 1 \right) \right] \\ \frac{dR_3}{dt} &= \rho_3 - \eta_3 R_3 + N_1 \min(R_1 C_{11}, R_2 C_{21}) \\ \frac{dN_1}{dt} &= N_1 \min(R_1 C_{11}, R_2 C_{21}) - \mu_1 N_1. \end{aligned} \quad (7)$$

Hence, we recover Liebig’s law (i.e. the net growth rate of the consumer is $\min(R_1 C_{11}, R_2 C_{21})$). Using these equations in numerical simulations, we observe that the consumer abundance saturates (Fig. 2A), and the final abundance depends on the inflow rate of the more limiting resource (Fig. 2B). As such, the model in this limit behaves similar to classical consumer-resource models. In addition, we generate novel terms for the byproduct production rate, which in this fast reaction rate limit is $\simeq N_1 \min(R_1 C_{11}, R_2 C_{21})$, and we find corresponding equations for the uptake and export of glucose and phosphate.

Fig. 2 Single Species Simulation. We used numerical simulations of our single-species model to determine how the resource landscape changes population dynamics. In our model, N_1 is the population, R_1 and R_2 are the consumed resources (i.e., sugar and phosphate), R_3 is a metabolic byproduct, ρ is the resource inflow rate, and β is the internal metabolic rate of the consumer. **A & B:** In the limit where β is high relative to other rates (Eq. 7), we recover Liebig's Law of the Minimum. Consumer abundance saturates, but the final abundance depends on the inflow rate, ρ_α , of the more limiting resource. When the inflow of R_2 (ρ_2) is lower than the inflow of R_1 (ρ_1), R_2 becomes depleted. At saturation, R_1 is in excess while R_2 is limiting (A). Changes in the inflow rates alters the final abundance (B), but the final abundance will always be determined by the more limiting resource. When $\rho_1 = 100$ (blue) the final abundance will increase until $\rho_2 = 100$ (blue dashed line), and when $\rho_1 = 108$ (red) the final abundance will increase until $\rho_2 = 108$ (red dashed line). In contrast, in the limit where β is low relative to other rates (Eq. 9), we recover multiplicative co-limitation. Consumer abundance still saturates, albeit at a lower final abundance given the same resource inflow rates and with a growth lag-phase (C). As consumer abundance increases, both R_1 and R_2 are depleted. In addition, the final abundance increases as inflow rates increase (D). When $\rho_1 = 100$ (blue) the final abundance will continue to increase past $\rho_2 = 100$, and when $\rho_1 = 108$ (red) the final abundance will continue to increase past $\rho_2 = 108$. Simulations based on Eq. 5 using the following parameters: $\eta_\alpha = 0.01$, $C_{\alpha 1} = 0.01$, $\mu_1 = 0.01$, $\lambda_\alpha = 0.1$, $\beta_1 = 10$ (A) or 10^{-6} (B), $\rho_1 = 100$ or 108, and ρ_2 varies between 80 and 120 (A & C: $\rho_1 = 100$, $\rho_2 = 80$). Resource concentrations are expressed in cell equivalents (i.e., moles required to generate a cell) and we assume 1:1 resource stoichiometry.

145 The second limit is where β (i.e., the internal metabolic rate) is small relative to the other rates,

$$\begin{aligned}M_1 &\simeq \frac{k_1}{\lambda_1} - \frac{\beta k_1 k_2}{\lambda_1^2 \lambda_2} + O(\beta^2) \\M_2 &\simeq \frac{k_2}{\lambda_2} - \frac{\beta k_1 k_2}{\lambda_1 \lambda_2^2} + O(\beta^2) \\v_1 &= \lambda_1 M_1 \\v_2 &= \lambda_2 M_2 \\v_3 &= v_{bio} \simeq \beta \frac{k_1 k_2}{\lambda_1 \lambda_2} + O(\beta^2),\end{aligned}\tag{8}$$

146 so that the consumer-resource equations become:

$$\begin{aligned}\frac{dR_1}{dt} &= \rho_1 - \eta_1 R_1 - R_1 C_{11} N_1 + N_1 \left(C_{11} R_1 - \frac{\beta C_{11} R_1 C_{21} R_2}{\lambda_1 \lambda_2} \right) \\ \frac{dR_2}{dt} &= \rho_2 - \eta_2 R_2 - R_2 C_{21} N_1 + N_1 \left(C_{21} R_2 - \frac{\beta C_{11} R_1 C_{21} R_2}{\lambda_1 \lambda_2} \right) \\ \frac{dR_3}{dt} &= \rho_3 - \eta_3 R_3 + N_1 \frac{\beta C_{11} R_1 C_{21} R_2}{\lambda_1 \lambda_2} \\ \frac{dN_1}{dt} &= N_1 \frac{\beta C_{11} R_1 C_{21} R_2}{\lambda_1 \lambda_2} - \mu_1 N_1.\end{aligned}\tag{9}$$

147 Hence, we recover multiplicative co-limitation by the two resources (i.e., the net growth rate of the consumer is
148 $\frac{\beta C_{11} R_1 C_{21} R_2}{\lambda_1 \lambda_2}$). Using numerical simulations, we observe that consumer abundance, N_1 saturates as expected but now
149 includes a growth lag-phase (Fig. 2C). The lag-phase (i.e., period with no net population growth: $\frac{dN}{dt} = 0$) occurs while
150 $\beta < \frac{\mu_1 \lambda_1 \lambda_2}{C_{11} R_1 C_{21} R_2}$. When R_1 and R_2 accumulate (Fig. 2C), the denominator increases and the inequality flips which yields
151 population growth. Due to the parameters used in the simulation above, the equation for the lag-phase simplifies to
152 $\beta = \frac{1}{R_1 R_2}$; therefore, our simulation demonstrates how the lag-phase is controlled by an interaction between the resource
153 environment and the internal metabolic rates. Finally, the final abundance depends on the inflow rate of both resources
154 (Fig. 2D). An increase in the inflow rate of either resource will yield a higher final population abundance.

155 In summary, from this coarse-grained representation, we recover two classic outcomes of consumer-resource theory
156 by taking limits of the internal reaction rate β . We can also generalize these classic limits for intermediate β , in a
157 way that is not currently used in consumer-resource models and falls neither into the category of Liebig's Law nor
158 multiplicative co-limitation. Finally, we find functional forms for the production rate of metabolic byproducts (and
159 excretion of other resources) in each of these limits and in between. Our model therefore demonstrates how we can
160 generalize the functional form of consumer-resource models by considering realistic, simplified intracellular processes.

Fig. 3 Two Species Model. In the two species model, two species interact through competition and facilitation. For each species, we use a simplified metabolic networks to describe resource uptake rates ($k_{\alpha i}$), biomass generation rates ($\beta_i M_{\alpha i} M_{\alpha i}$), and metabolic byproduct export rates ($v_{\alpha i}$) for species i and resource α . Species N_1 consumes glucose (R_1) and phosphate (R_2) and produces lactate (R_3). Species N_2 competes for phosphate (R_2) but also uses the lactate produced by species N_1 . In addition, all internal resources (byproducts and unused resources) are exported at rate v_{α} (not shown). As such, facilitation is modeled by cross-feeding (i.e., production and consumption of lactate) between species N_1 and N_2 .

The Two Species Model. We now modify the model above to incorporate a second species. Here, both species use R_2 (phosphate), but the second species, N_2 , uses a combination of R_2 and R_3 (lactate) to generate new biomass (Fig. 3). While we are using this as a model with both competition (e.g., shared resources) and facilitation (e.g., metabolic cross-feeding) interactions, it also represents the natural cross-feeding interaction between organisms. For example, the model could represent the interactions between the lactate producing (e.g., *Bifidobacterium adolescentis*) and lactate consuming (e.g., *Eubacterium hallii*) bacteria found in human and animal digestive systems (Duncan et al., 2004). Likewise, other examples of these interactions would include the exchange of vitamin precursors and amino acids by auxotrophic organisms (Carini et al., 2014; Embree et al., 2015). These metabolic cross-feeding interactions are common in microbial systems (Pfeiffer and Bonhoeffer, 2004; Morris et al., 2012; Mee et al., 2014; Morris, 2015; Tasoff et al., 2015; Zelezniak et al., 2015; Sun et al., 2019) and have industrial applications (Jiao et al., 2012). Cross-feeding has been shown to promote species diversity and increase net ecosystem production (Morris et al., 2012; Pande et al., 2014). In our model, productivity (and efficiency) increase due to the second species using the metabolic byproducts of the first (Fig. 3). As such, our model demonstrates how competition (i.e., shared resources) and facilitation (i.e., production and consumption of intermediate resources) mediate species dynamics and coexistence conditions and can be used to understand natural and engineered microbial systems.

First, we define two distinct internal metabolic processes, one for each consumer species:

Consumer N_2 may also produce a metabolic byproduct, but we have not included such a process here because we are focusing on competition for R_2 and facilitation through the production of R_3 by species N_1 (Fig. 3). Importantly, we now have two internal reaction rates, β_1 and β_2 . Here we focus on how these rates, both relative to each other and also to the other rates in the model, affect species coexistence. This approach allows us to use the coarse-grained metabolic processes as the mechanisms underlying species interactions. Furthermore, it allows us to explore the potential for changes in species coexistence due to metabolic (i.e., reaction rates) and landscape (i.e., inflow rates) factors.

Generalizing the approach in the previous section, and again assuming internal flux balance, we can define the fluxes for an individual belonging to species N_1 as:

$$\begin{aligned} v_{bio,1} &= \beta_1 M_{11} M_{21} \\ v_{11} &= k_{11} - \beta_1 M_{11} M_{21} \\ v_{21} &= k_{21} - \beta_1 M_{11} M_{21} \\ v_{31} &= \beta_1 M_{11} M_{21}, \end{aligned} \quad (11)$$

for internal concentrations $M_{\alpha 1}$ and uptake rates $k_{\alpha 1}$. While for an individual of species N_2 we have:

$$\begin{aligned} v_{bio,2} &= \beta_2 M_{22} M_{32} \\ v_{22} &= k_{22} - \beta_2 M_{22} M_{32} \\ v_{32} &= k_{32} - \beta_2 M_{22} M_{32}, \end{aligned} \quad (12)$$

for internal concentrations $M_{\alpha 2}$ and uptake rates $k_{\alpha 2}$. We also assume that all resources can be excreted from both consumers, but to simplify the model slightly we will assume that the specific export rates are equal, λ , broadly consistent with passive diffusion across sufficiently similar cell wall types. Based on these assumptions, we can solve for internal equilibrium in both cell types to obtain:

$$\begin{aligned} v_{bio,1} &= \frac{\lambda^2}{2\beta_1} - \frac{F(\beta_1, k_{11}, k_{21})}{2} + \frac{k_{11} + k_{21}}{2} \\ v_{bio,2} &= \frac{\lambda^2}{2\beta_2} - \frac{F(\beta_2, k_{22}, k_{32})}{2} + \frac{k_{22} + k_{32}}{2}, \end{aligned} \quad (13)$$

where, similarly to the one species case, the function $F(x, a, b) = \sqrt{\left(a + b + \frac{\lambda^2}{x}\right)^2 - 4ab}$ (see: Eqns: 3 & 4). Finally, we can put all of this together to generate a set of equations for the dynamics of both populations and the external concentrations of all three resources.

To focus on the effects of internal reaction rates and the resource landscape, we will further simplify our model by making a few assumptions. We will assume that the outflow rates for each resource are the same, so that $\eta_i = \eta$, and that the per capita mortality rates for each consumer are equal, so that $\mu_i = \mu$. We will also again assume that the per capita uptake rate of resource α by species i can be written as $C_{\alpha i} R_{\alpha}$. We will then determine the effects of internal reaction rate by independently changing β (i.e., the internal metabolic rate) for each species. In addition, we will change the landscape conditions by exploring the inflow rates for each resource ρ_{α} . Given these assumptions, our general two species consumer-resource model is:

$$\begin{aligned} \frac{dR_1}{dt} &= \rho_1 - \eta R_1 - R_1 C_{11} N_1 + v_{11} N_1 \\ &= \rho_1 - \eta R_1 - N_1 \left(\frac{\lambda^2}{2\beta_1} - \frac{F(\beta_1, C_{11} R_1, C_{21} R_2)}{2} + \frac{C_{11} R_1 + C_{21} R_2}{2} \right) \end{aligned} \quad (14a)$$

$$\begin{aligned} \frac{dR_2}{dt} &= \rho_2 - \eta R_2 - R_2 C_{21} N_1 + v_{21} N_1 - R_2 C_{22} N_2 + v_{22} N_2 \\ &= \rho_2 - \eta R_2 \\ &\quad - N_1 \left(\frac{\lambda^2}{2\beta_1} - \frac{F(\beta_1, C_{11} R_1, C_{21} R_2)}{2} + \frac{C_{11} R_1 + C_{21} R_2}{2} \right) \\ &\quad - N_2 \left(\frac{\lambda^2}{2\beta_2} - \frac{F(\beta_2, C_{22} R_2, C_{32} R_3)}{2} + \frac{C_{22} R_2 + C_{32} R_3}{2} \right) \end{aligned} \quad (14b)$$

$$\begin{aligned} \frac{dR_3}{dt} &= \rho_3 - \eta R_3 + v_{31} N_1 - R_3 C_{32} N_2 + v_{32} N_2 \\ &= \rho_3 - \eta R_3 \\ &\quad + N_1 \left(\frac{\lambda^2}{2\beta_1} - \frac{F(\beta_1, C_{11} R_1, C_{21} R_2)}{2} + \frac{C_{11} R_1 + C_{21} R_2}{2} \right) \\ &\quad - N_2 \left(\frac{\lambda^2}{2\beta_2} - \frac{F(\beta_2, C_{22} R_2, C_{32} R_3)}{2} + \frac{C_{22} R_2 + C_{32} R_3}{2} \right) \end{aligned} \quad (14c)$$

$$\begin{aligned} \frac{dN_1}{dt} &= N_1 v_{bio,1} - \mu N_1 \\ &= N_1 \left(\frac{\lambda^2}{2\beta_1} - \frac{F(\beta_1, C_{11} R_1, C_{21} R_2)}{2} + \frac{C_{11} R_1 + C_{21} R_2}{2} \right) - \mu N_1 \end{aligned} \quad (14d)$$

$$\begin{aligned} \frac{dN_2}{dt} &= N_2 v_{bio,2} - \mu N_2 \\ &= N_2 \left(\frac{\lambda^2}{2\beta_2} - \frac{F(\beta_2, C_{22} R_2, C_{32} R_3)}{2} + \frac{C_{22} R_2 + C_{32} R_3}{2} \right) - \mu N_2, \end{aligned} \quad (14e)$$

where ρ_{α} and η_{α} are inflow and outflow rates for each of the three resources, and μ_1 is the mortality rate of the consumer (see Table 1). For the resources (Eqs. 14a–14c), the extracellular resource concentrations are determined by the inflow

and outflow rates minus the consumer-density dependent components of each consumer which include uptake, leakage, and conversion into biomass (Eq. 13). For the consumers (Eqs. 14d & 14e), the production of new biomass is determined by the density dependent uptake, leakage, and conversion of resources into biomass (Eq. 13) minus the density dependent mortality.

Using numerical simulations, we model the resource and consumer dynamics and determine equilibrium conditions (Figs. 4 & 5). First we consider when the internal metabolic rates, β_i , are the same. When internal metabolic rates are both high, species coexist at a density determined by the shared resource inflow rate (i.e., ρ_2) until the inflow rate of the unshared resource, R_1 , exceeds the inflow rate of the shared resource, R_2 . When ρ_1 is greater than ρ_2 (i.e., $\rho_1/\rho_2 > 1$), species N_1 will outcompete species N_2 for R_2 , and N_2 will become rare (Fig. 4A). In our simulations, species N_2 remains in the system with rare abundance due to resource leakage from species N_1 , but would likely go extinct due to stochastic fluctuations, and so coexistence in this limit is somewhat tenuous. These findings expand the expectations of Liebig's Law to two cross-feeding species and demonstrate both species dynamics and equilibrium abundances across various resource landscapes. In contrast, when internal metabolic rates are low, species relative abundances are determined by their respective required resources based on multiplicative co-limitation and both species now demonstrate growth lag-phases. These findings expand the expectations of multiplicative co-limitation to two cross-feeding species. However, since species N_1 will not be resource limited when ρ_1 is greater than ρ_2 , then species N_2 will maintain a higher relative abundance across wider resource landscape (Fig. 4B, see Eq. 9). Together, we find that if internal rates are the same but either high or low compared to the other rates in our model, then our results expand the expectations of Liebig's Law and multiplicative co-limitation to the two-species system with a metabolic dependency. Between these limits, we find a smooth transition between the two scenarios as coexistence conditions become decoupled from classical resource limitation predictions when the internal metabolic rates decrease (Fig. 4C). In addition, we find that coexistence depends on both the internal metabolic rates and the resource inflow rates even when uptake rates are the same. Furthermore, while species N_2 depends on species N_1 for the production of R_3 , external inputs of R_3 (i.e., $\rho_3 > 0$) alone are not enough to allow species N_2 to outcompete species N_1 but will result in a lower final density (see Supplemental).

Finally, we consider the dynamics and equilibria when the internal metabolic rates, β_i , are different. We find that, when the rates differ the outcome depends on which species has the higher metabolic rate. When the byproduct producer, N_1 , has the higher rate, then the results are similar to when both species have high internal metabolic rates (Fig. 5A). We do note, however, two important differences: 1) species N_2 alone exhibits a growth lag-phase, and 2) both R_1 and R_2 are depleted as the species reach an equilibrium. However, when species N_2 has the higher rate the coexistence

Fig. 4 Two Species with the Same Metabolic Rates. We simulated species dynamics in the two-species model with equal internal metabolic rates, β_i . In this model, two species compete for a shared resource (R_2), but N_1 consumes R_1 and produces R_3 as a metabolic byproduct which is consumed by N_2 . When metabolic rates are high, species abundances saturate and species can coexist provided the appropriate resource conditions (A). When the shared resource inflow rate (ρ_2) is higher than the inflow rate of R_1 (ρ_1), then the two species coexist at a relatively high abundance because N_1 is more limited by its exclusive resource (R_1). However, when ρ_1 is higher than ρ_2 (i.e., $\rho_1/\rho_2 > 1$), then N_1 can outcompete N_2 for the shared resource (R_2) and the final abundance of N_2 will be reduced. Even when the shared resource inflow rate is increased ($\rho_2 = 100$ vs. 80), the final abundance of N_2 does not increase when $\rho_1/\rho_2 > 1$. When metabolic rates are low, species still coexist, but the range of coexistence is greatly expanded (B). While the final abundance of N_2 still decreases as the resource inflow ratio (ρ_1/ρ_2) approaches 1, it does not decrease at the same rate and the final abundance remains relatively high even when the resource inflow ratio is > 1 . In this simulation, the resource inflow ratio needs to be greater than around ~ 1.5 before N_1 outcompetes N_2 and N_2 becomes rare. Between these metabolic rates, there is a smooth transition (C). When both resources have equal inflow rates, the final abundance of N_1 decreases and N_2 increases as β drops below 1. Simulations based on Eqs. 14a–14e using the following parameters: $\eta_\alpha = 0.01$, $C_{ai} = 0.01$, $\mu_i = 0.01$, $\lambda_\alpha = 0.1$, $\beta_i = 10$ (high) or 10^{-6} (low), $\rho_2 = 80$ or 100, and ρ_1/ρ_2 varies between 0.5 and 1.7 (A_{left} & B_{left}: $\rho_1 = 64$, $\rho_2 = 80$). Resource concentrations are expressed in cell equivalents (i.e., moles required to generate a cell) and we assume 1:1 resource stoichiometry.

conditions and high relative abundances for both species are greatly expanded. In fact, we find coexistence with moderate abundances along all inflow rates tested and the final abundances of both species are determined only by the shared resource, R_2 (Fig. 5B). In addition, we find that both species exhibit growth lag-phases and that R_2 and R_3 are now the depleted resources. Between these limits, we find that the transition from classical resource limitation prediction to complete coexistence depends on both metabolic rates decreasing (Fig. 5C). These findings highlight how variation in both the internal metabolic reaction rates and in the environmental conditions can influence species interactions and change coexistence expectations.

Discussion

Here, we reformulated the classic consumer-resource model by independently including resource uptake, internal metabolic rates, and byproduct export. We found that when internal metabolic dynamics are included – in addition to uptake – two common models of resource limitation (Liebig’s Law and multiplicative co-limitation) appear in particular limits of the internal reaction rates. In this model, we make predictions for the functional form of the production of metabolic byproducts which can be used in metabolic exchanges, and we balance the requirements for growth, energy, storage, and export. Because our model includes an independent term for internal metabolic rates, it provides a flexible and more general approach to understand how the resource landscape and other physical variables affect species dynamics in microbial communities. We further expand our model to include a second species which uses the metabolic byproduct of the first species. In this metabolically-informed approach, both species interactions (competition and facilitation) and resource use efficiency are emergent properties of the system. In addition, we show how internal metabolic reaction rates and the resource landscape determine species dynamics and equilibria. We find that the growth dynamics change when resources are *metabolically* limiting which depends on the internal metabolic rates. Therefore our model shows how metabolic rates and the resource landscape change the interactions between cross-feeding species which can shift the contributions of facilitation and competition at equilibrium. We further show that changes in the environmental landscape and/or metabolic rates can change resource and consumer dynamics in ways that alter species-species and species-resource correlations (see resource insets in Figs. 4 & 5); therefore, correlations may not accurately reflect interactions (see: Barner et al., 2018).

Classic formulations for pairwise interactions and consumer-resource dynamics have each led to insights regarding species coexistence, community stability, population self-regulation (Schoener, 1983; Barabás et al., 2017; Allesina and Tang, 2012; Leibold and McPeck, 2006). Lotka-Volterra (and related) equations provide a flexible approach to modeling

Fig. 5 Two Species with Different Metabolic Rates. We simulated species dynamics in the two-species model where the species have different internal metabolic rates, β_i . In this model, one of the species has high and the other low metabolic rates compared to the other rates in the model. The species compete for R_2 , and species N_1 consumes R_1 and produces R_3 which is used by species N_2 . When N_1 has the higher metabolic rate, the species dynamics are similar and the coexistence conditions are identical to the model where both species have high metabolic rates (A). However, there are now two important differences. First, N_2 (gray line) has a growth lag-phase. Second, both R_1 and R_2 are now depleted as the population abundance saturates. When N_2 has the higher metabolic rates, the coexistence conditions are drastically different (B). N_1 and N_2 have growth lag-phases, and R_2 and R_3 are now the depleted resources. In addition, both species coexist at a final abundance determined only by the shared resource inflow rate (ρ_2) across all resource inflow rates (ρ_1/ρ_2) tested. When we independently change β_1 and β_2 , the final abundances depend on both values (C). As β_1 decreases from 1 to 10^{-4} (blocks), the final abundance of N_1 decreases and N_2 increases for a given resource inflow ratio. In addition, as β_2 increases from 10^{-4} to 1 (within each block), the final abundance of N_1 decreases more and N_2 increases for a given inflow ratio. Simulations based on Eqs. 14a–14e using the following parameters: $\eta_\alpha = 0.01$, $C_{ai} = 0.01$, $\mu_i = 0.01$, $\lambda_\alpha = 0.1$ and $\beta_i = 10$ (high) or 10^{-6} (low), $\rho_2 = 80$ or 100 , and ρ_1/ρ_2 varies between 0.5 and 1.7 (A_{left} & B_{left}: $\rho_1 = 64$, $\rho_2 = 80$). Resource concentrations are expressed in cell equivalents (i.e., moles required to generate a cell) and we assume 1:1 resource stoichiometry.

a range of interactions between species but are unable to generalize across environmental variation because they do not provide an unambiguous way to include the resource landscape. While it may be possible to modify per capita growth rates to be a function of environmental conditions (e.g., temperature) and the resource landscape, how to incorporate these effects in the species interaction rates is ambiguous, at best. However, these models allow positive and negative species interactions to be explored straightforwardly, with far-reaching implications for coexistence and stability. Consumer-resource models explicitly include the interaction between the resource landscape and consumers, but at the expense of introducing more explicit mechanism, and therefore more choices in the way interactions are implemented (O'Dwyer, 2018). Including positive interactions through the production of resources has led to new predictions regarding the stability of communities (Butler and O'Dwyer, 2018, 2019), indicating that incorporating resource exchange (i.e., cross-feeding) may be important for understanding the dynamics of natural microbial communities. However, we do not know how sensitive these results are to the precise way consumption and exchange are formulated. Here, we incorporated metabolism more explicitly into consumer-resource dynamics and it allowed us to explain a broader range of community dynamics and natural phenomena, with less ambiguity in the functional form of interactions, and these metabolic rates may also reveal how environmental conditions that can alter metabolic rates (e.g., temperature) will contribute to species dynamics.

Our model used a simplified metabolic network to describe the internal physiology of cells and this allowed us to determine how uptake, growth, and export contribute to the dynamics and coexistence of interacting species. Specifically, we were able to incorporate competition via shared resource consumption and facilitation via the production and consumption of a metabolic byproduct (i.e., cross-feeding). Our model accomplishes this by using a consumer-resource framework with multiple inputs and outputs and by explicitly incorporating internal resource dynamics (see Figs. 1 & 3). Because we incorporate multiple resources as required inputs, there are similarities between our model and the synthesizing unit concept used in the dynamic energy budget theory (Kooijman, 1998, 2001). For example, both are able to produce limiting resource dynamics based on Liebig's Law. However, our model was also able to produce multiplicative co-limitation, and it allowed us to explore metabolic cross-feeding by providing a mechanism for the production of cellular byproducts. We accomplish this by allowing internal resources, including luxury uptake and metabolic byproducts, to leak out of cells, which establishes strong conceptual similarities to Droop's model (Droop, 1974). Because our model uses this internal flux balance approach, efficiency becomes an emergent property rather than a fixed term. This contrasts with approaches that include fixed efficiency terms that account for the internal processes without providing physiological mechanism (Moore et al., 1993; Schimel and Weintraub, 2003; Moore et al., 2005).

Likewise, our model differs from others that include explicit resource transfer pathways and corresponding efficiencies between species (Wiegert and Owen, 1971; DeAngelis et al., 1975; De Ruiter et al., 1995; Moore et al., 2005) because the byproducts and internal resource excesses are returned to the environment as 'public goods' rather than transferred directly. As such, our model uniquely allows us to incorporate a mixture of species interactions in addition to exploring how each population responds to a complex resource landscape because we include a simplified metabolic network.

Our model allows us to address scenarios that are unique to microbial communities – metabolic cross feeding and internal cellular metabolic rates – and this provided novel insights. First, by providing a mechanism for the production and subsequent consumption of metabolic byproducts (i.e., metabolic cross-feeding) our model provides a way to explore the role of facilitation interactions in a consumer-resource framework. Cross-feeding is a widely observed phenomenon in microbial communities (Morris et al., 2012; Morris, 2015; D'Souza et al., 2018). However, while cross-feeding appears to be ubiquitous, there are very few hypotheses or theoretical prediction for how it will shape the dynamics of interacting species. Our model provides a novel approach to explore this phenomenon and how it contributes to the diversity, complexity, and dynamics of microbial communities. In our model, the coexistence of species N_2 is dependent on the production of a metabolic byproduct, R_3 , from species N_1 . Assuming there is no external input of the byproduct, the coexistence of N_2 is dependent on N_1 . However, due to the way interactions are formulated, external inputs of byproduct alone are not enough to let N_2 outcompete N_1 (see Supplemental). As such, our model provides a new tool to explore the benefits and consequences of cross-feeding in microbial communities. Second, by separating the uptake and internal metabolic rates, our model is able to produce unique species dynamics and it recapitulates two common models of resource limitation (Liebig's Law and multiplicative co-limitation). One major finding based on our model is that internal metabolic rates can have a strong impact on the dynamics of interacting species. Specifically, changes in the internal metabolic rates can alter species coexistence (see Figs. 4 & 5). There are multiple mechanisms that could modify these metabolic rates, including temperature dependence, cofactors, and redox status (Price and Sowers, 2004; Glass and Orphan, 2012; Falkowski et al., 2016; Ramírez-Flandes et al., 2019; Russell and Cook, 1995). Therefore, our model demonstrates how environmental conditions can radically change the relative abundances of interacting species even when resource inflow rates are fixed. Together, these two components of the metabolically-informed consumer-resource model highlight the potential to address questions and scenarios that may be unique to microbial communities.

Our model provides a mechanistic approach to include diverse species interaction in a consumer resource framework and generates novel insights, but it also provides a solid foundation to explore new questions. For example, we have added mechanisms to describe both competition and facilitation based species interactions. These interactions are not

set parameters as in Lotka-Volterra and therefore may be more realistic in terms of modeling natural systems. However, because we have proposed mechanisms for these interactions, some of our conclusions may not be robust due to the precise way we have implemented the specific interactions. This provides an opportunity to explore how changing the implementation yields additional insights and tests new hypotheses. Likewise, we assumed that uptake and export rates increase linearly with resource concentration, but these parts of our equations could easily be modified to incorporate Monod dynamics (as noted above). These simplifications provide a useful starting point from which other hypotheses can be generated and tested. For example, our simplified metabolic models may not capture important idiosyncrasies regarding how metabolism changes along environmental gradients. Perhaps a multi-species genome-scale metabolic model is needed to better describe the metabolic transformations and exchanges between species (Kauffman et al., 2003; Orth et al., 2010; Embree et al., 2015). However, we believe that our simplified version is more tractable, especially when complete genomic information is lacking, and similar approaches have been successfully implemented to describe natural systems (Kempes et al., 2012; Litchman et al., 2015; Chakraborty et al., 2017). The aim of our approach was to find a balance that allows us to more mechanistically describe species interactions while still being tractable for describing natural systems and explaining observed variation.

We propose that when extended more broadly, this approach will lead to more mechanistic predictions for the role of positive interactions along stress gradients (Callaway and Walker, 1997; Brooker and Callaghan, 1998; Koffel et al., 2018), the possibility of species interactions to stabilize or de-stabilize communities (Butler and O'Dwyer, 2018; Allesina and Tang, 2012), and the mechanisms underlying biodiversity-ecosystem function relationships (Duffy et al., 2007; Flynn et al., 2011; Cardinale et al., 2012). Species interactions often change across environmental gradients (Brooker and Callaghan, 1998; He et al., 2013) and a shift that promotes facilitation may increase ecosystem function (e.g., productivity; see: Cardinale et al., 2002). Likewise, increases in nutrient availability has been shown to favor competition over facilitation (Koffel et al., 2018) and can decrease the importance of mutualisms (Keller and Lau, 2018). Our model provides a mechanistic approach to understand these shifts based on the resource landscape and internal metabolic rates. In addition, our model demonstrates that even when competition is favored over facilitation, two species can still coexist. Other studies suggest that positive species interactions, such as facilitation mediated through cross-feeding, can promote coexistence via negative frequency dependence (Morris, 2015). Our model demonstrates how even when competition is favored between cross-feeding species, the poorer competitor can remain as a rare member of the community due to the leakiness of the cross-feeder interaction; although, we predict that stochastic fluctuations could lead to exclusion. As such, our model can be used to better understand why some systems favor facilitation versus

competition and how facilitation promotes coexistence and potentially the coevolution between species. In short, we propose that metabolically-informed consumer-resource dynamics will provide a platform to explore the consequences of cooperative and competitive interactions across environmental contexts.

Acknowledgements We thank Jake McKinlay, Thomas Koffel, and two anonymous reviewers for critical feedback on an earlier version of this manuscript. This work was supported by the Simons Foundation Grant # 376199 and the James S. McDonnell Foundation Grant # 220020439.

Conflict of interest

The authors declare that they have no conflict of interest.

References

- Abrams PA (1982) Functional Responses of Optimal Foragers. *The American Naturalist* 120(3):382–390, DOI 10.1086/283996
- Abrams PA (1983) Arguments in Favor of Higher Order Interactions. *The American Naturalist* 121(6):887–891, DOI 10.1086/284111
- Abrams PA (2009) Determining the Functional Form of Density Dependence: Deductive Approaches for Consumer-Resource Systems Having a Single Resource. *The American Naturalist* 174(3):321–330, DOI 10.1086/603627
- Allesina S, Tang S (2012) Stability criteria for complex ecosystems. *Nature* 483(7388):205–8, DOI 10.1038/nature10832
- Barabás G, Michalska-Smith MJ, Allesina S (2017) Self-regulation and the stability of large ecological networks. *Nature Ecology & Evolution* 1(12):1870–1875, DOI 10.1038/s41559-017-0357-6
- Barner AK, Coblenz KE, Hacker SD, Menge BA (2018) Fundamental contradictions among observational and experimental estimates of non-trophic species interactions. *Ecology* 99(3):557–566, DOI 10.1002/ecs.2133
- Brooker RW, Callaghan TV (1998) The balance between positive and negative plant interactions and its relationship to environmental gradients: a model. *Oikos* 81(1):196, DOI 10.2307/3546481
- Butler S, O’Dwyer JP (2018) Stability criteria for complex microbial communities. *Nature Communications* 9(1):2970, DOI 10.1038/s41467-018-05308-z
- Butler S, O’Dwyer JP (2019) Cooperation and Stability for Complex Systems in Resource Limited Environments. *Theoretical Ecology* In Press, DOI xxx

Callaway RM, Walker LR (1997) Competition and facilitation: A synthetic approach to interactions in plant communities. *Ecology* 78(7):1958–1965, DOI 10.1890/0012-9658(1997)078[1958:CAFASA]2.0.CO;2

Cardinale BJ, Palmer MA, Collins SL (2002) Species diversity enhances ecosystem functioning through interspecific facilitation. *Nature* 415(6870):426–9, DOI 10.1038/415426a

Cardinale BJ, Duffy JE, Gonzalez A, Hooper DU, Perrings C, Venail P, Narwani A, MacE GM, Tilman D, Wardle DA, Kinzig AP, Daily GC, Loreau M, Grace JB, Larigauderie A, Srivastava DS, Naeem S (2012) Biodiversity loss and its impact on humanity. *Nature* 486(7401):59–67, DOI 10.1038/nature11148, 9605103

Carini P, Campbell EO, Morré J and Sañudo-Wilhelmy SA, Thrash JC, Bennett SE, Temperton B, Begley T, Giovannoni SJ (2014) Discovery of a SAR11 growth requirement for thiamin’s pyrimidine precursor and its distribution in the Sargasso Sea. *ISME J* 8(8):1727–1738, DOI 10.1038/ismej.2014.61

Carrara F, Giometto A, Seymour M, Rinaldo A, Altermatt F (2015) Inferring species interactions in ecological communities: A comparison of methods at different levels of complexity. *Methods in Ecology and Evolution* 6(8):895–906, DOI 10.1111/2041-210X.12363

Chakraborty S, Nielsen LT, Andersen KH (2017) Trophic Strategies of Unicellular Plankton. *The American Naturalist* 189(4):E77–E90, DOI 10.1086/690764

Cherif M, Loreau M (2007) Stoichiometric constraints on resource use, competitive interactions, and elemental cycling in microbial decomposers. *The American Naturalist* 169(6):709–724, DOI 10.1086/516844

Cherif M, Loreau M (2010) Towards a more biologically realistic use of Droop’s equations to model growth under multiple nutrient limitation. *Oikos* 119(6):897–907, DOI 10.1111/j.1600-0706.2010.18397.x

De Ruiter PC, Neutel AM, Moore JC (1995) Energetics, patterns of interaction strengths, and stability in real ecosystems. *Science* (80-) 269(5228):1257–1260, DOI 10.1126/science.269.5228.1257

DeAngelis DL, Goldstein RA, O’Neill RV (1975) A Model for Tropic Interaction. *Ecology* 56(4):881–892, DOI 10.2307/1936298

DeAngelis DL, Mulholland PJ, Palumbo AV, Steinman AD, Huston MA, Elwood JW (1989) Nutrient Dynamics and Food-Web Stability. *Annual Review of Ecology and Systematics* 20:71–95, DOI 10.1146/annurev.es.20.110189.000443

Droop MR (1974) The nutrient status of algal cells in continuous culture. *Journal of the Marine Biological Association of the United Kingdom* 54(04):825, DOI 10.1017/S002531540005760X

- D'Souza G, Shitut S, Preussger D, Yousif G, Waschina S, Kost C (2018) Ecology and evolution of metabolic cross-feeding interactions in bacteria. *Nat Prod Rep* 35(5):455–488, DOI 10.1039/c8np00009c
- Duffy JE, Cardinale BJ, France KE, McIntyre PB, Thébault E, Loreau M (2007) The functional role of biodiversity in ecosystems: incorporating trophic complexity. *Ecology Letters* 10(6):522–38, DOI 10.1111/j.1461-0248.2007.01037.x
- Duncan SH, Louis P, Flint HJ (2004) Lactate-Utilizing Bacteria, Isolated from Human Feces, That Produce Butyrate as a Major Fermentation Product. *Applied and Environmental Microbiology* 70(10):5810–5817, DOI 10.1128/AEM.70.10.5810-5817.2004
- Embree M, Liu JK, Al-Bassam MM, Zengler K (2015) Networks of energetic and metabolic interactions define dynamics in microbial communities. *Proceedings of the National Academy of Sciences* 112(50):15450–15455, DOI 10.1073/pnas.1506034112, 1011.1669v3
- Falkowski PG, Jelen B, Giovannelli D (2016) The Role of Microbial Electron Transfer in the Coevolution of the Geosphere and Biosphere. *Annu Rev Microbiol* 70(1), DOI doi:10.1146/annurev-micro-102215-095521, URL <http://www.annualreviews.org/doi/abs/10.1146/annurev-micro-102215-095521>
- Flynn DFB, Mirotchnick N, Jain M, Palmer MI, Naeem S (2011) Functional and phylogenetic diversity as predictors of biodiversity–ecosystem-function relationships. *Ecology* 92(8):1573–1581, DOI 10.1890/10-1245.1, arXiv:1011.1669v3
- Freilich S, Zarecki R, Eilam O, Segal ES, Henry CS, Kupiec M, Gophna U, Sharan R, Ruppin E (2011) Competitive and cooperative metabolic interactions in bacterial communities. *Nature Communications* 2:589, DOI 10.1038/ncomms1597
- Gause GF, Witt AA (1935) Behavior of Mixed Populations and the Problem of Natural Selection. *The American Naturalist* 69(725):596–609, DOI 10.1086/280628
- Glass JB, Orphan VJ (2012) Trace Metal Requirements for Microbial Enzymes Involved in the Production and Consumption of Methane and Nitrous Oxide. *Front Microbiol* 3(FEB):1–20, DOI 10.3389/fmicb.2012.00061, URL <http://journal.frontiersin.org/article/10.3389/fmicb.2012.00061/abstract>
- Gottschalk G (1986) *Bacterial Metabolism*, 2nd edn. Springer-Verlag, New York, NY
- Grilli J, Barabás G, Michalska-Smith MJ, Allesina S (2017) Higher-order interactions stabilize dynamics in competitive network models. *Nature* 548(DOI: 10.1038/nature23273):210–213, DOI 10.1038/nature23273

- Grover JP (1990) Resource Competition in a Variable Environment: Phytoplankton Growing According to Monod's Model. *The American Naturalist* 136(6):771–789, DOI 10.1086/285131
- Grover JP (2011) Resource storage and competition with spatial and temporal variation in resource availability. *The American naturalist* 178(5):E124–48, DOI 10.1086/662163
- Hall SR (2009) Stoichiometrically Explicit Food Webs: Feedbacks between Resource Supply, Elemental Constraints, and Species Diversity. *Annual Review of Ecology, Evolution, and Systematics* 40(1):503–528, DOI 10.1146/annurev.ecolsys.39.110707.173518
- Harpole WS, Ngai JT, Cleland EE, Seabloom EW, Borer ET, Bracken ME, Elser JJ, Gruner DS, Hillebrand H, Shurin JB, Smith JE (2011) Nutrient co-limitation of primary producer communities. *Ecology Letters* 14(9):852–862, DOI 10.1111/j.1461-0248.2011.01651.x
- He Q, Bertness MD, Altieri AH (2013) Global shifts towards positive species interactions with increasing environmental stress. *Ecology Letters* 16(5):695–706, DOI 10.1111/ele.12080, 2666
- Holling CS (1959) The Components of Predation as Revealed by a Study of Small-Mammal Predation of the European Pine Sawfly. *The Canadian Entomologist* 91(05):293–320, DOI 10.4039/Ent91293-5
- Holt RD (1977) Predation, Apparent Competition, and the Structure of Prey Communities. *Theoretical Population Biology* 12:197–229, DOI 10.1016/0040-5809(77)90042-9
- Ives A, Dennis B, Cottingham K, Carpenter S (2003) Estimating community stability and ecological interactions from time-series data. *Ecological Monographs* 73(2):301–330, DOI 10.1890/0012-9615(2003)073[0301:ECSAEI]2.0.CO;2
- Jiao Y, Navid A, Stewart BJ, McKinlay JB, Thelen MP, Pett-Ridge J (2012) Syntrophic metabolism of a co-culture containing *Clostridium cellulolyticum* and *Rhodospseudomonas palustris* for hydrogen production. *International Journal of Hydrogen Energy* 37(16):11719–11726, DOI 10.1016/j.ijhydene.2012.05.100
- Kauffman KJ, Prakash P, Edwards JS (2003) Advances in flux balance analysis. *Current Opinion in Biotechnology* 14(5):491–496, DOI 10.1016/j.copbio.2003.08.001, /www.nature.com/nature/journal/v473/n7346/abs/10.1038-nature10011-unlocked.html#supplementary-information
- Keller KR, Lau JA (2018) When mutualisms matter: Rhizobia effects on plant communities depend on host plant population and soil nitrogen availability. *Journal of Ecology* 106(3):1046–1056, DOI 10.1111/1365-2745.12938
- Kempes CP, Dutkiewicz S, Follows MJ (2012) Growth, metabolic partitioning, and the size of microorganisms. *Proceedings of the National Academy of Sciences* 109(2):495–500, DOI 10.1073/pnas.1115585109

- Koffel T, Boudsocq S, Loeuille N, Daufresne T (2018) Facilitation- vs. competition-driven succession: the key role of resource-ratio. *Ecology Letters* 21(7):1010–1021, DOI 10.1111/ele.12966
- Kooijman SA (1998) The Synthesizing Unit as model for the stoichiometric fusion and branching of metabolic fluxes. *Biophys Chem* 73(1-2):179–188, DOI 10.1016/S0301-4622(98)00162-8
- Kooijman SA (2001) Quantitative aspects of metabolic organization: A discussion of concepts. *Philosophical Transactions of the Royal Society B: Biological Sciences* 356(1407):331–349, DOI 10.1098/rstb.2000.0771
- Leibold M, McPeck M (2006) Coexistence of the niche and neutral perspectives in community ecology. *Ecology* 87(6):1399–1410, DOI 10.1890/0012-9658(2006)87[1399:COTNAN]2.0.CO;2
- von Liebig JF, Gregory W (1842) *Animal chemistry: or, Organic chemistry in its application to physiology and pathology*. John Owen
- Litchman E (2003) Competition and coexistence of phytoplankton under fluctuating light: experiments with two cyanobacteria. *Aquatic Microbial Ecology* 31:241–248, DOI 10.3354/ame031241
- Litchman E, Edwards KF, Klausmeier CA (2015) Microbial resource utilization traits and trade-offs: implications for community structure, functioning, and biogeochemical impacts at present and in the future. *Frontiers in Microbiology* 06(April):1–10, DOI 10.3389/fmicb.2015.00254
- Loreau M (2001) Microbial diversity, producer-decomposer interactions and ecosystem processes: a theoretical model. *Proceedings of the Royal Society* 268(1464):303–9, DOI 10.1098/rspb.2000.1366
- Loreau M (2010) *From Populations to Ecosystems: Theoretical Foundations for a New Ecological Synthesis*. Princeton University Press, Princeton, NJ
- Lotka AJ (1932) The growth of mixed populations: Two species competing for a common food supply. *Journal of the Washington Academy of Sciences* 22(16/17):461–469, DOI 10.1007/978-3-642-50151-7_12
- MacArthur R (1970) Species packing and competitive equilibrium for many species. *Theoretical Population Biology* 1(1):1–11, DOI 10.1016/0040-5809(70)90039-0
- Marino S, Baxter NT, Huffnagle GB, Petrosino JF, Schloss PD (2014) Mathematical modeling of primary succession of murine intestinal microbiota. *Proceedings of the National Academy of Sciences* 111(1):439–444, DOI 10.1073/pnas.1311322111, arXiv:1011.1669v3
- Mee MT, Collins JJ, Church GM, Wang HH (2014) Syntrophic exchange in synthetic microbial communities. *Proceedings of the National Academy of Sciences of the United States of America* 111(20):E2149–56, DOI 10.1073/pnas.1405641111

- Moore JC, Ruiter PCD, Hunt HW (1993) Influence of Productivity on the Stability of Real and Model Ecosystems. Science (80-) 261(5123):906–908, URL <http://www.jstor.org/stable/2882122>
- Moore JC, McCann K, De Ruiter PC (2005) Modeling trophic pathways, nutrient cycling, and dynamic stability in soils. Pedobiologia (Jena) 49(6):499–510, DOI 10.1016/j.pedobi.2005.05.008
- Morris JJ (2015) Black Queen evolution: The role of leakiness in structuring microbial communities. Trends in Genetics 31(8):475–482, DOI 10.1016/j.tig.2015.05.004
- Morris JJ, Lenski RE, Zinser ER (2012) The black queen hypothesis: evolution of dependencies through adaptive gene loss. MBio 3(2):e00036–12, DOI 10.1128/mBio.00036-12
- Mougi A, Kondoh M (2012) Diversity of Interaction Types and Ecological Community Stability. Science 337(6092):349–351, DOI 10.1126/science.1220529
- Murdoch WW, Briggs CJ, Nisbet RM (2003) Consumer-resource dynamics, vol 36. Princeton University Press
- Nisbet RM, Muller EB, Lika K, Kooijman SaLM (2008) From molecules to ecosystems through dynamic energy budget models. Journal of Animal Ecology 69(6):913–926, DOI 10.1111/j.1365-2656.2000.00448.x
- Odum EP (1959) Fundamentals of ecology. WB Saunders company
- O’Dwyer JP (2018) Whence Lotka-Volterra? Theoretical Ecology pp 1–12, DOI 10.1007/s12080-018-0377-0
- Orth JD, Thiele I, Palsson BO (2010) What is flux balance analysis? Nature Biotechnology 28(3):245–248, DOI 10.1038/nbt.1614, NIHMS150003
- Pacheco AR, Moel M, Segrè D (2019) Costless metabolic secretions as drivers of interspecies interactions in microbial ecosystems. Nature Communications 10(1):103, DOI 10.1038/s41467-018-07946-9
- Pande S, Merker H, Bohl K, Reichelt M, Schuster S, de Figueiredo LF, Kaleta C, Kost C (2014) Fitness and stability of obligate cross-feeding interactions that emerge upon gene loss in bacteria. ISME J 8:953–62, DOI 10.1038/ismej.2013.211, URL <http://www.ncbi.nlm.nih.gov/pubmed/24285359>
- Pfeiffer T, Bonhoeffer S (2004) Evolution of Cross-Feeding in Microbial Populations. The American Naturalist 163(6):E126–E135, DOI 10.1086/383593
- Price PB, Sowers T (2004) Temperature dependence of metabolic rates for microbial growth, maintenance, and survival. Proceedings of the National Academy of Sciences 101(13):4631–4636, DOI 10.1073/pnas.0400522101
- Ramírez-Flandes S, González B, Ulloa O (2019) Redox traits characterize the organization of global microbial communities. Proc Natl Acad Sci 116(9):3630–3635, DOI 10.1073/pnas.1817554116

Russell JB, Cook GM (1995) Energetics of Bacterial Growth: Balance of Anabolic and Catabolic Reactions. *Microbiology and Molecular Biology Reviews* 59(1):48–62

Schimel JP, Weintraub MN (2003) The implications of exoenzyme activity on microbial carbon and nitrogen limitation in soil: A theoretical model. *Soil Biol Biochem* 35(4):549–563, DOI 10.1016/S0038-0717(03)00015-4

Schoener TW (1983) Field Experiments on Interspecific Competition. *The American Naturalist* 122(2):240–285, DOI 10.1086/284133

Sterner RW, Elser JJ (2002) *Ecological stoichiometry: the biology of elements from molecules to the biosphere*. Princeton University Press, Princeton, NJ

Sun Z, Koffel T, Stump SM, Grimaud GM, Klausmeier CA (2019) Microbial cross-feeding promotes multiple stable states and species coexistence, but also susceptibility to cheaters. *Journal of Theoretical Biology* 465:63–77, DOI 10.1016/j.jtbi.2019.01.009

Tasoff J, Mee MT, Wang HH (2015) An economic framework of microbial trade. *PLoS ONE* 10(7):1–20, DOI 10.1371/journal.pone.0132907

Terry JCD, Morris RJ, Bonsall MB (2017) Trophic interaction modifications: an empirical and theoretical framework. *Ecology Letters* 20(10):1219–1230, DOI 10.1111/ele.12824

Tilman D (1977) Resource competition between plankton algae: an experimental and theoretical approach. *Ecology* 58(2):338–348

Tilman D (1980) Resources: a graphical-mechanistic approach to competition and predation. *The American Naturalist* 116(3):362–393

Tilman D (1987) The Importance of the Mechanisms of Interspecific Competition. *The American Naturalist* 129(5):769–774, DOI 10.1086/284672

Tilman D, Kilham SS, Kilham P (1982) Phytoplankton community ecology: the role of limiting nutrients. *Annual Review of Ecology and Systematics* 13(1):349–372, DOI 10.1146/annurev.es.13.110182.002025

Vellend M (2010) Conceptual synthesis in community ecology. *The Quarterly review of biology* 85(2):183–206, DOI 10.1086/652373

Vellend M (2016) *The theory of ecological communities*. Princeton University Press

Volterra V (1926) Fluctuations in the Abundance of a Species considered Mathematically. *Nature* 118(2972):558–560, DOI 10.1038/118558a0

Wiegert RG, Owen DF (1971) Trophic structure, available resources and population density in terrestrial vs. aquatic ecosystems. *J Theor Biol* 30(1):69–81, DOI 10.1016/0022-5193(71)90037-3

Xiao Y, Angulo MT, Friedman J, Waldor MK, Weiss ST, Liu YY (2017) Mapping the ecological networks of microbial communities. *Nature Communications* 8(1):2042, DOI 10.1038/s41467-017-02090-2

Zelezniak A, Andrejev S, Ponomarova O, Mende DR, Bork P, Patil KR (2015) Metabolic dependencies drive species co-occurrence in diverse microbial communities. *Proceedings of the National Academy of Sciences* 112(20):6449–6454, DOI 10.1073/pnas.1421834112, 1408.1149

Zomorodi AR, Segrè D (2016) Synthetic Ecology of Microbes: Mathematical Models and Applications. *Journal of Molecular Biology* 428(5):837–861, DOI 10.1016/j.jmb.2015.10.019