1 Inhibitor of Differentiation 4 (ID4) represses myoepithelial differentiation of
2 mammary stem cells through its interaction with HEB

3 Holly Holliday^{1,2}, Daniel Roden^{1,2}, Simon Junankar^{1,2}, Sunny Z. Wu^{1,2}, Laura A.
 4 Baker^{1,2}, Christoph Krisp^{3,4}, Chia-Ling Chan¹, Andrea McFarland¹, Joanna N. Skhinas¹,
 5 Thomas R. Cox^{1,2}, Bhupinder Pal⁵, Nicholas Huntington⁶, Christopher J. Ormandy^{1,2},
 6 Jason S. Carroll⁷, Jane Visvader⁵, Mark P. Molloy³, Alexander Swarbrick^{1,2}

- 7 1. The Kinghorn Cancer Centre, Garvan Institute of Medical Research, Sydney,
8 NSW 2010, Australia
- 9 2. St Vincent's Clinical School, Faculty of Medicine, UNSW Sydney, Sydney, NSW
10 2010, Australia
- 11 3. Australian Proteome Analysis Facility, Macquarie University, Sydney, NSW
12 2109, Australia
- 13 4. Institute of Clinical Chemistry and Laboratory Medicine, Mass Spectrometric
14 Proteomics, University Medical Center Hamburg-Eppendorf, Hamburg 20251,
15 Germany
- 16 5. ACRF Stem Cells and Cancer Division, The Walter and Eliza Hall Institute of
17 Medical Research, Parkville, Victoria 3052, Australia
- 18 6. Biomedicine Discovery Institute, Department of Biochemistry and Molecular
19 Biology, Monash University, Clayton, VIC 3168, Australia
- 20 7. Cancer Research UK Cambridge Institute, University of Cambridge, Robinson
21 Way, Cambridge CB2 0RE, UK

23 Keywords

24 Differentiation / Epithelium / Mammary gland / Myoepithelial / Transcription factor

25 Abstract

26 Differentiation of stem cells embedded within the mammary epithelium is
 27 orchestrated by lineage-specifying transcription factors. Unlike the well-defined
 28 luminal hierarchy, dissection of the basal lineage has been hindered by a lack of
 29 specific markers. Inhibitor of Differentiation 4 (ID4) is a basally-restricted helix-loop-
 30 helix (HLH) transcription factor essential for mammary development. Here we show
 31 that ID4 is highly expressed in basal stem cells and decreases during myoepithelial
 32 differentiation. By integrating transcriptomic, proteomic, and ChIP-sequencing data,
 33 we reveal that ID4 is required to suppress myoepithelial gene expression and cell
 34 fate. We identify the bHLH protein HEB as a direct binding partner of ID4, and
 35 describe a previously-unknown role for this regulator in mammary development.
 36 HEB binds to E-boxes in regulatory elements of developmental genes, negatively
 37 regulated by ID4, involved in extracellular matrix synthesis and cytoskeletal
 38 contraction. Together our findings support a model whereby ID4 binds to HEB and
 39 blocks it from promoting myoepithelial specialisation. These new insights expand our
 40 current understanding into control of myoepithelial differentiation and mammary
 41 gland morphogenesis.

42 Introduction

43 The mammary gland undergoes marked tissue remodelling throughout life. During
 44 murine pubertal development, terminal end buds (TEBS) located at the tips of the

ducts invade into the surrounding stromal fat pad (Macias & Hinck, 2012, Williams & Daniel, 1983). This process is driven by collective migration and rapid proliferation of outer cap cells, which surround multiple layers of inner body cells (Williams & Daniel, 1983). As the ducts elongate, the cap cells differentiate into the basal cell layer and the body cells adjacent to the basal cells give rise to the luminal cell layer (Williams & Daniel, 1983), while the innermost body cells undergo apoptosis to sculpt the bilayered ductal tree present in the adult gland (Paine & Lewis, 2017). During pregnancy the gland undergoes alveolar morphogenesis in preparation for production and secretion of milk at lactation. The luminal cells differentiate into milk-producing alveolar cells and the milk is ejected from the gland by the contractile action of specialised myoepithelial cells, smooth muscle-like epithelial cells which differentiate from basal cells (Macias & Hinck, 2012). From hereon ‘basal’ refers to the basal lineage, encompassing cap cells, duct basal cells, and myoepithelial cells. Initial transplantation experiments demonstrated that the basal lineage contains bipotent mammary stem cells (MaSCs), capable of forming both myoepithelial and luminal populations, while luminal cells are restricted in their differentiation potential (Shackleton, Vaillant et al., 2006, Stingl, Eirew et al., 2006). Subsequent lineage tracing experiments support the existence of a small population of basally-restricted bipotent stem cells (Rios, Fu et al., 2014, Wang, Cai et al., 2015), with a majority of unipotent basal and luminal mammary stem cells (Davis, Lloyd-Lewis et al., 2016, Lilja, Rodilla et al., 2018, Lloyd-Lewis, Davis et al., 2018, Prater, Petit et al., 2014, Scheele, Hannezo et al., 2017, van Amerongen, Bowman et al., 2012, Van Keymeulen, Rocha et al., 2011, Wuidart, Ousset et al., 2016, Wuidart, Sifrim et al., 2018) that replenish the myoepithelial and luminal lineages throughout life.

69 Lineage specifying transcription factors are responsible for directing luminal and
70 myoepithelial differentiation, and also for maintaining the self-renewal of
71 uncommitted stem cells upstream in the mammary epithelial hierarchy.
72 Transcriptomic profiling of sorted epithelial subpopulations has been instrumental in
73 identifying such lineage specifying transcription factors that regulate each step of
74 luminal-alveolar differentiation (Asselin-Labat, Sutherland et al., 2007, Bouras, Pal et
75 al., 2008, Buchwalter, Hickey et al., 2013, Carr, Kiefer et al., 2012, Chakrabarti, Wei
76 et al., 2012, Kouros-Mehr, Slorach et al., 2006, Liu, Ginestier et al., 2008, Oakes,
77 Naylor et al., 2008, Yamaji, Na et al., 2009). However, due to lack of specific cell
78 markers that can resolve stem and myoepithelial populations, it has been
79 challenging to dissect molecular regulators of basal differentiation. While a number
80 of basal-specific transcription factors have been identified, such as P63, SLUG, SOX9,
81 SRF and MRTFA (Guo, Keckesova et al., 2012, Li, Chang et al., 2006, Mills, Zheng et
82 al., 1999, Sun, Boyd et al., 2006, Yang, Schweitzer et al., 1999), their role in the basal
83 compartment and myoepithelial specialisation is poorly understood. ID proteins
84 (ID1-4) are helix-loop-helix (HLH) transcriptional regulators that lack a DNA binding
85 domain. They function by dimerizing with basic HLH (bHLH) transcription factors and
86 preventing them from binding to E-box DNA motifs and regulating transcription
87 (Benezra, Davis et al., 1990). E-box motifs are found in regulatory regions of genes
88 involved in lineage specification and as such ID proteins and bHLH transcription
89 factors are critical regulators of stemness and differentiation across diverse cellular
90 lineages (Massari & Murre, 2000). The expression of ID4 in mouse and human
91 mammary epithelium is exclusive to the basal population (Lim, Wu et al., 2010). We
92 and others have demonstrated that ID4 is a key regulator of mammary stem cells,

93 required for ductal elongation during puberty (Best, Hutt et al., 2014, Dong, Huang
94 et al., 2011, Junankar, Baker et al., 2015). ID4 was also shown to have a role in
95 blocking luminal differentiation (Best et al., 2014, Junankar et al., 2015). The precise
96 molecular mechanism by which ID4 functions in the mammary gland, including its
97 full repertoire of transcriptional targets and interacting partners, have yet to be
98 determined. Here we show that ID4 marks immature basal cells and demonstrate
99 that ID4 inhibits myoepithelial differentiation. Moreover, using unbiased interaction
100 proteomics, we identify a novel player in the mammary differentiation hierarchy –
101 the bHLH transcription factor HEB. By mapping the genome-wide binding sites of
102 HEB we show that it directly binds to regulatory elements of ID4 target genes
103 involved in myoepithelial functions such as contraction and extracellular matrix
104 (ECM) synthesis.

105 Results

106 Loss of ID4 causes upregulation of myoepithelial genes in basal cells

107 In order to determine the genes regulated by ID4 in mammary epithelial cells, we
108 FACS-enriched basal, luminal progenitor and mature luminal cells from wild type
109 (WT) and ID4 knockout (KO) mice (Yun, Mantani et al., 2004) and performed RNA-
110 sequencing (RNA-seq) (Fig. 1a). No significant differences were observed in the
111 proportion of mammary epithelial subpopulations between WT and KO mice (Fig.
112 S1a). 96 genes were significantly ($FDR < 0.05$) upregulated and 141 genes
113 downregulated in ID4 KO basal cells. In contrast, only 2 genes were differentially
114 expressed in the two luminal subpopulations (Fig. 1b-c and Table S1) suggesting that
115 ID4 predominantly regulates gene expression within basal cells *in vivo*.

116 In line with the known role of ID4 in promoting proliferation of mammary epithelial
 117 cells (Dong et al., 2011, Junankar et al., 2015), the genes downregulated in ID4 KO
 118 basal cells were enriched for pathways involved in cell growth such as translation
 119 and metabolism (Fig. 1d-e and Fig. S1b). Conversely, the genes upregulated in KO
 120 basal cells were enriched for pathways related to basal cells and smooth muscle
 121 function. These included a muscle-derived leiomyosarcoma signature (Nielsen, West
 122 et al., 2002), smooth muscle contraction genes, epithelial-mesenchymal transition
 123 (EMT) and Serum Response Factor (SRF) targets (Fig. 1d and Fig. S1c), which shared
 124 common genes as indicated by the connecting edges in the enrichment map network
 125 (Fig. 1e). SRF target genes are of relevance as SRF is a master regulator of
 126 cytoskeletal contraction, and is one of the only transcription factors implicated in
 127 myoepithelial differentiation (Li et al., 2006, Miano, Long et al., 2007, Sun et al.,
 128 2006). Taken together, loss of ID4 causes basal cells to decrease proliferation and
 129 adopt a more differentiated myoepithelial and mesenchymal gene expression
 130 program, implicating ID4 in the repression of basal cell specialisation.

131

132 **ID4 marks basal stem cells and is dynamically regulated during myoepithelial** 133 **differentiation**

134 We previously showed that ID4 is heterogeneously expressed in basal cells and that
 135 ID4-positive cells have enhanced mammary reconstitution activity (Junankar et al.,
 136 2015). To further characterise the phenotype of ID4-positive basal cells, we used an
 137 ID4-GFP reporter mouse in which the ID4 promoter drives GFP expression (Best et
 138 al., 2014). Basal cells with high expression of the stem cell marker CD49f (also known
 139 as ITGA6) and the epithelial marker EPCAM have been shown to be enriched for

140 MaSC activity (Prater et al., 2014, Stingl et al., 2006). ID4-GFP expression was
 141 maximal in this EPCAM^{hi} CD49f^{hi} subset (Fig. 2a). ID4-GFP expression within the basal
 142 gate was binned into 3 groups: bright (top 10%), intermediate (middle 80%) and dim
 143 (bottom 10%) and the median fluorescent intensity (MFI) of EPCAM and CD49f was
 144 analysed within these groups (Fig. 2b-c). ID4-bright cells had significantly higher
 145 EPCAM and CD49f MFI than ID4-dim and intermediate cells, suggesting ID4 marks
 146 basal stem cells.

147 We next sought to locate ID4-high and ID4-low populations in a tissue context and to
 148 further investigate the association between ID4 and markers of myoepithelial
 149 differentiation. During ductal elongation, the cap cells differentiate at the neck of
 150 the TEBs and mature into myoepithelial cells which form the outer basal layer of the
 151 ducts (Paine & Lewis, 2017). Mammary gland sections from pubertal mice, in which
 152 TEBs and ducts are both present, were stained for ID4 and the myoepithelial marker
 153 alpha-smooth muscle actin (α -SMA). ID4 expression was highest in the nuclei of cap
 154 cells at the extremity of the TEBs and expressed at lower levels in basal cells of ducts
 155 (Fig. 2d and Fig. S2). Conversely, α -SMA expression was higher in ductal basal cells
 156 and lower in cap cells. ID4-high cap cells had a compact cuboidal epithelial
 157 appearance compared to the more separated elongated morphology of the ID4-low
 158 duct cells (Fig. 2d and Fig. S2). Quantification of fluorescence demonstrated that ID4
 159 was more highly expressed in cap cells of TEBs than in basal cells of ducts, while the
 160 opposite was true for α -SMA expression (Fig. 2e-f). A negative correlation between
 161 α -SMA and ID4 was observed, with clear separation between cap (dark blue Fig. 2g)
 162 and duct basal cells (light blue Fig. 2g). As a negative control, α -SMA fluorescence
 163 was compared with nuclear stain Hoechst and no correlation or separation based on

164 region was observed (Fig. 2h). Thus, based on marker expression, morphology and
165 spatial localization, ID4 expression is high in epithelial-like cap cells and is lower in
166 myoepithelial cells.

167 Terminal differentiation of basal cells into contractile myoepithelial cells occurs
168 during lactation. We interrogated a published single cell RNA-seq (scRNA-seq) data-
169 set (Bach, Pensa et al., 2017) to examine *Id4* expression dynamics over postnatal
170 murine mammary gland development. In this study, individual EPCAM+ mammary
171 epithelial cells from four developmental stages: nulliparous (8 week), gestation (day
172 14.5), lactation (day 6), and involution (day 11) were captured and profiled. We
173 limited our analysis to basal cell clusters (9,663 cells), defined by expression of both
174 *Krt5* and *Krt14*. Basal cells broadly clustered by developmental time point (Fig. 2i).
175 Increased differentiation of myoepithelial cells appears to proceed from nulliparous
176 to gestation to lactation, associated with decreased expression of epithelial marker
177 *Epcam* and increased expression of *Acta2* (encoding α -SMA) (Fig. 2j-k), consistent
178 with gradual acquisition of a smooth muscle phenotype and loss of adherent
179 epithelial features (Deugnier, Moiseyeva et al., 1995). Like *Epcam*, *Id4* expression
180 was highest in basal cells from nulliparous mice, and decreased in basal cells of
181 pregnant and lactating mice (Fig. 2j-k). This result was validated on the protein level
182 by immunohistochemical staining for ID4 on mammary gland sections at different
183 developmental time points (Fig. S2b). Using the Monocle 2 package (Trapnell,
184 Cacchiarelli et al., 2014), we performed pseudotemporal ordering of all basal cells to
185 form a myoepithelial differentiation trajectory (Fig. S2c). As expected, the
186 nulliparous and involution cells clustered together in pseudotime space in the least
187 differentiated part of the trajectory. Basal cells from gestating mice were dispersed

188 between the nulliparous and lactation stages, while basal cells at lactation were the
 189 most differentiated. *Id4* expression decreased, while several myoepithelial markers
 190 (*Acta2*, *Cnn1*, *Mylk*, *Myh11* and *Oxtr*) increased over pseudotime (Fig. S2d).
 191 We next sought to identify transcriptional signatures associated with high and low
 192 *Id4* expressing cells in the mammary basal epithelium across all developmental time
 193 points (Fig 2l and Table S2). Basal cells with high *Id4* expression were enriched for
 194 genes involved in RNA binding, metabolic processes, translation and ribosome
 195 biogenesis (Fig. 2m). Conversely, genes enriched in the *Id4*-low basal cells were
 196 involved in muscle contraction and response to cytokine and hormone stimuli, and
 197 circulatory system development (Fig. 2n). Notably, myoepithelial genes such as *Oxtr*,
 198 encoding the oxytocin receptor, and contractile genes *Mylk*, *Cnn1*, *Cav1* and *Myh11*,
 199 were among the top differentially expressed genes in the cells with low *Id4*
 200 expression (Fig. 2l and Table S2). Thus, ID4 downregulation during pregnancy and
 201 lactation is associated with terminal differentiation into functionally mature
 202 contractile myoepithelial cells.

203

204 **Loss of ID4 results in myoepithelial differentiation of mammary organoids**

205 To functionally validate the role of ID4 in suppressing the myoepithelial
 206 differentiation of basal cells we generated a primary basal cell organoid model (Fig.
 207 3a). The organoid model system complements and expands upon the findings from
 208 the KO mouse, as the acute consequence of ID4 loss on basal cell phenotype can be
 209 determined. Furthermore, organoids are less complex cellular systems than tissue,
 210 thus cell-autonomous effects can be isolated more precisely. ID4-positive basal cells
 211 from mice in which exons 1 and 2 of *Id4* are floxed (*Id4^{fl/fl}*) (Best et al., 2014) were

212 isolated using cell sorting. To overcome culture-induced senescence, basal cells were
 213 conditionally reprogrammed into a proliferative stem/progenitor state using an
 214 established protocol utilizing irradiated 3T3 fibroblast feeders and Rho Kinase
 215 (ROCK) inhibition (Liu, Ory et al., 2012, Prater et al., 2014) (Fig. 3a). Conditionally
 216 reprogrammed basal cells adopted an epithelial cobblestone morphology and
 217 expressed high levels of ID4 (Fig. S3a-b). Cells cultured in the absence of ROCK
 218 inhibitor or feeders adopted a flattened differentiated/senescent cell morphology
 219 and had reduced ID4 expression (Fig. S3a-b). Reprogrammed cells maintained basal
 220 marker expression of P63 and KRT14 (Fig. S3c) and could be grown as 3D organoids
 221 with basal marker KRT14 on the outer cell layer and luminal marker KRT8 on the
 222 inner cells of the organoids (Fig. S3d).

223 ID4 expression was limited to the outer basal cells (Fig. 3b), recapitulating expression
 224 of ID4 in cap cells of TEBs (Fig. 2d). To test whether ID4 regulates differentiation of
 225 basal cells we deleted ID4 with Adenoviral-Cre and compared these to cells treated
 226 with control Adenoviral-GFP. ID4 protein was markedly downregulated in organoids
 227 infected with Cre adenovirus confirming successful gene deletion (Fig. 3b-c). Loss of
 228 ID4 resulted in increased α -SMA fluorescent intensity (Fig. 3d-e). This finding
 229 demonstrates that as well as marking undifferentiated basal cells, ID4 has a cell
 230 autonomous role in impeding maturation of basal cells into myoepithelial cells.

231

232 **ID4 inhibits expression of contractile and ECM genes *in vitro***

233 Given the negative association between ID4 and the differentiated myoepithelial
 234 phenotype, we next sought to determine direct ID4 target genes by performing RNA-
 235 seq following siRNA-mediated ID4 depletion in the spontaneously immortalized

236 mouse mammary epithelial cell line Comma-D β (Danielson, Oborn et al., 1984). This
 237 normal-like cell line expresses basal markers and is commonly used as a model of
 238 mammary stem/progenitors cells as they retain the capacity of multi-lineage
 239 differentiation (Best et al., 2014, Danielson et al., 1984, Deugnier, Faraldo et al.,
 240 2006, Idoux-Gillet, Nassour et al., 2018, Junankar et al., 2015). They can also be
 241 readily expanded *in vitro* to produce a large amount of material for investigation of
 242 mechanisms by molecular biology and biochemistry. Western blotting confirmed 70-
 243 80% knockdown (KD) of ID4 protein 48 hr after siRNA transfection (Fig. 4a). RNA-seq
 244 analysis of ID4 KD cells resulted in 471 and 421 (FDR<0.05) down and upregulated
 245 genes, respectively, compared to the non-targeting siRNA control (Table S3).
 246 Downregulated genes were predominantly involved in cell proliferation and growth
 247 pathways (Fig. 4b-c; blue) including hallmark gene sets such as E2F targets, MYC
 248 targets, and G2M checkpoint (Fig. S4a-b). Driving enrichment were several key cell
 249 cycle genes such as *Mki67*, *Cdk2*, *Cdk6* and *Cdk17* (Table S3). This result is consistent
 250 with the loss of cell growth gene expression programs in ID4 KO basal cells *in vivo*
 251 (Fig. 1d-e).
 252 Genes acutely upregulated upon ID4 depletion were involved in development,
 253 morphogenesis, ECM remodelling, and immune signalling (Fig. 4b-c; red). Consistent
 254 with ID4 repressing myoepithelial specialisation, loss of ID4 resulted in upregulation
 255 of SRF targets, actomyosin cytoskeleton, EMT and myogenesis gene signatures (Fig.
 256 4d and Fig. S4a-b). Several contractile genes were increased in ID4 depleted cells
 257 including *Cnn1*, *Cnn2*, *Tagln*, *Lmod1* and *Acta2* (FDR=0.06) (Table S3), many of which
 258 were inversely correlated with *Id4* expression in the scRNA-seq analysis (Fig. 2l and
 259 Table S2).

Several ECM genes encoding collagens (e.g. *Col1a1*, *Col1a2*, and *Col5a1*), basement membrane laminins (e.g. *Lamc1* and *Lama4*), and matricellular proteins (e.g. *Sparc*) were also upregulated upon ID4 depletion (Fig. 4c-d and Table S3). The ECM provides physical support to the mammary gland and is a source of biochemical signals important for coordinating morphogenesis (Muschler & Streuli, 2010). Changes in ECM gene expression are associated with EMT and cellular contractility (Kierner, Takeuchi et al., 2001, Liu, Eischeid et al., 2012). To functionally validate the role of ID4 in regulating ECM proteins we examined whether ID4 represses ECM deposition *in vivo* using picrosirius red staining of mammary glands from ID4 WT and KO mice to visualise fibrillar collagen (Fig. 4e). ID4 KO TEBs in pubertal mammary glands were surrounded by a thickened collagen-dense ECM when compared to ID4 WT TEBs (Fig. 4e i-ii), indicating that ID4 normally restrains collagen expression by cap cells during puberty. In addition to total abundance, the thickness of bundled collagen fibres can be further distinguished using polarized light. Analysis of birefringence signal revealed an increase in thick fibres and a decrease in thin fibres in the ECM surrounding ID4 KO TEBs (Fig. 4e iii) signifying a redistribution of collagen composition, as well as an overall increase in collagen abundance, in the absence of ID4.

Hence, ID4 positively regulates proliferative genes and negatively regulates genes involved in myoepithelial functions such as contraction and ECM synthesis in mammary epithelial cells. These ID4-regulated functions are likely to be important for morphogenesis of the ductal tree during pubertal development.

282

283 **ID4 interacts with E-proteins in mammary epithelial cells**

284 As ID4 lacks a DNA binding domain, it inevitably influences transcription through its
 285 interaction with other DNA-binding proteins. We used Rapid Immunoprecipitation
 286 Mass spectrometry of Endogenous proteins (RIME) to discover the binding partners
 287 of ID4 in Comma-D β cells (Mohammed, D'Santos et al., 2013). We identified 48
 288 proteins that were significantly ($p < 0.05$) more abundant in the ID4 IPs compared to
 289 the IgG negative control IPs in three independent RIME experiments (Fig. 5a and
 290 Table S4). ID4 was consistently identified in all replicates and was the top hit (Table
 291 S4), verifying the validity of the technique. Among the putative ID4 binding partners
 292 were many DNA and RNA interacting proteins (Fig. 5a).

293 The E-proteins E2A and HEB were identified as ID4 interactors in our RIME analysis.
 294 There are three members of the E-protein family (E2A, HEB, and ITF-2), that dimerise
 295 with other E-proteins or tissue-specific bHLH transcription factors (e.g. MyoD and
 296 NeuroD) to regulate expression of lineage commitment genes (Wang & Baker, 2015).
 297 E2A has been implicated in branching morphogenesis of mammary organoids (Lee,
 298 Gjorevski et al., 2011), however, there are no previous studies implicating HEB in
 299 mammary gland development. Given this, and that E-proteins are the canonical
 300 binding partners of ID proteins in other lineages, we chose to pursue the ID4-HEB
 301 interaction further.

302 Reciprocal co-immunoprecipitation followed by western blotting (co-IP WB)
 303 experiments were performed to validate the ID4-E-protein interactions in the same
 304 Comma-D β cell line, as well as in normal human mammary epithelial cell lines
 305 PMC42 and MCF10A. IP of ID4 resulted in co-IP of E2A and HEB and correspondingly,
 306 IP of E2A and HEB resulted in co-IP of ID4 (Fig. 5b and S5a-b). E2A and HEB did not
 307 form heterodimers with each other in Comma-D β cells (Fig. 5b), implying that E-

308 proteins either function as homodimers, or heterodimers with other bHLH proteins
 309 in this context. To identify mammary specific bHLH factors binding to HEB, we
 310 performed another RIME experiment by immunoprecipitating HEB protein. HEB
 311 (HTF4_MOUSE in Table S4) was identified as the top hit, and ID4 was also identified
 312 (Fig. S5c). However, we did not identify any other bHLH transcription factors in this
 313 experiment (Fig. S5c and Table S4). This suggests that HEB either binds DNA as a
 314 homodimer or binds another factor that was not detected by this assay. To
 315 independently validate the interaction between ID4 and HEB, the Proximity Ligation
 316 Assay (PLA) was used to visualize protein-protein interactions *in situ*. Multiple PLA
 317 foci were detected in the nuclei of Comma-D β cells co-stained with ID4 and HEB
 318 antibodies (Fig. 5c-d).

319 To test if ID4 and HEB interacted *in vivo*, we engineered a tagged ID4 mouse model
 320 in which a FlagV5 tag, a very efficient and specific target for immunoprecipitation,
 321 was integrated into the *Id4* locus downstream of the open reading frame. The result
 322 was FlagV5-tagged ID4 protein under the control of endogenous regulatory
 323 elements. We first ensured that ID4 and HEB were co-expressed by performing co-
 324 immunofluorescent staining for V5 and HEB. ID4-FlagV5 expression was tightly
 325 restricted to the cap and ductal basal cells, while HEB was more ubiquitous in its
 326 expression; detected in basal and luminal epithelial cells and surrounding stromal
 327 cells (Fig. 5e). ID4 and HEB co-localized in the nuclei of cap and duct basal cells
 328 (insets; Fig. 5e). Reciprocal co-IP was carried out from digested mammary glands of
 329 ID4-FlagV5 mice using antibodies raised against ID4, Flag, V5 (two different
 330 antibodies) and HEB (Fig. 5f). Each antibody tested was able to precipitate both ID4
 331 and HEB, confirming interaction between the two transcription factors *in vivo*.

332

333 **HEB binds to regulatory elements of a subset of ID4 differentially expressed genes**

334 To establish if ID4 regulates gene expression through HEB, we sought to determine if
 335 HEB directly binds to genes regulated by ID4 by performing ChIP-seq for HEB in
 336 Comma-D β cells. Across four biological replicates there were a total of 2752 HEB
 337 peaks identified (FDR<0.05). This was narrowed down to 956 consensus peaks which
 338 were present in at least 2 replicates. Transcription factor motif enrichment was
 339 carried out using MEME-ChIP (Machanick & Bailey, 2011) and the top enriched
 340 motifs were canonical E-box motifs (CANNTG), which are the binding sites for E-
 341 proteins (Fig. 6a). The majority of peaks were mapped to intergenic and intronic
 342 regions, and approximately 5% of the peaks occurred at gene promoters (Fig. S6a).
 343 We used the Genomics Regions Enrichment of Annotations Tool (GREAT) to analyse
 344 the functional significance of the regions bound by HEB (McLean, Bristor et al.,
 345 2010). In this unbiased analysis, top enriched pathways were related to actin
 346 cytoskeleton and ECM organization (Fig. 6b), resembling the pathways negatively
 347 regulated by ID4 in the gene expression profiling (Fig. 4b). This suggests that ID4
 348 mediates repression through its physical interaction with HEB.

349 The consensus peaks were annotated to 1320 genes, using the default GREAT basal
 350 plus extension gene annotation rule (McLean et al., 2010). We overlapped these
 351 genes with those regulated following ID4 KD to determine the genes directly
 352 regulated by HEB (Fig. 6c). Approximately 10% of these genes had an associated HEB
 353 peak, which is more than expected by chance ($p < 5.65E-13$; hypergeometric test).
 354 The remainder of the genes regulated by ID4 KD are likely due to secondary effects
 355 of ID4 KD or HEB independent mechanisms. The fact that there was a similar number

356 of genes overlapping in both upregulated (46) and downregulated (43) genes,
357 suggests that HEB can bind to sites both negatively and positively regulated by ID4.
358 In line with this, E-proteins have previously been demonstrated to act as both
359 activators and repressors of gene transcription by recruiting different co-factors
360 (Bayly, Chuen et al., 2004, Zhang, Kalkum et al., 2004).

361 In parallel we performed ChIP-seq for three histone modifications; H3K4Me3 (active
362 promoter mark), H3K27Ac (active enhancer mark) and H3K27Me3 (repressive
363 chromatin mark) to elucidate the chromatin context of HEB-bound regions. A
364 number of HEB peaks associated with ID4 regulated genes demonstrated a bimodal
365 distribution of H3K27Ac signal (Fig. 6d and Fig. S6b) suggesting that HEB binds to
366 enhancers. Some of the peaks were also localized to active promoters (Fig. 6d and
367 Fig. S6b). HEB peaks were observed at enhancer-marked chromatin upstream of ID4-
368 repressed ECM genes expressed in myoepithelial cells (Barsky & Karlin, 2005) such as
369 *Sparc*, *Col1a1*, *Col1a2*, *Col3a1* and *Col5a1* (Fig. 6e and Fig. S6c). HEB binding was also
370 observed near the promoter of contractile gene *Cnn2* (Fig. 6e). Of relevance *Cnn2*,
371 encoding Calponin 2, was recently discovered to be regulated by a super enhancer
372 specifically accessible in myoepithelial cells (Pervolarakis, Nguyen et al., 2019). This
373 further indicates that ID4 inhibits HEB's ability to activate transcription of genes that
374 define the myoepithelial fate.

375 We next performed HEB ChIP-seq on Comma-D β cells in which ID4 had been
376 depleted by siRNA in order to determine whether HEB DNA binding is augmented
377 when released of inhibition by ID4. Western blotting revealed that ID4 protein was
378 reduced to approximately 20% of control levels, while HEB expression was
379 unchanged (Fig. S6d). E-box motifs were again enriched in both conditions (Fig. S6e).

Differential binding analysis revealed a total of 290 regions changing upon ID4 depletion ($p < 0.05$) (Table S5). More peaks were increased than decreased (263 compared to 27) suggesting that, as expected, depletion of ID4 increased HEB's DNA binding activity (Fig. 6f and Fig. S6f). GREAT analysis revealed that the peaks that increased were involved in processes such as gland morphogenesis, skeletal development, and branching morphogenesis (Fig. S6g). No pathways were enriched in the regions that were decreased when ID4 was knocked down. Finally, we observed an increase in HEB binding in cells depleted of ID4, specifically at genes that were differentially expressed by ID4 KD (Fig. 6g). Together, our ChIP-seq analysis suggests that HEB binds to E-box motifs in regulatory elements of basal developmental genes involved in ECM and the contractile cytoskeleton, and this is antagonised by its interaction with ID4 (Fig. 6h).

Discussion

The mammary gland is the defining morphological feature of mammals and is also the source of one of the most common human cancers. Beyond their role in milk ejection at lactation, myoepithelial cells can also sense, synthesize and remodel the ECM, and control the polarity of neighbouring luminal cells (Barsky & Karlin, 2005, Gudjonsson, Rønnov-Jessen et al., 2002, Ingthorsson, Hilmarsdottir et al., 2015). Furthermore, myoepithelial cells are “natural tumour suppressors”; acting as physical barriers to prevent cancer cell invasion, secreting factors that limit growth and rarely undergoing malignant transformation (Gudjonsson, Adriance et al., 2005, Sternlicht, Kedeshian et al., 1997). Thus, understanding the regulation of the basal lineage has implications for fundamental mammary gland biology as well as breast

403 cancer.

404

405 Both bipotent and unipotent stem cells have been identified within the mammary
 406 epithelium (reviewed in (Visvader & Clevers, 2016)). ID4 has previously been
 407 demonstrated to block luminal commitment of basal cells via inhibition of key
 408 luminal driver genes including *Elf5*, *Notch*, *Brca1*, *Esr1*, *PR* and *FoxA1* (Best et al.,
 409 2014, Junankar et al., 2015). We now show that ID4 can also block genes associated
 410 with myoepithelial differentiation, in part through the inhibition of the E-protein
 411 HEB. The dual inhibition of both luminal and myoepithelial differentiation by ID4
 412 may protect the stem-cell phenotype of uncommitted basal cells during
 413 development. Subsequent downregulation of ID4, through a currently unknown
 414 mechanism, may then allow basal cells to adopt a luminal or myoepithelial fate
 415 depending on the cellular context.

416

417 HEB has been implicated in the specification of lymphocyte (Braunstein & Anderson,
 418 2012), haematopoietic (Li, Brauer et al., 2017), mesodermal (Yoon, Foley et al.,
 419 2015), neuronal (Mesman & Smidt, 2017), and skeletal muscle (Conway, Pin et al.,
 420 2004) lineages. It however has not been associated with lineage commitment of
 421 epithelial tissues. Our proteogenomic analyses support the model outlined in Fig. 6h.
 422 When ID4 is highly expressed, such as in cap cells, it sequesters HEB off chromatin,
 423 preventing expression of differentiation genes, thus determining a stem-like state
 424 (Fig. 6h; left). When ID4 expression is low, HEB is able to bind to E-box DNA motifs at
 425 promoters/enhancers to activate transcription of developmental genes that specify
 426 functional myoepithelial cells (Fig. 6h; right). Further functional studies are needed

427 to demonstrate HEB's requirement in promoting myoepithelial differentiation.

428 Compared with the luminal lineage, the molecular regulators controlling the basal

429 lineage remain poorly understood. One of the few transcription factors known to

430 promote myoepithelial differentiation is MADS-box protein SRF and its associated

431 co-activator MRTFA (Li et al., 2006, Sun et al., 2006). Interestingly, HEB and SRF

432 cooperatively activate transcription of *Acta2* in cultured fibroblasts (Kumar, Hendrix

433 et al., 2003). This occurred in an E-box dependent manner and was inhibited by ID1

434 and ID2 overexpression (Kumar et al., 2003). While we show that ID4 suppresses

435 *Acta2*, we did not observe HEB binding to the *Acta2* promoter. However, it is

436 possible that in the absence of ID4, HEB and SRF cooperate to drive expression of

437 other myoepithelial genes. This is supported by the positive enrichment of SRF

438 targets upon ID4 depletion. Subsequent studies should test whether HEB and SRF

439 cooperate in mammary epithelial cells.

440

441 The newly elucidated roles of ID4 in regulating myoepithelial commitment and ECM

442 deposition expand upon why ID4 is required for pubertal mammary gland

443 morphogenesis (Best et al., 2014, Dong et al., 2011, Junankar et al., 2015). TEBs

444 undergo collective migration, enabling the coordinated movement of adherent cells

445 into the stromal fat pad (Ewald, Brenot et al., 2008). We hypothesise that the high

446 levels of ID4 in cap cells prevent epithelial cells from acquiring a

447 mesenchymal/myoepithelial phenotype. Similar mechanisms have been observed

448 for the transcriptional repressor OVOL2, which inhibits EMT to allow for collective

449 migration (Watanabe, Villarreal-Ponce et al., 2014), and for C/EBP α , which maintains

450 epithelial homeostasis of human mammary epithelial cells (Lourenço, Roukens et al.,

2020). During ductal elongation, collagenous stromal ECM is absent directly in front of the invading TEBs (Paine & Lewis, 2017, Silberstein, Strickland et al., 1990, Sternlicht, 2006). We show that ID4 suppresses collagen synthesis and deposition around TEBs, which may otherwise act as a physical barrier to impede invasion. In support of this idea, ectopic deposition of collagen by mammary epithelial cells induced by exogenous TGF- β , or forced expression of recombinant type I collagen which is resistant to collagenase attack, causes ensheathment of TEBs by collagen, and retardation of ductal elongation (Feinberg, Zheng et al., 2018, Silberstein et al., 1990).

Developmental transcription factors are often dysregulated in cancer. ID4 is highly expressed in ~50% of basal-like breast cancer (BLBC) cases and associates with poor prognosis (Baker, Holliday et al., 2016, Junankar et al., 2015). Given the cell-intrinsic role of ID4 in promoting growth/proliferation and inhibiting differentiation in the mammary gland, it is easy to envision how overexpression of ID4 could lead to an aggressive breast cancer phenotype. Additionally, the suppression of ECM synthesis may allow tumour cells to easily invade into the surrounding stroma, akin to the invasion of cap cells during ductal elongation. Future work should test whether the mechanisms discovered here are conserved in cancer. If so, targeting ID4 may represent an attractive therapeutic strategy for ID4-positive BLBCs. Excitingly, a small-molecule Pan-ID inhibitor has recently been identified by the Benezra laboratory, which works by inhibiting the interaction between ID and E-proteins (Wojnarowicz, Lima E Silva et al., 2019).

475 To conclude, these new insights into ID4 and HEB function have helped unravel
476 regulation within the elusive basal differentiation hierarchy, with broader
477 implications for the regulation of epithelial stem cells in general, and also in tumour
478 progression.

479 **Materials and methods**

480 **Mice**

481 All mice experiments were performed in accordance with the ethical regulations of
482 the Garvan Institute Animal Experimentation Committee. ID4 KO mice were
483 generated as previously described (Yun et al., 2004). The founder ID4-FlagV5 mouse
484 was engineered using Crispr/Cas9 gene editing by the Mouse Engineering at
485 Garvan/ABR (MEGA) facility. The ID4floxGFP mice were generated as previously
486 described (Best et al., 2014). All mice used were on the FVB/N background. For the
487 gene expression profiling experiment, mice were synchronised in estrus to reduce
488 hormone-induced gene expression variation (Dalal, Estep et al., 2001), and checked
489 by vaginal swab cytology.

490

491 **Mammary epithelial cell preparations**

492 Mammary epithelial cells were prepared from freshly harvested 3rd and 4th
493 mammary glands pooled from 4-8 female mice at the indicated ages. Glands were
494 mechanically disrupted using a McIlwain tissue chopper then digested with 15,000 U
495 collagenase (Sigma-Aldrich) and 500 U hyaluronidase (Sigma-Aldrich) in FV media
496 (DMEM/F12 (Gibco), 5% (v/v) FBS (HyClone) 10 mM HEPES (Gibco), 0.14 IU/mL
497 Insulin (Novo Nordisk), 500 ng/mL Hydrocortisone (Sigma-Aldrich), 20 ng/mL Cholera

498 Toxin (Sigma-Aldrich), 2 ng/mL mEGF (Life Technologies)). Digestion was carried out
 499 in a shaking incubator for 1 hr at 37°C 225 rpm. The resulting organoids were further
 500 digested with warm 0.25% trypsin for 1 min with constant pipetting followed by
 501 treatment with 5 mg/mL Dispase (Roche) for 5 min in at 37°C. Cells were incubated
 502 with 1X red blood cell lysis buffer (BD Biosciences) for 5 min at room temperature.
 503 Cells were passed through a 70 µm strainer followed by a 40 µm strainer to remove
 504 clumps and debris. Live cells were counted using trypan blue (Thermo Fisher
 505 Scientific) and a haemocytometer prior to use in downstream applications.

506

507 Flow cytometry and FACS

508 Single-cell suspensions of mouse mammary epithelial cells were blocked in Fc block
 509 cocktail (FACS buffer, 6.25 µg/mL Mouse BD Fc block (BD Bioscience) and 200 µg/mL
 510 Rat Gamma Globulin (Jackson ImmunoResearch)) for 10 min on ice. Fluorophore- or
 511 biotin-conjugated antibodies were diluted in Fc block cocktail and incubated on the
 512 cells on ice for 20 min in the dark. Antibodies used were CD24-PE (1:200, BD
 513 Biosciences, Clone M1/69), EPCAM-PerCP/Cy5.5 (1:200, Biolegend, Clone G8.8),
 514 CD29-APC/Cy7 (BioLegend, Clone HMβ1-1), CD61-APC (1:50, Thermo Fisher
 515 Scientific, Clone HMβ3-1), CD49f-APC (1:100, BioLegend, Clone GoH3), Ter119-biotin
 516 (1:80, BD Biosciences, Clone TER119), CD45-biotin (1:100, BD Biosciences, Clone 30-
 517 F11), CD31-biotin (1:40, Biolegend, Clone 390) and BP1-biotin (1:500, Thermo Fisher
 518 Scientific, Clone 6C3). Cells were washed and stained with streptavidin-BV421
 519 (1:100, Biolegend). Following washing, cells were resuspended in FACS buffer at a
 520 density of 1×10^7 cells/mL and DAPI was added (1:1000, Invitrogen). FACS was
 521 performed on a FACS Aria III 4 laser 15 colour sorter with BD FACS DIVA software.

For analytical flow cytometry, a CytoFLEX 3 laser 13 colour flow cytometer (Beckman Coulter) was used. All flow cytometry data was analysed using FlowJo software version 10 (Tree Star Inc).

RNA-seq

RNA was extracted using the Qiagen miRNease mini kit (Qiagen) for Comma-D β cells or the Qiagen miRNease micro kit (Qiagen) for FACS-sorted mammary epithelial cells. Cells were sorted directly into 700 μ L Qiazol lysis reagent to minimise cell loss, with a maximum of 5×10^4 cells per 1.5 mL Eppendorf tube. RNA was eluted in RNase free water. The Qubit RNA BR Assay kit (Thermo Fisher Scientific) was used to measure RNA concentration and RNA integrity was determined using the Agilent Bioanalyser 2100 with the 6000 Nano Assay (Agilent Technologies).

For low input RNA extracted from FACS sorted mammary glands, the Ovation RNA-seq System V2 kit (NuGEN) was used to synthesise cDNA with RNA inputs ranging from 0.5-2 ng as per the manufacturers protocol. The Ovation Ultralow System V2 kit was then used to prepare libraries from the cDNA. cDNA libraries were sequenced on a HiSeq 2500 system (high output mode) (Illumina), with 125 bp paired-end reads. For RNA extracted from Comma-D β cells, The Illumina TruSeq Stranded mRNA Library Prep Kit (Illumina) was used with 1 μ g of input RNA following the manufacturer's instructions. cDNA libraries were sequenced on the NextSeq system (Illumina), with 75 bp paired-end reads.

Quality control was checked using FastQC (Andrews, 2010) to remove poor quality reads. Illumina sequencing adapters were then trimmed using Cutadapt (Martin, 2011). Reads were then aligned to the mouse reference genome mm10 using STAR

546 ultrafast universal RNA-seq aligner (Dobin, Davis et al., 2013). RSEM accurate
547 transcript quantification for RNA-seq data was used to generate a gene read count
548 table and to filter genes with read counts of 0 (Li & Dewey, 2011). Differential gene
549 expression analysis was performed using edgeR (McCarthy, Chen et al., 2012,
550 Robinson, McCarthy et al., 2010) and voom/limma (Ritchie, Phipson et al., 2015) R
551 packages. Genes were ranked based on the limma moderated t-statistic and this was
552 used as input GSEA pre-ranked (Subramanian, Tamayo et al., 2005) using Molecular
553 Signature Database (MSigDB) collections (v7.1). The EnrichmentMap Cytoscape
554 plugin (Merico, Isserlin et al., 2010) was used to visualise the GSEA results.

555

556 **Analysis of public single-cell RNA-seq data**

557 Raw Unique Molecular Identifier (UMI) expression data was taken from Bach *et al.*
558 and re-processed using Seurat v2 using default parameters, as recommended by the
559 developers (Satija, Farrell et al., 2015). All basal cell clusters, defined by *Krt5* and
560 *Krt14* expression, were extracted for subsequent analysis. Developmental
561 trajectories were inferred using Monocle 2 (Qiu, Mao et al., 2017). Clustering was
562 first performed using the 'densityPeak' and 'DDRTree' methods, with a delta local
563 distance threshold of 2, and a rho local density threshold of 3. For inferring
564 pseudotime, we first selected all genes with a mean expression greater than 0.5, and
565 an empirical dispersion greater than 1, and proceeded with the top 1000 significant
566 genes for ordering. Differential gene expression between the two branch states were
567 performed using the MAST algorithm through the Seurat package (Finak, McDavid et
568 al., 2015, Satija et al., 2015). For comparisons between *Id4*-high and *Id4*-low states,
569 we first filtered for cells with the detection of *Id4* to avoid technical biases from gene

570 drop out. Differential gene expression was performed (as described above) between
571 the top and bottom 200 cells grouped based on log normalised gene expression of
572 *Id4*.

573

574 **Immunohistochemistry and immunofluorescence on mammary tissue sections**

575 IHC staining was performed on 4 µm sections of formalin-fixed paraffin-embedded
576 (FFPE) tissue. Slides were dewaxed with xylene and hydrated through graded
577 alcohols. Antigen retrieval was performed for 1 min in a pressure cooker in DAKO
578 target retrieval reagent s1699. A DAKO Autostainer was used for subsequent steps.
579 Briefly, slides were incubated with DAKO peroxide block for 5 min and blocked with
580 DAKO protein block for 30 min. ID4 Primary antibody (1:400, Biocheck BCH9/82-12)
581 was incubated on the slides for 60 min. Following washing, slides were incubated
582 with Envision rabbit secondary antibody (Agilent Technologies, Santa Clara, CA, USA)
583 for 30 min. Slides were washed then incubated with DAKO DAB+ (Agilent
584 Technologies) reagent for 10 min. Slides were then rinsed and counterstained with
585 haematoxylin for 20-30 sec and dehydrated through graded alcohols, cleared using
586 xylene and mounted using Ultramount #4 (Fronine). Bright-field images were
587 captured using a Leica DM 4000 microscope with high-resolution colour camera
588 (DFC450).

589 Immunofluorescence was performed manually on antigen retrieved tissue sections
590 prepared as described for IHC. Slides were blocked with Mouse on Mouse (MOM)
591 blocking buffer (Vector Biolabs) for 1 hour. Primary antibody was diluted in MOM
592 diluent and incubated overnight at 4°C. Antibodies used were ID4 (1:400, Biocheck
593 BCH9/82-12), α-SMA (1:200, Sigma-Aldrich A5228), HEB (1:200, Protein Tech 14419-

1-AP), V5 (1:200, Santa Cruz sc-58052). Following washing, slides were incubated with fluorescent secondary antibody (1:500, Jackson ImmunoResearch) for 1 hr at room temperature. Nuclei were stained using Hoechst 33342 (Sigma-Aldrich). Slides were mounted using Prolong Diamond mounting media (Thermo Fisher Scientific). Fluorescent images were captured using a Leica DM 5500 microscope. For quantification of fluorescence of ID4 and α -SMA in cap and ductal basal cells, 3 representative ducts and TEBs were imaged from each mouse (n=9). Using FIJI software, a region of interest was drawn around 10 randomly selected basal cells from each image. The mean fluorescence of ID4 and α -SMA was determined within each cell.

Picrosirius red staining

Picrosirius red staining was performed following as per (Vennin, Chin et al., 2017, Vennin, Méléneec et al., 2019). Briefly, dewaxed FFPE sections were stained with haematoxylin. Sections were then treated with 0.2% Phosphomolybdic acid (Sigma-Aldrich) followed by 0.1% Sirius red (Sigma-Aldrich) Picric Acid-Saturated Solution (Sigma-Aldrich). Slides were then rinsed with acidified water (90 mM glacial acetic acid) then with 70% ethanol. A Leica DM 4000 microscope was used to image total collagen staining in the tissue sections by phase contrast. To image birefringence of collagen fibres, 2 polarised filters were used. Matched phase-contrast and polarised images were taken for each region of interest. Image analysis was performed using FIJI image analysis software.

Proximity Ligation Assay

618 Comma-D β cells were grown on glass coverslips in 6 well plates until ~80% confluent.
 619 Cells were fixed in 4% Paraformaldehyde (PFA) (ProSciTech) for 15 min then
 620 permeabilised in 0.2% Triton-X-100 for 15 min. The Duolink Proximity Ligation Assay
 621 (PLA) (Sigma-Aldrich) was performed as per the manufacturer's protocol and using
 622 the following antibodies: ID4 (1:50, Santa Cruz sc-365656) and HEB (1:200, Protein
 623 Tech 14419-1-AP), and equivalent concentrations of IgG negative controls (sc-2027
 624 and sc-2025). DAPI (Life Technologies) and Phalloidin (Life Technologies) were added
 625 to the final wash step. Coverslips were mounted onto glass slides with Prolong
 626 Diamond mountant (Thermo Fisher Scientific).
 627 Cells were imaged using a Leica DMI Sp8 confocal microscope (63X oil objective). Six
 628 random fields of view per image were captured, each with approximately 50 cells,
 629 and images were quantified using Andy's algorithms Fiji package (Law, Yin et al.,
 630 2017) to enumerate the number of PLA foci per nuclei.

631

632 **Cell lines**

633 The mouse mammary epithelial cell line Comma-D β was a gift was a gift from Joseph
 634 Jeffery (University of Massachusetts, Amherst, MA, USA). Comma-D β cells were
 635 maintained in DMEM/F12 media (Gibco) supplemented with 2% FBS (HyClone), 10
 636 mM HEPES (Gibco), 0.125 IU/mL Insulin (Novo Nordisk) and 5 ng/mL mEGF (Life
 637 Technologies). The human mammary epithelial cell line PMC42 was a gift from
 638 Professor Leigh Ackland (Deakin University, Melbourne, Victoria, Australia). PMC42
 639 cells were maintained in RPMI 1640 (Gibco) supplemented with 10% FBS (HyClone).
 640 The MCF10A cell line was obtained from the American Type Culture Collection and
 641 were maintained in DMEM/F12 (Gibco), 5% Horse Serum (Thermo Fisher Scientific),

642 20 ng/mL hEGF (In Vitro Technologies), 0.5 mg/mL Hydrocortison (Sigma-Aldrich),
643 100 ng/mL Cholera Toxin (Sigma-Aldrich) and 0.125 IU/mL Insulin (Novo Nordisk).

644

645 **Conditional reprogramming of primary mouse basal cells**

646 Viable basal cells were purified by FACS from 10-12 week old mice as described
647 above. Cells were collected into FAD media (DMEM/F12 3:1 (Gibco) supplemented
648 with 10% FBS, 0.18 mM Adenine (Sigma-Aldrich), 500 ng/mL Hydrocortisone (Sigma-
649 Aldrich), 8.5 ng/mL Cholera Toxin (Sigma-Aldrich), 10 ng/mL mEGF (Life
650 Technologies), 0.14 IU/mL Insulin (Novo Nordisk), 5 μ M Y-27632 (Seleckchem), 1X
651 AB/AM (Gibco) and 50 μ g/mL Gentamicin (Life Technologies). Basal cells were
652 maintained in culture using as per (Prater et al., 2014). Tissue culture flasks coated
653 with Growth Factor-reduced Matrigel (Corning) diluted 1:60 in PBS for 30 min to 1 hr
654 at 37°C. Excess Matrigel/PBS solution was aspirated from the flasks and basal cells
655 were seeded at a density of $\sim$ 5000 cells/cm². Basal cells were co-cultured with
656 irradiated (50 Gy) NIH-3T3 feeder cells at a density of 10,000 cells/cm². Cells were
657 maintained in a 37°C low 5% oxygen 5% CO₂ incubator. Differential trypsinisation
658 was used to first remove the less-adherent 3T3 cells when passaging.

659

660 **Organoid culture**

661 A 40 μ L plug of Matrigel (Corning) was added to each well of 8-well chamber slides
662 (Corning) and allowed to set for 30 min in a 37°C incubator. Primary cells were
663 resuspended in Epicult-B (Stemcell Technologies) media containing 2% Matrigel and
664 12,000 cells were seeded into each chamber in a total volume of 400 μ L. Chambers

665 were observed using a light microscope to ensure that cells were not aggregated.

666 Organoids were allowed to form over 1 week and media was changed every 3 days.

667

668 **Immunofluorescent staining of organoids**

669 Organoids were stained within the chamber slides. Media was aspirated and

670 organoids were fixed with 2% PFA diluted in PBS for 20 min at room temperature.

671 Organoids rinsed with PBS for 5 min following permeabilisation with 0.5% Triton X-

672 100 in PBS for 10 min at 4°C then rinsed 3 times with 100 mM Glycine (Astral

673 Scientific) in PBS for 10 min each. Organoids were blocked for 1 hr in IF buffer with

674 10% goat serum (Vector Laboratories). Primary antibody made up in blocking buffer

675 was incubated in the chambers overnight at 4°C in a humidified chamber. Antibodies

676 used were ID4 (1:200, Biocheck BCH9/82-12), α -SMA (1:100, Abcam ab5694), KRT14

677 (1:1000, Covance PRB-155P), KRT8 (1:500, DSHB TROMA1) and P63 (1:100, Novus

678 NB100-691). The following day, slides were equilibrated to room temperature for 1-2

679 hr. Organoids were rinsed 2 times with IF buffer (0.1% BSA, 0.2% Triton-X-100, 0.05%

680 Tween-20 in PBS) for 20 min each. Fluorescent secondary antibody (Jackson

681 ImmunoResearch) diluted in blocking buffer (1:500) was added to the chambers and

682 incubated for 45 min in the dark. Secondary antibody was rinsed off for 20 min in IF

683 buffer, then 2 times with PBS for 10 min each. DAPI (Life Technologies) was

684 incubated in the chambers for 15 min followed by a 10 min PBS rinse. The walls of

685 the chamber slides were removed and one drop of Prolong Diamond mounting

686 media (Thermo Fisher Scientific) was added to each well. Slides were coverslipped

687 and edges sealed with clear nail varnish. Slides were allowed to dry for 24 hr at room

688 temperature protected from light.

689 Organoids were imaged using a Leica DMI Sp8 confocal microscope using the 40X oil
690 objective and 3X optical zoom. The FIJI software package was used to quantify the
691 fluorescence within the organoids. For α -SMA, a circle was drawn around the entire
692 organoid and fluorescence measured. For ID4, a mask was made using the DAPI
693 channel and signal was measured within the nuclear mask. Multiple representative
694 regions of background staining were quantified. Corrected fluorescence (CF) was
695 calculated as described in (McCloy, Rogers et al., 2014). CF = integrated density –
696 (area of selected organoid * mean fluorescence of all background readings)

697

698 **Cre mediated deletion of ID4 from primary cells**

699 Following differential trypsinisation to remove irradiated 3T3 feeder cells, 1.0-
700 1.5×10^5 ID4floxGFP primary basal cells were seeded into a T75 with 7.5×10^5 fresh
701 irradiated 3T3 cells in 20 mL FAD media. Adenovirus (Vector Biolabs) was added to
702 the media at a multiplicity of infection (MOI) of 100. Adenoviruses' used were codon
703 optimised Cre (iCre) and GFP adenovirus (#1772) and control eGFP adenovirus
704 (#1060). Infection efficiency was assessed the following day using a fluorescence
705 microscope to check GFP expression within the cells. Cells were harvested after 72 hr
706 for downstream analysis.

707

708 **siRNA transfections**

709 Comma-D β cells were seeded at a density of 1.5×10^4 cells/cm² into 6 well plates
710 (Corning) for RNA-seq or 100 mm dishes (Corning) for ChIP-seq in antibiotic-free
711 media. The following day siRNA constructs (Dharmacon) were transfected into the
712 cells using Dharmafect-4 transfection reagent (Dharmacon) as per the

713 manufacturer's instructions at 20 nM. siRNA used were siGENOME Mouse Id4
714 SMARTpool (M-043687) and ON-TARGETplus Non-targeting Control Pool (D-001810).
715 Media was changed the following day and cells were harvested 48 hr post-
716 transfection.

717

718 Western blotting

719 Cells were lysed in RIPA buffer (50 mM Tris-HCl pH 7.4, 1% NP-40, 0.5% Sodium
720 Deoxycholate, 0.1% SDS, 1% Glycerol, 137.5 mM NaCl, 100 μ M Sodium
721 Orthovanadate, 20 μ M MG132, 1 mM DTT and 1x cOmplete ULTRA Tablet (Roche))
722 and protein concentration was quantified using the Pierce BCA Protein Assay Kit
723 (Thermo Fisher Scientific) according to the manufacturer's instructions. Protein
724 lysates (15-30 μ g protein) were mixed with 1X NuPage loading buffer (Life
725 Technologies) and 1X NuPage reducing agent (Life Technologies) and denatured by
726 heating at 85°C for 5 min. Samples were run on a 4-12% Bis/Tris gels (Life
727 Technologies) in MES or MOPS running buffer (Life Technologies). Protein was
728 transferred to a 0.45 μ m PVDF membrane (Merck Millipore) using BioRad transfer
729 modules in transfer buffer (25 mM Tris, 192 mM Glycine, 20% Methanol).
730 Membranes were blocked in Odyssey blocking buffer (LiCOR) for 1 hr at room
731 temperature then incubated in primary antibody diluted in 5% BSA/TBS overnight at
732 4°C. The following antibodies were used for western blotting: ID4 (1:20,000,
733 Biocheck BCH9/82-12), E2A (1:1000, from Dr Nicolas Huntington and Prof Lynn
734 Corcoran from the Walter Eliza Hall Institute (WEHI)), HEB (1:1000, from Dr Nicolas
735 Huntington and Prof Lynn Corcoran, WEHI or 1:1000 ProteinTech 144191-1-AP), Flag
736 (1:5000, Sigma-Aldrich F1804), V5 (1:200, sc-58052), β -Actin (1:1000, Sigma-Aldrich

737 A5441) and α -Tubulin (1:1000, Santa Cruz sc-5286). Fluorescent secondary antibody
738 conjugated to IRDye680 or IRDye800 (LiCOR) diluted in Odyssey blocking buffer
739 (1:15,000 - 1:20,000) were used for detection. An Odyssey CLx Infrared Imaging
740 System (LiCOR) was used to image and quantify the western blots.

741

742 **Co-immunoprecipitation**

743 Pierce Protein A/G magnetic beads (Thermo Fisher Scientific) were incubated with
744 antibody for a minimum of 4 hr at 4°C on a rotating platform. For each IP 15 μ L of
745 beads and 2.5 μ g antibody were used. Immunoprecipitating antibodies used were
746 ID4 (pool of Santa Cruz sc-491 and sc-13047), E2A (from Dr Nicolas Huntington and
747 Prof Lynn Corcoran, WEHI), HEB (Santa Cruz sc-357 and antibody from Dr Nicolas
748 Huntington and Prof Lynn Corcoran, WEHI), Flag (Sigma-Aldrich F1804), V5 (Santa
749 Cruz sc-58052 and Thermo Fisher R960-25), species matched IgG negative controls
750 (sc-2027, sc-2025 and BioLegend 400602). Protein was extracted in IP lysis buffer
751 (10% Glycerol, 0.03% $MgCl_2$, 1.2% HEPES, 1% Sodium acid pyrophosphate, 1% Triton-
752 X, 0.8% NaCl, 0.4% NaF, 0.04% EGTA, 1x cOmplete ULTRA Tablet (Roche), 100 μ M
753 Sodium Orthovanadate, 20 μ M MG132, 1 mM DTT and cells were passed 5 times
754 through a 23 g needle to aid in lysis. Protein lysates (0.5-2 mg per IP) were added to
755 washed beads and incubated overnight at 4°C on a rotating platform. Following
756 washing, beads were resuspended in 2X NuPage loading buffer (Life Technologies)
757 and 2X reducing agent (Life Technologies) in IP lysis buffer. Samples were incubated
758 at 85°C for 5 min. Beads were separated on a magnetic rack and supernatant loaded
759 onto NuPage gel for SDS-PAGE and western blotting. For detection of protein,
760 fluorescent TrueBlot secondary antibodies were used (Jomar Life Research).

761

762 **RIME**

763 The RIME protocol was adapted from the protocol developed by Mohammed *et al.*
 764 (Mohammed et al., 2013). Comma-D β cells were grown in 150 mm tissue culture
 765 dishes (Corning) until 80-90% confluent. A total of 8x 150 mm dishes were used per
 766 RIME sample. Cells were fixed with 1% PFA (ProSciTech) in DMEM/F12 (Gibco) for 7
 767 min at room temperature on a rocking platform. Cross-linking was quenched by
 768 addition of molecular grade glycine (Astral Scientific) to a final concentration of 125
 769 mM for 2 min on a rocking platform. Cells were washed twice with ice-cold PBS
 770 (Gibco) and scraped in 1 mL of ice-cold PBS containing magnesium and calcium salts
 771 (Gibco). Cells were centrifuged at 4°C for 5 min at 1200 rpm. Cell nuclei were
 772 enriched through a series of lysis buffers. Cells were first resuspended in 10 mL lysis
 773 buffer 1 (50 mM HEPES-KOH pH 7.5, 140 mM Sodium, 1 mM EDTA, 10% Glycerol,
 774 0.5% NP-40 or Igepal CA-630, 0.25% Triton X-100, 100 μ M Sodium Orthovanadate,
 775 20 μ M MG132, 1 mM DTT and 1x cOmplete ULTRA Tablet (Roche) and incubated for
 776 30 min at 4°C on a rotating platform. Cells were then centrifuged at 3000 g for 3 min
 777 at 4°C and supernatant removed. The pellet was then resuspended in 10 mL lysis
 778 buffer 2 (10 mM Tris-HCl pH 8.0, 200 mM NaCl, 1 mM EDTA, 0.5 mM EGTA, 100 μ M
 779 Sodium Orthovanadate, 20 μ M MG132, 1 mM DTT and 1x cOmplete ULTRA Tablet
 780 (Roche)) and incubated for a further 30 min at 4°C on a rotating platform. Following
 781 centrifugation, the resulting nuclei pellet was lysed in 2.5 mL lysis buffer 3 (10 mM
 782 Tris-HCl pH 8.0, 100 mM NaCl, 1 mM EDTA, 0.5 mM EGTA, 0.1% Sodium
 783 Deoxycholate, 0.5% N-lauroylsarcosine, 100 μ M Sodium Orthovanadate, 20 μ M
 784 MG132, 1 mM DTT and 1x cOmplete ULTRA Tablet (Roche)). The nuclear lysate was

785 sonicated using a Bioruptor sonicator (Diagenode) for 20 cycles of 30 sec on/30 sec
786 off. Triton-X-100 was added to a final concentration of 1% and the sample was then
787 centrifuged at 4700 g for 5 min at 4°C to remove cellular debris.
788 Immunoprecipitation was carried out from the sheared nuclear supernatant
789 described above with 20 µg of antibody and 100 µL of magnetic beads per sample.
790 The following day, the beads were washed 10 times with RIPA buffer then 5 times
791 with 100 mM ammonium hydrogen carbonate (AMBIC) (Sigma-Aldrich) solution to
792 remove salts and detergents, resuspended in 50 µL AMBIC solution and transferred
793 into clean tubes.

794

795 **Mass Spectrometry**

796 Samples were processed as described in (Huang, Yang et al., 2015) for liquid
797 chromatography coupled mass spectrometry (LC-MS/MS) using Sequential
798 Windowed Acquisition of all THEoretical fragment-ion spectra (SWATH) acquisition.
799 Briefly, samples were denatured in 100 mM triethylammonium bicarbonate and 1%
800 sodium deoxycholate, disulfide bonds were reduced in 10 mM DTT, alkylated in 20
801 mM iodo acetamide, and proteins digested on the magnetic beads using trypsin.
802 After C18 reversed phase (RP) StageTip sample clean up, peptides were analysed
803 using a TripleToF 6600 mass spectrometer (SCIEX, MA, USA) coupled to a nanoLC
804 Ultra 2D HPLC system (SCIEX). Peptides were separated for 60 min using a 15 cm chip
805 column (ChromXP C18, 3 µm, 120 Å) (SCIEX) with an acetonitrile gradient from 3-
806 35%. The MS was operated in positive ion mode using either a data dependent
807 acquisition method (DDA) or SWATH acquisition mode. DDA was performed of the
808 top 20 most intense precursors with charge stages from 2+ - 4+ with a dynamic

809 exclusion of 30 s. SWATH-MS was acquired using 100 variable size precursor
810 windows. DDA files were searched using ProteinPilot 5.0 (SCIEX) against the
811 reviewed UniProt *Mus musculus* protein database (release February 2016) using an
812 unused score of 0.05 with decoy search strategy enabled. These search outputs were
813 used to generate a spectral library for targeted information extraction from SWATH-
814 MS data files using PeakView v2.1 with SWATH MicroApp v2.0 (SCIEX) importing only
815 peptides with < 1% FDR. Protein areas, summed chromatographic area under the
816 curve of peptides with extraction FDR $\leq$ 1%, were calculated and used to compare
817 protein abundances between bait and control IPs.

818

819 **ChIP-seq**

820 Our ChIP-seq protocol was adapted from (Khoury, Achinger-Kawecka et al., 2020).
821 Comma-D β cells were grown in 150 mm or 100 mm dishes (Corning) until 80-90%
822 confluent. For the ChIP-seq experiment on unperturbed cells 2x 150 mm dishes were
823 used for HEB ChIP and 1x 150 mm dishes were used for histone marks. For the ID4
824 siRNA experiment 3x 100 mm dishes per condition were used. Cells were scraped in
825 ice-cold PBS (Gibco) and resuspended by passing 10 times through a 19 g syringe.
826 Cells were fixed in 1% PFA (ProSciTech) in PBS (Gibco) for 15 min at room
827 temperature followed by quenching with glycine to a final concentration of 125 mM
828 for 5 minutes followed by two PBS washes. Nuclei was extracted by resuspending
829 cells in nuclei extraction buffer (10 mM Tris-HCl pH 7.5, 10 mM NaCl, 3 mM MgCl₂,
830 0.1 mM EDTA, 0.5% IGEPAL, 1x cComplete ULTRA Tablet (Roche)) and incubated on
831 ice for 10 min followed by 20 passes on a tight Dounce homogeniser, or until nuclei
832 were extracted. Nuclei were visually inspected using trypan blue and a

833 haemocytometer. Nuclei were pelleted by centrifugation and resuspended in
834 sonication buffer (50 mM Tris-HCl pH 8, 1% SDS, 10 mM EDTA, 1x cOmplete ULTRA
835 Tablet (Roche)) and sonicated using a Bioruptor sonicator (Diagenode) to achieve
836 DNA fragments between 100-500 bp with a mean fragment size of approximately
837 300 bp. In the siRNA experiment, protein was quantified using the Pierce BCA assay
838 to load equal amounts of input material for NT and ID4 siRNA for
839 immunoprecipitation. Sonicated sample was diluted with IP dilution buffer to 1 mL
840 (16.7 mM Tris-HCl pH 8, 0.01% SDS, 1% Triton X-100, 167 mM NaCl, 1.2 mM EDTA).
841 Samples were cleared with protein A/G magnetic beads for 1.5 hr at 4°C. 1% of the
842 cleared nuclear lysate was removed for the input control and immunoprecipitation
843 was performed as described previously. For HEB ChIP 100 µL of beads and 20 µg of
844 HEB antibody (sc-58052) was used with chromatin from approximately 20-30x10⁶
845 cells. For histone mark ChIP, 50 µL beads with 10 µg of the following antibodies –
846 H3K4Me3 (Active Motif 28431), H3K27Me3 (Merck Millipore 07-449), was used with
847 chromatin from approximately 10-15x10⁶ cells. The following day, beads were
848 washed for 5 min each with 1 mL of the following buffers: Low salt buffer (0.1% SDS,
849 1% Triton X-100, 2 mM EDTA, 20 mM Tris-HCl, pH 8, 150 mM NaCl), High salt buffer
850 (0.1% SDS, 1% Triton X-100, 2 mM EDTA, 20 mM Tris-HCl, pH 8, 500 mM NaCl), LiCl
851 buffer (0.25 M LiCl, 1% IGEPAL, 1% deoxycholic acid (sodium salt), 1 mM EDTA, 10
852 mM Tris, pH 8) followed by 2 washes with TE buffer (10 mM Tris-HCl, 1 mM EDTA,
853 pH 8). DNA was eluted twice in 100 µL ChIP elution buffer (1% SDS, 0.1 M Sodium
854 Biocarbonate) for 15 minutes at room temperature each. Cross-linking was reversed
855 overnight by treating samples with 200 mM NaCl and 250 µg/mL Proteinase K (New
856 England Biolabs) and incubating overnight at 65°C. The following day samples were

857 treated with 100 µg/mL RNase A (Qiagen) for 1 hr at 37°C. DNA was purified using
858 Phase Lock Gel Light tubes (Quantabio 5Prime) according to the manufacturer's
859 instructions. DNA was eluted in 20 uL nuclease-free TE buffer, pH8 (Qiagen).
860 DNA concentration was measured using the Qubit HS Assay Kit (Thermo Fisher
861 Scientific). Libraries were prepared using the Illumina TruSeq ChIP library prep kit
862 (Illumina) following the manufacturer's instructions except the gel purification step
863 was replaced with a two-sided AMPure XP bead (Beckman Coulter) size selection to
864 obtain libraries between 200-500 bp. Library sizes were verified using the 4200
865 TapeStation System (Agilent) with a D1000 ScreenTape (Agilent) and concentration
866 was determined using the Qubit HS Assay Kit (Thermo Fisher Scientific). ChIP
867 libraries were sequenced on the NextSeq system (Illumina), with 75 bp paired-end
868 reads.

869 Reads were aligned with BWA (Li & Durbin, 2009) and all reads with a MAPQ<15
870 were removed. Alignment statistics were generated using Samtools Flagstat (Li,
871 Handsaker et al., 2009). ChIP-seq peaks were called using the peak calling algorithm
872 MACS (Zhang, Liu et al., 2008) and ENCODE blacklist regions were removed. For
873 unperturbed cells, consensus peaks present in at least 2 of 4 replicates were used for
874 downstream analysis. Due to lower amount of input material available from the
875 siRNA experiment, less consensus peaks were called and for this reason analysis was
876 conducted on merged peaks from the 3 replicates. Motif enrichment analysis was
877 performed using MEME-ChIP (Machanick & Bailey, 2011). Peaks were annotated to
878 genomic features using HOMER (Heinz, Benner et al., 2010). GREAT was used for
879 functional enrichment analysis (v4.0.4) and gene annotation using the default
880 parameters (McLean et al., 2010). SeqPlots (Stempor & Ahringer, 2016) and IGV

(Robinson, Thorvaldsdóttir et al., 2011) software were used for data visualisation. Differential binding analysis was performed using the DiffBind package (Stark, 2011). For overlapping of of siLD4 RNA-seq and HEB ChIP-seq genes, a hypergeometric test was used to determine if overlap was significant, assuming 30,000 genes in the mouse genome.

Data availability

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository with the dataset identifier PXD017517. Processed data files for the RNA-seq (counts tables) and ChIP-seq (bed files) have been uploaded as Figure Source Data. Raw data files are in the process of being uploaded onto the GEO public repository and will be made available prior to publication. We can provide the raw data files upon request.

Acknowledgements

We are very grateful for Amanda Khoury for her extensive advice on the ChIP-seq experiments, Samantha Oakes for providing the mammary gland FFPE blocks from different developmental stages, David Gallego Ortega for advice on tissue dissociation and sorting and William Hughes for help with microscopy image analysis. We thankfully acknowledge the support of the Australia Postgraduate Award and The National Health and Medical Research Council of Australia (1107671). Aspects of this research were supported by access to the Australian Proteome Analysis Facility, funded by the Australian Government's National

903 Collaborative Research Infrastructure Scheme. T.R.C and J.N.S are supported by the
904 Cancer Institute NSW (171105) and a Career Catalyst grant from Susan G. Komen.

905 Author Contributions

906 H.H designed and performed the experiments and wrote the manuscript. D.R
907 analysed the RNA-seq and ChIP-seq data. S.J co-supervised the project, contributed
908 to experiments and discussions and extensively revised the manuscript. S.Z.W
909 analysed public scRNA-seq data and revised the manuscript. L.A.B advised on RIME
910 experiments and contributed to discussions. C.K and M.P.M performed mass
911 spectrometry and analysis. C.H performed RNA-seq library preps and analysis. A.M
912 assisted with mouse experiments. J.N.S and T.R.C performed picosirius red image
913 analysis. B.P assisted with ChIP-seq experiments. N.H contributed IP antibodies. C.J.O
914 co-supervised the project and revised the manuscript. J.S.C helped to conceptualise
915 the project and revised the manuscript. J.V contributed the reporter mice and
916 revised the manuscript. A.S conceived the study and was in charge of overall
917 direction and planning.

918 Conflicts of Interest

919 NH has ownership and stock options in oNKO-Innate Pty Ltd.

920 Figure legends

921 **Figure 1. Loss of ID4 results in upregulation of myoepithelial genes and**
922 **downregulation of cell growth genes in sorted mammary basal cells.**

923 **a)** Live lineage negative CD24⁺CD29^{hi}CD61⁺ basal, CD24⁺CD29^{lo}CD61⁺ luminal
924 progenitor and CD24⁺CD29^{lo}CD61⁻ mature luminal cells were isolated by FACS from

adult (10-12 weeks) ID4 wild type (WT) and knockout (KO) mice at estrus for RNA-seq. Representative FACS plots shown from 4 experiments. **b)** Number of significantly (FDR<0.05) differentially expressed genes (DEGs) upregulated (red) or downregulated (blue) in ID4 KO epithelial subpopulations compared to ID4 WT. LP = luminal progenitor, ML = mature luminal. **c)** Heat map displaying the 237 significant differentially expressed genes between ID4 WT and ID4 KO basal cells. **d)** Genes were ranked based on the limma t-statistic comparing ID4 WT and KO basal cells and GSEA was carried out using the C2all, C3TF, C5 and Hallmark gene sets. The top 10 positive and negatively enriched pathways with an FDR<0.1 are displayed. Only 3 pathways were negatively enriched with an FDR<0.1. **e)** GSEA results were visualised using the EnrichmentMap plugin on Cytoscape. Nodes represent gene sets and edges represent overlap. Red nodes are positively enriched and blue nodes are negatively enriched. Gene sets with an FDR<0.25 are shown.

938

Figure 2. ID4 marks basal stem cells and decreases in terminally differentiated myoepithelial cells.

a) FACS analysis of EPCAM and CD49f in live lin⁻ mammary cells from adult (10-14 week) Id4floxGFP reporter mice. ID4-GFP expression is indicated in the heatmap scale. Representative plots from 5 experiments shown. Basal cells were binned into 3 groups based on ID4-GFP expression, ID4-bright (red), ID4-intermediate (orange) and ID4-dim (green) and the median fluorescence intensity (MFI) of EPCAM **(b)** and CD49f **(c)** were compared between the 3 gates. MFI expressed as a fold change relative to the ID4-low basal cells. Ordinary one-way ANOVA test was used to test significance. n=5. Error bars represent SEM. ** p<0.01, *** p<0.001, **** p<0.0001.

949 **d)** Representative co-immunofluorescent staining of ID4 and α -SMA in TEB and duct
 950 from a pubertal (6 week) mammary gland. Scale bar = 20 μ m. Comparison of ID4 **(e)**
 951 and α -SMA **(f)** mean fluorescence between individual cap cells (dark blue) and basal
 952 duct cells (light blue). Unpaired two-tailed students t-test. **** p<0.0001.
 953 Correlation between α -SMA and ID4 **(g)** and Hoescht **(h)** mean fluorescence in
 954 individual cap cells (dark blue) and basal duct cells (light blue). R^2 and p values are
 955 displayed. Data is pooled from 9 mice. Approximately 20 TEB cap cells and 20 ductal
 956 basal cells were analysed per mouse. **i)** tSNE plot of 9663 Krt5+/Krt14+ basal cells
 957 from (Bach et al., 2017). 2 mice were analysed per developmental stage. NP =
 958 Nulliparous (8 week), G= Gestation (Day 14.5), L = Lactation (Day 6), I = Involution
 959 (Day 11). Feature plots **(j)** and Violin plots **(k)** displaying expression of *Id4*, *Epcam*
 960 and *Acta2* in single cells in the different developmental stages. **l)** Heatmap displaying
 961 top and bottom 30 differentially expressed genes between the top and bottom 200
 962 *Id4* high and *Id4* low basal cells across all stages. Top 10 GO terms enriched in the
 963 top 50 genes upregulated in *Id4* high **(m)** and low **(n)** basal cells.

964

965 **Figure 3. ID4 inhibits myoepithelial differentiation of organoids.**

966 **a)** Schematic diagram of 3D Matrigel organoid assay. ID4-GFP+ basal cells were FACS
 967 purified from adult (10-11 weeks) ID4-GFP reporter mice. Exon 1 and 2 of *Id4* are
 968 floxed and a GFP reporter cassette introduced. Basal cells are reprogrammed in
 969 culture using ROCK inhibitor Y-27632 and irradiated NIH-3T3 feeder cells. Adenoviral
 970 Cre is used to knock out ID4 as shown by western blotting. Single cells are then
 971 seeded on top of a Matrigel plug and grown for 6 days followed by
 972 immunofluorescent staining and quantification. Organoids grown from conditionally

973 reprogrammed basal cells were treated with control GFP adenovirus (ID4 WT) or
 974 with Cre Adeno virus (ID4 KO) then stained for ID4 **(b)** and α -SMA **(d)**. Scale bar = 10
 975 μ m. Fluorescence was quantified for ID4 **(c)** and α -SMA **(e)** in approximately 10
 976 organoids per experiment. n=4. Unpaired two-tailed students t-test. *** $p<0.001$,
 977 **** $p<0.0001$.

978

979 **Figure 4. ID4 negatively regulates EMT and ECM production in mammary epithelial**
 980 **cells.**

981 **a)** Western blot analysis of ID4 expression in Comma-D β cells treated with non-
 982 targeting (NT) or ID4-targeting siRNA. Representative results from 3 western blots
 983 shown **(i)**. Densitometry quantification of ID4 bands. Band intensity was normalised
 984 to β -Actin and expressed as fold change relative to NT control. N=3. Unpaired two-
 985 tailed students t-test . **** $p<0.0001$. Error bars represent SEM. **(ii). b)** Genes were
 986 ranked based on the limma t-statistic comparing NT and siID4 cells and GSEA was
 987 carried out using Gene Ontology (G) gene sets. The top 10 positively (red) and
 988 negatively (blue) enriched pathways are displayed **c)** GO GSEA results were
 989 visualised using the EnrichmentMap plugin on Cytoscape. Nodes represent gene sets
 990 and edges represent overlap. Red nodes are positively enriched and blue nodes are
 991 negatively enriched in siID4 cells compared to control NT cells. Gene sets with an
 992 FDR<0.1 are shown. **d)** Representative GSEA enrichment plots displaying the profile
 993 of the running Enrichment Score (green) and positions of gene set members on the
 994 rank ordered list. **e)** Collagen fibres were visualised by picrosirius red staining of TEBs
 995 from 6-week-old ID4 WT and KO mice. Scale bar = 50 μ m **(i)**. Total collagen staining
 996 **(ii)** and birefringence signal **(iii)** were quantified from approximately 3 TEBs from

each mammary gland section. N=9 for WT and N=6 for KO mice. Unpaired two-tailed students t-test. * $p < 0.05$. *** $p < 0.001$. Error bars represent SEM.

999

Figure 5. ID4 interacts with E-proteins E2A and HEB.

a) Unsupervised hierarchical clustering heat map of SWATH RIME data from Comma-D β cells. Proteins with significantly higher abundance (p -value <0.05) in the ID4 IPs compared to IgG IPs are displayed. Log2 protein area was used to generate the heatmap. Data from 3 independent experiments shown, each with 2-3 technical replicates. **b)** Co-immunoprecipitation (co-IP) and western blotting of ID4, E2A and HEB and IgG negative controls from uncrosslinked Comma-D β lysates. **c)** Proximity ligation assay (PLA) in Comma-D β cells for ID4 and HEB or corresponding negative control IgG. The cytoskeleton is stained with phalloidin and nucleus with DAPI. Scale bar = 20 μ m. High power insets are shown below. **d)** Quantification of PLA foci from 6 random fields of view for each condition, each with approximately 50 nuclei per image. Ordinary one-way ANOVA test was used to test significance. **** $p < 0.001$. N=3. **e)** Co-immunofluorescent staining of V5 and HEB in TEB (upper) and duct (lower) from ID4-FlagV5 mouse mammary glands. High-power insets feature cells positive for both proteins. Scale bar = 20 μ m. **f)** co-IP and western blotting from ID4-FlagV5 mammary gland protein extracts. ID4 was immunoprecipitated using antibodies raised against ID4 and Flag, and two independent V5 antibodies. ID4, V5 and HEB were detected by western blotting.

1018

Figure 6. HEB directly binds to a subset of ID4 target genes

a) Top 4 enriched transcription factor binding motifs determined using MEME-ChIP

1020

1021 for consensus HEB ChIP-seq peaks in Comma-D β cells. E-values are displayed. **b)**
1022 GREAT pathway analysis of consensus HEB peaks. Top 16 Gene Ontologies (Biological
1023 process, cellular component, molecular function) are displayed. **c)** Venn diagram
1024 showing overlap between genes associated with a HEB peak (GREAT basal plus
1025 extension annotation) and ID4 RNA-seq differentially expressed genes. **d)** Profile
1026 plots of average HEB, H3K4Me3, H3K27Ac, and H3K27Me3 signal intensity at regions
1027 associated with silD4 downregulated (left) and upregulated (right) RNA-seq
1028 differentially expressed genes. **e)** Examples of HEB and histone mark peaks occurring
1029 upstream of silD4 upregulated genes *Sparc*, *Col1a1* and *Cnn2* from the Integrative
1030 Genomics Viewer (IGV). Bars beneath peaks represent consensus MACS call
1031 (FDR<0.05) in at least 2 of 4 biological replicates. Input was used as a negative
1032 control. Purple boxes highlight HEB binding regions. Refseq genes shown in blue.
1033 Data scales for each track are indicated. **f)** Volcano plot of Differential Binding
1034 Analysis. Analysis using EdgeR of HEB binding in silD4 verses NT of 3 biological
1035 replicates. Regions with an FDR less than 0.05 are indicated in pink. **g)** Profile plots of
1036 average HEB signal intensity in NT control (pink) and silD4 (purple) conditions at
1037 regions associated with RNA-seq silD4 downregulated (left), upregulated (middle),
1038 and unchanged (right) genes. **h)** Model of ID4 and HEB action in mammary epithelial
1039 cells. Left: when ID4 is expressed it interacts with HEB, antagonising its
1040 transcriptional activity. Right: when ID4 is depleted, HEB dimerises and binds to E-
1041 box motifs in the promoters and enhancers of developmental genes involved in
1042 contraction and ECM. Below are hypothetical ChIP signals for H3K27Ac (blue),
1043 H3K4Me3 (green), and HEB (pink).

1044 **Supplementary Figure legends**

1045

1046 **Figure S1 related to Figure 1.**

1047 **a)** Proportions of basal, luminal progenitor (LP), and mature luminal (ML)
 1048 subpopulations between ID4 WT and KO mammary epithelial cells. Unpaired two-
 1049 tailed students t-test. ns = not significant. Error bars represent SEM. N=4.
 1050 Representative GSEA enrichment plots displaying the profile of the running
 1051 Enrichment Score (green) and positions of gene set members on the rank ordered
 1052 list for pathways related to growth **(b)** and myoepithelial function **(c)**.

1053

1054 **Figure S2 related to Figure 2.**

1055 **a)** Co-immunofluorescent staining of ID4 and α -SMA in a section containing both a
 1056 TEB and a duct from a pubertal (6 week) mammary gland. Scale bar = 20 μ m. **b)**
 1057 Mouse mammary glands at different developmental stages were stained by IHC for
 1058 ID4. Scale bar = 100 μ m. Representative images from 3 animals per stage shown.
 1059 High power insets are displayed below images. **c)** Differentiation trajectory of basal
 1060 cells from (Bach et al., 2017) coloured by developmental stage (upper) and
 1061 pseudotime (lower). Low values (dark blue) represent undifferentiated cells. NP =
 1062 Nulliparous (8 week), G= Gestation (Day 14.5), L = Lactation (Day 6), I = Involution
 1063 (Day 11). **d)** Expression of *Id4*, *Cnn1*, *Myh11*, *Acta2*, *Mylk* and *Oxtr* over pseudotime.

1064

1065 **Figure S3 related to Figure 3.**

1066 **a)** Phase contrast images of passage 11 conditionally reprogrammed cells grown in
 1067 the presence and absence of ROCK inhibitor Y-27632, irradiated NIH-3T3 feeder cells,

1068 and Matrigel (MG) coated tissue culture flasks. Scale bar = 100 μ m. **b)** ID4 western
 1069 blot in cell lysates collected from cells shown in panel (a). ID4 from basal cells runs at
 1070 a higher molecular weight to ID4 expressed by NIH-3T3 cells. **c)** Co-
 1071 immunofluorescent staining of primary basal cells with basal markers P63 and
 1072 KRT14. Scale bar = 20 μ m. **d)** Basal cells grown in 3D as organoids on top a plug of
 1073 Matrigel. Confocal image of an organoid stained for luminal marker KRT8 and basal
 1074 marker KRT14. Scale bar = 10 μ m.

1075

1076 **Figure S4 related to Figure 4.**

1077 **a)** Genes were ranked based on the limma t-statistic comparing NT and siID4 cells
 1078 and GSEA was carried out using the Hallmark gene sets. The top 10 positively (red)
 1079 and negatively (blue) enriched pathways are displayed. **b)** Representative Hallmark
 1080 GSEA enrichment plots displaying the profile of the running Enrichment Score
 1081 (green) and positions of gene set members on the rank ordered list.

1082

1083 **Figure S5 related to Figure 5.**

1084 Co-immunoprecipitation and western blotting of ID4, E2A and HEB from human
 1085 normal-like mammary epithelial cell lines PMC42 **(a)** and MCF10A cells
 1086 overexpressing Flag-tagged ID4 protein **(b)**. **c)** Unsupervised hierarchical clustering
 1087 heat map of SWATH RIME data from Comma-D β cells. Proteins with significantly
 1088 higher abundance (p-value<0.05) in the HEB IPs compared to IgG IPs are displayed.
 1089 Log2 protein area was used to generate the heatmap.

1090

1091 **Figure S6 related to Figure 6.**

1092 **a)** Genomic distribution of HEB consensus peaks. TSS = transcriptional start site. TTS
1093 = transcriptional termination site. **b)** Heatmaps of HEB, H3K4Me3, H3K27Ac, and
1094 H3K27Me3 ChIP-seq signal at HEB-bound regions. **c)** Examples of HEB and histone
1095 mark peaks occurring upstream of ID4 repressed genes *Col1a2*, *Col3a1*, and *Col5a1*
1096 (multiple HEB peaks) from the Integrative Genomics Viewer (IGV). Bars beneath
1097 peaks represent consensus MACS call (FDR<0.05) in at least 2 of 4 biological
1098 replicates. Input was used as a negative control. Purple boxes highlight HEB binding
1099 regions. Refseq genes shown in blue. Data scales for each track are indicated. **d)**
1100 Western blot analysis of ID4 and HEB expression in Comma-D β cells treated with NT
1101 or ID4-targeting siRNA. Densitometry quantification of ID4 and HEB bands. Band
1102 intensity was normalised to β -Actin and expressed as fold change relative to NT
1103 control. N=3. Unpaired two-tailed students t-test. **** p<0.0001. ns = not significant
1104 Error bars represent SEM. **e)** Top 4 enriched transcription factor binding motifs
1105 determined using MEME-ChIP for consensus HEB ChIP-seq peaks in Comma-D β cells
1106 treated with NT or ID4-targeting siRNA. E-values are displayed. **f)** Example of HEB
1107 peaks in control NT cells and siID4 cells upstream of *Col5a1* from the Integrative
1108 Genomics Viewer (IGV). Green boxes highlight HEB binding regions. Individual
1109 replicates and merged tracks are displayed. Refseq genes shown in blue. Data scales
1110 indicated. **g)** GREAT pathway analysis of 263 peaks increased in siID4 compared to
1111 NT from Fig. 6f. Top 16 Gene Ontologies (Biological process, cellular component,
1112 molecular function) are displayed.

1113 Supplementary Tables

- 1114 **Table S1.** All differentially expressed genes between ID4 wild type and knockout
 1115 sorted mammary populations (Basal, luminal progenitor and mature luminal).
 1116 **Table S2.** Top 50 differentially expressed genes between *Id4*-high and *Id4*-lo basal
 1117 cells from Bach et al. single cell RNA-seq analysis.
 1118 **Table S3.** Top 50 differentially expressed genes between siID4 and Non-targeting
 1119 control Comma-Dβ cells.
 1120 **Table S4.** Proteins identified in ID4 and HEB RIME experiments.
 1121 **Table S5.** Genomic regions of Differentially bound HEB ChIP-Seq peaks between siID4
 1122 and NT control Comma-Dβ cells.

1123 References

- 1124 Andrews S (2010) FastQC: A quality control tool for high throughput sequence data. .
 1125 In
 1126 Asselin-Labat ML, Sutherland KD, Barker H, Thomas R, Shackleton M, Forrest NC,
 1127 Hartley L, Robb L, Grosveld FG, van der Wees J, Lindeman GJ, Visvader JE (2007)
 1128 Gata-3 is an essential regulator of mammary-gland morphogenesis and luminal-cell
 1129 differentiation. *Nature cell biology* 9: 201-9
 1130 Bach K, Pensa S, Grzelak M, Hadfield J, Adams DJ, Marioni JC, Khaled WT (2017)
 1131 Differentiation dynamics of mammary epithelial cells revealed by single-cell RNA
 1132 sequencing. *Nature communications* 8: 2128
 1133 Baker LA, Holliday H, Swarbrick A (2016) ID4 Controls Mammary Lineage
 1134 Commitment and Inhibits BRCA1 in Basal Breast Cancer. *Endocrine-related cancer*
 1135 23: R381-92
 1136 Barsky SH, Karlin NJ (2005) Myoepithelial cells: autocrine and paracrine suppressors
 1137 of breast cancer progression. *Journal of mammary gland biology and neoplasia* 10:
 1138 249-260
 1139 Bayly R, Chuen L, Currie RA, Hyndman BD, Casselman R, Blobel GA, LeBrun DP (2004)
 1140 E2A-PBX1 interacts directly with the KIX domain of CBP/p300 in the induction of
 1141 proliferation in primary hematopoietic cells. *The Journal of biological chemistry* 279:
 1142 55362-71
 1143 Benezra R, Davis RL, Lockshon D, Turner DL, Weintraub H (1990) The protein Id: a
 1144 negative regulator of helix-loop-helix DNA binding proteins. *Cell* 61: 49-59

Best SA, Hutt KJ, Fu NY, Vaillant F, Liew SH, Hartley L, Scott CL, Lindeman GJ, Visvader JE (2014) Dual roles for Id4 in the regulation of estrogen signaling in the mammary gland and ovary. *Development (Cambridge, England)* 141: 3159-64

Bouras T, Pal B, Vaillant F, Harburg G, Asselin-Labat ML, Oakes SR, Lindeman GJ, Visvader JE (2008) Notch signaling regulates mammary stem cell function and luminal cell-fate commitment. *Cell stem cell* 3: 429-41

Braunstein M, Anderson MK (2012) HEB in the spotlight: Transcriptional regulation of T-cell specification, commitment, and developmental plasticity. *Clinical & developmental immunology* 2012: 678705

Buchwalter G, Hickey MM, Cromer A, Selfors LM, Gunawardane RN, Frishman J, Jeselsohn R, Lim E, Chi D, Fu X, Schiff R, Brown M, Brugge JS (2013) PDEF promotes luminal differentiation and acts as a survival factor for ER-positive breast cancer cells. *Cancer cell* 23: 753-67

Carr JR, Kiefer MM, Park HJ, Li J, Wang Z, Fontanarosa J, DeWaal D, Kopanja D, Benevolenskaya EV, Guzman G, Raychaudhuri P (2012) FoxM1 regulates mammary luminal cell fate. *Cell reports* 1: 715-29

Chakrabarti R, Wei Y, Romano RA, DeCoste C, Kang Y, Sinha S (2012) Elf5 regulates mammary gland stem/progenitor cell fate by influencing notch signaling. *Stem cells (Dayton, Ohio)* 30: 1496-508

Conway K, Pin C, Kiernan JA, Merrifield P (2004) The E protein HEB is preferentially expressed in developing muscle. *Differentiation; research in biological diversity* 72: 327-340

Dalal SJ, Estep JS, Valentin-Bon IE, Jerse AE (2001) Standardization of the Whitten Effect to induce susceptibility to *Neisseria gonorrhoeae* in female mice. *Contemporary topics in laboratory animal science / American Association for Laboratory Animal Science* 40: 13-7

Danielson KG, Oborn CJ, Durban EM, Butel JS, Medina D (1984) Epithelial mouse mammary cell line exhibiting normal morphogenesis in vivo and functional differentiation in vitro. *Proceedings of the National Academy of Sciences of the United States of America* 81: 3756-60

Davis FM, Lloyd-Lewis B, Harris OB, Kozar S, Winton DJ, Muresan L, Watson CJ (2016) Single-cell lineage tracing in the mammary gland reveals stochastic clonal dispersion of stem/progenitor cell progeny. *Nature communications* 7: 13053

Deugnier MA, Faraldo MM, Teuliere J, Thiery JP, Medina D, Glukhova MA (2006) Isolation of mouse mammary epithelial progenitor cells with basal characteristics from the Comma-Dbeta cell line. *Developmental biology* 293: 414-25

Deugnier MA, Moiseyeva EP, Thiery JP, Glukhova M (1995) Myoepithelial cell differentiation in the developing mammary gland: progressive acquisition of smooth muscle phenotype. *Developmental dynamics : an official publication of the American Association of Anatomists* 204: 107-17

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR (2013) STAR: ultrafast universal RNA-seq aligner. *Bioinformatics (Oxford, England)* 29: 15-21

Dong J, Huang S, Caikovski M, Ji S, McGrath A, Custorio MG, Creighton CJ, Maliakkal P, Bogoslovskaya E, Du Z, Zhang X, Lewis MT, Sablitzky F, Brisken C, Li Y (2011) ID4 regulates mammary gland development by suppressing p38MAPK activity. *Development (Cambridge, England)* 138: 5247-56

1192 Ewald AJ, Brenot A, Duong M, Chan BS, Werb Z (2008) Collective epithelial migration
1193 and cell rearrangements drive mammary branching morphogenesis. *Developmental*
1194 *cell* 14: 570-581

1195 Feinberg TY, Zheng H, Liu R, Wicha MS, Yu SM, Weiss SJ (2018) Divergent Matrix-
1196 Remodeling Strategies Distinguish Developmental from Neoplastic Mammary
1197 Epithelial Cell Invasion Programs. *Developmental cell* 47: 145-160.e6

1198 Finak G, McDavid A, Yajima M, Deng J, Gersuk V, Shalek AK, Slichter CK, Miller HW,
1199 McElrath MJ, Prlic M, Linsley PS, Gottardo R (2015) MAST: a flexible statistical
1200 framework for assessing transcriptional changes and characterizing heterogeneity in
1201 single-cell RNA sequencing data. *Genome biology* 16: 278

1202 Gudjonsson T, Adriance MC, Sternlicht MD, Petersen OW, Bissell MJ (2005)
1203 Myoepithelial cells: their origin and function in breast morphogenesis and neoplasia.
1204 *Journal of mammary gland biology and neoplasia* 10: 261-72

1205 Gudjonsson T, Rønnov-Jessen L, Villadsen R, Rank F, Bissell MJ, Petersen OW (2002)
1206 Normal and tumor-derived myoepithelial cells differ in their ability to interact with
1207 luminal breast epithelial cells for polarity and basement membrane deposition.
1208 *Journal of cell science* 115: 39-50

1209 Guo W, Keckesova Z, Donaher JL, Shibue T, Tischler V, Reinhardt F, Itzkovitz S, Noske
1210 A, Zurrer-Hardi U, Bell G, Tam WL, Mani SA, van Oudenaarden A, Weinberg RA
1211 (2012) Slug and Sox9 cooperatively determine the mammary stem cell state. *Cell*
1212 148: 1015-28

1213 Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, Cheng JX, Murre C, Singh H,
1214 Glass CK (2010) Simple combinations of lineage-determining transcription factors
1215 prime cis-regulatory elements required for macrophage and B cell identities.
1216 *Molecular cell* 38: 576-589

1217 Huang Q, Yang L, Luo J, Guo L, Wang Z, Yang X, Jin W, Fang Y, Ye J, Shan B, Zhang Y
1218 (2015) SWATH enables precise label-free quantification on proteome scale.
1219 *Proteomics* 15: 1215-23

1220 Idoux-Gillet Y, Nassour M, Lakis E, Bonini F, Theillet C, Du Manoir S, Savagner P
1221 (2018) Slug/Pcad pathway controls epithelial cell dynamics in mammary gland and
1222 breast carcinoma. *Oncogene* 37: 578-588

1223 Ingthorsson S, Hilmarsdottir B, Krickler J, Magnusson MK, Gudjonsson T (2015)
1224 Context-Dependent Function of Myoepithelial Cells in Breast Morphogenesis and
1225 Neoplasia. *Current molecular biology reports* 1: 168-174

1226 Junankar S, Baker LA, Roden DL, Nair R, Elsworth B, Gallego-Ortega D, Lacaze P, Cazet
1227 A, Nikolic I, Teo WS, Yang J, McFarland A, Harvey K, Naylor MJ, Lakhani SR, Simpson
1228 PT, Raghavendra A, Saunus J, Madore J, Kaplan W et al. (2015) ID4 controls
1229 mammary stem cells and marks breast cancers with a stem cell-like phenotype.
1230 *Nature communications* 6: 6548

1231 Khoury A, Achinger-Kawecka J, Bert SA, Smith GC, French HJ, Luu P-L, Peters TJ, Du Q,
1232 Parry AJ, Valdes-Mora F, Taberlay PC, Stirzaker C, Statham AL, Clark SJ (2020)
1233 Constitutively bound CTCF sites maintain 3D chromatin architecture and long-range
1234 epigenetically regulated domains. *Nature communications* 11: 54-54

1235 Kiemer AK, Takeuchi K, Quinlan MP (2001) Identification of genes involved in
1236 epithelial-mesenchymal transition and tumor progression. *Oncogene* 20: 6679-6688

1237 Kouros-Mehr H, Slorach EM, Sternlicht MD, Werb Z (2006) GATA-3 maintains the
1238 differentiation of the luminal cell fate in the mammary gland. *Cell* 127: 1041-55

1239 Kumar MS, Hendrix JA, Johnson AD, Owens GK (2003) Smooth muscle alpha-actin
1240 gene requires two E-boxes for proper expression in vivo and is a target of class I
1241 basic helix-loop-helix proteins. *Circulation research* 92: 840-7
1242 Law AMK, Yin JXM, Castillo L, Young AJ, Piggan C, Rogers S, Caldon CE, Burgess A,
1243 Millar EKA, O'Toole SA, Gallego-Ortega D, Ormandy CJ, Oakes SR (2017) Andy's
1244 Algorithms: new automated digital image analysis pipelines for Fiji. *Scientific reports*
1245 7: 15717
1246 Lee K, Gjorevski N, Boghaert E, Radisky DC, Nelson CM (2011) Snail1, Snail2, and E47
1247 promote mammary epithelial branching morphogenesis. *The EMBO journal* 30:
1248 2662-74
1249 Li B, Dewey CN (2011) RSEM: accurate transcript quantification from RNA-Seq data
1250 with or without a reference genome. *BMC bioinformatics* 12: 323
1251 Li H, Durbin R (2009) Fast and accurate short read alignment with Burrows-Wheeler
1252 transform. *Bioinformatics (Oxford, England)* 25: 1754-60
1253 Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G,
1254 Durbin R, Genome Project Data Processing S (2009) The Sequence Alignment/Map
1255 format and SAMtools. *Bioinformatics (Oxford, England)* 25: 2078-2079
1256 Li S, Chang S, Qi X, Richardson JA, Olson EN (2006) Requirement of a myocardin-
1257 related transcription factor for development of mammary myoepithelial cells.
1258 *Molecular and cellular biology* 26: 5797-808
1259 Li Y, Brauer PM, Singh J, Xhiku S, Yoganathan K, Zúñiga-Pflücker JC, Anderson MK
1260 (2017) Targeted Disruption of TCF12 Reveals HEB as Essential in Human Mesodermal
1261 Specification and Hematopoiesis. *Stem cell reports* 9: 779-795
1262 Lilja AM, Rodilla V, Huyghe M, Hannezo E, Landragin C, Renaud O, Leroy O, Rulands
1263 S, Simons BD, Fre S (2018) Clonal analysis of Notch1-expressing cells reveals the
1264 existence of unipotent stem cells that retain long-term plasticity in the embryonic
1265 mammary gland. *Nature cell biology* 20: 677-687
1266 Lim E, Wu D, Pal B, Bouras T, Asselin-Labat ML, Vaillant F, Yagita H, Lindeman GJ,
1267 Smyth GK, Visvader JE (2010) Transcriptome analyses of mouse and human
1268 mammary cell subpopulations reveal multiple conserved genes and pathways. *Breast*
1269 *cancer research : BCR* 12: R21
1270 Liu J, Eischeid AN, Chen XM (2012) Col1A1 production and apoptotic resistance in
1271 TGF-β1-induced epithelial-to-mesenchymal transition-like phenotype of 603B cells.
1272 *PloS one* 7: e51371
1273 Liu S, Ginestier C, Charafe-Jauffret E, Foco H, Kleer CG, Merajver SD, Dontu G, Wicha
1274 MS (2008) BRCA1 regulates human mammary stem/progenitor cell fate. *Proceedings*
1275 *of the National Academy of Sciences of the United States of America* 105: 1680-5
1276 Liu X, Ory V, Chapman S, Yuan H, Albanese C, Kallakury B, Timofeeva OA, Nealon C,
1277 Dakic A, Simic V, Haddad BR, Rhim JS, Dritschilo A, Riegel A, McBride A, Schlegel R
1278 (2012) ROCK inhibitor and feeder cells induce the conditional reprogramming of
1279 epithelial cells. *The American journal of pathology* 180: 599-607
1280 Lloyd-Lewis B, Davis FM, Harris OB, Hitchcock JR, Watson CJ (2018) Neutral lineage
1281 tracing of proliferative embryonic and adult mammary stem/progenitor cells.
1282 *Development (Cambridge, England)* 145
1283 Lourenço AR, Roukens MG, Seinstra D, Frederiks CL, Pals CE, Vervoort SJ, Margarido
1284 AS, van Rheenen J, Coffey PJ (2020) C/EBPβ is crucial determinant of epithelial

1285 maintenance by preventing epithelial-to-mesenchymal transition. *Nature*
1286 *communications* 11: 785
1287 Machanick P, Bailey TL (2011) MEME-ChIP: motif analysis of large DNA datasets.
1288 *Bioinformatics (Oxford, England)* 27: 1696-1697
1289 Macias H, Hinck L (2012) Mammary gland development. *Wiley interdisciplinary*
1290 *reviews Developmental biology* 1: 533-57
1291 Martin M (2011) Cutadapt removes adapter sequences from high-throughput
1292 sequencing reads. *2011* 17: 3
1293 Massari ME, Murre C (2000) Helix-loop-helix proteins: regulators of transcription in
1294 eucaryotic organisms. *Molecular and cellular biology* 20: 429-40
1295 McCarthy DJ, Chen Y, Smyth GK (2012) Differential expression analysis of multifactor
1296 RNA-Seq experiments with respect to biological variation. *Nucleic acids research* 40:
1297 4288-97
1298 McCloy RA, Rogers S, Caldon CE, Lorca T, Castro A, Burgess A (2014) Partial inhibition
1299 of Cdk1 in G 2 phase overrides the SAC and decouples mitotic events. *Cell cycle*
1300 *(Georgetown, Tex)* 13: 1400-12
1301 McLean CY, Bristor D, Hiller M, Clarke SL, Schaar BT, Lowe CB, Wenger AM, Bejerano
1302 G (2010) GREAT improves functional interpretation of cis-regulatory regions. *Nature*
1303 *biotechnology* 28: 495-501
1304 Merico D, Isserlin R, Stueker O, Emili A, Bader GD (2010) Enrichment map: a
1305 network-based method for gene-set enrichment visualization and interpretation.
1306 *PloS one* 5: e13984
1307 Mesman S, Smidt MP (2017) Tcf12 Is Involved in Early Cell-Fate Determination and
1308 Subset Specification of Midbrain Dopamine Neurons. *Frontiers in Molecular*
1309 *Neuroscience* 10
1310 Miano JM, Long X, Fujiwara K (2007) Serum response factor: master regulator of the
1311 actin cytoskeleton and contractile apparatus. *American journal of physiology Cell*
1312 *physiology* 292: C70-C81
1313 Mills AA, Zheng B, Wang XJ, Vogel H, Roop DR, Bradley A (1999) p63 is a p53
1314 homologue required for limb and epidermal morphogenesis. *Nature* 398: 708-13
1315 Mohammed H, D'Santos C, Serandour AA, Ali HR, Brown GD, Atkins A, Rueda OM,
1316 Holmes KA, Theodorou V, Robinson JL, Zwart W, Saadi A, Ross-Innes CS, Chin SF,
1317 Menon S, Stingl J, Palmieri C, Caldas C, Carroll JS (2013) Endogenous purification
1318 reveals GREB1 as a key estrogen receptor regulatory factor. *Cell reports* 3: 342-9
1319 Muschler J, Streuli CH (2010) Cell-matrix interactions in mammary gland
1320 development and breast cancer. *Cold Spring Harbor perspectives in biology* 2:
1321 a003202
1322 Nielsen TO, West RB, Linn SC, Alter O, Knowling MA, O'Connell JX, Zhu S, Fero M,
1323 Sherlock G, Pollack JR, Brown PO, Botstein D, van de Rijn M (2002) Molecular
1324 characterisation of soft tissue tumours: a gene expression study. *Lancet (London,*
1325 *England)* 359: 1301-7
1326 Oakes SR, Naylor MJ, Asselin-Labat ML, Blazek KD, Gardiner-Garden M, Hilton HN,
1327 Kazlauskas M, Pritchard MA, Chodosh LA, Pfeffer PL, Lindeman GJ, Visvader JE,
1328 Ormandy CJ (2008) The Ets transcription factor Elf5 specifies mammary alveolar cell
1329 fate. *Genes & development* 22: 581-6
1330 Paine IS, Lewis MT (2017) The Terminal End Bud: the Little Engine that Could. *Journal*
1331 *of mammary gland biology and neoplasia* 22: 93-108

1332 Pervolarakis N, Nguyen Q, Gutierrez G, Sun P, Jhutti D, Zheng GX, Nemec CM, Dai X,
1333 Watanabe K, Kessenbrock K (2019) Integrated single-cell transcriptomics and
1334 chromatin accessibility analysis reveals novel regulators of mammary epithelial cell
1335 identity. *bioRxiv*: 740746

1336 Prater MD, Petit V, Alasdair Russell I, Giraddi RR, Shehata M, Menon S, Schulte R,
1337 Kalajic I, Rath N, Olson MF, Metzger D, Faraldo MM, Deugnier MA, Glukhova MA,
1338 Stingl J (2014) Mammary stem cells have myoepithelial cell properties. *Nature cell
1339 biology* 16: 942-50, 1-7

1340 Qiu X, Mao Q, Tang Y, Wang L, Chawla R, Pliner HA, Trapnell C (2017) Reversed graph
1341 embedding resolves complex single-cell trajectories. *Nature methods* 14: 979-982

1342 Rios AC, Fu NY, Lindeman GJ, Visvader JE (2014) In situ identification of bipotent
1343 stem cells in the mammary gland. *Nature* 506: 322-7

1344 Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK (2015) limma powers
1345 differential expression analyses for RNA-sequencing and microarray studies. *Nucleic
1346 acids research* 43: e47

1347 Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, Mesirov
1348 JP (2011) Integrative genomics viewer. *Nature biotechnology* 29: 24-26

1349 Robinson MD, McCarthy DJ, Smyth GK (2010) edgeR: a Bioconductor package for
1350 differential expression analysis of digital gene expression data. *Bioinformatics
1351 (Oxford, England)* 26: 139-40

1352 Satija R, Farrell JA, Gennert D, Schier AF, Regev A (2015) Spatial reconstruction of
1353 single-cell gene expression data. *Nature biotechnology* 33: 495-502

1354 Scheele CL, Hannezo E, Muraro MJ, Zomer A, Langedijk NS, van Oudenaarden A,
1355 Simons BD, van Rheenen J (2017) Identity and dynamics of mammary stem cells
1356 during branching morphogenesis. *Nature* 542: 313-317

1357 Shackleton M, Vaillant F, Simpson KJ, Stingl J, Smyth GK, Asselin-Labat ML, Wu L,
1358 Lindeman GJ, Visvader JE (2006) Generation of a functional mammary gland from a
1359 single stem cell. *Nature* 439: 84-8

1360 Silberstein GB, Strickland P, Coleman S, Daniel CW (1990) Epithelium-dependent
1361 extracellular matrix synthesis in transforming growth factor-beta 1-growth-inhibited
1362 mouse mammary gland. *The Journal of cell biology* 110: 2209-19

1363 Stark RB, G. D. (2011) DiffBind: differential binding analysis of ChIP-seq peak data. In
1364 Bioconductor:

1365 Stempor P, Ahringer J (2016) SeqPlots - Interactive software for exploratory data
1366 analyses, pattern discovery and visualization in genomics [version 1; peer review: 2
1367 approved, 1 approved with reservations]. *Wellcome Open Research* 1

1368 Sternlicht MD (2006) Key stages in mammary gland development: the cues that
1369 regulate ductal branching morphogenesis. *Breast cancer research : BCR* 8: 201

1370 Sternlicht MD, Kedeshian P, Shao ZM, Safarians S, Barsky SH (1997) The human
1371 myoepithelial cell is a natural tumor suppressor. *Clinical cancer research : an official
1372 journal of the American Association for Cancer Research* 3: 1949-58

1373 Stingl J, Eirew P, Ricketson I, Shackleton M, Vaillant F, Choi D, Li HI, Eaves CJ (2006)
1374 Purification and unique properties of mammary epithelial stem cells. *Nature* 439:
1375 993-7

1376 Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA,
1377 Paulovich A, Pomeroy SL, Golub TR, Lander ES, Mesirov JP (2005) Gene set

1378 enrichment analysis: A knowledge-based approach for interpreting genome-wide
1379 expression profiles. *Proceedings of the National Academy of Sciences* 102: 15545
1380 Sun Y, Boyd K, Xu W, Ma J, Jackson CW, Fu A, Shillingford JM, Robinson GW,
1381 Hennighausen L, Hitzler JK, Ma Z, Morris SW (2006) Acute myeloid leukemia-
1382 associated Mkl1 (Mrtf-a) is a key regulator of mammary gland function. *Molecular*
1383 *and cellular biology* 26: 5809-26
1384 Trapnell C, Cacchiarelli D, Grimsby J, Pokharel P, Li S, Morse M, Lennon NJ, Livak KJ,
1385 Mikkelsen TS, Rinn JL (2014) The dynamics and regulators of cell fate decisions are
1386 revealed by pseudotemporal ordering of single cells. *Nature biotechnology* 32: 381-
1387 386
1388 van Amerongen R, Bowman AN, Nusse R (2012) Developmental stage and time
1389 dictate the fate of Wnt/beta-catenin-responsive stem cells in the mammary gland.
1390 *Cell stem cell* 11: 387-400
1391 Van Keymeulen A, Rocha AS, Ousset M, Beck B, Bouvencourt G, Rock J, Sharma N,
1392 Dekoninck S, Blanpain C (2011) Distinct stem cells contribute to mammary gland
1393 development and maintenance. *Nature* 479: 189-93
1394 Vennin C, Chin VT, Warren SC, Lucas MC, Herrmann D, Magenau A, Melenec P,
1395 Walters SN, Del Monte-Nieto G, Conway JRW, Nobis M, Allam AH, McCloy RA, Currey
1396 N, Pinese M, Boulghourjian A, Zaratzian A, Adam AAS, Heu C, Nagrial AM et al. (2017)
1397 Transient tissue priming via ROCK inhibition uncouples pancreatic cancer
1398 progression, sensitivity to chemotherapy, and metastasis. *Science translational*
1399 *medicine* 9: eaai8504
1400 Vennin C, Méléne P, Rouet R, Nobis M, Cazet AS, Murphy KJ, Herrmann D, Reed DA,
1401 Lucas MC, Warren SC, Elgundi Z, Pinese M, Kalna G, Roden D, Samuel M, Zaratzian A,
1402 Grey ST, Da Silva A, Leung W, Australian Pancreatic Genome I et al. (2019) CAF
1403 hierarchy driven by pancreatic cancer cell p53-status creates a pro-metastatic and
1404 chemoresistant environment via perlecan. *Nature communications* 10: 3637-3637
1405 Visvader JE, Clevers H (2016) Tissue-specific designs of stem cell hierarchies. *Nature*
1406 *cell biology* 18: 349-55
1407 Wang D, Cai C, Dong X, Yu QC, Zhang XO, Yang L, Zeng YA (2015) Identification of
1408 multipotent mammary stem cells by protein C receptor expression. *Nature* 517: 81-4
1409 Wang LH, Baker NE (2015) E Proteins and ID Proteins: Helix-Loop-Helix Partners in
1410 Development and Disease. *Developmental cell* 35: 269-80
1411 Watanabe K, Villarreal-Ponce A, Sun P, Salmans ML, Fallahi M, Andersen B, Dai X
1412 (2014) Mammary morphogenesis and regeneration require the inhibition of EMT at
1413 terminal end buds by Ovol2 transcriptional repressor. *Developmental cell* 29: 59-74
1414 Williams JM, Daniel CW (1983) Mammary ductal elongation: differentiation of
1415 myoepithelium and basal lamina during branching morphogenesis. *Developmental*
1416 *biology* 97: 274-90
1417 Wojnarowicz PM, Lima E Silva R, Ohnaka M, Lee SB, Chin Y, Kulukian A, Chang S-H,
1418 Desai B, Garcia Escolano M, Shah R, Garcia-Cao M, Xu S, Kadam R, Goldgur Y, Miller
1419 MA, Ouerfelli O, Yang G, Arakawa T, Albanese SK, Garland WA et al. (2019) A Small-
1420 Molecule Pan-Id Antagonist Inhibits Pathologic Ocular Neovascularization. *Cell*
1421 *reports* 29: 62-75.e7
1422 Wuidart A, Ousset M, Rulands S, Simons BD, Van Keymeulen A, Blanpain C (2016)
1423 Quantitative lineage tracing strategies to resolve multipotency in tissue-specific stem
1424 cells. *Genes & development* 30: 1261-77

1425 Wuidart A, Sifrim A, Fioramonti M, Matsumura S, Brisebarre A, Brown D, Centonze A,
1426 Dannau A, Dubois C, Van Keymeulen A, Voet T, Blanpain C (2018) Early lineage
1427 segregation of multipotent embryonic mammary gland progenitors. *Nature cell*
1428 *biology* 20: 666-676
1429 Yamaji D, Na R, Feuermann Y, Pechhold S, Chen W, Robinson GW, Hennighausen L
1430 (2009) Development of mammary luminal progenitor cells is controlled by the
1431 transcription factor STAT5A. *Genes & development* 23: 2382-7
1432 Yang A, Schweitzer R, Sun D, Kaghad M, Walker N, Bronson RT, Tabin C, Sharpe A,
1433 Caput D, Crum C, McKeon F (1999) p63 is essential for regenerative proliferation in
1434 limb, craniofacial and epithelial development. *Nature* 398: 714-8
1435 Yoon SJ, Foley JW, Baker JC (2015) HEB associates with PRC2 and SMAD2/3 to
1436 regulate developmental fates. *Nature communications* 6: 6546
1437 Yun K, Mantani A, Garel S, Rubenstein J, Israel MA (2004) Id4 regulates neural
1438 progenitor proliferation and differentiation in vivo. *Development (Cambridge,*
1439 *England)* 131: 5441-8
1440 Zhang J, Kalkum M, Yamamura S, Chait BT, Roeder RG (2004) E protein silencing by
1441 the leukemogenic AML1-ETO fusion protein. *Science (New York, NY)* 305: 1286-9
1442 Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, Nusbaum C, Myers
1443 RM, Brown M, Li W, Liu XS (2008) Model-based analysis of ChIP-Seq (MACS).
1444 *Genome biology* 9: R137-R137
1445

Figure 1. Loss of ID4 results in upregulation of myoepithelial genes and downregulation of cell growth genes in sorted mammary basal cells.

a) Live lineage negative CD24⁺CD29^{hi}CD61⁺ basal, CD24⁺CD29^{lo}CD61⁺ luminal progenitor and CD24⁺CD29^{lo}CD61⁻ mature luminal cells were isolated by FACS from adult (10-12 weeks) ID4 wild type (WT) and knockout (KO) mice at estrus for RNA-seq. Representative FACS plots shown from 4 experiments. b) Number of significantly (FDR<0.05) differentially expressed genes (DEGs) upregulated (red) or downregulated (blue) in ID4 KO epithelial subpopulations compared to ID4 WT. LP = luminal progenitor, ML = mature luminal. c) Heat map displaying the 237 significant differentially expressed genes between ID4 WT and ID4 KO basal cells. d) Genes were ranked based on the limma t-statistic comparing ID4 WT and KO basal cells and GSEA was carried out using the C2all, C3TF, C5 and Hallmark gene sets. The top 10 positive and negatively enriched pathways with an FDR<0.1 are displayed. Only 3 pathways were negatively enriched with an FDR<0.1. e) GSEA results were visualised using the EnrichmentMap plugin on Cytoscape. Nodes represent gene sets and edges represent overlap. Red nodes are positively enriched and blue nodes are negatively enriched. Gene sets with an FDR<0.25 are shown.

Figure 2. ID4 marks basal stem cells and decreases in terminally differentiated myoepithelial cells.

a

b

c

d

e

Figure 3. ID4 inhibits myoepithelial differentiation of organoids.

a) Schematic diagram of 3D Matrigel organoid assay. ID4-GFP⁺ basal cells were FACS purified from adult (10-11 weeks) ID4-GFP reporter mice. Exon 1 and 2 of *Id4* are floxed and a GFP reporter cassette introduced. Basal cells are reprogrammed in culture using ROCK inhibitor Y-27632 and irradiated NIH-3T3 feeder cells. Adenoviral Cre is used to knock out ID4 as shown by western blotting. Single cells are then seeded on top of a Matrigel plug and grown for 6 days followed by immunofluorescent staining and quantification. Organoids grown from conditionally reprogrammed basal cells were treated with control GFP adenovirus (ID4 WT) or with Cre Adeno virus (ID4 KO) then stained for ID4 (b) and α-SMA (d). Scale bar = 10 μm. Fluorescence was quantified for ID4 (c) and α-SMA (e) in approximately 10 organoids per experiment. n=4. Unpaired two-tailed students t-test. *** p<0.001, **** p<0.0001.

Figure 4. ID4 negatively regulates EMT and ECM production in mammary epithelial cells.

a) Western blot analysis of ID4 expression in Comma-D β cells treated with non-targeting (NT) or ID4-targeting siRNA. Representative results from 3 western blots shown (i). Densitometry quantification of ID4 bands. Band intensity was normalised to β -Actin and expressed as fold change relative to NT control. N=3. Unpaired two-tailed students t-test. **** $p < 0.0001$. Error bars represent SEM. (ii). b) Genes were ranked based on the limma t-statistic comparing NT and siID4 cells and GSEA was carried out using Gene Ontology (G) gene sets. The top 10 positively (red) and negatively (blue) enriched pathways are displayed c) GO GSEA results were visualised using the EnrichmentMap plugin on Cytoscape. Nodes represent gene sets and edges represent overlap. Red nodes are positively enriched and blue nodes are negatively enriched in siID4 cells compared to control NT cells. Gene sets with an $FDR < 0.1$ are shown. d) Representative GSEA enrichment plots displaying the profile of the running Enrichment Score (green) and positions of gene set members on the rank ordered list. e) Collagen fibres were visualised by picrosirius red staining of TEBs from 6-week-old ID4 WT and KO mice. Scale bar = 50 μ m (i). Total collagen staining (ii) and birefringence signal (iii) were quantified from approximately 3 TEBs from each mammary gland section. N=9 for WT and N=6 for KO mice. Unpaired two-tailed students t-test. * $p < 0.05$. *** $p < 0.001$. Error bars represent SEM.

Figure 5. ID4 interacts with E-proteins E2A and HEB.

a) Unsupervised hierarchical clustering heat map of SWATH RIME data from Comma-D β cells. Proteins with significantly higher abundance (p-value < 0.05) in the ID4 IPs compared to IgG IPs are displayed. Log2 protein area was used to generate the heatmap. Data from 3 independent experiments shown, each with 2-3 technical replicates. b) Co-immunoprecipitation (co-IP) and western blotting of ID4, E2A and HEB and IgG negative controls from uncrosslinked Comma-D β lysates. c) Proximity ligation assay (PLA) in Comma-D β cells for ID4 and HEB or corresponding negative control IgG. The cytoskeleton is stained with phalloidin and nucleus with DAPI. Scale bar = 20 μ m. High power insets are shown below. d) Quantification of PLA foci from 6 random fields of view for each condition, each with approximately 50 nuclei per image. Ordinary one-way ANOVA test was used to test significance. **** p < 0.001. N=3. e) Co-immunofluorescent staining of V5 and HEB in TEB (upper) and duct (lower) from ID4-FlagV5 mouse mammary glands. High-power insets feature cells positive for both proteins. Scale bar = 20 μ m. f) co-IP and western blotting from ID4-FlagV5 mammary gland protein extracts. ID4 was immunoprecipitated using antibodies raised against ID4 and Flag, and two independent V5 antibodies. ID4, V5 and HEB were detected by western blotting.

Figure 6. HEB directly binds to a subset of ID4 target genes

a) Top 4 enriched transcription factor binding motifs determined using MEME-ChIP for consensus HEB ChIP-seq peaks in Comma-Dβ cells. E-values are displayed. b) GREAT pathway analysis of consensus HEB peaks. Top 16 Gene Ontologies (Biological process, cellular component, molecular function) are displayed. c) Venn diagram showing overlap between genes associated with a HEB peak (GREAT basal plus extension annotation) and ID4 RNA-seq differentially expressed genes. d) Profile plots of average HEB, H3K4Me3, H3K27Ac, and H3K27Me3 signal intensity at regions associated with siID4 downregulated (left) and upregulated (right) RNA-seq differentially expressed genes. e) Examples of HEB and histone mark peaks occurring upstream of siID4 upregulated genes Sparc, Col1a1 and Cnn2 from the Integrative Genomics Viewer (IGV). Bars beneath peaks represent consensus MACS call (FDR<0.05) in at least 2 of 4 biological replicates. Input was used as a negative control. Purple boxes highlight HEB binding regions. Refseq genes shown in blue. Data scales for each track are indicated. f) Volcano plot of Differential Binding Analysis. Analysis using EdgeR of HEB binding in siID4 versus NT of 3 biological replicates. Regions with an FDR less than 0.05 are indicated in pink. g) Profile plots of average HEB signal intensity in NT control (pink) and siID4 (purple) conditions at regions associated with RNA-seq siID4 downregulated (left), upregulated (middle), and unchanged (right) genes. h) Model of ID4 and HEB action in mammary epithelial cells. Left: when ID4 is expressed it interacts with HEB, antagonising its transcriptional activity. Right: when ID4 is depleted, HEB dimerises and binds to E-box motifs in the promoters and enhancers of developmental genes involved in contraction and ECM. Below are hypothetical ChIP signals for H3K27Ac (blue), H3K4Me3 (green), and HEB (pink).