

1 **MARCH8 inhibits viral infection by two different mechanisms**

2

3 Yanzhao Zhang^{1,4}, Takuya Tada^{1,4,5}, Seiya Ozono^{1,2}, Satoshi Kishigami², Hideaki Fujita³, and
4 Kenzo Tokunaga^{1,2*}

5

6 ¹Department of Pathology, National Institute of Infectious Diseases, Tokyo 162-8640, Japan;

7 ²Faculty of Life and Environmental Sciences, University of Yamanashi, Yamanashi, Japan;

8 ³Faculty of Pharmaceutical Sciences, Nagasaki International University, Nagasaki 859-3298,

9 Japan

10

11

12 Correspondence: Kenzo Tokunaga, Department of Pathology, National Institute of Infectious

13 Diseases, Shinjuku-ku, Tokyo 162-8640, Japan, Tel: +81 3 5285 1111; Fax: +81 3 5285 1189; E-

14 mail: tokunaga@nih.go.jp

15

16 ⁴These authors contributed equally to this work.

17 ⁵Present address: Dept. of Microbiology, New York University School of Medicine, New York,

18 New York.

19

20 Total number of words in the abstract: 176

21 Total number of words in the text: 2,558

22 Key words: HIV-1, MARCH8, envelope, antiviral factor, ubiquitination, tyrosine motif

1 **ABSTRACT**

2 Membrane-associated RING-CH (MARCH) family members are cellular transmembrane proteins
3 that downregulate several host membrane proteins. We have recently reported that one family
4 member, MARCH8, inhibits infection with both HIV-1 and vesicular stomatitis virus G-
5 glycoprotein (VSV-G)-pseudotyped viruses by reducing virion incorporation of envelope
6 glycoproteins. The molecular mechanisms by which MARCH8 targets envelope glycoproteins
7 remain unknown. Here, we show two different mechanisms by which MARCH8 inhibits viral
8 infection. Viruses pseudotyped with the VSV-G mutant, in which cytoplasmic lysine residues were
9 mutated, were insensitive to the inhibitory effect of MARCH8, whereas those with a similar lysine
10 mutant of HIV-1 Env remained sensitive to it. Ubiquitination assays showed that wild-type VSV-
11 G, but not its lysine mutant, was ubiquitinated by MARCH8. Furthermore, the MARCH8 mutant,
12 which had a disrupted cytoplasmic tyrosine motif that is critical for intracellular protein sorting,
13 did not inhibit HIV-1 Env-mediated infection, while it still impaired infection by VSV-G-
14 pseudotyped viruses. Overall, we conclude that MARCH8 impairs viral infectivity by
15 downregulating envelope glycoproteins through two different mechanisms mediated by a
16 ubiquitination-dependent or tyrosine motif-dependent pathway.

17

18

INTRODUCTION

1
2 Membrane-associated RING-CH (MARCH) 8 is one of 11 members of the MARCH family of
3 RING-finger E3 ubiquitin ligases, which consist of an N-terminal cytoplasmic tail (CT) domain
4 containing a C4HC3 RING finger (RING-CH finger) motif, two transmembrane (TM) domains,
5 between which a short ectodomain is located, and a C-terminal CT domain^{1,2}. MARCH8
6 downregulates a variety of cellular transmembrane proteins, such as MHC-II³, CD86⁴, CD81⁵,
7 CD44⁶, TRAIL receptor 1⁷, CD98⁶, IL-1 receptor accessory protein⁸, and transferrin receptor⁹. We
8 have recently reported that MARCH8 reduces HIV-1 infectivity by downregulating HIV-1
9 envelope glycoproteins (Env) from the cell surface, resulting in a reduced incorporation of Env
10 into virions¹⁰. Intriguingly, vesicular stomatitis virus G-glycoprotein (VSV-G) was even more
11 sensitive to the inhibitory effect of MARCH8. In the case of HIV-1 Env, it is retained
12 intracellularly without degradation after cell-surface downregulation. In contrast, VSV-G is not
13 only downregulated from the cell surface but also undergoes lysosomal degradation by
14 MARCH8¹⁰. In this regard, we hypothesized that VSV-G, whose cytoplasmic tail is lysine-rich (5
15 out of 29 amino acids), could be readily ubiquitinated by the E3 ubiquitin ligase MARCH8 and
16 therefore undergo lysosomal degradation, whereas HIV-1 Env carries only two lysines (out of 151
17 amino acids) in its cytoplasmic tail and may rarely undergo degradation after being trapped by
18 MARCH8. In this study, we created lysine mutants of both HIV-1 Env and VSV-G, together with
19 newly generated MARCH8 mutants to explore the hypothesis described above. The results with
20 these mutants show that MARCH8 targets HIV-1 Env and VSV-G by two different inhibitory
21 mechanisms.

22

RESULTS AND DISCUSSION

1
2 We have recently reported that MARCH8 inhibits lentiviral infection by reducing virion
3 incorporation of both HIV-1 Env and VSV-G in a RING-CH domain-dependent manner. Because
4 the RING-CH domain is known to be essential for the E3 ubiquitin ligase activity of MARCH8,
5 we asked whether these envelope glycoproteins are susceptible to MARCH8-mediated
6 ubiquitination. To investigate this, we first created the VSV-G mutant CT5K/R in which five
7 arginine residues were introduced in place of cytoplasmic lysine residues that could be
8 ubiquitination targets (Fig. 1A, upper). We also generated the HIV-1 Env gp41 mutant CT2K/R
9 harboring two arginines in place of the cytoplasmic lysines (Fig. 1A, lower). Then, we prepared
10 HIV-1 luciferase reporter viruses pseudotyped with the mutant envelope glycoproteins (VSV-G
11 CT5K/R and HIV-1 Env CT2K/R) from 293T cells transiently expressing MARCH8, and
12 compared their viral infectivity with that of control viruses pseudotyped with wild-type (WT)
13 envelope glycoproteins. The infectivity of viruses harboring either VSV-G CT5K/R or HIV-1 gp41
14 Env CT2K/R was almost comparable to WT-enveloped viruses (Figs. 1B and 1C). As expected,
15 the virus pseudotyped with VSV-G CT5K/R was completely resistant to MARCH8 (Fig. 1B). In
16 contrast, the HIV-1 Env CT2K/R-pseudotyped virus was still susceptible to the inhibitory effect
17 of MARCH8 (Fig. 1C). Consistent with these results, immunofluorescence staining showed that
18 the five lysine mutations in the CT domain of VSV-G conferred resistance to MARCH8-mediated
19 intracellular degradation (Fig. 1D), whereas the two lysine mutations in the CT domain of HIV-1
20 Env had no effect on its cell-surface downregulation by MARCH8 (Fig. 1E). It should be noted
21 that MARCH8-resistant VSV-G CT5K/R colocalized with MARCH8 (Fig. 1D). We thus

1 speculated that MARCH8 ubiquitinates lysine residues of VSV-G but not of HIV-1 Env at their
2 CT domains.

3 To investigate this possibility, we performed immunoprecipitation (IP)/Western-based
4 ubiquitination assays. In cells coexpressing WT HA-MARCH8, VSV-G was efficiently
5 ubiquitinated, whereas in cells coexpressing the RING-CH mutant of HA-MARCH8, the
6 ubiquitination of VSV-G was lost. More importantly, the VSV-G lysine mutant CT5K/R did not
7 undergo MARCH8-mediated ubiquitination, as expected (Fig. 1F), suggesting that the lysine
8 residues at the CT domain of VSV-G are specifically ubiquitinated by MARCH8. These findings
9 are consistent with the immunofluorescence results (Figs. 1D and 1E). We therefore conclude that
10 lysine residues at the CT domain of VSV-G are ubiquitinated by MARCH8, which determines the
11 difference in the MARCH8-mediated intracellular fate between these viral glycoproteins.

12 Because unlike VSV-G, HIV-1 Env was retained intracellularly without degradation, as we
13 previously reported¹⁰, we hypothesized that these viral envelope glycoproteins might undergo
14 endocytosis with different mechanisms of action. It has been reported that another MARCH family
15 member, MARCH11, has a conserved tyrosine-based motif, YXX ϕ , which is known to be
16 recognized by the adaptor protein (AP) μ -subunits in the C-terminal CT domain¹¹. We therefore
17 looked for the same motif(s) in MARCH8 and found the ²²²YxxL²²⁵ and ²³²YxxV²³⁵ sequences in
18 the CT domain at the C-terminus (Fig. 2A). Based on this finding, we generated tyrosine motif
19 mutants of MARCH8, in which either ²²²Y or ²³²Y was mutated to alanine (designated ²²²AxxL²²⁵
20 or ²³²AxxV²³⁵). The protein expression in cells transfected with each MARCH8 plasmid was
21 confirmed by immunoblotting using an anti-hemagglutinin (HA) antibody (Fig. 2B). Then, we
22 examined whether these YXX ϕ motifs are important for the antiviral activity of MARCH8. The

1 infectivity of VSV-G–pseudotyped viruses was still inhibited by the expression of both ²²²AxxL²²⁵
2 and ²³²AxxV²³⁵ MARCH8 mutants (Fig. 2C). In contrast, the infectivity of HIV-1 Env–
3 pseudotyped viruses was not impaired by ²²²AxxL²²⁵ MARCH8 expression but was reduced by
4 that of ²³²AxxV²³⁵ and WT MARCH8 (Fig. 2D), suggesting that the first tyrosine motif
5 (²²²YxxL²²⁵) is involved in the antiviral activity of MARCH8 against HIV-1 Env but not VSV-G.

6 We previously reported that the MARCH8-mediated reduction in viral infectivity was due to
7 reduced entry efficiency, resulting from a decreased virion incorporation of envelope
8 glycoproteins through their downregulation from the cell surface¹⁰. Therefore, by performing a β-
9 lactamase (BlaM)-fused viral protein R (Vpr)-based entry assay, we first focused on whether the
10 loss of function of ²²²AxxL²²⁵ MARCH8 against HIV-1 Env but not VSV-G would indeed be due
11 to the loss of its inhibitory activity on viral entry. Whereas the entry of VSV-G–pseudotyped HIV-
12 1 prepared from cells expressing either WT or ²²²AxxL²²⁵ MARCH8 was reduced compared with
13 that of the control virus (Fig. 2E), the inhibition of the entry of whole HIV-1 virions was abrogated
14 in viruses produced from cells expressing ²²²AxxL²²⁵, as expected (Fig. 2D). We further analyzed
15 whether this motif of MARCH8 is indeed involved in the reduced virion incorporation of HIV-1
16 Env, which results from its cell-surface downregulation. To address this, we conducted flow
17 cytometric analysis and quantified the levels of cell-surface and intracellular expression of Env
18 glycoproteins. In accordance with the results obtained in infectivity assays (Fig. 2C), ²²²AxxL²²⁵
19 MARCH8 still reduced intracellular VSV-G expression as well as WT MARCH8 did (Fig. 2G).
20 On the other hand, the mutant MARCH8 had a completely abrogated ability to downregulate cell-
21 surface HIV-1 Env, whereas WT expression led to the downregulation of HIV-1 Env from the cell
22 surface and its intracellular retention (Fig. 2H), as we previously observed¹². The results were

1 consistent with those of the inhibitory activity of MARCH8 against the virion incorporation of
2 HIV-1 Env (Fig. 2I). We therefore conclude that ²²²YxxL²²⁵ is critical for the MARCH8-mediated
3 downregulation of HIV-1 Env but not VSV-G, which results in its reduced virion incorporation
4 leading to impaired viral entry.

5 In summary, we first show the two different mechanisms by which MARCH8 inhibits viral
6 infections, one being a ubiquitin-dependent downregulation that mediates lysosomal degradation
7 of VSV-G whose cytoplasmic lysine residues are recognized by the RING-CH domain of
8 MARCH8 (Fig. 3, left), and the other, a YxxΦ motif-dependent downregulation that could explain
9 the intracellular retention of HIV-1 Env without degradation after cell-surface downregulation (Fig.
10 3, right). In terms of the latter mechanism, although this could be attributed to the AP-dependent
11 trafficking that requires the YxxΦ motif, which binds to AP μ-subunits, we were unable to prove
12 the interaction between these subunits and MARCH8 (data not shown). We have recently reported
13 that MARCH1 and MARCH2 are also antiviral MARCH family members that inhibit HIV-1
14 infection, although their antiviral activity is less robust, which is probably due to the lower protein
15 expression and/or stability than that of MARCH8¹². Because MARCH1 and MARCH2 also harbor
16 the Yxxφ motif in their C-terminal CT domains, it would be intriguing to verify the importance of
17 the motif in these proteins. Overall, our present findings are consistent with our previous studies
18 showing that MARCH8-induced downregulation of VSV-G leads to lysosomal degradation, while
19 that of HIV-1 Env results in intracellular retention without degradation. Thus, we conclude that
20 MARCH8 targets HIV-1 Env and VSV-G by two different inhibitory mechanisms (either
21 ubiquitin-dependent or YxxΦ motif-dependent downregulation). Further investigations will clarify
22 the more detailed host defense mechanisms of this protein.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22

METHODS

DNA constructs. The Env-deficient HIV-1 proviral indicator construct pNL-Luc2-E(-), HIV-1 Gag-Pol expression plasmid pC-GagPol-RRE, HIV-1 Env-expression vector pC-NLenv, HIV-1 Rev expression plasmid pCa-Rev, HIV-1 Tat expression plasmid pLTR-Tat, lentiviral packaging plasmid pMDLg/pRRE, VSV-G expression plasmid pC-VSVg and its C-terminally T7-epitope-tagged version pVSVg-T7E, GFP expression plasmid pCa-EGFP, Vpr/ β -lactamase (BlaM) expression plasmid pMM310, and MARCH8 expression plasmid either pC-MARCH8 or pC-HA-MARCH8 and its RING-CH mutant pC-HA-MARCH8-CS have previously been described elsewhere¹⁰. The HIV-1 Env gp41 mutant CT2K/R, in which cytoplasmic lysine residues at positions 792 and 806 were mutated to arginine, and the VSV-G mutant CT5K/R, in which cytoplasmic lysine residues at positions 491, 493, 496, 497, and 511 were mutated to arginine residues, were created by inserting overlapping PCR fragments into *MfeI/XhoI*-digested pC-NLenv and *KpnI/XhoI*-digested pVSVg-T7E, respectively. The hemagglutinin (HA)-tagged version of the WT and its lysine mutant NL Env expression plasmid were created by inserting the HA-tag linker into the *BamHI* site located at the gp41 cytoplasmic domain of pC-NLenv and its CT2K/R mutant. The MARCH8 mutant ²²²AxxL²²⁵ or ²³²AxxV²³⁵, in which a tyrosine residue at position 222 or 232 was mutated to an alanine residue, was generated by inserting overlapping PCR fragments into the *KpnI/XhoI*-digested pCAGGS or *XhoI-NotI*-digested pCAGGS-NHA to create an untagged or N-terminally HA-tagged expression plasmid. All constructs were verified by a DNA sequencing service (FASMAC).

1 **Cell maintenance and generation of stable cell lines.** 293T, MT4, HeLa, MAGIC5 (HeLa
2 derivative¹³), and HOS cells were maintained under standard conditions. Cells were originally
3 obtained from ATCC (except MAGIC5 cells) and routinely tested negative for mycoplasma
4 contamination (PCR Mycoplasma Detection kit, Takara).

5

6 **Virion production and infectivity assays.** To prepare VSV-G-pseudotyped or HIV-1-Env-
7 pseudotyped luciferase reporter viruses, 2.5×10^5 293T cells were cotransfected with 120 ng of
8 the MARCH expression plasmid (WT or RING-CH mutants), 20 ng of pC-VSVg, pC-VSVg-
9 CT5K/R, pC-NLenv, or pC-NLenv-CT2K/R, 500 ng of pNL-Luc2-E(-), and an empty vector up
10 to 1 μ g of total DNA, using FuGENE 6. Sixteen hours later, the cells were washed with phosphate-
11 buffered saline, and 1 ml of fresh complete medium was added. After 24 h, supernatants were
12 treated as described above and then harvested. The p24 antigen levels in viral supernatants were
13 measured by an HIV-1 p24 antigen capture ELISA (XpressBio). Transfection efficiencies were
14 normalized to the luciferase activity. To determine viral infectivity, 1×10^4 MAGIC5 cells were
15 incubated with 1 ng of p24 antigen from the HIV-1 supernatants. After 48 h, cells were lysed in
16 100 μ l of One-Glo Luciferase Assay Reagent (Promega). The firefly luciferase activity was
17 determined with a Centro LB960 (Berthold) luminometer.

18

19 **Immunoblotting assays.** Protein expression of constructs was confirmed by Western blot analyses
20 as described elsewhere^{10,12}. Briefly, cells transfected as described above were lysed in 500 μ l of
21 lysis buffer containing 1.25% n-octyl- β -D-glucoside, and Complete protease inhibitor mixture
22 (Roche Applied Science). Cell extracts were then subjected to gel electrophoresis and transferred

1 to a nitrocellulose membrane, followed by probing with an anti-HA mouse monoclonal antibody
2 (Sigma-Aldrich, H9658) or an anti- β -actin mouse monoclonal antibody (Sigma-Aldrich, A5316).
3 Proteins were then visualized by chemiluminescence using an ECL Western blotting detection
4 system (GE Healthcare) and monitored by using a LAS-3000 imaging system (FujiFilm).

5

6 **Viral entry assays.** A β -lactamase (BlaM)-fused viral protein R (Vpr)-based entry assay was
7 performed as described elsewhere^{10,12}. Briefly, HIV-1 particles containing a fusion protein of Vpr
8 and BlaM-Vpr were produced by cotransfection of 293T cells with pNL4-3, pMM310¹⁴ encoding
9 BlaM-Vpr, and either pC-MARCH8, pC-MARCH8-²²²AxxL²²⁵, pC-MARCH8-²³²AxxV²³⁵, or the
10 control vector. Similarly, VSV-G-pseudotyped HIV-1 particles containing BlaM-Vpr were
11 prepared by cotransfection with pNL-Luc-E(-), pC-VSVg, pMM310, and either pC-MARCH8,
12 pC-MARCH8-²²²AxxL²²⁵, pC-MARCH8-²³²AxxV²³⁵, or the control vector. The produced viruses
13 were normalized to the p24 antigen level (100 ng) and used for infection of the CD4⁺ T cell line
14 MT4 (5 x 10⁵ cells) at 37 °C for 4 h to allow viral entry. After extensive washing with Hank's
15 balanced salt solution (HBSS; Invitrogen), cells were incubated with 1 μ M CCF2-AM dye
16 (Invitrogen), a fluorescent substrate of BlaM, in HBSS containing 1 mg ml⁻¹ Pluronic F-127
17 surfactant (Invitrogen) and 0.001% acetic acid for 1 h at room temperature and then washed with
18 HBSS. Cells were further incubated for 14 h at room temperature in HBSS supplemented with
19 10% FBS, washed three times with PBS and fixed in a 1.2% paraformaldehyde solution.
20 Fluorescence was monitored at 520 and 447 nm by flow cytometry using BD FACS Canto II (BD
21 Bioscience), and the data were collected and analyzed with BD FACS Diva Software (BD
22 Bioscience).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22

Env incorporation assays. HIV-1 gp120 ELISA-based Env incorporation assays were described elsewhere¹².

Ubiquitination assays. 293T cells (5×10^5) were cotransfected with 0.8 μ g of pC-VSVg-T7E, 0.2 μ g of the empty vector, and 0.8 μ g of pC-HA-MARCH8. After 48 h, cells were lysed in TBS-T buffer (50 mM Tris-HCl buffer (pH 7.5), 0.15 M NaCl, 1% Triton X-100, and 0.5% deoxycholic acid) containing a protease inhibitor cocktail and 10 mM N-ethylmaleimide, as an inhibitor of deubiquitination enzymes. The mixture was centrifuged at $21,500 \times g$ for 15 min, and the supernatant was used as total cell lysate for immunoblotting or immunoprecipitation. Fifty microliters of Protein A-coupled Sepharose 4B (GE Healthcare, 17-0780-01) was preincubated for 2 h at 4 °C with 4 μ g of the appropriate antibody (anti-T7 epitope rabbit polyclonal antibody; MBL, PM022, H6908, anti-ubiquitin mouse monoclonal antibody; Clone FK2, Cayman, 14220). Total cell lysate was incubated with antibody-coupled Sepharose for 20 h at 4 °C. The Sepharose was washed three times with TBS-T buffer and one time with PBS before the immunoprecipitated proteins were eluted with SDS sample buffer. To evaluate the ubiquitination states of the immunoprecipitated proteins, proteins immunoprecipitated with an anti-T7 epitope rabbit antibody were subjected to Western blotting with an anti-ubiquitin mouse antibody, whereas proteins immunoprecipitated with the anti-ubiquitin mouse antibody were subjected to Western blotting with the anti-T7 epitope rabbit antibody. Immunoreactive bands were detected using an ECL detection kit (ATTO, EzWestLumi plus, WSE-7120) with a ChemiDoc imaging system (Bio-Rad).

1 **Immunofluorescence microscopy.** HOS cells were plated on 13-mm coverslips, cotransfected
2 with the indicated plasmids, 0.5 µg of either pC-NLenv or pC-VSVg-T7E, 0.1 µg of pC-Gag-Pol,
3 0.05 µg of pCa-Rev, and 0.3 µg of the pC-HA-MARCH8 expression plasmids (WT or CS-mutant)
4 using FuGENE6 and cultured for 24 h. For the total staining of both VSV-G and MARCH8, cells
5 were fixed with 4% paraformaldehyde for 30 min on ice and permeabilized with 0.05% saponin.
6 The fixed cells were incubated with both primary antibodies anti-T7 epitope mouse monoclonal
7 antibody (Novagen, 69522-4) and anti-HA goat polyclonal antibody (GenScript, A00168-40). The
8 secondary antibodies, Alexa 488 donkey anti-mouse IgG (Molecular Probes, A-21202) and Alexa
9 647 donkey anti-goat IgG (Molecular Probes, A-21447) were used for the double staining assay.
10 For the cell surface staining of gp120 protein, cells were incubated with an anti-gp120 goat
11 polyclonal antibody (Abcam, Ab21179) at 4 °C for 5 min and washed with PBS at 4 °C before
12 fixation. Fixation was performed with 4% paraformaldehyde for 30 min on ice, and fixed cells
13 were permeabilized with 0.05% saponin (Sigma-Aldrich) to detect the intracellular expression and
14 localization of MARCH8 proteins. The coverslips were incubated with the anti-HA mouse
15 monoclonal antibody (Sigma Aldrich, H9658) for 1 h, washed with PBS and incubated for 30 min
16 with the secondary antibodies Alexa 488 donkey anti-goat IgG (Molecular Probes, A-11055) and
17 Alexa 647 donkey anti-mouse IgG (Molecular Probes, A-31571) for the double staining assay.
18 Confocal images were obtained with a FluoView FV10i automated confocal laser-scanning
19 microscope (Olympus; Tokyo).

20

21

ACKNOWLEDGMENTS

1 This work was supported by grants from the Japan Society for the Promotion of Science
2 (KAKENHI, 18K07156) to K.T.

3

4

AUTHOR CONTRIBUTIONS

5 Y.Z., T.T., S.O., H.F. and K.T. performed the experiments and analyzed the data. Y.Z., H.F. and
6 K.T. discussed the data. S.K. provided reagents. K.T. conceived the study, supervised the work
7 and wrote the paper. All authors read and approved the final manuscript.

8

9

COMPETING FINANCIAL INTERESTS

10 The authors declare no competing financial interests.

11

12

1 **REFERENCES**

- 2 1. Bartee, E., Mansouri, M., Hovey Nerenberg, B. T., Gouveia, K. & Fruh, K.
3 Downregulation of major histocompatibility complex class I by human ubiquitin ligases
4 related to viral immune evasion proteins. *J Virol.* **78**, 1109-1120 (2004).
- 5 2. Goto, E. *et al.* c-MIR, a human E3 ubiquitin ligase, is a functional homolog of herpesvirus
6 proteins MIR1 and MIR2 and has similar activity. *J Biol Chem.* **278**, 14657-14668 (2003).
- 7 3. Ohmura-Hoshino, M. *et al.* Inhibition of MHC class II expression and immune responses
8 by c-MIR. *J Immunol.* **177**, 341-354 (2006).
- 9 4. Tze, L. E. *et al.* CD83 increases MHC II and CD86 on dendritic cells by opposing IL-10-
10 driven MARCH1-mediated ubiquitination and degradation. *J Exp Med.* **208**, 149-165
11 (2011).
- 12 5. Bartee, E. *et al.* Membrane-Associated RING-CH proteins associate with Bap31 and target
13 CD81 and CD44 to lysosomes. *PLoS One.* **5**, e15132 (2010).
- 14 6. Eyster, C. A. *et al.* MARCH ubiquitin ligases alter the itinerary of clathrin-independent
15 cargo from recycling to degradation. *Mol Biol Cell.* **22**, 3218-3230 (2011).
- 16 7. van de Kooij, B. *et al.* Ubiquitination by the membrane-associated RING-CH-8 (MARCH-
17 8) ligase controls steady-state cell surface expression of tumor necrosis factor-related
18 apoptosis inducing ligand (TRAIL) receptor 1. *J Biol Chem.* **288**, 6617-6628 (2013).
- 19 8. Chen, R., Li, M., Zhang, Y., Zhou, Q. & Shu, H. B. The E3 ubiquitin ligase MARCH8
20 negatively regulates IL-1beta-induced NF-kappaB activation by targeting the IL1RAP
21 coreceptor for ubiquitination and degradation. *Proc Natl Acad Sci U S A.* **109**, 14128-14133
22 (2012).

- 1 9. Fujita, H., Iwabu, Y., Tokunaga, K. & Tanaka, Y. Membrane-associated RING-CH
2 (MARCH) 8 mediates the ubiquitination and lysosomal degradation of the transferrin
3 receptor. *J Cell Sci.* **126**, 2798-2809 (2013).
- 4 10. Tada, T. *et al.* MARCH8 inhibits HIV-1 infection by reducing virion incorporation of
5 envelope glycoproteins. *Nat Med.* **21**, 1502-1507 (2015).
- 6 11. Morokuma, Y. *et al.* MARCH-XI, a novel transmembrane ubiquitin ligase implicated in
7 ubiquitin-dependent protein sorting in developing spermatids. *J Biol Chem.* **282**, 24806-
8 24815 (2007).
- 9 12. Zhang, Y. *et al.* Membrane-associated RING-CH (MARCH) 1 and 2 are MARCH family
10 members that inhibit HIV-1 infection. *J Biol Chem.* **294**, 3397-3405 (2019).
- 11 13. Mochizuki, N. *et al.* An infectious DNA clone of HIV type 1 subtype C. *AIDS Res Hum*
12 *Retroviruses.* **15**, 1321-1324 (1999).
- 13 14. Tobiume, M., Lineberger, J. E., Lundquist, C. A., Miller, M. D. & Aiken, C. Nef does not
14 affect the efficiency of human immunodeficiency virus type 1 fusion with target cells. *J*
15 *Virol.* **77**, 10645-10650 (2003).

16

17

1 **FIGURE LEGENDS**

2 **Figure 1 (A)** Schematic structure of the lysine mutants of VSV-G (CT5K/R; upper) and HIV-1
3 Env (CT2K/R; lower). SP, signal peptide; EC, extracellular domain; TM, transmembrane domain;
4 CT, cytoplasmic tail; SU, surface subunit. **(B)** Infectivity of viruses prepared from 293T cells
5 cotransfected with Env-defective HIV-1 luciferase (luc) reporter proviral DNA and either a control
6 (Ctrl) or HA-MARCH8 plasmid together with either the VSV-G wild-type (WT) (black) or
7 CT5K/R mutant (gray) plasmid. Data are shown as a percentage of the viral infectivity in the
8 absence of MARCH8 when WT VSV-G was used (mean + s.d. from three independent
9 experiments). **(C)** Infectivity of viruses prepared as shown in *B*, except for using either the WT
10 HIV-1 Env (black) or its CT2K/R mutant (gray) plasmid (mean + s.d. from three independent
11 experiments). **(D)** The VSV-G lysine mutant is resistant to MARCH8-mediated intracellular
12 degradation. Shown are immunofluorescence-based analyses of the intracellular levels of either
13 the WT or lysine mutant VSV-G with or without MARCH8 in transfected HOS cells. Scale bars,
14 10 μ m. **(E)** The lysine mutant of HIV-1 Env is still sensitive to MARCH8-induced downregulation
15 from the cell surface. Immunofluorescence images show cell-surface levels of either the WT or
16 lysine mutant HIV-1 Env with or without MARCH8 in transfected HOS cells. Scale bars, 10 μ m.
17 **(F)** Lysine residues at the CT domain of VSV-G are ubiquitinated by MARCH8. The
18 ubiquitination of the WT or lysine mutant VSV-G tagged with T7-epitope (T7e) in cells expressing
19 control or MARCH8 (WT or RING-CH mutant (CS)) was examined by immunoprecipitation of
20 either ubiquitinated proteins with an anti-ubiquitin antibody (left panel) or of T7e-tagged VSV-G
21 with an anti-T7e antibody (right panel), followed by immunoblotting with an antibody to either
22 T7e (left panel) or ubiquitin (right panel), respectively.

1
2 **Figure 2** (A) Schematic structure of YxxΦ motif mutants of MARCH8 (²²²YxxL²²⁵ and
3 ²³²YxxV²³⁵). (B) Western blot analysis was performed by using extracts from 293T cells
4 transfected with HA-tagged MARCH8 expression plasmids. Antibodies specific for HA were used
5 to detect MARCH8 proteins. (C) Infectivity of viruses prepared from 293T cells cotransfected
6 with Env-defective HIV-1 luciferase (luc) reporter proviral DNA and either a control (Ctrl), HA-
7 WT, HA-²²²AxxL²²⁵ or HA-²³²AxxV²³⁵ MARCH8 plasmid, together with either the VSV-G
8 expression plasmid. Data are shown as a percentage of the viral infectivity in the absence of
9 MARCH8 (mean + s.d. from three independent experiments). (D) Infectivity of viruses prepared
10 as shown in C, except for using the HIV-1 Env expression plasmid (mean + s.d. from three
11 independent experiments). (E, F) BlaM-Vpr-based viral entry assay using VSV-G-pseudotyped
12 viruses (E) or NL4-3 whole viruses (F) produced from cells expressing either control, WT
13 MARCH8, or the ²²²AxxL²²⁵ mutant. Representative FACS dot plots are shown from four
14 independent experiments. (G) VSV-G is downregulated by both WT and ²²²AxxL²²⁵ mutant
15 MARCH8, (H) whereas the cell-surface expression of HIV-1 Env is not affected by the mutant
16 MARCH8. (I) ²²²AxxL²²⁵ MARCH8 expression in producer cells is unable to decrease HIV-1
17 gp120 levels in viral supernatants. ELISA-based levels of Env gp120 in viral supernatants from
18 293T cells cotransfected with luc reporter proviral DNA and NL-Env plasmid, together with either
19 MARCH8 WT or its ²²²AxxL²²⁵ mutant. Representative data from three independent experiments
20 are shown as percent gp120 Env/p24 Gag in the supernatants relative to that from control cells.
21
22 **Figure 3** Schematic diagram of two different molecular mechanisms by which MARCH8

1 inhibits viral infection. *Left*, MARCH8 (red) downregulates VSV-G (violet) in a ubiquitin-
2 dependent manner. The RING-CH domain (pink) of MARCH8 recognizes VSV-G's
3 cytoplasmic lysine residues, which results in ubiquitin conjugation (shown as orange beads),
4 leading to lysosomal degradation; *Right*, MARCH8 downregulates HIV-1 Env (green) in a
5 Yxx Φ motif-dependent manner. The tyrosine motif located in the C-terminal CT of
6 MARCH8 likely interacts with the adaptor protein μ -subunits (navy) (if this is the case with
7 μ 2 or μ 1, clathrin (brown) is involved in this step), resulting in the intracellular retention of
8 HIV-1 Env without degradation. It should be noted that the downregulation of these viral
9 glycoproteins might not necessarily occur at the plasma membrane. The nucleus and other
10 organelles are not shown.

Figure 1




Figure 2




Figure 3

