1 **Complete Coding Genome Sequence for a Novel Multicomponent Kindia Tick Virus**
2 **Detected from Ticks Collected in Guinea**

3

4 Running title: Genome Sequence for Kindia Tick Virus from Guinea

5

6 Vladimir A. Ternovoi,^a Elena V. Protopopova,^a Alexander N. Shvalov,^a Mikhail Yu. Kartashov,^a
7 Roman B. Bayandin,^a Tatyana V. Tregubchak,^a Sergey A. Yakovlev,^b Konstantin A. Nikiforov,^b
8 Svetlana N. Konovalova,^a Valery B. Loktev,^a Alexander P. Agafonov,^a Rinat A. Maksyutov,^a
9 Anna Yu. Popova^c

10

11 ^aState Research Center for Virology and Biotechnology “Vector”, Koltsovo, Novosibirsk
12 Region, Russia

13 ^bFederal Government Health Institution Russian Research Anti-Plaque Institute “Microbe”,
14 Saratov, Russia

15 ^cRussian Federal Service for Surveillance on Consumer Rights Protection and Human
16 Wellbeing, Moscow, Russia

17

18 **Corresponding contributor:** Valery B Loktev, State Research Center of Virology and
19 Biotechnology «Vector», Rospotrebnadzor, Koltsovo, Novosibirsk Region, 630559, Russia,
20 Phone number: +7 383 3634700 add 2453; Facsimile number: +7 383 3367409; email addresses:
21 valeryloktev@gmail.com and loktev@vector.nsc.ru

22

23

24

25 ABSTRACT

26 Kindia tick virus (KITV) is a novel multicomponent virus first detected by direct sequencing of
27 *Rhipicephalus geigy* ticks in Guinea in 2017. Here, we present a complete coding genome
28 sequence for all four segments of KITV/2017/1. This virus appears to be evolutionarily related to
29 unclassified flaviviruses, such as Alongshan virus.

30

31 ANNOUNCEMENT

32 The multicomponent (segmented) tick flaviviruses evolutionarily related to the unsegmented
33 viruses of the genus *Flavivirus* have been recognized since only 2014 (1-4). This group of
34 unclassified segmented tick-borne flaviviruses now includes Alongshan virus, Jingmen tick virus
35 (JMTV) and Mogiana tick virus (MGTV), which have been detected in Asia, Europe and South
36 America, respectively. These diverse and globally distributed viruses are capable of infecting a
37 wide range of hosts, such as ticks, animals and humans (1–8). However, only a few complete
38 genome sequences have been reported for JMTV (Kosovo and China), Alongshan virus (China)
39 and MGTV (Brasilia).

40 Here, we report the complete coding genome sequences for the first African isolate, named
41 Kindia tick virus (KITV), from *Rhipicephalus geigy* collected from domestic cattle (*Bos taurus*)
42 in Kindia, Guinea, West Africa. Twenty-six pools of 5 ticks each were frozen in liquid nitrogen,
43 crushed by plastic pestles, homogenized in phosphate-buffered saline and used for RNA isolation
44 with TRIzol reagent (Invitrogen Co., USA). Total RNA was quantified with a Qubit RNA Assay
45 Kit (Invitrogen Co., USA) following the manufacturer's instructions. RNA-seq libraries were
46 constructed with an NEBNext Ultra RNA Library Prep Kit for Illumina (New England Biolabs).
47 Sequencing was performed using a MiSeq Reagent Kit v3 for 600 cycles. Cutadapt (version
48 1.18) and SAMtools (version 0.1.18) were used to remove the Illumina adaptors and duplicate
49 reads. After removing adapters, the read length was 108-118 bases, with the numbers of reads

per pool ranging from 151,282 to 540,929. The contigs were assembled *de novo* using the MIRA assembler with default parameters (version 4.9.6). In five pools, we found Mogiana-like fragments using BLASTN. These fragments were aligned to the reference genome segments of Mogiana tick virus isolate MGTV/V4/11 (4). The average coverage of the four segments was 35, 18, 52, and 30, respectively. The complete sequences for KITV/2017/1 were verified by Sanger sequencing of Seg 2 (2 short fragments), Seg 3 (1 sf) and Seg 4 (2 sf), with gaps or low coverage, as previously described (9). Total RNA from tick pools positive for KITV/2017/1 was used for first-strand DNA synthesis using a Reverta-L Kit (Interlabservice, Russia). The KITV/2017/1 genetic material was amplified by PCR with specific primers designed based on draft NGS sequences (available on request), with subsequent PCR fragment isolation and sequencing. The sequences for two detected KITVs (KITV/2017/1 and KITV/2017/2) were also deposited in GenBank. The nucleotide identity between KITV/2017/1 and KITV/2017/2 was 99.7% (Seg 1), 99.4% (Seg 2), 98.1% (Seg 3), and 99.3% (Seg 4). KITV contains putative open reading frames (ORFs) congruent with JMTV and MGTV, namely, nonstructural protein 1 (Seg 1), VP1 (Seg 2), NSP2 (Seg 3), VP2 and VP3 (Seg 4). The sizes of the sequenced segments (ORFs) are 2968 (2743), 2805 (2262), 2667 (2427) and 2725 (2351) bases, with CG contents of 52.3%, 54.9%, 54.4% and 54.2% for each segment, respectively. The lengths of the 5' UTR and 3' UTR were different for each segment and were 97-156 and 121-387 bases, respectively. The 5' UTR conservative motif GCAAGTGCA typical for JMTV was found in four segments, and the 3' UTR conservative motifs GGCAAGTGC and CAAGTG were also found in Seg 2 and Seg 4 of KITV. The divergence between KITV and MGTV/JMTV based on nucleotide sequence was 7–28% (Seg 1), 7.7–20.6% (Seg 2), 2.3–29.3% (Seg 3), and 0.4–22% (Seg 4), and based on amino acid sequences, it was 3.1–21.8% (Seg 1), 3.4–20.3% (Seg 2), 1.3–20.8% (Seg 3), and 0.2–15% (Seg 4), as evaluated by BLAST. Given these similarities, we suggest that KITV is a new member of the multicomponent (segmented) tick-borne flavivirus group and possibly represents a new species together with MGTV.

Data availability. GenBank accession numbers for the viral sequences: [MH678723](#) and [MH678727](#) (Seg 1), [MH678724](#) and [MH678728](#) (Seg 2), [MH678725](#) and [MH678729](#) (Seg 3), and [MH678726](#) and [MH678730](#) (Seg 4); complete sequences [MK673133](#) (Seg 1), [MK673134](#) (Seg 2), [MK673135](#) (Seg 3), and [MK673136](#) (Seg 4); and Sequence Read Archive accessions [SRX5930668](#) to [SRX5930671](#) under project [PRJNA545394](#). The annotations have also been deposited into GenBank for these sequences.

ACKNOWLEDGMENTS

The authors would like to gratefully acknowledge support from the Russian Federal Service for Surveillance on Consumer Rights Protection and Human Wellbeing, Project RPN₂2904, from 12/22/2017.

REFERENCES

1. Qin XC, Shi M, Tian JH, Lin XD, Gao DY, He JR, Wang JB, Li CX, Kang YJ, Yu B, Zhou DJ, Xu J, Plyusnin A, Holmes EC, Zhang YZ. 2014. A tick-borne segmented RNA virus contains genome segments derived from unsegmented viral ancestors. *Proc Natl Acad Sci U S A* 111:6744–6749. <https://doi.org/10.1073/pnas.1324194111>.
2. Shi M, Lin XD, Vasilakis N, Tian JH, Li CX, Chen LJ, Eastwood G, Diao XN, Chen MH, Chen X, Qin XC, Widen SG, Wood TG, Tesh RB, Xu J, Holmes EC, Zhang YZ. 2015. Divergent viruses discovered in arthropods and vertebrates revise the evolutionary history of the *Flaviviridae* and related viruses. *J Virol* 90:659–669. <https://doi.org/10.1128/JVI.02036-15>.
3. Ladner JT, Wiley MR, Beitzel IB, Auguste AJ, Dupuis AP, Jr, Lindquist ME, Sibley SD, Kota KP, Fetterer D, Eastwood G, Kimmel D, Prieto K, Guzman H, Aliota MT, Reyes D, Brueggemann EE, StJohn L, Hyeroba D, Lauck M, Friedrich TC, O'Connor DH, Gestole MC, Cazares LH, Popov VL, Castro-Llanos F, Kochel TJ, Kenny T,

- 102 White B, Ward MD, Loaiza JR, Goldberg TL, Weaver SC, Kramer LD, Tesh RB,
103 Palacios G. 2016. A multicomponent animal virus isolated from mosquitoes. *Cell Host*
104 *Microbe* 20:357–367. <https://doi.org/10.1016/j.chom.2016.07.011>.
- 105 4. Villa EC, Maruyama SR, de Miranda-Santos IKF, Palacios G, Ladner JT. 2017.
106 Complete Coding Genome Sequence for Mogiana Tick Virus, a Jingmenvirus Isolated
107 from Ticks in Brazil. *Genome Announc* 5(18). pii: e00232-17.
108 <https://doi.org/10.1128/genomeA.00232-17>.
- 109 5. Shi M, Lin XD, Vasilakis N, Tian JH, Li CX, Chen LJ, Eastwood G, Diao XN, Chen
110 MH, Chen X, Qin XC, Widen SG, Wood TG, Tesh RB, Xu J, Holmes EC, Zhang YZ.
111 2015. Divergent viruses discovered in arthropods and vertebrates revise the
112 evolutionary history of the *Flaviviridae* and related viruses. *J Virol* 90:659–669.
113 <https://doi.org/10.1128/JVI.02036-15>
- 114 6. Meng F, Ding M, Tan Z, Zhao Z, Xu L, Wu J, He B, Tu C. 2019. Virome analysis of tick-
115 borne viruses in Heilongjiang Province, China. *Ticks Tick Borne Dis* 10(2):412-420.
116 <https://doi.org/10.1016/j.ttbdis.2018.12.002>.
- 117 7. Emmerich P, Jakupi X, von Possel R, Berisha L, Halili B, Günther S, Cadar D, Ahmeti S,
118 Schmidt-Chanasit J. 2018. Viral metagenomics, genetic and evolutionary characteristics
119 of Crimean-Congo hemorrhagic fever orthonairovirus in humans, Kosovo. *Infect Genet*
120 *Evol* 65:6-11. <https://doi.org/10.1016/j.meegid.2018.07.010>.
- 121 8. Wang ZD, Wang B, Wei F, Han SZ, Zhang L, Yang ZT, Yan Y, Lv XL, Li L, Wang SC,
122 Song MX, Zhang HJ, Huang SJ, Chen J, Huang FQ, Li S, Liu HH, Hong J, Jin YL, Wang
123 W, Zhou JY, Liu Q.A. 2019. New Segmented Virus Associated with Human Febrile
124 Illness in China. *N Engl J Med* 380(22):2116-2125.
125 <https://doi.org/10.1056/NEJMoa1805068>.
- 126 9. Ponomareva EP, Ternovoi VA, Mikryukova TP, Protopopova EV, Gladysheva AV,
127 Shvalov AN, Konovalova SN, Chausov EV, Loktev VB. 2017. Adaptation of tick-borne

- 128 encephalitis virus from human brain to different cell cultures induces multiple genomic
- 129 substitutions. Arch Virol 162(10):3151-3156. <https://doi.org/10.1007/s00705-017-3442-x>