Functional screening of β -Glucanase Producing Actinomycetes Strains from Western Ghats ecosystems of Kerala, India

Lekshmi K. Edison¹ and N. S. Pradeep^{2}*

¹Microbiology Division, KSCSTE-Jawaharlal Nehru Tropical Botanic Garden and Research Institute, Palode, Thiruvananthapuram, Kerala- 695 562, India

^{2*}KSCSTE-Malabar Botanical Garden and Institute of Plant Sciences, Kozhikode, Kerala- 673014, India, Email: drnspradeep@gmail.com, Tel: +91 9446093865

Abstract

Screening of potential soil actinomycetes is static at infant phase because less than one part of soil biodiversity has been explored. An important factor considered before isolating microorganisms with potential application is understanding the biodiversity and environmental features associated with growth. Search of distinctive enzymes from unusual ecological habitats are highly fascinating and have great opportunities that may also pointed the developments in high throughput screening programs. In the present study Western Ghats hot spot regions of Kerala has been explored for the actinomycetes strains with beta glucanase activity. A total of 127 actinomycetes strains were isolated. After qualitative primary screening 106 strains (83%) produced exo- β -1,4-glucanase enzyme and 79 strains (62%) produced endo- β -1,3-glucanase enzyme. The quantitative secondary screening confirmed the strains TBG-MR17 and TBG-AL13 recognised as respective dominant producers of exo- β -1,4-glucanase and endo- β -1,3-glucanase enzymes. The study reveals the richness of the Western Ghats soils with innumerable actinomycetes having potential β -glucanase activities.

Keywords: Enzymatic index, Avicel, CM-curdlan, AZCL-pachyman

Introduction

Western Ghats in India are well-known biodiversity hot spot of rich flora and fauna, also recognised highly productive ecosystems. It is a forested strip of relatively old mountain ranges, beginning from Central Maharashtra and stretched up to the Southern tip of Kerala. These areas are granted with a “heritage tag” by UNESCO as a gene pool, sheltering millions of species of animals, plants and microbes. Western Ghats regions are domicile to immense collection of unexplored and novel microbial diversity including actinomycetes species. Exploitation of unique, natural, highly documented and less explored biodiversity ecosystems

for actinomycetes is highly necessary for the discovery of novel bioactive metabolites with prospective applications (Jalaja et al., 2011; Mohandas et al., 2012; Balachandran et al., 2012; Nampoothiri et al., 2013).

Microbial β -glucanase have been isolated from variety of microbes and well characterized (Velho-Pereira and Kamat, 2013). Actinomycetes have been extensively recognised as a source of β -glucan degrading enzymes. Among the wide genus, Streptomyces are most prevalent group of enzyme producers (Wu et al., 2018). Some of them includes Streptomyces sioyaensis (Hong et al., 2008), Streptomyces sp. 9X166 (Sakdapetsiri et al., 2016), Streptomyces sp. S27 (Shi et al., 2010), Streptomyces matensis ATCC 23935 (Woo et al., 2014), Streptomyces rochei (Wu et al., 2002) and Streptomyces sp. EF-14 (Fayad et al., 2001).

Nevertheless β -glucanase are less characterized in actinomycetes strains within the Western Ghats regions. These regions remain less explored, and also due to inextricable altered habitat the chances of obtaining potential strains of actinomycetes including Streptomyces species with exceptional β -glucanase activities are much higher. This chapter deals with the isolation of actinomycetes strains by exploring selected Western Ghats regions of Kerala for quantitative and qualitative screening of two β -glucanase enzymes, exo- β -1,4-glucanase and endo- β -1,3-glucanase. The synergic action of both β -glucanases requires the complete degradation of barley and oat β -glucans, which is essential for industrial applications such as brewing industry and feed enzyme industry.

Materials and Methods

Sample Collection

Soil samples were collected from Western Ghats of Kerala; include Wayandu, Munnar, Chinnar, Marayoor, Anamala, Neryamangalam, Nelliampathi, Agasthyarkoodum, Palode and Kulathupuzha. After removing the surface layer (approximately top 4 cm), the soil samples were taken from a depth of 5 to 10 cm of the superficial layers in each location. Three different samples were collected from each areas.

Pre-treatment of Soil Samples and Isolation of Actinomycetes Strains

Soil samples were pre-treated with 1% CaCO₃ incubated at 28°C for 10 days before use (El-Nakeeb and Lechevalier, 1963). One gram of soil taken in 100mL Erlenmeyer flask was pre-treated by heating the soil at 50°C for 1 h. Standard serial dilution plate method was

employed for the isolation of actinomycetes strains. 9mL sterile distilled water was added to the 1g of previously oven dried soil samples and mixed thoroughly in a rotary shaker for 30 min at 150 rpm at room temperature. The suspension was serially diluted to obtain 10⁻³ to 10⁻⁷ dilutions. 1.0mL of each dilution was pour plated on starch casein agar (SCA) plates in triplicate. After incubation at 28°C for 7 days the actinomycetes colonies were counted and represented as colony-forming units per gram (CFU.g⁻¹) of soil. For the purification of single isolated colonies, streak plate method was used.

Primary Screening for Exo-β-1,4-Glucanase and Endo-β-1,3-Glucanase Activities

Plate assay method was used for primary screening of enzymes (Meddeb-Mouelhi et al., 2014). Isolated actinomycetes strains were spot inoculated on a modified agar screening media with 1% (w/v) Avicel® PH-101 (Sigma) as a substrate for exo-β-1,4-glucanase and 0.2% (w/v) AZCL- Pachyman (Megazyme, Germany) as substrate for endo-β-1,3-glucanase enzymes and incubated for 5 days at 28°C. The presence of clear zone around the growth indicated the exo-β-1,4-glucanase activity. The enzymatic index (EI) of each strain was calculated as follows by measuring the diameter of hydrolysis zone and diameter of colony.

$$EI = \frac{\text{Diameter of hydrolysis zone } (\phi_h)}{\text{Diameter of colony } (\phi_c)}$$

The experiment was performed using three replicates for each strain. The average EI of three experiments were calculated, together with standard deviation.

Secondary Screening for Exo-β-1,4-Glucanase and Endo-β-1,3-Glucanase Activities

Modified liquid media with 0.5% (w/v) Avicel (for exo-β-1,4-Glucanase) and 0.2% (w/v) CM-curdlan (for endo-β-1,3-Glucanase) were used. The enzyme activities were quantitatively estimated by DNS assay method by measuring the released reducing sugars (Miller, 1959). One unit of beta glucanase activity is defined as the amount of enzyme required to release 1 μmol reducing sugar (as glucose equivalence) in one minute under defined conditions (Fulop and Ponyi, 1997).

Results

Isolation of Actinomycetes Strains

The study explored Western Ghats biodiversity for the isolation of β-glucanase producing actinomycetes strains. 10 different areas of Western Ghats of Kerala such as Wayanad,

Nelliyampathy, Neriya Mangalam, Munnar, Chinnar, Anamalai, Marayoor, Kulathupuzha, Palode and Agasthyarkoodam, were explored. All the selected sample collection spots were mountain and natural forest hot spot areas. A total of 127 morphologically different actinomycetes strains were isolated. The isolates formed well branched substrate mycelia and ample aerial hyphae that segregated into well-developed spores chains. The number of actinomycetes strains obtained per gram of soil samples from these areas are shown in Table 1.

Primary Screening of Exo- β -1,4-Glucanase and Endo- β -1,3- Glucanase Activities

All the isolated actinomycetes strains were evaluated for semi-quantitative production of exo- β -1,4-glucanase and endo- β -1,3-glucanase activities using plate assay method. Exo- β -1,4-glucanase production was screened using 1% Avicel (microcrystalline cellulose) as sole carbon source. The presence of a pale halo around the colonies after congo red dye staining indicated the production of enzyme exo- β -1,4-glucanase. This zone of clearance was due to hydrolysis of Avicel by exo- β -1,4-glucanase to glucose residues, it does not have any affinity to congo red. Out of 127 isolates, 106 strains (83% of total strains) produced exo- β -1,4-glucanase enzyme activity (Table 2).

The endo- β -1,3-glucanase activity of isolated actinomycetes strains were screened using 0.2% AZCL- Pachyman as a carbon source in plate assay. AZCL- Pachyman is an insoluble blue coloured azurine dye cross-linked substrate specifically for the determination of endo- β -1,3-glucanase activity. The endo- β -1,3-glucanolytic activity of actinomycetes strains were produced a blue halo around the colonies indicating the zone of hydrolysis. This is due to the activity of enzyme on insoluble AZCL-Pachyman released water soluble blue colour dyed fragments. Among the total 127 isolated strains, only 79 strains (62%) showed endo- β -1,3-glucanase activity (Table 3). The EI value of exo- β -1,4-glucanase was shown in between 1.7 to 7.3 and that of endo- β -1,3-glucanase was in between 1.8 to 8.5. Strains showed high EI values (in and above 4.5) were considered as potential enzyme producers and were selected for secondary screening (quantitative screening).

Secondary Screening of Exo- β -1,4-Glucanase and Endo- β -1,3-Glucanase Activities

Potential isolates selected by primary screening of both exo- β -1,4-glucanase and endo- β -1,3-glucanase enzymes were subcultured and prepared spore inoculum. Secondary or quantitative enzyme screening was performed by submerged fermentation. 14 strains with exo- β -1,4-glucanase and 8 strains with endo- β -1,3-glucanase activities were selected for quantitative

enzyme screening. Prepared 3×10^8 spores.mL⁻¹ were inoculated into liquid media for both enzymes. After 5 days of incubation at 28 °C in the particular screening media, the actinomycetes strain TBG-MR17 showed highest exo-β-1,4-glucanase activity and TBG-AL13 produced highest endo-β-1,3-glucanase activity. The exo-β-1,4-glucanase and endo-β-1,3-glucanase activities obtained in secondary screening are shown in table 4. The strains produced highest enzyme activities, TBG-MR17 for exo-β-1,4-glucanase activity (141 U.mL⁻¹) and TBG-AL13 for endo-β-1,3-glucanase activity (892 U.mL⁻¹), where considered as potent beta glucanase enzyme producers.

Discussion

Western Ghats are one of the treasurable natural resources of earth (Gadgil, 1979). Mostly it has protected and conserved unique biodiversity, point out the chance to discover unidentified microorganisms. Study on actinobacterial diversity for β-glucanase enzymes from 10 different less explored unusual and unique ecological niche in Western Ghats regions, has led to the isolation of 127 morphologically different actinomycetes strains. We have selected unexploited Western Ghats regions in Kerala mainly natural mountain and forest areas of Wayanad, Nelliampathy, Neriya Mangalam, Munnar, Chinnar, Anamalai, Marayoor, Kulathupuzha, Palode and Agasthyarkoodam. According to Nampoothiri et al., (2013), Western Ghats soil samples were immensely utilized for the isolation of industrially important novel enzyme producing microorganisms. Forest soils are the huge domicile of taxonomically diverse actinomycetes strains especially *Streptomyces* sp. Despite the presence of rich recalcitrant biopolymers they are actively involved forest nutrient turnover (Bontemps et al., 2013).

Pre-treatment of soil samples with CaCO₃ (1%) produced better colony counts. CaCO₃ accelerates the growth of actinomycetes spores in the collected soils. It is one of the good method for enriching actinomycetes propagules, which significantly yielded high total plate counting. This method was also proved effective in increasing the number of rare isolates. It yielded two-fold higher number of microbes than from untreated soil samples (Tiwari and Gupta, 2012). According to Otoguro et al. (2001), calcium carbonate soil treatment yielded good colony count (2.9×10^5 CFU.g⁻¹) of dried soil and leaf litter samples. CaCO₃ alter the pH of soil, thus favours the growth of spores and stimulate the formation of aerial mycelia (Natsume et al., 1989).

Current study also intended to screen and quantify two different β -glucanase, exo- β -1,4-glucanase and endo- β -1,3-glucanase, from isolated actinomycetes strains. Avicel (microcrystalline cellulose) was used as the substrate for exo- β -1,4-glucanase. As per earlier reports, Avicel is the specific substrate for exo-acting β -1,4-glucanase (Florencio et al., 2012; Annamalai et al., 2016b). Exo-glucanase or avicelase produced by *Streptomyces reticuli* was efficiently utilize Avicel (crystalline cellulose), when provided Avicel as a sole carbon source (Wachinger et al., 1989; Walter and Schrempf, 1996).

Primary screening is based on the clear halo formation around the isolated colonies which directly indicated the region of enzyme action to produce glucose units. The congo red dye remain attached to areas where the presence of β -1,4-D-glucanohydrolase bonds (Lamb and Loy, 2005). Out of total 127 isolates, 106 strains (83%) produced exo- β -1,4-glucanase enzyme activity, indicated majority of Western Ghats actinomycetes isolates are good producers of exo- β -1,4-glucanase. AZCL- Pachyman was used as substrate for screening endo- β -1,3-glucanase activity. According to previous reports, pachyman is a purely β -1,3-linked substrate so can be used for determining precise endo- β -1,3-glucanase activity (Zantinge et al., 2002; Sakamoto et al., 2006). Azurine cross-linked (AZCL) polysaccharide substrates are widely used for screening glycosyl hydrolases (Li et al., 2011; Nyysönen et al., 2013). Out of total 127 isolated strains, only 79 strains (62%) produced blue halo by the action of the endo- β -1,3-glucanase which degraded dye linked polysaccharides to monosaccharide and subsequently released dye.

Enzyme degradation in agar media was calculated in terms of enzymatic index. It is modest and fastest methodology to screen the strains prospective for enzyme production within the same genus (Ruegger and Tauk-Tornisielo, 2004). Ten et al. (2004), indicated that cellulase, xylanase and amylase producing strains effectively selected using halo zone diameter and colony diameter based enzymatic index. Strains showed high EI values (≥ 4.5) were considered as potential enzyme producers and selected for quantitative enzyme assays. Bhaturiwal et al., (2017), reported enzyme activity profiling of 20 actinomycetes strains such as cellulase, lipase, chitinase, β -mannanase, amylase, caseinase, caffeinase and xylanase activities by determining enzymatic index. Highest EI values were observed in TBG-NR3 (7.27 ± 0.225) and TBG-CH22 (8.53 ± 0.058) respectively for exo- β -1,4-glucanase and endo- β -1,3-glucanase plate assays.

Based on EI values 14 strains with exo- β -1,4-glucanase activity and 8 strains with endo- β -1,3-glucanase activity were selected for quantitative evaluation using submerged fermentation. After five days of incubation, actinomycetes strain TBG-MR17 showed highest exo- β -1,4-glucanase activity (95 U.mL⁻¹) and TBG-AL13 showed the highest endo- β -1,3-glucanase activity (219 U.mL⁻¹). Florencio et al. (2012), reported that no straight correlation was obtained between quantitative enzyme activity and enzymatic indexes. The same results were observed in our study. However quantitative method considered as final validation, as it deliver more exact data with slight variability, greater sensitivity and allow to compare relative amount of enzymes (Bisswanger, 2014; Farris et al., 2016).

Conclusion

Microbial communities from diverse Western Ghats ecological niches are almost unexplored and rich reservoirs of valuable metabolites, likely to provide extensive applications beneficial to humanity. Especially fast growing prerequisites for enzymes in diverse extents demands an urgent need to explore actinomycetes as a treasured source of marketable enzymes. Our exploration revealed that Western Ghats ecosystems are unusual habitat for promising actinobacterial diversity with extraordinary β -glucanase activity. Extensive range of climatic environments, rich woodland areas and less discrepancies in soil type with less acidic to alkaline pH and low EC, are relatively favourable for the largest distribution of actinobacteria with high β -glucanase activity. Total of 127 actinomycetes isolates were documented during the course of study. Qualitative enzyme assay revealed, 106 strains (83%) showed exo- β -1,4-glucanase enzyme activity and only 79 strains (62%) produced endo- β -1,3-glucanase activity. According to quantitative activity profiling, the actinomycetes strains TBG-MR17 and TBG-AL13 recognised as dominant exo- β -1,4-glucanase and endo- β -1,3-glucanase producers with 141 and 892 U.mL⁻¹ of respective activities. This is the first report of exploration of Western Ghats actinomycetes for both exo- β -1,4-glucanase and endo- β -1,3-glucanase enzymes with tremendously high activities.

References

- Balachandran, C., Duraipandiyan, V., & Ignacimuthu, S. (2012). Cytotoxic (A549) and antimicrobial effects of *Methylobacterium* sp. isolate (ERI-135) from Nilgiris forest soil, India. *Asian Pacific Journal of Tropical Biomedicine*, 2, 712-716.
- Bhaturiwala, R. A., Jha, S. C., Jain, N. K., & Modi, H. A. (2017). Enzyme profiling of selected chitinase producing Actinomycetes. *European Journal of Biotechnology and Bioscience*, 5(1), 39-43.
- Bisswanger, H. (2014). Enzyme assays. *Perspectives in Science*, 1(1-6), 41-55.

- Bontemps, C., Toussaint, M., Revol, P. V., Hotel, L., Jeanbille, M., Uroz, S., Turpault, M. P., Blaudez, D., & Leblond P. (2013). Taxonomic and functional diversity of Streptomyces in a forest soil. FEMS Microbiology Letters, 342(2), 157–167.
- El-Nakeeb, M. A., & Lechevalier, H. A. (1963). Selective isolation of aerobic actinomycetes. Applied Microbiology, 11(2), 75-77.
- Farris, M. H., Ford, K. A., & Doyle, R. C. (2016). Qualitative and quantitative assays for detection and characterization of protein antimicrobials. Journal of Visualized Experiments, 10(110), e53819.
- Fayad, K. P., Simao-Beaunoir, A. M., Gauthier, A., Leclerc, C., Mamady, H., Beaulieu, C., & Brzezinski, R. (2001). Purification and properties of a beta-1,6-glucanase from Streptomyces sp. EF-14, an actinomycete antagonistic to Phytophthora spp. Applied Microbiology and Biotechnology, 57(1-2), 117-23.
- Florencio, C., Couri, S., & Farinas, C. S. (2012). Correlation between agar plate screening and solid-state fermentation for the prediction of cellulase production by Trichoderma Strains. Enzyme research, 793708, 1-7.
- Florencio, C., Couri, S., & Farinas, C. S. (2012). Correlation between agar plate screening and solid-state fermentation for the prediction of cellulase production by Trichoderma Strains. Enzyme research, 793708, 1-7.
- Fulop, L., & Ponyi, T. (1997). Rapid screening for endo-β-1,4-glucanase and endo-β-1,4-mannanase activities and specific measurement using soluble dye-labelled substrates. Journal of Microbiological Methods, 29(1), 15-21.
- Gadgil, M. (1979). Hills, dams and forests. Some field observations from the Western Ghats. Proceedings of the Indian Academy of Science, 2(3), 291-303.
- Hong, T. Y., Hsiao, Y. Y., Meng, M., & Li, T. T. (2008). The 1.5 Å structure of endo-1,3-beta-glucanase from Streptomyces sioyaensis: evolution of the active-site structure for 1,3-beta-glucan-binding specificity and hydrolysis. Acta Crystallographica D Biological Crystallography, 64(9), 964-70.
- Jalaja, V., Swetha, S., Kumar, S. S., Deepthi, A., Kumar, R. S., & Pandey, A. (2011). Isolation and characterization of alpha amylase from a metagenomic library of Western Ghats of Kerala, India. Biologia, 66, 939-944.
- Lamb, J., & Loy, T. (2005). Seeing red: the use of Congo red dye to identify cooked and damaged starch grains in archaeological residues. Journal of Archaeological Science, 32(10), 1433–1440.
- Li, L. L., Taghavi, S., McCorkle, S. M., Zhang, Y. B., Blewitt, M. G., Brunecky, R., Adney, W. S., Himmel, M. E., Brumm, P., Drinkwater, C., Mead, D. A., Tringe, S. G., ... Lelie, D. V. (2011). Bioprospecting metagenomics of decaying wood: mining for new glycoside hydrolases. Biotechnology for biofuels, 4(1), 23.
- Meddeb-Mouelhi, F., Moisan, F. J., & Beauregard, M. (2014). A comparison of plate assay methods for detecting extracellular cellulase and xylanase activity. Enzyme and Microbial Technology, 66, 16-19.
- Miller, G.L. (1959). Use of dinitrosalicylic acid reagent for determination of reducing sugar. Analytical Chemistry, 31, 426-428.
- Mohandas, S. P., Ravikumar, S., Menachery, S. J., Suseelan, G., Narayanan, S.S., Nandanwar, H., & Nampoothiri, K.M. (2012). Bioactives of microbes isolated from Western Ghats belt of Kerala show

lactamase inhibition along with wide spectrum antimicrobial activity. *Applied Biochemistry and Biotechnology*, 167 (6), 1753- 1762.

Nampoothiri, K. M., Ramkumar, B., & Pandey, A. (2013). Western Ghats of India: Rich source of microbial diversity. *Journal of Scientific and Industrial Research*, 72, 617-623.

Natsume, M., Yasui, K., & Marumo, S. (1989). Calcium ion regulates aerial mycelium formation in actinomycetes. *The Journal of Antibiotics*, 42, 440-447.

Nyyssonen, M., Tran, H. M., Karaoz, U., Weihe, C., Hadi, M. Z., Martiny, J. B., Martiny, A. C., Brodie, E. L. (2013). Coupled high-throughput functional screening and next generation sequencing for identification of plant polymer decomposing enzymes in metagenomic libraries. *Frontiers in Microbiology*, 4, 282.

Otoguro, M., Hayakawa, M., Yamazaki, T., & Iimura, Y. (2001). An integrated method for the enrichment and selective isolation of *Actinokineospora* spp. in soil and plant litter. *Journal of applied microbiology*, 91(1), 118-130.

Ruegger, M. J. S., & Tauk-Tornisielo, S. M. (2004). Cellulase activity of fungi isolated from soil of the ecological station of Jureia- Itatins, Sao Paulo, Brazil. *Brazilian Journal Botany*, 27, 205-211.

Sakamoto, Y., Watanabe, H., Nagai, M., Nakade, K., Takahashi, M., & Sato, T. (2006). *Lentinula edodes* tlg1 encodes a thaumatin-like protein that is involved in lentinan degradation and fruiting body senescence. *Plant Physiology*, 141(2), 793-801.

Sakdapetsiri, C., Fukuta, Y., Aramsirujiwet, Y., Shirasaka, N., & Kitpreechavanich, V. (2016). Antagonistic activity of endo- β -1,3-glucanase from a novel isolate, *Streptomyces* sp. 9X166, against black rot in orchids. *Journal of Basic Microbiology*, 56(5), 469-79.

Shi, P., Yao, G., Yang, P., Li, N., Luo, H., Bai, Y., Wang, Y., & Yao, B. (2010). Cloning, characterization, and antifungal activity of an endo-1,3-beta-D: -glucanase from *Streptomyces* sp. S27. *Applied Microbiology and Biotechnology*, 85 (5), 1483-1490.

Ten, L. N., Im, W. T., Kim, M. K., Kang, M. S., & Lee, S. T. (2004). Development of a plate technique for screening of polysaccharide-degrading microorganisms by using a mixture of insoluble chromogenic substrates. *Journal of Microbiological Methods*, 56(3), 375–382.

Tiwari, K., & Gupta, R. K. (2012). Diversity and isolation of rare actinomycetes: an overview. *Critical Reviews in Microbiology*, 39(3), 256-294.

Velho-Pereira, S. & Kamat, N. M. (2013). Actinobacterial research in India. *Indian Journal of Experimental Biology*, 51, 573-596.

Wachinger, G., Bronnenmeier, K., Staudenbauer, W. L., & Schrempf, H. (1989). Identification of mycelium-associated cellulase from *Streptomyces reticuli*. *Applied Environmental Microbiology*, 55, 2653-2657.

Walter, S., & Schrempf, H., (1996). Physiological studies of cellulase (avicelase) synthesis in *Streptomyces reticuli*. *Applied Environmental Microbiology*, 62(3), 1065-1069.

Woo, J. B., Kang, H. N., Woo, E. J., & Lee, S. B. (2014). Molecular cloning and functional characterization of an endo- β -1,3-glucanase from *Streptomyces matensis* ATCC 23935. *Food Chemistry*, 1(148), 184-187.

Wu, H., Shimoi, H., & Ito, K. (2002). Purification and characterization of beta-1,6-glucanase of *Streptomyces rochei* application in the study of yeast cell wall proteins. *Bioscience Biotechnology and Biochemistry*, 66(11), 2515-2519.

Wu, Q., Dou, X., Wang, Q., Guan, Z., Cai, Y., & Liao, X. (2018). Isolation of β -1,3-Glucanase-Producing Microorganisms from *Poria cocos* Cultivation Soil via Molecular Biology. *Molecules*, 23(7), 1555.

Zantinge, J. L., Huang, H. C., & Cheng, K. J. (2002). Microplate diffusion assay for screening of beta-glucanase-producing microorganisms. *Biotechniques*, 33(4), 798-806.

Table 1. Description of soil sampling areas and number of colonies

Area	Description	Actinomycetes Colonies (CFU.g ⁻¹)
Wayanad	Mountain forest	15x10 ⁻³
Nelliyampathy	Hill range	10x10 ⁻³
Neryamangalam	Natural forest	7x10 ⁻³
Munnar	Hill station	11x10 ⁻³
Chinnar	Lower mountain forest	23x10 ⁻³
Anaimalai	Mountain	24x10 ⁻³
Marayur	Natural sandal wood forest	5x10 ⁻³
Kulathupuzha	Natural forest	13x10 ⁻³
Palode	Reserve forest	5x10 ⁻³
Agasthyakoodam	Hill range	14x10 ⁻³

Table 2. Primary screening of *exo-β-1,4-glucanase* activity

No	Strain	Mean Ø _h	Mean Ø _c	Mean EI	SD
1	TBG-CH1	2.9	0.7	4.10	0.089
2	TBG-CH2	2.4	0.7	3.40	0.089
3	TBG-CH3	3	0.7	4.30	0.155
4	TBG-CH4	3	0.9	3.30	0.179
5	TBG-CH5	3.3	0.9	3.73	0.137
6	TBG-CH6	3.2	1	3.23	0.186
7	TBG-CH7	3.3	0.8	4.13	0.137
8	TBG-CH8	3.4	0.9	3.80	0.089
9	TBG-CH9	No zone			
10	TBG-CH10	No zone			
11	TBG-CH11	3	0.9	3.33	0.258
12	TBG-CH12	2.4	0.7	3.40	0.089
13	TBG-CH13	2.3	1	2.30	0.089
14	TBG-CH14	2	0.6	3.30	0.089
15	TBG-CH15	2.9	0.7	4.13	0.052
16	TBG-CH16	2.5	0.6	4.20	0.089
17	TBG-CH17	3.3	0.9	3.63	0.052
18	TBG-CH18	3.1	0.8	4.10	0.322
19	TBG-CH19	1.8	0.8	2.23	0.052
20	TBG-CH20	3.1	0.8	4.00	0.089
21	TBG-CH21	2.5	0.6	4.20	0.179
22	TBG-CH22*	2.7	0.6	4.50	0.089
23	TBG-CH23	3.2	0.8	4.00	0.089
24	TBG-AL1	1.5	0.9	1.73	0.052
25	TBG-AL2	No zone			
26	TBG-AL3	2.3	1.2	1.87	0.052
27	TBG-AL4	1.5	0.8	1.90	0.179
28	TBG-AL5	2.9	1.2	2.40	0.155
29	TBG-AL6	3	0.9	3.30	0.089
30	TBG-AL7	3.5	1.2	2.90	0.089
31	TBG-AL8	1.75	0.7	2.50	0.089
32	TBG-AL9	2.8	0.7	4.03	0.052
33	TBG-AL10	3.15	0.8	3.87	0.052
34	TBG-AL11	2.1	0.8	2.60	0.089

35	TBG-AL12	3	0.9	3.30	0.089
36	TBG-AL13	3.5	1.3	2.73	0.137
37	TBG-AL14	1.8	0.8	2.30	0.155
38	TBG-AL16	2.5	1.2	2.10	0.155
39	TBG-AL17	1.5	0.5	3.03	0.052
40	TBG-AL18	2.7	0.7	3.93	0.186
41	TBG-AL19*	3.6	0.6	6.03	0.137
42	TBG-AL20	2.6	0.7	3.67	0.052
43	TBG-AL21	2.8	1	2.80	0.089
44	TBG-AL22	2.5	1	2.50	0.089
45	TBG-AL23	2.5	0.9	2.83	0.052
46	TBG-AL24	3.1	0.9	3.40	0.089
47	TBG-AL25	2.1	0.8	2.60	0.089
48	TBG-NR1*	3.7	0.8	4.63	0.052
49	TBG-NR2*	3.7	0.8	4.63	0.137
50	TBG-NR3*	2.9	0.4	7.27	0.225
51	TBG-NR4	1.7	0.5	3.37	0.137
52	TBG-NR6	3.9	1	3.90	0.089
53	TBG-NR7	3	1	2.97	0.186
54	TBG-NR11	3.6	0.9	4.00	0.179
55	TBG-NR14	3.6	0.9	3.90	0.089
56	TBG-NR16	2.8	0.9	3.07	0.052
57	TBG-NR18	3.7	0.9	4.07	0.103
58	TBG-NR19*	3.6	0.8	4.47	0.052
59	TBG-NR21*	3.8	0.8	4.80	0.179
60	TBG-NR22	No zone			
61	TBG-NR23*	3.6	0.7	5.10	0.089
62	TBG-NR24*	3.6	0.8	4.50	0.089
63	TBG-MN1	3.4	0.9	3.83	0.137
64	TBG-MN2	3.3	0.8	4.07	0.207
65	TBG-MN3	3.7	1.2	3.13	0.137
66	TBG-MN5*	3.3	0.7	4.67	0.186
67	TBG-MN6	3.5	1	3.47	0.137
68	TBG-MN7	3	0.8	3.80	0.089
69	TBG-MN8	3.3	0.8	4.13	0.137
70	TBG-MN9	3.4	0.8	4.27	0.137
71	TBG-MN10	3.1	0.9	3.40	0.089
72	TBG-MN11	3.5	1	3.53	0.137
73	TBG-MN12	3.3	0.8	4.13	0.137
74	TBG-MR1	3.7	0.9	4.07	0.137
75	TBG-MR2	2.9	0.8	3.57	0.052
76	TBG-MR3*	3.8	0.8	4.77	0.137
77	TBG-MR4	3.5	1	3.47	0.052
78	TBG-MR5	2.8	1	2.77	0.052
79	TBG-MR7	2.2	1	2.17	0.137
80	TBG-MR8*	4	0.8	5.03	0.052
81	TBG-MR9*	4.2	0.9	4.67	0.186
82	TBG-MR12	3.5	1	3.50	0.089

83	TBG-MR17*	3.5	0.7	5.03	0.137
84	TBG-MR18	2.8	0.8	3.47	0.052
85	TBG-MR19	3	0.8	3.75	0.089
86	TBG-I5II	3.7	0.9	4.10	0.089
87	TBG-N1	3.3	0.8	4.12	0.207
88	TBG-N2	3.4	0.8	4.20	0.155
89	TBG-N3	No zone			
90	TBG-N5	No zone			
91	TBG-N6	3.6	0.8	4.33	0.273
92	TBG-N7	No zone			
93	TBG-N8	2.7	0.8	3.40	0.155
94	TBG-NMP6	No zone			
95	TBG-NMP5	3.4	1.2	2.80	0.089
96	TBG-NMP7	No zone			
97	TBG-AM1A	2.5	0.9	2.80	0.089
98	TBG-AM5	3.5	0.8	4.40	0.089
99	TBG-AM31	3.7	1.1	3.40	0.155
100	TBG-AM21	2.4	0.8	3.00	0.390
101	TBG-AM27	No zone			
102	TBG201	2.8	0.8	3.50	0.390
103	TBG19	No zone			
104	TBG3-6	3.4	1.2	2.80	0.179
105	TBG3-8	No zone			
106	TBG3-12	No zone			
107	TBG-AM47	No zone			
108	TBG-AM83	3.3	0.8	4.10	0.473
109	TBG-AM2	2.3	0.4	3.80	0.237
110	TBG-AM42	No zone			
111	TBG-AM55	3.5	1	3.50	0.179
112	TBG-AM57	No zone			
113	TBG-AG21	3.7	0.6	4.03	0.186
114	TBG-AG20	No zone			
115	TBG-AG28	3.7	0.9	4.10	0.089
116	TBG-B1	No zone			
117	TBG-B2	2.3	1.3	1.80	0.237
118	TBG-B3	No zone			
119	TBG-B4	No zone			
120	TBG-B5	No zone			
121	TBG-WY1	No zone			
122	TBG-WY2	3.1	0.9	3.40	0.358
123	TBG-WY5	3.1	0.9	3.40	0.358
124	TBG-WY8	2.7	0.8	3.40	0.358
125	TBG-WY9	2.7	0.6	4.50	0.322
126	TBG-WY17	2.5	0.9	2.80	0.237
127	TBG-WY19	No zone			

*strains selected for secondary screening

Table 3. Primary screening of endo- β -1,3-glucanase activity

No	Strain	Mean Øh	Mean Øc	Mean EI	SD
1	TBG-CH1	2.2	0.6	3.60	0.200
2	TBG-CH2	No zone			
3	TBG-CH3	2	0.8	2.50	0.300
4	TBG-CH4	No zone			
5	TBG-CH5	1.9	0.9	2.13	0.252
6	TBG-CH6	2.2	0.6	3.60	0.300
7	TBG-CH7	1.5	0.5	3.00	0.300
8	TBG-CH8	No zone			
9	TBG-CH9	2.4	1	2.43	0.058
10	TBG-CH10	No zone			
11	TBG-CH11	No zone			
12	TBG-CH12	1.4	0.7	2.03	0.058
13	TBG-CH13	No zone			
14	TBG-CH14	No zone			
15	TBG-CH15	No zone			
16	TBG-CH16	No zone			
17	TBG-CH17	No zone			
18	TBG-CH18	No zone			
19	TBG-CH19	No zone			
20	TBG-CH20	2.2	0.8	2.80	0.100
21	TBG-CH21	2.4	0.7	3.40	0.458
22	TBG-CH22*	1.7	0.2	8.53	0.058
23	TBG-CH23	1.8	0.6	3.00	0.346
24	TBG-AL1	2.2	0.7	3.10	0.265
25	TBG-AL2	2.5	0.8	3.10	0.346
26	TBG-AL3	2.4	0.9	2.70	0.265
27	TBG-AL4	3.2	1.1	2.90	0.200
28	TBG-AL5	1.8	0.7	2.60	0.361
29	TBG-AL6	No zone			
30	TBG-AL7	No zone			
31	TBG-AL8	1.8	0.5	3.60	0.173
32	TBG-AL9	No zone			
33	TBG-AL10	No zone			
34	TBG-AL11	No zone			

35	TBG-AL12	2.3	0.8	2.90	0.200
36	TBG-AL13*	2.6	0.4	6.90	0.100
37	TBG-AL14*	3.4	0.8	4.60	0.458
38	TBG-AL16	No zone			
39	TBG-AL17	No zone			
40	TBG-AL18	No zone			
41	TBG-AL19	2.8	0.9	3.10	0.265
42	TBG-AL20	2.2	0.8	2.80	0.200
43	TBG-AL21	No zone			
44	TBG-AL22	2	1	2.00	0.300
45	TBG-AL23	2.5	0.7	3.90	0.265
46	TBG-AL24	2.5	0.6	4.10	0.173
47	TBG-AL25	2.3	0.8	2.80	0.200
48	TBG-NR1	3.2	0.9	3.50	0.361
49	TBG-NR2*	2.3	0.5	4.60	0.346
50	TBG-NR3	No zone			
51	TBG-NR4	No zone			
52	TBG-NR6	3	0.9	3.30	0.300
53	TBG-NR7	No zone			
54	TBG-NR11	3.5	0.8	4.30	0.200
55	TBG-NR14*	1.5	0.3	5.00	0.600
56	TBG-NR16	No zone			
57	TBG-NR18	3	0.9	3.30	0.265
58	TBG-NR19	2.8	1	2.80	0.100
59	TBG-NR21	3.1	0.9	3.40	0.361
60	TBG-NR22	No zone			
61	TBG-NR23	3.6	1.1	3.20	0.361
62	TBG-NR24*	2.5	0.5	5.00	0.600
63	TBG-MN1	2.2	0.7	3.10	0.200
64	TBG-MN2	1.9	0.9	2.10	0.200
65	TBG-MN3	3.5	0.9	3.80	0.265
66	TBG-MN5	2.7	0.7	3.80	0.265
67	TBG-MN6	2.6	1	2.60	0.300
68	TBG-MN7	1.9	0.5	3.80	0.200
69	TBG-MN8	2.5	0.6	4.10	0.361
70	TBG-MN9	2.2	0.8	2.75	0.466
71	TBG-MN10	2.5	0.8	3.10	0.265
72	TBG-MN11	2.5	1	2.50	0.265
73	TBG-MN12	No zone			
74	TBG-MR1	2	0.8	2.57	0.306
75	TBG-MR2*	2.3	0.5	4.60	0.100
76	TBG-MR3	2.5	0.8	3.10	0.100
77	TBG-MR4	2.1	0.7	3.00	0.173
78	TBG-MR5	1.3	0.6	2.10	0.100
79	TBG-MR7*	2.4	0.5	4.80	0.300
80	TBG-MR8	2.9	1	2.90	0.200
81	TBG-MR9	No zone			
82	TBG-MR12	2.6	0.8	3.20	0.400

83	TBG-MR17	2.8	0.9	3.10	0.265
84	TBG-MR18	No zone			
85	TBG-MR19	1.5	0.6	2.50	0.265
86	TBG-I5II	1.8	0.7	2.60	0.200
87	TBG-N1	2.2	0.7	3.10	0.100
88	TBG-N2	2.1	0.6	3.50	0.300
89	TBG-N3	2.7	0.7	3.90	0.265
90	TBG-N5	No zone			
91	TBG-N6	3	0.9	3.30	0.436
92	TBG-N7	1.5	0.6	2.50	0.300
93	TBG-N8	2.8	0.8	3.50	0.300
94	TBG-NMP6	No zone			
95	TBG-NMP5	No zone			
96	TBG-NMP7	2.7	0.8	3.30	0.265
97	TBG-AM1A	1.7	0.6	2.80	0.100
98	TBG-AM5	3.5	0.9	3.90	0.100
99	TBG-AM31	2	0.8	2.50	0.265
100	TBG-AM21	No zone			
101	TBG-AM27	No zone			
102	TBG201	1.7	0.7	2.40	0.100
103	TBG19	3.2	0.9	3.50	0.500
104	TBG3-6	No zone			
105	TBG3-8	No zone			
106	TBG3-12	No zone			
107	TBG-AM47	No zone			
108	TBG-AM83	1.3	0.3	4.30	0.265
109	TBG-A6-2	2.4	0.7	3.33	0.416
110	TBG-S14A2	No zone			
111	TBG-S40A5	3	1	3.00	0.200
112	TBG-S13A5	No zone			
113	TBG-AG21	No zone			
114	TBG-AG20	No zone			
115	TBG-AG28	2	0.8	2.50	0.300
116	TBG-B1	No zone			
117	TBG-B2	2.3	1.3	1.80	0.300
118	TBG-B3	No zone			
119	TBG-B4	No zone			
120	TBG-B5	No zone			
121	TBG-WY1	No zone			
122	TBG-WY2	3.7	1.1	3.40	0.173
123	TBG-WY5	3.3	0.9	3.70	0.200
124	TBG-WY8	2.1	0.5	4.20	0.361
125	TBG-WY9	3.2	0.9	3.60	0.265
126	TBG-WY17	No zone			
127	TBG-WY19	2.7	0.7	3.90	0.265

*strains selected for secondary screening

Table 4. Secondary screening of enzyme activities

<i>Exo-β-1,4-glucanase</i>			<i>Endo-β-1,3-glucanase</i>		
No	Strain	Enzyme (U. mL ⁻¹)	No	Strain	Enzyme (U. mL ⁻¹)
1	TBG-MR17	141±0.11	1	TBG-AL13	892±0.06
2	TBG-AL19	90±0.13	2	TBG-NR2	217±0.14
3	TBG-NR3	83±0.16	3	TBG-MR7	217±0.12
4	TBG-MR3	77±0.09	4	TBG-CH22	199±0.09
5	TBG-CH22	76±0.14	5	TBG-AL14	196±0.11
6	TBG-NR1	75±0.13	6	TBG-MR2	186±0.13
7	TBG-NR2	72±0.19	7	TBG-NR24	17.4±0.15
8	TBG-NR23	71±0.17	8	TBG-NR14	17.2±0.13
9	TBG-NR24	70±0.16			
10	TBG-MR9	70±0.15			
11	TBG-MR8	61±0.14			
12	TBG-NR19	58±0.16			
13	TBG-MN5	56±0.08			
14	TBG-NR21	55±0.17			