Elemental analysis of *Moringa oleifera* seeds by Laser Induced Breakdown Spectroscopy (LIBS) and its anti-cancer and anti-microbial studies

R. K. Aldakheel^{a,b}, S. Rehman^c, Firdos A. Khan^d, A. Mostafa^e, M. A. Almessiere^a,
M. A. Gondal^{f*1}, A. Baykal^g

^aDepartment of Biophysics, ^cDepartment of Epidemic Disease Research, ^dDepartment of Stem Cell Research, ^eDepartment of Nanomedicine, Institute for Research & Medical Consultations (IRMC), Imam Abdulrahman Bin Faisal University, Dammam, 31441, Saudi Arabia.

^bDepartment of Physics, College of Science, Imam Abdulrahman Bin Faisal University, Dammam, 31441, Saudi Arabia.

^eDepartment of Pharmaceutical Chemistry, College of Clinical Pharmacy, Imam Abdulrahman Bin Faisal University, King Faisal Road, P.O. Box 1982, Dammam 31441, Saudi Arabia

^fLaser Research Group, Physics Department, King Fahd University of Petroleum & Minerals, Dhahran 31261, Saudi Arabia

¹ **Abbreviations:** MOSs, *Moringa oleifera* seed; LIBS, Laser Induced Breakdown Spectroscopy; GC-MS, Gas chromatography–mass spectrometry.

Corresponding author: magondal@kfupm.edu.sa

19 **Abstract**

20 Moringa oleifera plant contains a numerous antioxidants, antibiotics and nutrients (vitamins and minerals)
 21 which makes it prospective for diverse biomedical applications. This report investigated the anti-
 22 cancerous and anti-microbial potential of the Moringa oleifera seeds (MOSs) due to the bioactive
 23 components a detail elemental analysis. MOSs in the form of pellets were used in elemental analysis.
 24 Alcoholic extraction of were utilized in anti-cancerous and anti-microbial activity.To clarify the anti-
 25 cancerous and anti-microbial potential of the (MOSs) because of the bioactive components. An elemental
 26 analysis was conducted using Laser Induced Breakdown Spectroscopy (LIBS) . The GC-MS used for the
 27 validation of the LIBS outcome. Additionally, the anti-cancerous and anti-microbial activity of the MOSs
 28 was evaluated. Herein, the human colorectal carcinoma cells (HCT-116) were treated with the seeds
 29 aqueous extracts for 48 h and the cell viability plus the DNA nuclear morphology were measured via the
 30 MTT assay and DAPI staining. The cell viability of the normal human embryonic stem cells (HEK-293)
 31 was also examined after the treatment. Percentage of cell viability and inhibitory concentration (IC₅₀) of
 32 both normal and treated cells were determined.The recorded LIBS signal of the MOSs revealed the
 33 existence of elements like Ca, K, Mg, P, S, Fe, Mn, Zn, Na and Se (vital for human health). The results of
 34 the MTT assay revealed a profound inhibitory action of the MOSs extracts against the HCT-116 cells
 35 growth. On top, such extracts did not affect the HEK-293 cells growth, indicating the specificity of the
 36 proposed extracts towards cancer cells escalation hindrance. The anti-microbial activity of the ethanolic
 37 MOSs extracts was tested on the S. aureus and E. coli bacteria using the Agar well diffusion assay. The
 38 observed anti-cancerous and the anti-microbial activities of the MOSs extracts can be attributed to the
 39 charisma of various bioactive compounds including the oleic acid, palmitic acid, hexadecenoic acid ethyl
 40 ester and D-allose.

41 *Conclusion:* Current observations may contribute towards the development of the MOSs-based
42 biomedicine (organic without any side effects, cheap, plentiful, pure and sustainable) effective for the
43 cancer and bacterial infection cure.

44 **Keywords:** *Moringa oleifera*; seed extracts; LIBS; GC-MS ;anti-cancer; anti-microbial

45

46 **1. Introduction**

47 Generally, the moringa tree with the scientific name of *Moringa oleifera* Lam. (MOL) is the part of
48 Moringaceae family [1] . Being originated from the Northern India and Pakistan region alongside the
49 Himalaya Mountains, moringa tree also grows in different parts of African, Asian, and South American
50 tropics [2-4] . In Pakistan, India, Thailand and Philippines the moringa leaves and fruits are usually
51 consumed as the vegetables [5]. In recent times, several countries (Mexico, Hawaii, Cambodia, and the
52 Caribbean Islands) started cultivating it due to its numerous health benefits and nutritional values [6,7].
53 Apart from the staple organic food with enriched vitamins, proteins and minerals, the natural moringa
54 leaves and fruits have several biological applications [5,8]. The MO as a food is recommended by the
55 WHO due to its ramified nutritional benefits to human health [5].

56 In the past, several emperors worldwide used to consume the fruits and leaves of the *Moringa* plants
57 for keeping the skin healthy, acquiring energy, relieving the tension/stress and pain during the battles [8]
58 . Intensive studies revealed that the MO have many notable attributes including anti-oxidant, anti-
59 microbial, anti-cancer, anti-inflammatory, anti-ulcer, anti-hypertensive, anti-urolithiatic, anti-diabetic,
60 anti-asthmatic, anti-aging, cardiovascular, analgesic, immunomodulation, hepatoprotective, diuretic,
61 anthelmintic, and hypoglycemic characteristics [2,9-19] . In fact, the presence of some distinctive
62 chemical compounds and elements such as the 4-(4'-O-acetyl- α -L-rhamnopyranosyloxy) benzyl
63 isothiocyanate, niazimicin, 4-(α -L-rhamnopyranosyloxy) benzyl isothiocyanate, pterygospermin, 4-(α -

64 Lrhamnopyranosyloxy), benzyl isothiocyanate, and benzyl glucosinolate impart the hypo-tensive, anti-
65 bacterial and anti-cancer efficacies to the MO leaves and fruits [20]

66 To take the advantages of the MO plant products as the food supplements and medicines as well as
67 to determine its other health benefits various techniques have been developed for extracting the contents
68 from the MO leaves and fruits. These methods include pressurized liquid extraction [21], ultrasound [22],
69 microwave-based extraction [23], and supercritical fluid extraction [24]. However, all these techniques
70 suffer from some limitations in terms of the requirement of a large quantity of toxic organic solvents,
71 cumbersome sample preparation procedures and costly [25]. To overcome such drawbacks the LIBS has
72 been emerged as an efficient approach to determine the elemental compositions (various trace elements)
73 in the extracts obtained from different medicinal plants.

74 Currently, the laser induced breakdown spectroscopy (LIBS) technique has widely been exploited
75 for elemental analyses of different types of samples. It is a versatile analytical technique with many distinct
76 features such as the real-time measurement, rapidity, cost-effectiveness, sensitiveness, and in-situ
77 elemental analysis with microanalyses capacity for diverse types of materials and phases (solid, liquid, or
78 gas) [26-31]. In addition, this technique is simple, eco-friendly, and does not need cumbersome sample
79 preparation protocol because the sample treatments frequently tend to induce errors through contamination
80 and losses. Depending on the varieties of the inorganic constituents present in the medicinal plants, the
81 LIBS technique can accurately disclose their secrecy in a scientific manner. It is needless to mention that,
82 besides the occurrences of diverse organic molecular components most of the edible plants (herbs and
83 vines) are enriched with inorganic molecules and elements (in the form of vitamins, proteins and
84 minerals), signifying their immense health benefits. Since ancient times, the Ayurvedic medicinal
85 practices have been relied on these crude chemical components extracted from the medicinal plants

86 [32,33]. The screening of the glycemic index of various elements in diverse medicinal plants was
87 performed to evaluate their anti-diabetic potency [34,35].

88 Based on the abovementioned factors, for the first time we used the LIBS technique to analyze the
89 presence of various trace elements in the MO seeds (MOSs) extracts. The gas chromatography - mass
90 spectroscopy (GC-MS) measurement of the ethanolic MOSs extract was carried out to validate the LIBS
91 results. In addition, the anti-cancerous and anti-microbial characteristics of the MOSs extracts were
92 evaluated. The human colorectal carcinoma (HCT-116) cancerous cells and embryonic kidney (HEK-293)
93 normal cells were chosen for the anti-cancerous assessment (using MTT assays) of the MOSs extracts.
94 The *gram-positive Staphylococcus aureus* (*S. aureus*) and *Escherichia coli* (*E. coli*) bacterial strains were
95 selected for the bactericidal evaluation (using Agar well diffusion) of the MOSs extracts. Results were
96 interpreted and discussed to understand the mechanism of the anti-cancerous and anti-microbial
97 effectiveness of such extracts.

98

99 **2. Experimental procedure**

100 **2.1. Seeds collection and extract synthesis**

101 The good quality MOSs were purchased from the local market (originally imported from India) and
102 used for the extract preparation in few steps. The coatings of the seeds were first removed before being
103 crushed and ground to fine powder. Next, the resultant powder was compressed in the form of pellets.
104 Fig.1 (a)-(d) shows various stages of the sample preparation for further analyses using the LIBS technique.
105 Various amounts of aliquots (5, 10, 15, 20 and 25 g) were mixed separately in 200 ml of deionized water
106 (DIW). The obtained solution was heated at 60 °C with constant stirring and then kept overnight before
107 being filtered (Whatman filter paper No.1, Sigma-Aldrich). The obtained filtrate was maintained at 4 °C
108 for the anti-cancerous activity analyses of the MOSs. An alcoholic extraction was prepared for the

109 antimicrobial activity test. In this process, the fine powder of MOSs of various contents (5, 10, 15, 20 and
110 25 gm) was mixed in ethanol (200 ml) separately and stirred for 5 h. Later, the solutions containing the
111 MOSs were filtered and placed inside a rotary evaporator.

112

113

114

115 **Fig 1.** *Moringa oleifera* seeds sample: **a)** seed, **b)** seeds without coat, **c)** crushed seeds, **d)** seeds in the form
116 of pellets.

117

118 2.2. LIBS Setup Details

119 Fig 2 depicts the customized LIBS setup used to quantify the constituent elements present in the
 120 MOSs samples. It consisted of quadrupled Q-switched Nd:YAG laser (model QUV-266-5) operated at the
 121 wavelength of 266 nm, repetition rate of 20 Hz, maximum output energy of 30 mJ and pulse width of 8
 122 ns. The collimated laser pulses were focused onto the MOSs pellet (target) using a UV convex lens of 30
 123 mm focal length for ablation. Upon ablating the target pellets a plasma plume was generated. The emitted
 124 plasma was detected/collected using the fiber optical system equipped with a miniaturized lens placed at
 125 an angle of 45°. The other end of fiber was connected to the 500 mm spectrograph (Andor SR 500i-A)
 126 with the grating groove of approximately 1200 lines/mm. The pellet was mounted on a 2-D motorized
 127 sample holder capable of moving in the X-Y direction to avoid the crust formation on the target surface
 128 due to multiple laser shots on the same spot. The emission spectra were recorded by a ICCD camera
 129 (model iStar 320T, 690 X 255 pixels with delay time setting at 300 ns) and transferred to the interfaced
 130 on line personal computer (PC).

131

132

Fig. 2. Schematic of our LIBS setup, where the inset shows an image of *Moringa oleifera* seed sample.

2.3 GC-MS measurement

The ethanolic MOSs extract was analyzed using GC-MS (Shimadzu GC-2010 Plus) which was equipped with a split/spitless auto-injector (AOC-20i series) and coupled to a QP2010 Ultra single quadrupole (Shimadzu Corporation, Kyoto, Japan). The GC separation was achieved via an Rxi-5MS fused silica capillary column (Restek, USA) of dimension (30 m × 0.25 mm id × 100 μm film thickness). The temperature was raised from 60 °C (kept for 0.5 min) to 280 °C at the rate of 5 °C/min (hold for 5 min). The inlet was operated in the splitless mode at the temperature of 270 °C. Helium was flown (purity

of 99.9999%) as the carrier gas at the rate of 1 ml/min and the temperature of the MS transfer line was maintained at 280 °C. The ion source was worked in the electron impact mode at the energy of 70 eV and temperature of 250 °C. The full scan mass spectra were collected from 33 to 550 m/z. The spectral data was obtained by controlling the GC-MS and processed using the GC-MS solution (version 4.52, Shimadzu Corporation, Japan). The detected volatile compounds were identified using the NIST 11 and WILEY 9 libraries and relative area of each compound was calculated.

150

2.4. Anti-cancerous activity evaluation of MOSs extract

2.4.1. In vitro cell culture and cell viability test

In this study, the cancerous and normal (healthy) cell line including the respective human colorectal carcinoma (HCT-116) and embryonic kidney (HEK-293) cells were used to evaluate the anti-cancerous activity of various MOSs extracts on these cells. Following the earlier prescribed method [36,37], the proposed cells were seeded in 96 well plates and grown in the Dulbecco's Modified Eagle Medium (DMEM) supplemented with the reagents such as selenium chloride, L-glutamine, fetal bovine serum, antibiotic penicillin and streptomycin. One group of these cells were cultured inside a CO₂ incubator at the temperature of 37 °C and treated at various concentrations (0.030 mg/ml to 0.1 mg/ml) of the MOSs extracts. Another group was treated at different concentrations (0.030 mg/ml to 0.1 mg/ml) of the MOSs extracts. The control group was devoid of the MOSs extracts. After treating the cells using the MOSs extracts for 48 hrs they were treated with 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyl-tetrazolium bromide (MTT assay, Molecules, New Zealand) for 4 hrs. Next, the growth medium was eliminated from the plates and dimethyl sulfoxide (DMSO) was incorporated in every well where MTT created the Formazan crystals. Later, the culture plates were inspected at the wavelength of 570 nm using a microplate

reader (Bio-Rad Laboratories, Hercules, CA, USA). Finally, the GraphPad Prism Software was used to analyze the acquired data where a one-way analysis of variance (ANOVA) statistical tool was utilized.

2.4.2. Nuclear staining via DAPI

The cancer cells (HCT-116) were cultured inside the CO₂ incubator at 37 °C and treated for 48 hrs using the MOSs extract of concentration 0.066 mg/ml. The control group was made without adding any such extract. The effects of the MOSs extracts on the cells nuclei were studied after staining by DAPI. Next, the cold paraformaldehyde (4%) was used to pre-treat these cells and washed using Triton X-100 (0.1%) prepared in the phosphate buffer saline (PBS). Then, both control cell and those treated with the MOSs extracts were stained using DAPI (1.0 µg/ml) made in the PBS. Lastly, the cells were rinsed using Triton X-100 (0.1%) made in PBS. The morphologies of the control cells and those treated by MOSs extracts were scanned via a confocal scanning microscope (CSM, Zeiss, Germany).

2.5. Anti-microbial activity assessment of MOSs extract

The bactericidal efficacy of the proposed MOSs extracts at various concentrations (50, 100, 150, 200, 250 mg/ml) was assessed using the Agar well diffusion method. The *gram-positive S. aureus* and *E. coli* bacterial strain were selected. For the preparation of inoculum, bacteria were freshly grown in the nutrient broth (NB) at 37 °C by incubating overnight and regulated to the 0.5 McFarland standards. The Mueller Hinton agar solution was prepared by adding 100 µl of freshly adjusted inoculum. The inoculum was spread uniformly over the surface of the MHA plates and left to dry under the aseptic conditions. Consequently, the sterile cork-borer was used to punch the inoculated plates for the wells of 6 mm. Approximately 50 µl of the prepared extract was placed in the wells at 37 °C for overnight incubation. At the end of the incubation period, the bacterial zone

of inhibition (ZOI) diameter around the wells was measured to evaluate the inhibition effectiveness of the studied MOSs extracts [38].

3. Results and Discussion

3.1. Qualitative analysis of MOSs using LIBS

The LIBS spectra (Fig 3) of the MOSs were recorded in range of 200 to 800 nm. For detecting the major elements, the spectra were collected from different regions of the pellets each of scan length 50 nm. In order to reduce the background noise that may affect the quality of the LIBS signal, the LIBS parameters including the delay times, number of accumulations and laser energies were optimized. The recorded LIBS signal of the MOSs comprised of various significant spectral peaks of the atomic and ionic lines (with varying intensities) were related to abundant elements. Based on the NIST database the observed spectral lines were identified and categorized in terms of the characteristic elements present in the MOSs, suggesting their significant role towards the anti-microbial and anti-cancerous activities. Clearly, the measured LIBS spectra (Fig 3) showed the presence of Ca, K, Mg, P, S, Fe, Mn, Zn, Na and Se.

Fig 3. LIBS spectra of *Moringa oleifera* seeds.

205

206 Table 1 presents the LIBS signal intensities of the spectral transition lines corresponding to various
 207 detected elements in MOSs. In accordance to the Boltzmann distribution, the intensities of the LIBS
 208 spectral lines have direct relationship with elemental contents in MOSs [39]. This correlation was attained
 209 by considering the intensity ratio of the detected elemental lines with that of the C line taken as the
 210 reference (247.8 nm). The achieved intensity ratio of the elements Ca, K, Mg, P, S, Fe, Mn, Zn, Na and
 211 Se were ranged from 2.7 - 1.7, 1.2 - 1.4, 0.9, 1.1, 1.1, 1.1 - 0.9, 1.0, 0.9, 1.2 and 1.1, respectively which
 212 were consistent with those reported in the literature [40]. These results approved that the MOSs are rich
 213 in different minerals which are useful to human as food and medicine, indicating their remarkable
 214 influence in regulating the level of blood pressure, blood lipids, regulating the stomach, protecting the
 215 liver, strengthening bones, generate protein and enhancing the immunity of the body [41]. Furthermore,
 216 the existence of Se in the MOSs seeds plays a vital role by protecting from cancer, cardiovascular disease,
 217 cognitive decline and thyroid disease. Moreover, these seeds exhibit a powerful antibacterial activity due
 218 to the availability of the detected elements including Ca, K, Mg, P, S, Fe, Mn, Zn, Na and Se [42].

219

220 **Table 1.** Some of the detected spectral lines of the elemental analytes present in the Moringa seeds
 221 recorded by using our LIBS system.

Elements	Wavelength (nm)	Transition Configuration	LIBS Signal Intensity (a.u.)
Ca	422.6	$3p^64s^2\ ^1S_0 \rightarrow 3p^64s4p\ ^1P^{\circ}_1$	4960.9
	518.8	$3p^64s4p\ ^1P^{\circ}_1 \rightarrow 3p^64s5d\ ^1D_2$	2967.1
	527.0	$3p^63d4s\ ^3D_3 \rightarrow 3p^63d4p\ ^3P^{\circ}_2$	3196.3

222	Fe	248.3	$3d^6 4s^2 {}^5D_4 \rightarrow 3d^6({}^5D) 4s 4p ({}^1P^o) {}^5F^o_5$	2114.1
223		252.2	$3d^6 4s^2 {}^5D_4 \rightarrow 3d^6({}^5D) 4s 4p ({}^1P^o) {}^5D^o_4$	1810.1
224		259.9	$3d^6 ({}^5D) 4s {}^6D_{9/2} \rightarrow 3d^6({}^5D) 4p {}^6D^o_{9/2}$	2061.9
225	K	414.9	$3p^5 3d {}^3P^o_0 \rightarrow 3p^5 4p {}^3D_1$	2347.6
226		430.5	$3p^5 3d {}^3P^o_2 \rightarrow 3p^5 4p {}^1D_2$	2270.9
227		532.3	$3p^6 4p {}^2P^o_{1/2} \rightarrow 3p^6 8s {}^2S_{1/2}$	2639.7
228	Mg	518.3	$3s 3p {}^3P^o_2 \rightarrow 3s 4s {}^3S_1$	1753.5
229	Mn	259.3	$3d^5 ({}^6S) 4s {}^7S_3 \rightarrow 3d^5({}^6S) 4p ({}^7P^o_3)$	1843.9
230	Na	249.3	$2s^2 2p^5 3s {}^1P^o_1 \rightarrow 2s^2 2p^5 3p {}^1S_0$	2128.4
231		251.5	$2s^2 2p^5 3p {}^3S_1 \rightarrow 2s^2 2p^5 ({}^2P^o_{1/2}) 3d {}^2[{}^3/2]_2$	1834.7
232	P	253.5	$3s^2 3p^3 {}^2P^o_{3/2} \rightarrow 3s^2 3p^2({}^3P) 4s {}^2P_{3/2}$	2100.0
233		255.3	$3s^2 3p^3 {}^2P^o_{1/2} \rightarrow 3s^2 3p^2({}^3P) 4s {}^2P_{1/2}$	2032.1
234	S	416.2	$3s^2 3p^2({}^3P) 4p {}^4D^o_{7/2} \rightarrow 3s^2 3p^2({}^3P) 4d {}^4F_{9/2}$	1987.8
235	Se	418.09	$4s^2 4p^2 ({}^3P) 5p {}^4D^o_{7/2} \rightarrow 4s^2 4p^2 ({}^3P) 5d {}^4F_{9/2}$	1969.7
236		522.7	$4s^2 4p^2 ({}^3P) 5s {}^4P^o_{5/2} \rightarrow 4s^2 4p^2 ({}^3P) 5p {}^4D^o_{7/2}$	2002.0
237	Zn	250.1	$3d^{10} 4p {}^2P^o_{1/2} \rightarrow 3d^{10} 5s {}^2S_{1/2}$	1751.4
238		255.7	$3d^{10} 4p {}^2P^o_{3/2} \rightarrow 3d^{10} 5s {}^2S_{1/2}$	1762.3
239	C	426.7	$2s^2 3d {}^2D^o_{5/2} \rightarrow 2s^2 4f {}^2F^o_{7/2}$	2379.0
240		247.8	$2s^2 2p^2 {}^1S_0 \rightarrow 2s^2 2p 3s {}^1P^o_1$	1831.6
241	O	418.9	$2s^2 2p^2 ({}^1D) 3p {}^2F^o_{7/2} \rightarrow 2s^2 2p^2 ({}^1D) 3d {}^2G_{9/2}$	2003.4

3.2. Volatile content analyses of MOSs using GC-MS

A total of 114 volatile compounds were detected in the proposed MOSs via GC-MS data analyses.

These compounds included different chemical groups including varieties of fatty acid, aldehyde, ester,

243 alcohol, ketone and hydrocarbon. Table 2 and Fig 4 show the retention times and percentage composition
244 of all the identified chemical compounds existed in the MOSs. The detected major constituents were the
245 oleic acid (22.53%), 2,3-dihydroxypropyl elaidate (13.48%), 9-octadecenoic acid (Z)-, 2,3-
246 dihydroxypropyl ester (11.35%), docosenamide (6.04%), ethyl oleate (6.03%), 1,3-propanediol, 2-ethyl-
247 2-(hydroxymethyl) (5.52%), oleic anhydride (3.96%) and 2-Propanone, 1,1-dimethoxy (3.86%). These
248 MOSs are rich in the fatty acids and their ester derivatives (65.45%) followed by alcohols (9.4%), nitrogen
249 containing compounds (9.09%), ketones (5.34%) and aldehydes (2.88%). The fatty acids and alcohols in
250 plants may undergo esterification to form esters [43]. The GC-MS measurement revealed the occurrences
251 of several fatty acids and their esters such as oleic, n-hexadecenoic (palmitic), cis-9-Hexadecenoic acid
252 (Palmitoleic), octadecanoic (stearic) acids and their alcohols and esters. In fact, most of such compounds
253 have been reported to possess anti-cancer activity. For example, the D-allose was reported to inhibit cancer
254 cells growth at G1 phase [44]. The palmitic acid disclosed a selective cytotoxicity against the leukemic
255 cells in humans [45] . In addition, such fatty acids are known to have antifungal and antibacterial effects
256 [46]. The oleic acid that was identified as the major compound in the MOSs is an omega-9-fatty acid with
257 numerous benefits for the human health. This can effectively be used to prevent the ulcerative colitis [47]
258 and reduce the blood pressure [48] . On top, it has remarkable antioxidant efficiency [49].
259

Fig 4. GC-MS chromatograph of *Moringa oleifera* seeds.

Table 2: Volatile compounds identified in *Moringa oleifera* seeds using GC-MS analysis

No.	Compounds	RT	Peak area (%)	No. on Chromatogram
Esters				
1	Propanoic acid, 2-oxo-, methyl ester	6.29	0.76	3
2	Acetic acid, ethoxyhydroxy-, ethyl ester	7.04	0.04	5
3	Diethoxymethyl acetate	12.83	0.09	28
4	2-Propenoic acid, 2-methyl-, 2-hydroxypropyl ester	13.13	0.02	29
5	6,9,12-Octadecatrienoic acid, phenylmethyl ester, (Z,Z,Z)-	17.80	0.01	42
6	Cyclopentanecarboxylic acid, 4-tridecyl ester	19.18	0.03	47
7	1-Cyclohexene-1-carboxylic acid, 2,6,6-trimethyl-, methyl ester	20.06	0.13	49
8	Hexanoic acid, 4-hexadecyl ester	28.29	0.07	64

9	Phthalic acid, diethyl ester	29.57	0.14	68
10	2-Propenoic acid, pentadecyl ester	31.63	0.59	70
11	Acetic acid, 3,7,11,15-tetramethyl-hexadecyl ester	33.63	0.02	72
12	Phthalic acid, dibutyl ester	35.69	0.07	77
13	Phthalic acid, diisobutyl ester	35.76	0.15	78
14	Palmitoleic acid, methyl ester	36.29	0.03	79
15	Pentadecanoic acid, 13-methyl-, methyl ester	36.64	0.09	80
16	9-Hexadecenoic acid, ethyl ester	37.65	0.15	86
17	Hexadecanoic acid, ethyl ester (Ethyl palmitate)	37.99	0.78	88
18	Heptadecanoic acid, ethyl ester	38.42	0.46	90
19	Propanoic acid, 3-mercapto-, dodecyl ester	38.59	0.25	91
20	9-Octadecenoic acid, methyl ester, (E)- (Methyl elaidate)	40.16	1.52	94
21	Oleic acid, methyl ester (Methyl Oleate)	40.26	0.15	95
22	L-(+)-Ascorbic acid 2,6-dihexadecanoate	40.56	0.10	96
23	Ethyl Oleate	41.39	6.03	98
24	Octadecanoic acid, ethyl ester (Ethyl stearate)	41.77	0.53	100
25	9-octadecenyl ester (Oleoyl oleate)	43.52	0.34	102
26	2,3-dihydroxypropyl elaidate	43.79	13.48	104
27	9-Octadecenoic acid, 1,2,3-propanetriyl ester	44.29	0.90	105
28	Docosanoic acid, ethyl ester	45.27	0.37	107
29	Oleoyl chloride	46.07	0.65	109
30	9-Octadecenoic acid (Z)-, 2,3-dihydroxypropyl ester	47.10	11.35	111
31	Glycidol stearate	47.52	1.76	113
32	Hexadecanoic acid, 2-hydroxy-1-(hydroxymethyl)ethyl ester	47.80	1.57	114
Alcohols				
33	1,4-Cyclohexanediol, trans-	7.22	0.02	6
34	1,2-Propanediol, 3-methoxy-	7.43	0.05	7
35	Ethanol, 2,2-diethoxy-	7.54	0.31	8
36	1,2,4-Butanetriol	9.50	0.04	15
37	Glycerin	11.18	1.35	21
38	1-Butanol, 4-(ethylthio)-	11.74	0.02	22
39	Methoxyacetaldehyde diethyl acetal	11.88	0.02	24
40	1-Dodecanol	18.43	0.48	43
41	1-Tetradecanol	24.15	0.42	58
42	1,3-Propanediol, 2-ethyl-2-(hydroxymethyl)-	25.44	5.52	60
43	1-Tridecanol	26.33	1.06	61
44	3-Hexadecanol	31.35	0.05	69
45	3-Heptadecanol	35.56	0.06	76
Aldehydes				
46	Butanal, 3-hydroxy-	8.38	0.00	12
47	Heptanal	10.01	0.02	17
48	Octanal	12.57	0.04	27
49	Nonanal	15.79	0.09	36
50	2-Methyl-oct-2-enal	20.61	0.04	51
51	Undecanal	23.52	0.02	54

52	Tridecanal	34.46	0.01	74
53	10-Octadecenal	37.42	0.09	84
54	9-Octadecenamide	37.55	0.09	85
55	cis-9-Hexadecenal	38.24	0.55	89
56	cis-13-Octadecenal	47.23	1.92	112
Ketones				
57	2-Propanone, 1,1-dimethoxy-	6.84	3.86	4
58	2-Butanone	8.15	0.03	10
59	Dihydroxyacetone	8.97	0.14	13
60	1,2-Cyclopentanedione	10.17	0.14	18
61	2-Propanone, 1-(1,3-dioxolan-2-yl)-	10.87	0.04	20
62	1,3-Dioxol-2-one,4,5-dimethyl-	13.53	0.28	30
63	2-Heptanol, 5-ethyl-	14.31	0.01	33
64	2-Methyl-4-octanone	16.47	0.03	37
65	2-Pentanone, 3,4-epoxy-	18.93	0.03	45
66	1-Oxa-spiro[4.5]deca-6,9-diene-2,8-dione	37.01	0.05	82
67	Z-11-Pentadecenol	39.21	0.05	92
68	Cyclopentadecanone, 2-hydroxy-	39.82	0.13	93
69	Cyclopentadecanone	43.43	0.21	101
70	2-Tetradecanone	46.69	0.34	110
Acids				
71	Acetic acid, (acetyloxy)-	5.88	0.18	2
72	Butanoic acid, 3-hydroxy-	10.59	0.03	19
73	Octanoic acid	17.42	0.03	41
74	Nonanoic acid	20.36	0.02	50
75	n-Hexadecanoic acid (Palmitic acid)	37.31	0.05	83
76	Oleic Acid	40.94	22.53	97
Furans and lactones				
77	Furfural	7.62	0.02	9
78	2(5H)-Furanone	9.84	0.04	16
79	2-Hydroxy-gamma-butyrolactone	12.08	0.23	25
80	2,5-Dimethyl-4-hydroxy-3(2H)-furanone	14.10	0.19	32
81	1,2-Ethanediol, 1-(2-furanyl)	19.10	0.03	46
82	5-Hydroxymethylfurfural	19.45	0.36	48
83	3-Deoxy-d-mannonic lactone	28.92	0.52	65
Nitrogen containing compounds				
84	N,N-Dimethylaminoethanol	5.22	0.12	1
85	1,3,5-Triazine-2,4,6-triamine	14.78	0.68	34
86	Acetic acid, 2-(N-methyl-N-phosphonomethyl)amino-	15.62	0.11	35
87	1-Heptadecanamine	18.67	0.07	44
88	Nonanamide	23.03	0.02	52
89	Dodecanamide (Lauryl amide)	33.33	0.04	71
90	Tetradecanamide (Myristic amide)	37.72	0.44	87
91	Hexadecanamide (Palmitic amide)	41.72	1.26	99
92	Docosenamide	44.99	6.04	106

93	Nonadecanamide	45.35	0.32	108
	<i>Sulfur containing compounds</i>			
94	Sulfurous acid, cyclohexylmethyl hexadecyl ester	23.17	0.67	53
	<i>Hydrocarbons</i>			
95	1-Butene, 4,4-diethoxy-2-methyl-	9.43	0.07	14
96	1-Methyl-2-octylcyclopropane	12.17	0.07	26
97	2-Trifluoroacetoxytridecane	13.92	0.03	31
98	trans-2,3-Epoxy-nonane	23.64	0.03	55
99	2-Heptafluorobutyroxypentadecane	23.89	0.04	56
100	1,2-Epoxyundecane	24.02	0.02	57
101	Heptacosane	24.36	0.03	59
102	1-Heptadecene	29.26	0.33	66
103	Octadecane, 1,1'-[(1-methyl-1,2-ethanediyl)bis(oxy)]bis-	29.43	0.01	67
104	1-Nonadecene	33.85	0.22	73
	<i>Pyrans</i>			
105	Tetrahydro-4H-pyran-4-ol	17.07	0.03	38
106	4H-Pyran-4-one, 2,3-dihydro-3,5-dihydroxy-6-methyl-	17.20	0.02	39
	<i>Others</i>			
107	5,6-Dihydroxypiperazine-2,3-dione dioxime	8.32	0.02	11
108	Tetraethyl silicate	11.81	0.04	23
109	Silanediol, dimethyl-, diacetate	17.30	0.07	40
110	D-Allose	26.61	0.19	62
111	Phenol, 2,4-bis(1,1-dimethylethyl)-	27.43	0.51	63
112	Oxirane, hexadecyl-	34.90	0.02	75
113	Ethyl iso-allocholate	36.94	0.06	81
114	Oleic anhydride	43.70	3.96	103

* RT: Retention time in minutes.

3.3. Anti-proliferative activities of MOSs

The anti-proliferative effectiveness of the MOSs extracts were examined on the colon cancer cells via the MTT assay. After 48 hrs of treatments, the cell viability status was examined in both normal (HEK-293) and cancerous (HCT-116) cells. The percentage of cell viability and inhibitory concentration (IC₅₀) values of both normal and treated cells were calculated. The MTT assay results after 48 hrs of treatment revealed the cytotoxic effects of the MOSs extracts, indicating their inhibitory action on the HCT-116 (cancerous) cells. The average of cell viability data showed (Fig 5) that the treatment using MOSs extracts

could induce a significant decrease in the cancer cells viability compared to the control cells (without treatment using MOSs extracts).

Fig 5. Impact of *Moringa oleifera* seed extracts on the colon cancer cells. The HCT-116 treated with moringa seed extracts for 48 hrs with different concentrations showing the average cell viability done by MTT assay. (**p < 0.01)

The specificity of the MOSs extracts (at different concentrations ranged from 0.03 to 0.1 mg/ml) on the normal and healthy (HEK-293) cells were inspected after 48 hrs of treatment using the MTT assays (Fig 6). The results showed that the inhibitory activity of the MOSs extracts (irrespective of their concentration) on the normal cells was insignificant, thereby suggesting the specificity of the MOSs extract towards the colon cancer cells only.

Fig 6. Impact of *Moringa oleifera* seed extracts on HEK-293 which were treated with moringa seed extracts for 48h showing the average cell viability done by MTT assay. The cells were treated with different concentrations (0.03 to 0.1 mg/ml) and. Data are the means $\pm$ SD of three different experiments. Difference between two treatment groups were analysed by student's t test

3.4. Nuclear disintegration due to treatment of Moringa seed extracts

The morphologies of the untreated and MOSs extracts treated cells were imaged using the CSM as shown in Fig 7. The cancerous cells (HCT-116) treated with MOSs extracts exhibited strong inhibitory action (Fig 7(b)) than the control sample (Fig 7(a)).

Fig 7. Nuclear staining by DAPI. The cancerous cells HCT-116 cells. (A) Control (nontreated) (B) treated with seed extracts of *Moringa oleifera* (0.066 mg/mL). DAPI stained cells visualized through confocal microscope and figure B show significant loss (arrows) of staining due to treatment. 200x magnifications.

It was inferred that the observed strong anti-cancerous activities of the MOSs extracts on the colon cancer cells may be due to the presence of high level of oleic acid and fatty acid in the extracts as supported by the GC-MS analysis. Yet, only few studies have been performed on the MOSs extracts indicating the enhanced anti-cancerous potency of such extracts correlated to the occurrences of high oleic acid and fatty acid contents [50-52].

3.5. Anti-microbial efficacy of MOSs extracts

The Agar well diffusion assay was used to determine the anti-microbial effectiveness of the proposed MOSs extracts where the inhibited areas around the inoculated wells were measured. This zone of inhibition was caused by the diffusion of the active chemical constituents of the MOSs extract around

the inoculated wells. The results revealed the considerable impact of the MOSs extracts seed on the bacterial strains (both gram-positive and gram-negative). However, the anti-bacterial action of the MOSs extracts was better against the gram-positive *S. aureus* bacteria. Fig 8 shows the MOSs extracts concentration dependent inhibition zones of the *S. aureus* bacteria (ranged from 18 to 24 mm). Conversely, the inhibition zones of the *E. coli* were ranged from 6 to 20 mm wherein the maximum and minimum inhibition zone diameter was obtained with the extracts concentration of 250 and 50 mg/ml, respectively (Fig 9). The observed discrepancy in the bactericidal action of the MOSs extracts for two different types of test bacteria may be due to the occurrence of their varying cell components [42].

Fig 8. Agar well diffusion plates showing the zone of inhibition. (A) *S. aureus* (B) *E. coli*. 1: 50 mg/ml, 2: 100 g/ml, 3: 150 mg/ml, 4: 200 mg/ml, 5: 250 mg/ml of seed extract of *Moringa oleifera*.

Fig 9. Zone of inhibition in millimeters (mm) against *S. aureus* and *E. coli* using the different concentrations of *Moringa oleifera* seed extract.

4. Conclusions

The LIBS and GC-MS technique was used to identify and quantify the elemental compositions of the MOSs. The anti-cancerous and anti-microbial effectiveness of the MOSs extracts were evaluated for the first time. The LIBS spectra revealed the presence of diverse elements in the MOSs useful for many health benefits. The GC-MS analysis reconfirmed the presence of several anti-cancerous and anti-microbial compounds in the MOSs extracts. The cell viability and DNA nuclear morphologies of the HEK-293 and HCT-116 cells treated with the MOSs extracts were measured using the MTT assay and DAPI staining, respectively. For the anti-cancer efficacy evaluation the HCT-116 were treated with MOSs extracts for 48 hrs before the test. The viability of the HEK-293 cells was examined after treating those using MOSs extracts. The cell viability percentage and IC_{50} values for both normal and cancerous cells were assessed. The MTT assay showed a significant impact of the MOSs extract for inhibiting the growth of the HCT-116 cells and insignificant inhibitory action of such extracts on the HEK-293 cells, indicating

the excellent specificity of the extracts towards cancer cells. The MOSs extracts showed strong anti-microbial activity in terms of the growth inhibition when tested with *S. aureus* and *E. coli* bacteria using Agar well diffusion assay. It is established that MOSs extract can be a prospective anti-cancer and anti-microbial agent for the functional biomedical drug formulation.

Availability of data and material: The data generated during the current study are available from the corresponding author upon request.

Author Declaration: We wish to confirm that there are no known conflicts of interest associated with this publication and there has been no significant financial support for this work that could have influenced its outcome.

Authors' Consent: All authors have approved the final version of the manuscript.

Acknowledgements: Authors are thankful to the Deanship of Scientific Research (DSR) King Fahd University of Petroleum and Minerals and Institute for Research & Medical Consultations (IMRC), Imam Abdulrahman Bin Faisal University, Dammam, Saudi Arabia for providing the financial assistance [Project number 2019-015-IRMC] in carrying out the experiments. We also thank the help of Mr. Dionezio Jr. Bagon Dela Roca to carry out cell culture and bioassay. Authors are also grateful to Dr. Khaldoun M. Alsamman, Clinical Laboratory Science, College of Applied Medical Science, Imam Abdulrahman Bin Faisal University for providing HCT-116 and HEK-293 cell lines.

References

- 379 1. Khor K.Z., Lim V., Moses E.J., Samad N.A., The in vitro and in vivo anticancer properties of Moringa
380 oleifera. Evid. Based Complement. Alternat. Med. 2018;2018:1-14.
- 381 2. Al-Zahrani A.A., Ibrahim A.H., Changes in 2S albumin gene expression in Moringa oleifera under
382 drought stress and expected allergenic reactivity in silico analysis. Theor. Exp. Plant Physiol.
383 2018;30:19–27.
- 384 3. Gandji K., Chadare F.J., Idohou R., Salako V.K., Assogbadjo A.E., Kakaï R.L.G., Status and
385 utilization of Moringa oleifera Lam: A review. Afr. Crop Sci. J. 2018;26:137–156.
- 386 4. Oduro I., Ellis W.O., Owusu D., Nutritional potential of two leafy vegetables: Moringa oleifera and
387 Ipomoea batatas leaves. Sci. Res. Essays. 2008 ;3:57–60.
- 388 5. Lin M., Zhang J., Chen X., Bioactive flavonoids in Moringa oleifera and their health-promoting
389 properties. J. Funct. Foods. 2018;47:469–479.
- 390 6. Amao A.O., Echeckwu C.A., Aba D.A., Katung M.D., Odeseye A.O., Diversity study of Drumstick
391 (Moringa oleifera Lam.) using Microsatellite markers. Int. J. Environ. Agric. Biotech.
392 2017;2:2380–2386.
- 393 7. Stohs S.J., Hartman M.J., Review of the safety and efficacy of Moringa oleifera. Phytother. Res.
394 2015;29:796–804.
- 395 8. Yalew T.A., Mekonnen Y., Retta N., Comparison of total phenolic content, free radical scavenging
396 potential and antihyperglycemic condition from leaves extract of Moringa stenopetala and
397 Moringa oleifera. Ethio. J. Pub. Health Nutr 2019;1:20–27.
- 398 9. Sandeep G., Anitha T., Vijayalatha K.R., Sadasakthi A., Moringa for nutritional security (Moringa
399 oleifera Lam.) Int. J. Bot. Stud 2019;4: 21–24.

- 400 10. Arévalo-Hijar L., Aguilar-Luis M.Á., Caballero-García S., Gonzáles-Soto N., Del Valle-Mendoza
401 J. ,Antibacterial and cytotoxic effects of *Moringa oleifera* (Moringa) and *Azadirachta indica*
402 (Neem) methanolic extracts against strains of *Enterococcus faecalis*. *Int. J. Dent.* 2018;3:1-5.
- 403 11. Abd El-Hack M.E., Alagawany M., Elrys A.S., Desoky E. S.M., Tolba H.M.N., Elnahal A.S.M, et
404 al. Effect of forage *Moringa oleifera* L. (Moringa) on animal health and nutrition and its beneficial
405 applications in soil, plants and water purification. *Agriculture.* 2018;145:1-22.
- 406 12. Uphadek B., Shinkar D.M., Patil P.B., Saudagar R.B., *Moringa oleifera* as a pharmaceutical
407 excipient. *Int. J. Curr. Pharm. Res.* 2018;10:13–16.
- 408 13. Swati S., Virk A.K., Kumari C., Ali A., Garg P., Thakur P. et al. *Moringa oleifera*—A never die
409 tree: An overview. *Asian J. Pharm. Clin. Res.* 2018 ;11:57–65.
- 410 14. Sultana S., Akram M., Shah S.M.A., Ayaz S., Rehman M.R.U., Ahmad H. et al. *Moringa oleifera*
411 Lam. A miraculous medicinal plant: Review. *Pak. J. Med. Biol. Sci.* 2018;2:60–63.
- 412 15. Ma Z.F., Ahmad J., Zhang H., Khan I., Muhammad S., Evaluation of phytochemical and medicinal
413 properties of *Moringa* (*Moringa oleifera*) as a potential functional food. *S. Afr. J. Bot.* 2019;
414 <https://doi.org/10.1016/j.sajb.2018.12.002>.
- 415 16. Bezerra K., Silva N.G., de Souza A., Sá G., de Araújo F., *Moringa oleifera*: A new perspective for
416 the synthesis of natural products. *Mol2net.* 2016 ;2:1–5.
- 417 17. Ali A., Akhtar N., Chowdhary F., Enhancement of human skin facial revitalization by *Moringa* leaf
418 extract cream. *Postep. Derm. Alergol.* 2014;31:71–76.

419

- 420 18. Manzoor M., Anwar F., Iqbal T., Bhanger M.I., Physico-chemical characterization of Moringa
421 concanensis seeds and seed oil. J. Amer. Oil Chem. Soc. 2007;84:413–419.
- 422 19. Sanchez-Machado D.I., Nunez-Gastelum J.A., Reyes-Moreno C., Ramirez-Wong B., Lopez-
423 Cervantes J. , Nutritional Quality of edible Parts of Moringa oleifera. Food Anal. Methods. 2010;
424 3:175–180.
- 425 20. Hermawan A., Nur K.A., Sarmoko Dewi D., Putri P., Meiyanto E., Ethanolic extract of Moringa
426 oleifera increased cytotoxic effect of doxorubicinon HeLa cancer cells. J. Nat. Remedies
427 2012 ;12:108–114.
- 428 21. Rodríguez-Pérez C, Mendiola J., Quirantes-Piné R, Ibáñez E, Segura-Carretero A., Green
429 downstream processing using supercritical carbon dioxide, CO₂-expanded ethanol and pressurized
430 hot water extractions for recovering bioactive compounds from Moringa oleifera leaves. J
431 Supercrit Fluids. 2016;116: 90–100.
- 432 22.Da Porto C, Decorti D, Natolino A., Microwave pretreatment of Moringa oleifera seed: effect on oil
433 obtained by pilot-scale supercritical carbon dioxide extraction and Soxhlet apparatus. J Supercrit
434 Fluids. 2016;107:38–43.
- 435 23. Zhong J, Wang Y, Yang R, Liu X, Yang Q, Qin X. , The application of ultrasound and microwave
436 to increase oil extraction from *Moringa oleifera* seeds. Ind Crop Prod. 2018;120:1–10.
- 437 24. Rai, A, Mohanty, B, Bhargava (Late) , R, 2017. Experimental modeling and simulation of
438 supercritical fluid extraction of Moringa oleifera seed oil by carbon dioxide. Chem Eng Commun.
439 204,957–964.
- 440 25. Belo Y., Al-Hamimi S, Chimuka L, Turner C., Ultrahigh-pressure supercritical fluid extraction and
441 chromatography of Moringa oleifera and Moringa peregrina seed lipids. Anal Bioanal Chem.
442 2019;411:3685-3693.

26. Almessiere M.A., Altuwiriqi R., Gondal M.A., AlDakheel R.K., Alotaibi H.F., Qualitative and quantitative analysis of human nails to find correlation between nutrients and vitamin D deficiency using LIBS and ICP-AES. *Talanta*. 2018;185:61–70.
27. Gondal M.A., Siddiqui M.N., Nasr M.M., Detection of Trace Metals in Asphaltenes Using an Advanced Laser-Induced Breakdown Spectroscopy (LIBS) Technique. *Energy & Fuels*. 2010 ; 24 :1099–1105.
28. Corsi M., Cristoforetti G., Hidalgo M., Legnaioli S., Palleschi V., Salvetti A et al. Application of laser-induced breakdown spectroscopy technique to hair tissue mineral analysis. *Applied Optics*. 2003;42: 6133-6137.
29. Bonta M., Gonzalez J.J., Quarles C.D., Russo R.E., Hegedus B., Limbeck A., Elemental mapping of biological samples by the combined use of LIBS and LA-ICP-MS. *Anal Atom Spectrom*. 2016; 31:252–258.
30. Chen X., Li X., Yang S., Yu X., Liu A., Discrimination of lymphoma using laser-induced breakdown spectroscopy conducted on whole blood samples. *Biomedical Optics Express*. 2018 ;9:1057–1068.
31. Mehari F., Rohde M., Knipfer C., Kanawade R., Klämpfl F., Adler W. et al. 2014. Laser induced breakdown spectroscopy for bone and cartilage differentiation - ex vivo study as a prospect for a laser surgery feedback mechanism. 2018;5:4013–23.
32. Anjula P., Pradheep K., Rita G., Eroshini N., Bhandari D.C. , ‘Drumstick tree’ (*Moringa oleifera* Lam.): A multipurpose potential species in India. *Genet. Resour. Crop Evol*. 2011 ;58:453–460.
33. Pasquini C, Cortez J, Silva L., Gonzaga F., Laser induced breakdown spectroscopy. *Braz Chem Soc*. 2007;18:463–512.

- 465 34. Rai A.K., Yueh F.Y., Singh J.P. , Rai D.K. , Laser induced breakdown spectroscopy of solid and
466 molten materials. Elsevier Science, B.V. Chemistry and Chemical Engineering, The Netherlands,
467 2007.pp. 255–285.
- 468 35. Rai P.K., Jaiswal D., Diwaker S., Watal G. , Antihyperglycemic profile of *Trichosanthes dioica*
469 seeds in experimental models. *Pharm Biol* 2008;46:1–6.
- 470 36. Khan F.A., Akhtar S., Almoft, S.A., Almohazey D., Alomari M., FMSP-Nanoparticles Induced Cell
471 Death on Human Breast Adenocarcinoma Cell Line (MCF-7 Cells): Morphometric Analysis.
472 *Biomolecules* 2018;8:1-13.
- 473 37. Khan F.A., Akhtar S., Almohazey D., Alomari M., Almoft S.A., Badr I. et al. Targeted delivery of
474 poly (methyl methacrylate) particles in colon cancer cells selectively attenuates cancer cell
475 proliferation. *Artif. Cells Nanomed. Biotechnol.* 2019;47:1533-1542.
- 476 38. Rehman S., Asiri M., Khan F. A., Rabindran J., Khan H., Akhtar S. et al. Biocompatible Tin Oxide
477 Nanoparticles: Synthesis, Antibacterial, Anticandidal and Cytotoxic Activities. *ChemistrySelect*
478 2019;4:4013-17.
- 479 39. Sabsabi M., Cielo P., Quantitative Analysis of Aluminum Alloys by Laser-Induced Breakdown
480 Spectroscopy and Plasma Characterization . *Appl. Spectrosc.* 1995;49:499–507 .
- 481 40. Mehta S., Rai P. K., Rai N. K., Rai A. K., Bicanic D., Watal G., Role of Spectral Studies in
482 Detection of Antibacterial Phytoelements and Phytochemicals of *Moringa oleifera*. *Food Biophys.*
483 2011;6:497–502.

484

- 485 41. Sharma S., Shukla N., Bharti A. S., Uttam K. N., Simultaneous Multielemental Analysis of the Leaf
486 of *Moringa oleifera* by Direct Current Arc Optical Emission Spectroscopy. Natl. Acad. Sci. Lett.
487 2018;41:65–68.
- 488 42. Pasha S. N., Shafi K. M., Joshi A.G. , Meenakshi I. , Harini K., Mahita J, et al. The transcriptome
489 enables the identification of candidate genes behind medicinal value of Drumstick tree (*Moringa*
490 *oleifera*). Genomics 2020;112:621-628.
- 491 43. Akinmoladun A.C., Ibukun E., Afor E., Akinrinlola B., Onibon T., Akinboboye A, et al. Chemical
492 constituents and antioxidant activity of *Alstonia boonei*. Afr. J. Biotechnol. 2007;6: 1197-1201.
- 493 44. Yamaguchi F., Takata M., Kamitori K., Nonaka M., Dong Y., Sui L, et al. Rare sugar D-allose
494 induces specific up-regulation of TXNIP and subsequent G1 cell cycle arrest in hepatocellular
495 carcinoma cells by stabilization of p27kip1. Int. J. Oncol. 2008;32:377-385.
- 496 45. Al-Asmari A. K., Albalawi S. M., Athar M. T. , Khan A. Q., Al-Shahrani H., Islam M., *Moringa*
497 *oleifera* as an anti-cancer agent against breast and colorectal cancer cell lines. PLoS One 2015;10:
498 0135814-0135814.
- 499 46. Ogunlesi M., Okiei W., Osibote E., Analysis of the essential oil from the leaves of *Sesamum radiatum*,
500 a potential medication for male infertility factor, by gas chromatography-mass spectrometry. Afr.
501 J. Biotechnol 2010;9:1060-1067.
- 502 47. De Silva P.S., Luben R., Shrestha S.S., Khaw K.T., Hart A. R., hepatology, Dietary arachidonic and
503 oleic acid intake in ulcerative colitis etiology: a prospective cohort study using 7-day food diaries.
504 Eur. J. Gastroenterol. Hepatol. 2014;26:11-18.
- 505 48. Ruz-gutierrez V., Muriana F J, Guerrero A., Cert A M, Villar J, Plasma lipids, erythrocyte membrane
506 lipids and blood pressure of hypertensive women after ingestion of dietary oleic acid from two
507 different sources. Int. J. Hypertens. 1996;14:1483-1490.

- 508 49. Haug A., Høstmark A.T., Harstad O.M., Bovine milk in human nutrition—a review. *Lipids Health Dis*
509 2007;6:1-16.
- 510 50. Menendez J.A., Lupu R., Mediterranean dietary traditions for the molecular treatment of human
511 cancer: anti-oncogenic actions of the main olive oil's monounsaturated fatty acid oleic acid (18:1n-
512 9). *Curr Pharm Biotechnol*. 2006;7:495–502.
- 513 51. Fernández E., Gallus S., Vecchia C. L., Nutrition and cancer risk: an overview. *J Br Menopause Soc*.
514 2006;12:139–142.
- 515 52. Menendez J.A., Papadimitropoulou A., Vellon L., Lupu R., A genomic explanation connecting
516 "Mediterranean diet", olive oil and cancer: oleic acid, the main monounsaturated fatty acid of olive
517 oil, induces formation of inhibitory "PEA3 transcription factor-PEA3 DNA binding site"
518 complexes at the Her-2/neu (erbB-2) oncogene promoter in breast, ovarian and stomach cancer
519 cells. *Eur J Cancer*. 2006;42:2425–2432.