Title

Areas of global importance for terrestrial biodiversity, carbon, and water

Martin Jung¹, Andy Arnell², Xavier de Lamo², Shaenandhoa García-Rangel², Matthew Lewis³, Jennifer Mark², Cory Merow⁴, Lera Miles², Ian Ondo⁵, Samuel Pironon⁵, Corinna Ravilious², Malin Rivers⁶, Dmitry Schepashenko^{1 7}, Oliver Tallowin², Arnout van Soesbergen², Rafaël Govaerts⁵, Bradley L. Boyle⁸, Brian J. Enquist⁸, Xiao Feng^{8 9}, Rachael V. Gallagher¹⁰, Brian Maitner⁸, Shai Meiri¹¹, Mark Mulligan¹², Gali Ofer¹¹, Jeffrey O. Hanson¹³, Walter Jetz^{14 15}, Moreno Di Marco¹⁶, Jennifer McGowan¹⁷, D. Scott Rinnan^{14 15}, Jeffrey D. Sachs¹⁸, Myroslava Lesiv¹, Vanessa Adams¹⁹, Samuel C. Andrew²⁰, Joseph R. Burger⁹, Lee Hannah²¹, Pablo A. Marquet^{22 23 24 25 26}, James K. McCarthy²⁷, Naia Morueta-Holme²⁸, Erica A. Newman⁸, Daniel S. Park²⁹, Patrick R. Roehrdanz²¹, Jens-Christian Svenning^{30 31}, Cyrille Violle³², Jan J. Wieringa³³, Graham Wynne³⁴, Steffen Fritz¹, Bernardo B.N. Strassburg^{35 36 37 38}, Michael Obersteiner^{1 39}, Valerie Kapos², Neil Burgess², Guido Schmidt-Traub⁴⁰ and Piero Visconti¹

Affiliations

- 1 Ecosystems Services and Management Program (ESM), International Institute for Applied Systems Analysis (IIASA), Schlossplatz 1, A-2361 Laxenburg, Austria
- 2 UN Environment Programme World Conservation Monitoring Centre (UNEP-WCMC), 219 Huntingdon Road, Cambridge CB3 0DL, United Kingdom
- 3 Department of Zoology, University of Cambridge, Downing Street, Cambridge CB2 3EJ, United Kingdom
- 4 Department of Ecology and Evolutionary Biology, University of Connecticut, CT 06269, USA
- 5 Royal Botanic Gardens, Kew, Richmond FPH3+FQ, United Kingdom
- 6 Botanic Gardens Conservation International, Richmond, Surrey TW9 3BW, United Kingdom
- 7 Siberian Federal University, Krasnoyarsk 660041, Russia
- 8 Department of Ecology and Evolutionary Biology, University of Arizona, Tucson, AZ 85721, USA.
- 9 Institute of the Environment, University of Arizona, Tucson, AZ 85721, USA
- 10 Department of Biological Sciences, Macquarie University, North Ryde, NSW 2019, Australia
- 11 School of Zoology, Faculty of Life Sciences, Tel Aviv University, Tel Aviv 6997801, Israel
- 12 Department of Geography, King's College London, London WC2B 4BG, United Kingdom
- 13 CIBIO/InBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos da Universidade do Porto, 4485-661 Vairão, Portugal
- 14 Department of Ecology and Evolutionary Biology, Yale University, New Haven, CT, 06520, USA
- 15 Center for Biodiversity and Global Change, Yale University, New Haven, CT, 06520, USA
- 16 Department of Biology and Biotechnologies, Sapienza University of Rome, viale dell'Università 32, 6 I-00185 Rome, Italy
- 17 The Nature Conservancy, 4245 Fairfax Drive, Arlington, VA, 22203, USA
- 18 Columbia University, New York, NY 10027, USA
- 19 Discipline of Geography and Spatial Sciences, University of Tasmania, Hobart TAS 7005, Australia
- 20 CSIRO Land and Water, Canberra, Australian Capital Territory, Australia
- 21 Betty and Gordon Moore Center for Science, Conservation International, 2011 Crystal Dr., Arlington, VA 22202, USA.
- 22 Departamento de Ecología, Facultad de Ciencias Biológicas, Pontificia Universidad Católica de Chile, CP 8331150, Santiago, Chile
- 23 Instituto de Ecología y Biodiversidad (IEB), Santiago, Chile
- 24 Laboratorio Internacional en Cambio Global (LINCGlobal) y Centro de Cambio Global UC, Facultad de Ciencias Biológicas, Pontificia Universidad Católica de Chile, CP 8331150, Santiago, Chile
- 25 The Santa Fe Institute, 1399 Hyde Park Road, Santa Fe NM 87501, USA
- 26 Instituto de Sistemas Complejos de Valparaíso (ISCV), Artillería 470, Valparaíso
- 27 Manaaki Whenua – Landcare Research, Lincoln 7640, New Zealand
- 28 Center for Macroecology, Evolution and Climate, GLOBE Institute, University of Copenhagen, Universitetsparken 15, build. 3, DK-2100 Copenhagen Ø, Denmark
- 29 Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, Massachusetts, 02138 USA
- 30 Center for Biodiversity Dynamics in a Changing World (BIOCHANGE), Department of Biology, Aarhus University, Ny Munkegade 114, DK-8000 Aarhus C, Denmark
- 31 Section for Ecoinformatics and Biodiversity, Department of Biology, Aarhus University, Ny Munkegade 114, DK-8000 Aarhus C, Denmark
- 32 CEFE, Univ. Montpellier, CNRS, EPHE, IRD, Univ. Paul Valéry Montpellier 3, Montpellier, France
- 33 Naturalis Biodiversity Center, Darwinweg 2, Leiden, The Netherlands
- 34 World Resources Institute

35 Rio Conservation and Sustainability Science Centre, Department of Geography and the Environment, Pontifical Catholic University, 2453900,
 36 Rio de Janeiro, Brazil
 37 Programa de Pós Graduação em Ecologia, Universidade Federal do Rio de Janeiro, 21941-590, Rio de Janeiro, Brazil
 38 Botanical Garden Research Institute of Rio de Janeiro, Rio de Janeiro, Brazil
 39 Environmental Change Institute Oxford University Centre for the Environment South Parks Road Oxford OX1 3QY
 40 Sustainable Development Solutions Network, Paris, France

Summary paragraph

To meet the ambitious objectives of biodiversity and climate conventions, countries and the international community require clarity on how these objectives can be operationalized spatially, and multiple targets be pursued concurrently¹. To support governments and political conventions, spatial guidance is needed to identify which areas should be managed for conservation to generate the greatest synergies between biodiversity and nature's contribution to people (NCP). Here we present results from a joint optimization that maximizes improvements in species conservation status, carbon retention and water provisioning and rank terrestrial conservation priorities globally. We found that, selecting the top-ranked 30% (respectively 50%) of areas would conserve 62.4% (86.8%) of the estimated total carbon stock and 67.8% (90.7%) of all clean water provisioning, in addition to improving the conservation status for 69.7% (83.8%) of all species considered. If priority was given to biodiversity only, managing 30% of optimally located land area for conservation may be sufficient to improve the conservation status of 86.3% of plant and vertebrate species on Earth. Our results provide a global baseline on where land could be managed for conservation. We discuss how such a spatial prioritisation framework can support the implementation of the biodiversity and climate conventions.

Introduction

Biodiversity and nature's contributions to people (NCP) are in peril, requiring an increasing level of ambition to avert further decline¹. Existing global biodiversity conservation targets are unlikely to be met by the end of 2020². Similarly, the world is falling short of mobilizing the full climate mitigation potential of nature-based climate solutions, estimated at around a third of mitigation effort under the Paris Agreement³. A new global biodiversity framework is scheduled to be adopted by the Convention on Biological Diversity (CBD) in Kunming, China, in October 2020⁴, and there are growing calls to integrate nature-based solutions into climate strategies⁵.

Targets for site-based conservation actions, hereafter area-based conservation targets, will likely remain important for the new global biodiversity framework⁴. Several calls have been made for such targets, including suggestions that at least 30% of land and oceans be protected for conservation and an additional 20% for climate mitigation⁶ and that the value of areas of global importance for conservation is maintained or restored⁷. The Sustainable Development Goals

(SDGs), the United Nations Framework Convention on Climate Change (UNFCCC) and the CBD emphasize that habitat conservation and restoration should contribute simultaneously to biodiversity conservation and climate change mitigation⁴. Recent analyses of conservation priorities for biodiversity and carbon have spatially overlaid areas of importance for both assets, effectively treating the two goals as to be pursued separately (e.g.^{6,9}). However, multi-criteria spatial optimization approaches applied to conservation and restoration prioritisation have shown that carbon sequestration could be doubled, and the number of extinctions prevented tripled, if priority areas were jointly identified rather than independently^{10,11}. Yet, no comparable optimization analyses exist at a global scale.

A number of recent studies have attempted to map spatial conservation priorities on land¹², relying on spatial conservation prioritisation (SCP) methods^{13–1617}. However, these approaches are limited, in that: they (i) are limited by geographic extent²² or focus on only a subset of global biodiversity, notably ignoring either reptiles or plant species, which show considerable variation in areas of importance compared to other taxa^{18,19}; (ii) focus on species representation only, rather than reducing extinction risk, as per international biodiversity targets, and often ignore other dimensions of biodiversity, e.g. evolutionary distinctiveness^{20,21}; (iii) do not investigate the extent to which synergies between biodiversity and NCPs, such as carbon sequestration or clean water provisioning²², can be maximised²¹; and (iv) they use a-priori defined, and subjective measures of importance, such as intactness^{8,17}, or area-based conservation targets, such as 30% or 50% of the Earth^{6,24} instead of objectively delineating the relative importance of biodiversity and NCPs across the whole world irrespective of such constraints.

The aim of this study is to identify the most important areas for biodiversity - here focussing on species conservation - as well as NCPs including carbon storage and water provisioning, to be managed for conservation globally. We define managing an area for conservation as any site-based action that is appropriate for the local context (considering pressures, tenure, land-use, etc.), and that is commensurate with retaining or restoring the desirable assets (e.g. species, habitat types, soil or biomass carbon, clean water). This management may sometimes require legal protection to be effective, but not necessarily in the form of protected areas.

We obtained fine-scale distribution maps for the world's terrestrial vertebrates as well as the largest sample of plant distribution data ever considered in global species-level analysis, ~41% of all accepted species names in this group. As NCPs we use the latest global spatial data on above- and below-ground biomass carbon, and vulnerable soil carbon, as well as the volume of potential clean water by river basin. We applied a multicriteria spatial optimization framework to investigate synergies between these assets and explore how priority ranks change depending on how much weight is given to either carbon sequestration, water provisioning or biodiversity, and examined whether priorities vary if species evolutionary distinctiveness and threat status are considered.

Results

We found large potential synergies between managing land for biodiversity conservation, storing soil and biomass carbon, and maintaining clean water provisioning. Managing the top-ranked 10% of land, i.e. those areas with the highest priority, to achieve these objectives simultaneously (Fig.

1, SI Fig. 1), has the potential to improve the conservation status of 46.1% of all species considered, of which 51.1% are plant species, as well as conserve 27.1% of the total carbon and 24.1% of the potential clean water globally. Areas of biodiversity importance notably include mountain ranges of the world, large parts of Mediterranean biomes and South-East Asia (SI Fig. 2) and were overall mostly comparable to previous expert-based delineations of conservation hotspots¹⁶, while also highlighting additional areas of importance for biodiversity only, such as the West African Coast, Papua New-Guinea and East Australian Rainforest (SI Fig. 2). The Hudson Bay area, the Congo Basin and Papua New Guinea were among the top-ranked 10% areas for global carbon storage (SI Fig. 3a), while the Eastern United States of America, the Congo, European Russia and Eastern India were among the areas with the greatest importance for clean water provisioning (SI Fig. 3b). Overall, top-ranked areas of joint importance of biodiversity, carbon and water were spatially distributed across all continents, latitudes and biomes.

Fig. 1: Global areas of importance for terrestrial biodiversity, carbon and water. All assets were jointly optimized with equal weighting given to each asset (central point in the series of segments in Fig. 2) and ranked by the most (1-10%) to least (90-100%) important areas to conserve globally. The triangle plot shows the extent to which protecting the top-ranked 10% and 30% of land (dark brown and yellow areas on the map) contributes to improving species conservation status, storing carbon and providing clean water. The map is at 10 km resolution in Mollweide projection. A map highlighting the uncertainty in priority ranks can be found in SI Fig 1.

Synergies and trade-offs depend on the relative importance given to conservation of terrestrial biodiversity, carbon storage and water provisioning (Fig. 2a). We explored an array of conservation scenarios each with a range of possible outcomes: at one extreme, priority is given to conserving biodiversity and carbon only, and with equal weight (Fig. 2b). At the other extreme are scenarios that prioritize conserving only biodiversity and water (Fig. 2c). Intermediate options include giving equal weighting to all three assets (Fig. 1). Similar to earlier assessments^{9,26,27}, we found synergies between the conservation of biodiversity and carbon storage (Fig. 2b). However

we also discovered similar synergies for biodiversity and water provisioning (Fig. 2c). Conserving the top-ranked 10% of land for biodiversity and carbon can only protect up to 23.6% of the global total carbon and 45.8% of all species (Fig. 2a), while maintaining 17.8% of all global water provisioning as co-benefit (Fig. 2b). In contrast, conserving the top-ranked 10% of land for biodiversity and water only can protect 21.7% of water and 43.6% of all species (Fig. 2a), while maintaining 18% as carbon co-benefit (Fig. 2c). The implications of assigning different relative preferences to conserving NCPs magnify with increasing amounts of land dedicated to conservation. For example, with 10% and 30% of land managed for conservation the range of carbon conserved is 18% to 23.6% and 49.2% to 63.1% respectively, and the range in water conserved is 17.8% to 21.7% and 51.8% to 66.4% (Fig. 2a). Our results suggest that there is ample scope for identifying co-benefits from conserving these three assets, if explicit targets for each are considered, areas of importance for each asset are identified through multi-criteria optimization, and the range of relative weights given to each asset is comprehensively explored.

Fig. 2: Implications of different relative weights given to carbon or water over improving species conservation status. (a) Each 'boomerang-shaped' segment of dots represents a series of conservation prioritisation scenarios with a common area budget (from 10% of land bottom left to 100% at top-right). Axes indicate the proportion of all carbon and water provisioning assets conserved, colours represent the proportion of species for which conservation status could be improved in a given conservation scenario, and the point size indicates the difference in weighting given to carbon or water relative to biodiversity, ranking from none to equal weighting. (b-c) Global areas of importance if 10% (dark-brown), or 30% (yellow), of land area is managed for conservation while preferring (b) carbon protection over water or (c) water protection over carbon.

The amount of land necessary to exclusively protect global biodiversity continues to be debated^{15,28,29} In our analysis we found that, in the absence of any socio-economic constraints and ignoring other NCPs (here water and carbon), at least ~67% of land needs to be managed for

conservation globally, to improve the conservation status for terrestrial plants and vertebrates (Fig. 3a). This is robust to the number of species included in the analyses, provided that they are a representative subset (see methods), with the variation typically being ~0.1% around the mean accumulation curves (Fig. 3a).

Optimally placing areas managed for conservation on 30% of the world's land is already sufficient to conserve 86.3% of all species considered in this analysis (ignoring existing protected areas, socio-economic constraints and other NCPs). Currently protected areas (PAs) are potentially sufficient to achieve persistence targets for 16.3% of the species analysed (SI Fig. 5, SI Fig. 6). However, by building on the current PA estate to increase areas managed for biodiversity conservation up to 30% of land, the conservation status of an additional 60.8% of the species could be improved (for a total of 77.1% of the species analysed). Therefore, there is an efficiency gap of only ~9.2% between re-designing global conservation efforts and optimally building on existing efforts.

When jointly optimizing target achievement for biodiversity, carbon and water (Fig. 3a), we found that selecting the top-ranked 30% (respectively 50%) of areas, a popular proposal for area-based conservation targets⁶, would conserve 62.4% (86.8%) of the estimated total carbon stock and 67.8% (90.7%) of all clean water provisioning, in addition to improving the conservation status for 69.7% (83.8%) of all species considered.

When optimizing conservation efforts for biodiversity only, we found that the groups that benefited the most were amphibian and plant species (Fig. 3b) and threatened species (Fig. 3c). The latter tend to have smaller range sizes and smaller absolute area targets than other groups and are inherently prioritized with area budgets $\leq 30\%$ of land.

Fig. 3: Accumulation curves showing how the number of species targets met increases with amount of land optimally allocated to conservation. Confidence bounds of accumulation curves indicate the uncertainty among representative sets and were generally found to be very small (~0.1%). This analysis ignores current protected areas and a version including those areas can be

found in the SI Fig. 6. (a) Target accumulation curves for analysis variants including other assets; (b) for different taxonomic groups when optimizing biodiversity only to conservation; (c) for species classified by IUCN as threatened or not (see Methods) when optimizing for biodiversity only.

Our analysis included, for the first time in a global prioritisation analysis, a representative subset of plant distribution data totalling ~41% of described vascular plant species³² (Fig. 4). Incorporating data on plants resulted in spatial shifts in areas of importance for conservation, particularly in the western United States of America, West-Central and South Africa, South-West Australia, Central Brazil, as well as northern Europe and central Asian steppes and mountains compared to an analysis where plants are ignored (Fig. 4a). Overall we found montane and temperate grasslands, Mediterranean savannas and shrublands biomes to increase in importance when considering plants, whereas flooded grasslands and mangroves lost relative importance (Fig. 4b). The accumulation curves of species targets achieved were comparable between analysis variants with and without plants (Fig. 4c). Overall this indicates high surrogacy between vertebrate and plant species, despite spatial shifts in areas of importance (Fig. 4a).

Fig. 4: Change in global areas of biodiversity importance after adding plant species. (a) Calculated as the difference in areas of biodiversity importance with either plant species included or excluded. Positive changes (yellow to dark green) in rank imply an increase in priority if plant species are considered, while negative changes (light to dark blue) show a decrease in priority ranks. The map is at 10 km resolution in a Mollweide projection. (b) Average change in ranks per biome after plants have been added. (c) Representation curves of areas necessary to be managed for conservation with (solid) and without plants (dashed) included.

Areas of importance can vary spatially if species are given different weights, prioritising for instance the protection of threatened or more evolutionarily distinct species^{20,21}. We tested the implication of prioritising the improvement of conservation status for these groups of species by weighting them by current conservation status or evolutionary distinctiveness. We found that doing so has only small inefficiency implications compared to a prioritisation without these weights (0.7% fewer biodiversity targets achieved when prioritising threatened species and 1.7% fewer

when prioritising evolutionarily distinct species with 10% of land). Yet, overall spatial patterns of the top-ranked 10% of areas of importance were comparable, with only minor differences, notably highlighting the importance of New Zealand and the Brazilian Amazon for conserving threatened species, the Mediterranean Basin, North-West USA, Florida and fringes of the Amazon Basin for conserving evolutionarily distinct species (SI Fig. 10). These results highlight that threatened or more evolutionary distinct species are well covered by other species³⁰, and their full conservation can be achieved at minimal extra cost.

Discussion

How much area and where it should be managed for conservation is one of the key questions underpinning global biodiversity conventions and conservation planning discussions^{4,29}. Our analyses suggest that even ambitious objectives such as ‘Half Earth’²⁴ or ‘30 by 30’⁶ are insufficient to ensure that the conservation status of threatened species is improved and that non-threatened species remain so (Fig. 3). However, managing for conservation the top-ranked 30% of areas of importance for biodiversity, as identified here, can bring over 86% of the world's terrestrial vertebrate and a representative sample of plant species (of ~41% of all plant species) to a non-threatened conservation status, with further increases in area offering minor additional returns (Fig. 3). Depending on the level of political ambition, an extra 20% of land could be dedicated to carbon storage as a contribution to climate regulation⁶ and sustainable management of natural resources. However, our analysis shows that considerable co-benefits can already be achieved by managing an optimally placed 30% of land, if conservation of biodiversity, carbon and water is planned for with spatial optimization approaches (Fig. 2). We caution that these estimates, and equally those from previous studies^{6,14,16,23}, can vary with different data and methods applied.

We ranked priority areas in order of importance for conservation management; but we note that specific forms of management are highly contextual and will depend on local anthropogenic pressures, governance and opportunity costs. Areas of biodiversity importance that require strict protection and active management, e.g. where narrow-ranging and threatened species occur might be suitable for protected area expansion³¹. Other effective area-based conservation measures³², such as watersheds managed primarily for water resource management or community-managed forests, might be more suitable in areas where biodiversity, carbon and water benefits are high but threats to species conservation remain low.

Our analyses does not impose any constraint on feasibility or equity among countries³³, some of which contain over half of their territory in the top-ranked 10% of global importance for biodiversity, carbon and water provision (Fig. 1). Thus, there is a need for fair resourcing of the required management actions to offset the financial burden on some, predominantly tropical, countries^{33,34}. Existing funding mechanisms should further explore opportunities to synergistically benefit both biodiversity and NCPs, as has been shown in the case of carbon²⁶. Future, synergistic conservation prioritization efforts should particularly focus on incorporating socio-economic constraints³⁵, consider integrated scenarios of the projected distribution of biodiversity, carbon and

water, support countries in identifying conservation actions at finer scale to maximize the achievement of national and global targets.

Our work also reveals research and data gaps in determining global areas of importance for terrestrial biodiversity conservation and NCPs. As NCPs we choose carbon and water because of their relevance to international conventions, but there are others we did not consider²² such as food provisioning or cultural relevance. Similarly, many aspects of biodiversity remain under-represented - although we consider a significant portion of plant species on Earth, and we developed a framework to remove spatial bias in priority setting resulting from incomplete taxonomic coverage - there is a need to expand available data on other groups such as freshwater, soil and invertebrate species^{36,37}. We also only investigated the influence of evolutionary history on vertebrate, but not plant species, for whom hotspots of evolutionary history might differ, and ignored other dimensions such as functional rarity³⁸. Despite remaining gaps in taxonomic coverage and species checklists, our analysis also confirms the results of previous, broad-scale studies^{18,19,39} that found high congruence between vertebrate and plant areas of importance, but we also highlight areas that would be overlooked if plants were not considered, especially so in dry grasslands, savannahs and Mediterranean shrublands (Fig. 4).

Our analyses highlight global areas of conservation importance that can maximize synergies across conventions (e.g. CBD, UNFCCC) and the SDGs. Particularly, our integrated maps could support governments in translating set targets (such as area-based conservation measures proposed for the 2021-2030 Strategic Plan of the CBD⁴) into national policies and actions on the ground and demonstrate how integrated spatial planning can be used to assist national biodiversity strategies. Meeting the SDGs requires real, transformative commitments that are yet to be enacted¹, however, by maximizing synergies in efforts and resources, a pathway towards effective biodiversity conservation can be laid out for the next decade.

Methods

Biodiversity data

We utilized best available global species distribution data (overview in SI Table 1), including all extant terrestrial vertebrates and a representative proportion (41.31%) of all accepted plant species according to Plants of the World Online⁴⁰. Extant mammal (5,685 species) and amphibian (6,660) distribution data were obtained from the International Union for Conservation of Nature Red List database (IUCN ver. 2019_2⁴¹), while bird (10,953) range maps were obtained from Birdlife International⁴². Data on the distribution of reptiles were obtained from the IUCN database when available (6,830 species), otherwise from the Global Assessment of Reptile Distributions (GARD) database (3,755⁴³). We obtained native plant range maps (193,954 species) from a variety of sources, including IUCN, Botanic Gardens Conservation International (BGCI) and the Botanical Information and Ecology Network (BIEN). The IUCN and BGCI data contains expert-based range maps and alpha-hulls (see Supporting Information), while the BIEN data consists mainly of herbarium collections, ecological plots and surveys⁴⁴⁻⁵², that were used to construct conservative estimates of species ranges using species distribution models (SDMs). We benefited from version 4.1 of BIEN, which includes data from RAINBIO⁵³, TEAM⁵⁴, The Royal Botanical Garden of

Sydney, Australia, and NeoTropTree⁵⁵. Additional plant plot data from a number of networks and datasets have been included in BIEN and a full listing of the herbaria data used can be found in the extended acknowledgements and online (<http://bien.nceas.ucsb.edu/bien/data-contributors/all/>). In cases where multiple data sources were available for the same plant species, we preferentially used expert-based range maps to characterize a species' spatial distribution. A full description of the preparation and processing of the plant data can be found in the Supporting Information.

All vertebrate range maps were pre-processed following common practice⁵⁶ by selecting only those parts of a species' range where 1) it is extant or possibly extinct, 2) where it is native or reintroduced and 3) where the species is seasonally resident, breeding, non-breeding, migratory or where the seasonal occurrence is uncertain. We acknowledge that these ranges can contain some areas where the species is possibly extinct.

Suitable habitat refinement

Where data on species habitat and elevational preferences were available, we refined each species' range to obtain the area of habitat (AOH) in which the species could potentially persist^{57,58}. Data on species habitat preferences and suitable elevational range were obtained from the IUCN Red List database⁴¹ and, for an additional 1,452 reptile species in the GARD database, habitat preferences were compiled from an extensive literature search. For seasonally migrating birds and mammal species we ensured that separate habitat refinements were conducted for permanent and seasonally occupied areas of their range, that is, the breeding and non-breeding range. Whenever no habitat or elevation preferences were available for a given species, we used the full range except for areas considered to be artificial habitat type classes, such as arable or pasture land, plantations and built-up areas, noting that this could exclude areas suitable for some generalist species. For the AOH refinement we used a newly-developed global map (see Supporting Information) that follows the IUCN habitat classification system, thereby avoiding crosswalks between habitat preferences and land cover maps⁵⁹. This data product integrates the best available land cover and climate data, while also using newly developed land-use data such as data on global forest management⁶⁰. Finally, for each species and grid cell, we calculated the fractional amount (> 0-100%) of suitable habitat to include in the prioritisation analysis. Development of the habitat type map and all AOH refinement was performed on Google Earth Engine⁶¹.

Global representativeness

There is considerable bias and variability in the completeness of biodiversity records globally, particularly so for plant species⁶². To estimate the amount of geographic bias in completeness of distribution data among plants, we first estimated the proportion of species for which we had distribution data relative to the number of species known to occur in the regional checklists of World Checklist of Vascular Plants database⁴⁰, which provides for each accepted species name its native regions from the World Geographical Scheme for Recording Plant Distributions (WGSRPD,⁶⁴). We used geographic delineations for 50 WGSRPD level 2 regions⁶⁴, but excluded Antarctica and mid-Atlantic islands (Saint Helena and Ascension) for which we had no plant

records. The proportion of species for which we had range data varied from 11% in islands of the North Pacific up to 100% in the Russian far east (mean 60.1% $\pm$ 24.5 SD). However, for 48 of the 50 WCSP regions we had distribution data for over >10% of all described plants known to occur natively in that region, (the exception being islands in the South-West and South-Central Pacific). For 44 of these 50 regions we had distribution data for >40% of described plants in those regions.

Having identified 10% as the minimum common denominator of completeness across most regions, we then used an iterative heuristic algorithm, to construct ‘representative’ subsets consisting of random samples that approximated 10% of species from each WGSRPD level 2 region while accounting for the fact that some species occur across multiple regions. To test if this approach yielded sets representative of biogeographic patterns of the full dataset, we compared the spatial patterns of scaled vertebrate species richness to the 10% sets of these species for each WGSRPD level 2 regions, random subsets of 10% of all vertebrates and for all vertebrates combined. We performed the test on vertebrates because we had range maps for ~95% of terrestrial vertebrates described, therefore we can assess if our subsampling to representative sets can replicate “true” patterns in species richness obtained with a complete sample of species in a taxonomic group. Spatial patterns of scaled species richness were identical across those sets, suggesting that this sampling approach can account for incomplete coverage (SI Fig 7a).

We also checked if the frequency distribution of range sizes within our subsets matched the range size distribution of the entire set using mammals as a test group, and found very modest differences between the full set and multiple subsets (SI Fig 7b). Having confirmed that this procedure recreates correct patterns of conservation priorities and it does not alter the range-size distribution (SI Fig 7), we proceeded to create 10 subsets of ~10% of plant species known to occur in each WGSRPD level 2 region and ten non-overlapping subsets of 10% of vertebrate species for all of our analyses. We found little difference among representation curves regardless of whether multiple representative subsets or all species were included in the SCP, although there was greater efficiency in the latter (SI Fig. 8).

Carbon data

We used spatial estimates of the density of aboveground and belowground biomass carbon and vulnerable soil carbon⁹. Estimates for aboveground carbon (AGC) were created by selecting the best available carbon maps for different types of vegetation classes, identified spatially using the Copernicus Land Cover map in 2015⁶⁵. We used Santoro *et al.* as a baseline for a global carbon biomass map^{66,67}, which has been shown to be the most accurate, especially so for ‘tree’ covered land. In addition, we used more detailed estimates of above-ground biomass for African “open forest” and “shrubland” land cover⁶⁸, global “herbaceous vegetation” and “moss and lichen” land cover⁶⁹ and “cropland” and “bare/sparse vegetation” land-cover classes⁷⁰. To map below-ground carbon, we applied corrected root-to-shoot ratios⁷¹ obtained from the Intergovernmental Panel on Climate Change (IPCC) technical guidance documents⁷². A newly developed forest management layer⁶⁰ was used to update biomass density, by averaging estimates from 2010 and 2017⁶⁶ in the most dynamic tree-covered classes (e.g. short rotation plantations, agroforestry).

The map of vulnerable soil organic carbon was created following IPCC Guidelines for National Greenhouse Inventories to estimate emissions and removals associated with changes in land use⁷². Vulnerable soil organic carbon was defined as those carbon stocks that could potentially be lost during the coming 30 years as a result of land use. We used recently published data on baseline soil organic carbon stocks⁷³, and vulnerable stocks were estimated separately for mineral and organic soils. Organic soils were defined as those soils with $\geq 5\%$ probability of being a Histosols according to USDA soil orders taxonomy⁷⁴. All other soils were considered to be mineral soils. A 30cm depth was used to estimate vulnerable carbon stocks on mineral soils, while 200cm depth was used for organic soils. IPCC change factors (mineral soils) and emission factors (for organic soils) were used to estimate vulnerable soil organic carbon stocks according to IPCC land cover categories and climate zones. To be consistent with biomass carbon estimations, we created a crosswalk between the Copernicus global land cover map⁶⁵ and IPCC land cover classes. The newly developed forest management layer⁶⁰ was used to refine vulnerable carbon stock estimates for mineral soils, whilst managed forest with organic soils were excluded from this assessment given that due to drainage, these areas would be more suitable for restoration than for conservation action. Finally, all global carbon estimates were reprojected, summed and aggregated (arithmetic mean) to 10 km to match the biodiversity data in scale.

Water data

For capturing water provisioning, we used estimates of potential clean water provision calculated by WaterWorld⁷⁵ and Co\$ting Nature⁷⁶. This quantity calculates for each grid cell the volume of water available, as the accumulated water balance from upstream based on rainfall, fog and snowmelt sources minus actual evapotranspiration. Second, clean water was assessed using the Human Footprint on Water Quality (HFWQ) index, which is a measure of the extent to which water runoff is drawn from contaminating human land uses: both point (urban, roads, mining, oil and gas) and nonpoint (unprotected cropland, unprotected pasture) sources. The HFWQ index is calculated by cumulating the downstream runoff from polluting and non-polluting land uses and expressing the former runoff as a proportion of the total runoff. This is calculated by assigning an associated percentage (or dilution) intensity fraction to each land-use class (default values taken from⁷⁶). The potential clean water provisioning service is calculated for each cell as the inverse of clean water (i.e. 100 - HFWQ) available from upstream. For the analysis we ranked each grid per river basin⁷⁷ to determine their relative importance in delivering clean water within the basin.

Prioritisation analysis

We determined global areas of importance to be managed for conserving biodiversity, carbon and water by using a spatial conservation prioritisation approach (SCP⁷⁸). We divided the world in 10 km resolution ‘planning units’ (PUs, the cells of the land-surface area grids), in which ‘features’ are distributed (each species, plus carbon stocks and water provision), for which we establish conservation targets⁷⁹. Each PU had an area ‘cost’ subject to ‘budget’ constraints (the total amount of the terrestrial land-surface within a PU). For biodiversity, we defined species-specific targets

aimed at conserving the area of habitat (AOH) for a species to improve in conservation status (¹⁵, see Supporting Information) and for each species we calculated the amount of suitable habitat within each PU. For tonnes of carbon storage ($\frac{tC}{km^2}$) and/or volume of water ($\frac{Mm^3}{km^2}$), we maximized the total amount present in each PU. All PUs had a cost equivalent to the amount of land within them ($\{0 < c \leq 1\}$), which we calculated from Copernicus land-cover data⁶⁵. As global budget (B) we set different percentages of the terrestrial land surface area starting at 10%, then increasing by 10% increments up until all targets were met.

Problem formulation

Areas of importance for the conservation of biodiversity, carbon and water were determined by solving a global optimization problem. For each feature j included in the analysis we aimed to minimize the proportional shortfall⁸⁰ in achieving each representation target t_j given a planning unit cost c and an area budget B (10, 20, ..., 100% of $\sum_{i=1}^I c_i$ the planet). For all species, t is the target shortfall, that is, the difference between the part of an AOH that is included in the solution, and the amount that is necessary to be conserved for the species to improve in conservation status (¹⁵, Supporting Information), while for carbon storage and water provisioning t is the total amount available on the terrestrial land (the target is 100%). The problem is formulated as follows:

$$\text{Minimize } \sum_{j=1}^J w_j \frac{y_j}{t_j}$$

Subject to

$$\sum_{i=1}^I x_i r_{ij} + y_j \geq t_j \forall j \in J$$

$$\sum_{i=1}^I x_i c_i \leq B, \text{ where } 0 \leq x_i \leq 1 \forall i \in I$$

where r_{ij} is the amount (suitable habitat in km^2 , total tons of carbon $\frac{tC}{km^2}$ or volume of water $\frac{Mm^3}{km^2}$) of feature j in planning unit i , y_j is the shortfall for feature j , t_j is the target for feature j , c_i is the cost of grid cell i (the fractional area within the planning unit), B is the budget of the problem, x_i is a proportional decision variable [0-1], where 1 means that the full PU and values ≥ 0 a fraction of the PU is selected, and W_j is the weight assigned to feature j . We tested different W_j of carbon, respectively water, relative to biodiversity and different weights among species based on their global threat status and/or evolutionary distinctiveness (Supporting Information). The problem is then solved for each budget incrementally, by ‘locking in’ previous solutions with lower area-budget prior to running the next prioritisation, effectively building nested sets of priorities with increasing budget B .

Analysis variants

For a separate analysis, we constrained the optimization by locking in the fraction of currently protected areas and adjusted the starting budget accordingly (Supporting Information). We then

jointly optimized globally for biodiversity, carbon and water by minimizing the proportional shortfall⁸⁰ in reaching the targets for each given area budget B (10, 20, ..., 100% of the planet).

We furthermore considered a number of optimization variants in which we modified either the targets or weights assigned to each feature (biodiversity, carbon and/or water). For biodiversity, we also considered variants distinguishing between species intraspecific variation, threat status and evolutionary distinctiveness (SI Table 2). To capture intraspecific variation, we considered each part of a species range occurring in geographically separate biomes as a separate feature with its own target²⁸, e.g. the Tiger (*Panthera tigris*) was split into five separate features, one for each of the five biomes overlapping the tiger range (Supporting Information). However, we only considered a split for features in which at least 2,200 km² of AOH (the minimum absolute target area) was contained within a different biome compared to the biome with the majority of the species range. Compared to a version without these splits and when optimizing for biodiversity, carbon and water, overall differences were relatively minor (SI Fig. 11), but potentially locally important. We also collated data on species current threat status and, for vertebrates, data on their evolutionary distinctiveness (Supporting Information), and then calculated weights for each species following¹³. We then optimized all variants by minimizing the target-weighted shortfalls across all biodiversity features, subject to budget constraints.

We set weights for carbon storage and water provisioning relative to biodiversity in all analyses variants that included these assets. To do so we assigned sequences of weights from ‘none’ up to ‘equal’ importance by weighting carbon and water as follows: $w_k = 1 + \sum_{j=1}^J w_j$, w_k is the weight for carbon and water, J is the total number of species in the analysis, and $\sum_{j=1}^J w_j$ is the cumulative sum of all species weights. This weighting ensures that carbon is given equal importance to all species combined and that feature targets are treated equally in the optimization. We also created separate scenarios where w_k is set to $\frac{1}{10}, \frac{2}{10}, \dots$ of the equal weighting relative to the cumulative shortfall for biodiversity. We visualized all scenarios with increasing budget and by the shortfall in carbon, water and improvement in species conservation status (Fig. 2) Because of the high computational cost of calculating $(2N_w - 1) * N_B$ prioritizations, where N_w is the number of weights and N_B the number of budgets, for each of the 10 representative sets, we assessed differing weights at 50 km rather than 10 km resolution. However, we note that compared to a 10 km resolution, both spatial patterns and accumulation curves were highly similar (See Supporting Information and SI Fig. 9) and we don’t expect results to differ because of differences in resolution.

Optimization algorithm and ranking

All SCP variants were solved using an integer linear programming (ILP) approach. Compared to other conservation planning solutions that rely on simulated annealing or heuristics⁸¹, ILP has been shown to outcompete those approaches in both speed and solution performance, being able to reliably find optimal solutions^{82,83}. We ran all problem variants under each budgetary constraints (10%, 20%...100% of land), each with a representative set of species and solved them to optimality using proportional decisions (e.g. asking which fraction of a grid cell is part of the solution). For each problem variant, we therefore obtained 10 nested sets of priorities (priority ranks), each

resulting from solving all budgetary constraints with a representative set of species. We summarized these priority ranks through an arithmetic mean while also separately calculating the coefficient of variation as a measure of uncertainty in priorities across representative subsets (SI Fig. 1). Selected planning units in the obtained solutions were investigated for the representation of input features by taxonomic group, threatened species and biomes.

All data preparation and analysis was conducted in R⁸⁴ mainly relying on the ‘prioritizr’ package⁸⁵ with the Gurobi solver enabled (ver 8.11,⁸⁶).

Data availability All produced integrated maps will be made available through <https://unbiodiversitylab.org/> and a data repository upon acceptance. The raw input data can be requested from the respective data providers, namely IUCN, GARD, Birdlife International, Kew Gardens and predicted plant distribution data will be made available as part of the BIEN initiative⁴⁴. The IUCN habitat type map used to construct the AOH is made available in the Supporting Information. Any additional data not listed can be made available from the authors upon reasonable request or will be openly published separately.

Code availability Code to reproduce the main results will be made available upon acceptance.

Acknowledgements This work was conducted by the NatureMap consortium. We thank Richard Corlett & Tom Brooks who provided feedback on an earlier version of the manuscript. We furthermore thank Tom Hengl (OpenLandMap) for his advice on the Soil Organic Carbon analysis. This study has benefited from a number of data providers and networks. We explicitly acknowledge all data providers in a separate Extended Acknowledgements owing to their length (Supporting Information). The NatureMap project acknowledges funding from Norway's International Climate and Forest Initiative (NICFI). Collection of the plant data used in this analysis has benefited from funding in form of a GEF grant 5810-SPARC 'Spatial Planning for Area Conservation in Response to Climate Change'. CM acknowledges funding from NSF (National Science Foundation) grant DBI-1913673. RVG was supported by Australian Research Council DECRA Fellowship (DE170100208). Furthermore EAN and XF were funded by the Bridging Biodiversity and Conservation Science Program of the University of Arizona. NMH was supported by the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement no. 746334. JCS considers this work a contribution to his VILLUM Investigator project “Biodiversity Dynamics in a Changing World” funded by VILLUM FONDEN (grant 16549) and his Independent Research Fund Denmark | Natural Sciences project TREECHANGE (grant 6108-00078B).

Author contributions MJ and PV designed the study, MJ led the analysis and interpretation of the data and has drafted the manuscript; PV conceived the study, contributed to the analysis and drafting of the manuscript; JH, BB, CM contributed to creating software used in the work; AA, CR, SGR, ML, DS, AvS, MM, JM, SP, IO, BS, CM, BJE, XF, PRR, BB, BM, RVG contributed to acquisition, analysis and interpretation of data; JH, MDM, JM, WJ, SR, JM, MO, MR, XDL contributed to interpretation of the data; GO, SM, ML, RG, MyL, OT contributed to acquisition and interpretation of data; XDL, VK, LM, NB, GW, JDS, GST contributed to conception of the study;

VA,SPA,SCA,JRB,RTC,LH,PAM,JKM,DMN,NMH,EAN,DSP,PRR,JCS,CV,JJW provided data and contributed to interpretation of the data. All authors contributed to revising the manuscript. Correspondence and requests for materials should be addressed to MJ & PV.

Bibliography

1. Díaz, S. *et al.* Pervasive human-driven decline of life on Earth points to the need for transformative change. *Science* **366**, eaax3100 (2019).
2. Butchart, S. H. M., Miloslavich, P., Reyers, B. & Subramanian, S. M. Chapter 3. Assessing progress towards meeting major international objectives related to nature and nature's contributions to people. in *IPBES Global Assessment on Biodiversity and Ecosystem Services* 1–355 (IPBES, 2019).
3. Griscom, B. W. *et al.* Natural climate solutions. *Proc. Natl. Acad. Sci.* **114**, 11645–11650 (2017).
4. CBD. Zero draft of the Post-2020 Global Biodiversity Framework. (2020).
5. Anderson, C. M. *et al.* Natural climate solutions are not enough. *Science* **363**, 933–934 (2019).
6. Dinerstein, E. *et al.* A Global Deal For Nature: Guiding principles, milestones, and targets. *Sci. Adv.* **5**, eaaw2869 (2019).
7. Visconti, P. *et al.* Protected area targets post-2020. *Science* **364**, eaav6886 (2019).
8. Locke, H. *et al.* Three Global Conditions for Biodiversity Conservation and Sustainable Use: an implementation framework. *Natl. Sci. Rev.* (2019) doi:10.1093/nsr/nwz136.
9. Soto-Navarro, C. *et al.* Mapping co-benefits for carbon storage and biodiversity to inform conservation policy and action. *Philos. Trans. R. Soc. B Biol. Sci.* **375**, 20190128 (2020).
10. Strassburg, B. B. N. *et al.* Strategic approaches to restoring ecosystems can triple conservation gains and halve costs. *Nat. Ecol. Evol.* **3**, 62–70 (2019).
11. Greve, M., Reyers, B., Mette Lykke, A. & Svenning, J.-C. Spatial optimization of carbon-

- 597 stocking projects across Africa integrating stocking potential with co-benefits and feasibility.
- 598 *Nat. Commun.* **4**, 2975 (2013).
- 599 12. Brooks, T. M. *et al.* Global Biodiversity Conservation Priorities. *Science* **313**, 58–61 (2006).
- 600 13. Pouzols, F. M. *et al.* Global protected area expansion is compromised by projected land-use
- 601 and parochialism. *Nature* **516**, 383–386 (2014).
- 602 14. Allan, J. R. *et al.* Conservation attention necessary across at least 44% of Earth’s terrestrial
- 603 area to safeguard biodiversity. *bioRxiv* (2019) doi:10.1101/839977.
- 604 15. Fastre, S., Mogg, C., Jung, M. & Visconti, P. Targeted expansion of Protected Areas to
- 605 maximise the persistence of terrestrial mammals. *bioRxiv* **3124**, 1–19 (2019).
- 606 16. Rinnan, D. S. & Jetz, W. Terrestrial conservation opportunities and inequities revealed by
- 607 global multi-scale prioritization. *bioRxiv* (2020) doi:10.1101/2020.02.05.936047.
- 608 17. Newbold, T. *et al.* Has land use pushed terrestrial biodiversity beyond the planetary
- 609 boundary? A global assessment. *Science* **353**, 288–291 (2016).
- 610 18. Kier, G. *et al.* A global assessment of endemism and species richness across island and
- 611 mainland regions. *Proc. Natl. Acad. Sci.* **106**, 9322–9327 (2009).
- 612 19. McInnes, L. *et al.* Do Global Diversity Patterns of Vertebrates Reflect Those of Monocots?
- 613 *PLoS ONE* **8**, e56979 (2013).
- 614 20. Pollock, L. J., Thuiller, W. & Jetz, W. Large conservation gains possible for global
- 615 biodiversity facets. *Nature* **546**, 141–144 (2017).
- 616 21. Daru, B. H. *et al.* Spatial overlaps between the global protected areas network and terrestrial
- 617 hotspots of evolutionary diversity. *Glob. Ecol. Biogeogr.* **28**, 757–766 (2019).
- 618 22. Chaplin-Kramer, R. *et al.* Global modeling of nature’s contributions to people. *Science* **366**,
- 619 255–258 (2019).
- 620 23. Hannah, L. *et al.* 30% land conservation and climate action reduces tropical extinction risk
- 621 by more than 50%. *Ecography* *ecog.05166* (2020) doi:10.1111/ecog.05166.

- 622 24. Wilson, E. O. *Half-earth: our planet's fight for life*. (WW Norton & Company, 2016).
- 623 25. Myers, N., Mittermeier, R. A., Mittermeier, C. G., da Fonseca, G. A. B. & Kent, J.
624 Biodiversity hotspots for conservation priorities. *Nature* **403**, 853–858 (2000).
- 625 26. Venter, O. *et al.* Harnessing Carbon Payments to Protect Biodiversity. *Science* **326**, 1368–
626 1368 (2009).
- 627 27. Strassburg, B. B. N. *et al.* Global congruence of carbon storage and biodiversity in terrestrial
628 ecosystems. *Conserv. Lett.* **3**, 98–105 (2010).
- 629 28. Dinerstein, E. *et al.* An Ecoregion-Based Approach to Protecting Half the Terrestrial Realm.
630 *BioScience* **67**, 534–545 (2017).
- 631 29. Woodley, S. *et al.* A Review of Evidence for Area-based Conservation Targets for the Post-
632 2020 Global Biodiversity Framework. *Parks* **25**, 31–46 (2019).
- 633 30. Rapacciuolo, G. *et al.* Species diversity as a surrogate for conservation of phylogenetic and
634 functional diversity in terrestrial vertebrates across the Americas. *Nat. Ecol. Evol.* **3**, 53–61
635 (2019).
- 636 31. Venter, O. *et al.* Targeting Global Protected Area Expansion for Imperiled Biodiversity.
637 *PLoS Biol.* **12**, e1001891 (2014).
- 638 32. Laffoley, D. *et al.* An introduction to ‘other effective area-based conservation measures’
639 under Aichi Target 11 of the Convention on Biological Diversity: Origin, interpretation and
640 emerging ocean issues’. *Aquat. Conserv. Mar. Freshw. Ecosyst.* **27**, 130–137 (2017).
- 641 33. Chauvenet, A. L. M., Kuempel, C. D., McGowan, J., Beger, M. & Possingham, H. P.
642 Methods for calculating Protection Equality for conservation planning. *PLoS ONE* (2017)
643 doi:10.1371/journal.pone.0171591.
- 644 34. Waldron, A. *et al.* Reductions in global biodiversity loss predicted from conservation
645 spending. *Nature* **551**, 364–367 (2017).
- 646 35. Adams, V. M., Pressey, R. L. & Naidoo, R. Opportunity costs: Who really pays for

- 647 conservation? *Biol. Conserv.* **143**, 439–448 (2010).
- 648 36. Cameron, E. K. *et al.* Global gaps in soil biodiversity data. *Nat. Ecol. Evol.* **in press**, 6–7
- 649 (2018).
- 650 37. Jetz, W. *et al.* Essential biodiversity variables for mapping and monitoring species
- 651 populations. *Nat. Ecol. Evol.* **3**, 539–551 (2019).
- 652 38. Violle, C. *et al.* Functional Rarity: The Ecology of Outliers. *Trends Ecol. Evol.* **32**, 356–367
- 653 (2017).
- 654 39. Di Marco, M., Ferrier, S., Harwood, T. D., Hoskins, A. J. & Watson, J. E. M. Wilderness
- 655 areas halve the extinction risk of terrestrial biodiversity. *Nature* **573**, 582–585 (2019).
- 656 40. WCVF. World Checklist of Vascular Plants. *World Checklist of Vascular Plants*
- 657 <http://wcvf.science.kew.org/> (2020).
- 658 41. IUCN. IUCN 2019. The IUCN Red List of Threatened Species. Version 2019.2. *IUCN*
- 659 *Redlist* www.iucnredlist.org (2019).
- 660 42. Birdlife International. *Digital boundaries of Key Biodiversity Areas from the World*
- 661 *Database of Key Biodiversity Areas*. (2019).
- 662 43. Roll, U. *et al.* The global distribution of tetrapods reveals a need for targeted reptile
- 663 conservation. *Nat. Ecol. Evol.* **1**, 1677–1682 (2017).
- 664 44. Enquist, B., Condit, R., Peet, R., Schildhauer, M. & Thiers, B. Cyberinfrastructure for an
- 665 integrated botanical information network to investigate the ecological impacts of global
- 666 climate change on plant biodiversity. *PeerJ Prepr.* (2016) doi:10.7287/peerj.preprints.2615.
- 667 45. Enquist, B. J. *et al.* The commonness of rarity: Global and future distribution of rarity across
- 668 land plants. *Sci. Adv.* **5**, eaaz0414 (2019).
- 669 46. Maitner, B. S. *et al.* The bien r package: A tool to access the Botanical Information and
- 670 Ecology Network (BIEN) database. *Methods Ecol. Evol.* **9**, 373–379 (2018).
- 671 47. Anderson-Teixeira, K. J. *et al.* CTFS-ForestGEO: a worldwide network monitoring forests in

- 672 an era of global change. *Glob. Change Biol.* **21**, 528–549 (2015).
- 673 48. Forest Inventory and Analysis National Program. www.fia.fs.fed.us/ (2013).
- 674 49. Peet, R., Lee, M., Jennings, M. & Faber-Langendoen, D. VegBank – a permanent, open-
675 access archive for vegetation-plot data. *Biodivers. Ecol.* **4**, 233–241 (2012).
- 676 50. Boyle, B. & Enquist, B. SALVIAS – the SALVIAS vegetation inventory database.
677 *Biodivers. Ecol.* (2012) doi:10.7809/b-e.00086.
- 678 51. Wiser, S. K., Bellingham, P. J. & Burrows, L. E. Managing biodiversity information:
679 Development of New Zealand’s National Vegetation Survey databank. *N. Z. J. Ecol.* (2001).
- 680 52. DeWalt, S. J., Bourdy, G., ChÁvez de Michel, L. R. & Quenevo, C. Ethnobotany of the
681 Tacana: Quantitative inventories of two permanent plots of Northwestern Bolivia. *Econ. Bot.*
682 **53**, 237–260 (1999).
- 683 53. Dauby, G. *et al.* RAINBIO: A mega-database of tropical African vascular plants
684 distributions. *PhytoKeys* (2016) doi:10.3897/phytokeys.74.9723.
- 685 54. Fegraus, E. Tropical Ecology Assessment and Monitoring Network (TEAM Network).
686 *Biodivers. Ecol.* **4**, 287–287 (2012).
- 687 55. Oliveira-Filho, A. T. Um sistema de classificação fisionômico-ecológico da vegetação
688 neotropical: segunda aproximação. in *Fitossociologia no Brasil - Métodos e estudos de caso*
689 - *Volume II* (2015). doi:10.1016/j.jns.2007.02.032.
- 690 56. Butchart, S. H. M. *et al.* Shortfalls and Solutions for Meeting National and Global
691 Conservation Area Targets. *Conserv. Lett.* **8**, 329–337 (2015).
- 692 57. Rondinini, C., Stuart, S. & Boitani, L. Habitat suitability models and the shortfall in
693 conservation planning for African vertebrates. *Conserv. Biol.* **19**, 1488–1497 (2005).
- 694 58. Brooks, T. M. *et al.* Measuring Terrestrial Area of Habitat (AOH) and Its Utility for the
695 IUCN Red List. *Trends Ecol. Evol.* **34**, 977–986 (2019).
- 696 59. IUCN. Habitats Classification Scheme, Version 3.1. 1–14 (2012).

- 697 60. Lesiv et al., M. Mapping Human Impact On Forests At Global Scale. (2020).
- 698 61. Gorelick, N. *et al.* Google Earth Engine: Planetary-scale geospatial analysis for everyone.
699 *Remote Sens. Environ.* **202**, 18–27 (2017).
- 700 62. Meyer, C., Weigelt, P. & Kreft, H. Multidimensional biases, gaps and uncertainties in global
701 plant occurrence information. *Ecol. Lett.* **19**, 992–1006 (2016).
- 702 63. Meyer, C., Kreft, H., Guralnick, R. & Jetz, W. Global priorities for an effective information
703 basis of biodiversity distributions. *Nat. Commun.* **6**, 8221 (2015).
- 704 64. Brummitt, R. K. *World geographical scheme for recording plant distributions*. (International
705 Working Group on Taxonomic Databases for Plant Sciences (TDWG), 2001).
- 706 65. Buchhorn, M. *et al.* Copernicus Global Land Service: Land Cover 100m: Collection 2: epoch
707 2015. in (2019). doi:10.5281/zenodo.3243509.
- 708 66. Santoro, M. *GlobBiomass - global datasets of forest biomass*. (PANGAEA, 2018).
709 doi:10.1594/PANGAEA.894711.
- 710 67. Santoro, M. & Cartus, O. ESA Biomass Climate Change Initiative (Biomass_cci): Global
711 datasets of forest above-ground biomass for the year 2017. (2019)
712 doi:10.5285/bedc59f37c9545c981a839eb552e4084.
- 713 68. Bouvet, A. *et al.* An above-ground biomass map of African savannahs and woodlands at 25
714 m resolution derived from ALOS PALSAR. *Remote Sens. Environ.* **206**, 156–173 (2018).
- 715 69. Xia, J. *et al.* Spatio-Temporal Patterns and Climate Variables Controlling of Biomass Carbon
716 Stock of Global Grassland Ecosystems from 1982 to 2006. *Remote Sens.* **6**, 1783–1802
717 (2014).
- 718 70. Spawn, S. A., Lark, T. J. & Gibbs, H. K. A New Global Biomass Map for the Year 2010. in
719 *A New Global Biomass Map for the Year 2010* (2017).
- 720 71. Schepaschenko, D. *et al.* Improved estimates of biomass expansion factors for Russian
721 forests. *Forests* (2018) doi:10.3390/f9060312.

- 722 72. Eggleston, S., Buendia, L., Miwa, K., Ngara, T. & Tanabe, K. *2006 IPCC guidelines for*
723 *national greenhouse gas inventories*. vol. 5 (2006).
- 724 73. Hengl, T. & Wheeler, I. Soil Organic Carbon Stock in kg/m² for 5 Standard Depth Intervals
725 (0–10, 10–30, 30–60, 60–100 and 100–200 Cm) at 250 M Resolution. (2018)
726 doi:10.5281/ZENODO.2536040.
- 727 74. Hengl, T. & Nauman, T. Predicted USDA soil orders at 250 m (probabilities) (Version v0.1).
728 <http://doi.org/10.5281/zenodo.2658183> (2019).
- 729 75. Mulligan, M. WaterWorld: a self-parameterising, physically based model for application in
730 data-poor but problem-rich environments globally. *Hydrol. Res.* **44**, 748–769 (2013).
- 731 76. Mulligan, M. Trading off agriculture with nature’s other benefits, spatially. in *The Impacts of*
732 *Climate Change on Water Resources in Agriculture* 184–204 (CRC Press, 2014).
- 733 77. Linke, S. *et al.* Global hydro-environmental sub-basin and river reach characteristics at high
734 spatial resolution. *Sci. Data* **6**, 283 (2019).
- 735 78. Kukkala, A. S. & Moilanen, A. Core concepts of spatial prioritisation in systematic
736 conservation planning. *Biol. Rev.* **88**, 443–464 (2013).
- 737 79. Cabeza, M. & Moilanen, A. Design of reserve networks and the persistence of biodiversity.
738 *Trends Ecol. Evol.* **16**, 242–248 (2001).
- 739 80. Arponen, A., Heikkinen, R., Thomas, C. D. & Moilanen, A. The Value of Biodiversity in
740 Reserve Selection: Representation, Species Weighting, and Benefit Functions. *Conserv. Biol.*
741 **19**, 2009–2014 (2005).
- 742 81. Lehtomäki, J. & Moilanen, A. Methods and workflow for spatial conservation prioritization
743 using Zonation. *Environ. Model. Softw.* **47**, 128–137 (2013).
- 744 82. Beyer, H. L., Dujardin, Y., Watts, M. E. & Possingham, H. P. Solving conservation planning
745 problems with integer linear programming. *Ecol. Model.* **328**, 14–22 (2016).
- 746 83. Hanson, J. O., Schuster, R., Strimas-Mackey, M. & Bennett, J. R. Optimality in prioritizing

747 conservation projects. *Methods Ecol. Evol.* 2041–210X.13264 (2019) doi:10.1111/2041-
748 210X.13264.

749 84. R Core Team. *R: A Language and Environment for Statistical Computing*. (R Foundation for
750 Statistical Computing, 2019).

751 85. Hanson, J. *et al. prioritizr: Systematic Conservation Prioritization in R*. (2019).

752 86. Gurobi Optimization, L. *Gurobi Optimizer Reference Manual*. (2019).

753

Supporting online material

Material and Methods

Choice of resolution

We chose a spatial resolution of 10km to adequately capture global biodiversity and nature's contribution to people per grid cell. For the biodiversity data we used estimates of a species global range. Previous studies have recommended coarser spatial resolution (~110km) when using species range maps as such, to better match equally downscaled atlas data considered to be the 'true' distribution of a species¹, however, this can result in more costly prioritisations due to commission errors, without meaningful reductions in spatial biases². In this study we refined a species range to an Area of Habitat (AOH,³) to minimize commission errors (false presences). This was done at a spatial resolution similar or even coarser than in comparable studies relying on the same range data⁴⁻⁷. Lastly, we also created separate maps of all analyses at 50km resolution to investigate differences on identified areas of biodiversity importance (SI Fig. 9), and found overall little to no difference between analyses done at these different resolutions. Nevertheless, we caution that the identified global areas of importance should not be used to inform conservation decisions on local or landscape scales.

Plant data preparation

To this date, there does not exist a single and consistent data source for species range data of all described plant species globally⁸⁻¹⁰. The total number of plant species globally is still unknown, with existing estimates ranging between 352,282 species¹¹ and over 434,934 species⁹. To obtain a representative subset of described plant species, the NatureMap consortium gathered the best available plant distribution data from a variety of sources and types, acknowledging that none of them are without errors and biases, which we addressed by calculating spatially representative sets, each approximating the same proportion of species known to exist in an area, across the planet.

We first relied on expert-based global range estimates created by the International Union for Conservation of Nature (IUCN), Royal Botanic Gardens, Kew, and Botanic Gardens Conservation International (BGCI). For many plant species only curated point estimates of their range were available. Based on this data, range estimates were constructed using alpha-hulls, a generalization of convex hulls that are particularly useful for estimating species ranges whose habitat is irregularly shaped¹² or where populations are spatially structured¹³. Parameters for alpha-hulls creation were adaptively selected, starting with initial alpha values - a parameter constraining the hull triangulation - of 2 or 3 recommended by the IUCN Red List categories and criteria, but adjusted for the distribution of records so that at least 95% of the records were included within the estimated range. The value of alpha ranges from zero (i.e. the finest resolution defined by the given set of points) to infinity (i.e. the coarsest resolution defined by the convex-hull). Since variations in alpha can also affect subpopulation structure (i.e. number of subpopulations), we combined alpha-hulls with the "1/10th max" circular buffer method (i.e. the buffer size is set to the tenth of the maximum interpoint distance) to better capture subpopulation structure¹³. Finally, we limited the number of subpopulations to maximum of 10 and if the conditions above are not met (i.e. >=

95% of records inside the estimated range and ≤ 10 subpopulations), a minimum convex hull or a buffer built with the “1/10th max” method is drawn around each record¹³. We split the occurrence records geographically into separate parts in cases the alpha hulls could not be constructed (for instance close to 180° longitude). In these cases, we applied the alpha-hull method to each individual dataset and merged the calculated hulls back into one unique range. All alpha hulls and “1/10th max” buffers were created using the *rangeBuilder* package¹⁴. In total, data for 8,702 plant species ranges could be obtained through both sources, including 4,598 tree species from BGCI and 4,104 plant species from IUCN.

For plant species not yet assessed by IUCN or BGCI, we relied on modelled range estimates derived from occurrence records acquired through the Botanical Information and Ecology Network (BIEN) initiative, the Global Biodiversity Information Facility (GBIF.org 2019, <https://doi.org/10.15468/dl.gvt20i>) and from iNaturalist (www.inaturalist.org). Not all research grade observations from iNaturalist are transferred to GBIF and we thus downloaded all research grade iNaturalist plant data separately and merged them with the GBIF data, while removing duplicate observations.

The observations in the BIEN database are the product of contributions by 1,076 different data contributors, including numerous individual herbaria, and data indexers of herbaria (550+ are listed in Index Herbariorum), that were used to construct conservative estimates of species ranges using species distribution models (SDMs). For details of specimen data sources see^{9,16}. We benefited from version 4.1 of BIEN, which includes data from RAINBIO¹⁷, TEAM¹⁸, The Royal Botanical Garden of Sydney, Australia, and NeoTropTree¹⁹. Additional plant plot data from a number of networks and datasets have been included in BIEN^{8,9,16,20–25} and a full listing of the herbaria data used can be found in the extended acknowledgements below and online (<http://bien.nceas.ucsb.edu/bien/data-contributors/all/>).

Taxon names associated with BIEN occurrence records were first corrected for misspellings, homonyms (e.g. plant and animal species with identical names) and synonyms. Afterwards all taxon names were standardized using TNRS v4.0 at default settings with checklists from Tropicos, The Plant List, USDA Plants, Global Compositae Checklist, ILDIS²⁶. Standard BIEN preprocessing procedures furthermore ensure that species outside their native ranges were removed using lists of endemic taxa and the Native Species Resolver (NSR; <https://github.com/ojalaquellueva/nsr>). Observations were furthermore flagged and removed as cultivated based on keywords in the original observation metadata.

We applied the following preprocessing steps to all plant occurrence records from BIEN, GBIF and iNaturalist. We removed all occurrence records that (1) had no or impossible coordinates (e.g. $< 90^\circ$ S latitude or longitude $> 180^\circ$ or $< -180^\circ$), (2) had a coordinate uncertainty greater than 10 km, (3) had identical latitude or longitude coordinates, duplicate records or where coordinates had a precision smaller than one digit, (4) removed occurrence records in the vicinity (10 km distance) of country capitals or outside the lowest declared political division in the case of BIEN using the Geographic Name Resolution Service (GNRS; <http://bien.nceas.ucsb.edu/bien/tools/gnrs/>), near country or province centroids (1 km), or in the vicinity (1 km) of known zoos, botanical gardens or herbaria and (5) removed all occurrence points that fell into the open ocean²⁷. For the modelling, we merged plant occurrence records from GBIF and iNaturalist into one dataset per species and only included those records from BIEN that were not already present in other data sources.

Plant species can have varying uncertainties in taxonomies and geographic spread and quite commonly occur in regions where the species is not considered native. In this study we relied on taxonomic and geographic information from the Plants of the World online (POWO) database, which provides for each accepted species name its native World Geographical Scheme for Recording Plant Distributions (WGSRPD) regions^{28,29}. We only included plant species in the

analysis whose name could be matched to POWO taxonomy (either as accepted name or as synonym) and which had at least one occupied grid cell in all WGSRPD level 2 regions in which the species is known to be native, to reduce influences of sampling biases. Lastly, we post-hoc removed from each predicted distribution all unconnected isolated patches outside native WGSRPD regions, which we identified through connected component labeling³⁰.

For modelling plant species distributions we used a number of environmental covariates, which are adequate for the spatial scale (global at 10 km) of our modelling approach³¹. Data on present (1979-2013) climatic conditions (Annual Mean Temperature, Mean Diurnal Range, Annual precipitation, Precipitation seasonality, Precipitation of Warmest Quarter, Precipitation of Coldest Quarter, maximum accumulated Aridity (consecutive water deficit during months where potential evapotranspiration exceed precipitation) & estimated relative Precipitation of Warmest

$$\text{Quarter} = \frac{\text{Precipitation of Warmest Quarter}}{(\text{Precipitation of Warmest Quarter} + \text{Precipitation of Coldest Quarter})}$$

Quarter = were obtained from CHELSA (<http://chelsa-climate.org/>,³²). Data on global aridity³³ and soil conditions (bulk density, % clay content, depth to bedrock, pH & % silt content all averaged over full depth to 200cm) from <https://soilgrids.org>³⁴. These covariates were chosen based on their ecological relevance for plant species and on having global correlations < 0.7 with each other³⁵. All environmental covariates were aggregated (arithmetic mean) to 10 km globally and projected to an equal-area Mollweide projection.

Point process modelling

For all plant species with 10 or more records available we fitted Poisson point process models (closely related to Maxent) using regularized down weighted Poisson regression models³⁶, fitted with the R package glmnet³⁷. We used up to a maximum of 20,000 background points in total, adjusted based on the total number of grid cells within the domain, and chose a spatial domain for predictions based on the biomes a species occurred in³⁸. All candidate predictors were further filtered for collinearity for each individual species separately³⁵, with highly collinear covariates (Pearson' $r > 0.7$) within the domain removed.

Five independent folds were trained for cross validation, where folds were assigned based on spatial clusters to remove the influence of spatial autocorrelation on cross-validated performance statistics. Linear (all species), quadratic (species with >100 records), and product (species with >200 records) features were used. Regularization parameters for each model were determined based on one standard deviation below the minimum variance³⁷. This resulted in five models per species which were then combined in an unweighted ensemble by calculating the arithmetic mean and standard deviation of the folds. Finally, the continuous predictions were thresholded to obtain binary presence/absence predictions based on the 5th percentile of the ensemble predictions.

Range-bagging models

For all plant species with between five and lower than ten records we utilized a 'range bagging' approach, which is a stochastic, hull-based method that can estimate climate niches from an ensemble of underfit models^{39,40}, and is therefore well suited for smaller datasets. We randomly sampled 100 times a proportion p of records ($p = 0.33$, based on recommendations in³⁹) and a subset d of environmental variables ($d = 2$,³⁹). A convex hull is then projected around the subsampled records in environmental space, with a record considered part of the species range if its environmental conditions fall within the hull. We then chose a voting threshold of 0.165 ($=0.33/2$), implying that the grid cell is part of the species range at least half the time for each subsample. Upon visual inspection we generally found that this threshold leads to relatively

conservative predictions. All range bagging records and environmental predictors were subjected to the same selection rules as for the point process models discussed above.

Grid cell data

For plant species with less than three covered grid cells records we used only those grid cells the points fall, which often describe the full distribution of the species known to science, many of which are globally rare⁹.

Ancillary data

To account for current areas managed for conservation, we included data on current global protected areas from the global World Database on Protected Areas (WDPA, April 2019 version, IUCN and UNEP-WCMC 2019). Following commonly used WDPA preparation standards⁴¹, we excluded protected areas whose status was ‘proposed’ or ‘not reported’ and furthermore removed UNESCO Man & Biosphere reserves. This figure, however, does not include data from countries that have restricted the sharing of their dataset through the WDPA, such as China, Estonia, Saint Helena, Ascension and Tristan da Cunha⁴¹. All layers were first rasterized at 1 km, then aggregated to 10 km by calculating the relative fraction of area protected, so that small PAs were not lost. As a result, ~15% of the land surface was identified as being protected in the prioritisation analysis. Lastly, we prepared data on terrestrial biomes and ecoregions (<http://ecoregions2017.appspot.com>,³⁸), which were likewise rasterized to 10 km resolution using a modal aggregation.

Habitat types map

Not all parts of a species range are equally suitable to allow a species to persist, thus requiring a refinement to an area of suitable habitat (AOH,^{3,5}). In the past this refinement has commonly been attempted using a crosswalk⁴² between land-cover legends and habitat type information from the IUCN habitat type classification⁴³. Crosswalks between different thematic legends can potentially cause issues such as inseparability of habitat types that are identical in land cover but different in climatic and soil conditions (e.g. tropical moist lowland forest and tropical mangrove forest). We developed a new global habitat type layer that follows the IUCN habitat classification system⁴³. This layer is an intersection of the best currently available land cover dataset⁴⁴, data on climate⁴⁵ and other ancillary datasets, such as a novel data product on the distribution of global anthropogenically modified forests including tropical and temperate plantations (Lesiv et al. unpublished). Using this layer we refined all species ranges (see methods) at 1 km globally and calculated the fraction of suitable habitat per 10 km grid cell. We make a version of this global layer available as part of this manuscript⁴⁶.

Prioritisation analysis

Target setting

One of the most impactful decisions in spatial conservation planning frameworks is the definition of feature targets. In the past, many studies set targets for species representation according to rules^{47–49} or area-based policies (e.g. 30% of a species range), which run the risk of leading to an excess of area for wide-ranging species and arbitrariness. We set targets relative to the amount of habitat necessary to improve a species conservation status as inspired by IUCN criteria⁵⁰. We recognise that this only takes the range (area of suitable habitat) into account, and ignores other

factors of extinction risk, such as population size and trends, but the purpose is to provide ecologically credible area-based conservation targets, rather than estimating extinction risk. For all species, these targets were defined as

$$t_i = \frac{\min(\max(2200 \text{ km}^2, 0.8 * A_{AOH_i}), 1e^6 \text{ km}^2)}{A_{AOH_i}},$$

where t_i is the relative target for a given species i and A_{AOH_i} the total area of suitable habitat for the species⁵⁰. Whenever the numerator exceeded the A_{AOH_i} (e.g. is smaller than 2200 km²), the target was set to the whole AOH (100%), following³⁷. In the prioritisation analysis we ranked each PU after formulating and solving a budget limited formulation of the reserve selection problem that aims to maximize conservation benefits.

Species-specific weights

Areas of biodiversity importance can vary depending on whether greater weight is placed on evolutionarily distinct⁵¹ and/or threatened species⁵². For this analysis we obtained data on the evolutionary distinctiveness (ED) scores for amphibians (99.7% of all species considered), birds (100%), mammals (100%) and reptiles (71.9%) from the EDGE program (EDGE 2019 list,⁵³). For plant species there does not yet exist a species-resolved phylogeny⁵⁴ and further research is necessary to fill that gap. Whenever ED scores could not be matched to species names, we used the congeneric or family-wide ED average⁵⁵. ED scores represent the amount of unique evolutionary history of a species^{56,57}, thus placing greater weight on evolutionary older and most distinctive lineages in a phylogeny. For example, Cuba and Hispaniola have evolutionary significance because these were the only two species of *Solenodon* that exist; the only members of the mammal family *Solenodontidae* which diverged from all other mammals over 60 million years ago, thus representing a disproportionate amount of evolutionary history. Data on the threat category (TC) of species was obtained from IUCN and encoded as numerical weight. In addition, for plant species we used data from the ThreatSearch online database⁵⁸. We followed Pouzols et al. (2014) and assigned a weight of 8 to Critically Endangered species (CR), 6 to Endangered (EN), 4 to Vulnerable (VU), 2 to Near Threatened (NT) and 1 to species of Least Concern (1). Plant species without a standardized IUCN threat category, but which are considered threatened according to BGCI, were assigned a weight of 6. Species without sufficient current TC information or that were Data Deficient (DD) were assigned a conservative score of 2, given that many Data Deficient species are likely threatened with extinction^{59,60}, especially so for plant species¹¹. We separately incorporated for each species either the evolutionary distinctiveness (ED) score or the threat category (TC) as weight in the prioritisation, using weight from TC weights⁵². In total, we included data on ED weights for 34,308 species, TC weights for 43,211 species and calculated separated problem variants where data for both (29,780 species) is available (SI Fig. 10).

Supplementary figures and tables

SI Fig. 1: Uncertainty in ranks of areas of importance for biodiversity, carbon and water. Calculated as coefficient of variation across optimal solutions with different representative sets. Expressed as percentage with lower values indicating higher precision of ranks. Map can be interpreted as overall confidence in the mapped ranks (Fig. 1), given existing biases in species range data. Map is at 10 km resolution in Mollweide projection.

SI Fig. 2: Global areas of importance for biodiversity only. Ranked hierarchical maps by the most (1-10%) and least important areas (90-100%) to conserve all of biodiversity globally. Map is at 10 km resolution in Mollweide projection.

991

992

993

994

995

996

997

998

SI Fig. 3: Global areas of importance for carbon and water. Normalized ranking for carbon (a) and water (b) presented as the most (1-10%) and least important areas (90-100%) to conserve globally. Map is at 10 km resolution in Mollweide projection.

SI Fig. 4: Global areas of importance for biodiversity and carbon or biodiversity and water. Showing an optimization across 10 representative sets for either (a) biodiversity and carbon or (b) biodiversity and water. All assets were jointly optimized and ranked hierarchical by the most (1-10%) and least important areas (90-100%) to conserve globally. Map is at 10 km resolution in Mollweide projection.

SI Fig. 5: Global areas of importance for biodiversity, carbon and water considering current protected areas. All assets were jointly optimized and ranked hierarchical by the most (1-10%) and least important areas (90-100%) to conserve globally. The fraction of grid cells currently managed for conservation (<https://www.protectedplanet.net>) are considered to be part of the most important areas. Map is at 10 km resolution in Mollweide projection.

SI Fig. 6: Accumulation curves showing how the number of species targets met increases with amount of land optimally allocated to conservation considering current protected areas. Shows the amount of land necessary for all assets to reach all persistence targets, defined as the amount of area needed for a species to be considered at reduced risk of extinction (see Methods). Uncertainty bands (~0.1% around the mean) show the standard deviation among representative sets. Estimates shown for species (a) overall and split by additional number of assets, (b) by taxonomic group, and (c) by current IUCN assessment of threat.

a

b

SI Fig. 7: Comparison of representative sets spatially and in range size distributions. Compared to a full dataset, both subsampling at random and per WGSRPD region produces similar patterns in space and species area-size distributions. (a) Spatial map in Mollweide projection showing aggregated richness layers of all vertebrate species for the full dataset, a random sample and a representative sample by WGSRPD level 2 regions, (b) Shows the log10-transformed Area of Habitat (AOH) of all species in the full dataset (dark blue) compared to representative subsets of species (other colours).

SI Fig. 8: Accumulation curves showing how the number of species targets met increases with amount of land optimally allocated to conservation. Estimates shown for representative subsets (dotted line) and for all species included (solid line).

SI Fig. 9: Comparison of global areas of importance at 10 km and 50 km areas. Comparisons in variants of areas of importance for biodiversity only; biodiversity and carbon; and biodiversity, carbon and water. Inset graphs show how the number of species targets met increases with amount of land optimally allocated to conservation for both 10 km (blue) and 50 km (orange).

SI Fig. 10: Difference in the top-ranked 10% solution for varying species weights. For each biodiversity feature a weight was assigned equating to either no differential weight (red), current

threat category (green) or evolutionary distinctiveness (ED) (blue). Comparison was made only for species with data on both threat category and evolutionary distinctiveness. Grid cells coloured in black were selected in all three solutions. Map in Mollweide projection at 10 km resolution. The line plot shows the amount of land necessary for all species to reach all persistence targets, defined as the amount of area needed for a species to improve in conservation status (see Methods). Shown for either no weight (red), species weighted by threat status (green) and weighted by evolutionary distinctiveness (blue). The inset zoom highlights the difference among solutions at a budget of 10% terrestrial land area. The confidence bounds of accumulation curves indicate the uncertainty among representative sets.

SI Fig. 11: Global areas of importance for terrestrial biodiversity, carbon and water without biome splits. All assets were jointly optimized with equal weighting and ranked hierarchical by the most (1-10%) and least (90-100%) important areas to conserve globally. The map is at 10 km resolution in Mollweide projection.

SI Table 1: List of data sources included in the analysis. Shown is the source, taxonomic group and number of species ranges from that source. For the analysis we preferentially used species range data from IUCN and Birdlife International. Subsequently we relied on GARD, Kew and BGCI data and used BIEN estimates of species ranges for all other plant species not already included. Details on data preparation can be found in the methods and supporting information.

Data source	Taxonomic group	Total number of species
IUCN Mammal ranges	Mammals	5,685
IUCN Amphibian ranges	Amphibians	6,660
Birdlife International	Birds	10,953
IUCN Reptiles	Reptiles	6,830
GARD Reptiles	Reptiles	3,755

IUCN Plants	Plants	8,172
IUCN Plants (new alpha hulls)	Plants	4,090
BGCI Plants (new alpha hulls)	Plants	4,571
BIEN Plant SDMs	Plants	105,336
BIEN Plant Rangebags	Plants	31,634
BIEN Plant Grid cells	Plants	40,151

SI Table 2: Problem variants created as part of the analyses.

<uploaded separately>

Extended acknowledgements

This study has benefited from data made available through a number of providers and networks. We would like to thank the IUCN Red List GIS Unit and Birdlife International for making vertebrate and plant species ranges available for scientific research. We thank the IUCN redlist and all species assessors globally for making habitat preference data available (<https://iucnredlist.org>). We thank Rikki Gumps and Claudia Gray for pointing us to the latest EDGE data (<https://www.edgeofexistence.org/edge-lists/>). We thank BGCI for making available up-to-date threat assessments of plant species via ‘threat search’ (https://tools.bgci.org/threat_search.php) and Royal Botanic Gardens, Kew for creating and making available the Global Plants of The World database (www.plantsoftheworldonline.org/).

We specifically thank Manaaki Whenua – Landcare Research and the New Zealand National Herbarium Network for making data available from the New Zealand National Vegetation Survey databank (NVS, <https://nvs.landcareresearch.co.nz/>) and the New Zealand Virtual Herbarium Network (NZVH). We thank iNaturalist for support in raising awareness and systematically collecting further data for countries with few plant occurrence records in a dedicated project (<https://www.inaturalist.org/projects/naturemap-plants>).

We thank all the 1076 data contributors to BIEN. This includes RAINBIO, TEAM, The Royal Botanical Garden of Sydney, Australia, and NeoTropTree. Plot data within BIEN are from the CVS, NVS, SALVIAS, VEGBANK, CTFS, FIA, MADIDI, and TEAM data networks and datasets (<http://bien.nceas.ucsb.edu/bien/data-contributors/all>). We acknowledge the herbaria that contributed data to BIEN: HA, FCO, DUKE, MFU, UNEX, VDB, ASDM, AMD, BPI, BRI, BRM, CLF, CNPO, L, LPB, AD, A, TAES, FEN, FHO, ANSM, ASU, B, BCMEX, RAS, RB, TRH, AAH, ACOR, UI, AK, CAS, ALCB, AKPM, EA, AAU, ALTA, ALU, AMES, AMNH, AMO, CHAPA, GH, ANGU, ANSP, ARAN, AS, CICY, BAI, CIMI, AUT, BA, BAA, BAB, CMMEX, BACP, BAF, BAJ, BAL, COCA, CODAGEM, BARC, BAS, BBS, BC, BCN, BCRU, BERE, BG, BH, BIO, BISH, SEV, BLA, BM, BOCH, MJG, BOL, CVRD, BOLV, BONN, DAV, BOUM, BR, DES, BREM, BRLU, BSB, BUT, C, DS, CALI, CAN, CANB, CAY, EBUM, CBM, CEN, CEPEC, CESJ, CHR, ENCB, CIIDIR, CINC, CLEMS, F, COA, COAH, FCME, COFC, CP, COL, COLO, CONC, CORD, CPAP, CPUN, CR, CRAI, FURB, CU, G,

CRP, CS, CSU, CTES, CTESN, CUZ, DAO, HB, DBN, DLF, DNA, DR, DUSS, E, HUA, EAC, EIF, EIU, GES, GI, GLM, GMNHJ, K, GOET, GUA, EMMA, HUAZ, ERA, ESA, FAA, FAU, FB, UVIC, FI, GZU, H, FLAS, FLOR, HCIB, FR, FTG, FUEL, GB, HNT, GDA, HPL, GENT, HUAA, HUJ, CGE, HAL, HAM, IAC, HAMAB, HAO, HAS, IB, HASU, HBG, IBUG, HBR, HEID, IEB, HIP, IBGE, ICEL, ICN, ILL, SF, HO, HRCB, HRP, HSS, HU, HUAL, HUEFS, HUEM, HUFU, HUSA, HUT, IAA, HXBH, HYO, IAN, ILLS, HAC, IPRN, IMSSM, FCQ, ABH, INEGI, INIF, BAFC, BBB, INPA, IPA, NAS, INB, INM, MW, EAN, IZTA, ISKW, ISC, ISL, GAT, JEPS, IBSC, UCSB, ISTC, ISU, IZAC, JACA, JBAG, JE, SD, JUA, JYV, KIEL, ECON, KSC, TOYA, MPN, USF, TALL, RELC, CATA, AQP, KMN, KMNH, KOELN, KOR, FRU, KPM, KSTC, LAGU, TRTE, KSU, UESC, GRA, IBK, KTU, ACAD, MISSA, KU, PSU, KYO, LA, LOMA, LW, SUU, UNITEC, TASH, NAC, UBC, IEA, GMDRC, LD, M, LE, LEB, LIL, LINN, AV, HUCP, QFA, LISE, MBML, NM, MT, FAUC, MACF, CATIE, LTB, LISI, LISU, MEXU, LL, LOJA, LP, LPAG, MGC, LPD, LPS, IRVC, MICH, JOTR, LSU, LBG, WOLL, LTR, MNHN, CDBI, LYJB, MOL, DBG, AWH, NH, HSC, LMS, MELU, NZFRI, MA, UU, MU, CSUSB, MAF, MAK, MB, KUN, MARY, MASS, MBK, MBM, UCSC, UCS, JBGP, DSM, OBI, BESA, LSUM, FULD, MCNS, ICESI, MEL, MEN, TUB, MERL, CGMS, MFA, FSU, MG, HIB, MIL, DPU, TRT, BABY, ETH, YAMA, SCFS, SACT, ER, JCT, JROH, SBBG, SAV, PDD, MIN, SJSU, MMMN, PAMP, MNHM, OS, SDSU, BOTU, OXF, P, MOR, POM, MPU, MPUC, MSB, MSC, CANU, SFV, RSA, CNS, WIN, MSUN, CIB, MUR, MTMG, VIT, MUB, MVFA, SLPM, MVFQ, PGM, MVJB, MVM, MY, PASA, N, UCMH, HGM, TAM, BOON, UFS, MARS, CMM, NA, NU, UADY, UAMIZ, UC, NE, NHM, NHMC, NHT, UFMA, NLH, UFRJ, UFRN, ULS, UMO, UNL, UNM, US, NMB, NMNL, USP, NMR, NMSU, WIS, NSPM, XAL, NSW, NT, ZMT, BRIT, MO, NCU, NY, TEX, U, UNCC, NUM, O, CHSC, LINC, CHAS, ODU, CDA, OSA, OSC, OSH, OULU, OWU, PACA, PAR, UPS, PE, PEL, SGO, PEUFR, PFC, PH, PKDC, SI, PLAT, PMA, PORT, PR, QM, PRC, TRA, PRE, PY, QCA, TROM, QCNE, QRS, UH, QUE, R, SAM, RBR, REG, RFA, RIOG, RM, RNG, RYU, S, SALA, SANT, SAPS, SASK, SBT, SEL, SIU, SJRP, SMDB, SMF, SNM, SOM, SP, SRFA, SPF, SPSF, SQF, STL, STU, SVG, TAI, TAIF, TAMU, TAN, TEF, TENN, TEPB, TFC, TI, TKPM, TNS, TO, TU, UAM, UB, UCR, UEC, UFG, UFMT, UFP, UGDA, UJAT, ULM, UME, UNA, UNB, UNR, UNSL, UPCB, UPEI, UPNA, USAS, USJ, USM, USNC, USZ, UT, UTC, UTEP, UWO, V, VAL, VALD, VEN, VMSL, VT, W, WAG, WAT, WII, WELT, WFU, WMNH, WS, WTU, WU, Z, ZSS, ZT, CUVC, LZ, AAS, AFS, BHCB, CHAM, FM, PERTH, SAN.

1150 Bibliography

- 1151 1. Hurlbert, A. H. & Jetz, W. Species richness, hotspots, and the scale dependence of range
1152 maps in ecology and conservation. *Proc. Natl. Acad. Sci.* **104**, 13384–13389 (2007).
- 1153 2. Di Marco, M., Watson, J. E. M., Possingham, H. P. & Venter, O. Limitations and trade-offs
1154 in the use of species distribution maps for protected area planning. *J. Appl. Ecol.* **54**, 402–
1155 411 (2017).
- 1156 3. Brooks, T. M. *et al.* Measuring Terrestrial Area of Habitat (AOH) and Its Utility for the
1157 IUCN Red List. *Trends Ecol. Evol.* **34**, 977–986 (2019).
- 1158

4. Buchanan, G. M., Donald, P. F. & Butchart, S. H. M. Identifying Priority Areas for Conservation: A Global Assessment for Forest-Dependent Birds. *PLoS ONE* **6**, e29080 (2011).
5. Powers, R. P. & Jetz, W. Global habitat loss and extinction risk of terrestrial vertebrates under future land-use-change scenarios. *Nat. Clim. Change* (2019) doi:10.1038/s41558-019-0406-z.
6. Soto-Navarro, C. *et al.* Mapping co-benefits for carbon storage and biodiversity to inform conservation policy and action. *Philos. Trans. R. Soc. B Biol. Sci.* **375**, 20190128 (2020).
7. Hanson, J. O. *et al.* Global conservation of species' niches. *Nature* (2020) doi:10.1038/s41586-020-2138-7.
8. Enquist, B., Condit, R., Peet, R., Schildhauer, M. & Thiers, B. Cyberinfrastructure for an integrated botanical information network to investigate the ecological impacts of global climate change on plant biodiversity. *PeerJ Prepr.* (2016) doi:10.7287/peerj.preprints.2615.
9. Enquist, B. J. *et al.* The commonness of rarity: Global and future distribution of rarity across land plants. *Sci. Adv.* **5**, eaaz0414 (2019).
10. Weigelt, P., König, C. & Kreft, H. GIFT – A Global Inventory of Floras and Traits for macroecology and biogeography. *J. Biogeogr.* jbi.13623 (2019) doi:10.1111/jbi.13623.
11. Joppa, L. N., Roberts, D. L. & Pimm, S. L. How many species of flowering plants are there? *Proc. R. Soc. B Biol. Sci.* **278**, 554–559 (2011).
12. Burgman, M. A. & Fox, J. C. Bias in species range estimates from minimum convex polygons: implications for conservation and options for improved planning. *Anim. Conserv.* **6**, 19–28 (2003).
13. Rivers, M. C., Bachman, S. P., Meagher, T. R., Nic Lughadha, E. & Brummitt, N. A. Subpopulations, locations and fragmentation: applying IUCN red list criteria to herbarium specimen data. *Biodivers. Conserv.* **19**, 2071–2085 (2010).

- 1184 14. Davis Rabosky, A. R. *et al.* Coral snakes predict the evolution of mimicry across New World
1185 snakes. *Nat. Commun.* (2016) doi:10.1038/ncomms11484.
- 1186 15. GBIF.org. GBIF Occurrence Download. (2019) doi:<https://doi.org/10.15468/dl.gvt20i>.
- 1187 16. Maitner, B. S. *et al.* The bien r package: A tool to access the Botanical Information and
1188 Ecology Network (BIEN) database. *Methods Ecol. Evol.* **9**, 373–379 (2018).
- 1189 17. Dauby, G. *et al.* RAINBIO: A mega-database of tropical African vascular plants
1190 distributions. *PhytoKeys* (2016) doi:10.3897/phytokeys.74.9723.
- 1191 18. Fegraus, E. Tropical Ecology Assessment and Monitoring Network (TEAM Network).
1192 *Biodivers. Ecol.* **4**, 287–287 (2012).
- 1193 19. Oliveira-Filho, A. T. Um sistema de classificação fisionômico-ecológico da vegetação
1194 neotropical: segunda aproximação. in *Fitossociologia no Brasil - Métodos e estudos de caso*
1195 - *Volume II* (2015). doi:10.1016/j.jns.2007.02.032.
- 1196 20. Anderson-Teixeira, K. J. *et al.* CTFS-ForestGEO: a worldwide network monitoring forests in
1197 an era of global change. *Glob. Change Biol.* **21**, 528–549 (2015).
- 1198 21. Forest Inventory and Analysis National Program. www.fia.fs.fed.us/ (2013).
- 1199 22. Peet, R., Lee, M., Jennings, M. & Faber-Langendoen, D. VegBank – a permanent, open-
1200 access archive for vegetation-plot data. *Biodivers. Ecol.* **4**, 233–241 (2012).
- 1201 23. Boyle, B. & Enquist, B. SALVIAS – the SALVIAS vegetation inventory database.
1202 *Biodivers. Ecol.* (2012) doi:10.7809/b-e.00086.
- 1203 24. Wiser, S. K., Bellingham, P. J. & Burrows, L. E. Managing biodiversity information:
1204 Development of New Zealand’s National Vegetation Survey databank. *N. Z. J. Ecol.* (2001).
- 1205 25. DeWalt, S. J., Bourdy, G., Chávez de Michel, L. R. & Quenevo, C. Ethnobotany of the
1206 Tacana: Quantitative inventories of two permanent plots of Northwestern Bolivia. *Econ. Bot.*
1207 **53**, 237–260 (1999).
- 1208 26. Boyle, B. *et al.* The taxonomic name resolution service: an online tool for automated

1209 standardization of plant names. *BMC Bioinformatics* **14**, 16 (2013).

1210 27. Zizka, A. *et al.* CoordinateCleaner: Standardized cleaning of occurrence records from
1211 biological collection databases. *Methods Ecol. Evol.* **10**, 744–751 (2019).

1212 28. Brummitt, R. K. *World geographical scheme for recording plant distributions*. (International
1213 Working Group on Taxonomic Databases for Plant Sciences (TDWG), 2001).

1214 29. WCVP. World Checklist of Vascular Plants. *World Checklist of Vascular Plants*
1215 <http://wcvp.science.kew.org/> (2020).

1216 30. Hesselbarth, M. H. K., Sciaini, M., With, K. A., Wiegand, K. & Nowosad, J.
1217 landscapemetrics : an open-source R tool to calculate landscape metrics. *Ecography*
1218 *ecog.04617* (2019) doi:10.1111/ecog.04617.

1219 31. Thuiller, W., Araújo, M. B. & Lavorel, S. Do we need land-cover data to model species
1220 distributions in Europe? *J. Biogeogr.* **31**, 353–361 (2004).

1221 32. Karger, D. N. *et al.* Climatologies at high resolution for the earth’s land surface areas. *Sci.*
1222 *Data* **4**, 170122 (2017).

1223 33. Antonio Trabucco & Robert Zomer. Global Aridity Index and Potential Evapotranspiration
1224 (ET0) Climate Database v2. (2019) doi:10.6084/m9.figshare.7504448.v3.

1225 34. Hengl, T. *et al.* SoilGrids250m: Global gridded soil information based on machine learning.
1226 *PLoS ONE* (2017) doi:10.1371/journal.pone.0169748.

1227 35. Dormann, C. F. *et al.* Collinearity: a review of methods to deal with it and a simulation study
1228 evaluating their performance. *Ecography* **36**, 27–46 (2013).

1229 36. Renner, I. W. *et al.* Point process models for presence-only analysis. *Methods Ecol. Evol.* **6**,
1230 366–379 (2015).

1231 37. Friedman, J., Hastie, T. & Tibshirani, R. Regularization Paths for Generalized Linear Models
1232 via Coordinate Descent. *J. Stat. Softw.* **33**, 1–22 (2010).

1233 38. Dinerstein, E. *et al.* An Ecoregion-Based Approach to Protecting Half the Terrestrial Realm.

BioScience **67**, 534–545 (2017).

39. Drake, J. M. Range bagging: a new method for ecological niche modelling from presence-only data. *J. R. Soc. Interface* **12**, 20150086 (2015).

40. Drake, J. M. & Richards, R. L. Estimating environmental suitability. *Ecosphere* **9**, e02373 (2018).

41. Bingham, H. C. *et al.* Sixty years of tracking conservation progress using the World Database on Protected Areas. *Nat. Ecol. Evol.* (2019) doi:10.1038/s41559-019-0869-3.

42. Santini, L. *et al.* Applying habitat and population-density models to land-cover time series to inform IUCN Red List assessments. *Conserv. Biol.* **00**, cob1.13279 (2019).

43. IUCN. Habitats Classification Scheme, Version 3.1. 1–14 (2012).

44. Buchhorn, M. *et al.* Copernicus Global Land Service: Land Cover 100m: Collection 2: epoch 2015. in (2019). doi:10.5281/zenodo.3243509.

45. Beck, H. E. *et al.* Present and future Köppen-Geiger climate classification maps at 1-km resolution. *Sci. Data* **5**, 180214 (2018).

46. Jung, M. *et al.* A global map of terrestrial habitat types (Version 001). *Zenodo* (2020) doi:10.5281/zenodo.3666246.

47. Rodrigues, A. S. L. *et al.* Effectiveness of the global protected area network in representing species diversity. *Nature* **428**, 640–643 (2004).

48. Carwardine, J. *et al.* Cost-effective priorities for global mammal conservation. *Proc. Natl. Acad. Sci.* **105**, 11446–11450 (2008).

49. Venter, O. *et al.* Targeting Global Protected Area Expansion for Imperiled Biodiversity. *PLoS Biol.* **12**, e1001891 (2014).

50. Fastre, S., Mogg, C., Jung, M. & Visconti, P. Targeted expansion of Protected Areas to maximise the persistence of terrestrial mammals. *bioRxiv* **3124**, 1–19 (2019).

51. Pollock, L. J., Thuiller, W. & Jetz, W. Large conservation gains possible for global

biodiversity facets. *Nature* **546**, 141–144 (2017).

52. Pouzols, F. M. *et al.* Global protected area expansion is compromised by projected land-use and parochialism. *Nature* **516**, 383–386 (2014).

53. Gumbs, R., Gray, C. L., Wearn, O. R. & Owen, N. R. Tetrapods on the EDGE: Overcoming data limitations to identify phylogenetic conservation priorities. *PLoS ONE* **13**, 1–19 (2018).

54. Eiserhardt, W. L. *et al.* A roadmap for global synthesis of the plant tree of life. *Am. J. Bot.* **105**, 614–622 (2018).

55. Curnick, D. J. *et al.* Setting evolutionary-based conservation priorities for a phylogenetically data-poor taxonomic group (Scleractinia). *Anim. Conserv.* **18**, 303–312 (2015).

56. Isaac, N. J. B., Turvey, S. T., Collen, B., Waterman, C. & Baillie, J. E. M. Mammals on the EDGE: Conservation Priorities Based on Threat and Phylogeny. *PLoS ONE* **2**, e296 (2007).

57. Tucker, C. M. *et al.* A guide to phylogenetic metrics for conservation, community ecology and macroecology. *Biol. Rev.* n/a–n/a (2016) doi:10.1111/brev.12252.

58. BGCI. ThreatSearch online database. *BGCI Threat search* www.bgci.org/threat_search.php (2019).

59. Bland, L. M., Collen, B., Orme, C. D. L. & Bielby, J. Predicting the conservation status of data-deficient species. *Conserv. Biol.* **29**, 250–259 (2015).

60. Veron, S. *et al.* Integrating data-deficient species in analyses of evolutionary history loss. *Ecol. Evol.* **6**, 8502–8514 (2016).