Immune modulation to improve survival of respiratory virus infections in mice

Authors: Shradha Wali^{1,2}, Jose R. Flores², Ana Maria Jaramillo², David L. Goldblatt², Jezreel Pantaleón García², Michael J. Tuvim², Burton F. Dickey², Scott E. Evans^{1,2}

Author affiliations: ¹University of Texas MD Anderson Cancer Center UTHealth Graduate School of Biomedical Sciences, Houston Texas 77030

²University of Texas MD Anderson Cancer Center, Department of Pulmonary Medicine, Houston Texas 77030

To whom correspondence should be addressed: Scott E. Evans, M.D.

6565 MD Anderson Blvd

Houston TX 77030

Email: seevans@mdanderson.org

Telephone: 713-563-7433

Author contributions: S.W. designed and performed the experiments, analyzed the data, and wrote the manuscript. J.R.F., A.M.J., D.L.G. and J.P.G performed experiments. M.J.T., B.F.D. conceptualized the project and critically reviewed the data. S.E.E. conceptualized the project, designed experiments, provided critical evaluation of data and edited the manuscript. ¹

¹ This study was supported by NIH grants R01 HL117976, DP2 HL123229 and R35 HL144805 to S.E.E.

Running Title: Epithelial defence against viral immunopathology

Abstract

Viral pneumonia remains a global health threat requiring novel treatment strategies, as strikingly exemplified in the SARS-CoV-2 pandemic of 2019-2020. We have reported that mice treated with a combination of inhaled Toll-like receptor (TLR) 2/6 and TLR 9 agonists (Pam2-ODN) to stimulate innate immunity are broadly protected against respiratory pathogens, but the mechanisms underlying this protection remain incompletely elucidated. Here, we show in a lethal paramyxovirus model that Pam2-ODN-enhanced survival is associated with robust virus inactivation by reactive oxygen species (ROS), which occurs prior to internalization by lung epithelial cells. However, we also found that mortality in sham-treated mice temporally corresponded with CD8⁺ T cell-enriched lung inflammation that peaks on days 11-12 after viral challenge, when the viral burden has waned to a scarcely detectable level. Pam2-ODN treatment blocked this injurious inflammation by reducing the viral burden, and alternatively, depleting CD8⁺ T cells 8 days after viral challenge also decreased mortality. These findings reveal opportunities for targeted immunomodulation to protect susceptible individuals against the morbidity and mortality of respiratory viral infections.

Keywords: Immunomodulation, immunopathology, CD8⁺ T cells, viral pneumonia, inducible epithelial resistance

Introduction

Viruses are the most frequent cause of community acquired pneumonia in children and adults, resulting in significant morbidity in vulnerable subjects and exerting a tremendous health care burden (1-5). Seasonal influenza and emergent pandemic viruses, such as SARS-CoV-2, inflict particular mortality in susceptible individuals, with clinicians frequently lacking effective interventions to improve patient outcomes (6-9). Moreover, in addition to causing acute disease, respiratory virus infections are often complicated by chronic lung pathologies, such as asthma induction, progression and exacerbation (10-12). Therefore, development of novel therapeutic anti-viral strategies is required to effectively prevent and treat respiratory infections and their associated chronic complications (13, 14).

While lung epithelial cells are the principal targets of most respiratory viruses (15), there is expanding evidence that lung epithelia themselves are capable of generating anti-microbial responses (12, 16, 17). We hypothesized that lung epithelial cells can be harnessed to control virus replication, thereby enhancing acute survival and reducing chronic complications of virus infections (18-21). Our group has previously described the phenomenon of inducible epithelial resistance wherein the lungs' mucosal defenses can be broadly stimulated to protect against a wide range of respiratory pathogens, including viruses (18-23). This protection is induced by a single inhalation of a combination treatment consisting of Toll like receptor (TLR) 2/6 and 9 agonists (Pam2-ODN) shortly before or after viral challenge. While no individual leukocyte populations have been identified as critical for Pam2-ODN-induced resistance, lung epithelial cells are essential to the inducible anti-viral response (18). Further, we have shown that

Pam2-ODN mediated protection is dependent upon epithelial generation of reactive oxygen species (ROS) but, interestingly, does not require Type I interferons (22, 23). More recently, we have shown prevention of chronic virus-induced asthma in mice treated with Pam2-ODN but we have not clarified the anti-viral mechanisms (24).

In this study, we investigated the mechanisms of Pam2-ODN enhanced mouse survival of pneumonia caused by a paramyxovirus, Sendai virus (SeV). We found that Pam2-ODN treatment not only reduced lung SeV burden but also decreased epithelial cell injury and host immunopathologic leukocyte responses to SeV infections. While CD8⁺ T cells are known to contribute to anti-viral immunity, it is shown here that CD8⁺ T cells contribute substantially to mortality, and this effect can be prevented by Pam2-ODN treatment early in the course of infection or CD8⁺ depletion late in the course. Further, we demonstrate anti-viral mechanisms of inducible epithelial resistance, where virus particles are inactivated in a ROS-dependent manner prior to internalization by their epithelial targets.

Materials and Methods

Mice: All *in vivo* experiments were performed using 6- to 10-week-old C57BL/6J mice of a single sex and were handled according to the Institutional Animal Care and Use Committee of MD Anderson Cancer Center, protocol 00000907-RN01.

Cells: Mouse lung epithelial (MLE-15) cells were kindly provided by Jeffrey Whitsett, Cincinnati Children's Hospital Medical Center. Mouse tracheal epithelial cells were harvested and cultured as previously described (22, 25). See *Supplemental Methods* for additional details.

TLR treatments and viral challenges: Cells were treated with Pam2CSK₄ (2.2 μM) and ODN M362 (0.55 μM) as previously described (22, 23). Mice were treated with 10 ml of Pam2CSK₄ (4 μM) and ODN M362 (1 μM) by nebulization as previously described (22, 23). For *in vitro* challenges, SeV at multiplicity of infection (MOI) = 1 was used. Unless otherwise stated, mice were challenged with 1 x 10⁸ plaque forming units (pfu) inserted into the oropharynx as described (24). See *Supplemental Methods* for additional details.

Flow cytometry: Single cells from disaggregated lungs or cell culture were stained as indicated in the antibody table (Table 1), fixed, and acquired on a BD LSRII (BD Biosciences). See *Supplemental Methods* for additional details.

Epithelial proliferation assays: Epithelial proliferation was determined by staining lung sections for EdU 24 h after intraperitoneal injection. See *Supplemental Methods* for details.

CD8⁺ T cell depletion: Anti-CD8- β antibody (200 μ g/mouse, clone 53-5.8, Bioxell) was delivered to mice intraperitoneally at indicated time points. CD8⁺ T cell depletion was confirmed by flow cytometry analysis 24 to 48 h after depletion.

Viral burden quantification: Viral burden was determined by reverse transcription quantitative PCR (RT-qPCR) of the Sendai Matrix (M) protein normalized to host house-keeping gene 18SRNA. For *in vivo* experiments, mouse lungs were collected 5 days after SeV challenge. For *in vitro* experiments, cell lysates were collected 24 h after infection, unless otherwise indicated. See *Supplemental Methods* for additional details.

ROS inhibition *in vitro* and *in vivo*: NADPH oxidase activity was inhibited using GKT137831 (Selleckchem). Mitochondrial ROS production was inhibited using the combination of FCCP (Cayman Chemicals) and TTFA (Cayman Chemicals). See *Supplemental Methods* for additional details.

Viral attachment assays: For most enveloped viruses, internalization into epithelial cells is inhibited at 4° C without affecting viral binding to epithelial cells (26-28). MLE-15 cells were infected with SeV at 4° C for 4 h, washed to remove unattached virus, then assessed for uninternalized SeV burden using immunofluorescence or flow cytometry. See *Supplemental Methods* for additional details.

Results

Enhanced mouse survival of SeV infection by Pam2-ODN treatment

Aerosolized Pam2-ODN treatment one day prior to SeV challenge increased mouse survival of SeV challenge (Figure 1A), similar to the protection observed against lethal influenza pneumonia (18, 21, 22). The survival benefit was associated with reduced lung SeV burden, as measured by SeV M gene expression (Figure 1B). Investigating the natural progression of infection revealed that SeV lung burden was maximal on day 5 and gradually decreased until falling below the limit of quantification (LOQ) by day 11 (Figure 1C). Pam2-ODN pretreatment reduced SeV burden on all assessed days (Figure 1C). Although the lethality of SeV infection was exquisitely dependent on the inoculum size, we strikingly found that peak mortality paradoxically occurred around days 10 to 12 after infection irrespective of inoculum size, despite the fact that SeV is essentially undetectable that long after challenge (Figure 1A, C, and D).

Pam2-ODN treatment attenuates SeV-induced epithelial injury

This temporal dissociation between peak virus burden and peak mortality led to the hypothesis that SeV-induced mortality may not be exclusively driven by excessive virus burden but may also result from untoward SeV-induced host immune response. Therefore, the acute changes in mouse lungs following SeV infection were characterized. We found increases in lung epithelial cleaved caspase 3 (cCasp3), a marker for programmed cell death, on days 7 to 11 after SeV infection (Figure 2A, upper panel). Virus infection-related epithelial cell injury and death is typically associated with

proliferative repair mechanisms (29, 30). Staining the infected mouse lung tissue for Ki67 and EdU revealed maximum signals for both markers in the second week after infection (Figure 2B-E, upper panel). These events of lung epithelial cell death and proliferation coincided with the peak of mortality (day 12, Figure 1E). Further, hematoxylin and eosin staining of lung tissues infected with SeV showed profound increases in inflammatory cells from days 7 to 10 with evidence of damaged airway and parenchymal tissue (Figure 2F). However, Pam2-ODN pretreatment of mice reduced epithelial cell injury and proliferation (Figure 2A-E, lower panel). This temporal association of epithelial injury and death after viral clearance supported our hypothesis that mouse mortality caused by SeV infection is due in part to the host immune response to SeV infections.

Pam2-ODN attenuates SeV-induced lymphocytic lung inflammation.

To explore this hypothesis, the host leukocyte response to SeV infection was characterized. Differential Giemsa staining of bronchoalveolar lavage (BAL) cells revealed increased neutrophils on days 2 to 5 and increased macrophages on days 5 to 8 (Figure 3A, left and middle panel, solid grey line) after SeV challenge. Congruent with our prior studies, inhaled treatment with Pam2-ODN in the absence of infection led to a rapid rise in neutrophils that was resolved within 5 days (Figure 3A, dashed line) (31). The neutrophil response to SeV challenge was modestly increased among mice pretreated with Pam2-ODN (Figure 3A, left panel, solid dark line). Pam2-ODN-treated, SeV-challenged mice showed almost no difference in macrophage number compared to PBS-treated, SeV-challenged mice (Figure 3A, middle panel, solid dark line). A rise in

lymphocytes was observed on days 8 to 11 in PBS-treated, SeV-challenged mice (Figure 3A, right panel, solid grey line), temporally corresponding with peak mortality. However, Pam2-ODN treated, SeV-challenged mice displayed significantly reduced lymphocyte numbers at every time point assessed (Figure 3A, right panel, solid dark line). The gating strategy for lymphocyte subsets by flow cytometry is shown in Supplementary Figure 1. A modest reduction in CD4⁺ T cells was observed in Pam2-ODN-treated, SeV-challenged mice compared to PBS-treated, SeV-challenged mice (Supplementary Figure 2). We also found the percentage of CD19⁺ B220⁺ B cells reduced after SeV infection in comparison to Pam2-ODN treated and uninfected mice (Supplementary Figure 2), as has been seen with other viral models (32, 33). However, the biggest difference between groups was in CD8⁺ T cells, with Pam2-ODN-treated, SeV-challenged mice displaying a significantly lower number and percentage of CD8⁺ T cells than PBS treated, SeV-challenged mice (Figure 3B, C). Since the greatest difference after Pam2-ODN treatment was in CD8⁺ T cell levels and there was a tight correlation between peak mortality and the increase in lung CD8⁺ T cells on days 8 to 11, we investigated the role of CD8⁺ T cells in SeV-induced mortality.

Depleting CD8⁺ T cells after viral clearance enhances survival of SeV infection

To understand the apparent contributions of host immunopathology to mouse outcomes, we depleted CD8⁺ T cells on day 8 -- after virus burden was substantially reduced but before peak mouse mortality (Figures 1 and 4A). Mice depleted of CD8⁺ T cells displayed significantly enhanced survival of SeV challenge compared to mice with intact CD8⁺ T cells (Figure 4B). Depletion of CD8⁺ T cells was confirmed by flow

cytometry in disaggregated lung cells 10 days after SeV challenge (Figure 4C, Supplementary Figure 3A). We also assessed lung injury by hematoxylin and eosin staining of lung tissue 10 days after SeV challenge and found increased inflammation and epithelial cell damage in undepleted mice compared to CD8⁺ T cell-depleted mice (Figure 4D). This supported our hypothesis that CD8⁺ T cells contribute to fatal SeV-induced immunopathology.

To assess the role of CD8⁺ T cells throughout the course of infection, mouse CD8⁺ T cells were depleted prior to and during SeV challenge (Figure 4A, Supplementary Figure. 3A, B). This depletion resulted in significantly reduced survival of SeV infection (Supplementary Figure 3C), compatible with the known antiviral functions of CD8⁺ T cells (34-36). However, it is notable that Pam2-ODN treatment still significantly enhanced survival of SeV challenge even in the absence of CD8⁺ T cells (Supplementary Figure 3C). This finding was congruent with our previous studies showing Pam2-ODN inducible resistance against bacterial pneumonia despite the lack of mature lymphocytes (*Rag1*^{-/-}) (18).

Pam2-ODN treatment leads to extracellular inactivation of virus particles

As the antiviral protection consistently correlated with reduced viral burden *in vivo*, and as the reduced virus burden likely contributes to the reduced CD8⁺ T cell levels, we sought to determine how Pam2-ODN-induced responses cause antiviral effects. Assessing the effect of Pam2-ODN on SeV burden in immortalized mouse epithelial

cells (MLE-15) and primary mouse tracheal epithelial cells (mTEC), we found that Pam2-ODN treatment reduced SeV burden at every time point measured, reflecting the inducible antiviral capacity of isolated epithelial cells (Supplementary Figure 4). Further, we investigated whether the principal Pam2-ODN effect occurred before (extracellular) or after (intracellular) virus internalization into their epithelial targets. SeV inoculation was carried out at 4° C preventing SeV internalization while allowing SeV attachment to epithelial cells (26-28). Using multiple methods to determine the effect of Pam2-ODN on SeV attachment, we found no differences in attachment (Figure 5A-D). However, even though similar numbers of virus particles were attached to epithelial cells, when these attached virus particles were liberated from the epithelial cell targets, virus particles from Pam2-ODN-treated epithelial cells were less able to subsequently infect other naive epithelial cells (Figure 5E, F). As the number of attached virus particles was the same, this difference in SeV burden in cells that received liberated virus particles from PBS vs Pam2-ODN treated cells indicated that SeV is inactivated prior to epithelial internalization.

Pam2-ODN-induced epithelial ROS protect against SeV infection and CD8⁺ T cell immunopathology

The anti-influenza response initiated by Pam2-ODN requires epithelial generation of ROS from both NADPH-dependent dual oxidase and mitochondrial sources (22, 23). Extending these findings to the SeV model, an NADPH oxidase inhibitor (GKT 137831) fully abrogated the Pam2-ODN-induced anti-SeV response (Figure 6A). Similarly, treatment with a combination of FCCP (an uncoupler of oxidative phosphorylation) and

TTFA (a complex II inhibitor) obviated the Pam2-ODN-induced anti-SeV response (Figure 6B) (22, 23). Further, it was found that Pam2-ODN induced epithelial generation of ROS were required for inactivation of SeV prior to epithelial entry (Figure 6C). Congruent with these *in vitro* and *ex vivo* studies, mice treated with FCCP and TTFA before Pam2-ODN treatment and SeV challenge (Figure 6D) demonstrated reduced survival (Figure 6E), increased SeV burden (Figure 6F), and increased CD8⁺ T cells on day 10 (Figure 6G).

Discussion

In this study, we demonstrate that therapeutic stimulation of lung epithelial cells enhances mouse survival of acute SeV infections by both reducing the virus burden and attenuating host immunopathology. While our group has demonstrated inducible resistance against multiple respiratory pathogens including viruses (18-23, 31), these studies demonstrate for the first time when in the virus lifecycle the anti-viral effects begin (viz., prior to internalization), and substantiate the role of ROS in protection against SeV.

While Pam2-ODN treatment provided a significant host survival benefit in SeV infection, we observed this survival benefit occurring after the time when PBS-treated mice had cleared the virus. This observation prompted the hypothesis that host mortality is not the exclusive result of direct viral injury to the lungs, but due at least in part to the host response to the virus infections. We observed enhanced survival of SeV infections in

mice depleted of CD8⁺ T cells 8 days after infection (Figure 4A,B), revealing the importance of balancing the dual functions of CD8⁺ T cells in anti-viral immunity and in causing fatal immunopathology. Our findings suggest that the surge in CD8⁺ T cells within the lungs after most virus has been cleared causes physiologic impairment via lung injury and cell death (Figure 4D). These observations demonstrate an advantage of early immune stimulation to enhance viral clearance and late immune suppression to prevent immunopathology and enhance overall outcome of respiratory infections. This is potentially informative in the context of treating pneumonia in immunocompromised patients, and is likely applicable to broader clinical populations, including those suffering lung injury associated with SARS-CoV-2. Based on this reasoning, clinical trials of the use of Pam2-ODN to prevent or treat early COVID-19 have been launched (NCT04313023, NCT04312997), and we suggest that therapeutic targeting of CD8⁺ T cells later in COVID-19 be considered.

Previous reports support the concept of counter-balanced immune protection and immunopathology by CD8⁺ T cells during virus infections (36-41). Some reports have shown that antigen-experienced memory CD8⁺ T cells enhance respiratory syncytial virus (RSV) clearance, but also mediate severe immunopathology (39, 42). However, our study is the first to demonstrate the survival advantage in paramyxovirus respiratory infection of either stimulating the lungs' mucosal defenses early in the infection or of suppressing the CD8⁺ T cells later in the infection. Our findings are also congruent with reports on the role of CD8⁺ T cells in non-respiratory viral infection models, such as in West Nile virus infection, where CD8⁺ T cell deficient mice display decreased mortality

(40). While findings from that study and others reveal that the harmful effects of CD8⁺ T cell mediated immunopathology may supersede the benefits of T cell mediated viral clearance, the question arises of what might be the adaptive value of the vigorous late CD8⁺ T cell response. One possibility is that it might ensure that the infection does not flare again, but that seems implausible since the host has successfully defended itself against the initial infection, and innate immune mechanisms presumably remain intact and are possibly primed (43, 44), in addition to the multiple adaptive immune mechanisms that increasingly come into play. The possibility that the immunopathology simply results from an error on the part of the immune system also seems implausible in view of the substantial rate of host mortality, suggesting there is likely an adaptive value to the response. A third possibility, that the persistence of pockets of low level infection might lead to chronic lung pathology, is supported by a recent study showing that sites of viral RNA remnants following influenza infection are linked to chronic lung disease (45). Thus, a trade-off may exist between the adaptive value of a vigorous CD8⁺ T cell response to prevent chronic lung disease and the acute mortality it can cause. Manipulating this balance therapeutically will need to account for both the benefits and costs of the response. It is particularly appealing to develop inducible anti-microbial strategies that do not rely on conventional T cell-mediated microbial clearance and are also effective in vulnerable immune deficient populations (18, 22, 25, 46).

Although the CD8⁺ T cell depletion studies enhanced our understanding of immunopathology in virus infections, much of the survival benefit against SeV infection was mediated by rapid anti-viral effects induced by Pam2-ODN. This led us to

investigate the mechanisms of these inducible anti-viral effects. Given the multiple steps in the virus life cycle, it was not known at what stage Pam2-ODN exerted its anti-viral effect. Exploring this, we found no difference in number of SeV particles attached to the cells between PBS and Pam2-ODN treatment (Figure 5A-D). However, attached virus particles that were liberated from Pam2-ODN treated cells retained less infective capacity when added to naïve epithelial cells, revealing pre-internalization virus inactivation by Pam2-ODN treatment (Figure 5E, F).

Knowing that Pam2-ODN inducible resistance required ROS production to protect against influenza (22), we studied the role of ROS in Pam2-ODN-mediated reduction in SeV burden. ROS inhibition not only led to attenuation of Pam2-ODN's anti-viral effect but allowed increased lung CD8⁺ T cell numbers, implicating Pam2-ODN-induced ROS in preventing both identified mechanisms of mouse mortality in SeV pneumonia (Figure 6). ROS inhibition led to loss of Pam2-ODN-inducible *in vitro* inactivation of SeV prior to epithelial internalization (Figure 6C), demonstrating for the first time that epithelial ROS directly contribute to virus inactivation.

Production of ROS as a microbicidal mechanism has been widely reported in phagocytic cells (47-49). However, this mechanism has not been extensively studied in non-phagocytic cells (50), where it apparently acts predominantly extracellularly rather than intracellularly as in phagocytes. (Figure 5F, G). These findings of viral inactivation

by epithelial ROS production reveal an essential component of inducible epithelial resistance.

Taken together, these findings provide mechanistic insights into the antiviral responses generated by the lung epithelium and the prevention of host immunopathology that may inform future therapeutics to target immunomodulation as a means to improve the survival of respiratory infections in vulnerable populations.

Acknowledgments and Disclosures:

The authors would like to thank Dr. Yongxing Wang for optimization of ROS inhibition experiments. M.J.T., B.F.D., and S.E.E. are authors on U.S. patent 8,883,174, “Stimulation of Innate Resistance of the Lungs to Infection with Synthetic Ligands.” M.J.T., B.F.D., and S.E.E. own stock in Pulmotect, Inc., which holds the commercial options on these patent disclosures. All other authors declare that no conflict of interest exists.

References:

1. Mizgerd JP. Lung infection--a public health priority. *PLoS medicine* 2006;3(2):e76.
2. Luckhaupt SE, Sweeney MH, Funk R, Calvert GM, Nowell M, D'Mello T, Reingold A, Meek J, Yousey-Hindes K, Arnold KE, et al. Influenza-associated

hospitalizations by industry, 2009-10 influenza season, united states. *Emerging infectious diseases* 2012;18(4):556-562.

3. Thompson WW, Shay DK, Weintraub E, Brammer L, Cox N, Anderson LJ, Fukuda K. Mortality associated with influenza and respiratory syncytial virus in the united states. *Jama* 2003;289(2):179-186.

4. Jain S, Self WH, Wunderink RG, Fakhran S, Balk R, Bramley AM, Reed C, Grijalva CG, Anderson EJ, Courtney DM, et al. Community-acquired pneumonia requiring hospitalization among u.S. Adults. *N Engl J Med* 2015;373(5):415-427.

5. Jain S, Williams DJ, Arnold SR, Ampofo K, Bramley AM, Reed C, Stockmann C, Anderson EJ, Grijalva CG, Self WH, et al. Community-acquired pneumonia requiring hospitalization among u.S. Children. *N Engl J Med* 2015;372(9):835-845.

6. Fauci AS, Lane HC, Redfield RR. Covid-19 - navigating the uncharted. *N Engl J Med* 2020;382(13):1268-1269.

7. Taubenberger JK, Kash JC, Morens DM. The 1918 influenza pandemic: 100 years of questions answered and unanswered. *Sci Transl Med* 2019;11(502).

8. Wang X, Li Y, O'Brien KL, Madhi SA, Widdowson MA, Byass P, Omer SB, Abbas Q, Ali A, Amu A, et al. Global burden of respiratory infections associated with seasonal influenza in children under 5 years in 2018: A systematic review and modelling study. *Lancet Glob Health* 2020;8(4):e497-e510.

9. Li Q, Guan X, Wu P, Wang X, Zhou L, Tong Y, Ren R, Leung KSM, Lau EHY, Wong JY, et al. Early transmission dynamics in wuhan, china, of novel coronavirus-infected pneumonia. *N Engl J Med* 2020;382(13):1199-1207.

10. Busse WW, Lemanske RF, Jr., Gern JE. Role of viral respiratory infections in asthma and asthma exacerbations. *Lancet* 2010;376(9743):826-834.
11. Folkerts G, Busse WW, Nijkamp FP, Sorkness R, Gern JE. Virus-induced airway hyperresponsiveness and asthma. *American journal of respiratory and critical care medicine* 1998;157(6 Pt 1):1708-1720.
12. Holtzman MJ, Byers DE, Brett JA, Patel AC, Agapov E, Jin X, Wu K. Linking acute infection to chronic lung disease. The role of il-33-expressing epithelial progenitor cells. *Annals of the American Thoracic Society* 2014;11 Suppl 5:S287-291.
13. Olin JT, Wechsler ME. Asthma: Pathogenesis and novel drugs for treatment. *Bmj* 2014;349:g5517.
14. Shaw DE, Green RH, Bradding P. Asthma exacerbations: Prevention is better than cure. *Therapeutics and clinical risk management* 2005;1(4):273-277.
15. Ibricevic A, Pekosz A, Walter MJ, Newby C, Battaile JT, Brown EG, Holtzman MJ, Brody SL. Influenza virus receptor specificity and cell tropism in mouse and human airway epithelial cells. *J Virol* 2006;80(15):7469-7480.
16. Byers DE, Alexander-Brett J, Patel AC, Agapov E, Dang-Vu G, Jin X, Wu K, You Y, Alevy Y, Girard JP, et al. Long-term il-33-producing epithelial progenitor cells in chronic obstructive lung disease. *The Journal of clinical investigation* 2013;123(9):3967-3982.
17. Leiva-Juarez MM, Kolls JK, Evans SE. Lung epithelial cells: Therapeutically inducible effectors of antimicrobial defense. *Mucosal immunology* 2018;11(1):21-34.

- 393 18. Cleaver JO, You D, Michaud DR, Pruneda FA, Juarez MM, Zhang J, Weill PM,
394 Adachi R, Gong L, Moghaddam SJ, et al. Lung epithelial cells are essential effectors of
395 inducible resistance to pneumonia. *Mucosal immunology* 2014;7(1):78-88.
- 396 19. Duggan JM, You D, Cleaver JO, Larson DT, Garza RJ, Guzman Pruneda FA,
397 Tuvim MJ, Zhang J, Dickey BF, Evans SE. Synergistic interactions of tlr2/6 and tlr9
398 induce a high level of resistance to lung infection in mice. *Journal of immunology*
399 2011;186(10):5916-5926.
- 400 20. Evans SE, Tuvim MJ, Fox CJ, Sachdev N, Gibiansky L, Dickey BF. Inhaled
401 innate immune ligands to prevent pneumonia. *British journal of pharmacology*
402 2011;163(1):195-206.
- 403 21. Tuvim MJ, Gilbert BE, Dickey BF, Evans SE. Synergistic tlr2/6 and tlr9 activation
404 protects mice against lethal influenza pneumonia. *PLoS One* 2012;7(1):e30596.
- 405 22. Kirkpatrick CT, Wang Y, Leiva Juarez MM, Shivshankar P, Pantaleon Garcia J,
406 Plumer AK, Kulkarni VV, Ware HH, Gulraiz F, Chavez Cavasos MA, et al. Inducible lung
407 epithelial resistance requires multisource reactive oxygen species generation to protect
408 against viral infections. *mBio* 2018;9(3).
- 409 23. Ware HH, Kulkarni VV, Wang Y, Pantaleon Garcia J, Leiva Juarez M, Kirkpatrick
410 CT, Wali S, Syed S, Kontoyiannis AD, Sikkema WKA, et al. Inducible lung epithelial
411 resistance requires multisource reactive oxygen species generation to protect against
412 bacterial infections. *PLoS One* 2019;14(2):e0208216.
- 413 24. Goldblatt DL, Flores JR, Valverde Ha G, Jaramillo AM, Tkachman S, Kirkpatrick
414 CT, Wali S, Hernandez B, Ost DE, Scott BL, et al. Inducible epithelial resistance against

415 acute sendai virus infection prevents chronic asthma-like lung disease in mice. *British*
416 *journal of pharmacology* 2020.

417 25. Leiva-Juarez MM, Ware HH, Kulkarni VV, Zweidler-McKay PA, Tuvim MJ, Evans
418 SE. Inducible epithelial resistance protects mice against leukemia-associated
419 pneumonia. *Blood* 2016;128(7):982-992.

420 26. Tscherne DM, Jones CT, Evans MJ, Lindenbach BD, McKeating JA, Rice CM.
421 Time- and temperature-dependent activation of hepatitis c virus for low-ph-triggered
422 entry. *J Virol* 2006;80(4):1734-1741.

423 27. Haywood AM, Boyer BP. Sendai virus membrane fusion: Time course and effect
424 of temperature, ph, calcium, and receptor concentration. *Biochemistry*
425 1982;21(24):6041-6046.

426 28. Tai CJ, Li CL, Tai CJ, Wang CK, Lin LT. Early viral entry assays for the
427 identification and evaluation of antiviral compounds. *J Vis Exp* 2015(105):e53124.

428 29. Hines EA, Szakaly RJ, Leng N, Webster AT, Verheyden JM, Lashua AJ,
429 Kendzioriski C, Rosenthal LA, Gern JE, Sorkness RL, et al. Comparison of temporal
430 transcriptomic profiles from immature lungs of two rat strains reveals a viral response
431 signature associated with chronic lung dysfunction. *PLoS One* 2014;9(12):e112997.

432 30. Look DC, Walter MJ, Williamson MR, Pang L, You Y, Sreshta JN, Johnson JE,
433 Zander DS, Brody SL. Effects of paramyxoviral infection on airway epithelial cell foxj1
434 expression, ciliogenesis, and mucociliary function. *Am J Pathol* 2001;159(6):2055-2069.

435 31. Alfaro VY, Goldblatt DL, Valverde GR, Munsell MF, Quinton LJ, Walker AK,
436 Dantzer R, Varadhachary A, Scott BL, Evans SE, et al. Safety, tolerability, and

437 biomarkers of the treatment of mice with aerosolized toll-like receptor ligands. *Front*
438 *Pharmacol* 2014;5:8.

439 32. Bekker V, Scherpbier H, Beld M, Piriou E, van Breda A, Lange J, van Leth F,
440 Jurriaans S, Alders S, Wertheim-van Dillen P, et al. Epstein-barr virus infects b and non-
441 b lymphocytes in hiv-1-infected children and adolescents. *J Infect Dis*
442 2006;194(9):1323-1330.

443 33. Shearer WT, Easley KA, Goldfarb J, Rosenblatt HM, Jenson HB, Kovacs A,
444 McIntosh K. Prospective 5-year study of peripheral blood cd4, cd8, and cd19/cd20
445 lymphocytes and serum igs in children born to hiv-1 women. The p(2)c(2) hiv study
446 group. *J Allergy Clin Immunol* 2000;106(3):559-566.

447 34. Cannon MJ, Stott EJ, Taylor G, Askonas BA. Clearance of persistent respiratory
448 syncytial virus infections in immunodeficient mice following transfer of primed t cells.
449 *Immunology* 1987;62(1):133-138.

450 35. Graham BS, Bunton LA, Wright PF, Karzon DT. Role of t lymphocyte subsets in
451 the pathogenesis of primary infection and rechallenge with respiratory syncytial virus in
452 mice. *The Journal of clinical investigation* 1991;88(3):1026-1033.

453 36. Schmidt ME, Varga SM. Cytokines and cd8 t cell immunity during respiratory
454 syncytial virus infection. *Cytokine* 2018.

455 37. Duan S, Thomas PG. Balancing immune protection and immune pathology by
456 cd8(+) t-cell responses to influenza infection. *Front Immunol* 2016;7:25.

457 38. Hou S, Doherty PC, Zijlstra M, Jaenisch R, Katz JM. Delayed clearance of sendai
458 virus in mice lacking class i mhc-restricted cd8+ t cells. *Journal of immunology*
459 1992;149(4):1319-1325.

39. Schmidt ME, Knudson CJ, Hartwig SM, Pewe LL, Meyerholz DK, Langlois RA, Harty JT, Varga SM. Memory cd8 t cells mediate severe immunopathology following respiratory syncytial virus infection. *PLoS Pathog* 2018;14(1):e1006810.
40. Wang Y, Lobigs M, Lee E, Mullbacher A. Cd8+ t cells mediate recovery and immunopathology in west nile virus encephalitis. *J Virol* 2003;77(24):13323-13334.
41. Connors TJ, Ravindranath TM, Bickham KL, Gordon CL, Zhang F, Levin B, Baird JS, Farber DL. Airway cd8(+) t cells are associated with lung injury during infant viral respiratory tract infection. *Am J Respir Cell Mol Biol* 2016;54(6):822-830.
42. Cannon MJ, Openshaw PJ, Askonas BA. Cytotoxic t cells clear virus but augment lung pathology in mice infected with respiratory syncytial virus. *J Exp Med* 1988;168(3):1163-1168.
43. Kaufmann E, Sanz J, Dunn JL, Khan N, Mendonca LE, Pacis A, Tzelepis F, Pernet E, Dumaine A, Grenier JC, et al. Bcg educates hematopoietic stem cells to generate protective innate immunity against tuberculosis. *Cell* 2018;172(1-2):176-190 e119.
44. Netea MG, Joosten LAB. Trained immunity and local innate immune memory in the lung. *Cell* 2018;175(6):1463-1465.
45. Keeler SP, Agapov EV, Hinojosa ME, Letvin AN, Wu K, Holtzman MJ. Influenza a virus infection causes chronic lung disease linked to sites of active viral rna remnants. *Journal of immunology* 2018;201(8):2354-2368.
46. Evans SE, Xu Y, Tuvim MJ, Dickey BF. Inducible innate resistance of lung epithelium to infection. *Annu Rev Physiol* 2010;72:413-435.

47. Forman HJ, Torres M. Reactive oxygen species and cell signaling: Respiratory burst in macrophage signaling. *American journal of respiratory and critical care medicine* 2002;166(12 Pt 2):S4-8.
48. Huang J, Canadien V, Lam GY, Steinberg BE, Dinanuer MC, Magalhaes MA, Glogauer M, Grinstein S, Brumell JH. Activation of antibacterial autophagy by nadph oxidases. *Proc Natl Acad Sci U S A* 2009;106(15):6226-6231.
49. Yang CS, Shin DM, Kim KH, Lee ZW, Lee CH, Park SG, Bae YS, Jo EK. Nadph oxidase 2 interaction with tlr2 is required for efficient innate immune responses to mycobacteria via cathelicidin expression. *Journal of immunology* 2009;182(6):3696-3705.
50. Paiva CN, Bozza MT. Are reactive oxygen species always detrimental to pathogens? *Antioxid Redox Signal* 2014;20(6):1000-1037.

Table 1

Antibodies	Vendor	Catalogue numbers
CD3	Tonbo	65-0031-U100
CD4	Tonbo	60-0042-U100
CD8	Tonbo	25-0081-U100
Live dead	Tonbo	13-0870-T500
CD25	Biolegend	102038
Foxp3 Treg kit	eBiosciences	72-5775
CD8-Depleting Ab	Bioxell	BE0223-A025
CD19	Biolegend	115507
B220	BD Biosciences	562922
Anti-SeV virus Ab	MBL International	PD029
Ki67	Invitrogen	MA5-14520
cCasp3	Cell signaling	9662S

Figure Legends

Figure 1. Pam2-ODN enhances mouse survival of SeV infection and reduces lung virus burden. (A) Survival of mice treated with PBS or Pam2-ODN one day prior to SeV virus challenge. **(B)** Mouse lung SeV burden 5 days after infection assessed by qPCR for Sendai Matrix (M) gene (Relative quantification, RQ to 18S) relative to 18S.

(C) Time course of lung SeV burden in mice treated with PBS or Pam2-ODN. **(D)** SeV inoculum dependent mouse survival. Data are representative from three independent experiments. n=10 mice per group in survival plots, n=4 mice/group in virus burden experiments. LOQ, limit of quantification. * $p<0.05$, ** $p<0.005$.

Figure 2. Pam2-ODN pretreatment reduces epithelial cell death and proliferation during acute SeV infection. Cleaved caspase 3 (cCasp3) **(A)** or Ki67 **(B)** positive cells in mouse lung epithelium after SeV infection with or without Pam2-ODN treatment (lower panel). EdU positive cells in axial **(C)**, small airways **(D)** and parenchyma **(E)** after SeV infection with or without Pam2-ODN (lower panel). **(F)** Mouse lung histology following SeV challenge with or without Pam2-ODN. n=5 mice per condition. Data are representative from two independent experiments. Scale bar = 100 μm . * $p<0.05$.

Figure 3. Pam2-ODN pretreatment reduces SeV induced lung CD8⁺ T cells. (A) Differential Giemsa staining of BAL cells from mice challenged with SeV with or without Pam2-ODN pretreatment. **(B)** Flow cytometry for CD8⁺ T cells from disaggregated mouse lungs 11 days after SeV infection with or without Pam2-ODN. **(C)** Lung CD8⁺ T cells 11 days after SeV challenge in mice pretreated with PBS or Pam2-ODN. Data are representative of three independent experiments for (A) and of five independent experiments for (B) and (C). * $p<0.05$ compared to PBS+SeV

Figure 4. Pam2-ODN treatment reduces CD8⁺ T cell associated SeV induced

immunopathology. Experimental outline **(A)**, survival **(B)** and percentage of CD8⁺ T

cells **(C)** from disaggregated mouse lungs 10 days after SeV challenge following

pretreatment with PBS or Pam2-ODN and with or without CD8⁺ T cells depleted on day

8 of SeV challenge. **(D)** Lung histology 10 days after SeV challenged with or without

Pam2-ODN treatment and/or CD8⁺ T cells. Data are representative of two independent

experiments. Scale bar = 100 μ m. n=16 mice/group for survival in experiment A and

n=4 mice/group in experiment B. **** p <0.0001 compared to PBS in (c), *** p <0.0005

compared to PBS in (B) and (C), † p <0.05 compared to PBS, * p <0.05 compared to PBS.

Figure 5. Pam2-ODN inhibits SeV without altering attachment. (A) Flow cytometry

to measure virus attachment to epithelial cells 4 h after SeV challenge. **(B)** Percentage

of SeV positive epithelial cells from **(A)**. **(C)** Representative examples of

immunofluorescence for virus attachment. **(D)** Mean fluorescence intensity of SeV-

exposed epithelial cells 4 h after SeV challenge. **(E)** Experimental outline showing viral

attachment and prevention of virus internalization by epithelial cells. **(F)** SeV M gene

expression in untreated MLE-15 cells (left) or primary tracheal epithelial cells (right)

challenged with liberated virus (uninternalized virus particles) from cultures that had

been pretreated with PBS or Pam2-ODN prior to SeV infection 24 h after transfer of

liberated virus to new cells. Data are representative of five independent experiments.

* p <0.05

Figure 6. Pam2-ODN induced reactive oxygen species protects against acute SeV virus infections and immunopathology. SeV burden in MLE-15 cells with or without treatment with Pam2-ODN and/or NADPH inhibitors **(A)** or mitoROS inhibitors **(B)**. **(C)** SeV M gene expression in untreated MLE-15 cells challenged with liberated virus from cells that had been pretreated with PBS or Pam2-ODN with or without mitoROS inhibition. **(D)** Experimental outline. **(E)** Survival of SeV challenge in mice treated with PBS or Pam2-ODN and/or mtROS inhibitors. **(F)** Lung SeV burden measured on day 5 and **(G)** lung CD8⁺ T cells assessed on day 10. Data are representative of three independent experiments. n=13 mice/group in experiment **D** and **E**. *** $p < 0.0001$, ** $p < 0.005$, * $p < 0.01$ compared to PBS, † $p < 0.05$ compared to Pam2-ODN-treated mice without ROS inhibition, * $p < 0.05$.

Wali, Figure 1

Wali, Figure 2

Wali, Figure 3

A

B

C

D

Wali, Figure 6