

Species	Gene(s)	5' Incomplete	3' Incomplete	Approach	Reference
Human	<i>FLOT1</i>	5'	3'	CRISPR/Cas9	[1]
Mouse	<i>Gli3</i>	-	3'	CRISPR/Cas9	[2]
Human	<i>TEAD1</i>	5'	3'	Non-AUG start	[3]
Human	<i>MRVI1</i>	5'	3'	Non-AUG start	[3]
Human	<i>PIM1</i>	5'	3'	Non-AUG start	[3]
Human	<i>EIF4G2</i>	5'	3'	Non-AUG start	[3]
Human	<i>MYC</i>	-	3'	Non-AUG start	[3]
Human	<i>RARB</i>	-	3'	Non-AUG start	[3]
Human	<i>TEAD4</i>	5'	3'	Non-AUG start	[3]
Human	<i>R3HCC1</i>	5'	3'	Non-AUG start	[3]
Mouse	<i>Cttnb1</i>	5'	3'	CRISPR/Cas9	[4]
Human	<i>LMNA</i>	5'	3'	CRISPR/Cas9	[4]
Zebrafish	<i>cd36</i>	5'	3'	ENU	[5]
Zebrafish	<i>pla2g12b</i>	5'	-	ENU	[5]
Zebrafish	<i>smyd1</i>	5'	-	CRISPR/Cas9	[5]
Human	<i>PHACTR1</i>	5'	3'	CRISPR/Cas9	[6]
Human	<i>CTNNB1</i>	5'	3'	CRISPR/Cas9	[7]
Human	<i>AXIN1</i>	5'	3'	CRISPR/Cas9	[7]
Human	<i>LRP5</i>	5'	-	CRISPR/Cas9	[7]
Human	<i>LRP6</i>	5'	3'	CRISPR/Cas9	[7]
Human	<i>PTEN</i>	5'	-	CRISPR/Cas9	[7]
Human	<i>TBK1</i>	-	3'	CRISPR/Cas9	[7]
Human	<i>BAP1</i>	5'	-	CRISPR/Cas9	[7]
Human	<i>TLE3</i>	5'	3'	CRISPR/Cas9	[7]
Human	<i>PPM1A</i>	-	3'	CRISPR/Cas9	[7]
Human	<i>BCL2L2</i>	5'	3'	CRISPR/Cas9	[7]
Human	<i>NGLY1</i>	5'	3'	CRISPR/Cas9	[8]
Human	<i>PI4KB</i>	5'	3'	CRISPR/Cas9	[8]
Human	<i>BRD4</i>	5'	3'	CRISPR/Cas9	[8]
Human	<i>BRD3</i>	-	3'	CRISPR/Cas9	[8]
Human	<i>MTF2</i>	5'	3'	CRISPR/Cas9	[8]
Human	<i>BAZ2A</i>	5'	3'	CRISPR/Cas9	[8]

References:

1. Kapahnke M, Banning A, Tikkanen R: **Random Splicing of Several Exons Caused by a Single Base Change in the Target Exon of CRISPR/Cas9 Mediated Gene Knockout.** *Cells* 2016, **5**.
2. Makino S, Fukumura R, Gondo Y: **Illegitimate translation causes unexpected gene expression from on-target out-of-frame alleles created by CRISPR-Cas9.** *Sci Rep* 2016, **6**:39608.
3. Diaz de Arce AJ, Noderer WL, Wang CL: **Complete motif analysis of sequence requirements for translation initiation at non-AUG start codons.** *Nucleic Acids Res* 2018, **46**:985-994.
4. Mou H, Smith JL, Peng L, Yin H, Moore J, Zhang XO, Song CQ, Sheel A, Wu Q, Ozata DM, et al: **CRISPR/Cas9-mediated genome editing induces exon skipping by alternative splicing or exon deletion.** *Genome Biol* 2017, **18**:108.
5. Anderson JL, Mulligan TS, Shen MC, Wang H, Scahill CM, Tan FJ, Du SJ, Busch-Nentwich EM, Farber SA: **mRNA processing in mutant zebrafish lines generated by chemical and CRISPR-mediated mutagenesis produces unexpected transcripts that escape nonsense-mediated decay.** *PLoS Genet* 2017, **13**:e1007105.
6. Lalonde S, Stone OA, Lessard S, Lavertu A, Desjardins J, Beaudoin M, Rivas M, Stainier DYR, Lettre G: **Frameshift indels introduced by genome editing can lead to in-frame exon skipping.** *PLoS One* 2017, **12**:e0178700.
7. Tuladhar R, Yeu Y, Tyler Piazza J, Tan Z, Rene Clemenceau J, Wu X, Barrett Q, Herbert J, Mathews DH, Kim J, et al: **CRISPR-Cas9-based mutagenesis frequently provokes on-target mRNA misregulation.** *Nat Commun* 2019, **10**:4056.
8. Smits AH, Ziebell F, Joberty G, Zinn N, Mueller WF, Clauder-Munster S, Eberhard D, Falth Savitski M, Grandi P, Jakob P, et al: **Biological plasticity rescues target activity in CRISPR knock outs.** *Nat Methods* 2019, **16**:1087-1093.