Ribosome profiling reveals ribosome stalling on tryptophan codons upon oxidative stress in fission yeast

Angela Rubio¹, Sanjay Ghosh^{1,2}, Michael Mülleder³, Markus Ralser^{3,4}, Juan Mata^{1,5}

¹ Department of Biochemistry, University of Cambridge, UK

² Current address: Institute of Bioinformatics and Applied Biotechnology, Bengaluru, India

³ The Molecular Biology of Metabolism Laboratory, The Francis Crick Institute, London, UK

⁴ Department of Biochemistry, Charité University Medicine, Berlin, Germany

⁵ Corresponding author

ABSTRACT

Modulation of translation is an essential response to stress conditions. We have investigated the translational programmes launched by the fission yeast *Schizosaccharomyces pombe* subject to five environmental stresses: oxidative stress, heavy metal, heat shock, osmotic shock and DNA damage. We also explored the contribution of two major defence pathways to these programmes: The Integrated Stress Response, which directly regulates translation initiation, and the stress-response MAPK pathway. To obtain a genome-wide and high-resolution view of this phenomenon, we performed ribosome profiling of control cells and of cells subject to each of the five stresses mentioned above, both in wild type background and in cells in which the Integrated Stress Response or the MAPK pathway were inactivated.

Translational changes were partially dependent on the integrity of both signalling pathways. Interestingly, we found that the transcription factor Fil1, a functional homologue of the Gcn4 and Atf4 proteins (from budding yeast and mammals, respectively), was highly upregulated in most stresses. Consistent with this result, Fil1 was required for the normal response to most stresses. A large group of mRNAs were translationally downregulated, including many required for ribosome biogenesis. Overall, our data suggest that severe stresses lead to the implementation of a universal translational response, which includes energy-saving measures (reduction of ribosome production) and induction of a Fil1-mediated transcriptional programme.

Surprisingly, ribosomes stalled on tryptophan codons specifically upon oxidative stress, a phenomenon that is likely caused by a decrease in charged tRNA-Tryptophan. Tryptophan stalling led to a mild translation elongation reduction and contributed to the inhibition of initiation by the Integrated Stress Response. Taken together, our results show that different stresses elicit common and specific translational responses, revealing a special and so far unknown role in Tryptophan-tRNA availability.

INTRODUCTION

Cells react to stress situations such as starvation, changes in temperature, or the presence of toxic substances in their environment, by transcriptome and translation remodelling, as well as by the reconfiguration of their metabolism. Translation is the most energy-consuming process of the cell. Therefore, translational control plays an essential role by determining the rate of protein synthesis, which helps shape the composition of the proteome. Compared to transcriptional regulation, translational control of existing mRNAs allows for more rapid changes in protein levels, making this process particularly important upon stress exposure [1]. In addition, as protein synthesis requires a large proportion of the cell energy, its regulation is related to the metabolic status of the cell [2]. Therefore, a tight control of translation is essential to cope with stress situations, and misregulation of this process often leads to disease [3].

In eukaryotes, translational control is often performed at the initiation stage [4], when the AUG start codon is identified and decoded by the initiator tRNA (Met-tRNAⁱ Met). A key regulator of this process is the translation initiation factor eIF2, which is part of the so-called ternary complex (TC) together with the initiator tRNA and GTP. The TC, in complex with other translation initiation factors, binds to the 40S ribosomal subunit to form the 43S preinitiation complex (PIC). The 43S PIC is recruited to the 5' cap of the mRNA by additional initiation factors, leading to the formation of the 48S PIC, which scans the mRNA until the initiation codon is reached. At this step, the 60S subunit binds the complex and GTP is hydrolysed. eIF2-GDP must then be recycled to eIF2-GTP by the GTP/GDP-exchange factor eIF2B. After stress exposure, the eIF2 α subunit is phosphorylated at a specific serine residue (serine 51 in mammals and 52 in fission yeast), and binds to eIF2B acting as a competitive inhibitor, which reduces levels of the ternary complex and triggers a global down-regulation of translation [5,6].

The pathway that regulates translation through eIF2 α phosphorylation is called the Integrated Stress Response. In mammals, there are four eIF2 α kinases (Hri, Gcn2, Pek/Perk and Pkr) that are activated by different stresses and inhibit translation initiation through the phosphorylation of eIF2 α [4]. In the yeast *Saccharomyces cerevisiae*, Gcn2 is the sole eIF2 α kinase [7,8]. In the fission yeast *Schizosaccharomyces pombe*, three eIF2 α kinases (Gcn2, Hri1 and Hri2) show distinct and overlapping activation patterns in response to cellular stresses [9,10]. Whereas Hri2 is mainly activated in response to heat shock and Hri1 at stationary phase in response to nutritional limitation, Gcn2 is the main eIF2 α kinase activated in early exposure to H₂O₂ and MMS [9–11].

In parallel to the general downregulation of translation upon stress, there is an induction of the translation of specific mRNAs, some of them encoding transcription factors, which in turn promote the transcriptional response. Recently, we have shown that amino acid starvation in

S. pombe increases the translation of the transcription factor Fil1, the functional orthologue of Atf4 in mammals and Gcn4 in budding yeast, through the activation of the Gcn2-eIF2 α pathway [12]. Fil1 is required for the transcriptional response to amino acid starvation as well as for normal growth in minimal medium lacking amino acids. Furthermore, Fil1 is regulated in a similar manner through inhibitory upstream ORFs (uORFs) located at the 5'-leader sequence (six uORFs in *fil1*, four in *GCN4* and two in *ATF4*) [12]. In budding yeast and mammalian cells, eIF2 α phosphorylation reduces the abundance of active ternary complexes and the reinitiation of translation occurs after bypassing the inhibitory uORFs, which allows the scanning subunit to reach the main coding sequence [8,13]. Notably, Fil1 does not show sequence similarity to either Gcn4 or Atf4 [12].

Gene expression programs in response to stress are also regulated by Stress Activated Protein Kinases (SAPK). A key player in this pathway in *S. pombe* is the mitogen-activated protein kinase (MAPK) Sty1/Spc1 [14,15], which is homologous to the Hog1 osmo-sensing MAPK in *S. cerevisiae* and to the mammalian and *Drosophila* JNK and p38 SAPKs [16].

Stress signals activate and phosphorylate Sty1, promoting its transient accumulation in the nucleus, where it triggers a wide transcriptional shift of the gene expression program. This transcriptional response is mediated by the Atf1 transcription factor [17–20]. Sty1 also has a role in the translational response to stress. First, cells show higher levels of eIF2 α phosphorylation in the absence of Sty1 [10,21]. Second, Sty1 associates *in vivo* with the translation elongation factor 2 (eEF2) and the translation initiation factor 3a (eIF3a) [22]. Finally, the presence of Sty1 is required to maintain the levels and the phosphorylation of eIF3a, and the recovery of translation levels after stress is less efficient in the absence of Sty1 [21,22].

Translation upon stress can be also regulated at the elongation step through the abundance, modification, and charging levels of transfer RNA (tRNA) [23]. tRNA is the most extensively modified RNA and many post-transcriptional nucleoside modifications occur at the anticodon loop [24]. These modifications can change the stability or localization of tRNAs [25], as well as the fidelity and efficiency of translation [26]. It has also been shown that tRNA fragmentation occurs upon stress in eukaryotes [27] and, in human cells, that tRNA fragments have a role in translation repression [28].

Ribosome profiling (ribo-seq) provides a genome-wide and high-resolution view of translational control. The approach is based on the treatment of translating ribosome–mRNA complexes with a ribonuclease (RNase), in such a way that only RNA fragments protected by bound ribosome survive the treatment. These fragments are then isolated and analysed by high-throughput sequencing. The number of sequence reads that map to a coding sequence, normalized by mRNA levels, provides an estimate of the efficiency of translation for every cellular mRNA [29]. In addition, the position of the reads on the genome identifies the location

of ribosomes on the mRNA. This information can be used to determine relative ribosome occupancies on each codon, which allow the detection of ribosomes stalled on specific codons [12,30,31].

Although some stress-induced gene expression programs have been studied in detail at the genome-wide level [14,15,32–36], the effects of different stresses have been examined in isolation, making comparisons across stresses difficult. Moreover, to our knowledge, there are no systematic studies of the role of major stress-response pathways on genome-wide translation programmes. Here we use *S. pombe* to investigate similarities and differences among the translational responses to five commonly studied stress situations: exposure to H₂O₂ (oxidative stress), cadmium (heavy metal), and methyl methanesulfonate (MMS, genotoxic stress), sorbitol treatment to induce osmotic shock, and heat shock. We perform ribosome profiling in control and stressed cells, under highly controlled and comparable conditions. We also investigate the contribution of the Integrated Stress Response (mediated by eIF2 α phosphorylation) and the stress-responsive MAPK pathway (Sty1) to these translation programs. Overall, we found that translation is typically downregulated, but activated for very few genes. Interestingly, the Fil1 transcription factor is strongly induced at the translational level upon several stresses, in a manner strongly dependent on eIF2 α phosphorylation. Moreover, Fil1 is required for the full implementation of stress-responsive transcription programs. These stresses also cause a rapid translational downregulation of genes involved in ribosome biogenesis and ribosomal proteins.

Surprisingly, we found that ribosomes stall selectively on tryptophan codons upon oxidative stress, a phenomenon that is likely to be caused by a decrease in the levels of charged tRNA-Tryptophan (tRNA-Trp). This specific stalling on tryptophan led to a mild elongation defect. Our results suggest that different stresses elicit common and specific translational responses, both at the initiation and the elongation levels, and uncovers a novel and specific role to tryptophan tRNA availability.

RESULTS

Transcriptomic responses to stress

To investigate genome-wide effects of stress on gene expression, we carried out two independent ribosome profiling experiments (with parallel RNA-seq) of *S. pombe* cells subject to five different stress-inducing treatments for 15 minutes: 0.5 mM H₂O₂ (oxidative stress), 0.5 mM CdSO₄ (heavy metal exposure), temperature shift from 32°C to 39°C (heat shock), 1 M sorbitol (osmotic shock) and 0.02% methyl methanesulfonate (MMS, an alkylating agent that causes DNA damage). To study the contribution of the eIF2 α phosphorylation and Sty1 pathways we performed parallel experiments with the mutant strain *eIF2 α -S52A*, which expresses a non-phosphorylatable version of eIF2 α , and a *sty1 Δ* strain, in which the main

stress-responsive MAPK pathway is inactive. These conditions have been used in the past for microarray-based transcriptomics, and have been shown to elicit robust transcriptional responses within a similar timeframe while maintaining high cell viability [14]. All experiments were carried out using rich medium (YE), as *sty1Δ* cells grow very poorly in minimal medium.

We first examined the transcriptional responses to stress using the RNA-seq data. As we did not perform a global correction for total mRNA abundance, the discussion below refers to relative changes in gene expression. There were more genes significantly up-regulated than down-regulated in every strain and condition (Table 1). For the specific stress conditions that we employed (e.g. concentration of stressor and time), the changes were stronger after heat and cadmium stress, but genes induced by the five stress treatments overlapped significantly with one another. A similar overlap across stresses was observed for repressed genes (highest P value < 2×10^{-5} in wild type cells). Figure S1A shows a heat map with over 1,200 genes that are differentially expressed in at least one condition in wild-type cells. Most of the regulated genes behave similarly in wild type and *eIF2α-S52A* cells. By contrast, the transcriptional response to stress in *sty1Δ* cells was substantially weaker although not completely abolished [14] (Fig. S1A). Consistently, the numbers of induced and repressed genes were comparable among wild type and the *eIF2α-S52A* mutant but were lower in *sty1Δ* cells (Table 1).

The Core Environmental Stress Response (CESR) was defined in a microarray study as those genes induced or repressed two-fold or greater in most of the five stresses analysed [14]. Consistent with the microarray data, the majority of induced and repressed CESR genes, as well as stress-specific genes, were also differentially expressed in our RNA-seq experiments (Fig. 1A, B and Fig. S1B, S1C). Overall, our data show that transcriptomic responses to stress are largely independent of *eIF2α* phosphorylation, but strongly reliant on the MAPK pathway.

General translational responses to stress

Activation of the Integrated Stress Response by stress leads to general translational down-regulation, mediated through *eIF2α* phosphorylation. To investigate the level of activation of this pathway and its effect on global translation levels, we monitored *eIF2α* phosphorylation levels by immunoblotting and analysis of polysome profiles on sucrose gradients. For the specific conditions we used (single stressor concentration and 15-minute exposures), cadmium, heat shock and H_2O_2 treatments led to increased *eIF2α* phosphorylation levels (Fig. 1C) and lower polysomes to subpolysomes ratios (Fig. 1D-G), the latter being indicative of a decrease in translation initiation. Both effects were also detected, although less pronounced, after the MMS and sorbitol treatments (Fig. 1C, 1G). *eIF2α-S52A* cells reduced the polysomes to subpolysomes ratio in response to stress, albeit to a lesser extent than wild-type cells (Fig.

1D-G). This partial dependency on eIF2 α phosphorylation is in agreement with recent reports of *S. pombe* responses to UV and oxidative stress [35,37,38]. By contrast, the Sty1 protein was not required for the downregulation of translation initiation (Fig. 1F, G). Indeed, in some cases (H₂O₂ and sorbitol), *sty1* Δ cells reduced translation more than wild type cells (Fig. 1G). Thus, since polysome profiling cannot distinguish between unaltered active translation and slowed-down elongation, our polysome profiling data suggest that translation is affected at least at the initiation step upon cadmium, heat shock and H₂O₂ exposure. This effect is only partially dependent on eIF2 α phosphorylation. In addition, the Sty1 MAPK pathway may also modulate translation initiation under some stresses [10,21].

Gene-specific translational regulation upon stress exposure

We used ribosome profiling to investigate gene-specific translational control in response to stress. Ribosome-protected fragments (RPFs) were isolated and analysed by high-throughput sequencing, whereas mRNA levels were estimated in parallel by RNA-seq. The number of RPFs mapped to each coding gene, normalized by the corresponding number of RNA-seq reads, was used to calculate relative translation efficiencies (TEs) (Fig. 2A). Genes showing significant changes in TE upon stress treatment were identified using RiboDiff (see Methods), with thresholds of a minimum 1.5-fold change and adjusted P value < 0.01. This approach does not consider global changes in gene expression and, therefore, may overestimate TE values. Nevertheless, these TE values reflect relative changes among conditions and among genetic backgrounds, and identifies genes that behave differently from the majority of transcripts.

Table 1. Number of transcripts regulated after stress exposure on each strain.

	mRNA up			mRNA down			Translation up			Translation down		
	<i>wt</i>	<i>S52A</i>	<i>sty1</i> Δ	<i>wt</i>	<i>S52A</i>	<i>sty1</i> Δ	<i>wt</i>	<i>S52A</i>	<i>sty1</i> Δ	<i>wt</i>	<i>S52A</i>	<i>sty1</i> Δ
CdSO₄	443	410	176	302	269	111	57	37	17	320	212	103
HS	621	468	306	404	364	149	39	18	1	229	124	5
H₂O₂	186	120	25	36	50	18	7	1	3	49	0	65
MMS	142	147	0	8	22	0	4	2	0	1	0	0
Sorbitol	193	198	0	113	109	0	0	0	0	1	1	7

Numbers of mRNAs regulated during five stress conditions in three genetic backgrounds: wild type, *eIF2 α -S52A* and *sty1* Δ cells. mRNA up/down, and translation up/down indicate numbers of mRNAs whose expression is significantly up- or down-regulated at the transcriptome or

translational levels, respectively (see Methods for details). Annotated lists containing these genes are presented in Datasets S1/ S2.

Upon stress, the number of translationally regulated genes was much smaller than those affected at the transcript level (Table 1). In relative terms, unlike transcriptomic responses, the numbers of translationally down-regulated genes were higher than those up-regulated (Table 1). Thus, the gene-specific translational response seems to be targeted to reduce mRNA translation. The treatment that had the strongest effect was cadmium, followed by heat shock and H₂O₂, while fewer changes were detected upon the applied MMS and sorbitol concentrations (Table 1). This trend is generally similar to transcriptional and eIF2 α phosphorylation changes (Fig. 1), indicating that gene expression programs in response to the latter stresses are generally weaker.

We found 82 translationally upregulated genes in wild type cells in at least one stress treatment, 28 of which overlapped with the CESR induced genes (P value < 5x 10⁻¹⁵). Of those 82 genes, four were induced in all the stress conditions except for sorbitol, and 15 were shared across cadmium and heat shock. We could not find any specific GO category enriched within the 82 genes (see Methods). Yet, the data included individual genes that have previously linked to the stress response. An example of cadmium-upregulated gene (both transcriptionally and translationally) was *prf1*, which encodes a transcription factor involved in the oxidative stress response and sexual differentiation [39–41] (Fig. S2A). This regulation is consistent with the generation of reactive oxygen species after cadmium stress [42]. Other interesting genes were related to protein catabolism (*ubp3*, in cadmium and heat shock), autophagy (*atg3*, in cadmium), cell cycle (*srk1*, in cadmium), DNA repair (*dna2*, in heat shock and *uve1*, in H₂O₂ and cadmium), transmembrane transport, carbohydrate and amino acid metabolism. The most translationally upregulated gene was *fil1* [12], which encodes a transcription factor essential for the response to amino acid starvation (Fig. 2A). The role of this gene in stress responses is discussed in detail below. Finally, we found that upon cadmium and heat shock treatments, more than 45% of translationally upregulated genes were also transcriptionally induced, whereas less than 14% of translationally downregulated were also transcriptionally repressed. This coordination of transcriptomic and translational induction (potentiation) has been observed by polysome profiling in budding and fission yeast responses to stress [36,43].

We also identified 382 translationally down-regulated genes (translation efficiency repressed in at least one condition), 149 of which were part of the CESR-repressed list (P value < 5x 10⁻⁷³). Like the induced genes, cadmium, heat shock and H₂O₂ led to stronger effects, whereas the consequences of MMS and sorbitol exposure were very mild (Table 1). The most

extensive overlap was between cadmium and heat shock stress, with 170 genes shared. These genes included *cdr2*, which is involved in the regulation of the G2/M transition through the inhibition of the Wee1 kinase [44] (Fig. S2B). Repression of this gene is consistent with the block of cell cycle progression in response to stress. There was also a substantial overlap (38 genes) among cadmium, heat shock and H₂O₂ stresses. These data indicate that heavy metal exposure, oxidative conditions and heat shock are strong stresses under the conditions used in this work, and that they cause similar translational responses.

Translationally repressed genes were mainly associated with cytoplasmic translation (GO:0002181; $P = 10^{-89}$), ribosome biogenesis (GO:0042254; $P = 10^{-28}$) and ribosome assembly (GO:0042255; $P = 10^{-11}$). Consistently, 110 of the 382 down-regulated genes encoded ribosomal proteins. (Fig. 2A, B). We have previously observed this effect upon nitrogen depletion [30] and amino acid starvation [12], demonstrating that this is a widespread translational response to stress.

Finally, we investigated whether translational responses were dependent on the major stress response pathways. Many TE-upregulated genes were partially induced in both *eIF2 α -S52A* and *sty1 Δ* mutants (Fig. 2C). However, a direct comparison of the induction levels in wild type and mutant cells (Fig. 2E), revealed that the upregulation in the mutant was impaired for most genes. A very similar dependency was observed for downregulated genes (Fig. 2D, F), which were generally less repressed in both mutant backgrounds. Therefore, both the eIF2 α and the MAPK pathways contribute to a normal translational response, but neither of them is sufficient to explain the response in its entirety.

***Fil1* is the major translational responder to stress**

The transcription factor *fil1* was induced very strongly at the translation efficiency level by cadmium, heat shock, and H₂O₂ treatments, while showing small reductions or no changes in mRNA levels (Fig. 3A, B). There was also a weak translational induction after MMS treatment, but none after the sorbitol doses applied. These data mirror the change in eIF2 α phosphorylation in these stresses (Fig. 1C). Consistently, the translational induction of *fil1* was completely dependent on eIF2 α phosphorylation, and very weakly on Sty1 signalling (Fig. 3A). We investigated if this increase in TE was accompanied by higher protein levels. To do this, cells expressing Fil1-TAP from their endogenous locus were used to compare protein levels by immunoblot in the five stress conditions. Consistent with the TE data, there was a strong increase in protein levels after cadmium, heat shock and H₂O₂ treatments, weaker after MMS, and none upon sorbitol (Fig. 3C, D and Fig. S2 C-E). The kinetics of the *fil1* induction was stress-specific, with cadmium and heat shock showing a transient response (peaking at 15 minutes) and H₂O₂ and MMS showing slower responses. In particular, the increase at 60 minutes after H₂O₂ and MMS (Fig. 3D and Fig. S2D) was analogous to the behaviour of CESR-

induced genes under the same conditions, whose induction persisted for an hour [14]. The variety in induction kinetics suggests that Fil1 protein induction may be regulated at other levels in addition to translation. Consistent with the ribosome profiling data, no Fil1 protein induction was detected in *eIF2 α -S52A* cells (Fig. 3C, D and Fig. S2C-E). Surprisingly, Fil1 protein levels were higher in *sty1 Δ* than in wild type cells, even in the absence of stress (Fig. 3C, D and Fig. S2C-E). This may relate to the fact that *sty1 Δ* cells are sensitive to stress [45–48]. Indeed cells lacking this protein show evidence of an induced stress response even under normal laboratory growth conditions [14]. Taken together, these data indicate that *fil1* translational induction upon stress leads to an increase in Fil1 protein levels in an *eIF2 α* -dependent manner, and that the Sty1 MAPK pathway modulates this effect. This response takes place in rich medium, where Fil1 is not required for normal growth [12]. Therefore, Fil1 behaves as a general stress-responsive transcription factor.

We then investigated the role of Fil1 in the transcriptional responses to stress. We performed RNA-seq experiments in the five stress conditions in cells lacking *fil1*, and monitored the behaviour of 165 previously identified Fil1 targets [12]. Note that Fil1 targets were defined as genes that showed lower expression in *fil1 Δ* cells in minimal medium and in the absence of stress [12]. In unstressed cells growing in rich medium, Fil1 targets were expressed at slightly lower levels in the *fil1 Δ* mutant (Fig. 4A-C and Fig. S3A, B). In response to cadmium, heat shock and H₂O₂ treatments, Fil1 targets were expressed at substantially decreased levels in the mutant (Fig. 4A-C and Fig. S3A, B). Consistently, Fil1 target genes overlapped significantly with genes underexpressed in *fil1 Δ* cells at 15 minutes after heat shock (Fig. 4D), and with genes underexpressed 60 minutes after H₂O₂ treatment (Fig. 4E). These data demonstrate that Fil1 promotes the expression of a common group of genes in response to strong stresses (cadmium, heat shock and H₂O₂ treatments).

These results suggest that Fil1 may be important for survival to stress in rich medium. We explored this hypothesis by performing viability assays of wild type and *fil1 Δ* mutant under the five stress treatments. *fil1 Δ* cells were sensitive to high temperature, H₂O₂ and MMS, whereas no difference to wild-type was observed at the sorbitol concentrations applied. Surprisingly, cells lacking Fil1 were resistant to cadmium treatment (Fig. 4F). Although the reason for this phenotype is unclear, deletion of genes encoding other transcription factors involved in stress response (*atf1*) [39,49], and of genes encoding several RNA-binding proteins [50] show similar resistance to cadmium.

As mentioned above, Fil1 is necessary for the normal expression of its targets in response to several stresses (Fig. 4A-C), and Fil1 expression was induced under the same conditions in an *eIF2 α* -dependent manner (Fig. 3 C, D and Fig. S2C). To investigate if *fil1* induction is required for the normal expression of Fil1 targets, we compared the expression levels of *fil1* targets upon stress in wild type and *eIF2 α -S52A* mutants. After 15 minutes of treatment, Fil1

target levels were mildly increased by heat shock and H₂O₂ (but not by cadmium) in an eIF2 α -dependent manner (Fig. S3C). As Fil1 targets tend to be induced more strongly by H₂O₂ at later time points (Fig. 4C), we repeated the experiment upon a 60-minute H₂O₂ exposure. Indeed, this led to a late and stronger induction of Fil1 targets that was almost completely dependent on eIF2 α phosphorylation (Fig. S3D). These results indicate that, at least in some conditions, *fil1* translational upregulation plays a role in the implementation of the normal transcriptional responses to stress.

In addition, cells lacking Sty1 showed increased mRNA levels of Fil1 targets in both stressed and unstressed cells (Fig. S3E). Indeed, the overlap between upregulated genes in unstressed *sty1* Δ cells and Fil1-dependent genes was significant ($P=3.5 \times 10^{-11}$). These data are also consistent with the increased Fil1 protein levels in unstressed *sty1* Δ cells (Fig. 3C-D and Fig. S2C-E). These data suggest that the Sty1 MAPK pathway cross-talks with the Integrated Stress Response, consistent with previous observations [10,21].

Ribosome occupancy on tryptophan codons is increased upon oxidative stress

Ribosome profiling provides information on ribosome locations with codon-level resolution and can thus be used to detect codon-specific ribosome stalling caused by stress conditions. To investigate this level of regulation, we quantified the fraction of ribosomes translating each of the 61 amino acid-encoding codons, normalized by the abundance of the corresponding codon in the transcriptome. This 'relative codon occupancy' reflects the average time spent by the ribosome on each codon. Strikingly, the single codon for tryptophan (TGG) showed strongly increased ribosome occupancy upon H₂O₂ treatment (Fig. 5A, S4A). This enrichment was highly specific, as it was not observed for any other codon, and in any other stress condition (Fig. S4B).

A possible explanation for this observation was that cellular tryptophan levels decreased because of H₂O₂ treatment. We therefore measured intracellular amino acid concentrations by mass spectrometry. We observed a reduction of approximately 20% in tryptophan levels. However, this change was borderline of statistical significance ($P=0.04$), and was smaller than that of other amino acids that did not show any difference in ribosome occupancy of their cognate codons (Fig. 5F, S4C). In addition, mRNA levels and TE of the tRNA-Trp ligase, encoded by *wrs1* gene, remained unaffected.

A second hypothesis was that the levels of tRNA charging could be affected by oxidative stress. To investigate this possibility, we compared the levels of amino-acylated (charged) and deacylated (uncharged) tRNA-Trp. tRNAs were first subjected to periodate oxidation. This treatment leads to the removal of the 3' nucleotide of uncharged tRNAs through β -

elimination, whereas the charged fraction is protected by the amino acid and remains unaltered. The tRNAs are then deacylated at high pH. Thus, uncharged and charged tRNAs show a difference of one nucleotide in size, which can be detected by polyacrylamide gel electrophoresis and Northern blotting [51] (Fig. 5B). tRNA-Trp deacylated samples were used as controls and showed that total levels of tRNA-Trp were not affected by oxidative stress, as well as ruling out the fragmentation of tRNAs [27] (Fig. 5B). By contrast, tRNA-Trp charging levels were reduced more than two-fold after H₂O₂ treatment (Fig. 5B, C). To confirm that this effect is specific of tRNA-Trp, we verified that charged tRNA-His levels remained identical upon H₂O₂ treatment (Fig. 5D). The addition of supplemental tryptophan to rich medium two hours before, or only during the H₂O₂ treatment, prevented the charged tRNA-Trp drop and increased the charging levels (Fig. 5E). Thus, these data indicate that the increase in ribosome occupancy at the tryptophan codon in H₂O₂ stress condition reflects a reduced charged tRNA-Trp fraction.

Decreased tRNA-Trp charging may affect eIF2 α phosphorylation

In *S. cerevisiae* uncharged tRNAs activate the Gcn2 kinase, which phosphorylates eIF2 α to downregulate global translation initiation in response to amino acid starvation [7]. Recently, it has been shown that Gcn2 in *S. pombe* is activated in response to UV and oxidative stress through a mechanism that involves Gcn1 and most likely the binding of tRNAs [52]. Thus, we reasoned that increased levels of uncharged tRNA-Trp upon H₂O₂ treatment might contribute to eIF2 α phosphorylation. To explore this possibility, we compared eIF2 α phosphorylation levels upon H₂O₂ stress in the presence and absence of supplemental tryptophan in rich medium. Consistent with this idea, addition of tryptophan (which increases tRNA-Trp charging levels, see above) caused a reduction of eIF2 α phosphorylation (Fig. 6C). These data suggest that uncharged tRNA-Trp could promote Integrated Stress Response signalling upon H₂O₂ exposure in rich medium, and thus affect both translation initiation and translation elongation.

Differences in charged tRNA-Trp levels cannot affect codon usage because tryptophan is encoded by only one codon, TGG. However, as tryptophan is the least frequent amino acid in proteins (less than 0.015% on average), we asked whether the translation efficiency of genes containing more tryptophan might be affected by lower charged tRNA-Trp levels upon H₂O₂ stress. Genes were ranked by the percentage of tryptophan codons in their coding sequences, and assigned to 11 bins. We then measured the apparent change in translation efficiency upon oxidative stress for each bin. The data showed a trend towards increased translation efficiency changes after H₂O₂ treatment with higher tryptophan content, which was not observed in other stresses (Fig. 6A and Fig. S5). We also ruled out that this trend was determined by transcript lengths (Fig. 6B). The group of genes lacking tryptophan is enriched in genes encoding ribosomal proteins ($P = 10^{-27}$), which have a small transcript size (Fig. 6B).

By contrast, the group with the most tryptophan (0.023-0.056%) showed an enrichment in genes related to lipid biosynthesis (GO:0008610; $P = 6.6 \times 10^{-10}$), protein glycosylation (GO:0006486; $P = 4 \times 10^{-7}$) and cell wall biogenesis (GO:0071554; $P = 1.5 \times 10^{-5}$). This enrichment is consistent with the fact that tryptophan is an amphipathic amino acid, often found in transmembrane domains. Given that translation efficiency as defined above is a relative measurement of the number of ribosomes per transcript (normalised to mRNA abundance), an increase in this parameter in tryptophan-rich genes is likely to reflect a slowdown of translation elongation of these genes rather than an increase in their protein synthesis. Thus, we propose that oxidative stress modulates translation at two levels: at initiation, through the Integrated Stress Response, and at elongation, through tryptophan stalling.

DISCUSSION

The translational landscape of the response to multiple stresses

We present a genome-wide analysis of the translational response of the fission yeast *S. pombe* to five different stresses using highly standardized and comparable conditions. We found that heavy metal exposure, heat shock and oxidative stress led to stronger transcriptional and translational changes than DNA damage and osmotic stress. Moreover, increased levels of eIF2 α phosphorylation correlate with pronounced global downregulation of translation and with strong induction of *fil1* translation. Of course, as we monitored a single time point and condition for each stress, the conclusions about the relative strength of the effects are only valid to the specific experimental conditions used. Despite this caveat, our results indicate that the translational response to stress is mainly directed to repress the translational machinery, with few genes upregulated at this level. Moreover, we found that many differentially translated genes are often regulated in multiple stresses. Thus, our study identifies the key players of a stress response at the translational level and provides insight into the biological response to stress.

Fil1 regulation and role in stress responses

The induction of the Fil1 transcription factor in multiple stress conditions in rich medium was surprising, as Fil1 is a master regulator of the amino acid starvation response (analogous to Gcn4 and Atf4), and is required to maintain a normal growth rate in minimal medium [12]. Cells lacking Fil1 showed similar growth than wild type cells in rich medium, suggesting that Fil1 does not have a role in unstressed cells in rich medium. Strikingly, the *fil1* gene showed much higher translational induction upon cadmium, heat shock and H₂O₂ in rich medium than in amino acid starvation induced by 3-AT in minimal medium (17-fold, 12-fold, 13-fold and 3.8-

fold, respectively). A possible explanation is that *fil1* translation levels may already be higher in minimal medium in unstressed cells. Similarly, the induction of the Fil1 orthologue in *S. cerevisiae* (Gcn4) has been reported not only in amino acid starvation or glucose limitation, but also after MMS (although the conditions were different from the ones used in this work) and H₂O₂ treatments [33,34,53]. These results suggest that metabolic adaptation, mediated by translationally controlled transcription factors such as Fil1 and Gcn4, is an evolutionary conserved part of many stress responses (and not just amino acid starvation).

Fil1-dependent genes are mostly related to amino acid metabolism and transmembrane transport, and show a significant overlap with CESR induced genes. In rich medium, cells lacking Fil1 are unable to regulate the expression of Fil1 targets after cadmium, heat shock and H₂O₂ treatments. In addition, this regulation is also impaired in *eIF2α-S52A* cells, consistent with the complete absence of *fil1* induction. Surprisingly, Fil1 and its targets were upregulated in *sty1Δ* cells (compared to wild type) in unstressed conditions, possibly because the lack of a normal transcriptional response in these cells may lead to stress [14]. Moreover, it has been described that *sty1Δ* cells are sensitive to stress [45–48]. This may also reflect that Sty1 may have a role in the modulation of eIF2α kinases after stress. Indeed, *sty1Δ* cells show increased eIF2α phosphorylation after oxidative stress, which might result in induction of Fil1 [10,21]. Given that Sty1 is also important in adaptation to stress and that nothing is known about Fil1 protein stability or post-translational modifications, it could be also involved in downregulation of Fil1 after stress.

Strikingly, upon cadmium treatment, despite the induction of Fil1, the expression of Fil1-dependent genes in wild-type cells is very weak and *fil1Δ* cells are resistant to this stress. Cadmium stress increase ROS and intracellular oxidative stress [42], but the response is very different to H₂O₂ treatment. For example, cadmium is imported through specific transporters, whereas H₂O₂ freely diffuses into cells. Moreover, H₂O₂ converts into superoxide, and is a substrate for the peroxiredoxin system that is responsible for the oxidation of most proteins. Indeed, *S. pombe* cells lacking transcription factors like Atf1 or Prr1 are sensitive to H₂O₂, but resistant or insensitive to cadmium [39,54]. Additionally, it was reported that Fil1 might drive the response to amino acid starvation partially through the action of downstream transcription factors [12]. Thus, different kinetics in the induction and the expression of Fil1-dependent genes suggest that Fil1 might be modulating transcriptional changes depending on the stress through other transcription factors. We propose that Fil1 acts as a master regulator of several stress conditions, promoting a distinct response for each situation. Further work will be required to unveil the direct targets of Fil1 under each stress condition, and whether Fil1 can activated them in different ways to modulate specific responses.

Ribosomes stall on tryptophan codons upon H₂O₂ treatment

Ribo-seq experiments revealed that oxidative stress caused ribosome stalling on tryptophan codons, which correlated with decreased levels of charged tRNA-Trp. By contrast, intracellular levels of tryptophan were not changed significantly under these conditions, suggesting that tryptophan metabolism is not directly affected by the stress. This suggests that changes in tRNA modifications, which are poorly described in *S. pombe*, or in the tRNA-Trp synthetase activity, may affect the levels of charged tRNA-Trp and cause stalling.

The main eIF2 α kinase activated in response to early exposure to H₂O₂ in *S. pombe* is Gcn2 [10]. The molecular pathway leading to activation of Gcn2 upon nutrition limitation is well understood: Gcn2 is activated by uncharged tRNAs, which bind the histidyl-tRNA synthetase-related domain of Gcn2 at the C-terminus [55–57]. However, Gcn2 is activated by other stresses that are not expected to cause uncharged tRNA accumulation [52], such as UV irradiation or oxidative stress. Under these conditions, the tRNA binding domain of Gcn2 is still required for its activation [52]. Here, we provide evidence for the first time that a nutrient-unrelated stress like oxidative stress causes uncharged tRNA accumulation. In addition, we show that eIF2 α phosphorylation after oxidative stress correlates with uncharged tRNA-Trp accumulation; addition of tryptophan results in a reduction in the levels of both uncharged tRNA-Trp and of eIF2 α phosphorylation. Thus, we propose that the accumulation of uncharged tRNA-Trp upon oxidative stress, which causes a mild elongation defect, is used to regulate initiation through Gcn2 activation.

METHODS

Strains, Growth Conditions, and Experimental Design.

All strains used were prototrophic. Table S1 presents a full list of strains. Standard methods and media were used for *S. pombe* [58]. For all genome-wide stress experiments, *S. pombe* cells were grown in YES medium (supplemented with leucine, uracil and adenine) at 32 °C. Cells were treated for 15 minutes as described below. Heavy metal stress: cadmium sulphate (CdSO₄; 481882; Sigma) was added to a final concentration of 0.5 mM. Oxidative stress: hydrogen peroxide (H₂O₂; H1009; Sigma) was added to a final concentration 0.5 mM. Heat stress: cells were quickly transferred from 32°C to a prewarmed flask in a 39°C water bath. Alkylating agent: methyl methanesulfonate (MMS, 129925, Sigma) at a final concentration of 0.02%. Osmotic stress: cells were grown to OD₆₀₀ = 0.7, and diluted with prewarmed YES 3 M sorbitol (BP-439-500, Fisher) to a final concentration 1 M sorbitol.

For plate drop assays, cells were grown in YES to exponential phase at 32 °C and plated in 10-fold dilutions. Plates were incubated for two days at 32 °C (except the plate incubated at 39 °C for heat shock assays). For tryptophan charging experiments, supplemental tryptophan (DOC0188, Formedium) was added to the culture from a stock 8 g/L in YES to a final concentration of 200 mg/L.

All repeats of genome-wide experiments were independent biological replicates carried out on separate days (see ArrayExpress deposition footnote for a complete list). The following sequencing experiments were performed: (i) ribosome profiling and matching RNA-seq in five stress conditions of three strains (wild-type, *eIF2α-S52A* and *sty1Δ*), and (ii) RNA-seq of *fil1Δ* and cells in five stress conditions.

Amino acid analysis. Amino acid quantification was performed by liquid chromatography selective reaction monitoring (LC-SRM) as described [59].

Protein Analyses. To prepare samples for Western blotting, cells were harvested by filtration, washed with 20% TCA, resuspended in 100 µL of 20% TCA, and frozen. Cell pellets were lysed with 1 mL of acid-treated glass beads in a bead beater (FastPrep-5; MP Biomedicals) at level 7.5 for 15 s, and 400 µL of 5% TCA was added before eluting from the glass beads. Lysates were frozen on dry ice and spun at 18,000 relative centrifugal force for 10 min. Pellets were resuspended in SDS-Tris solution (2% SDS and Tris 0.3M pH 10.7), boiled during 5 min and cleared by centrifugation at maximum speed for 2 min. Protein extract concentrations were measured using Pierce BCA Protein Assay solution (Thermo), and 60-90 µg of total protein were loaded with Laemmli SDS Sample Buffer (reducing) (Alfa Aesar). For western blot analysis, the following antibodies were used: anti-eIF2α (1:500; 9722, Cell

Signalling), anti-Phospho-eIF2 α (1:1,000; 9721, Cell Signalling), and anti-tubulin (1:10,000; sc-23948, Santa Cruz). The secondary antibodies were HRP-conjugated goat anti-mouse IgG (H+L) (1:10,000; 31430, Thermo Fisher) and anti-rabbit IgG (1:10,000; ab-6721, Abcam). TAP tag was detected with peroxide–anti-peroxide complexes (P1291; Sigma). Detection was performed using the enhanced chemiluminescence procedure (ECL kit).

Northern analysis of aminoacyl-tRNA charging. RNA from the indicated conditions was prepared by phenol extraction under acidic conditions using acid buffer (0.3 M sodium acetate pH 5, 10 mM Na₂EDTA) for resuspension of frozen cells, and final acid RNA buffer (10 mM sodium acetate pH 5, 1mM EDTA). As control samples, RNA was deacylated with 0.2 M Tris pH 9.0 for 2h at 37 °C. Periodate oxidation and β -elimination were performed as described [51] using 2 μ g of total RNA. In the final step, all samples were deacylated. tRNAs were detected by Northern blotting in non-acidic conditions: RNA samples were loaded onto a 6.5 % denaturing urea polyacrylamide gel, electrophoresed at 17 W for 70 min and transferred onto an Amersham Hybond-N membrane (GE Healthcare) using the Trans-blot SD semi-dry transfer cell (Biorad) as described [60], but using non-acidic conditions. The oligonucleotide probes used were: tRNA Trp CCA (5'-TGACCCCTAAGTGACTTGAACACTTGA-3'), tRNA His GUG (5'-TGCCCACACCAGGAATCGAACCTGGGT-3') and U5 snRNA as loading control (5'-GCACACCTTACAAACGGCTGTTTCTG-3') [51]. The oligonucleotides were labelled with infrared dyes (IRD-700 or IRD-800) at the 5' end for fluorescence detection using the LICOR Odyssey system. The tRNA loading was quantified as the ratio of upper to lower bands relative to the unstressed condition.

Polysome profiling, ribosome profiling, library preparation and sequencing.

Ribosome-protected fragment (RPF) analyses, preparation of cell extracts, RNase treatment, separation of samples by centrifugation through sucrose gradients, and isolation of protected RNA fragments were performed as described [61]. For polysome profiles, there is no RNase I digestion step, and lysis buffer and sucrose solutions were prepared with double concentration of MgCl₂ (10 mM). Polysome-subpolysome ratio was quantified by measuring the area under the curve relative to the unstressed conditions using Image J software (NIH).

For all RPF samples, gel-purified RNA fragments of around 17-30 nucleotides were treated with 10 units of T4 PNK (Thermo Fisher) in a low-pH buffer (700 mM Tris, pH 7, 50 mM DTT, and 100 mM MgCl₂) for 30 min at 37 °C. ATP and buffer A (Thermo Fisher) were then added for an additional 30 min incubation. RNA fragments were column-purified (PureLink RNA microcolumns; Life Technologies). A total of 100 ng was used as input for the NEXTFLEX

Small RNA Sequencing Kit (Version 3; Bioo Scientific), and libraries were generated following the manufacturer's protocol. For mRNA analyses, total RNA was isolated as described [61]. Total RNA was then depleted from rRNA by using Ribo-Zero Gold rRNA Removal Kit Yeast (Illumina) with 4 µg as input. Finally, 30 ng of rRNA-depleted RNA was used as starting material for the NEXTflex Rapid Directional qRNA-Seq Kit (Bioo Scientific). Libraries were sequenced in an Illumina HiSeq4000 or Novaseq6000 as indicated (ArrayExpress submission).

Data Analysis. Data processing and read alignment were performed as described [12]. Data quantification (number of reads per coding sequence) was carried out by using in-house Perl scripts as described [12]. All statistical analyses were performed using R.

Differential expression analysis was performed by using the Bioconductor DESeq2 package [62]. Raw counts were directly fed to the program, and no filtering was applied. Unless otherwise indicated, a threshold of 10^{-2} was chosen for the adjusted P value, and a cut-off of 2-fold minimal change in RNA levels.

For codon usage analyses, RPF reads were aligned to nucleotide 13 (corresponding to position 1 of the codon in the ribosome P site). Only codons after 90 were used. For each coding sequence, the following calculations were performed: (i) determination of the fraction of RPFs that occupy each codon (RPFs in a given codon divided by total RPFs); (ii) quantification of the relative abundance of each codon on the coding sequence (number of times each codon is present divided by total codon number); and (iii) definition of the normalized codon occupancy by dividing parameter 1 by parameter 2. The average codon enrichments (Fig. 5, S4) were then calculated with data from all coding sequences.

For the analysis of translational efficiencies we used RiboDiff [63]. RiboDiff was provided with raw read counts for each gene, from ribosome profiling and from RNA-seq. To select differentially translated genes, a threshold of 10^{-2} for the adjusted P value, and a cut-off of 1.5-fold were chosen.

Translation efficiency and mRNA ratios were median-centred for plotting. The list of Fil1 targets for Figures 4 and S3 was obtained from, Dataset_S01, repressed genes in *fil1Δ* versus wild type without stress (no 3AT) [12] and only those genes from the lists with at least 20 counts in 80% or more samples were used. Gene set enrichment was performed with AnGeLi [64]. The significance of the overlap between gene lists was calculated using Fisher's exact test.

Data availability. All raw data files have been deposited in ArrayExpress [65] [\[https://www.ebi.ac.uk/arrayexpress/\]](https://www.ebi.ac.uk/arrayexpress/) under accessions: E-MTAB-8746, E-MTAB-8686, E-MTAB-8744, E-MTAB-8745, E-MTAB-8602 and E-MTAB-8583.

ACKNOWLEDGMENTS

This work was supported by a Biotechnology and Biological Sciences (BBSRC) grant to Juan Mata (BB/N007697/1). Markus Ralser was supported by the Francis Crick Institute, which receives its core funding from Cancer Research UK (FC001134), the UK Medical Research Council (FC001134), and the Wellcome Trust (FC001134), as well as specific project funding from the Wellcome Trust (IA 200829/Z/16/Z to M.R.), as well as the German Federal Ministry of Education and Research (BMBF) as part of the National Research Node Mass Spectrometry in Systems Medicine (MSCoreSys 031L0220A). We thank Samuel Marguerat and Caia Duncan for comments on the manuscript and Mathew Peacey and Anno Koetje for help with experiments.

AUTOR CONTRIBUTIONS

A.R., S.G. and J.M. designed the study. A.R., S.G. and M.M. performed and analysed experiments. A.R., M.R. and J.M. analysed data. A.R. and J.M. wrote the manuscript.

CONFLICT OF INTEREST

The authors declare no competing financial interest.

626 **Table S1: Strain and experiment list**

Genotype	Origin	Reference	Usage
972 h-	Lab collection	JU96	Ribo-seq, RNA-seq, drop assay, eIF2 α phosphorylation
<i>eIF2α-S52A::ura4⁺ h-</i>	Sandra López-Avilés	JU1502	Ribo-seq, eIF2 α phosphorylation
<i>sty1::ura4⁺ ura4D18 h-</i>	Sandra López-Avilés, <i>crossed out markers</i>	JU1503	Ribo-seq, eIF2 α phosphorylation
<i>fil1::kanMX6 h-</i>	Lab collection	JU1594	RNA-seq, drop assay
<i>fil1:TAP h+</i>	Lab collection	JU1525	Fil1 expression
<i>fil1:TAP eIF2α S52A::ura4⁺ h-</i>	This work	JU1865	Fil1 expression
<i>fil1:TAP::CRISPR sty1::ura4⁺ h-</i>	This work	JU1749	Fil1 expression

REFERENCES

1. Spriggs KA, Bushell M, Willis AE. Translational Regulation of Gene Expression during Conditions of Cell Stress. *Mol Cell*. 2010;40: 228–237.
doi:10.1016/j.molcel.2010.09.028
2. Lindqvist LM, Tandoc K, Topisirovic I, Furic L. Cross-talk between protein synthesis, energy metabolism and autophagy in cancer. *Curr Opin Genet Dev*. 2018;48: 104–111. doi:10.1016/j.gde.2017.11.003
3. Tahmasebi S, Khoutorsky A, Mathews MB, Sonenberg N. Translation deregulation in human disease. *Nat Rev Mol Cell Biol*. 2018;19: 791–807. doi:10.1038/s41580-018-0034-x
4. Sonenberg N, Hinnebusch AG. Regulation of Translation Initiation in Eukaryotes: Mechanisms and Biological Targets. *Cell*. 2009;136: 731–745.
doi:10.1016/j.cell.2009.01.042
5. Pavitt GD, Ramaiah KVA, Kimball SR, Hinnebusch AG. eIF2 independently binds two distinct eIF2b subcomplexes that catalyze and regulate guanine-nucleotide exchange. *Genes Dev*. 1998;12: 514–526. doi:10.1101/gad.12.4.514
6. Dever TE, Kinzy TG, Pavitt GD. Mechanism and regulation of protein synthesis in *Saccharomyces cerevisiae*. *Genetics*. 2016;203: 65–107.
doi:10.1534/genetics.115.186221
7. Dever TE, Feng L, Wek RC, Cigan AM, Donahue TF, Hinnebusch AG. Phosphorylation of initiation factor 2 α by protein kinase GCN2 mediates gene-specific translational control of GCN4 in yeast. *Cell*. 1992;68: 585–596. doi:10.1016/0092-8674(92)90193-G
8. Hinnebusch AG. Translational Regulation of Gcn4 and the General Amino Acid Control of Yeast . *Annu Rev Microbiol*. 2005;59: 407–450.
doi:10.1146/annurev.micro.59.031805.133833
9. Zhan K, Narasimhan J, Wek RC. Differential activation of eIF2 kinases in response to cellular stresses in *Schizosaccharomyces pombe*. *Genetics*. 2004;168: 1867–1875.
doi:10.1534/genetics.104.031443
10. Berlanga JJ, Rivero D, Martín R, Herrero S, Moreno S, De Haro C. Role of Mitogen-Activated Protein Kinase Sty1 in Regulation of Eukaryotic Initiation Factor 2 α Kinases in Response to Environmental Stress in *Schizosaccharomyces pombe*. *Eukaryot Cell*. 2010;9: 194–207. doi:10.1128/EC.00185-09
11. Martin R, Berlanga JJ, de Haro C. New roles of the fission yeast eIF2 kinases Hri1 and Gcn2 in response to nutritional stress. *J Cell Sci*. 2013;126: 3010–3020.
Available: <http://jcs.biologists.org/cgi/doi/10.1242/jcs.118067>

12. Duncan CDS, Rodríguez-López M, Ruis P, Bähler J, Mata J. General amino acid control in fission yeast is regulated by a nonconserved transcription factor, with functions analogous to Gcn4/Atf4. *Proc Natl Acad Sci*. 2018;115: E1829–E1838. doi:10.1073/pnas.1713991115
13. Vatter KM, Wek RC. Reinitiation involving upstream ORFs regulates ATF4 mRNA translation in mammalian cells. *Proc Natl Acad Sci U S A*. 2004;101: 11269–11274. doi:10.1073/pnas.0400541101
14. Chen D, Toone WM, Mata J, Lyne R, Burns G, Kivinen K, et al. Global transcriptional responses of fission yeast to environmental stress. *Mol Biol Cell*. 2003;14: 214–29. doi:10.1091/mbc.e02-08-0499
15. Chen D, Wilkinson CRM, Watt S, Penkett CJ, Toone WM, Jones N, et al. Multiple Pathways Differentially Regulate Global Oxidative Stress Responses in Fission Yeast. *Mol Biol Cell*. 2008;19: 308–317. doi:10.1091/mbc.E07
16. Toone WM, Jones N. Stress-activated signalling pathways in yeast. *Genes to Cells*. 1998;3: 485–498. doi:10.1046/j.1365-2443.1998.00211.x
17. Shiozaki K, Russell P. Conjugation, meiosis, and the osmotic stress response are regulated by Spc1 kinase through Atf1 transcription factor in fission yeast. *Genes Dev*. 1996;10: 2276–2288. doi:10.1101/gad.10.18.2276
18. Wilkinson MG, Samuels M, Takeda T, Mark Toone W, Shieh JC, Toda T, et al. The Atf1 transcription factor is a target for the Sty1 stress-activated MAP kinase pathway in fission yeast. *Genes Dev*. 1996;10: 2289–2301. doi:10.1101/gad.10.18.2289
19. Gaits F, Degols G, Shiozaki K, Russell P. Phosphorylation and association with the transcription factor Atf1 regulate localization of Spc1/Sty1 stress-activated kinase in fission yeast. *Genes Dev*. 1998;12: 1464–1473. doi:10.1101/gad.12.10.1464
20. Perez P, Cansado J. Cell Integrity Signaling and Response to Stress in Fission Yeast. *Curr Protein Pept Sci*. 2010;11: 680–692. doi:10.2174/138920310794557718
21. Dunand-Sauthier I, Walker CA, Narasimhan J, Pearce AK, Wek RC, Humphrey TC. Stress-activated protein kinase pathway functions to support protein synthesis and translational adaptation in response to environmental stress in fission yeast. *Eukaryot Cell*. 2005;4: 1785–1793. doi:10.1128/EC.4.11.1785-1793.2005
22. Asp E, Nilsson D, Sunnerhagen P. Fission yeast mitogen-activated protein kinase Sty1 interacts with translation factors. *Eukaryot Cell*. 2008;7: 328–338. doi:10.1128/EC.00358-07
23. Pan T. Modifications and functional genomics of human transfer RNA. *Cell Research*. Nature Publishing Group; 2018. pp. 395–404. doi:10.1038/s41422-018-0013-y
24. Jackman JE, Alfonzo JD. Transfer RNA modifications: nature's combinatorial chemistry playground. *Wiley Interdiscip Rev RNA*. 2013;4: 35–48.

- doi:10.1002/wrna.1144
25. Kadaba S, Krueger A, Trice T, Krecic AM, Hinnebusch AG, Anderson J. Nuclear surveillance and degradation of hypomodified initiator tRNA^{Met} in *S. cerevisiae*. *Genes Dev.* 2004;18: 1227–40. doi:10.1101/gad.1183804
26. Patil A, Chan CTY, Dyavaiah M, Rooney JP, Dedon PC, Begley TJ. Translational infidelity-induced protein stress results from a deficiency in Trm9-catalyzed tRNA modifications. *RNA Biol.* 2012;9: 990–1001. doi:10.4161/rna.20531
27. Thompson DM, Lu C, Green PJ, Parker R. tRNA cleavage is a conserved response to oxidative stress in eukaryotes. *RNA.* 2008;14: 2095–2103. doi:10.1261/rna.1232808
28. Ivanov P, Emara MM, Villen J, Gygi SP, Anderson P. Angiogenin-induced tRNA fragments inhibit translation initiation. *Mol Cell.* 2011;43: 613–23. doi:10.1016/j.molcel.2011.06.022
29. Ingolia NT, Ghaemmaghami S, Newman JRS, Weissman JS. Genome-wide analysis in vivo of translation with nucleotide resolution using ribosome profiling. *Science.* 2009;324: 218–223. doi:10.1126/science.1168978
30. Duncan CDS, Mata J. Effects of cycloheximide on the interpretation of ribosome profiling experiments in *Schizosaccharomyces pombe*. *Sci Rep.* 2017;7: 10331. doi:10.1038/s41598-017-10650-1
31. Gerashchenko M V., Lobanov A V., Gladyshev VN. Genome-wide ribosome profiling reveals complex translational regulation in response to oxidative stress. *Proc Natl Acad Sci U S A.* 2012;109: 17394–17399. doi:10.1073/pnas.1120799109
32. Lackner DH, Beilharz TH, Marguerat S, Mata J, Watt S, Schubert F, et al. A network of multiple regulatory layers shapes gene expression in fission yeast. *Mol Cell.* 2007;26: 145–55. doi:10.1016/j.molcel.2007.03.002
33. Claire Mascarenhas, Laura C. Edwards-Ingram, Leo Zeef, Daniel Shenton, Mark P. Ashe and CMG. Gcn4 Is Required for the Response to Peroxide Stress in the Yeast *Saccharomyces cerevisiae*. *Mol Biol Cell.* 2008;19: 2995–3007. doi:10.1091/mbc.E07
34. Natarajan K, Meyer MR, Jackson BM, Slade D, Roberts C, Hinnebusch AG, et al. Transcriptional Profiling Shows that Gcn4p Is a Master Regulator of Gene Expression during Amino Acid Starvation in Yeast. *Mol Cell Biol.* 2001;21: 4347–4368. doi:10.1128/MCB.21.13.4347-4368.2001
35. Shenton D, Smirnova JB, Selley JN, Carroll K, Hubbard SJ, Pavitt GD, et al. Global translational responses to oxidative stress impact upon multiple levels of protein synthesis. *J Biol Chem.* 2006;281: 29011–29021. doi:10.1074/jbc.M601545200
36. Preiss T, Baron-Benhamou J, Ansorge W, Hentze MW. Homodirectional changes in transcriptome composition and mRNA translation induced by rapamycin and heat shock. *Nat Struct Biol.* 2003;10: 1039–1047. doi:10.1038/nsb1015

37. Knutsen JHJ, Rødland GE, Bøe CA, Håland TW, Sunnerhagen P, Grallert B, et al. Stress-induced inhibition of translation independently of eIF2 α phosphorylation. *J Cell Sci.* 2015;128: 4420–4427. doi:10.1242/jcs.176545
38. Boye E, Grallert B. eIF2 α phosphorylation and the regulation of translation. *Current Genetics.* Springer Verlag; 2019. doi:10.1007/s00294-019-01026-1
39. Ohmiya R, Kato C, Yamada H, Aiba H, Mizuno T. A fission yeast gene (prr1+) that encodes a response regulator implicated in oxidative stress response. *J Biochem.* 1999;125: 1061–1066. doi:10.1093/oxfordjournals.jbchem.a022387
40. Nakamichi N, Yanada H, Aiba H, Aoyama K, Ohmiya R, Mizuno T. Characterization of the prr1 response regulator with special reference to sexual development in *Schizosaccharomyces pombe*. *Biosci Biotechnol Biochem.* 2003;67: 547–555. doi:10.1271/bbb.67.547
41. Calvo IA, García P, Ayté J, Hidalgo E. The transcription factors Pap1 and Prr1 collaborate to activate antioxidant, but not drug tolerance, genes in response to H₂O₂. *Nucleic Acids Res.* 2012;40: 4816–4824. doi:10.1093/nar/gks141
42. Cuypers A, Plusquin M, Remans T, Jozefczak M, Keunen E, Gielen H, et al. Cadmium stress: An oxidative challenge. *BioMetals.* 2010;23: 927–940. doi:10.1007/s10534-010-9329-x
43. Lackner DH, Schmidt MW, Wu S, Wolf DA, Bähler J. Regulation of transcriptome, translation, and proteome in response to environmental stress in fission yeast. *Genome Biol.* 2012;13: R25. doi:10.1186/gb-2012-13-4-r25
44. Kanoh J, Russell P. The protein kinase Cdr2, related to Nim1/Cdr1 mitotic inducer, regulates the onset of mitosis in fission yeast. *Mol Biol Cell.* 1998;9: 3321–3334. doi:10.1091/mbc.9.12.3321
45. Millar JB, Buck V, Wilkinson MG. Pyp1 and Pyp2 PTPases dephosphorylate an osmosensing MAP kinase controlling cell size at division in fission yeast. *Genes Dev.* 1995;9: 2117–30.
46. Shiozaki K, Russell P. Cell-cycle control linked to extracellular environment by MAP kinase pathway in fission yeast. *Nature.* 1995;378: 739–743.
47. Degols G, Shiozaki K, Russell P. Activation and regulation of the Spc1 stress-activated protein kinase in *Schizosaccharomyces pombe*. *Mol Cell Biol.* 1996;16: 2870–2877. doi:10.1128/mcb.16.6.2870
48. Degols G, Russell P. Discrete roles of the Spc1 kinase and the Atf1 transcription factor in the UV response of *Schizosaccharomyces pombe*. *Mol Cell Biol.* 1997;17: 3356–3363. doi:10.1128/mcb.17.6.3356
49. Greenall A, Hadcroft AP, Malakasi P, Jones N, Morgan BA, Hoffman CS, et al. Role of fission yeast Tup1-like repressors and Prr1 transcription factor in response to salt

- stress. *Mol Biol Cell*. 2002;13: 2977–89. doi:10.1091/mbc.01-12-0568
50. Hasan A, Cotobal C, Duncan CDS, Mata J, Copenhaver GP. Systematic Analysis of the Role of RNA-Binding Proteins in the Regulation of RNA Stability. *PLoS Genet*. 2014;10. doi:10.1371/journal.pgen.1004684
51. Vakiloroyaei A, Shah NS, Oeffinger M, Bayfield MA. The RNA chaperone La promotes pre-tRNA maturation via indiscriminate binding of both native and misfolded targets. *Nucleic Acids Res*. 2017;45: 11341–11355. doi:10.1093/nar/gkx764
52. Anda S, Zach R, Grallert B. Activation of Gcn2 in response to different stresses. *PLoS One*. 2017;12: e0182143. doi:10.1371/journal.pone.0182143
53. Yang R, Wek SA, Wek RC. Glucose Limitation Induces GCN4 Translation by Activation of Gcn2 Protein Kinase. *Mol Cell Biol*. 2000;20: 2706–2717. doi:10.1128/mcb.20.8.2706-2717.2000
54. Guo L, Ganguly A, Sun L, Suo F, Du L-L, Russell P. Global Fitness Profiling Identifies Arsenic and Cadmium Tolerance Mechanisms in Fission Yeast. *G3 (Bethesda)*. 2016;6: 3317–3333. doi:10.1534/g3.116.033829
55. Dong J, Qiu H, Garcia-Barrio M, Anderson J, Hinnebusch AG. Uncharged tRNA activates GCN2 by displacing the protein kinase moiety from a bipartite tRNA-binding domain. *Mol Cell*. 2000;6: 269–279. doi:10.1016/S1097-2765(00)00028-9
56. Wek SA, Zhu S, Wek RC. The histidyl-tRNA synthetase-related sequence in the eIF2 alpha protein kinase GCN2 interacts with tRNA and is required for activation in response to starvation for different amino acids. *Mol Cell Biol*. 1995;15: 4497–4506. doi:10.1128/mcb.15.8.4497
57. Zhu S, Sobolev AY, Wek RC. Histidyl-tRNA synthetase-related sequences in GCN2 protein kinase regulate in vitro phosphorylation of eIF2. *J Biol Chem*. 1996;271: 24989–24994. doi:10.1074/jbc.271.40.24989
58. Moreno S, Klar A, Nurse P. Molecular genetic analysis of fission yeast *Schizosaccharomyces pombe*. *Methods Enzymol*. 1991;194: 795–823.
59. Mülleder M, Bluemlein K, Ralser M. A high-throughput method for the quantitative determination of free amino acids in *saccharomyces cerevisiae* by hydrophilic interaction chromatography-tandem mass spectrometry. *Cold Spring Harb Protoc*. 2017;2017: 729–734. doi:10.1101/pdb.prot089094
60. Janssen BD, Diner EJ, Hayes CS. Analysis of Aminoacyl- and Peptidyl-tRNAs by Gel Electrophoresis. In: Keiler KC, editor. *Bacterial Regulatory RNA: Methods and Protocols*. Totowa, NJ: Humana Press; 2012. pp. 291–309. doi:10.1007/978-1-61779-949-5_19
61. Duncan CDS, Mata J. The translational landscape of fission-yeast meiosis and sporulation. *Nat Struct Mol Biol*. 2014;21: 641–647. doi:10.1038/nsmb.2843

62. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* 2014;15. doi:10.1186/s13059-014-0550-8
63. Zhong Y, Karaletsos T, Drewe P, Sreedharan VT, Kuo D, Singh K, et al. RiboDiff: Detecting changes of mRNA translation efficiency from ribosome footprints. *Bioinformatics.* 2017;33: 139–141. doi:10.1093/bioinformatics/btw585
64. Bitton DA, Schubert F, Dey S, Okoniewski M, Smith GC, Khadayate S, et al. AnGeLi: A Tool for the Analysis of Gene Lists from Fission Yeast. *Front Genet.* 2015;6: 330. doi:10.3389/fgene.2015.00330
65. Athar A, Füllgrabe A, George N, Iqbal H, Huerta L, Ali A, et al. ArrayExpress update - from bulk to single-cell expression data. *Nucleic Acids Res.* 2019;47: D711–D715. doi:10.1093/nar/gky964

Figure 1

Fig 1. General responses to stress. (A) Scatter plot comparing mRNA levels and translation efficiencies ($\log_2$ ratios stress/control) upon cadmium treatment (15 min) in wild type, eIF2 α -S52A and sty1 Δ genetic backgrounds. The results of a single experiment are shown. CESR-induced genes are plotted in red. **(B)** As in A, but CESR-repressed genes are plotted in blue. **(C)** Western blots comparing eIF2 α phosphorylation levels after the five treatments for the indicated times in wild type cells. In the first two lanes (left), eIF2 α -S52A cells were used as negative control for the anti-eIF2 α phosphorylation antibody. Tubulin was employed as a loading control. **(D to F)** Representative polysome profile traces before and after cadmium treatment (15 min) in wild type **(D)**, eIF2 α -S52A **(E)** and sty1 Δ cells **(F)**. **(G)** Estimation of polysome/subpolysome ratios after five stress treatments (15 min) relative to unstressed conditions.

Figure 2

Fig. 2. Translational regulation upon stress exposure. (A) Scatter plot comparing mRNA levels and translation efficiencies ($\log_2$ ratios stress/control) upon cadmium treatment (15 min) in wild type, *eIF2 α -S52A* and *sty1 Δ* genetic backgrounds. The results of a single experiment are shown. The *fil1* gene is plotted in red and genes encoding ribosomal proteins in blue. (B) Boxplots comparing translation efficiencies of stressed and control cells ($\log_2$ ratios stress/control) of genes encoding ribosomal proteins. Data are shown for wild type, *eIF2 α -S52A* and *sty1 Δ* cells. (C) Comparisons of translation efficiencies of stressed (15 min) and control cells ($\log_2$ ratios stress/control). Only genes that showed significant translational upregulation in wild type cells in at least one stress are displayed. Data are presented for wild type, *eIF2 α -S52A* and *sty1 Δ* cells. (D) As in C, but only genes that showed significant translational downregulation are displayed. (E) As in C, but TE changes have been normalised to those of wild type cells. Dots corresponding to *fil1* gene are shown in red. (F) As in E, but data are displayed for significantly downregulated genes.

Figure 3

Fig. 3. *Fil1* is the major translational responder to stress. (A) Comparison of translation efficiency of the *fil1* gene between stressed and control cells ($\log_2$ ratios stress/control). The dotted lines indicate 1.5-fold changes. Data are presented for wild type, *eIF2 α -S52A* and *sty1 Δ* cells. (B) As in C, but for *fil1* mRNA changes. The dotted lines indicate 2-fold changes (C) Western blots to measure *Fil1*-TAP protein levels after cadmium treatment for the indicated times. Data are presented for wild type, *eIF2 α -S52A* and *sty1 Δ* cells. Tubulin was used as a loading control. (D) As in C, but after H_2O_2 treatment.

Figure 4

Fig. 4. Role of Fil1 in the transcriptional responses to stress. (A to C) Boxplots comparing mRNA levels of stressed and control cells ($\log_2$ ratios stress/control) of Fil1 targets. Data are shown for wild type and $\text{fil}1\Delta$ cells at the indicated times and stresses **(D)** Venn diagram showing the overlap between genes expressed at low levels in $\text{fil}1\Delta$ mutant relative to wild type cells after heat shock (15 min), and Fil1 targets in unstressed cells. The P value of the observed overlap is shown. **(E)** As in D, but genes expressed at low levels in $\text{fil}1\Delta$ mutant relative to wild type after H_2O_2 treatment (60 min) were compared to Fil1 targets in unstressed cells. **(F)** Drop assays of wild type and $\text{fil}1\Delta$ cells plated in the indicated conditions.

Figure 5

Fig. 5. Levels of charged tRNA-Trp are affected by oxidative stress. **(A)** Scatter plots showing $\log_2$ relative codon enrichments before and after H_2O_2 treatment for 15 minutes in wild type cells. The TGG codon encoding tryptophan is plotted in black. **(B)** Representative northern blot for the determination of tRNA-Trp charging levels before and after H_2O_2 exposure. The top blot was hybridised with a probe against tRNA-Trp, and the bottom one with a probe against the U5 snRNA. In the upper blot, the top band corresponds to charged tRNA, and the bottom to the uncharged form. tRNA-Trp samples were either deacylated to remove the linked amino acid from charged tRNAs (sample D) or oxidised to remove the unprotected 3' nucleotides from uncharged tRNAs by beta-elimination (sample B-e) (see Methods for details). U5 snRNA was used as a loading control. **(C)** Quantification of tRNA-Trp charging ratios. Ratios between charged and uncharged tRNA were calculated, and normalised to the ratio in untreated cells. Each dot corresponds to an independent biological replicate ($n = 5$), and the horizontal line indicates the mean. Significance was calculated by using a paired Student's t test. **(D)** As in C, but using a probe against tRNA-His (top panel) or U5 snRNA (bottom). **(E)** Northern blot as in C, to explore the effects of supplementing the culture medium with tryptophan. Cells were grown in the presence of tryptophan for 0, 15 or 120 min, and H_2O_2 was added at the indicated times (0, 15 min) before the end of the incubation with tryptophan. Control deacylated RNA (sample D) is used to identify the location of uncharged tRNA. **(F)** Changes in intracellular tryptophan levels in response to H_2O_2 exposure. Tryptophan levels were measured at the indicated times after H_2O_2 addition to the culture medium. Each dot corresponds to an independent biological replicate ($n = 4$), and the horizontal lines indicate the means. Significance was calculated using a paired Student's t test. No adjustment for multiple testing was performed.

Figure 6

Fig. 6. Oxidative stress affects the translation efficiency of tryptophan-enriched genes, and decreased tRNA-Trp charging may affect eIF2 α phosphorylation. (A) Boxplots showing changes in translation efficiency upon oxidative stress (log₂ TE ratios stress/control, 15 min treatment) according to tryptophan content. Genes were binned into 11 categories based on the fraction of tryptophan in their coding sequences (the first group contains 269 genes without tryptophan, and the other 10 groups have 234 genes each). The horizontal red dashed line indicates the median of the second group. **(B)** As above, but displaying coding sequence lengths. **(C)** Western blots to investigate the effect of tryptophan on eIF2 α phosphorylation levels after H_2O_2 treatment. Cells were treated with H_2O_2 for 15 minutes and supplemental tryptophan was added as indicated. Tubulin was used as a loading control.

Figure S1

Fig. S1. Transcriptomic responses to stress. (A) Heat map of changes in mRNA levels in response to five stress conditions (log₂ ratio stress/control). Data are shown for 1,247 genes that are differentially expressed in at least one stress in wild type (see Methods for details). All data are normalised to the corresponding untreated wild type sample. (B) Heat map displaying enrichment analysis of previously described CCSR gene lists [14] and genes differentially expressed during five stress treatments (our dataset). (C) Comparison of the absolute numbers of induced and repressed genes in our experiments (>2-fold repressed or induced) with the CCSR-induced, CCSR-repressed and stress-specific genes [14].

Figure S2

Fig. S2. Gene-specific translational regulation. (A) Changes in translation efficiency and transcript levels ($\log_2$ ratios stress/control) obtained for the *prf1* gene. Data for two biological replicates are shown. (B) As in A, but for the *cdr2* gene. (C to E) Western blots to measure Fil1-TAP protein levels after heat shock, MMS and sorbitol at the indicated times and genetic backgrounds. Tubulin was used as a loading control.

Figure S3

Fig. S3. Transcription of Fil1 targets relies on eIF2α phosphorylation and is increased in unstressed *sty1Δ* cells. (A and B) Boxplot showing changes in mRNA levels ($\log_2$ ratio stress/control) of genes encoding Fil1 targets after MMS and sorbitol treatments, in wild type and *fil1Δ* strains. (C) As in A and B, but after cadmium, heat shock and H₂O₂ treatments, in wild type and *eIF2α-S52A* strains. (D) As in C, but after H₂O₂ treatment and with an additional time point. (E) As in C, but in the wild type and *sty1Δ* strains.

Figure S4

Fig. S4. Tryptophan codon enrichment is specific for H₂O₂ treatment. (A) Scatter plots showing log₂ relative codon enrichments before and after H₂O₂ treatment for 15 minutes in wild type cells (similar to Fig. 5A, but a different biological replicate). The TGG codon encoding tryptophan is plotted in black. (B) As in A, after cadmium treatment. TGG is displayed in black and highlighted with an arrow. (C) Change in amino acid levels (ratio stress/control) upon oxidative stress exposure. Data are from 4 independent biological replicates (means $\pm$ SD). Tryptophan (W) is highlighted by a red arrow.

Figure S5

Fig. S5. Translation efficiency of tryptophan-enriched genes after other stress treatments. (A to D) Boxplots displaying changes in translation efficiency upon the indicated stresses (log₂ stress/control ratios) according to tryptophan content. Genes were binned into 11 categories based on the fraction of tryptophan in their coding sequences (the first group contains 269 genes lacking tryptophan, and the other 10 groups have 234 genes each). The horizontal red dashed line indicates the median of the second group.