1 **Contrasting response of two *Lotus corniculatus* L. accessions to** 2 **combined waterlogging-saline stress**

3 Antonelli, C.J.^{1#}; Calzadilla, P.I.^{1□}; Campestre M.P.; Escaray F.J.; Ruiz, O.A.*

4 Instituto Tecnológico de Chascomús (IBB-INTECH), UNSAM-CONICET, Av. Intendente Marino
5 Km 8.2 CC 164 (B7130IWA), Chascomús, Bs As, Argentina.

6

7 ¹These authors contributed equally to this work

8 *Corresponding author: Oscar Adolfo Ruiz; ruiz@intech.gob.ar; Tel: (0054) - 2241 430323

9 [#]Present address: Instituto de Fisiología Vegetal (INFIVE), Universidad Nacional de La Plata
10 (UNLP) y Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Diagonal 113
11 No. 495, 1900 La Plata, Argentina

12 [□]Present address: Department of Earth and Environmental Sciences, Faculty of Science and
13 Engineering, University of Manchester, Manchester M13 9PT, UK

14

15 Abstract

16 Waterlogging and salinity impair crops growth and productivity worldwide, being
 17 their combined effects larger than the additive effects of both stresses separately. Recently,
 18 a new *Lotus corniculatus* accession has been collected from a coastal area with a high
 19 frequency of waterlogging-saline stress events. This population is diploid and has potential
 20 to increase nutritional values of *Lotus* cultivars used as forages. Due to its environmental
 21 niche, we hypothesize that this accession would show a better adaptation to combined
 22 waterlogging-saline stress compared to another commonly used tetraploid *L. corniculatus*
 23 (cv. San Gabriel). Shoot and root growth under waterlogging, salinity and combined
 24 waterlogging-saline treatments were addressed, together with chlorophyll fluorescence and
 25 gas exchange measurements. Results showed that salinity and waterlogging effects were
 26 more severe for the tetraploid accession, being the differences larger under the combined
 27 stress condition. In addition, Na^+ , Cl^- and K^+ concentrations were measured in young and old
 28 leaves, and in roots. A larger accumulation of Na^+ and Cl^- was observed under both saline
 29 and combined stress treatments for the tetraploid *L. corniculatus*, for which ion toxicity
 30 effects were evident. The expression of the *NHX1* and *CLC* genes, coding for Na^+ and Cl^-
 31 transporters respectively, was only increased in response to combined stress in the diploid
 32 *L. corniculatus* plants, suggesting that ion compartmentalization mechanisms were induced
 33 in this accession. As a conclusion, the recent characterized *L. corniculatus* might be used for
 34 the introduction of new tolerant traits to combined stresses, in other *Lotus* species used as
 35 forage.

36

37 **Key words:** Legumes; Flooding; Forage; Ion Transporters;
 38 Compartmentalization.

39

40

41

42 **1. Introduction**

43 Human agriculture is facing one of its most important challenges due to an increase
44 in food demand and the more severe consequences of global climate change (Thornton et
45 al., 2014). Under these circumstances, the decrease in biological diversity and widespread
46 monoculture add more uncertainties to the problem (Nunez et al., 2019). In this context,
47 discovering new germplasm adapted to restrictive soils or environments will be of
48 fundamental importance, allowing the introduction of new valuable traits to the existing
49 crop species.

50 Within climate change consequences, global warming alters the precipitations
51 regime, increasing the frequency of flooding and drought events worldwide (Hirabayashi et
52 al., 2013). This phenomenon, together with the sea-level rise in coastal areas (Carter et al.,
53 2006; Martin et al., 2011) and the expansion of high sodicity soils (Ghassemi et al., 1995;
54 Lavado and Taboada, 1987), also increase the frequency of combined waterlogging-saline
55 stress events (Bennett et al., 2009).

56 The effects of waterlogging and salinity stress have been extensively studied in
57 different plant species. Waterlogging limits oxygen diffusion in the rhizosphere (Armstrong
58 and Drew, 2002; Ponnampereuma, 1984), compromises mitochondrial respiration in roots
59 (Gupta et al., 2009; Zabalza et al., 2009) and leads to an ATP production failure for energy
60 demanding processes (Bailey-Serres and Voesenek, 2008; Geigenberger, 2003). Thus, plant
61 responses to waterlogging involve anatomical and morphological changes, such as
62 aerenchyma formation and development of adventitious roots, which aim to increase root
63 oxygenation (Colmer and Voesenek, 2009; McDonald et al., 2002). In particular, in the *Lotus*
64 genus, aerenchyma formation was shown to correlate with flooding tolerance in several
65 species (Antonelli et al., 2019; revised by Striker and Colmer, 2017).

66 Regarding salinity, salt causes osmotic stress and ionic toxicity in most crop plants
67 (Blumwald, 2000; Munns, 2002). Firstly, the osmotic stress occurs as a consequence of the
68 decrease of water availability in roots due to the increase in ion concentration in the soil.
69 Ion toxicity takes place in a second phase, and is caused when Na^+ and/or Cl^- are
70 accumulated in leaves, disrupting protein structure, organelles and affecting cell
71 metabolism (reviewed by Munns and Tester, 2008). While the osmotic stress effect is
72 immediate, the effect due to ionic toxicity is observed after longer exposure time (days or

73 weeks) (Munns, 2002). In addition, salinity may cause nutrient deficiencies or imbalances,
74 due to the competition of Na^+ and Cl^- with nutrients such as K^+ , Ca^{2+} and NO_3^- (Hu and
75 Schmidhalter, 2005).

76 Plants present different strategies to respond to salinity, such as ion exclusion or
77 compartmentalization mechanisms (Munns and Tester, 2008; Teakle and Tyerman, 2010). In
78 the case of the ion exclusion, these mechanisms can be achieved by exclusion transporters,
79 which avoid the entrance of toxic ions into root cells. By contrast, in the
80 compartmentalization mechanisms, ions are accumulated in sub-compartments within the
81 plant cells, avoiding toxic levels to be reached in the cytoplasm and maintaining ionic
82 homeostasis (Munns and Tester, 2008; Teakle and Tyerman, 2010). In both cases, the active
83 transport of ions, with ATP consumption, is required (Munns and Tester, 2008; Teakle and
84 Tyerman, 2010).

85 When waterlogging occurs together with salinity, the combined effects are larger
86 than the additive effects of both stresses separately (Barrett-Lennard, 2003; Teakle et al.,
87 2010). For instance, the energy deficit caused by waterlogging-induced hypoxia will have a
88 direct impact on the Na^+ transporters (Byrt et al., 2007; James et al., 2006), the Na^+/H^+
89 antiporters in the plasma membrane (Martínez-Atienza et al., 2007), and/or the Na^+/H^+
90 antiporters of the tonoplast (like NHX1 ; reviewed by Pardo et al., 2006; Xue et al., 2004). In
91 this sense, it was reported that combined stress largely increases the concentrations of Na^+
92 and Cl^- in plant shoots, as compared with salinity stress alone (Barrett-Lennard, 2003;
93 Barrett-Lennard and Shabala, 2013).

94 Between the regions affected by combined waterlogging-saline stress events, it is
95 possible to mention the Flooding Pampa (Argentina). This area comprises approximately 9
96 million hectares, being one of the most important cattle rearing area in South America
97 (Soriano et al., 1991). Its soil is characterized by a poor nutrient availability, high clay
98 content, salinity and alkalinity. Its frequent exposure to flooding periods, makes the
99 Flooding Pampa a very restrictive environment for crops growth and forage production. As a
100 consequence, the main forage source for cattle bearing consists of natural grasses which are
101 reduced in protein content (Perelman et al., 2001).

102 Different strategies to improve forage quality traits are carried out in regions like the
103 Flooding Pampa, using plant species adapted to constrain conditions. Within them, the use
104 of species of the *Lotus* genus is in the spotlight due to its high plasticity and nutritional

value, being relevant forage alternative in South America, Australia and Europe (Blumenthal and McGraw, 1999; Escaray et al., 2012). In particular, *L. corniculatus* has been extensively used due to its moderate level of proanthocyanidins (PA), which contributes positively to nutritional quality of ruminant diet, increasing protein fraction assimilation, avoiding cattle bloat and reducing intestinal parasites (Foo et al., 1996; McNabb et al., 1996; Min et al., 2003). Nevertheless, the extended use of commercial cultivars of *L. corniculatus* (all tetraploids) has failed due to its edaphic requirements and its susceptibility to different stress conditions such as flooding or salinity (Antonelli et al., 2019; Escaray et al., 2019). By contrast, *L. tenuis* developed a relatively greater tolerance to waterlogging and salinity conditions (revised by Striker and Colmer, 2017), becoming naturalized in the Flooding Pampa. This species presents forage quality comparable to that of other forage legumes (such as *Medicago spp.* or *Trifolium spp.*). However, the low productivity of *L. tenuis* and its PA absence at foliar level, make of *L. corniculatus* a better forage species (revised by Escaray et al., 2012).

Recently, a new *L. corniculatus* accession has been collected from an alkaline-salty area, in the Valencia Albufera in Spain (Escaray et al., 2014). This population is diploid and has been used to improve *Lotus* cultivars through inter-specific hybridization. Although this new germplasm was tested under waterlogging stress (Antonelli et al., 2019) and salinity (Escaray et al., 2019), its stress response under combined waterlogging-saline stress was not yet evaluated. In the present study, we aimed to compare the response to waterlogging, salinity and combined waterlogging-saline stress of the diploid accession of *L. corniculatus* (described by Escaray et al., 2014) and one of the most common *L. corniculatus* cultivars used as forage production in South America (*L. corniculatus* cv. San Gabriel). Due to its environmental niche, we hypothesize that the diploid accession would show a better adaptation to combined waterlogging-saline stress than the commercial *L. corniculatus* cultivar.

2. Materials and Methods

2.1 Plant material

The *L. corniculatus* diploid accession (LcD) corresponds to a wild population collected at the Devesa de El Saler, Valencia (Spain). The taxonomical identity of this population as

belonging to *L. corniculatus* was established and confirmed by different authors (Ballester Ramírez de Arellano, 2015; Escaray et al., 2014). *L. corniculatus* commercial cv. San Gabriel (LcT) is a germplasm obtained by INIA (Instituto de Investigación Agropecuaria de Uruguay, Uruguay).

2.2 Experimental design and growth conditions.

A completely randomized design was performed and the two mentioned plant materials were evaluated under four treatments: 1- control treatment, plants were irrigated with nutrient solution and free drainage (Ctrl); 2- waterlogging with nutrient solution, keeping the water column 3 cm above the pot substrate surface without drainage (WL); 3- salinity, nutrient solution with 150 mM of NaCl and free drainage (NaCl); and 4- combined waterlogging-saline, a combination of the two last treatments (no drainage) (WL+NaCl).

Experiments were initiated from seeds, which were scarified with concentrated sulfuric acid (98%) during 3 min, washed ten times with sterile distilled water and sown in Petri dishes containing water-agar (0.8%). Plates were incubated for 7 days in a growth chamber, with a 16/8 h photoperiod at $24/21 \pm 2^\circ\text{C}$ (day/night) and $55/65 \pm 5\%$ relative humidity. Light intensity ($250 \mu\text{mol photons m}^{-2} \text{s}^{-1}$) was provided by Gro-lux fluorescent lamps (F 40W). Seedlings at the full expanded cotyledon stage were transferred to 300 cm³ pots, containing a mixture of washed sand-perlite (1:1 V/V), and irrigated as explained above. An individual plant per pot was considered an experimental unit. After 21 days of growing, when plants showed five fully-developed leaves, the stress treatments were initiated. Irrigation was performed throughout the experiment with a modified 0.5 x Hoagland's nutrient solution (Hoagland and Arnon, 1950) containing 3 mM KNO₃; 2 mM Ca(NO₃)₂·4H₂O; 1 mM MgSO₄·7H₂O; 0.5 mM NH₄H₂PO₄; 50 μM NaFeO₈EDTA·2H₂O; 4.5 μM MnCl₂, 23 μM H₃BO₃, 0.16 μM CuSO₄·5H₂O, 0.09 μM ZnSO₄·7H₂O, and 0.06 μM Na₂MoO₄·2H₂O.

2.3 Stress treatments

In order to avoid any osmotic shock in saline and combined waterlogging-saline treatments, plants were treated with increasing concentration of NaCl (starting from 25 mM and reaching a final concentration of 150 mM) during 12 days (acclimation period). Overall, plants were subjected to NaCl during 33 days. In all pots subjected to waterlogging, nutrient

solution containing 0.1% (w/v) agar was bubbled with N₂ gas before irrigation. This procedure lowers the dissolved O₂ levels to less than approximately 10% of air-saturated solution (Gibbs and Greenway, 2003), whereas the dilute agar prevents convective movements in the solution (Wiengweera et al., 1997). There were five pots per treatment (n = 5), and the experiments were repeated at least once. Data shown corresponds to the most representative experiment.

2.4 Plant growth measurement

At harvest, plant tissues were divided in young leaves (upper 2 fully expanded leaves), old leaves, stems and roots, and their dry matter was determined after drying at 60°C, until constant weight. By adding the weight of different tissues, total dry weight and the shoot:root ratio were calculated.

2.5 Photosynthesis measurements

One day before harvest, net photosynthetic rate at light saturation (Asat) was measured on the second apical fully expanded leaf (1500 µmol photons m⁻² s⁻¹ illumination, LED light), using a portable photosynthesis system (TPS-2 Portable Photosynthesis System, MA, USA). Then, net photosynthetic rate was relativized to leaf area. For this purpose, leaves were scanned and their area estimated using an image analyser program (Image Pro Plus 4.5). Non-invasive OJIP test (Strasser and Srivastava, 1995) was also performed on the second fully expanded leaf using a Pocket PEA Chlorophyll Fluorimeter (Hansatech Instruments, UK). Leaves were dark adapted for 20 min before analysis and then exposed for 3 s to light at an intensity of 3500 µmol m⁻² s⁻¹. Data were processed by PEA Plus software (Hansatech Instruments, UK) and Windows Excel (Microsoft, WA, USA). The maximum quantum yield of primary photochemistry (Fv/Fm) was calculated.

2.6 Analytical determinations

An aliquot of 10 mg of dried material was used to estimate the concentration of Na⁺ and K⁺ by standard flame photometry (Proehl and Nelson, 1950). Chloride was determined by a thiocyanate-Hg-based colorimetric reaction (Iwasaki et al., 1956). For this, 12.5 mg of powdered dry plant material was extracted in 0.5 mL of a solution containing H₂O₂ (30%):concentrated HNO₃:isoamyl alcohol:H₂O at 1:1:0.08:7.9 (V/V). The extraction was

incubated at room temperature for 15 min, diluted to 5 mL with Milli-Q water and vigorously agitated in a Vortex. Then, 1.5 mL of the extraction mixture was centrifuged (10,000 rpm, 5 min) and the supernatant transferred to another tube. The colorimetric reaction solution contained polyethylene glycol dodecyl ether–water (Brij 35®, 4%):mercuric thiocyanate (4.17 g/L methanol):(NO₃)₃Fe (202 g/L Milli-Q water plus 21 mL concentrated HNO₃):Milli-Q water at 0.05:15:15:70 (V/V). For treatments not involving NaCl, one millilitre of colorimetric reaction was added to 320 µL of the sample supernatant. For treatments that include NaCl, 50 µL of the supernatant were previously diluted with extraction solution to 320 µL. Sample absorbance was determined at 450 nm with a spectrophotometer (Hitachi U-1100), and interpolated into a KCl calibration curve (0, 5, 10, 15, 20 ppm).

2.7 RNA isolation and cDNA synthesis

Total RNA was extracted from frozen apical leaves using a Plant Spectrum Total RNA Kit (Sigma), according to the manufacturer's instructions, and treated with DNase (TURBO DNA-free™ Kit, Ambion). The quality and quantity of RNA were verified by agarose gel electrophoresis and spectrophotometric analysis. The absence of DNA from the RNA samples was tested by the null PCR amplification of the universal rDNA primer pair ITS1/ITS4, as described in Paolocci et al (2006). Then, cDNA from *L. corniculatus* plants was synthesized from 3 µg of total RNA using a Moloney Murine Leukemia Virus Reverse Transcriptase (MMLV-RT) (Promega, WI, USA) and 100 pmol of random hexamers (Pharmacia Biotech), according to supplier's instructions.

2.8 Primer Design

The cDNA sequences related to the *CLC* genes from *L. japonicus* and other model species were downloaded from GenBank. Using own transcriptomic information from *L. corniculatus* species (manuscript in preparation), an *in silico* analysis was performed to obtain a *CLC* homologous sequence for *L. corniculatus*. Deduced *CLC* sequences from *L. corniculatus* and *L. japonicus* were aligned with the BioEdit program (Hall, 1999). The most conserved nucleotide sequence between species was used to design the corresponding primers for qRT-PCR, by the software Primer3Plus (<http://www.bioinformatics.nl/cgi-bin/primer3plus/primer3plus.cgi>). Additionally, a primer pair reported by Teakle et al.

(2010) was used to evaluate the relative expression of the *NHX1* gene. In all cases, the *EF-1 α* gene was used as housekeeping (Escaray et al., 2014).

The primer pairs were initially checked for their specificity and amplification efficiency in both *L. corniculatus* accessions. Only primer pairs that produced the expected amplicon and showed similar PCR efficiency were used in the present study. Primers used for qRT-PCR analysis are listed in Supplementary Table 1.

2.9 Quantitative RT-PCR

An aliquot of 5 μ L of 1:8 diluted cDNA was used in the qRT-PCR reactions, made using 15 μ L of the FastStart Universal SYBR-Green Master Mix (Rox, Roche) and 2.5 pmol of each primer, according to the supplier's instructions. Three biological replicates were performed per sample and gene. Cycling parameters were two initial steps of 50°C for 2 min and 95°C for 2 min, a two-step cycle of 95°C for 15 s and 60°C for 1 min repeated 50 times, and a final step of 10 min at 60°C. This was followed by the dissociation protocol. Amplifications were performed on Mx3005P qPCR System apparatus with the help of the MxPro qPCR Software 4.0 (Stratagene, La Jolla, CA, U.S.A.). For each transcript, the average threshold cycle (Ct) was determined. The gene quantification method was based on the relative expression of the target gene versus the reference *EF-1 α* gene, according to Paolucci et al. (2007).

2.10 Statistical analysis

The experimental results were analyzed using one-way ANOVA for each of the two accessions independently of each other, with four treatments (Ctrl, WL, NaCl and WL+NaCl), followed by Duncan's test ($p < 0.05$). Previously, the assumptions of variance homogeneity and normality were tested for all variables with Levene's and Shapiro-Wilk, respectively. In the case of the analysis of the maximum quantum yield of PSII (Fv/Fm), a t-test was performed. In some cases, comparison between the accessions was also performed using a t-test analysis, for the determined growth variables and ion accumulation measurements. Statistical analysis of gene relative expression was performed based on the pairwise fixed reallocation randomization test ($p < 0.05$) (Pfaffl et al., 2002). In all cases the Infostat software tool was used (Di Rienzo et al., 2010).

3. Results

3.1 Effect of salt, waterlogging and combined waterlogging-saline stress on the growth of the two *L. corniculatus* accessions

The stress treatments imposed visually affected both *L. corniculatus* accessions after 20 days (12 d of salt acclimation plus 8 d of full treatments) (Figure 1). Effects of saline and waterlogging-saline treatments were more evident for LcT plants, for which a smaller number of stems and leaves was observed when compared to control. Similar symptoms were also observed for this accession under waterlogging, although this effect was less strong when compared to the other stress treatments. Regarding LcD, the stress treatments did not reduce the growth of the plants during the first week of the experiment.

After 33 d since the beginning of the experiment (12 d of salt acclimation plus 21 d of full treatments; harvest date), the phenotype of the plants subjected to the combined waterlogging-saline stress was markedly different between the studied accessions (Figure 1), being LcT more severely affected than LcD. While dead leaves and chlorosis were clearly observed for LcT, none of these symptoms were registered for LcD. The phenotype of the others treatments followed the same trend observed for day 8 (not shown).

The effect of the different stress treatments on the growth of both *L. corniculatus* accessions can also be observed through their accumulated shoots and roots dry mass (Figure 2). Although the variability between biological replicates was larger for LcD than for LcT, no significant differences in shoots and roots dry mass accumulation were observed for LcD among stress treatments and its control (Figure 2A and C). By contrast, the stress treatments significantly reduced the shoots and roots dry mass of LcT. This effect was more pronounced in the combined stress treatment, when compared with the waterlogging and salinity conditions imposed independently. The dry mass reduction for LcT between the combined stress treatment and the control was of 75 % and 60 % for shoots and roots, respectively. It is worth mentioning that the total dry mass accumulation was higher for LcT than for LcD (t-test, $p < 0.05$), in control, waterlogging and saline treatments (Figure 2B). However, under the combined stress treatment, both accessions accumulated similar amount of total dry mass (t-test, $p > 0.05$).

The shoot:root ratio was calculated in the different treatments, from the dry mass accumulated for each accession (Figure 2D and Supplementary Figure 1). An increased in this parameter was observed for both LcD and LcT as a response to waterlogging, when

compared to controls. By contrast, a decreased in shoot:root ratio was determined under salinity for both accessions. Regarding the combined stress treatment, a significant decreased compared to controls, was only observed for LcT.

3.2 Photosynthetic response of the LcT and LcD accessions to the different stress treatments.

Net photosynthetic rate at saturating irradiance (Asat) and the maximum quantum yield of PSII (Fv/Fm) were measured in all plants one day before harvest date (Figure 3). The stomatal conductance (gs) was also measured in each case (Supplementary Figure 2). Asat values were only reduced for LcD under salt stress, while for LcT a decreased was observed in all the stress treatments. In LcT, the stronger effect was observed under the salt and the combined stress. Differences observed in gs were comparable to the ones showed by Asat. The larger reduction of gs was observed under the saline stress for both LcD and LcT. However, no effect was observed in gs in neither of the plant accessions under the waterlogging stress, when compared to controls. In the case of the combined stress treatment, the gs value decreased strongly for LcT, but not for LcD.

Regarding the Fv/Fm measurements, no differences were observed for LcD between the different treatments; meanwhile, the maximum yield of PSII was significantly reduced for the salt and combined stress condition in LcT, compared to its control (Figure 3B). Reduction of this parameter was more pronounced under the combined stress condition, reaching values lower than 0.8 (Figure 3B).

3.3 Ions accumulation in different tissues of LcT and LcD subjected to stress conditions

The concentration of Cl^- , Na^+ and K^+ were measured in apical and basal leaves and in roots of LcT and LcD, at the end of the experiment (harvest date) (Figure 4A-I). The Na^+/K^+ was also calculated as a salinity tolerance index, due to the fact that Na^+ can interfere with K^+ homeostasis, affecting several metabolic processes (Figure 4J-L) (Assaha et al., 2017; Shabala and Pottosin, 2014). Cl^- and Na^+ concentrations were not altered in the waterlogging treatment when compared with controls. By contrast, in both *L. corniculatus* accessions, Cl^- and Na^+ increased in the stress treatments where irrigation was supplemented with NaCl, both in leaves and roots (Figure 4A-I). Nevertheless, the ion accumulation in both fractions of leaves (young and old) was always higher than in roots,

showing that the ions were transported to shoots when plants were exposed to saline and waterlogging-saline stress.

For roots and basal leaves tissues, the increase in Cl^- and Na^+ concentration was larger under the combined stress condition, when compared to salinity, in both *L. corniculatus* accessions. Interestingly, Cl^- accumulation in apical leaves of LcD was similar between saline and waterlogging-saline stress (Figure 4A). Nevertheless, in general, a similar pattern of ion accumulation was observed between apical and basal leaves of both LcT and LcD; although Cl^- and Na^+ accumulation was higher for LcT, when compared to LcD, under the combined stress treatment (t-test, $p < 0.05$).

Regarding the K^+ concentration, a reduction in apical and basal leaves was observed in both saline and combined stress treatments, when compared with controls, for LcD. Meanwhile, a significant decrease was only measured under the waterlogging-saline treatment for LcT (Figure 4G and H). In the case of roots, salinity and waterlogging-saline stress reduced the concentration of K^+ in similar proportions for LcD and LcT, compared with their respective controls (Figure 4I). If we consider the changes in both K^+ and Na^+ concentrations, a decrease in K^+/Na^+ ratio was also observed under the salt and combined stress conditions for both LcT and LcD, compared to controls (Figure 4J-L). No differences were observed for the K^+/Na^+ ratio between the waterlogging stress and control treatments for LcD, neither in leaves nor roots, but differences were detected for LcT.

3.4 Effect of stress treatments on the expression of *CLC* and *NHX1* genes.

CLC and *NHX1* are genes coding for Cl^- and Na^+ transporters, respectively, that have been reported to participate in ion homeostasis and salt stress tolerance in different plant species (Bao et al., 2014; Diédhiou and Gollack, 2006; Jossier et al., 2010; Nakamura et al., 2006; Teakle and Tyerman, 2010). The relative expression of both genes was measured on the most contrasting treatments (control and combined waterlogging-saline stress) for both LcD and LcT (Figure 5). The relative expression of *CLC* and *NHX1* was not affected by waterlogging-salt treatment in LcT plants (Figure 5A and B, respectively). Nevertheless, in the LcD accession, the *CLC* gene expression increased three-folds under the combined stress condition, when compared to its control (Figure 5A). Regarding the *NHX1* expression, an

increased expression trend was observed for LcD, although in this case the difference was not significant (Figure 5B).

4. Discussion

In the present study, the stress tolerance of a recent described *L. corniculatus* diploid accession was addressed, with the hypothesis that, due to the characteristics of its ecological niche (Escaray et al., 2014), it would present a better performance under waterlogging-saline conditions than other *L. corniculatus* accession. The larger variability for LcD can be justified with the fact that it was collected from a natural population and did not go through a domestication process such as the commercial cultivar LcT (Escaray et al., 2014). It is worth mentioning that a high variability within individuals of the same population could be an advantage in the search of tolerant traits for forage breeding programs. In fact, the inter-specific hybridization of LcD with other species was already demonstrated to be a useful tool for improving forage legumes (Antonelli et al., 2019; Escaray et al., 2019, 2014).

As a general response, waterlogging stress significantly affected shoot and root dry mass accumulation and net photosynthetic rate in LcT, but did not cause a severe impairment in growth in neither of both accessions. The tolerance extent of *L. corniculatus* to waterlogging stress was previously described (Antonelli et al., 2019; Striker and Colmer, 2017), and is related with the ability of aerenchyma and adventitious root formation. These traits allow O₂ supply and respiration, sustaining metabolism and growth rates under the hypoxic conditions imposed by the water submergence (Colmer and Voesenek, 2009; McDonald et al., 2002). Another common plant response to waterlogging is the higher relative partition of photosynthates to shoots than to roots, resulting in an increased shoot:root ratio (Mendoza et al., 2005; Rubio et al., 1995). This was observed for both LcT and LcD under the waterlogging stress treatment.

On the contrary, plants were severely affected under the saline stress treatments. This effect was more evident in the tetraploid accession, showing a decrease in its dry biomass accumulation and in the photosynthetic parameters evaluated. Both measurements are consistent, showing that a reduction in net photosynthetic rate has a significant impact in dry biomass accumulation. Salinity has been described as a two-phase

stress, where a first osmotic component affects plant growth, followed by a toxic phase due to the accumulation of toxic ions (Munns, 2002). Due to the salinity acclimation period that was imposed before the salinity treatments (see Materials and Methods), the osmotic phase was probably reduced and the decrease in the photosynthetic capacity of the plants could be attributed to the accumulation of toxic levels of ions. Nevertheless, although an increase in the concentration of Cl^- and Na^+ was observed under salt stress in LcD and LcT, the maximum quantum yield of PSII was not severely affected under this condition. Similar results were reported in the halophyte *Suaeda salsa* (Lu et al., 2002), and for LcD under comparable saline stress conditions (Escaray et al., 2019).

Considering the decrease in stomatal conductivity, an osmotic effect due to salinity cannot be discarded, and the reduction in the photosynthetic capacity of both accessions could be also attributed to a lower CO_2 availability. Similar results were observed in other species, where it was concluded that stomatal aperture was the main factor limiting leaf photosynthetic capacity in NaCl-treated plants (Loreto et al., 2003; Meloni et al., 2003). The lack of alterations in PSII functionality under salt stress might be the result of an increase in photoprotection mechanisms, such as Non-Photochemical Quenching and cyclic electron flow (Bencke-Malato et al., 2014; Dionisio-Sese and Tobita, 2000), or the increase in the activity of ROS scavenging mechanisms in the chloroplast (Badawi et al., 2004; Meloni et al., 2003). The latter was previously reported to take place in other *Lotus* species, although under low temperature conditions (Calzadilla et al., 2016).

Despite salinity similarly affected both *L. corniculatus* accessions, the larger contrasting response between accessions was observed under the combined stress treatment. This condition mainly affected LcT plants; while for LcD, unexpectedly, waterlogging seemed to reduce the negative effects caused by salinity (Figure 1). The better performance of LcD compared to LcT could be, at least partially, explained by a lower accumulation of Cl^- and Na^+ ions in their leaves. Combined waterlogging-saline stress cause an increase in these ions concentrations when compared to salinity (Barrett-Lennard, 2003), which was, in general, more pronounced for LcT plants.

The saline stress tolerance has been correlated to the capacity of Cl^- exclusion in different legume species (Teakle and Tyerman, 2010), including some of the *Lotus* genus (Sanchez et al., 2010; Teakle et al., 2006). Moreover, the negative correlation between Cl^- concentration and stress tolerance has been reported to be even stronger than the one

existing for Na^+ , in *Trifolium* (Rogers et al., 1997), *Medicago* (Sibole et al., 2003), *Glycine* (Luo et al., 2005) and even *Lotus* (Teakle et al., 2007, 2006). Xu et al. (1999) defined critical values for Cl^- toxicity above $200 \mu\text{mol.g}^{-1}\text{DM}$ for sensitive crops species, such as rice, wheat, alfalfa and peanut; and above $1400 \mu\text{mol.g}^{-1}\text{DM}$ for the most tolerant ones, i.e. sugar beet and tomato. Under the combined stress condition, the values measured for LcT were twice as high as the one observed for LcD, although in both cases Cl^- concentration reached the levels defined as toxic (Xu et al., 1999). This toxic effect was clearly observed in the reduction of the maximum quantum yield of PSII in LcT, in the further reduction of dry biomass accumulation and in the appearance of chlorotic and senesced leaves in this accession.

By contrast, no symptoms of ion toxicity were observed in LcD under the waterlogging-saline treatment, despite the high levels of Cl^- concentration in leaves. Even more, surprisingly, a net photosynthetic rate increased was observed for this accession under the combined stress, when compared with the salinity treatment alone. Although it is known that waterlogging deepens the effects caused by salinity stress (Barrett-Lennard, 2003; Bennett et al., 2009), similar results regarding the amelioration of salinity were recently reported in *Mentha aquatic* (Haddadi et al., 2016). These results were suggested to be the consequence of the priming of an antioxidant response, which could help to increase membrane stability and reduce the toxic effects of NaCl (Haddadi et al., 2016). However, further studies are needed to understand how photosynthesis acclimates to the combined stress condition.

The better tolerance of LcD to the combined stress condition could be explained through a better compartmentalization of toxic ions within the cells. Different subcellular compartmentalization mechanisms have been described in the plant response to saline stress (revised by Munns and Tester, 2008; Teakle and Tyerman, 2010), including the participation of ion transporters such as *CLC* and *NHX1*, for Cl^- and Na^+ respectively (Bao et al., 2014; Jossier et al., 2010; Nakamura et al., 2006). In this sense, expression of *CLC* was strongly correlated with salinity tolerance in grapevine (Henderson et al., 2014), while overexpression of *NHX1* was demonstrated to enhance salt stress in *Arabidopsis* and even *L. corniculatus* (Liu et al., 2010; Sun et al., 2006). These transgenic plants showed a higher Na^+ accumulation in their tissues, but a higher photosynthetic capacity. The amelioration of the toxic effect of Na^+ was ascribed to its compartmentalization into the vacuole (Liu et al.,

2010), which provides an efficient way to alleviate Na^+ excess in the cytosol, and helps keeping cellular turgence under stress (Flowers et al., 1977). Furthermore, overexpression of tonoplast related proteins was also recently associated with a higher tolerance to salinity due to ion compartmentalization in legumes, such as *L. corniculatus* (Bao et al., 2014) and *M. sativa* (Bao et al., 2016).

The gene expression of *CLC* and *NHX1* was measured in order to approach a possible mechanism of waterlogging-saline tolerance in both *L. corniculatus* accessions. The expression of *CLC* was significantly increased under stress in LcD, while a similar expression trend was also observed for *NHX1*; meanwhile, plants of LcT showed no effect in relative expression of both genes between treatments. Chloride channels are involved into intracellular compartmentalization of Cl^- , sequestering this anion to prevent toxic levels in cytoplasm. Our results suggest that, in LcD, increasing the expression of *CLC* might favour the compartmentalization of Cl^- , improving its waterlogging-saline stress response. A similar effect might be taking place for Na^+ and its transporter *NHX1*. In this sense, a previous work showed that *NHX1* is also involved in *L. tenuis* response to waterlogging-salinity stress, and that its expression levels can justify the better tolerance of *L. tenuis* when compared to *L. corniculatus* (commercial cv. San Gabriel) (Teakle et al., 2010). Nevertheless, in this case, the differences in *NHX1* expression between *L. tenuis* and *L. corniculatus* were shown in root tissues.

It is worth mentioning that the active transport of ions against their concentration gradient implies the consumption of energy (Colmer and Flowers, 2008; Munns and Tester, 2008). Thus, the hypoxic condition imposed by flooding severely affects respiration, the production of ATP and, as a consequence, affects ions compartmentalization as a salinity stress response (Barrett-Lennard, 2003; Kotula et al., 2015). This is one of the reasons why a more severe effect of salinity is generally observed when is combined with waterlogging (Barrett-Lennard, 2003; Teakle et al., 2007). Wetland halophytes plants were reported to have a lower increase in Na^+ and Cl^- ions due to a higher oxygenation of their roots (Colmer and Flowers, 2008). Interestingly, Antonelli et al. (2019) measured the response of different *Lotus* species to waterlogging stress, including the ones addressed in the present study, and found that LcD shows almost three times the percentage of root aerenchyma than LcT. These results could imply a better root oxygenation in the first mentioned accession, which would allow respiration and ATP generation under hypoxic conditions. The higher root

aerenchyma formation of LcD, when compared to LcT, could also justify its better response to the waterlogging-saline stress condition. Similar results were obtained for *L. tenuis*, when compared to LcT, by Teakle et al. (2007).

There is a general agreement that cytoplasmic high K^+/Na^+ ratio is a good indicator of low salt damage and high salinity tolerance (Maathuis and Amtmann, 1999; Munns and Tester, 2008). The similar physicochemical characteristics between K^+ and Na^+ affect a wide range of metabolic processes, such as enzymatic reactions and protein synthesis, and maintaining the K^+/Na^+ ratio under stress conditions is of key importance to maintain K^+ homeostasis (Almeida et al., 2017; Maathuis and Amtmann, 1999; Shabala and Cuin, 2008). However, it was previously reported that for halophytes and some glycophytes plants, the K^+/Na^+ ratio is not a good parameter to assess salinity tolerance (Colmer and Voesenek, 2009).

Our results show that, in leaves, the K^+/Na^+ was only reduced in LcD when the irrigation solution was supplemented with NaCl; while in LcT, K^+/Na^+ was decreased in all of the stress treatments imposed. These results are in agreement with a better response of LcD to waterlogging, when compared to LcT (our own results; Antonelli et al., 2019). Nevertheless, for the salinity and combined stress treatments, the K^+/Na^+ ratio values did not differ between the evaluated *Lotus* accessions. Thus, the K^+/Na^+ , at least at tissue level, is not a strong indicator of salinity tolerance for species of the *Lotus* genus, in agreement with results obtained by other authors (Rejili et al., 2007).

Although salinity stress reduces K^+ and increases Na^+ concentration in plant tissues, this not necessarily implies changes in their cytoplasmic concentrations (Flowers et al., 2015). For instance, a significant proportion of K^+ of the leaves is located in the vacuole and is responsible of keeping cell turgence (Andrés et al., 2014; Barragán et al., 2012). In certain plant species, such as *Mesembryanthemum cristallinum* and *Suaeda maritima* (and probably *Lotus*), the role of K^+ could be replaced by Na^+ , which would allow maintaining ion homeostasis in the cytosol (Kronzucker et al., 2013; Leigh and Wyn Jones, 1986). As a consequence, for plants with a high salinity response implying subcellular ion compartmentalization, changes in the K^+/Na^+ ratio might not necessarily imply alteration in K^+ homeostasis and stress sensitivity.

5. Conclusions

In the present study, the stress tolerance to waterlogging, salinity and combined waterlogging-saline stress was evaluated in two accessions of *Lotus corniculatus*. Our results showed that the diploid accession, obtained from an environmental niche naturally affected by waterlogging and salinity, has a better response to all the stress conditions evaluated, when compared to LcT. This contrasting response was more evident under the combined stress treatment. A lower decrease in dry biomass accumulation and absence of stress symptoms were observed in treated LcD plants, when compared with LcT, which could be ascribe to a lower photoinhibitory effect and lower Cl^- and Na^+ accumulation in leaves. In addition, the better response could be justified through the triggering of ion subcellular compartmentalization mechanisms, which was suggested through the increased expression levels of the *CLC* and *NHX1* transporters genes. In this sense, we suggest that the K^+/Na^+ ratio is not a good indicator of salinity tolerance in plants where ion compartmentalization responses take place. As a conclusion, the higher adaptability of the *L. corniculatus* diploid accession to combined waterlogging-saline stress was demonstrated when compared to another *L. corniculatus* commercial cultivar. Thus, the recently characterized *L. corniculatus* accession could be used to introduce new tolerant traits to waterlogging-saline stress, in other *Lotus* species commonly used as forage.

Acknowledgements

This work was supported by grants from the *Agencia Nacional de Promoción Científica y Tecnológica* (ANPCYT-Argentina)/FONCyT-PICTs 1560, 1612, 3648 and 3718 and *Consejo Nacional de Investigaciones Científicas y Técnicas* (CONICET-Argentina)/ PIP 0980. C.J.A. and P.I.C. and M.P.C were doctoral fellows, whereas F.J.E. and O.A.R. are members of the Research Career of CONICET.

Authors contribution

C.J.A. designed and performed all the experiments, and analyzed data; P.I.C. analyzed data; M.P.C performed some experiments and analyzed data; F.J.E design experiments and analyzed data; O.A.R. conceived the project, designed and supervised all the experiments. The article was written by O.A.R, C.J.A. and P.I.C.

References

- Almeida, D.M., Oliveira, M.M., Saibo, N.J.M., 2017. Regulation of Na⁺ and K⁺ homeostasis in plants: towards improved salt stress tolerance in crop plants. *Genet. Mol. Biol.* 40, 326–345.
- Andrés, Z., Pérez-Hormaeche, J., Leidi, E.O., Schlücking, K., Steinhorst, L., McLachlan, D.H., Schumacher, K., Hetherington, A.M., Kudla, J., Cubero, B., 2014. Control of vacuolar dynamics and regulation of stomatal aperture by tonoplast potassium uptake. *Proc. Natl. Acad. Sci.* 111, E1806–E1814.
- Antonelli, C.J., Calzadilla, P.I., Vilas, J.M., Campestre, M.P., Escaray, F.J., Ruiz, O.A., 2019. Physiological and anatomical traits associated with tolerance to long-term partial submergence stress in the Lotus genus: responses of forage species, a model and an interspecific hybrid. *J. Agron. Crop Sci.* 205, 65–76.
- Armstrong, W., Drew, M.C., 2002. Root growth and metabolism under oxygen deficiency, in: *Plant Roots*. CRC Press, pp. 1139–1187.
- Assaha, D.V.M., Ueda, A., Saneoka, H., Al-Yahyai, R., Yaish, M.W., 2017. The role of Na⁺ and K⁺ transporters in salt stress adaptation in glycophytes. *Front. Physiol.* 8, 509.
- Badawi, G.H., Kawano, N., Yamauchi, Y., Shimada, E., Sasaki, R., Kubo, A., Tanaka, K., 2004. Over-expression of ascorbate peroxidase in tobacco chloroplasts enhances the tolerance to salt stress and water deficit. *Physiol. Plant.* 121, 231–238.
- Bailey-Serres, J., Voesenek, L., 2008. Flooding stress: acclimations and genetic diversity. *Annu. Rev. Plant Biol.* 59, 313–339.
- Ballester Ramírez de Arellano, J., 2015. Requisitos edáficos de algunas especies de interés especial de Las Malladas del Parque Natural de La Albufera.
- Bao, A.-K., Wang, Y.-W., Xi, J.-J., Liu, C., Zhang, J.-L., Wang, S.-M., 2014. Co-expression of xerophyte *Zygophyllum xanthoxylum* ZxNHX and ZxVP1-1 enhances salt and drought tolerance in transgenic *Lotus corniculatus* by increasing cations accumulation. *Funct. plant Biol.* 41, 203–214.
- Bao, A., Du, B., Touil, L., Kang, P., Wang, Q., Wang, S., 2016. Co-expression of tonoplast Cation/H⁺ antiporter and H⁺-pyrophosphatase from xerophyte *Zygophyllum xanthoxylum* improves alfalfa plant growth under salinity, drought and field conditions. *Plant Biotechnol. J.* 14, 964–975.
- Barragán, V., Leidi, E.O., Andrés, Z., Rubio, L., De Luca, A., Fernández, J.A., Cubero, B., Pardo, J.M., 2012. Ion exchangers NHX1 and NHX2 mediate active potassium uptake into vacuoles to regulate cell turgor and stomatal function in *Arabidopsis*. *Plant Cell* 24, 1127–1142.
- Barrett-Lennard, E.G., 2003. The interaction between waterlogging and salinity in higher plants: causes, consequences and implications. *Plant Soil* 253, 35–54.
- Barrett-Lennard, E.G., Shabala, S.N., 2013. The waterlogging/salinity interaction in higher plants revisited—focusing on the hypoxia-induced disturbance to K⁺ homeostasis. *Funct. Plant Biol.* 40, 872–882.

- 586 Bencke-Malato, M., Cabreira, C., Wiebke-Strohm, B., Bücken-Neto, L., Mancini, E., Osorio,
587 M.B., Homrich, M.S., Turchetto-Zolet, A.C., De Carvalho, M.C., Stolf, R., Weber, R.L.,
588 Westergaard, G., Castagnaro, A.P., Abdelnoor, R. V, Marcelino-Guimarães, F.C., Margis-
589 Pinheiro, M., Bodanese-Zanettini, M.H., 2014. Genome-wide annotation of the soybean
590 WRKY family and functional characterization of genes involved in response to
591 Phakopsora pachyrhizi infection. BMC Plant Biol. 14, 236.
592 <https://doi.org/10.1186/s12870-014-0236-0>
- 593 Bennett, S.J., Barrett-Lennard, E.G., Colmer, T.D., 2009. Salinity and waterlogging as
594 constraints to saltland pasture production: a review. Agric. Ecosyst. Environ. 129, 349–
595 360.
- 596 Blumenthal, M.J., McGraw, R.L., 1999. Lotus Adaptation, Use, and Management, in: Trefoil:
597 The Science and Technology of Lotus, CSSA Special Publication SV - 28. Crop Science
598 Society of America and American Society of Agronomy, Madison, WI, pp. 97–119.
599 <https://doi.org/10.2135/cssaspecpub28.c6>
- 600 Blumwald, E., 2000. Sodium transport and salt tolerance in plants. Curr. Opin. Cell Biol. 12,
601 431–434.
- 602 Byrt, C.S., Platten, J.D., Spielmeyer, W., James, R.A., Lagudah, E.S., Dennis, E.S., Tester, M.,
603 Munns, R., 2007. HKT1; 5-like cation transporters linked to Na⁺ exclusion loci in wheat,
604 Nax2 and Kna1. Plant Physiol. 143, 1918–1928.
- 605 Calzadilla, P.I., Signorelli, S., Escaray, F.J., Menéndez, A.B., Monza, J., Ruiz, O.A., Maiale, S.J.,
606 2016. Photosynthetic responses mediate the adaptation of two Lotus japonicus
607 ecotypes to low temperature. Plant Sci. 250.
608 <https://doi.org/10.1016/j.plantsci.2016.06.003>
- 609 Carter, J.L., Colmer, T.D., Veneklaas, E.J., 2006. Variable tolerance of wetland tree species to
610 combined salinity and waterlogging is related to regulation of ion uptake and
611 production of organic solutes. New Phytol. 169, 123–134.
- 612 Colmer, T.D., Flowers, T.J., 2008. Flooding tolerance in halophytes. New Phytol. 179, 964–
613 974.
- 614 Colmer, T.D., Voesenek, L., 2009. Flooding tolerance: suites of plant traits in variable
615 environments. Funct. Plant Biol. 36, 665–681.
- 616 Di Rienzo, J.A., Casanoves, F., Balzarini, M.G., Gonzalez, L., Tablada, M., Robledo, C.W., 2010.
617 InfoStat versión. Grup. InfoStat, FCA, Univ. Nac. Córdoba, Argentina.
- 618 Diédhiou, C.J., Gollack, D., 2006. Salt-dependent regulation of chloride channel transcripts
619 in rice. Plant Sci. 170, 793–800.
- 620 Dionisio-Sese, M.L., Tobita, S., 2000. Effects of salinity on sodium content and
621 photosynthetic responses of rice seedlings differing in salt tolerance. J. Plant Physiol.
622 157, 54–58.
- 623 Escaray, F.J., Antonelli, C.J., Carrasco, P., Ruiz, O.A., 2019. Interspecific hybridization
624 improves the performance of Lotus spp. under saline stress. Plant Sci. 283, 202–210.
- 625 Escaray, F.J., Menendez, A.B., Gárriz, A., Pieckenstain, F.L., Estrella, M.J., Castagno, L.N.,

- 626 Carrasco, P., Sanjuán, J., Ruiz, O.A., 2012. Ecological and agronomic importance of the
627 plant genus *Lotus*. Its application in grassland sustainability and the amelioration of
628 constrained and contaminated soils. *Plant Sci.* 182, 121–133.
- 629 Escaray, F.J., Passeri, V., Babuin, F.M., Marco, F., Carrasco, P., Damiani, F., Pieckenstain, F.L.,
630 Paolocci, F., Ruiz, O.A., 2014. *Lotus tenuis* x *L. corniculatus* interspecific hybridization as
631 a means to breed bloat-safe pastures and gain insight into the genetic control of
632 proanthocyanidin biosynthesis in legumes. *BMC Plant Biol.* 14, 1.
- 633 Flowers, T.J., Munns, R., Colmer, T.D., 2015. Sodium chloride toxicity and the cellular basis
634 of salt tolerance in halophytes. *Ann. Bot.* 115, 419–431.
- 635 Flowers, T.J., Troke, P.F., Yeo, A.R., 1977. The mechanism of salt tolerance in halophytes.
636 *Annu. Rev. Plant Physiol.* 28, 89–121.
- 637 Foo, L.Y., Newman, R., Waghorn, G., McNabb, W.C., Ulyatt, M.J., 1996. Proanthocyanidins
638 from *Lotus corniculatus*. *Phytochemistry* 41, 617–624.
- 639 Geigenberger, P., 2003. Response of plant metabolism to too little oxygen. *Curr. Opin. Plant*
640 *Biol.* 6, 247–256.
- 641 Ghassemi, F., Jakeman, A.J., Nix, H.A., 1995. Salinisation of land and water resources: human
642 causes, extent, management and case studies. CAB international.
- 643 Gibbs, J., Greenway, H., 2003. Mechanisms of anoxia tolerance in plants. I. Growth, survival
644 and anaerobic catabolism. *Funct. Plant Biol.* 30, 1–47.
- 645 Gupta, K.J., Zabalza, A., Van Dongen, J.T., 2009. Regulation of respiration when the oxygen
646 availability changes. *Physiol. Plant.* 137, 383–391.
- 647 Haddadi, B.S., Hassanpour, H., Niknam, V., 2016. Effect of salinity and waterlogging on
648 growth, anatomical and antioxidative responses in *Mentha aquatica* L. *Acta Physiol.*
649 *Plant.* 38, 119.
- 650 Hall, T.A., 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis
651 program for Windows 95/98/NT, in: *Nucleic Acids Symposium Series*. [London]:
652 Information Retrieval Ltd., c1979-c2000., pp. 95–98.
- 653 Henderson, S.W., Baumann, U., Blackmore, D.H., Walker, A.R., Walker, R.R., Gilliam, M.,
654 2014. Shoot chloride exclusion and salt tolerance in grapevine is associated with
655 differential ion transporter expression in roots. *BMC Plant Biol.* 14, 273.
- 656 Hirabayashi, Y., Mahendran, R., Koirala, S., Konoshima, L., Yamazaki, D., Watanabe, S., Kim,
657 H., Kanae, S., 2013. Global flood risk under climate change. *Nat. Clim. Chang.* 3, 816–
658 821.
- 659 Hoagland, D.R., Arnon, D.I., 1950. The water-culture method for growing plants without soil.
660 *Circ. Calif. Agric. Exp. Stn.* 347.
- 661 Hu, Y., Schmidhalter, U., 2005. Drought and salinity: a comparison of their effects on mineral
662 nutrition of plants. *J. Plant Nutr. Soil Sci.* 168, 541–549.
- 663 Iwasaki, I., Utsumi, S., Hagino, K., Ozawa, T., 1956. A new spectrophotometric method for
664 the determination of small amounts of chloride using the mercuric thiocyanate

method. Bull. Chem. Soc. Jpn. 29, 860–864.

James, R.A., Davenport, R.J., Munns, R., 2006. Physiological characterization of two genes for Na⁺ exclusion in durum wheat, Nax1 and Nax2. *Plant Physiol.* 142, 1537–1547.

Jossier, M., Kroniewicz, L., Dalmas, F., Le Thiec, D., Ephritikhine, G., Thomine, S., Barbier-Brygoo, H., Vavasseur, A., Filleur, S., Leonhardt, N., 2010. The Arabidopsis vacuolar anion transporter, AtCLCc, is involved in the regulation of stomatal movements and contributes to salt tolerance. *Plant J.* 64, 563–576.

Kotula, L., Clode, P.L., Striker, G.G., Pedersen, O., Läuchli, A., Shabala, S., Colmer, T.D., 2015. Oxygen deficiency and salinity affect cell-specific ion concentrations in adventitious roots of barley (*Hordeum vulgare*). *New Phytol.* 208, 1114–1125.

Kronzucker, H.J., Coskun, D., Schulze, L.M., Wong, J.R., Britto, D.T., 2013. Sodium as nutrient and toxicant. *Plant Soil* 369, 1–23.

Lavado, R.S., Taboada, M.A., 1987. Soil salinization as an effect of grazing in a native grassland soil in the Flooding Pampa of Argentina. *Soil Use Manag.* 3, 143–148.

Leigh, R.A., Wyn Jones, R.G., 1986. Cellular compartmentation in plant nutrition: the selective cytoplasm and the promiscuous vacuole. *Adv. plant Nutr.*

Liu, P., Yang, G.-D., Li, H., Zheng, C.-C., Wu, C.-A., 2010. Overexpression of NHX1s in transgenic Arabidopsis enhances photoprotection capacity in high salinity and drought conditions. *Acta Physiol. Plant.* 32, 81–90.

Loreto, F., Centritto, M., Chartzoulakis, K., 2003. Photosynthetic limitations in olive cultivars with different sensitivity to salt stress. *Plant. Cell Environ.* 26, 595–601.

Lu, C., Qiu, N., Lu, Q., Wang, B., Kuang, T., 2002. Does salt stress lead to increased susceptibility of photosystem II to photoinhibition and changes in photosynthetic pigment composition in halophyte *Suaeda salsa* grown outdoors? *Plant Sci.* 163, 1063–1068.

Luo, Q., Yu, B., Liu, Y., 2005. Differential sensitivity to chloride and sodium ions in seedlings of *Glycine max* and *G. soja* under NaCl stress. *J. Plant Physiol.* 162, 1003–1012.

Maathuis, F.J.M., Amtmann, A., 1999. K⁺ nutrition and Na⁺ toxicity: the basis of cellular K⁺/Na⁺ ratios. *Ann. Bot.* 84, 123–133.

Martin, J., Fackler, P.L., Nichols, J.D., Lubow, B.C., Eaton, M.J., Runge, M.C., Stith, B.M., Langtim, C.A., 2011. Structured decision making as a proactive approach to dealing with sea level rise in Florida. *Clim. Change* 107, 185–202.

Martínez-Atienza, J., Jiang, X., Garcíadeblas, B., Mendoza, I., Zhu, J.-K., Pardo, J.M., Quintero, F.J., 2007. Conservation of the salt overly sensitive pathway in rice. *Plant Physiol.* 143, 1001–1012.

McDonald, M.P., Galwey, N.W., Colmer, T.D., 2002. Similarity and diversity in adventitious root anatomy as related to root aeration among a range of wetland and dryland grass species. *Plant. Cell Environ.* 25, 441–451.

McNabb, W.C., Waghorn, G.C., Peters, J.S., Barry, T.N., 1996. The effect of condensed

704 tannins in *Lotus pedunculatus* on the solubilization and degradation of ribulose-1, 5-
705 bisphosphate carboxylase (EC 4.1. 1.39; Rubisco) protein in the rumen and the sites of
706 Rubisco digestion. *Br. J. Nutr.* 76, 535–549.

707 Meloni, D.A., Oliva, M.A., Martinez, C.A., Cambraia, J., 2003. Photosynthesis and activity of
708 superoxide dismutase, peroxidase and glutathione reductase in cotton under salt
709 stress. *Environ. Exp. Bot.* 49, 69–76.

710 Mendoza, R., Escudero, V., García, I., 2005. Plant growth, nutrient acquisition and
711 mycorrhizal symbioses of a waterlogging tolerant legume (*Lotus glaber* Mill.) in a
712 saline-sodic soil. *Plant Soil* 275, 305–315.

713 Min, B.R., Barry, T.N., Attwood, G.T., McNabb, W.C., 2003. The effect of condensed tannins
714 on the nutrition and health of ruminants fed fresh temperate forages: a review. *Anim.*
715 *Feed Sci. Technol.* 106, 3–19.

716 Munns, R., 2002. Comparative physiology of salt and water stress. *Plant. Cell Environ.* 25,
717 239–250.

718 Munns, R., Tester, M., 2008. Mechanisms of salinity tolerance. *Annu. Rev. Plant Biol.* 59,
719 651–681.

720 Nakamura, A., Fukuda, A., Sakai, S., Tanaka, Y., 2006. Molecular cloning, functional
721 expression and subcellular localization of two putative vacuolar voltage-gated chloride
722 channels in rice (*Oryza sativa* L.). *Plant Cell Physiol.* 47, 32–42.

723 Nunez, S., Arets, E., Alkemade, R., Verwer, C., Leemans, R., 2019. Assessing the impacts of
724 climate change on biodiversity: is below 2° C enough? *Clim. Change* 154, 351–365.

725 Paolocci, F., Robbins, M.P., Madeo, L., Arcioni, S., Martens, S., Damiani, F., 2007. Ectopic
726 expression of a basic helix-loop-helix gene transactivates parallel pathways of
727 proanthocyanidin biosynthesis. structure, expression analysis, and genetic control of
728 leucoanthocyanidin 4-reductase and anthocyanidin reductase genes in *Lotus cornicu.*
729 *Plant Physiol.* 143, 504–516.

730 Paolocci, F., Rubini, A., Riccioni, C., Arcioni, S., 2006. Reevaluation of the life cycle of *Tuber*
731 *magnatum*. *Appl. Environ. Microbiol.* 72, 2390–2393.

732 Pardo, J.M., Cubero, B., Leidi, E.O., Quintero, F.J., 2006. Alkali cation exchangers: roles in
733 cellular homeostasis and stress tolerance. *J. Exp. Bot.* 57, 1181–1199.

734 Perelman, S.B., León, R.J.C., Oesterheld, M., 2001. Cross-scale vegetation patterns of
735 Flooding Pampa grasslands. *J. Ecol.* 89, 562–577.

736 Pfaffl, M.W., Horgan, G.W., Dempfle, L., 2002. Relative expression software tool (REST©) for
737 group-wise comparison and statistical analysis of relative expression results in real-time
738 PCR. *Nucleic Acids Res.* 30, e36–e36.

739 Ponnamperuma, F.N., 1984. Effects of flooding on soils. *Flooding and plant growth* 9–45.

740 Proehl, E.C., Nelson, W.P., 1950. The flame photometer in determination of sodium and
741 potassium. *Am. J. Clin. Pathol.* 20, 806–813.

742 Rejili, M., Vadel, A.M., Guetet, A., Neffatti, M., 2007. Effect of NaCl on the growth and the

743 ionic balance K^+/Na^+ of two populations of *Lotus creticus* (L.)(Papilionaceae). South
744 African J. Bot. 73, 623–631.

745 Rogers, M.E., Noble, C.L., Halloran, G.M., Nicolas, M.E., 1997. Selecting for salt tolerance in
746 white clover (*Trifolium repens*): chloride ion exclusion and its heritability. New Phytol.
747 135, 645–654.

748 Rubio, G., Casasola, G., Lavado, R.S., 1995. Adaptations and biomass production of two
749 grasses in response to waterlogging and soil nutrient enrichment. Oecologia 102, 102–
750 105.

751 Sanchez, D.H., Szymanski, J., Erban, A., Udvardi, M.K., Kopka, J., 2010. Mining for robust
752 transcriptional and metabolic responses to long-term salt stress: a case study on the
753 model legume *Lotus japonicus*. Plant. Cell Environ. 33, 468–480.

754 Shabala, S., Cuin, T.A., 2008. Potassium transport and plant salt tolerance. Physiol. Plant.
755 133, 651–669.

756 Shabala, S., Pottosin, I., 2014. Regulation of potassium transport in plants under hostile
757 conditions: implications for abiotic and biotic stress tolerance. Physiol. Plant. 151, 257–
758 279.

759 Sibole, J. V, Cabot, C., Poschenrieder, C., Barceló, J., 2003. Ion allocation in two different
760 salt-tolerant Mediterranean *Medicago* species. J. Plant Physiol. 160, 1361–1365.

761 Soriano, A., León, R.J.C., Sala, O.E., Lavado, R.S., Deregibus, V.A., Cauhepe, M.A., Scaglia,
762 O.A., Velázquez, C.A., Lemcoff, J.H., 1991. Río de la Plata Grasslands. In 'Ecosystems of
763 the world 8A. Natural grasslands. Introduction and Western Hemisphere'.(Ed. RT
764 Coupland) pp. 367–407.

765 Strasser, R.J., Srivastava, A., 1995. Polyphasic chlorophyll a fluorescence transient in plants
766 and cyanobacteria*. Photochem. Photobiol. 61, 32–42.

767 Striker, G.G., Colmer, T.D., 2017. Flooding tolerance of forage legumes. J. Exp. Bot. 68,
768 1851–1872.

769 Sun, Y., Wang, D., Bai, Y., Wang, N., Wang, Y., 2006. Studies on the overexpression of the
770 soybean *GmNHX1* in *Lotus corniculatus*: the reduced Na^+ level is the basis of the
771 increased salt tolerance. Chinese Sci. Bull. 51, 1306–1315.

772 Teakle, N.L., Amtmann, A., Real, D., Colmer, T.D., 2010. *Lotus tenuis* tolerates combined
773 salinity and waterlogging: maintaining O_2 transport to roots and expression of an
774 *NHX1*-like gene contribute to regulation of Na^+ transport. Physiol. Plant. 139, 358–374.

775 Teakle, N.L., Flowers, T.J., Real, D., Colmer, T.D., 2007. *Lotus tenuis* tolerates the interactive
776 effects of salinity and waterlogging by 'excluding' Na^+ and Cl^- from the xylem. J. Exp.
777 Bot. 58, 2169–2180.

778 Teakle, N.L., Real, D., Colmer, T.D., 2006. Growth and ion relations in response to combined
779 salinity and waterlogging in the perennial forage legumes *Lotus corniculatus* and *Lotus*
780 *tenuis*. Plant Soil 289, 369.

781 Teakle, N.L., Tyerman, S.D., 2010. Mechanisms of Cl^- -transport contributing to salt tolerance.

- Plant. Cell Environ. 33, 566–589.
- Thornton, P.K., Ericksen, P.J., Herrero, M., Challinor, A.J., 2014. Climate variability and vulnerability to climate change: a review. Glob. Chang. Biol. 20, 3313–3328.
- Wiengweera, A., Greenway, H., Thomson, C.J., 1997. The use of agar nutrient solution to simulate lack of convection in waterlogged soils. Ann. Bot. 80, 115–123.
- Xu, G., Magen, H., Tarchitzky, J., Kafkafi, U., 1999. Advances in chloride nutrition of plants, in: Advances in Agronomy. Elsevier, pp. 97–150.
- Xue, Z.-Y., Zhi, D.-Y., Xue, G.-P., Zhang, H., Zhao, Y.-X., Xia, G.-M., 2004. Enhanced salt tolerance of transgenic wheat (*Triticum aestivum* L.) expressing a vacuolar Na⁺/H⁺ antiporter gene with improved grain yields in saline soils in the field and a reduced level of leaf Na⁺. Plant Sci. 167, 849–859.
- Zabalza, A., Van Dongen, J.T., Froehlich, A., Oliver, S.N., Faix, B., Gupta, K.J., Schmäzlin, E., Igal, M., Orcaay, L., Royuela, M., 2009. Regulation of respiration and fermentation to control the plant internal oxygen concentration. Plant Physiol. 149, 1087–1098.

Figure legends

Figure 1. Phenotypic characteristics of *L. corniculatus* accessions subjected to the different stress treatments. Plants were subjected to the different treatments for 21 days, after the salinity acclimation period imposed for those treatments involving salt stress. Pictures were taken after 8 and 21 d (harvest). For the control and waterlogging (WL) treatments, plants were irrigated periodically with Hoagland solution 0.5 x with or without free drainage, respectively. In the salt treatment (NaCl) and combined stress treatment (WL+NaCl), plants were irrigated with Hoagland solution 0.5 x supplemented with 150 mM of NaCl with or without drainage, respectively. When waterlogging stress was imposed, nutritive solution was previously bubbled with N₂ (g) to reduce O₂ to hypoxic levels. LcD, diploid *L. corniculatus* accession; LcT, tetraploid *L. corniculatus* accession.

Figure 2. Dry mass accumulation of *L. corniculatus* accessions subjected to the different stress treatments. The measured shoots (A) and roots (C) dry mass are shown as percentage of the control dry mass in each case. (B) Total dry mass (g) accumulation at the end of the experiment. (D) Shoot:Root ratio was calculated from the dry mass accumulation of shoots and roots, respectively. Means ($n = 5 \pm SD$) without common letters differ significantly within each of the accessions (one-way ANOVA; Duncan, $p < 0.05$). White bars, control conditions (Ctrl); light gray bars, waterlogging stress (WL); dark gray bars, saline stress (NaCl); black bars, combined stress (WL+NaCl).

Figure 3. Photosynthetic parameters of *L. corniculatus* accessions under the different stress treatments. Net photosynthetic rate at saturating irradiance (Asat) (A) and the

maximum quantum yield of PSII (Fv/Fm) (B) were measured for five independent biological replicates ($n = 5 \pm \text{SD}$). Means without common letters differ significantly within each of the accessions (one-way ANOVA; Duncan, $p < 0.05$) (A). Asterisks show significant differences of a stress treatment against the control treatment (Student t-test, * $p < 0.05$; ** $p < 0.01$) (B). White bars, control conditions (Ctrl); light gray bars, waterlogging stress (WL); dark gray bars, saline stress (NaCl); black bars, combined stress (WL+NaCl).

Figure 4. Ions concentrations in apical and basal leaves, and in roots. The concentration of Cl^- , Na^+ and K^+ ($\mu\text{mol.g}^{-1}.\text{dry mass}^{-1}$) were measured in apical leaves (A, D and G), basal leaves (B, E and H) and in roots (C, F and I) of both *L. corniculatus* accessions under the different treatments ($n = 5 \pm \text{SD}$). The K^+/Na^+ ratio was calculated for apical and basal leaves (J and K, respectively), and for roots (L), in each case. Means without common letters differ significantly within each of the accessions (one-way ANOVA; Duncan, $p < 0.05$). White bars, control conditions (Ctrl); light gray bars, waterlogging stress (WL); dark gray bars, saline stress (NaCl); black bars, combined stress (WL+NaCl).

Figure 5. Relative expression of *CLC* (A) and *NHX1* (B) genes. The values represent the mean $\pm \text{SD}$ ($n = 3$). Statistical analysis of relative expression was performed by comparing the relative expression of the genes based on the pairwise fixed reallocation randomization test ($p < 0.05$). Asterisk show significant differences ($p < 0.05$). White bars, control treatment (Ctrl); Black bars, combined stress (WL+NaCl).

Supplementary material

Supplementary Figure 1. Shoots and roots dry mass accumulation of *L. corniculatus* accessions subjected to the different stress treatments. The measured shoots (A) and roots (B) dry mass is shown in grams ($n = 5 \pm \text{SE}$). Means without common letters differ significantly within each of the accessions (one-way ANOVA; Duncan, $p < 0.05$). White bars, control conditions (Ctrl); light gray bars waterlogging stress (WL); dark gray bars, saline stress (NaCl); black bars, combined stress (WL+NaCl).

Supplementary Figure 2. Stomatal conductance of the *L. corniculatus* accessions under the different stress treatments. The stomatal conductance was measured for five independent biological replicates ($n = 5 \pm \text{SD}$). Means without common letters differ significantly within each of the accessions (one-way ANOVA; Duncan, $p < 0.05$). White bars, control conditions (Ctrl); light gray bars waterlogging stress (WL); dark gray bars, saline stress (NaCl); black bars, combined stress (WL+NaCl).

858 **Supplementary Table 1.** Primers used for qRT-PCR.

Primer name	Sequence (5' -> 3')	Reference
<i>NHX1</i> -Fw	TACTTCACTGCGGTCCAATG	Teakle et al., 2010
<i>NHX1</i> -Rv	GATCTAGGGAAGCCATGCTG	Teakle et al., 2010
<i>CLC</i> -Fw	TTAGTTGGAATGGCCGCTAC	Own design
<i>CLC</i> -Rv	ACTGAGGGAACCCATATTGC	Own design
<i>EF-1α</i> -Fw	TGACAAGCGTGTGATCGAGAGG	Escaray et al., 2014
<i>EF-1α</i> -Rv	GATACCTCTTTCACGCTCAGCCTT	Escaray et al., 2014

859

860

Figure 1. Phenotypic characteristics of *L. corniculatus* accessions subjected to the different stress treatments. Plants were subjected to the different treatments for 21 days, after the salinity acclimation period imposed for those treatments involving salt stress. Pictures were taken after 8 and 21 d (harvest). For the control and waterlogging (WL) treatments, plants were irrigated periodically with Hoagland solution 0.5 x with or without free drainage, respectively. In the salt treatment (NaCl) and combined stress treatment (WL+NaCl), plants were irrigated with Hoagland solution 0.5 x supplemented with 150 mM of NaCl with or without drainage, respectively. When waterlogging stress was imposed, nutritive solution was previously bubbled with N₂ (g) to reduce O₂ to hypoxic levels. LcD, diploid *L. corniculatus* accession; LcT, tetraploid *L. corniculatus* accession.

Figure 2. Dry mass accumulation of *L. corniculatus* accessions subjected to the different stress treatments. The measured shoots (A) and roots (C) dry mass are shown as percentage of the control dry mass in each case. (B) Total dry mass (g) accumulation at the end of the experiment. (D) Shoot:Root ratio was calculated from the dry mass accumulation of shoots and roots, respectively. Means ($n = 5 \pm \text{SD}$) without common letters differ significantly within each of the accessions (one-way ANOVA; Duncan, $p < 0.05$). White bars, control conditions (Ctrl); light gray bars, waterlogging stress (WL); dark gray bars, saline stress (NaCl); black bars, combined stress (WL+NaCl).

Figure 3. Photosynthetic parameters of *L. corniculatus* accessions under the different stress treatments. Net photosynthetic rate at saturating irradiance (Asat) (A) and the maximum quantum yield of PSII (Fv/Fm) (B) were measured for five independent biological replicates ($n = 5 \pm \text{SD}$). Means without common letters differ significantly within each of the accessions (one-way ANOVA; Duncan, $p < 0.05$) (A). Asterisks show significant differences of a stress treatment against the control treatment (Student t-test, * $p < 0.05$; ** $p < 0.01$) (B). White bars, control conditions (Ctrl); light gray bars, waterlogging stress (WL); dark gray bars, saline stress (NaCl); black bars, combined stress (WL+NaCl).

Figure 4. Ions concentrations in apical and basal leaves, and in roots. The concentration of Cl^- , Na^+ and K^+ ($\mu\text{mol}\cdot\text{g}^{-1}\cdot\text{dry mass}^{-1}$) were measured in apical leaves (A, D and G), basal leaves (B, E and H) and in roots (C, F and I) of both *L. corniculatus* accessions under the different treatments ($n = 5 \pm \text{SD}$). The K^+/Na^+ ratio was calculated for apical and basal leaves (J and K, respectively), and for roots (L), in each case. Means without common letters differ significantly within each of the accessions (one-way ANOVA; Duncan, $p < 0.05$). White bars, control conditions (Ctrl); light gray bars, waterlogging stress (WL); dark gray bars, saline stress (NaCl); black bars, combined stress (WL+NaCl).

Figure 5. Relative expression of *CLC* (A) and *NHX1* (B) genes. The values represent the mean $\pm$ SD (n = 3). Statistical analysis of relative expression was performed by comparing the relative expression of the genes based on the pairwise fixed reallocation randomization test ($p < 0.05$). Asterisk show significant differences ($p < 0.05$). White bars, control treatment (Ctrl); Black bars, combined stress (WL+NaCl).

Supplementary Figures

Supplementary Figure 1. Shoots and roots dry mass accumulation of *L. corniculatus* accessions subjected to the different stress treatments. The measured shoots (A) and roots (B) dry mass is shown in grams ($n = 5 \pm SE$). Means without common letters differ significantly within each of the accessions (one-way ANOVA; Duncan, $p < 0.05$). White bars, control conditions (Ctrl); light gray bars waterlogging stress (WL); dark gray bars, saline stress (NaCl); black bars, combined stress (WL+NaCl).

Supplementary Figure 2. Stomatal conductance of the *L. corniculatus* accessions under the different stress treatments. The stomatal conductance was measured for five independent biological replicates ($n = 5 \pm \text{SD}$). Means without common letters differ significantly within each of the accessions (one-way ANOVA; Duncan, $p < 0.05$). White bars, control conditions (Ctrl); light gray bars waterlogging stress (WL); dark gray bars, saline stress (NaCl); black bars, combined stress (WL+NaCl).