Traveling cortical networks compose a mindstream

Ernst Rudolf M. Hülsmann

26 Rue de Bonn, CH-3186 Guin, Switzerland.
ernsthuelsmann@gmail.com

ABSTRACT

The brain creates a physical response out of signals in a cascade of streaming transformations. These transformations occur over networks, which have been described in anatomical, cyto-, myeloarchitectonic and functional research. The totality of these networks has been modelled and synthesised in phases across a continuous time-space-function axis, through ascending and descending hierarchical levels of association¹⁻³ via changing coalitions of traveling networks⁴⁻⁶, where localised disorders might spread locally throughout the neighbouring tissues.

This study quantified the model empirically with time-resolving functional magnetic resonance imaging of an imperative, visually-triggered, self-delayed, therefor double-event related response task. The resulting time series unfold in the range of slow cortical potentials the spatio-temporal integrity of a cortical pathway from the source of perception to the mouth of reaction in and out of known functional, anatomical and cytoarchitectonic networks. These pathways are consolidated in phase images described by a small vector matrix, which leads to massive simplification of cortical field theory and even to simple technical applications.

INTRODUCTION

On a first sight it seems to be self-evident that a local perturbation within the cortical sheet should spread locally. The underpinning networks (see methods table II), dictating pathways, have been developed using different technologies and have varying numbers of nodes: anatomical³⁰⁻³⁸ cards are underpinned by fasciculi and smaller subcortical connectomes⁴⁸⁻⁵² and structured by cytoarchitectonic^{39-47,94-96,98-100,104,109,111,121-124}. These biological networks carry functions which, originally descriptively^{54,55} defined in psychological terms, lead, with integrated systems perspective^{56,57}, to the low-dimensional multifocal cards from dynamic functional magnetic resonance imaging (dfMRI) seen in resting states^{58,61,62,64-89}.

The networks are subject to a vectored dynamic system: in its multiple primary areas influxes of information flow inside and intermingle before running towards the only outlet in the primary motor area. This produces well defined temporal, spatial and functional boundary conditions. The system, even taking into account systemic feedforward-feedback-loops, yields an orientated direction and thus, using Hebbian¹ principles as a basis, Mesulam^{2,3,4} develops a model (figure I) where a linear thought stream is represented via four levels of association. A hierarchical vector is created across the four levels, integrating the above-mentioned boundary conditions. This four-levelled phase space illustrates not only effective but also causal and structural connectivity by using orthogonal, inter- and intra-associative projections:

A cycling phase might be started with the perception of a stimulus in a primary, idiosyncratic, unimodal association area, as e.g. within the visual cortex (ai); discrimination occurs further upstream in a secondary, still modality-specific association area (aai); heteromodal and often conscious integration follows in a tertiary, high-order association area (aiii); and as highest association area, a limbic, emotional response will upspring (I). Further ahead in time, the chain of ai-aai-aiii in reverse, falls down in descending order via a tertiary multimodal level of association, as in prefrontal thought processes (aiii); the anticipatory planning of movements, as in the vast secondary unimodal, supplementary-motor area (aai); to the execution of the movement in the primary level of association (ai), the motor cortex – a process of execution which ultimately triggers the ai-aai-aiii chain once again: this time through self-perception which takes place in the primary somatosensory area (ai) over unimodal discrimination in secondary parietal regions (aai) to multimodal parietal regions (aiii), resulting in temporal models as shown in figure I and more detailed in the methods figures IV – VII and movie I.

Independent of complex, dispersed processes, the order of the sequence ai-aai-aiii-I-aiii-aai-ai should be retained like a palindrome as a quasi-periodic pattern and be continually repeated in parallel and with varying phase lengths, just as the perception-reaction sequence involved in systematic thought processes also remains chronological and recurs rapidly throughout the day and gradually over the course of a lifetime in an ever-repeating self-similar loop.

Figure I. Hierarchies, longlasting waves and sensorimotor phases.

Above - The phasic hierarchy model of a visuo-motor task, adapted from Mesulam², figure 2, stretched over time.

The nodes named (within circles) following Mesulam (on the left), Jubrain^{45-47,63} respectively (on the right), 2π from visuo-perception to auto-perception.

AIT - anterior inferotemporal cortex; f - area specialized for face encoding; L - hippocampal-entorhinal or amygdaloid components of the limbic system; LIP - lateral intraparietal cortex; MST - medial superior temporal cortex; P - heteromodal posterior parietal cortex; Pf - lateral prefrontal cortex; s - area specialized for encoding spatial location; PIT - posterior inferotemporal cortex; T - heteromodal lateral temporal cortex; TF - part of medial inferotemporal cortex; V1 - primary visual cortex; V2, V3, V4, V5 - additional visual areas; W - Wernicke's area; wr - area specialized for encoding word-forms; 7a - part of dorsal parieto-occipital cortex.

Red lines – ventral streams; green lines - dorsal streams.

Details see methods, figures IV-VII and movie I.

Bottom - Event-related potentials ordered over time demonstrate like the rapid monosynaptic transmission-times in the range of 10 ms (from V1 to V5) to 50 ms (from V1 to AIT), as between Mesulam's hierarchical levels, add up to longlasting waves in the order from 2 Hertz down to 0.25 Hertz. Note that Hebbian monosynaptic feedforward-backward loops occur in parallel to the slow frequency oscillations in the range of some 10 ms.

SPN - Stimulus-preceding negativity, awaiting an informative stimulus; N100 – preattentive perceptual processing; N170 - faces; P200 – preattentive perceptual processing; N200 – stimulus detection; N250 – face identity; P300 – stimulus categorization and memory updating; N400 – semantic and conceptual processing; P600 – syntactic processing; SEP – Sensory evoked potential; ERN NE - Error negativity, false acts; Pe - Error positivity, false acts.

The model raises a series of prognoses regarding the where, what and when of a transformation whose content is ultimately arbitrary and of the likelihood of such a linear, successive combination occurring. Regulated by feedforward-feedback-loops, interferences appear over longer time and space periods as static or moving waves, described by Crick & Koch⁵ as

... a forward-traveling, ... propagating net-wave of neuronal activity, (but is not the same as a wave in a continuous medium) ... These net-waves represent at each point in time 'coalitions' ... There are coalitions, which achieve winning status and hence produce [a] conscious experience ... In the conscious mode, it seems likely that the flow is in both directions so that it resembles more of a standing net-wave.

Deco & al.⁶ ground this dynamic into a neural field model where the above described neural masses rise out of spiking neurons, concluding *in summary, for any model of neuronal dynamics, specified as a stochastic differential equation, there is a deterministic linear equation that can be integrated to generate ensemble dynamics.*

The corresponding empirical spatio-temporal matrix **M** and its physical node parameters (such as location, volume, point in time, phase, frequency, amplitude) and edge parameters (such as distance, direction, duration, hierarchy, functions) are determined, using time-resolved functional magnetic resonance imaging (fMRI), both separately and as arrays in tasking, i.e. double-event-related brains, where a visual instruction was followed by a considered motor-reaction (see methods).

RESULTS

The appearance of the resulting phase image (figure II, details supplementary figures VIII & IX, movie II) unfolds before the view a wave train, which reflects the flow of the visual stream, the dorsal and the ventral one, fulminant, and reproduces at first glance in great clarity the sensorimotor stream, ventral and dorsal stream well separated too.

A right-left hemispheric symmetry of the activation pattern allows to limit the detailed analysis to one hemisphere, which will be the right one in the following detailed descriptions.

There is a medio-dorsal cluster in the visual phase image, the medial supplementary motor area SMA (figure II, pink oval), the latest cluster in time to have been activated. The same cluster is the first to be activated in the motor pictures through the movement. The two clusters in the two different phase images overlap in the right hemisphere over 356 mm³ (table I) and are constructed out of the same microanatomical, functional and temporal elements. Activation begins in the two conditions in pre-SMA (visual 3.1 s post, motor 2.1 s pre) and flows over the SMA proper to the medial upper border of the cortical mantle (visual 3.9 s post, motor 1.6 s pre).

A joint time period is therefore consolidated in which a time arrow of 5.2 s represents the time from one perception over cognition to action and back to autoperception, the phase 2π in Mesulam's model out of figure I, with wave fronts covering expansive, anatomical, cyto- and myeloarchitectonical, functional and hierarchical areas, parallel and perpendicular, as shown in table I:

i) Anatomical network architecture is quantified (table I, column 3; both hemispheres supplementary table II) with Automated anatomical labeling (AAL)³² over 15 out of the 44 AAL-nodes (calcarine, temporal, fusiform, lingual, occipital inferior, parietal inferior, cuneus, parietal superior, supplementary motor area, middle cingulum, precentral, postcentral, Rolandic operculum and once more parietal inferior in each hemisphere). The AAL-algorithm gives activated volumes accumulating in total to almost 10000 mm³ in each hemisphere; with an outer layer thickness⁷ of 2.3 - 2.8 mm a neural coverage of about 250 cm² for a surface⁸ of 1200 cm². Identifying first, extreme and last point in time of correlation allows modelling of the regional *standing netwaves*, the frequencies moving with 0.3 – 0.5 Hz in the lower delta band. The entire set of *standing netwaves* results in a stream of continuously phase-shifted waves, where the individual regions develop in parallel as *winning coalitions* and form a *forward traveling netwave* across time and space.

ii) The cytoarchitectonic network (table I, columns 4 - 7) resolution augments the number of nodes up to 26 out of the 64 nodes shown in figure V. The ventral visual stream is here composed of 17⁹⁴, FG1⁹¹, ventral 18⁹⁴, hOc4v⁹⁵, hOc3v, FG2^{91,96}, BA 36, with a midtemporal branch hOc5⁹⁹, BA 21^{97,100}. The dorsal visual stream

Figure II. Phase images of a self-triggered response task.

Phase images from time series (supplementary figures VIII and IX, movie II) show the dynamic of an optically triggered, imperative, self-delayed response task.

Maximal-intensity-projections of the point-in-time of extreme activities as indicated by the colour bars. Left image event-related to the signal, and the two right pictures event-related to the respective reactions; stimulus-related from 1 second pre-stimulus (red) to 4 seconds post-stimulus (blue); reaction-related from 4 seconds prior to the reaction (blue) to 2 seconds after the reaction (red). The colour bar codes hierarchies^{2,3,4} too, separating unimodal (red/brown/green) from heteromodal (green/blue) areas.

Note the spatio-temporal gradient spanning from the calcarine sulcus over the medial, lateral and inferior extrastriate cortex to the precuneus and superior parietal cortex, and from the deep mid-frontal pre-SMA to the upper SMA proper (purple circle), where a spatio-temporal gradient starts in the motor images, followed within the lateral convexity by gradients from the central sulcus to anterior and to posterior areas, 7PC/mip short cutting to the latter visual superior parietal side (green circle).

is composed of 17, dorsal 18, hOc3d⁹⁸, ip1¹⁰⁹, mip¹⁰⁴, 7PC, 7A, and 7P^{105-108,124} and finally, a new observation, BA 6, pre-SMA^{63,110-116}.

The corresponding volitional motor response is made of a medial part from paleocortical aCMA¹¹³ over pre-SMA, SMA proper to 6aβ; and a lateral part from 4p¹²¹, a dorsal stream to prefrontal 6aα and to postcentral 3, to 1^{122,123} and further on to 7P¹²⁴, as well as a ventral stream to 6aα inferior⁶³ and to caudal PFt, Pfop¹⁰⁰.

iii) These macro- and microanatomical architectures port dfMRI-correlations (table I, column 2). Their temporal architecture is obvious in a sequence over seven networks with sixteen nodes, representing their hypothesized functional succession: At first, consecutive visual networks^{62,64,66,69,71,78-80} turn up - the occipital, lateral and medial visual network as per Smith & al.⁶⁴, with the latter two forming the high-level visual network, as per Zhang & al.⁷⁸, including the six hierarchical nodes of Yeo's & al.⁶² visual hierarchies.

Within the higher association areas it is the dorsal attention network^{62,70,72,75,78-81}, which follows parietal and ultimately the medial-precentral part of the sensory motor network^{64,65,78-81} originates with the latter.

This and the anterior cingulate cortex, a part of the ventral attention/salience network^{62,69,72,78,79}, is also the origin of the motor branch which subsequently employs the ventral motor network⁷⁸, and flows posteriorly and anteriorly back into the dorsal attention network, and inferiorly towards the Roland's motor operculum in the direction of the insular lobe, spreading in an almost elliptical fashion from the sulcus centralis.

M	dfMRI	AAL	BA	Jubrain	mm ³	s
aI, V1	vp	calcarinus	17	17	820	0.0
aII, V5 TF	vl	temporal mid.	21	hOc5	199	0.1
aII, V4 wr	vl	fusiformis	37	FG1	776	0.1
aII, V2	vl	lingualis	18	18v	856	0.3
aII	vm	occipitalis inf.	19	hOc4v	140	1.1
aIII, V5 T	vl	temporal mid.	21	21	125	1.2
aII, V3	vl	lingualis	19	hOc3v	468	1.5
aII, V4 wr	vl	fusiformis	37	FG2	272	1.8
I, TF		lingualis	36	36	60	2.1
M	dfMRI	AAL	BA	Jubrain	mm ³	s
aIII, LIP	dat	par. sup.	7	ip1	111	0.5
aII, V2	vm	cuneus	18	18d	200	1.5
aIII, LIP	dat	par. sup.	7	mip	148	2.0
aII, V2	vm	cuneus	19	hOc3d	170	2.4
aIII, LIP	dat	par. sup.	7	7PC	52	2.6
aIII, 7a	dat	par. sup.	7	7A	200	2.8
aIII, P	dat	precuneus	7	7P	686	2.8
M	dfMRI	AAL	BA	Jubrain	mm ³	s
I	vat	cingulum med.	32,24	aCMA	220	-2.2
aII	sm	SMA	6	pre-SMA	305	-2.1
aII	sm	SMA	6	SMA proper	195	-1.6
aI	sm	precentral	4	4p	1223	-1.6
aI	sm	postcentral	3	3a, 3b	444	-0.5
aII	sm	postcentral	1	eins	1190	-0.2
aII	sm	postcentral	5	5	300	0.4
aII	sm	SMA	6	6aβ	45	1.3
aII	sm	precentral	6	6aα sup	366	1.6
aII, LIP	dat	postcentral	7	7PC	207	1.7
aII	vm	rolandic op.	43	Pfop/6aα inf	253	1.8
aII	vm	par. inf.	40	PFt	202	1.9

ventral visual stream; signal t = 0

dorsal visual stream; signal t = 0

motor stream reaction t = 0

Table I. Hierarchical, functional, anatomical and histological visuo-motor nodes of the right hemisphere forming a mindstream.

Columns: M - Mesulam^{2,3} (abbreviations as in figure I); dfMRI - dynamic functional resonance imaging (networks: vp – visual primary; vl – visual lateral; vm – visual medial; dat – dorsal attention; vat – ventral attention; sm – sensorimotor; vm – ventral motor); AAL, automated anatomical labeling⁹; BA – Brodmann^{40,41} areas; Jubrain⁴⁵⁻⁴⁷, but the differentiation within the large surface region 18 between 18v (18 ventral, gyrus lingualis) and 18d (18 dorsal, cuneus and gyrus occipitalis); also the as yet in JuBrain uninvestigated Brodmann areas 36 and 21; also aCMA, a small portion of BA 32 and the differentiation of the large surface volume of SMA according to myelographic classification⁶³; mm³ – activated volume; s - the average points in time of maximal correlations for a given node in seconds related to the event.

iv) The hierarchical architecture (table I, column 1), including dynamic ideas, is of particular importance and shows the phenomenon of a linear *traveling netwave* over the visual perception, discrimination and higher association levels as delineated by Mesulam^{2,3}. This demonstrates the starting point in early idiosyncratic, primary areas and its evolution over time via secondary unimodal to multimodal high association areas, located midparietal. These midparietal high-association areas are omitted in a speeded response task, demonstrating their high hierarchical position (supplementary figure X).

The reversed hierarchy is present in the movement-linked development from limbic regions over secondary to primary unimodal motor, to primary, secondary unimodal somatosensory and tertiary heteromodal sensory areas. Over the course of time the process recruits all the four Mesulam hierarchies over nine nodes.

The complex concordance from i) - iv) is grouped into large clusters, which radically simplifies the description to the determination of local and temporal extreme values (figure III). Vectors (black in figure III) are constructed between these extremes over the extension of those clusters, which in their orientation cover the above described, ordered sets of anatomies, cytoarchitectures, functions and hierarchies. Putting it together leads to a vector matrix bridging the combined visual occipito-temporo-parietal-midprefrontal, visual occipito-lingual-fusiform and sensorimotor prefrontal-precentral-postcentral-parietal networks. The matrix spans a field from the source of the visual perception through the conscious reflexion to the action output into the newly emerging autoperception. By way of the constructed vectors, a wave develops like Hayne's *stream of consciousness*⁹.

Thus, the wanted matrix M is well approximated.

DISCUSSION

Waves are built from their amplitude and their duration in space and time, oscillate in different frequencies, appear time- and space-locked to events, can spread locally *and are believed to be fundamental mechanisms of neuronal computation and interregional communication*²⁷.

Neurons are the elemental building blocks of such cortical waves, spiking predominantly in the form of propagating, traveling waves, while at the meso- and macroscopic levels, mainly stationary waves have been described so far. Such stationary signals have been explored regarding the significance of their physical properties, as especially frequency. Slow cortical potentials, < 4 Hz, so in the time range of the here observed signals, have been seen by He & Raichle¹⁰ as optimally positioned for carrying out large-scale information integration in the brain, have even been linked by them to conscious processing, and have been inverted by neurofeedback¹¹.

In fMRI the temporal properties of oscillations could be used to describe the standing, anticorrelated bumps on unimodal networks superficially and on multimodal networks very profoundly. Time-locked event-related standing oscillations are fundamental in electrophysiology (see examples of event-related potentials in figure I) and functional MRI, and even more information seems to be linked to phase-coded signals. Montemurro & al.¹² calculate in macaques *that at low field potentials frequencies, the phase of firing conveyed 54% additional Shannon information beyond that conveyed by spike counts*. Glerean & al.⁵⁸ demonstrate by using instantaneous phase synchronization as a measure of dynamic, time-varying functional connectivity in the visual cortex of man the same regions to be activated as in the present study but the mid-precentral SMA.

Lakatos & al.¹³ show even – again in the same wave-length as in the present study - that *stimulus-locked phase-coding in the delta frequency range are present when attended stimuli are in a rhythmic stream. These ... oscillations ... entrain to the rhythm of the stream, resulting in increased response gain for task-relevant events and decreased reaction times*.

However, traveling netwaves have only since recently, with the dawn of methods as multielectrode array, electrocorticography, voltage-sensitive dye, low field potential and intracellular recordings, been in the focus of interest. Traveling netwaves over some hundred's milliseconds, observed in the cat's visual cortex, voltage-sensitive dye registration in V1, V2 and V4 of the awake monkey, traveling beta oscillations during movement planning in

Figure III. Dynamic topology of a visuo-motor vector field.

The right hemispheres from figure II combined to an entire stream, surface-rendered and the histological architecture as per the definitions of JuBrain⁴⁵⁻⁴⁷, with the exception of BA 21 and 36, which are not yet published. The subclassification of SMA/BA 6 as per Zilles & al.⁶³. 18m – medial (cuneus), 18l – lateral part (gyrus occipitalis) of dorsal JuBrain-area 18.

The colour bars provide the time scale in reference to visual perception and movement in seconds (supplementary movie III). Temporal extreme values were determined within the regions and vectors extrapolated (black arrows).

The table to the left provides the temporal extreme values (t in seconds, the time-arrow of the reaction transposed by -5.2 seconds to visual perception) and their point in space (x, y, z – Talairach²⁸-coordinates in mm).

The table to the right provides the vectors with their Euclidian distances in time (dt, in seconds) and space (ds, in mm).

macaques, and especially the electrocorticographic slow oscillations, 0.1 - 1.0 Hz, orchestrating large-scale neural spiking in human slow-wave sleep, traveling anterior-to-posterior with 1.2 - 7.0 m/s, lead Muller & al.¹⁴ to the conclusion *that stimulus-evoked travelling waves are a general and important feature of dynamics in the visual system ... so ... that dynamic wave fields introduce a general conceptual framework.*

All these individual biophysical properties - on standing waves, on frequencies, on rhythmic oscillations, on timelocking, on phase synchronization, on wandering waves – as observed in those studies are continuously and in overwhelming totality and simplicity present in the vector-matrix constructed in fig. III. The propagation speeds are from 0.1 to 0.25 m/s, the frequencies of 0.5 Hz, traveling across several centimeters, and adding up to a complete stream of about 50 cm length and 10 s duration. These waves develop parallel and dense in temporo-spatial continua, so that a current grows out of them. The phases of the current waves are temporally synchronized in a continuum along spatio-hierarchical-functional axes.

Methods to exploit the information encoded in time-varying connectivity have been developed in recent years in order to reach beyond the static measures of resting fMRI^{15,16}. Task positive paradigms are, in their nature, even more proximately linked to time-varying phenomena. While resting state might resemble testing an orchestra preparing for the concert, a task positive brain would be like observing a playing orchestra, where time synchronization will occur as the result of supervised Hebbian learning. Finally, it is the task-positive space where we look for orbits and attractors¹⁷. However, the solution space shown here can now be used as seed-point series for dfMRI.

In the course of these considerations, the phases cluster were understood as Hebbian layers, which adhere to each other spatially. The synchronized stream that was observed here would, however, not represent the Hebb's error minimization but it's result. Thus high synchronization, sharing information between entities of a system, might be, far away from concrete, local functions as the perception of colour, intention or emotion, the essential Hebbian supervisor.

CONCLUSION

Slow cortical potentials are ordered to a spatio-temporal matrix $\mathbf{M}$ out of a series of vectors, forming together a *stream of consciousness*⁹.

Well evaluated neuro-biological networks are reflected in $\mathbf{M}$ in a linear formation: The temporal order is reproduced i) between the dynamic functional networks visual, dorsal and ventral attention, and sensorimotor; ii) between the anatomical regions contained within these networks; iii) between the Brodmann and Zilles areas also found in these regions; iv) inside these areas and v) via functions and hierarchies.

Geometric waves are observed 'hovering' in the event of high over time synchronisation^{18,19} between input and output, in a similar fashion to a taut bow string. Crick & Koch's⁵ concept of consciousness as *a forward-traveling, propagating net-wave of neuronal activity* holds true and its quantification as phase-shifted synchronisation²⁰ includes the simplification of biophysical dynamic causal models^{21,22}, as well as cortical field theory⁶.

The Mesulam and more deeply stacked hierarchical models can be linearly parametrised in task-positive networks accordingly.

The discussed models of Hebb¹, Libet¹¹⁴, Kornhuber & Deecke¹¹⁵, Birbaumer¹⁰⁸, Crick & Koch⁵, Mesulam^{2,3}, He & Raichle¹⁰, Kjaer & al.¹⁰⁶ and Haynes⁹ are essentially about the formalization of consciousness. It seems on the first sight evident that a complex system like consciousness should be composed out of even few and simple elements like other natural phenomena are constructed from simple elements. These elements, vectors, compose $\mathbf{M}$, which can now be inverted²³ in a computational form leading, like Birbaumer & al.¹⁰⁸ did in neurofeedback with standing slow cortical potentials, to dynamic applications as multi-channel transcranial stimulation, inducing a virtual reality – spectra of cortical cycling phase continua forming a mindstream.

METHODS

The paradigm^{24,25,26} required a visual signal which prompted a test subject to push a button using either their left or right hand following a period they felt was “up to around 15 seconds”, with this process monitored using an fMRI scan:

Ten subjects (age range from 21 to 59 years, average age 30 years, 4 female, 6 male, all right-handers, as determined by the Edinburgh Inventory of Handedness) were studied. Informed written consents and approvals by the local ethic commission were obtained. The subjects held a button in each hand and watched visual stimuli presented from a video beamer onto a screen. The stimuli consisted of a short white flash (150 ms) on a black screen, followed by the appearance of a white arrow, which pointed in a pseudo-randomized sequence either to the right, or to the left. The arrow remained visible over the whole trial. The subjects were instructed to press the button with the corresponding thumb (left or right). They were furthermore instructed to choose, for each button press, varying temporal delays from zero up to approximately 15 seconds following the instruction flash. The whole experiment required 20 minutes, during which 87 trials of this delayed response task were presented with pseudo-randomized interstimulus-intervals of 14 +/- 2.5 seconds. The delayed task was contrasted with a speeded response task.

Event-related functional Magnetic Resonance Imaging was performed with a 1.5 Tesla MRI scanner (Siemens Vision, Erlangen, Germany). Twenty-four axial slices with 5 mm thickness and 3x3 mm² in-plane resolution, 64² matrix, 8/8 192 mm² FOV covering the whole brain were acquired using a gradient recalled single shot echo-planar-imaging sequence. To shorten examination time TE was set to 40 ms. The measurement time for one whole brain acquisition was 2.32 seconds and volume repetition time was 2.52 seconds. Thus, for each experiment, 480 whole brain measurements were performed during a time interval of 20 minutes resulting in a mean of 5.5 volume measurements for each trial.

The image evaluation was performed with the Statistical Parametric Mapping programme package (SPM99, Wellcome Department of Cognitive Neurology, London). Preprocessing of the single subject time series images consisted of i) slice-timing, referring to the first slice (top), ii) realignment with sinc interpolation and adjusting for sampling errors, iii) non-linear normalisation to the canonical echo planar target image in Montreal Neurological Imaging space, interpolated with 2 x 2 x 2 mm³ voxel size, in order to achieve maximal spatial signal-resolution in the group statistics, and iv) spatial smoothing with a Gaussian filter with full-width half-maximum of 6 mm.

To generate time-resolved statistical maps, a series of models was calculated with time-shifted event delays. For each delay, the BOLD response was modelled using a generic hrf (global scaling, specified high-frequency-filter, using the hrf as low-pass-filter, which corresponds to a Gaussian filter with a cut-off level of 4 seconds full-width half-maximum, and without correcting for further internal correlation), resulting in separate t-maps for time points prior to the signal and the self-paced movements (-2 s for the signal, -3.5 s for the movements) and following (6 s for the signal, 2.5 s for movements). T-values were scaled for each volume proportionately from threshold ($p < 0.001$) to individual volumes extreme t-value at point in time t_e . The resulting values were assigned to a baseline corrected²⁷ activity I. Spatial segmentation used the algorithm of Tzourio-Mazoyer³², leaving only regions which excised 10% volume activity, except supplementary, figure X, insert i.

Table II shows cortical networks that served for comparison with the classification of the results.

Figure IV shows the Jubrain-architecture⁴⁵⁻⁴⁷ aligned with the Mesulam model of hierarchies², simplified in figure V, used to sculpture in figures VI and VII models of cortical streams that served for comparison with the classification of the results.

Movie I shows the Mesulam model unfolded over time.

Object	Author	Brain coverage/nodes					Object	Author	Brain coverage/nodes				
		Whole brain	Cerebral cortex	Cerebellum	Subcortical	networks			Whole brain	Cerebral cortex	Cerebellum	Subcortical	networks
Space	Talairach Tournoux 1957	x					Function visual	Van Essen 1992		38			12
	Maziotta 1995	x						Ungerleider 1995		36			6
Macroanatomy	Ecker 1869		x				Function motoric	Glerean 2012		3			
	Smith 1907		x					Naser 2017		9			
	AAL Atlas 2002	116	83	23	10			Grandinetti 2013		12			
	Makris 2006	69						Pitzalis 2015		11			
	Desikan 2006		70					Yeo 2011		6			5
	Gousias 2008		83				Function rsfMRI	Zilles 1995		7(9)			
	Destrieux Atlas 2010		74					Smith 2009		18			10
	Mindboggle Klein 2012		50				Other	Andrews-Hanna 2010		11			3
	MarsAtlas Auzias 2016	x	41	7				Bellec 2010	7-444				
Cytoarchitecture	Campbell 1905		12					Power 2011	x				
	Brodman 1909		49					Choi 2012		17	x	x	7,17
	von Economo 1925		107					Craddock 2012	10 - 1000				
	Sarkisov 1949		64					Shen 2013	93,184,278				
	Bailey 1951		4					Spreng 2013		43			3
	Lacadie 2008		55					Wang 2015		x	x		18
	JuBrain Eickhoff 2005		100					AICHA Joliot 2015		122/50		20	
								Zhang 2015		15			9
Myeloarchitecture	Vogt 1919			x				Fan 2016			x		
	Flechsigs 1897 1920					5		Gordon 2016		333			
	Geschwind 1965		102		6	36		Braga 2017					3
	Catani 2005					5		Schaefer 2018		100-1000			
	Van Essen 2012		x	x	x			Kong 2018		x			17
Function descriptive	Gall 1791		33				Other	Huth 2016		160			
	Zilles 1994		49			13		Fan 2016		123		x	
	Van Essen 1998		49			7		Glasser 2016		180			
	Mesulam 2000		12			4		Varikuti 2018	2-500				

Table II. Nodes and networks in parcellations of the human brain.

The space, the anatomy, the cytoarchitecture, the myeloarchitecture and the functions supported by the aforementioned structures, for the whole cortex as well as for the visual and motoric subsystems. Given are the nodes (per hemisphere) and, if investigated, their organization in networks and hierarchies.

- Space – constituent Talairach²⁸ and MNI²⁹ spaces.
 - Macroanatomy - Different anatomical subdivisions, as per old observations such as Ecker³⁰, Smith³¹ and newer ones as Tzourio-Mazoyer & al.³², Harvard–Oxford Atlas Makris & al.³³, Desikan–Killiany Atlas Desikan & al.³⁴, Hammersmith Atlas Gousias & al.³⁵, Destrieux & al.³⁶, Klein & al.³⁷, Auzias & al.³⁸.
 - Cytoarchitecture – Campbell³⁹, Brodmann⁴⁰ and its transcription by the Yale Bioimage Suite Package Lacardie & al.⁴¹, von Economo & Koskinas⁴², Sarkisov & Filimonoff⁴³, Bailey & von Bonin⁴⁴, Zilles & al.^{45,46,47} Torso.
 - Myeloarchitecture - as per Vogt & Vogt⁴⁸, Flechsigs⁴⁹, Geschwind⁵⁰, Catani & Pfytych⁵¹ and the ongoing efforts of the Human-Connectome-Project⁵² on cortical parcelling.
 - Function descriptive – Gall⁵³, Zilles & Rehkaemper⁵⁴, Van Essen & al.⁵⁵, the two latter ones consolidating Brodmann's atlas.
 - Function visual – Van Essen & al.⁵⁶ (12 hierarchies), Ungerleider⁵⁷ (6 hierarchies), Glerean & al.⁵⁸ (dynamic regional phase synchrony), Naser & al.⁵⁹, Grandinetti & al.⁶⁰, Pitzalis & al.⁶¹, Yeo & al.⁶².
 - Function motoric - Zilles & al.⁶³ (integrating architectonic, transmitter receptor, MRI and PET data).
 - Function dfMRI – resting state functional magnetic resonance imaging observing networks (default mode network (consisting of a core, a dorsal medial and a medial temporal subsystem), executive control network, frontoparietal network, dorsal and ventral attention network, salience network, several visual and sensorimotor networks): Smith & al.⁶⁴ (ICA, 7000 subjects), Andrews-Hanna & al.^{65,66} (hierarchical relations within the default network), Bellec & al.⁶⁷, Power & al.⁶⁸, Yeo & al.⁶² (clustering ICA, 1000 subjects, separation of the early sensory and late motor cortices), Buckner & al.^{69,70} (seedbased ICA), Choi & al.^{71,72} (caudal-rostral parietal and rostro-caudal frontal hierarchies), Craddock & al.⁷³ (spectral clustering), Shen & al.⁷⁴, Spreng & al.⁷⁵ (hierarchical relations between three networks), Wang & al.⁷⁶ (interindividual), Joliot & al.⁷⁷, Zhang & al.⁷⁸ (directionality, subdividing two visual networks), Fan & al.⁷⁹ (eigenentropy), Gordon & al.⁸⁰ (individual), Braga & al.⁸¹ (within individual), Schaefer & al.⁸² (clustering and local boundary detection), Kong & al.⁸³ (individual, character-predictive), ix) Other – Huth & al.^{84,85} (language), Fan & al.⁸⁶ (multimodal, structure, connectivity, function), Glasser & al.⁸⁷ (multimodal), Varikuti & al.⁸⁸.
- See Eickhoff & al.⁸⁹ for a detailed discussion.

Figure IV. Temporal and spatial phasic hierarchies in Mesulam's² anatomical/histological/functional model (centre, adapted from Mesulam³) translate to the cytoarchitectures published in JuBrain⁴⁵⁻⁴⁷ (see methods movie I).

The right hemisphere, unfolded.

Waves might find their way in any direction, as, e.g., the top-down effect of the Bereitschaftspotential¹¹⁵⁻¹¹⁶ or the top-down effect of attention in early sensory cortex, largely invisible to spike recordings, but readily seen in the fMRI signal of monkeys^{13,90}.

Figure V. Mesulam^{2,3} hierarchies applied to the Brodmann^{40,41}/Zilles⁴⁵⁻⁴⁷-cytoarchitecture.

The right hemisphere of a normalized^{28,29} brain, unfolded. Monofocal histological regions named following Brodmann & Zilles. FG3, FG4, 2, hip3, Te2.1, Te2.2, TPJ, TI, HATA, amygdala, ID2-3, OP5-7, p24a, p24b, pd24sv, pd24cv, pd24cd, pv24c, s24b, s24a, 25p, 25a, p32, CH1-4 announced but not published.

8, 9, 38, 25, 34, 29, 26, 20, 35, 30, 23, 31, 24, 46, 47, 52 Brodmann-areas only.

The points give the centers of gravity and the lines give the long axes of the areas volumes out of figure IV.

The colors represent hierarchical hebbian layers, mono- or bisynaptic apart from each other.

Figure VI. The architecture of the ventral and dorsal visual stream.

Sculptured out of figure V.

Paragraphs: top row - concordance between Brodmann^{40,41}/Zilles⁴⁵⁻⁴⁷/Mesulam²/dfMRI^{62,64-66,70,72,75,78-81}/other; following rows - presumed functions^{2,56,57,59,60,62,61,58,91,97,104,108,105}, all using different number and localizations of nodes, as well as different numbers of hierarchical levels; bottom row - deficit-syndroms.

Red - The ventral stream ... shows a considerable functional heterogeneity from early visual processing (posterior) to higher, domain-specific processing (anterior)⁹¹, an ascending hierarchy with the sequence V1 – V2 (V3) – V4 – inferior temporal cortex (IT), starting with simpler features posteriorly (TEO/PIT) that increase in complexity as processing moves anteriorly (TE/AIT) to perform object recognition⁹², asymmetrical, tending to the right hemisphere⁹³:

17⁹⁴ - V1, primary visual, retinotopic two-dimensional picturing of the external world; 18⁹⁴ - gyrus lingualis, occipitalis inferior; hOc3v⁹⁵ - lingual cortex; FG1/FG2^{92,96} - fusiform cortex, orientation selective cells concerned with form, and direction selective cells concerned with motion. The fusiform gyrus hosts several "high-level" functional areas. It is involved in a core network subserving different cognitive tasks, that is, object recognition, visual language perception, or visual attention. It processes high-level visual information associated with faces, limbs, words, and places. FG2 is even a higher-order one⁹⁶;

Green - The dorsal stream comprises the sequence V1 – V2 (V3) – middle temporal cortex, MT, V5, always retinotopic – medial superior temporal cortex, MST – and finally intermediate object representations used by the decision-making circuitry further down the dorsal stream in lateral intraparietal sulcus, LIP⁹⁷; hOc3d⁹⁸, cuneus, V3, V3A. There is a medio-temporal component consisting of hOc5^{99,100} - temporo-occipito-parietal TOP-junction, related to the deficit symptom of neglect¹⁰¹; and BA 21 - medial temporal gyrus, it has been connected with processes as different as contemplating distance, recognition of known faces, and accessing word meaning while reading¹⁰²;

There is evidence to suggest that there is integration of both dorsal and ventral stream information into motion computation processes, giving rise to intermediate object representations, which facilitate object selection and decision-making mechanisms¹⁰³. The areas within the intraparietal sulcus, IPS, in particular, serve as interfaces between the perceptive and motor systems for controlling arm and eye movements in space¹⁰⁴. This region is functionally and histologically complex in so far as several association areas, default mode, executive control and dorsal attention^{105,106} overlap in here as well as the cytoarchitectonic areas 7A, 7PC, mip, Ip1, Ip2, PF and PFm, sulcus intraparietalis, concerned with the integration of multimodal

information for constructing a spatial representation of the external world in relation to the body or parts thereof, and planning and executing object-centered movements¹⁰⁴; 7P - precuneus, integration to a complex space out of proprioceptive and visual sources, information on objects in depth which are translating in space, because of the self-motion, object distance in a dynamic condition (as that created by self-motion) to orchestrate the eye and arm movements necessary to reach or avoid static and moving objects in the environment¹⁰⁷, self-perception¹⁰⁸. Premotor area engaged in the mental operation, the precuneus aids monitoring the operation in terms of internally represented visual images. Interfaces between the perceptive and motor systems for controlling arm and eye movements in space. Visuomotor tasks comprising target selections for arm and eye movements, object manipulation and visuospatial attention, deficit syndrome of apraxia, neglect, Bálint's syndrome¹⁰⁹.

Figure VII. The sensorimotor architecture.

Sculptured out of figure V.

Paragraphs: top row - concordance between Brodmann^{40,41}/Zilles⁴⁵⁻⁴⁷/Mesulam²/dfMRI^{62,64-66,70,72,75,78-81}/other; following rows - presumed functions^{12,60,104,110,118}.

aCMA - cingulum anterior, its proposed role in cognition and its relationship with the arousal/drive state of the organism placing it in a unique position to translate intentions to actions¹¹⁰, and being dedicated to the limbic portion of the cortex; 6¹¹¹ - pre-SMA and SMA proper^{112,113}, intention to act, generation of readiness potential^{114,115,116,12}, action planning and gesture recognition, the operation of the motor monitoring, namely detecting the mismatch between the actual motor output and intended action goal¹¹⁷. SMA proper and pre-SMA are active prior to volitional movement or action, as well as the cingulate motor area (CMA). This is now known as 'anterior mid-cingulate cortex (aMCC)'. It has been shown by integrating simultaneously acquired EEG and fMRI that SMA and aMCC have strong reciprocal connections that act to sustain each other's activity, and that this interaction is mediated during movement preparation according to the Bereitschaftspotential amplitude¹¹²; 6aβ - the supplementary eye field (SEF), which has a special role in high level aspects of saccade control, such as complex spatial transformations, learned transformations, and executive cognitive functions^{118,119}; 6aα superior - visuo-somatomotor area hosting the frontal eye field (FEF)¹²⁰, which constitutes together with the intraparietal sulcus (IPS) ... an important role in the control of visual attention and eye movements¹⁰⁴, the coding of object positions⁵⁹; 4p¹²¹ - central sulcus, primary area, involved in kinematic and dynamic parameters of voluntary movements¹²¹; 3a/3b, 1¹²² - primary area; 2¹²³ - the projection emphasising size and shape, secondary unimodal; 5¹²⁴ - secondary, unimodal.

Movie I. Traveling cortical Netwave, hypothesis¹²⁵. <https://youtu.be/cQVBLAThJqk>
 Jubrain⁴⁵⁻⁴⁷ regions aligned over time and neighbourhood from the visual perception to the voluntary motor reaction, and back over the autoperception and heteromodal association, assigned to the cortical hierarchies (coloured box bottom, right) of Mesulam², unfolded right cortical hemisphere.

SUPPLEMENTARY

Movie II. Traveling cortical Netwave, results¹²⁶. <https://youtu.be/efEaO79gBkM>

Part i - The time series for the visual perception, 2 seconds ante until 5 seconds post perception, perception at $t = 0$ s; last picture the resulting phase image.

Part ii - The right and left hand motor reaction, 3.5 seconds ante until 2 seconds post reaction, movement at $t = 0$ s; last picture the resulting phase image.

The red-white boxes at the bottom represent the time axis in seconds.

Activity strength (scaled t-values of hrf-correlation) following the colour bar at the bottom, but in frame 12 and 27, the resulting phase images, where colour codes time.

Figure VIII. Perception-related time series.

Time steps in second distances (2 seconds ante until 5 seconds post perception, white numbers). Perception at time step three. Activity strength following the colour bar (see supplementary movie II, part i).

Figure IX. Reaction-related time series.

Right hand - above; left hand - below. Time steps in half second distances (3 seconds ante until 2.5 seconds post reaction, white numbers). Reaction at time step eight. Activity strength following the colour bar (see supplementary movie II, part ii).

Table III. Right and left hemisphere in the AAL³²-network.

Visual signal, moving left and right hand (blue digits); AAL - Anatomical regions following Tzourio-Mazoyer³²; mm³ - activated volume; t_{mean} - mean-time of maximal activation over all voxels within the region in seconds, relating to the signal = 0 s (standard deviation 0.5 to 1 s); blue numbers for the movement, red for the signal. Black numbers indicate that this region activated under all conditions (the results for the SMA of the visual branch in the table, the results for the other conditions below); f - frequency in Hertz; lha - left hand; rha - right hand; nw - network. The colouration of the waves corresponds to Mesulam².

The individual *standing network waves* are approximated by

$$I_r = e^{-(t-t_{er})^2 f}$$

where r is the coordinate, its base-normalised regional nodes activity I_r , the time t , the mean time point t_{er} of the maximum regional activities, and frequency f .

Phase-shifted waves develop in parallel as *winning coalitions* and form a *forward traveling network wave* across time and space.

Figure X. Contrast of delayed and speeded responses, phase images.

The unique hierarchical position of the complex made of the parietal Brodmann area 7, a core area of the dorsal attention network^{62,70,72,75,78-81} becomes clear in contrast imaging of speeded and delayed-response tasks. The speeded responses (right) activate all of the same regions as the sum of the delayed responses (the visual phase image on the left, next to it both motor phase images) – with the exception of the high-association superior parietal regions (precuneus and parietalis superior, BA 7, pink rings). The high-association areas are apparently used only when they are involved in long-lasting feedforward-feedback-loops or, in short, in the case of accelerated responses, there is no time left for daydreaming.

Movie III. Traveling cortical Network, interpretation¹²⁷. <https://youtu.be/fvLWPrw2Ro0>

The fMRI time series for the visual perception and the following the left hand motor act, fused out of sup. movie II, using the medial supplementary motor area (resulting time axis 2 seconds ante until 7.5 seconds post perception, 7 seconds ante until 2.5 seconds post reaction),

The red-white boxes at the bottom represent the time axis, scaled in seconds above with respect to the visual signal and below with respect to the motor reaction.

Activity strength (scaled t-values of hrf-correlation) following the colour bar at the bottom.

The anatomical regions labelled following Jubrain^{45-47,91,94-96,98,99,100,109}.

The evolution of different standing networks, the resulting travelling network and winning coalitions⁵ following AAL³² assigned to the hierarchies of Mesulam^{2,3} in the insert above left (see sup. tab. III).

Frame 20 gives the surface rendered phase image, right hemisphere, as well as the extreme values in space and time (left table) and the resulting vector-matrix (right table).

Colour codes time, with respect to the signal (upper timescale) and to the reaction (lower timescale).

Labelling JuBrain. Temporo-spatial cortical vectors red. Their position in space (x, y, z, Talairach-space (blue boxes) in mm) and time (t in s) as shown in the left table. Their length in time (dt in s) and space (ds in mm) in the right table.

REFERENCES

- ¹ Hebb D. The Organization of Behavior. Wiley New York (1949).
- ² Mesulam MM. From sensation to cognition. *Brain* 121, 1013-1052 (1998).
- ³ Mesulam MM. Principles of Behavioral and Cognitive Neurology. *Oxford University Press* (2000).
- ⁴ Deco G. Tononi G. Boly M. Kringelbach ML. Rethinking segregation and integration: contributions of whole-brain modelling. *Nat Rev Neurosci.* 16, 430-439 (2015).
- ⁵ Crick F. Koch C. A framework of consciousness. *Nat Neurosci.* 6, 119-126 (2003).
- ⁶ Deco G. Jirsa VK. Robinson PA. Breakspear M. Friston K. The dynamic brain: from spiking neurons to neural masses and cortical fields. *PLoS Comput Biol.* 4, (2008).
- ⁷ Nieuwenhuys R. Donkelaar HJ. Nicholson C. The central nervous system of vertebrates, Volume 1. *Springer*, 2011–2012 (1998).
- ⁸ Toro R. Perron M. Pike B. Richer L. Veillette S. Pausova Z. Paus T. Brain Size and Folding of the Human Cerebral Cortex. *Cerebral Cortex* 18, 2352–2357 (2008).
- ⁹ Haynes JD. Rees G. Decoding mental states from brain activity in humans. *Nature Reviews Neuroscience* 7, 523-534 (2006).
- ¹⁰ He BJ. Raichle ME. The fMRI signal, slow cortical potential and consciousness. *Trends Cogn Sci.* 13, 302-309 (2009).
- ¹¹ Birbaumer N. Elbert T. Canavan AG. Rockstroh B. Slow potentials of the cerebral cortex and behavior. *Physiol Rev.* 70, 1-41 (1990).
- ¹² Montemurro MA. Rasch MJ. Murayama Y. Logothetis NK. Panzeri S. Phase-of-firing coding of natural visual stimuli in primary visual cortex. *Curr. Biol* 18, 375–380 (2008).
- ¹³ Lakatos P. Karmos G. Mehta AD. Ulbert I. Schroeder CE. Entrainment of neuronal oscillations as a mechanism of attentional selection. *Science* 320, 110–113 (2008).
- ¹⁴ Muller L. Chavane F. Reynolds J. Sejnowski TJ. Cortical travelling waves: mechanisms and computational principle. *Nat Rev Neurosci.* 19, 255-268 (2018).
- ¹⁵ Calhoun V. Liégeois R. Time-varying Connectivity in Resting-state fMRI: Methods, interpretations and clinical use. *OHBM* (2019).
- ¹⁶ Lurie D. Controversies and Open Questions in the Study of “Resting State” Time-Varying Functional Connectivity. *OHBM* (2019).
- ¹⁷ Breakspear M. Modelling the Multiscale Nature of Dynamic Functional Connectivity. *OHBM* (2019).
- ¹⁸ Breakspear M. Terry JR. Friston KJ. Modulation of excitatory synaptic coupling facilitates synchronization and complex dynamics in a biophysical model of neuronal dynamics. *Network* 14, 703-732 (2003).
- ¹⁹ Erb M. Aertsen A. Dynamics of Activity in Biology-Oriented Neural Network Models: Stability at Low Firing Rates. In: Information Processing in the Cortex. Experiment and Theory. Edited by Braitenberg V. & Aertsen A. *Springer* (1992).
- ²⁰ Omidvarnia A. Pedersen M. Walz JM. Vaughan DN. Abbott DF. Jackson GD. Dynamic regional phase synchrony (DRePS): An Instantaneous Measure of Local fMRI Connectivity Within Spatially Clustered Brain Areas. *Hum Brain Mapp.* 37, 1970-1985 (2016).
- ²¹ Friston KJ. Ashburner JT. Kiebel SJ. Nichols TE. Penny WD. Statistical Parametric Mapping. *Academic Press* (2008).
- ²² Fornito A. Zalesky Z. Bullmore E. Fundamentals of Brain Network Analysis. *Academic Press* (2016).
- ²³ Fernández-Corazza M. Turovets S. Luu P. Anderson E. Tucker D. Transcranial Electrical Neuromodulation Based on the Reciprocity Principle. *Front. Psychiatry* 27 May (2016).
- ²⁴ Huelsmann E. Erb M. Grodd W. From will to action: sequential cerebellar contributions to voluntary movement. *Neuroimage* 20, 1485-1492 (2003).

- ²⁵ Nolte J. (editor). The Human Brain: An Introduction to its functional Anatomy. *Elsevier*, 510-517 (2008).
- ²⁶ Bordier C. Macaluso E. Time-resolved detection of stimulus/task-related networks, via clustering of transient intersubject synchronization. *Hum Brain Mapp.* 36, 3404-3425 (2015).
- ²⁷ Cohen MX. Analysing Neural Time Series Data. *The MIT Press* (2014).
- ²⁸ Talairach J. David M. Tournoux P. Atlas d'anatomie stéréotaxique. *Masson Paris* (1957).
- ²⁹ Mazziotta JC. Toga AW. Evans AP. Fox A. Lancaster J. A Probabilistic Atlas of the Human Brain: Theory and Rationale for Its Development. *Neuroimage* 2, 89-101 (1995).
- ³⁰ Ecker A. Die Hirnwindungen des Menschen: nach eigenen Untersuchungen; insbesondere über die Entwicklung derselben beim Fötus und mit Rücksicht auf das Bedürfnis der Ärzte. *Vieweg Braunschweig* (1869).
- ³¹ Smith EG. A new topographical survey of the human cerebral cortex, being an account of the distribution of the anatomically distinct cortical areas and their relationship to the cerebral sulci. *J. Anat.* 41, 237-254 (1907).
- ³² Tzourio-Mazoyer N. Landeau B. Papathanassiou D. Crivello F. Etard O. Delcroix N. Mazoyer B. Joliot M. Automated anatomical labeling of activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject brain. *Neuroimage* 15, 273-289 (2002).
- ³³ Makris, N. Goldstein JM. Kennedy D. Hodge SM. Caviness VS. Faraone SV. Tsuang MT. Seidman LJ. Decreased volume of left and total anterior insular lobule in schizophrenia. *Schizophr. Res.* 83, 155-171 (2006).
- ³⁴ Desikan RS. Ségonne F. Fischl B. Quinn BT. Dickerson BC. Blacker D. Buckner RL. Dale AM. Maguire RP. Hyman BT. Albert MS. Killiany RJ. An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. *Neuroimage* 31, 968-980 (2006).
- ³⁵ Gousias IS. Rueckert D. Heckemann RA. Dyet LE. Boardman JP. Edwards AD. Hammers A. Automatic segmentation of brain MRIs of 2-year-olds into 83 regions of interest. *NeuroImage* 40, 672-684 (2008).
- ³⁶ Destrieux C. Fischl B. Dale A. Halgren E. Automatic parcellation of human cortical gyri and sulci using standard anatomical nomenclature. *Neuroimage* 53, 1-15 (2010).
- ³⁷ Klein A. Tourville J. 101 labeled brain images and a consistent human cortical labeling protocol. *Frontiers in Neuroscience* 6, 171 (2012).
- ³⁸ Auzias G. Coulon O. Brovelli A. MarsAtlas: A cortical parcellation atlas for functional mapping. *Human Brain Mapping* 37, 1573-1592 (2016).
- ³⁹ Campbell AW. Histological Studies on the Localisation of Cerebral Function. *Cambridge Univ. Press*, Cambridge, UK (1905).
- ⁴⁰ Brodmann K. Vergleichende Lokalisationslehre der Großhirnrinde in ihren Prinzipien dargestellt auf Grund ihres Zellenbaues. *Barth, Leipzig* (1909).
- ⁴¹ Lacadie CM. Fulbright RK. Arora J. Constable RT. Papademetris X. Brodmann Areas defined in MNI space using a new Tracing Tool in BioImage Suite. *Human Brain Mapping*, (2008).
- ⁴² von Economo C. Koskinas GN. Die Cytoarchitektonik der Hirnrinde des erwachsenen Menschen. *Springer Verlag Wien* (1925).
- ⁴³ Sarkisov SA. Filimonoff IN. Preobrashenskaya NS. Cytoarchitecture of the human cortex cerebri. *Medgiz Moscow* (1949).
- ⁴⁴ Bailey P. von Bonin G. The Isocortex of Man. *Univ. Illinois Press*, Urbana (1951).
- ⁴⁵ Amunts K. Zilles K. Architectonic Mapping of the Human Brain beyond Brodmann. *Neuron* 16, 1086-1107 (2015).
- ⁴⁶ Eickhoff SB. Stephan KE. Mohlberg H. Grefkes C. Fink GR. Amunts K. Zilles K. A new SPM toolbox for combining probabilistic cytoarchitectonic maps and functional imaging data. *Neuroimage* 25, 1325-1335 (2005).
- ⁴⁷ Mohlberg H. Eickhoff SB. Schleicher A. Zilles K. Amunts K. A new processing pipeline and release of cytoarchitectonic probabilistic maps – JuBrain. *OHBM Peking, China* (2012).

- ⁴⁸ Vogt C. Vogt O. Allgemeine Ergebnisse unserer Hirnforschung. *J Psychol Neural.* 25, 277-462 (1919).
- ⁴⁹ Flechsig PE. Anatomie des menschlichen Gehirns und Rückenmarks auf myelogenetischer Grundlage, Volume 1. Anatomie des menschlichen Gehirns und Rückenmarks auf myelogenetischer Grundlage. *G. Thieme* (1920).
- ⁵⁰ Geschwind N. Disconnexion syndromes in animals and man. I. *Brain* 88, 237–294 (1965).
- ⁵¹ Catani M. Ffytch DH. The rises and falls of disconnection syndromes. *Brain* 128, 2224–2239 (2005).
- ⁵² Van Essen DC. Ugurbil K. Auerbach E. Barch D. Behrens TE. Bucholz R. Chang A. Chen L. Corbetta M. Curtiss SW. Della Penna S. Feinberg D. Glasser MF. Harel N. Heath AC. Larson-Prior L. Marcus D. Michalareas G. Moeller S. Oostenveld R. Petersen SE. Prior F. Schlaggar BL. Smith SM. Snyder AZ Xu J. Yacoub E. The Human Connectome Project: a data acquisition perspective. *Neuroimage* 62, 2222-2231 (2012).
- ⁵³ Gall FJ. Philosophisch-medizinische Untersuchungen über Natur und Kunst im kranken und gesunden Zustand des Menschen. *Gräffer, Wien* (1791).
- ⁵⁴ Zilles K. Rehkaemper G. Funktionelle Neuroanatomie. *Springer* (1993).
- ⁵⁵ Van Essen DC. Drury HA. Joshi S. Miller MI. Functional and structural mapping of human cerebral cortex: solutions are in the surfaces. *Adv Neurol.* 84, 23-34 (2000).
- ⁵⁶ Van Essen DC. Anderson CH. Felleman DJ. Information processing in the primate visual system: an integrated systems perspective. *Science* 255, 419-423 (1992).
- ⁵⁷ Ungerleider LG. Functional brain imaging studies of cortical mechanisms for memory. *Science* 270, 769-775 (1995).
- ⁵⁸ Glerean E. Salmi J. Lahnakoski JM. Iiro P. Jääskeläinen IP. Mikko Sams M. Functional Magnetic Resonance Imaging Phase Synchronization as a Measure of Dynamic Functional Connectivity. *Brain Connect.* 2, 91–101 (2012).
- ⁵⁹ Naser K. Ricordel V. Le Callet P. in Benois-Pineau J. Le Callet P. Visual content indexing and retrieval with psycho-visual models. *Springer*, 20 (2017).
- ⁶⁰ Grandinetti L. Lippert T. Petkov N. Brain-inspired computing. *Springer*, 53 (2013).
- ⁶¹ Pitzalis S. Fattori P. Galletti C. The human cortical areas V6 and V6A. *Vis Neurosci.*, 32 (2015).
- ⁶² Yeo BT. Krienen FM. Sepulcre J. Sabuncu MR. Lashkari D. Hollinshead M. Roffman JL. Smoller JW. Zollei L. Polimeni JR. Fischl B. Liu H. Buckner RL. The organization of the human cerebral cortex estimated by intrinsic functional connectivity. *J Neurophysiol.* 106, 1125-1165 (2011).
- ⁶³ Zilles K. Schlaug G. Matelli M. Luppino G. Schleicher A. Qü M. Dabringhaus A. Seitz R. Roland PE. Mapping of human and macaque sensorimotor areas by integrating architectonic, transmitter receptor, MRI and PET data. *J Anat.* 187, 515-537 (1995).
- ⁶⁴ Smith SM. Fox PT. Miller KL. Glahn DC. Fox PM. Mackay CE. Filippini N. Watkins KE. Toro R. Laird AR. Beckmann CF. Correspondence of the brain's functional architecture during activation and rest. *Proc Natl Acad Sci USA.* 106, 13040-13045 (2009).
- ⁶⁵ Andrews-Hanna JR. Smallwood J. Spreng RN. The default network and self-generated thought: component processes, dynamic control, and clinical relevance. *Ann NY Acad Sci.* 1316, 29-52 (2014).
- ⁶⁶ Andrews-Hanna JR. Reidler JS. Sepulcre J. Poulin R. Buckner RL. Functional-anatomic fractionation of the brain's default network. *Neuron* 65, 550-562 (2010).
- ⁶⁷ Bellec P. Rosa-Neto P. Lyttelton OC. Benali H. Evans AC. Multi- level bootstrap analysis of stable clusters in resting- state fMRI. *NeuroImage* 51, 1126–1139 (2010).
- ⁶⁸ Power JD. Cohen AL. Nelson SM. Wig GS. Barnes KA. Church JA. Vogel AC. Laumann TO. Miezin FM. Schlaggar BL. Petersen SE. Functional network organization of the human brain. *Neuron* 72, 665–678 (2011).
- ⁶⁹ Buckner RL. Krienen FM. Castellanos A. Diaz JC. Yeo BT. The organization of the human cerebellum estimated by intrinsic functional connectivity. *J. Neurophysiol.* 106, 2322–2345 (2011).
- ⁷⁰ Buckner RL. Krienen FM. Yeo BTT. Opportunities and limitations of intrinsic functional connectivity MRI. *Nature Neuroscience* 16, 832–837 (2013).

- ⁷¹ Choi EY, Yeo BT, Buckner RL. The organization of the human striatum estimated by intrinsic functional connectivity. *J Neurophysiol.* 108, 2242–2263 (2012).
- ⁷² Choi EY, Drayna GK, Badre D. Evidence for a Functional Hierarchy of Association Networks. *J Cogn Neurosci.* 30, 722–736 (2018).
- ⁷³ Craddock RC, James GA, Holtzheimer PE, Hu XP, Mayberg HS. A whole brain fMRI atlas generated via spatially constrained spectral clustering. *Human Brain Mapping* 33, 1914–1928 (2012).
- ⁷⁴ Shen X, Tokoglu F, Papademetris X, Constable, RT. Groupwise whole-brain parcellation from resting-state fMRI data for network node identification. *NeuroImage* 82, 403–415 (2013).
- ⁷⁵ Spreng RN, Sepulcre J, Turner GR, Stevens WD, Schacter DL. Intrinsic architecture underlying the relations among the default, dorsal attention, and frontoparietal control networks of the human brain. *J Cogn Neurosci.* 25, 74–86 (2013).
- ⁷⁶ Wang D, Buckner RL, Fox MD, Holt DJ, Holmes AJ, Stoecklein S, Langs G, Pan R, Qian T, Li K, Baker JT, Stufflebeam SM, Wang K, Wang X, Hong B, Liu H. Parcellating cortical functional networks in individuals. *Nat Neurosci.* 18, 1853–1860 (2015).
- ⁷⁷ Joliot M, Jobard G, Naveau M, Delcroix N, Petit L, Zago L, Crivello F, Mellet E, Mazoyer B, Tzourio-Mazoyer N. AICHA: An atlas of intrinsic connectivity of homotopic areas. *Journal of Neuroscience Methods* 254, 46–59 (2015).
- ⁷⁸ Zhang D, Liang B, Wu X, Wang Z, Xu P, Chang S, Liu B, Liu M, Huang R. Directionality of large-scale resting-state brain networks during eyes open and eyes closed conditions. *Front Hum Neurosci.* 9, 81 (2015).
- ⁷⁹ Fan L, Li H, Zhuo J, Zhang Y, Wang J, Chen L, Yang Z, Chu C, Xie S, Laird AR, Fox PT, Eickhoff SB, Yu C, Jiang T. The Human Brainnetome Atlas: A New Brain Atlas Based on Connectional Architecture. *Cerebral Cortex* 26, 3508–3526 (2016).
- ⁸⁰ Gordon EM, Laumann TO, Adeyemo B, Huckins JF, Kelley WM, Petersen SE. Generation and Evaluation of a Cortical Area Parcellation from Resting-State Correlations. *Cerebral Cortex* 26, 288–303 (2016).
- ⁸¹ Braga RM, Buckner RL. Parallel interdigitated distributed networks within the individual estimated by intrinsic functional connectivity. *Neuron* 95, 457–472 (2017).
- ⁸² Schaefer A, Kong R, Gordon EM, Laumann TO, Zuo XN, Holmes AJ, Eickhoff SB, Yeo BTT. Local-global parcellation of the human cerebral cortex from intrinsic functional connectivity MRI. *Cerebral Cortex* 28, 3095–3114 (2017).
- ⁸³ Kong R, Li J, Sun N, Sabuncu MR, Liu H, Schaefer A, Orban C, Zuo XN, Holmes AJ, Eickhoff S, Yeo B.T. Spatial topography of individual-specific cortical networks predicts human cognition, personality and emotion. *Cerebral Cortex* <https://doi.org/10.1093/cercor/bhy123> (2018).
- ⁸⁴ Huth AG, de Heer WA, Griffiths TL, Theunissen FE, Gallant JL. Natural speech reveals the semantic maps that tile human cerebral cortex. *Nature* 532, 453–458 (2016).
- ⁸⁵ Huth AG, Griffiths TL, Theunissen FE, Gallant JL. PrAGMATiC: a probabilistic and generative model of areas tiling the cortex. *arXiv* (2015).
- ⁸⁶ Fan Y, Zeng LL, Shen H, Qin J, Li F, Hu D. Lifespan Development of the Human Brain Revealed by Large-Scale Network Eigen-Entropy. *Entropy* 19, 471 (2017).
- ⁸⁷ Glasser MF, Coalson TS, Robinson EC, Hacker CD, Harwell J, Yacoub E, Ugurbil K, Andersson J, Beckmann CF, Jenkinson M, Smith SM, Van Essen DC. A multi-modal parcellation of human cerebral cortex. *Nature* 536, 171–178 (2016).
- ⁸⁸ Varikuti DP, Genon S, Sotiras A, Schwender H, Hoffstaedter F, Patil KR, Jockwitz C, Caspers S, Moebus S, Amunts K, Davatzikos C, Eickhoff SB. Evaluation of non-negative matrix factorization of grey matter in age prediction. *NeuroImage* 173, 394–410 (2018).
- ⁸⁹ Eickhoff SB, Yeo BTT, Genon S. Imaging-based parcellations of the human brain. *Nat Rev Neurosci.* 19, 672–686 (2018).
- ⁹⁰ Logothetis NK. The neural basis of the blood-oxygen-level-dependent functional magnetic resonance imaging signal. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 357, 1003–1037 (2002).
- ⁹¹ Caspers J, Zilles K, Amunts K, Laird AR, Fox PT, Eickhoff SB. Functional characterization and differential coactivation patterns of two cytoarchitectonic visual areas on the human posterior fusiform gyrus. *Hum Brain Mapp.* 6, 2754–2767 (2014).

- ⁹² Cowey A. Weiskrantz L. A comparison of the effects of inferotemporal and striate cortex lesions on the visual behaviour of rhesus monkeys. *J Exp Psychol.* 3, 246-253 (1967).
- ⁹³ Karnath HO. Insights into the functions of the superior temporal cortex. *Nat Rev Neurosci.* 2, 568-575 (2001).
- ⁹⁴ Amunts K. Malikovic A. Mohlberg H. Schormann T. Zilles K. Brodmann's areas 17 and 18 brought into stereotaxic space-where and how variable? *Neuroimage* 11, 66-84 (2000).
- ⁹⁵ Rottschy C. Eickhoff SB. Schleicher A. Mohlberg H. Kujovic M. Zilles K. Amunts K. Ventral visual cortex in humans: cytoarchitectonic mapping of two extrastriate areas. *Human Brain Mapp.* 28, 1045-1059 (2007).
- ⁹⁶ Lorenz S. Weiner KS. Caspers J. Mohlberg H. Schleicher A. Bludau S. Eickhoff SB. Grill-Spector K. Zilles K. Amunts K. Two New Cytoarchitectonic Areas on the Human Mid-Fusiform Gyrus. *Cerebral Cortex.* 27, 373-385 (2017).
- ⁹⁷ van Polanen V. Davarea M. Interactions between dorsal and ventral streams for controlling skilled grasp. *Neuropsychologia* 79, 186-191 (2015).
- ⁹⁸ Kujovic M. Zilles K. Malikovic A. Schleicher A. Mohlberg H. Rottschy C. Eickhoff SB. Amunts K. Cytoarchitectonic mapping of the human dorsal extrastriate cortex. *Brain Struct Funct.* 218, 157-172 (2013).
- ⁹⁹ Malikovic A. Amunts K. Schleicher A. Mohlberg H. Eickhoff SB. Wilms M. Palomero-Gallagher N. Armstrong E. Zilles K. Cytoarchitectonic analysis of the human extrastriate cortex in the region of V5/MT+: a probabilistic, stereotaxic map of area hOc5. *Cereb Cortex* 17, 562-574 (2007).
- ^{100b} Malikovic A. Amunts K. Schleicher A. Mohlberg H. Kujovic M. Palomero-Gallagher N. Eickhoff SB. Zilles K. Cytoarchitecture of the human lateral occipital cortex: mapping of two extrastriate areas hOc4la and hOc4lp. *Brain Structure and Function*, 221,1877-1897 (2016).
- ¹⁰⁰ Caspers S. Eickhoff SB. Geyer S. Scheperjans F. Mohlberg H. Zilles K. Amunts K. The human inferior parietal lobule in stereotaxic space. *Brain Struct Funct.* 212, 481-495 (2008).
- ¹⁰¹ Vallar G. Spatial hemineglect in humans. *Trends Cogn Sci.* 2, 87-95 (1998).
- ¹⁰² Acheson DJ. Hagoort P. Stimulating the brain's language network: syntactic ambiguity resolution after TMS to the inferior frontal gyrus and middle temporal gyrus. *J Cogn Neurosci.*, 1664-1677 (2013).
- ¹⁰³ Perry CJ. Fallah M. Feature integration and object representations along the dorsal stream visual hierarchy. *Front Comput Neurosci.* 8, 84 (2014).
- ¹⁰⁴ Grefkes C. Fink GR. The functional organization of the intraparietal sulcus in humans and monkeys. *J Anat.* 207, 3-17 (2005).
- ¹⁰⁵ Oshio R. Tanaka S. Sadato N. Sokabe M. Hanakawa T. Honda M. Differential effect of double-pulse TMS applied to dorsal premotor cortex and precuneus during internal operation of visuospatial information. *Neuroimage* 49, 1108-1115 (2010).
- ¹⁰⁶ Kjaer TW. Nowak M. Lou HC. Reflective self-awareness and conscious states: PET evidence for a common midline parietofrontal core. *Neuroimage* 17, 1080-1086 (2002).
- ¹⁰⁷ Pitzalis S. Fattori P. Galletti C. The human cortical areas V6 and V6A. *Vis Neurosci.* 32 (2015).
- ¹⁰⁸ Birbaumer N. Sanyung K. Dein Gehirn weiss mehr als Du denkst. *Ullstein*, 230-233 (2016).
- ¹⁰⁹ Choi HJ. Zilles K. Mohlberg H. Schleicher A. Fink GE. Armstrong E. Amunts K. Cytoarchitectonic identification and probabilistic mapping of two distinct areas within the anterior ventral bank of the human intraparietal sulcus. *J Comp Neurol.* 495, 53-69 (2006).
- ¹¹⁰ Paus T. Primate anterior cingulate cortex: Where motor control, drive and cognition interface. *Nat Rev Neurosci.* 2, 417-424 (2001).
- ¹¹¹ Geyer S. Matelli M. Luppino G. Zilles K. Functional neuroanatomy of the primate isocortical motor system *Anat Embryol.* 202, 443-474 (2000).
- ¹¹² Nguyen VT. Breakspear M. Cunnington R. Reciprocal interactions of the SMA and cingulate cortex sustain pre-movement activity for voluntary actions. *J Neurosci.* 34, 16397-16407 (2014).

- ¹¹³ Picard N. PL. Motor areas of the medial wall: a review of their location and functional activation. *Cerebral Cortex* 6, 342-53 (1996).
- ¹¹⁴ Libet B. Gleason CA. Wright EW. Pearl DK. Time of conscious intention to act in relation to onset of cerebral activity (readiness-potential). The unconscious initiation of a freely voluntary act. *Brain* 106, 623-842 (1983).
- ¹¹⁵ Kornhuber HH. Deecke L. Changes in the brain potential in voluntary movements and passive movements in mean readiness potential and reafferent potentials. *Pflugers Arch Gesamte Physiologie Menschen Tiere*. 284, 1-17 (1965).
- ¹¹⁶ Ruan J. Bludau S. Palomero-Gallagher N. Caspers S. Mohlberg H. Eickhoff SB. Seitz Rj. Amunts K. Cytoarchitecture, probability maps, and functions of the human supplementary and pre-supplementary motor areas. *Brain Struct Funct*. 9, 4169-4186 (2018).
- ¹¹⁷ Bolognini N. Cognitive processes of motor behavior revealed by tDCS. *Front Neurol*. 8, 29 (2017).
- ¹¹⁸ Kirchner H. Barbeau EJ. Thorpe SJ. Régis J. Liégeois-Chauvel C. Ultra-rapid sensory responses in the human frontal eye field region. *Journal of Neuroscience* 29, 7599–7606 (2009).
- ¹¹⁹ Graziano MSA. The Intelligent Movement Machine. *Oxford University Press* (2008).
- ¹²⁰ Schall JD. On the role of frontal eye field in guiding attention and saccades. *Vision Research* 44, 1453–1467 (2004).
- ¹²¹ Geyer S. Ledberg A. Schleicher A. Kinomura S. Schormann T. Bürgel U. Klingberg T. Larsson J. Zilles K. Roland PE. Two different areas within the primary motor cortex of man. *Nature* 382, 805-807 (1996).
- ¹²² Geyer S. Schleicher A. Zilles K. Areas 3a, 3b, and 1 of human primary somatosensory cortex. *Neuroimage* 10, 63-83 (1999).
- ¹²³ Grefkes C. Geyer S. Schormann T. Roland P. Zilles K. Human somatosensory area 2: observer-independent cytoarchitectonic mapping, interindividual variability, and population map. *Neuroimage* 14, 617-31 (2001).
- ¹²⁴ Scheperjans F. Eickhoff SB. Hömke L. Mohlberg H. Hermann K. Amunts K. Zilles K. Probabilistic Maps, Morphometry, and Variability of Cytoarchitectonic Areas in the Human Superior Parietal Cortex. *Cereb. Cortex* 18, 2141–2157 (2008).
- ¹²⁵ Hülsmann ERM. Traveling cortical netwave, hypothesis. *Youtube* (2019).
- ¹²⁶ Hülsmann ERM. Traveling cortical netwave, results. *Youtube* (2019).
- ¹²⁷ Hülsmann ERM. Traveling cortical netwave, interpretation. *Youtube* (2019).