LxxIxE-like Motif in Spike Protein of SARS-CoV-2 that is Known to Recruit the Host PP2A-B56 Phosphatase Mimics Artepillin C, an Immunomodulator, of Brazilian Green Propolis

Halim Maaroufi

Institut de biologie intégrative et des systèmes (IBIS). Université Laval. Québec, Canada.

Halim.maaroufi@ibis.ulaval.ca

ABSTRACT

SARS-CoV-2 is highly contagious and can cause acute respiratory distress syndrome (ARDS) and multiple organ failure that are largely attributed to the cytokine storm. The surface coronavirus spike (S) glycoprotein is considered as a key factor in host specificity because it mediates infection by receptor-recognition and membrane fusion. Here, the analysis of SARS-CoV-2 S protein revealed two B56-binding LxxIxE-like motifs in S1 and S2 subunits that could recruit the host protein phosphatase 2A (PP2A). The motif in S1 subunit is absent in SARS-CoV and MERS-CoV. Phosphatases and kinases are major players in the regulation of pro-inflammatory responses during pathogenic infections. Moreover, studies have shown that viruses target PP2A in order to manipulate host's antiviral responses. Recent researches have indicated that SARS-CoV-2 is involved in sustained host inflammation. Therefore, by controlling acute inflammation, it is possible to eliminate its dangerous effects on the host. Among efforts to fight COVID-19, the interaction between LxxIxE-like motif and the PP2A-B56-binding pocket could be a target for the discovery and/or development of a bioactive ligand inhibitor for therapeutic purposes. Indeed, a small molecule called Artepillin C (ArtC), a main compound in Brazilian honeybee green propolis, mimics the side chains of LxxLxE motif. Importantly, ArtC is known, among other effects, to have anti-inflammatory activity that makes it an excellent candidate for future clinical trials in COVID-19 patients.

KEYWORDS SARS-CoV-2; spike glycoprotein; PP2A-B56 phosphatase; LxxIxE-like motif; anti-inflammation; Artepillin C.

INTRODUCTION

In March 11th 2020, the World Health Organization (WHO) announced that COVID-19 (Coronavirus Disease-2019) situation is a pandemic. The novel SARS-CoV-2 has had serious consequences for human health and socioeconomic stability worldwide. Coronaviruses (CoVs) are a large family of enveloped single positive-stranded RNA viruses that can infect both mammalian and avian species because their rapid mutation and recombination facilitate their adaptation to new hosts (Graham and Baric, 2010; Li, 2013). They can cause severe, often fatal Acute Respiratory Distress Syndrome (ARDS). CoVs are classified into *Alpha*-, *Beta*-, *Gamma*-, and *Deltacoronavirus* genetic genera. The novel betacoronavirus (betaCoVs) SARS-CoV-2 is relatively close to other betaCoVs: severe acute respiratory syndrome coronavirus (SARS-CoV), Middle East respiratory syndrome coronavirus (MERS-CoV), bat coronavirus HKU4, mouse hepatitis coronavirus (MHV), bovine coronavirus (BCoV), and human OC43 coronavirus (HCoV-OC43). SARS-CoV emerged in China (2002–2003) and spread to other countries (more than 8,000 infection cases and a fatality rate of ~10%) (Peiris et al., 2003). In 2012, MERS-CoV was detected in the Middle East. It spread to multiple countries, infecting more than 1,700 people with a fatality rate of ~36% (de Wit et al., 2016).

The surface-located SARS-CoV-2 spike glycoprotein S (S) is a 1273 amino acid residues. It is a homotrimeric, multidomain, and integral membrane protein that give coronaviruses the appearance of having crowns (*Corona* in Latin) (Li, 2016). It is a key piece of viral host recognition (receptor-recognition) and organ tropism and induces strongly the host immune reaction (Li, 2015). It is subdivided to S1 subunit that binds to a receptor on the host cell surface and S2 subunit that permits viral and host membranes fusion. S1 subunit is divided into two domains, an N-terminal domain (NTD) and a C-terminal receptor-binding domain (RBD) that can function as viral receptors-binding (Li, 2012). In addition, S1 subunit is normally more variable in sequence among different CoVs than is the S2 subunit (Masters, 2006).

Protein phosphatase 2A (PP2A) is a major family of Serine/Threonine phosphatases in eukaryotic cells and regulates diverse biological processes through dephosphorylation of numerous signaling molecules. PPA2 and phosphatase 1 (PP1), regulates over 90% of all Ser/Thr dephosphorylation

events in eukaryotic cells (Eichhorn et al., 2009). PP2A is a heterotrimeric holoenzyme composed of a stable heterodimer of the scaffold A-subunit (PP2A-A) and catalytic C-subunit (PP2A-C) and a variable mutually exclusive regulatory subunit from four families (B (B55), B' (B56), B'' and B''') which provide substrate specificity. The human B56 family consists of at least five different members (α , β , γ , δ and ϵ). Phosphatases and kinases are big players in the regulation of pro-inflammatory responses during microbial infections. Sun et al. (2017) showed that PP2A plays an important role in regulating inflammation by controlling the production of inflammatory cytokines/chemokines (Kozicky and Sly, 2015). In addition, PP2A is one of the phosphatases involved in negatively regulating the inflammatory response (Shanley et al., 2001). Moreover, studies have revealed that viruses use multiple strategies to target PP2A in the aim to manipulate host antiviral responses (Guergnon et al., 2011).

Artepillin C (ArtC) is uniquely found in Brazilian honeybee green propolis and is one of its major bioactive components (Marcucci et al., 2001; Park et al., 2004). It is a low-molecular weight phenolic single ring with two prenyl groups (3,5-diprenyl-4-hydroxycinnamic acid) (Szliszka et al., 2013). These properties suggest high oral bioavailability and cell-permeability allowing good biological activity of ArtC (Shimizu et al., 2004; Konishi et al. 2005; Konishi, 2005). Indeed, Paulino et al. (2008) showed that ArtC exhibited bioavailability by oral administration in mice. Interestingly, ArtC has many therapeutic effects, anti-microbial, anti-tumor, apoptosis-inductor, immunomodulatory, and anti-oxidant effects (Salomão et al., 2004; Kimoto et al., 2001; Orsolic et al., 2006; Matsuno et al., 1997; Gekker et al., 2005; Nakanishi et al., 2003). Many of these therapeutic effects can be attributed to its immunomodulatory functions (Chan et al., 2013; Cheung et al., 2011; Paulino, et al., 2008). Indeed, Szliszka et al., (2013) have tested the anti-inflammatory activity of ArtC in activated RAW264.7 macrophages. They found that ArtC exerted strong antioxidant activity and significantly inhibited the production of several pro-inflammatory cytokines, such as TNF- α , IL-1 β and IL-12, which makes ArtC an excellent anti-inflammatory drug. In addition, ArtC suppresses T cell proliferation and activation (Chan et al., 2013). Here, S protein was analyzed because of its importance in mediating infection. This analysis revealed two B56-binding LxxIx β E-like motifs in S1 and S2 subunits that could recruit the host PP2A. Interestingly, side chains of LxxLxE motif present similarity with a small molecule called Artepillin C (ArtC), a main compound in Brazilian honeybee green propolis. Moreover, ArtC has

anti-inflammatory activity that makes it an excellent candidate for future clinical trials in COVID-19 patients.

RESULTS AND DISCUSSION

Two LxxIxE-like motifs in S1 and S2 subunits of Spike protein

Sequence analysis of SARS-CoV-2 spike protein by the eukaryotic linear motif (ELM) resource (<http://elm.eu.org/>) revealed short linear motifs (SLiMs) known as LxxIxE-like motif, ²⁹³LDPLSE²⁹⁸ in S1 subunit and ¹¹⁹⁷LIDLQE¹²⁰² in S2 subunit (Fig. 1). SLiMs are few amino acid residues (3-15) in proteins that facilitate protein sequence modifications and protein-protein interactions (Davey et al., 2012; Van Roey et al., 2014). RNA viruses are known to mutate quickly and thus are able to create mimic motifs, on very short time scales, that could hijack biological processes in the host cell such as cell signaling networks (Davey et al., 2015; Via et al., 2015; Davey et al., 2011). Interestingly, ²⁹³LDPLSE²⁹⁸ is only present in SARS-CoV-2 and absent in S protein of coronaviruses analysed in this study (Fig. 1A). In order to interact with protein(s), ²⁹³LDPLSE²⁹⁸ must be present at the surface of S1 subunit. Indeed, this motif is exposed in the surface of S1 subunit in the end of NTD (Fig. 3B). A second motif ¹¹⁹⁷LIDLQE¹²⁰² is present in S2 subunit and conserved in SARS-CoV-2, SARS-CoV, SARS-like of bat from China and Kenya (Fig. 1B). These last betacoronaviruses are phylogenetically close (Fig. 2). Unfortunately, the region containing ¹¹⁹⁷LIDLQE¹²⁰² peptide has not been resolved in all known 3D structures of S protein to know if it is exposed in the surface. Probably, ¹¹⁹⁷LIDLQE¹²⁰² peptide is in an intrinsic disordered region of S protein.

Artepillin C, anti-inflammatory compound, mimics LxxLxE motif in S protein

In order to find small molecules with substituents that topologically and structurally resemble key amino acid side chains in LxxLxE motif, the method developed by Baran et al. (2007) has been used. This method allowed to discover a small molecule called Artepillin (ArtC) (Fig. 4A). It is a low-molecular weight phenolic single ring with two prenyl groups (3,5-diprenyl-4-hydroxycinnamic acid) (Szliszka et al., 2013). Figure 4B shows that the side chains of the two leucine and glutamic acid that constitute the key amino acid side chains are superimposed with the two prenyl groups and acid group of ArtC, respectively. ArtC is uniquely found in Brazilian

honeybee green propolis and is one of its major bioactive components (Marcucci et al., 2001; Park et al., 2004). In addition, it has many therapeutic effects, anti-microbial, anti-tumor, apoptosis-inductor, immunomodulatory, and anti-oxidant effects (Salomão et al., 2004; Kimoto et al., 2001; Orsolic et al., 2006; Matsuno et al., 1997; Gekker et al., 2005; Nakanishi et al., 2003). Many of these therapeutic effects can be attributed to its immunomodulatory functions (Chan et al., 2013; Cheung et al., 2011; Paulino, et al., 2008).

Interactions of ²⁹³LDPLSE²⁹⁸ and ArtC with B56 regulatory subunit

In order to determine the molecular interactions of ²⁹³LDPLSE²⁹⁸ and ArtC with B56 regulatory subunit of PP2A (PP2A-B56), molecular docking was performed with the software AutoDock vina (Trott and Olson, 2010). Figure 4C and D show that ²⁹³LDPLSE²⁹⁸ peptide is localized in the same region as pS-RepoMan peptide that contains the LxxIxE motif (PDBid: 5SW9_B) and important amino acids of LxxIxE-like motif are superimposed with those of pS-RepoMan peptide (Fig. 4D). That is confirmed the reliability of AutoDock vina peptide docking module. In addition, Leu293 of ²⁹³LDPLSE²⁹⁸ is docked into hydrophobic pocket and Glu298 form ionic interactions with amino acid residues in positive charged region of PP2A-B56 (Fig. 4C). Note that ²⁹³LDPLSE²⁹⁸ contains a serine that could be phosphorylated generating a negative charge that will interact with positive patch in B56 subunit, enhancing binding affinity (Nygren and Scott, 2015). In the case of ArtC, its two prenyl groups are docked into two pockets as already seen with ²⁹³LDPLSE²⁹⁸ peptide (Fig. 4E-F). Note that the side chain of carboxylic acid group of ArtC is short to form ionic interactions with amino acid residues in positive charged region of PP2A-B56 (Fig. 4E). Therefore, to enhance binding affinity of ArtC, it is necessary to increase the length of side chain of carboxylic acid group. Interestingly, by using PinaColada, a computational method (Zaidman and Wolfson, 2016) for the design and affinity improvement of peptides that preclude protein-protein interaction, the peptide predicted (²⁹³LIDLEE²⁹⁸) by the software is mutated in C-terminal to negative charge residue (glutamic acid), showing the importance of the negative charge of peptide to interact with PP2A-B56. According to Autodock software, predicted binding affinity of ²⁹³LDPLSE²⁹⁸ is -4.9 Kcal/mol, and this of ArtC is -6.1 Kcal/mol. It is known that the binding affinity of SLiMs is relatively weak (low μmolar range) (Gouw et al., 2018). This suggests that ArtC could compete with the virus to bind to PP2A-B56. To my knowledge, no compound has been identified to interact with regulatory subunits of PP2A, in addition it is the first time that a small molecule has been found that mimics

the LxxIxE motif. Despite of numerous studies on ArtC its target is not yet known. In this study, its target is predicted as B56-PP2A. In general, activation of PP2A appears to have a suppressive effect on the inflammatory response (Sun et al., 2017). This suggests according to anti-inflammatory effect of ArtC that it could activate, *in vivo*, B56-PP2A.

Protein phosphatase 2A and single RNA viruses

It has been shown in single RNA viruses, Ebola virus (EBOV) and Dengue fever virus (DENV) that they recruit the host PP2A through its regulatory subunit B56-binding LxxIxE motif to activate transcription and replication (Kruse et al., 2018; Oliveira et al., 2018). In addition, it has been shown an exacerbation of lung inflammation in mice infected with rhinovirus 1B (the most common viral infectious agent in humans). Administrating Salmeterol (beta-agonist) treatment to mice exerts anti-inflammatory effects by interacting with catalytic subunit PP2A, thus increasing its activity. It is probable that beta-agonists have the potential to target distinct pro-inflammatory pathways unresponsive to corticosteroids in patients with rhinovirus-induced exacerbations (Hatchwell et al., 2014). Treatment with Salmeterol drug may merit investigation for the possibility of using it in COVID-19's patients with sustained and dangerous inflammatory reaction.

Summary and conclusion

Analysis of SARS-CoV-2 proteome in the search of SLiMs that could be used by SARS-CoV-2 to manipulate host, allowed to discover in S protein an LxxIxE-like motif that is known to recruit the host PP2A-B56 phosphatase. Interestingly, PP2A is involved in the regulation of pro-inflammatory responses during pathogen infections. Well, recent researches have indicated that SARS-CoV-2 is involved in sustained host inflammation. Therefore, by controlling acute inflammation, it is possible to eliminate its dangerous effects on the host. LxxIxE motif of CoV-2 allowed to find a small molecule called Artepillin (ArtC), a main compound in Brazilian honeybee green propolis, which is known to have anti-inflammatory activity. ArtC, by its non-cytotoxicity in cells, high oral bioavailability, tested in mice, and cell-permeability, in addition that it can be synthesized in the laboratory (Uto et al., 2002; Yashiro et al., 2015) and produced in yeast by using synthetic biology (Munakata et al., 2019) makes it an ideal molecule for future clinical trials in COVID-19 patients.

MATERIALS AND METHODS

Sequence analysis

To search probable short linear motifs (SLiMs), SARS-CoV-2 spike protein sequence was scanned with the eukaryotic linear motif (ELM) resource (<http://elm.eu.org/>). In the aim to find small molecules containing amino acids substituents that mimic LxxIxE-like motif, method described by Baran et al. (2007) was used.

3D modeling and molecular docking

3D structure of Artepillin C (ArtC) was obtained from PubChem database: <https://pubchem.ncbi.nlm.nih.gov/compound/5472440#section=3D-Conformer>.

For docking, the coordinates of the ²⁹³LDPLSE²⁹⁸ peptide were extracted from spike S protein of CoV-2 structure (PDBid: 6VSB_A). Unfortunately, the region containing ¹¹⁹⁷LIDLQE¹²⁰² peptide has not been resolved in all known 3D structures of spike S protein. So, Pep-Fold (Thevenet et al., 2012) software was used to model *de novo* this peptide. The model quality of the peptide was assessed by analysis of a Ramachandran plot through PROCHECK (Vaguine et al., 1999).

The docking of the two peptides into B56 regulatory subunit of PPA2 (PDBid: 5SWF_A) was performed with the software AutoDock vina (Trott and Olson, 2010). The 3D complex containing B56 subunit and peptides was refined by using FlexPepDock (London et al., 2011), which allows full flexibility to the peptide and side-chain flexibility to the receptor. The electrostatic potential surface of the B56 subunit was realized with PyMOL software (<http://pymol.org/>).

PinaColada a computational method (Zaidman and Wolfson, 2016) for inhibitory peptide design was used to improve affinity of ²⁹³LDPLSE²⁹⁸ peptide to bind to B56 regulatory subunit of PPA2. The software mutates several times the input peptide in the aim to find the highest binding affinity.

Phylogeny

To establish the phylogenetic relationships between spike S protein of SARS-CoV-2 and representative betacoronaviruses, amino acid residues sequences were aligned with Clustal omega (Sievers et al., 2011) and a phylogenetic tree was constructed with MrBayes (Huelsenbeck and Ronquist, 2001) using: Likelihood model (Number of substitution types: 6(GTR); Substitution

model: Poisson; Rates variation across sites: Invariable + gamma); Markov Chain Monte Carlo parameters (Number of generations: 100 000; Sample a tree every: 1000 generations) and Discard first 500 trees sampled (burnin).

ACKNOWLEDGMENTS

I would like to thank the IBIS bioinformatics group for their help. I am grateful to Dr. Ahmad Abdel-Mawgoud SALEH, Université Laval, for the revision of the manuscript.

CONFLICT OF INTERESTED

The author declares that he has no conflicts of interest.

REFERENCES

- Baran, I., Varekova, R. S., Parthasarathi, L., Suchomel, S., Casey, F., & Shields, D. C. (2007). Identification of potential small molecule peptidomimetics similar to motifs in proteins. *J Chem Inf Model*, 47(2), 464-474. doi:10.1021/ci600404q
- Chan, G. C., Cheung, K. W., & Sze, D. M. (2013). The immunomodulatory and anticancer properties of propolis. *Clin Rev Allergy Immunol*, 44(3), 262-273. doi:10.1007/s12016-012-8322-2
- Cheung, K. W., Sze, D. M., Chan, W. K., Deng, R. X., Tu, W., & Chan, G. C. (2011). Brazilian green propolis and its constituent, Artepillin C inhibits allogeneic activated human CD4 T cells expansion and activation. *J Ethnopharmacol*, 138(2), 463-471. doi:10.1016/j.jep.2011.09.031
- Davey, N. E., Cyert, M. S., & Moses, A. M. (2015). Short linear motifs - ex nihilo evolution of protein regulation. *Cell Commun. Signal.*, 13, 43. doi:10.1186/s12964-015-0120-z
- Davey, N. E., Travé, G., & Gibson, T. J. (2011). How viruses hijack cell regulation. *Trends Biochem. Sci.*, 36(3), 159-169. doi:10.1016/j.tibs.2010.10.002
- Davey, N. E., Van Roey, K., Weatheritt, R. J., Toedt, G., Uyar, B., Altenberg, B., . . . Gibson, T. J. (2012). Attributes of short linear motifs. *Mol. Biosyst.*, 8(1), 268-281. doi:10.1039/c1mb05231d
- de Wit, E., van Doremalen, N., Falzarano, D., & Munster, V. J. (2016). SARS and MERS: recent insights into emerging coronaviruses. *Nat. Rev. Microbiol.*, 14(8), 523-534. doi:10.1038/nrmicro.2016.81
- Eichhorn, P. J. A., Creighton, M. P., & Bernards, R. (2009). Protein phosphatase 2A regulatory subunits and cancer. *Biochim. Biophys. Acta*, 1795(1), 1-15. doi:10.1016/j.bbcan.2008.05.005

255 Gekker, G., Hu, S., Spivak, M., Lokensgard, J. R., & Peterson, P. K. (2005). Anti-HIV-1 activity
256 of propolis in CD4(+) lymphocyte and microglial cell cultures. *J Ethnopharmacol*, 102(2), 158-
257 163. doi:10.1016/j.jep.2005.05.045

258 Gouw, M., Michael, S., Sámano-Sánchez, H., Kumar, M., Zeke, A., Lang, B., . . . Gibson, T. J.
259 (2018). The eukaryotic linear motif resource - 2018 update. *Nucleic Acids Res.*, 46(D1), D428-
260 D434. doi:10.1093/nar/gkx1077

261 Graham, R. L., & Baric, R. S. (2010). Recombination, reservoirs, and the modular spike:
262 mechanisms of coronavirus cross-species transmission. *J. Virol.*, 84(7), 3134-3146.
263 doi:10.1128/JVI.01394-09

264 Guernon, J., Godet, A. N., Galioot, A., Falanga, P. B., Colle, J.-H., Cayla, X., & Garcia, A. (2011).
265 PP2A targeting by viral proteins: a widespread biological strategy from DNA/RNA tumor viruses
266 to HIV-1. *Biochim. Biophys. Acta*, 1812(11), 1498-1507. doi:10.1016/j.bbadis.2011.07.001

267 Hatchwell, L., Girkin, J., Dun, M. D., Morten, M., Verrills, N., Toop, H. D., . . . Mattes, J. (2014).
268 Salmeterol attenuates chemotactic responses in rhinovirus-induced exacerbation of allergic airways
269 disease by modulating protein phosphatase 2A. *J. Allergy Clin. Immunol.*, 133(6), 1720-1727.
270 doi:10.1016/j.jaci.2013.11.014

271 Huelsenbeck, J. P., & Ronquist, F. (2001). MRBAYES: Bayesian inference of phylogenetic
272 trees. *Bioinformatics*, 17(8), 754-755. doi:10.1093/bioinformatics/17.8.754

273 Kimoto, T., Aga, M., Hino, K., Koya-Miyata, S., Yamamoto, Y., Micallef, M. J., . . . Kurimoto,
274 M. (2001). Apoptosis of human leukemia cells induced by Artepillin C, an active ingredient of
275 Brazilian propolis. *Anticancer Res*, 21(1A), 221-228.

276 Konishi, Y. (2005). Transepithelial transport of artepillin C in intestinal Caco-2 cell
277 monolayers. *Biochim Biophys Acta*, 1713(2), 138-144. doi:10.1016/j.bbamem.2005.05.011

278 Konishi, Y., Hitomi, Y., Yoshida, M., & Yoshioka, E. (2005a). Absorption and bioavailability of
279 artepillin C in rats after oral administration. *J Agric Food Chem*, 53(26), 9928-9933.
280 doi:10.1021/jf051962y

281 Kozicky, L. K., & Sly, L. M. (2015). Phosphatase regulation of macrophage activation. *Semin*
282 *Immunol*, 27(4), 276-285. doi:10.1016/j.smim.2015.07.001

283 Kruse, T., Biedenkopf, N., Hertz, E. P. T., Dietzel, E., Stalman, G., López-Méndez, B., . . . Becker,
284 S. (2018). The Ebola Virus Nucleoprotein Recruits the Host PP2A-B56 Phosphatase to Activate
285 Transcriptional Support Activity of VP30. *Mol. Cell*, 69(1), 136-145.e136.
286 doi:10.1016/j.molcel.2017.11.034

287 Li, F. (2012). Evidence for a common evolutionary origin of coronavirus spike protein receptor-
288 binding subunits. *J. Virol.*, 86(5), 2856-2858. doi:10.1128/JVI.06882-11

289 Li, F. (2013). Receptor recognition and cross-species infections of SARS coronavirus. *Antiviral*
290 *Res.*, 100(1), 246-254. doi:10.1016/j.antiviral.2013.08.014

291 Li, F. (2015). Receptor recognition mechanisms of coronaviruses: a decade of structural studies. *J.*
292 *Virol.*, 89(4), 1954-1964. doi:10.1128/JVI.02615-14

293 Li, F. (2016). Structure, Function, and Evolution of Coronavirus Spike Proteins. *Annu Rev Virol*,
294 3(1), 237-261. doi:10.1146/annurev-virology-110615-042301

295 London, N., Raveh, B., Cohen, E., Fathi, G., & Schueler-Furman, O. (2011). Rosetta FlexPepDock
296 web server--high resolution modeling of peptide-protein interactions. *Nucleic Acids Res.*, 39(Web
297 Server issue), W249-253. doi:10.1093/nar/gkr431

298 Marcucci, M. C., Ferreres, F., García-Viguera, C., Bankova, V. S., De Castro, S. L., Dantas, A. P.,
299 . . . Paulino, N. (2001). Phenolic compounds from Brazilian propolis with pharmacological
300 activities. *J Ethnopharmacol*, 74(2), 105-112. doi:10.1016/s0378-8741(00)00326-3

301 Masters, P. S. (2006). The molecular biology of coronaviruses. *Adv. Virus Res.*, 66, 193-292.
302 doi:10.1016/S0065-3527(06)66005-3

303 Matsuno, T., Jung, S. K., Matsumoto, Y., Saito, M., & Morikawa, J. (1997). Preferential
304 cytotoxicity to tumor cells of 3,5-diprenyl-4-hydroxycinnamic acid (artepillin C) isolated from
305 propolis. *Anticancer Res*, 17(5A), 3565-3568.

306 Munakata, R., Takemura, T., Tatsumi, K., Moriyoshi, E., Yanagihara, K., Sugiyama, A., . . .
307 Yazaki, K. (2019). Isolation of. *Commun Biol*, 2, 384. doi:10.1038/s42003-019-0630-0

308 Nakanishi, I., Uto, Y., Ohkubo, K., Miyazaki, K., Yakumaru, H., Urano, S., . . . Ikota, N. (2003).
309 Efficient radical scavenging ability of artemillin C, a major component of Brazilian propolis, and
310 the mechanism. *Org Biomol Chem*, 1(9), 1452-1454. doi:10.1039/b302098c

311 Nygren, P. J., & Scott, J. D. (2015). Therapeutic strategies for anchored kinases and phosphatases:
312 exploiting short linear motifs and intrinsic disorder. *Front. Pharmacol.*, 6, 158.
313 doi:10.3389/fphar.2015.00158

314 Oliveira, M., Lert-Itthiporn, W., Cavadas, B., Fernandes, V., Chuansumrit, A., Anunciação, O., . .
315 . Sakuntabhai, A. (2018). Joint ancestry and association test indicate two distinct pathogenic
316 pathways involved in classical dengue fever and dengue shock syndrome. *PLoS Negl. Trop. Dis.*,
317 12(2), e0006202. doi:10.1371/journal.pntd.0006202

318 Orsolić, N., Saranović, A. B., & Basić, I. (2006). Direct and indirect mechanism(s) of antitumour
319 activity of propolis and its polyphenolic compounds. *Planta Med*, 72(1), 20-27. doi:10.1055/s-
320 2005-873167

321 Park, Y. K., Paredes-Guzman, J. F., Aguiar, C. L., Alencar, S. M., & Fujiwara, F. Y. (2004).
322 Chemical constituents in *Baccharis dracunculifolia* as the main botanical origin of southeastern
323 Brazilian propolis. *J Agric Food Chem*, 52(5), 1100-1103. doi:10.1021/jf021060m

324 Paulino, N., Abreu, S. R., Uto, Y., Koyama, D., Nagasawa, H., Hori, H., . . . Bretz, W. A. (2008).
325 Anti-inflammatory effects of a bioavailable compound, Artemillin C, in Brazilian propolis. *Eur J*
326 *Pharmacol*, 587(1-3), 296-301. doi:10.1016/j.ejphar.2008.02.067

327 Peiris, J. S. M., Lai, S. T., Poon, L. L. M., Guan, Y., Yam, L. Y. C., Lim, W., . . . group, S. s.
328 (2003). Coronavirus as a possible cause of severe acute respiratory syndrome. *Lancet*, 361(9366),
329 1319-1325. doi:10.1016/s0140-6736(03)13077-2

330 Salomão, K., Dantas, A. P., Borba, C. M., Campos, L. C., Machado, D. G., Aquino Neto, F. R., &
331 de Castro, S. L. (2004). Chemical composition and microbicidal activity of extracts from Brazilian
332 and Bulgarian propolis. *Lett Appl Microbiol*, 38(2), 87-92. doi:10.1111/j.1472-765x.2003.01458.x

333 Shanley, T. P., Vasi, N., Denenberg, A., & Wong, H. R. (2001a). The serine/threonine phosphatase,
334 PP2A: endogenous regulator of inflammatory cell signaling. *J Immunol*, 166(2), 966-972.
335 doi:10.4049/jimmunol.166.2.966

336 Shimizu, K., Ashida, H., Matsuura, Y., & Kanazawa, K. (2004). Antioxidative bioavailability of
337 artepillin C in Brazilian propolis. *Arch Biochem Biophys*, 424(2), 181-188.
338 doi:10.1016/j.abb.2004.02.021

339 Sievers, F., Wilm, A., Dineen, D., Gibson, T. J., Karplus, K., Li, W., . . . Higgins, D. G. (2011).
340 Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal
341 Omega. *Mol. Syst. Biol.*, 7, 539. doi:10.1038/msb.2011.75

342 Sun, L., Pham, T. T., Cornell, T. T., McDonough, K. L., McHugh, W. M., Blatt, N. B., . . . Shanley,
343 T. P. (2017a). Myeloid-Specific Gene Deletion of Protein Phosphatase 2A Magnifies MyD88- and
344 TRIF-Dependent Inflammation following Endotoxin Challenge. *J Immunol*, 198(1), 404-416.
345 doi:10.4049/jimmunol.1600221

346 Szliszka, E., Mertas, A., Czuba, Z. P., & Król, W. (2013). Inhibition of Inflammatory Response by
347 Artepillin C in Activated RAW264.7 Macrophages. *Evid Based Complement Alternat Med*, 2013,
348 735176. doi:10.1155/2013/735176

349 Thévenet, P., Shen, Y., Maupetit, J., Guyon, F., Derreumaux, P., & Tufféry, P. (2012). PEP-FOLD:
350 an updated de novo structure prediction server for both linear and disulfide bonded cyclic
351 peptides. *Nucleic Acids Res.*, 40(Web Server issue), W288-293. doi:10.1093/nar/gks419

352 Trott, O., & Olson, A. J. (2010). AutoDock Vina: improving the speed and accuracy of docking
353 with a new scoring function, efficient optimization, and multithreading. *J. Comput. Chem.*, 31(2),
354 455-461. doi:10.1002/jcc.21334

355 Uto, Y., Hirata, A., Fujita, T., Takubo, S., Nagasawa, H., & Hori, H. (2002). First total synthesis
356 of artepillin C established by o,o'-diprenylation of p-halophenols in water. *J Org Chem*, 67(7),
357 2355-2357. doi:10.1021/jo0056904

358 Vaguine, A. A., Richelle, J., & Wodak, S. J. (1999). SFCHECK: a unified set of procedures for
359 evaluating the quality of macromolecular structure-factor data and their agreement with the atomic
360 model. *Acta Crystallogr. D Biol. Crystallogr.*, 55(Pt 1), 191-205.
361 doi:10.1107/S09074444998006684

- Van Roey, K., Uyar, B., Weatheritt, R. J., Dinkel, H., Seiler, M., Budd, A., . . . Davey, N. E. (2014). Short linear motifs: ubiquitous and functionally diverse protein interaction modules directing cell regulation. *Chem. Rev.*, 114(13), 6733-6778. doi:10.1021/cr400585q
- Via, A., Uyar, B., Brun, C., & Zanzoni, A. (2015). How pathogens use linear motifs to perturb host cell networks. *Trends Biochem. Sci.*, 40(1), 36-48. doi:10.1016/j.tibs.2014.11.001
- Yashiro, K., Hanaya, K., Shoji, M., & Sugai, T. (2015). New synthesis of artepillin C, a prenylated phenol, utilizing lipase-catalyzed regioselective deacetylation as the key step. *Biosci Biotechnol Biochem*, 79(12), 1926-1930. doi:10.1080/09168451.2015.1058704
- Zaidman, D., & Wolfson, H. J. (2016). PinaColada: peptide-inhibitor ant colony ad-hoc design algorithm. *Bioinformatics*, 32(15), 2289-2296. doi:10.1093/bioinformatics/btw133

FIGURES LEGEND

Figure 1. Multiple alignment of the spike glycoprotein of betacoronaviruses using Clustal omega. ²⁹³LDPLSE²⁹⁸ (A) and ¹¹⁹⁷LIDLQE¹²⁰² (B) motifs are indicated by green stars. GenBank and UniProt accession numbers are indicated at the start of each sequence. The figure was prepared with ESPript (<http://esprict.ibcp.fr>).

Figure 2. Unrooted phylogenetic tree of spike protein of representative betacoronaviruses. The tree was constructed using Mr Bayes method based on the multiple sequence alignment by Clustal omega. Red rectangle assembles betacoronaviruses with the same ¹¹⁹⁷LIDLQE¹²⁰². Green star indicated the only betacoronavirus with ²⁹³LDPLSE²⁹⁸. GenBank and UniProt accession numbers are indicated at the start of each sequence.

Figure 3. Spike (S) protein of SARS-CoV-2. (A) Diagram representation of S protein colored by domain. N-terminal domain (NTD), receptor-binding domain (RBD), subdomains 1 and 2 (SD1-2, orange), S1/S2 protease cleavage site, Fusion peptide (FP), heptad repeat 1 and 2 (HR1 and HR2), central helix (CH), connector domain (CD), transmembrane domain (TM), cytoplasmic tail (CT), and the localization of ²⁹³LDPLSE²⁹⁸ in the end of NTD and ¹¹⁹⁷LIDLQE¹²⁰² peptide in HR2. (B) Surface structure representation of the S1 subunit (PDBid: 6VSB_A). ²⁹³LDPLSE²⁹⁸ peptide is localized in the surface S1 subunit (red).

Figure 4. Stick representation of (A) Artepillin C (ArtC) and (B) superposition of ArtC (yellow) and ²⁹³LDPLSE²⁹⁸ peptide (green). Electrostatic potential surface representation of the region of the B56 regulatory subunit of PP2A (PDBid: 5SWF_A) with docked (C) ²⁹³LDPLSE²⁹⁸ peptide (green), (D) ²⁹³LDPLSE²⁹⁸ superimposed to pS-RepoMan (orange) (⁵⁸¹RDIASKKPLLPSPIPELPEVPE⁶⁰¹) peptide (PDBid: 5SW9_B), (E) ArtC (yellow) and (F) ArtC superimposed to ²⁹³LDPLSE²⁹⁸ peptide (green). The surfaces are colored by electrostatic potential with negative charge shown in red and positive charge in blue. Images were generated using PyMol (www.pymol.org).

Fig. 3

Fig. 4