Short title: Zn transport-related genes in biofortified alfalfa

Article title:

Transcriptional regulation of genes involved in Zn transport after foliar Zn application to *Medicago sativa*

Alessio Cardini^{a*}, Elisa Pellegrino^{a+*}, Philip J. White^b, Barbara Mazzolai^c, Marco C. Mascherpa^d,
Laura Ercoli^a

^aInstitute of Life Sciences, Scuola Superiore Sant'Anna, Piazza Martiri della Libertà 33, 56127 Pisa, Italy

^bDepartment of Ecological Science, The James Hutton Institute, Invergowrie, Dundee DD2 5DA, United Kingdom

^cCenter for Micro-BioRobotics, Istituto Italiano di Tecnologia, Viale Rinaldo Piaggio 34, 56025 Pontedera, Pisa, Italy

^dNational Research Council of Italy, C.N.R., Istituto di Chimica dei Composti OrganoMetallici - SS of Pisa, Italy

⁺Corresponding author: *Elisa Pellegrino*

Email: elisa.pellegrino@santannapisa.it

^{*}These authors contributed equally to the work

One-sentence summary:

Upregulation of *ZIP2*, *NAS1* and *HMA4* and downregulation of *ZIP3* are associated with Zn sequestration and shoot-to-root translocation in *Medicago sativa* following foliar Zn biofortification

ORCID

Alessio Cardini <https://orcid.org/0000-0001-9687-6047>

Elisa Pellegrino <https://orcid.org/0000-0002-2482-1327>

Philip J White <https://orcid.org/0000-0003-0827-288X>

30 Barbara Mazzolai <https://orcid.org/0000-0003-0722-8350>

31 Marco Carlo Mascherpa <https://orcid.org/0000-0001-5816-0953>

32 Laura Ercoli <https://orcid.org/0000-0002-7942-2120>

33

34 **Author contributions**

35 EP and LE conceived the ideas; AC, EP, LE designed the methodology; AC and BM designed and

36 performed real-time PCR; AC and CM performed chemical analyses; AC and EP performed data

37 analysis; AC, EP, PJW, LE led the writing of the manuscript; all authors contributed critically to the

38 drafts and gave final approval for publication.

39

ABSTRACT

Zinc (Zn) is an essential micronutrient for both plants and animals, and Zn deficiency is one of the most widespread problems for agricultural production. Although many studies have been performed on the biofortification of staple crops with Zn, few studies have focused on forage crops. In this study the molecular mechanisms of Zn transport-related in *Medicago sativa* L. were investigated following foliar Zn applications aimed at increasing the accumulation of Zn in edible tissues. Zinc uptake and redistribution between shoot and root were determined following the application of six Zn doses to leaves (0, 0.01, 0.1, 0.5, 1, 10 mg Zn plant⁻¹). Twelve putative genes encoding proteins involved in Zn transport (*MsZIP₁₋₇*, *MsZIF1*, *MsMTP1*, *MsYSL1*, *MsHMA4* and *MsNAS1*) were identified and the changes in their expression following foliar Zn application were quantified using newly designed RT-qPCR assays. Shoot and root Zn concentration was increased following foliar Zn applications ≥ 0.1 mg plant⁻¹. Increased expression of *MsZIP2*, *MsHMA4* and *MsNAS1* in shoots, and of *MsZIP2* and *MsHMA4* in roots, was observed with the largest Zn dose. By contrast, *MsZIP3* was downregulated in shoots at Zn doses ≥ 0.1 mg plant⁻¹. Three functional modules were identified in the *M. sativa* response to foliar Zn application: genes involved in Zn uptake by cells, genes involved in vacuolar Zn sequestration and genes involved in Zn redistribution within the plant. These results will inform genetic engineering strategies aimed at increasing the efficiency of crop Zn biofortification.

KEY WORDS:

Zinc, micronutrient, biofortification, alfalfa (*Medicago sativa*), ZIP transporters, Nicotianamine, Metal Tolerance Protein (MTP), Yellow Stripe-Like Protein (YSL), Zinc Induced facilitators (ZIF), Heavy Metal transporters (HMA), cellular zinc homeostasis.

INTRODUCTION

Food quality is a key factor for human health (Geissler and Powers, 2017). For their well-being, humans require sufficient quantities of at least 18 mineral elements, which have specific physiological roles and are irreplaceable in the diet (White, 2016). Eight essential macronutrients (i.e., N, P, S, Ca, Mg, K, Na, Cl) are required in large amounts in the diet ($> 100 \text{ mg day}^{-1}$), and 10 micronutrients are required in smaller amount (e.g., Zn, Fe, F, Mn). The main sources of these elements in the human diet include edible crops, animal products (e.g., meat, fish, eggs) and dairy products (e.g., milk, cheese, butter) as well as mineral supplements (Keen, 1990; Prasad, 2013; White, 2016). A large proportion of the world's population suffers from Zn related diseases (i.e., malabsorption syndrome, liver disease, chronic renal disease, sickle cell disease and other chronic diseases), since it relies on cereal-based diets with low Zn content due to poor soil Zn availability (WHO, 2005; Alloway, 2009; Prasad, 2013; Kumssa et al., 2015; Cakmak et al., 2017). Diversification of the human diet and biofortification of edible crops are therefore needed to alleviate Zn deficiency in humans. Similarly, increasing Zn concentrations in forage crops are important for maintaining livestock health and the quality of food products, which affect human health indirectly (McDonald et al., 2002; Ciccolini et al., 2017; Capstaff and Miller, 2018; Huma et al., 2019).

Zinc plays a major role as a co-factor of over 300 enzymes in plants and is an essential micronutrient (Broadley et al., 2007). Zinc is involved in various physiological functions, such as CO_2 fixation, protein synthesis, free radical capture, regulation of growth and development, and disease resistance (Sasaki et al., 1998; Broadley et al., 2007). Many structural motifs in transcriptional regulatory proteins are stabilized by Zn, such as Zn finger domains (Albert et al., 1998). Zinc deficiency reduces crop production, as does Zn excess (White and Pongrac, 2017). Excessive Zn^{2+} can compete with other cations in binding to enzymes and for transport across membranes, thereby impairing cellular activities (White and Pongrac, 2017). Thus, the uptake of Zn^{2+} by cells and its transport within the plant must be strictly regulated. Plant cells have evolved

several homeostatic mechanisms for avoiding Zn²⁺ toxicity when exposed to large Zn availability in their environment. These include the reduction of Zn influx to cells, the stimulation of Zn efflux from the cytosol, the sequestration of Zn in vacuoles, and the chelation of Zn by Zn binding ligands. In general, the concentration of Zn in plant tissues must be kept between 15 to 300 µg Zn g⁻¹ dry matter (DM) to maintain cell structure and function (Broadley et al., 2012; White and Pongrac, 2017). Although tolerance to large tissue Zn concentrations varies among species (Alloway, 2008; White and Pongrac, 2017), Zn concentrations above 400-500 µg g⁻¹ DM often cause toxicity symptoms including impaired root and shoot growth, chlorosis and necrosis of leaves, reduced photosynthesis, nutrient imbalance and ultimately loss of yield (Chaney, 1993; Broadley et al., 2007; Di Baccio et al., 2009; White and Pongrac, 2017).

The process of producing crops with greater mineral concentrations in edible tissues is called biofortification and provides a solution to the problem of mineral deficiencies in human and animal nutrition (White and Broadley, 2005). There are various approaches to Zn biofortification of edible crops, including agronomic strategies and conventional or transgenic breeding strategies. Agronomic biofortification aims to increase Zn concentrations in edible tissues through the application of Zn-fertilisers to the soil or to leaves. It is relatively inexpensive and efficient (Saltzman et al., 2013). Foliar application of Zn is generally more effective than the application of Zn fertilisers to soil, since Zn uptake by plant roots is often limited by the low solubility of Zn salts, its binding to organic substrates, and its immobilization in the microbial biomass (Gregory et al., 2017). Both agronomic and genetic biofortification strategies have been studied extensively in cereal staple crops, such as rice, wheat and maize, but less in legumes, such as beans, peas or lentils (White and Broadley, 2005, 2011; Rawat et al., 2013). An international program, the HarvestPlus Zinc Fertilizer Project, is exploring the potential of Zn fertilisers to enhance the yields and Zn concentrations in edible portions of staple crops in developing countries of Africa, Asia and South America (www.harvestzinc.org) (Cakmak, 2012), but this program does not include forage crops.

The natural direction of Zn flux in plants is from the soil via roots to the shoot and seeds (White

and Broadley, 2009). Various transport proteins and ligands that are responsible for Zn^{2+} uptake by roots and its transport and sequestration within the plant have been characterized (Olsen and Palmgreen, 2014; Caldelas and Weiss, 2017; White and Pongrac, 2017). Among these, ZRT-IRT-like Proteins (ZIPs), have been studied in several plants, including *Arabidopsis thaliana*, soybean (*Glycine max*), barley (*Hordeum vulgare*), barrel medic (*Medicago truncatula*) and rice (*Oryza sativa*) (Grotz et al., 1998; Zhao and Eide, 1996; López-Millán et al., 2004; Milner et al., 2013; Tiong et al., 2015). These proteins not only transport Zn^{2+} across membranes, but can also transport other transition metal cations, including Cd^{2+} , $\text{Fe}^{3+}/\text{Fe}^{2+}$, Mn^{2+} , Ni^{2+} , Co^{2+} and Cu^{2+} (Grotz et al., 1998; Mäser et al., 2001; Eckhardt et al., 2001). Generally, the expression of ZIP genes is upregulated when plants become Zn deficient (Ramesh et al., 2003; Ishimaru et al., 2006; Eide et al., 1996), facilitating Zn influx to cells and movement of Zn between organs, and also when plants become Fe or Mn deficient (Buglio et al., 2002; Vert et al., 2002; Ishimaru et al., 2006; Pedas et al., 2008). Other proteins that transport Zn include the Metal Tolerance Proteins (MTPs), which function as cation/proton antiporters and are thought to transport Zn into vacuoles (Kolaj-Robin et al., 2015) and the Yellow Stripe-Like Proteins (YSLs), which transport the Zn-Nicotianamine complex (NA-Zn) and load Zn into the xylem and phloem (Curie et al., 2009). The Zinc Induced Facilitators (ZIFs) and the Heavy Metal transporters (HMAs) are implicated in Zn influx to vacuoles and to the xylem, respectively (Olsen and Palmgren, 2014). Zinc is chelated by organic molecules, such as the carboxylic acid, citric acid, and nicotianamine (NA) in plants (Sinclair and Krämer, 2012). Nicotianamine is a non-proteinogenic amino acid with a high affinity for Fe, Cu and Zn, and is involved in their homeostasis (Deinlein et al., 2012). Nicotianamine mediates the intercellular and interorgan movement of Zn and was found to enable Zn hyperaccumulation in *Arabidopsis halleri* and *Noccaea caerulescens* (Deinlein et al., 2012; Foroughi et al., 2014). In general the functions of these transporters have been studied by expressing them in yeast, but to understand how the various Zn transport proteins and chelates act together to maintain appropriate cytosolic and tissue Zn concentrations it is important to study the responses of an intact plant to

fluctuations in Zn supply.

In this study the transcriptional responses of genes encoding Zn transport-related processes facilitating Zn uptake by cells, vacuolar sequestration and redistribution within the plant were studied following foliar Zn application to the most productive and widely cultivated forage legume, alfalfa (*Medicago sativa* L.). The study was designed to provide information on the molecular responses to Zn biofortification of forage crops (Foyer et al., 2016; Capstaff and Miller, 2018). The following hypotheses were tested: i) foliar application of Zn increases shoot and root Zn concentrations, which results in changes in the expression of genes involved in Zn transport-related processes to detoxify excess Zn; ii) genes encoding Zn transport-related processes are organized in functional modules, that act in a concerted manner to redistribute Zn within the plant to maintain non-toxic cytosolic and tissue Zn concentrations. Genes encoding putative Zn transport-related processes were identified in alfalfa through phylogenetic comparisons and their likely roles are discussed. Changes in the expression of these genes following foliar Zn application were determined and the possible effects of these on the redistribution of Zn within cells and between tissues are discussed. The knowledge gained from this study could help to optimize Zn biofortification strategies when using foliar Zn fertilisers and to provide strategies for breeding forage crops to addresses Zn deficiencies in livestock.

RESULTS

Shoot and root Zn concentrations

The application of Zn to leaves did not modify shoot or root biomass and all *M. sativa* plants had root nodules (data not shown). However, Zn concentrations in both shoots and roots were strongly affected by foliar Zn application ($F_{(5,17)}=32.61$, $P<0.001$; $F_{(5, 17)}=28.53$, $P<0.001$; respectively) (Fig. 1). A foliar Zn application of 0.01 mg Zn plant⁻¹ produced a shoot Zn concentration similar to that of the control (no-Zn addition), but shoot Zn concentrations were

increased progressively by larger doses ($0.1 < 0.5/1 < 10$ mg Zn plant⁻¹), from more than threefold to 35-fold more than that of the control (Fig. 1). Foliar applications of 0.01, 0.1 and 0.5 mg Zn plant⁻¹ did not produce root Zn concentrations greater than that of the control treatment, but foliar doses of 1 and 10 mg Zn plant⁻¹ increased root Zn concentrations to threefold and 11-fold more than the control treatment, respectively. Shoot and root Zn contents were also strongly affected by foliar Zn application ($F_{(5,17)}=53.73$, $P<0.001$; $F_{(5, 17)}=32.45$, $P<0.001$; respectively) and their responses to increasing foliar Zn applications followed the corresponding Zn concentrations (Supplemental Fig. S1).

Phylogenetic analysis

Phylogenetic analysis of the coding sequences of the *ZIP* genes revealed several distinct clades (Supplemental Fig S2). One clade contained sequences for *MsZIP2* and *MsZIP7*, which were similar to each other. In addition, the sequence of *MsZIP2* was closely related to those of *MtZIP2* and *GmZIP1-ZIP2* and the sequence of *MsZIP7* was closely related to those of *MtZIP7* and *AtZIP11*. Another clade contained the sequences of *MsZIP1*, *MsZIP3*, *MsZIP5* and *MsZIP6*. The sequence of *MsZIP1* clustered with that of *MtZIP1*. Sequences of *MsZIP3* and *MsZIP5* were similar to each other and clustered with the corresponding sequences for *M. truncatula* genes (Supplemental Fig. S2). Sequences for *MsZIP1*, *MsZIP3* and *MsZIP5* were closely related to each other, whereas that of *MsZIP6* formed a separate cluster with the sequences of *MtZIP6* and *AtZIP12*. The sequence of *MsZIP4* was distant from the sequences of other *M. sativa* *ZIPs* and formed a cluster with the sequences of *MtZIP4* and *AtZIP4*.

Phylogenetic analyses of the coding sequences of the other genes related to Zn transport processes revealed that they were all similar to their *M. truncatula* counterparts. As regards *ZIF*, the sequence of *MsZIF1* clustered with the sequences of *MtZIF1* and *GmZIF1* (Supplemental Fig. S3a). As regards *MTP*, the sequence of *MsMTP1* formed a cluster with *MtMTP1* and *GmMTP1* and was also related to *AtMTP1* and *AtMTPA1* (Supplemental Fig. S3b). Similarly, the sequence of *MsYSL1*

was most similar to those of *MtYSL1* and *GmYSL1* (Supplemental Fig. S3c) and the sequence of *MsHMA4* was most similar to those of *MtHMA4* and *GmHMA4* (Supplemental Fig. S3d). Finally, the sequence of *MsNAS1* was closely related to those of *MtNAS* and *GmNAS* (Supplemental Fig. S3e).

Gene expression analysis

The expression of *MsZIP3* was significantly downregulated at foliar doses of 0.1, 1 and 10 mg Zn plant⁻¹ ($F_{(3, 11)} = 28.46$, $P < 0.01$) (Fig. 2). By contrast, the expression of *MsZIP2* was significantly upregulated at the largest dose of 10 mg Zn plant⁻¹ ($F_{(3, 11)} = 5.59$, $P < 0.05$). The expression of *MtZIP1*, *MtZIP5* and *MtZIP6* in shoots was not significantly affected by foliar Zn application, although a general trend towards downregulation with increasing foliar Zn doses was observed. The expression of *MsZIP4* and *MsZIP7* in shoots was unaffected by foliar Zn application. The expression of *MsZIP2* was significantly upregulated in roots at the largest foliar dose of 10 mg Zn plant⁻¹ ($F_{(3, 11)} = 9.26$, $P < 0.01$) (Fig. 2). In roots, *ZIP* genes were not significantly affected by foliar Zn application, although a general trend of *MtZIP1*, *MtZIP3*, *MtZIP5* and *MtZIP7* towards upregulation with increasing foliar Zn doses was observed. Of the other genes related to Zn transport processes, the expression of *MsHMA4* was significantly upregulated in both shoots ($F_{(3, 11)} = 115.29$, $P < 0.01$) and roots ($F_{(3, 11)} = 14.23$, $P < 0.01$) following the application of 1 and 10 mg Zn plant⁻¹ (shoots: +63% and +424%, respectively; roots: +86% and +66%, respectively; Fig. 3). In shoots, the expression of *MsHMA4* was about 3-fold higher following a dose of 10 mg Zn plant⁻¹ than following a dose of 1 mg Zn plant⁻¹, whereas the expression of *MsHMA4* in roots was similar when 1 or 10 mg Zn plant⁻¹ was applied. The expression of *MsNAS1* was also significantly upregulated ($F_{(3, 11)} = 6.46$, $P < 0.05$) at the largest foliar Zn dose (10 mg plant⁻¹), whereas its expression in roots was unaltered following foliar Zn application (Fig. 3). In shoots, *MsYSL1* and *MsZIF1* were not significantly affected by foliar Zn application, although there was a trend towards upregulation of the expression with increasing Zn doses, while the expression of *MsMTP1* remained unaltered

following the application of Zn (Fig. 3). Finally, in roots, *MsMTP1* and *MsZIF1* were not significantly affected by foliar Zn application, although there was a trend towards upregulation of the expression with increasing Zn doses, whereas the expression of *MsYSL1* remained unchanged (Fig. 3).

Using correlation analysis to reveal functional modules of genes whose expression is co-regulated in plants, three functional modules for ZIP gene co-expression were observed (Fig. 4a; $r > 0.6$). In the first functional module, the expression of *MsZIP1*, *MsZIP3*, *MsZIP5* and *MsZIP6* in shoots were all strongly correlated. Although the expression of *MsZIP7* in shoots was also placed in this module, its expression was poorly correlated with the expression of other ZIP genes in either shoots or roots. The second functional module comprised correlations between the expression of *MsZIP4* in shoots and roots, and the expression of *MsZIP1*, *MsZIP3*, *MsZIP4*, *MsZIP5* and *MsZIP6* in roots. The third functional module comprised correlations in the expression of *MsZIP2* in shoots and roots and *MsZIP7* in roots. With respect to the expression of the other genes involved in Zn transport related processes, there was a strong divergence in their expression patterns in shoots and roots (Fig. 4b). In the root, the expression of *MsNAS1*, *MsYSL1*, *MsZIF1* and *MsMTP1* were strongly correlated (Fig. 4b). The expression of *MsYSL1*, *MsZIF1* and *MsMTP1* in roots showed good correlations with the expression of *MsMTP1* and *MsZIF1* in shoots. In the shoot the expression of *MsYSL1*, *MsMTP1* and *MsZIF1* were all strongly correlated. Also in shoots, the expression of *MsNAS1* and *MsHMA4* were well correlated and showed a good correlation with the expression of *MsMTP1* and *MsZIF1* in shoots. Interestingly, the expression of *MsNAS1* strongly correlated with that of *MsYSL1* in shoots.

PERMANOVA showed that the expression of ZIP genes was significantly affected by foliar Zn application dose and differed between shoots and roots, which explained 29% and 23% of the total variance, respectively (Table 1). The expression of other genes related to Zn transport processes that were studied (*MsZIF1*, *MsNAS1*, *MsHMA4*, *MsYSL1* and *MsMTP1*) were also affected by foliar Zn application dose and the organ examined. Zinc application dose explained 17% of the total variance,

while plant organ explained 19%. PERMANOVA on all studied genes highlighted a significant effect of Zn application dose, plant organ and their interaction on gene expression, explaining 68% of the total variance.

DISCUSSION

Plant Zn nutritional status after foliar Zn application

The critical leaf concentration for Zn deficiency approximates 15–20 $\mu\text{g Zn g}^{-1}$ dry weight and the critical leaf concentration for Zn toxicity approximates 400–500 $\mu\text{g Zn g}^{-1}$ (Broadley et al., 2012; White and Pongrac, 2017). Before foliar Zn application, the alfalfa plants used in the experiments reported here were probably Zn deficient, since their shoot Zn concentrations were below the critical leaf concentration for Zn deficiency (Fig. 1). After the application of the lowest foliar Zn dose (0.01 mg plant^{-1}) plants probably remained Zn deficient (7.6 $\mu\text{g Zn g}^{-1}$ dry weight), but all other foliar Zn doses increased Zn concentrations in shoots above the critical concentration for Zn deficiency (Fig. 1). Plants treated with 0.1 mg Zn plant^{-1} probably had an optimal Zn status for plant growth, whereas plants treated with 0.5 and 1 mg Zn plant^{-1} had shoot Zn concentrations close to the toxicity threshold. When a foliar dose of 10 mg Zn plant^{-1} was applied, shoot Zn concentrations greatly exceeding the threshold for Zn toxicity (Fig. 1). Plants often exhibit characteristic visual symptoms of Zn deficiency and Zn toxicity when these occur (Broadley et al., 2012; White and Pongrac, 2017), but five days after foliar Zn application no visual symptoms of Zn deficiency or toxicity, nor differences in plant biomass, were observed among plants receiving contrasting foliar Zn doses (data not shown). Foliar Zn doses larger than 0.1 mg Zn plant^{-1} resulted in incremental increases in the Zn concentration of roots (Fig. 1), despite Zn having limited mobility in the phloem (White and Broadley, 2011; White, 2012). This observation suggests that roots can act as a sink for Zn applied to leaves, thereby mitigating excessive Zn accumulation in shoot tissues. In previous work, foliar application of Zn was shown to increase Zn concentration in

phloem-fed tissues, such as fruits, seed, and tubers (Cakmak, 2004, 2008; Cakmak et al., 2010; White et al., 2017). The shoot to root Zn concentration ratio shifted from values below one in conditions of Zn deficiency (0.4) to values greater than one in Zn-replete or Zn-intoxicated plants (1.3 - 3.2) (Fig. 1). When the plants are Zn deficient the recirculation of Zn between organs via the xylem and phloem is required to meet minimal growth demands and the application of foliar Zn to Zn deficient plants must be effectively redistributed within the plant (Erenoglou et al., 2011; Sinclair and Kramer, 2012), whereas when excessive foliar Zn is applied, Zn must be chelated in the cytoplasm, sequestered in the vacuole and redistributed via the phloem or xylem to other organs to avoid toxicity (White and Pongrac, 2017).

Foliar Zn application alters the expression of genes involved in Zn transport-related processes

Despite several genes encoding Zn transporters having been identified in plants, and the encoded proteins characterized, the mechanisms of Zn uptake and transport in alfalfa are still largely unknown. However, the recently sequenced alfalfa genome has allowed the discovery of genes involved in Zn uptake and distribution within this species (O'Rourke et al., 2015). In the present study, 12 putative genes encoding proteins likely to be involved in processes related to Zn transport in the plant (*MsZIP1-7*, *MsZIP1*, *MsMTP1*, *MsYSL1*, *MsHMA4* and *MsNAS1*) were identified and their expression in shoots and roots quantified following foliar Zn applications (Fig. 2; Fig. 3). The expression of genes encoding proteins involved in Zn uptake by cells, vacuolar sequestration and redistribution within the plant responded differently to increasing foliar Zn dose. The expression of *MsZIP2*, *MsHMA4* and *MsNAS1* in shoots was increased by increasing foliar Zn dose, while only the expression of *MsZIP2* and *MsHMA4* in roots were upregulated by increasing foliar Zn dose (Fig. 2; Fig. 3). By contrast, *MsZIP3* was downregulated in shoots when foliar Zn doses ≥ 0.1 mg Zn plant⁻¹ were applied (Fig. 2). These changes in gene expression might produce a reduction in Zn uptake capacity (but not necessarily a reduction in actual Zn uptake since this will also be related to apoplastic Zn concentration) by cells in both the shoot and root (by reducing Zn

influx through *MsZIP3* and increasing Zn efflux through *MsHMA4*), chelation of Zn using Zn-NA in the shoot for Zn-detoxification and phloem transport, and greater recirculation of Zn within the plant in both the phloem and xylem (by increasing NA concentrations and *MsZIP2* and *MsHMA4* activities), and are, therefore, consistent with the observation that increasing foliar Zn dose increases both shoot and root Zn concentrations.

Regulation of genes encoding ZIP transporters

The influx and efflux of Zn across the plasma membrane of plant cells must be tightly controlled to allow optimal cell functioning and hence to ensure normal plant growth and development (Sinclair and Krämer, 2012). The expression of only two of the seven *ZIP* genes studied, *MsZIP2* and *MsZIP3*, showed statistically significant responses to foliar Zn application (Fig. 2). The expression of *MsZIP2* was significantly upregulated in both shoots and roots in response to the largest dose of foliar Zn applied (10 mg Zn plant⁻¹). It is likely that this dose is toxic to both shoot and root cells. The relative induction in the expression of *MsZIP2* was greatest in roots. The phylogenetic analysis of *ZIP* transporters revealed that *MsZIP2* is closely related to *MtZIP2* and *AtZIP2* (Supplemental Fig. S6). Thus, *MsZIP2* is probably located in the plasma membrane performing similar functions to *MtZIP2* and *AtZIP2*. Burleigh et al. (2003) reported that *M. truncatula* plants grown with adequate soil Zn availability expressed *MtZIP2* in roots and stems, but not in leaves. The expression of *MtZIP2* in roots increased with increasing Zn fertiliser applications to soil, with the greatest expression being found at toxic Zn doses (Burleigh et al., 2003). Similarly, Milner et al. (2013) found that the expression of *AtZIP2* was ~10-fold higher in roots than shoots in Zn-replete *Arabidopsis thaliana* plants and that Zn deficiency reduced the expression of *AtZIP2* in both roots and shoots. The localization of *ZIP2* at the plasma membrane was observed in both *M. truncatula* (Burleigh et al., 2003) and *A. thaliana* (Milner et al., 2013). The expression of *AtZIP2* was localized to the stele of the root (Milner et al., 2013), supporting a role of *AtZIP2* in long distance transport of Zn between roots and shoots. It is possible that the increased

expression of *MsZIP2* observed in our study when plants experience Zn toxicity might be a detoxification strategy, either through storing excess Zn in xylem parenchyma cells or recirculating Zn in the xylem.

The expression of *MsZIP3* was significantly downregulated in shoots following the foliar application of Zn (Fig. 2). The ZIP3 transporter is thought to mediate Zn influx to the cell from the apoplast (Sinclair and Kramer, 2012). Therefore, the downregulation of *MsZIP3* in shoots of plants receiving more Zn is consistent with the ability of plant cells to control their Zn uptake to effect cytoplasmic Zn homeostasis. Reduced expression of *MsZIP3* in plants with a greater Zn supply is also in agreement with previous studies of *M. truncatula* and *A. thaliana* (Grotz et al., 1998; López-Millán et al., 2004). However, although *AtZIP3* could restore growth to a Zn-uptake defective yeast (Milner et al., 2013), *MtZIP3* was not found to be able to restore the growth of a Zn-uptake defective yeast in Zn-limited media, although it did restore the growth of a Fe-uptake defective yeast in Fe-limited media (López-Millán et al., 2004). Thus, the *MsZIP3* transporter could have a higher affinity for Fe than Zn. In *O. sativa* ZIP3 gene is expressed in the xylem parenchyma and transfer cells and might be responsible for unloading transition metal cations from the xylem to the parenchyma in plants receiving an excessive Zn supply (Sasaki et al., 2015). The role of OsZIP3 in unloading Zn from the vascular tissues, suggests that the reduced expression of *MsZIP3* in shoots of *M. sativa* receiving an excessive foliar Zn dose might be a detoxification strategy to reduce Zn uptake by shoot cells.

The observation that foliar Zn applications had no effect on the expression of ZIP genes, except *MsZIP2* and *MsZIP3* (Fig. 2), might be explained by the roles of ZIP proteins in the transport of other transition metals. For example, evidence of Cu and Mn transport by ZIP4 were provided through yeast complementation studies (Wintz et al., 2003; López-Millán et al., 2004). Moreover, applying the same technique, a role of ZIP6 was highlighted in the transport of Fe by López-Millán et al. (2004), whereas Wintz et al. (2003) did not find any involvement of ZIP6 in the transport of Cu, Zn or Fe. Although the changes in the expression of *MsZIP1*, *MsZIP5* and *MsZIP6* following

foliar Zn application were not statistically significant, changes in their expression in shoots were positively correlated with changes in the expression of *MsZIP3*, showing a general trend for them to be downregulated following foliar Zn application and suggesting that these four ZIPs might act as a functional module in the shoot (Fig. 4). By contrast, the expression of *MsZIP1*, *MsZIP3*, *MSZIP4*, and *MsZIP5* were positively correlated in roots, suggesting that these genes behave as a functional module in roots (Fig. 4).

Regulation of genes encoding other Zn transport-related processes

The expression of *MsHMA4*, which is implicated in Zn redistribution within the plant (Hussain et al., 2004; Sinclair et al., 2018), was increased in both shoots and roots of plants whose shoot Zn concentration suggested they were close to, or experiencing, Zn toxicity (Fig. 3). The significant upregulation of *MsHMA4* following foliar application of ≥ 1 mg Zn plant⁻¹ might be related to the removal of excess Zn from both shoots and roots. This interpretation is consistent with the role of HMA4 in *A. thaliana* and in the metal hyperaccumulators *Arabidopsis halleri* and *Noccea caerulescens* (Baker and Whiting, 2002; Hussain et al., 2004; Hanikenne et al., 2008; Ò Lochlainn et al., 2011; White and Pongrac, 2017), in which greater expression of *HMA4* results in greater Zn flux to the xylem and Zn translocation to transpiring leaves. However, the phylogenetic similarity of *MsHMA4* to *MtHMA4* and, particularly, to *AtHMA5* (Supplemental Fig. S3d) suggest a role in Cu transport (Andrés-Colás et al., 2006; Sankaran et al., 2009; Hermand et al., 2014). This implication of the latter observation is unclear.

Since Zn²⁺ concentrations are low in the alkaline phloem sap, the transport of most Zn in the phloem is as Zn ligand complexes, such as zinc-nicotianamine (NA-Zn) (Deshpande et al., 2018). Nicotianamine is the main Zn chelate in phloem transport and is also important for Zn sequestration in vacuoles (Deinlein et al., 2012), and tolerance of excessive Zn uptake (Aarts et al., 2014). Nicotianamine concentrations generally correlate with those of *NAS* transcripts, and for this reason *NAS* expression can be used as a proxy for NA content (Talke et al., 2006; Haydon et al., 2012).

Accordingly, in the work reported here the increased expression of *MsNAS1* in shoots following the application of ≥ 1 mg Zn plant⁻¹ (Fig. 3) probably reflects the role of NA in Zn detoxification through its sequestration within vacuoles and its redistribution from shoot to root after excessive foliar Zn applications. This observation is consistent with Deshpande et al. (2018), who found that the expression of *NAS2* in the durum wheat (*Triticum durum* Desf.) increased following foliar Zn application and reports that *NAS* expression is constitutively high in plants that hyperaccumulate Zn (Becher et al., 2004; Weber et al., 2004; Haydon et al., 2012; White and Pongrac, 2017).

Homologs of *MsMTP1* and *MsZIF1* were previously found to encode transporters loading Zn and NA into the vacuoles of *Thlaspi geosingense* and *A. thaliana* cells, respectively (Gustin et al., 2009; Haydon et al., 2012). Unexpectedly, the expression of these genes was unaffected by foliar Zn application (Fig. 3). This observation suggests that the proteins encoded by these genes might not contribute to Zn detoxification in *M. sativa*. Nevertheless, only *MsZIF1* of all the genes studied here showed a trend towards increased expression in roots with increasing foliar Zn dose (Fig. 3), which might indicate a role in detoxification of excess Zn in roots through its sequestration with NA in the vacuole. The high correlation between the expression of *MsMTP1* and *MsZIF1* in both shoots and roots suggests that they might constitute a functional module and act synergistically to sequester Zn in the vacuole (Fig. 4), as supported by other studies (Gustin et al., 2009; Haydon et al., 2012; Sharma et al., 2016).

In *A. thaliana*, *AtYSL1* has a role in the long-distance transport of the NA-Zn complex and in loading Zn into seeds (Jean et al., 2005; Curie et al., 2009). For this reason, an increase in the expression of *AtYSL1* was expected to occur in parallel with the increased expression of *MsNAS1* in shoots. However, the expression of *MsYSL1* did not show any significant change in shoots or roots in response to foliar Zn application, although there was a trend towards greater *MsYSL1* expression in shoots with increasing foliar Zn doses (Fig. 3). In addition, the high correlation in the expression of *MsNAS1* and *MsYSL1* in shoots in response to foliar Zn applications (Fig. 4) supports the expectation that these genes are components of a functional module affecting the long-distance

transport of Zn in the plant, as it was previously highlighted in *A. thaliana* by Pita-Barbosa et al. (2019).

The responses of gene expression to foliar Zn applications suggest three functional modules that effect cytoplasmic Zn homeostasis through Zn transport related processes in *M. sativa*: genes involved in Zn influx to cells (shoots: *MsZIP1*, *MsZIP5*, and *MsZIP6*; roots: *MsZIP1*, *MSZIP4*, *MsZIP5*, and *MsZIP6*), genes involved in Zn sequestration in the vacuole (shoots and roots: *MsMTP1* and *MsZIF1*) and genes involved in Zn redistribution within the plant (shoots and roots: *MsHMA4* and *MsYSL1*).

In conclusion, this is the first study to characterise the expression of genes related to Zn transport processes following foliar Zn application to a forage legume and provides new molecular insights to the responses of Zn transport related processes to foliar Zn applications. A significant increase in the expression of *MsZIP2* as foliar Zn doses increase suggests the detoxification of excess Zn through the accumulation of Zn in xylem parenchyma cells. A decrease in the expression of *MsZIP3* as foliar Zn doses increase suggests a reduction in the Zn influx capacity of shoot cells to reduce Zn uptake. An increase in the expression of *MsHMA4* in roots and shoots as foliar Zn doses increase suggests an increase in the transport of Zn in the xylem when plants are subject to Zn toxicity, while an increase in the expression of *MsNAS1* in the shoot suggests the chelation of excess Zn in the shoot, enabling Zn sequestration in vacuoles or the redistribution of Zn to roots via the phloem. The elucidation of three functional modules of genes involved in (a) Zn influx to cells, (b) sequestration of Zn in the vacuole and (c) redistribution of Zn within the plant are fundamental to understanding the molecular mechanisms of cytoplasmic Zn homeostasis and might inform the selection of appropriate genotypes enabling greater Zn accumulation in edible portions or increased tolerance of Zn in the environment.

MATERIALS AND METHODS

Plant growth and experimental design

Surface sterilized seeds of alfalfa (*M. sativa* L.) were germinated on moist sterilized silica sand (1-4 mm size) in a climatic chamber at 24/21 °C day/night temperature, 16/18 h light/dark cycle and 200 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$. After two weeks of growth, three seedlings were transplanted to 1500 mL volume pots, filled with sterilized silica sand (number of pots 18) and *Sinorhizobium meliloti* was supplied as a filtrate to all plants to ensure that the plants produced nodules in all treatments. A Hoagland nutrient solution lacking Zn (Li et al., 2013) was used to fertilize the plants, with 10 mL solution being applied every week. After two months of growth, plants were treated with one of six doses of Zn (0, 0.01, 0.1, 0.5, 1 and 10 mg Zn plant⁻¹) (three replicates per dose). Six ZnSO₄·7H₂O solutions of 0, 0.05, 0.5, 2.5, 5, 50 g Zn L⁻¹ were prepared to supply these doses. A drop of Tween 20 detergent was added to the six solutions to break the surface tension of the leaves and enhance Zn uptake. Zinc was applied to the middle leaf laminae of the three plants in each pot as twenty 10 μL -droplets. The experiment was arranged in a fully randomized design, with three replicates for each Zn dose. The shoots and roots of the plants were harvested separately five days after Zn application. At harvest, 1 mM CaCl₂ solution and water were used to remove any residual Zn from the leaf surface (Yilmaz et al., 2017). Shoot and root fresh weight was measured, whereas shoot and root dry weight was determined on subsamples after oven drying at 70°C to constant weight.

Measurement of zinc concentrations

Approximately 100 mg of shoot or root dry biomass were carefully weighed and mineralized in a microwave medium pressure digester (Milestone Start D, FKV Srl, Torre Boldone, Italy) with 7 mL of 69% HNO₃ and 2 mL of 30% H₂O₂ (ultrapure grade). Zinc concentration in the resulting solutions was determined by inductively coupled plasma optical emission spectroscopy (ICP-OES) using an Optima 8000 spectrometer (Perkin Elmer, Waltham, MA, USA), following the procedure of Nölte (2003).

Gene selection and design and validation of new RT-qPCR assay

Seven genes encoding putative ZRT-IRT-like proteins (ZIP) were selected for investigation (i.e., *ZIP1-7*) (Fig. 5). The selection was based on information gathered by Burleigh et al. (2003) and López-Millán et al. (2004) on the expression of genes encoding Zn transporters in the model legume *Medicago truncatula* and on the structure of the neighbor joining (NJ) tree built using available ZIP sequences of several plant species. Five more genes, whose products are involved in Zn transport-related processes (Olsen and Palmgren, 2014), were also chosen for investigation based on information gathered by other authors and on sequence similarity with other plant species. The *NAS1* gene encoding nicotianamine synthase (NAS) was chosen because this enzyme synthesizes nicotianamine (NA), which is involved in long-distance Zn transport (Curie et al., 2009) (Fig. 5). The *HMA4* gene, which encodes a transmembrane P-type ATPase heavy metal transporter, was chosen because this transporter loads Zn into the xylem in roots for its transport to shoots (Palmer and Gueriot, 2009) (Fig. 5). The *MTP1* gene, which encodes a transporter of the CDF family, was selected because this transporter is implicated in the sequestration of excess Zn in the vacuole (Desbrosses-Fonrouge et al., 2005; Gustin et al., 2009) (Fig. 5). The *ZIF1* gene, which encodes the Zn-induced facilitator 1 transporter, was chosen because it transports NA into the vacuole to chelate vacuolar Zn (Haydon and Cobbett, 2007; Haydon et al., 2012) (Fig. 5). The *YSL1* gene, which encodes a transporter of Zn-NA complexes, was chosen because it is implicated in Zn loading and transport of Zn in the phloem (Palmer and Gueriot, 2009) (Fig. 5). To standardize the expression of genes encoding Zn transport-related processes, two reference genes were selected: actin (*ACT*) and elongation factor 1- α (*EF1- α*) (Nicot et al., 2005).

Using the draft genome sequence of alfalfa in the Alfalfa Gene Index and Expression Atlas Database (AEGD) (O'Rourke et al., 2015; <http://plantgrn.noble.org/AGED/index.jsp>), homologous gene sequences of *M. sativa* were retrieved by BLASTn similarity searches using the gene sequences of *M. truncatula*. The chosen genes for *M. sativa* were named *MsZIP1-7* for the seven ZIP genes and *MsNAS1*, *MsHMA4*, *MsZIF1*, *MsYSL1*, *MsMTP1* for the other selected genes. The

two reference genes were named *MsACT-101* and *MsEF1- α* . The gene sequences and their annotations have been deposited in NCBI under the Submission # 2338923.

Forward and reverse new PCR primers for the 12 Zn transport-related genes and the two reference genes suitable for SYBR[®] Green II RT-qPCR assays (Biorad, USA) were designed (Table 2). The Primer-BLAST online tool in the National Center for Biotechnology Information (NCBI; <https://www.ncbi.nlm.nih.gov/tools/primer-blast/>) was used to design primers. The newly designed RT-qPCR assays are suitable for both *M. sativa* and *M. truncatula*. The length of the fragment, the Sanger sequences of the PCR amplicons (Table 2) and the single melting temperature peaks confirmed the specificity of the new RT-qPCR assays (Supplemental Fig. S4). Sanger sequencing was performed on PCR amplicons of three cDNA samples (Supplemental Material and Methods S1). Examples of electropherograms of the sequences are reported in Supplemental Fig. S5. The sequences of the obtained PCR amplicons have been deposited in NCBI the Submission # 2338930. Amplification efficiencies (E) in the range of 96.1-111.0% are evidence of accurate quantification, while the coefficients of correlation ($R^2 > 0.998$) indicate high precision of measurements across concentration ranges of at least 3-4 orders of magnitude (Table 2; Supplemental Fig. S6). The concentration ranges over which the relationship between the relative fluorescence and the logarithm of the concentration is linear, and the precision of quantification (standard curves) as reflected in the coefficient of correlation (R^2), were determined using three independent 10-fold serial dilutions of a cDNA sample of *M. sativa*. The accuracy of quantification was determined by the efficiency (E) of each qPCR amplification, using the equation $E = [10^{-1/S} - 1] \times 100$, where S is the slope of the standard curve. The evaluation of the reference genes based on the cycle threshold (Ct) values made us choose the actin gene (*MsACT-101*) for quantifying relative gene expression in the shoots and the elongation factor 1- α (*MsEF1- α*) gene for quantifying relative gene expression in roots (Supplemental Fig. S7a,b). This choice was based on the observations that there was no statistical difference in the expression of the reference genes in tissues following foliar Zn

applications and that *MsACT-101* and *MsEF1-α* showed the smallest overall variation in the shoot and root, respectively (Supplemental Fig. S7c,d).

RNA extraction and gene expression analysis

Total RNA was extracted from 50 mg subsamples of fresh shoot and root tissue, using the RNeasy Mini Kit (Qiagen, Hilden, Germany). The extractions were performed from tissues of plants treated with the foliar Zn doses that produced a significant increase in Zn concentration in shoots (0.1, 1 and 10 mg Zn plant⁻¹) and the control plants to which no foliar Zn had been applied (24 RNA extractions). Any DNA in the RNA extracts was removed by a DNase treatment (Promega, USA). The purity of the RNA extracts was verified by spectroscopic light absorbance measurements at 230 nm, 260 nm and 280 nm using the NanoDrop 2000 (Termo Scientific, Warchesre, MA, USA) (Desjardins and Conkin, 2010). The integrity and approximate concentration of the extracted RNA was determined by electrophoresis of the RNA extracts in a 1% agarose gel containing Sybr Safe (Invitrogen, Carlsbad, CA). One microgram of total RNA was reverse transcribed to complementary DNA (cDNA) using the iScript cDNA Synthesis Kit (Biorad, Hercules, California) in a 20 µL reaction volume. The RT-qPCRs for gene expression analysis were run as three technical replicates with a final reaction volume of 20 µL, containing 10 µL of SYBR Green Supermix (Biorad), 5 µL of 100-fold diluted cDNA, and 0.4 µM final concentrations of the gene-specific PCR primers on a CFX Connect Real-Time System thermal cycler (Biorad, Hercules, California). The qPCR conditions were 95°C for 3', followed by 40 cycles of 95° C for 5', and 60° C for 30''. A dissociation curve of each reaction was performed (65° C to 95° C, 0.5° C increment every 5'') to check that PCR amplified only one product. The most suitable reference gene for relative gene expression analysis was determined by comparing the expression levels of the reference genes *MsACT-101* and *MsEF1-α* across all cDNA samples. Relative gene expression was calculated using the double standardisation (ΔΔCq) method that requires a reference gene and a control treatment (Livak and Schmittgen, 2001).

533

534 **Bioinformatic and statistical analyses**

535

536 A BLAST search was performed in the Alfalfa Gene Index and Expression Atlas database using
 537 the *ZIP1-7*, *ZIF1*, *MTP1*, *YSL1*, *HMA4* and *NAS* coding sequences from *M. truncatula*. This
 538 allowed the identification of gene sequences encoding potential metal transporters and chelators in
 539 the whole *M. sativa* genome. The sequences obtained were aligned with the corresponding
 540 sequences from *M. truncatula* and the length of the *M. sativa* genes were determined after removing
 541 the external unaligned nucleotides. The *M. sativa* and *M. truncatula* *ZIP* gene sequences were also
 542 aligned with those of other plant species (*A. thaliana*, *G. max*, *H. vulgare*, *O. sativa*, *Triticum*
 543 *aestivum*, and *Zea mays*) obtained from a search of GenBank. Similarly, the *M. sativa* and *M.*
 544 *truncatula* gene sequences of *ZIF1*, *MTP1*, *YSL1*, *HMA4* and *NAS* were aligned with their
 545 corresponding sequences of other plant species (*A. thaliana*, *G. max*, *H. vulgare*, *O. sativa*, *T.*
 546 *aestivum*, and *Z. mays*) obtained from a search of GenBank. Sequence alignments were performed
 547 using the algorithm ClustalW in MEGA X (Kumar et al., 2018). Phylogenetic comparisons were
 548 performed to infer the putative roles of the selected *M. sativa* Zn transport-related proteins. The
 549 phylogenetic trees were inferred by Neighbor-Joining (NJ) analysis (Saitou et al., 1987) in MEGA
 550 X and the evolutionary distances were calculated using the p-distance method (Nei and Kumar,
 551 2000). Branch support bootstrap values were derived from 500 bootstrap replicates. The
 552 phylograms were drawn by MEGA X and edited using Adobe Illustrator CC 2017.

553 The effect of the application of the foliar Zn on tissue Zn concentration and on the expression
 554 of the selected genes was analysed in shoots and roots separately by one-way analysis of variance
 555 (ANOVA), followed by a Tukey-B test in the case of significance of the response to foliar Zn
 556 application. When required, gene expression data were log-transformed to meet the ANOVA
 557 assumptions. The data displayed graphically are the means and associated standard errors of the
 558 untransformed raw data. All statistical analyses were performed using the software package SPSS

version 21.0 (SPSS Inc., Chicago, IL, USA). Permutational analysis of variance (PERMANOVA; Anderson, 2001) was used to test the effect of foliar Zn application and plant organ (shoot and root) on the expression of the seven *ZIP* genes and of the other five genes encoding Zn transport-related processes separately. In addition, the PERMANOVA was performed on the expression of all the genes together. The response data matrices were standardised by sample, and total and then Euclidean distances were calculated among samples. *P*-values were calculated using the Monte-Carlo test (Anderson and Braak, 2003). Since PERMANOVA is sensitive to differences in multivariate location and dispersion, analysis of homogeneity of multivariate dispersion (PERMDISP; Anderson, 2006) was performed to check the homogeneity of dispersion among groups. The analyses were performed using PRIMER 7 and PERMANOVA+ software (Clarke and Gorley, 2015). Finally, heatmaps were constructed to illustrate correlations in expression among *ZIP*s and among other genes encoding Zn transport-related processes using the R package ggplot2 (Wickham, 2011), using the average linkage clustering of the Pearson correlations calculated from relative gene expression following foliar Zn application.

Supplemental Materials

Supplemental Figure S1. Shoot and root zinc (Zn) content of alfalfa (*Medicago sativa* L.) five days after application of six doses of Zn to leaves.

Supplemental Figure S2. Neighbor-Joining phylogenetic tree of *ZIP* gene sequences of *Medicago sativa*, *Medicago truncatula* and other plant species.

Supplemental Figure S3. Neighbor-Joining phylogenetic trees of sequences of Zinc Induced Facilitator (*ZIF*), Metal Tollerance Protein (*MTP*), Yellow Stripe Like protein (*YSL*), Heavy Metal

585 Transporter (*HMA*) and Nicotianamine Synthase (*NAS*) genes of *Medicago sativa*, *Medicago*
586 *truncatula* and other plant species.

587

588 **Supplemental Figure S4.** Melting curve analysis of the qPCR products obtained by the newly
589 designed pair of primers.

590

591 **Supplemental Figure S5.** Electropherograms of PCR products obtained by the newly designed
592 primers for *MsZIP₁₋₇* transporters and for other Zn related genes.

593

594 **SupplementalFigure S6.** Standard curves for the newly designed qPCR primer pairs.

595

596 **Supplemental Figure S7.** Cycle threshold value of the reference genes, *actin 101* (*MsACT-101*)
597 and *elongation factor 1- α* (*MsEF1- α*) in shoots and roots for the no-Zn addition control and the
598 three Zn doses.

599

600 **Supplemental Material and Methods S1.** PCR amplification conditions.

601

602 **ACKNOWLEDGEMENTS**

603 We acknowledge the Italian Institute of Technology (IIT) for funding for PhD Fellowship of
604 AC under the PhD programme in Agrobiosciences at the Scuola Superiore Sant'Anna of Pisa, Italy.
605 PJW was supported by the Rural and Environment Science and Analytical Services Division
606 (RESAS) of the Scottish Government. We acknowledge Dr. Hannes A. Gamper for the technical
607 support in real-time RT-PCR.

608

609 **Tables**

610

Table 1 Permutation analyses of variance (PERMANOVAs) on the effect of application of three doses of zinc (Zn) (0.1, 1 and 10 mg Zn plant⁻¹) and plant compartment (shoot and root) on the expression of seven *MsZIP* genes and separately on the expression of other five genes (*MsZIF1*, *MsNAS1*, *MsHMA4*, *MsYSL1* and *MsMTP1*) (see Table 1, Fig. 1). A PERMANOVA was also performed on the response of all the genes. The analysis of homogeneity of multivariate dispersion (PERMDISP) was also performed. The studied plant was alfalfa (*Medicago sativa* L.). The analysis included also no-Zn addition control. Gene relative expression was studied on a total of 24 experimental units, corresponding to three replicates per each level of treatment.

Response variables	Explanatory variables	Zn application (Zn)	Plant compartment (Comp)	Zn x Comp	Residual
ZIP genes	Pseudo F	5.56	8.16	1.76	38.3
	P(perm)	0.002	0.001	0.082	
	Explained variance (%)	29.1	22.9	9.7	
	PERMDISP P(perm)	0.005	0.766		
Other genes	Pseudo F	3.06	5.59	1.76	50.55
	P(perm)	0.007	0.015	0.1	
	Explained variance (%)	17.3	19.35	12.78	
	PERMDISP P(perm)	0.412	0.852		
All genes	Pseudo F	4.27	10.49	3.41	31.9
	P(perm)	0.001	0.001	0.003	
	Explained variance (%)	17.3	25.2	25.6	
	PERMDISP P(perm)	0.152	0.030		

Table 2 Gene name, forward and reverse sequences of fourteen newly designed primer pairs for the quantification of the expression of genes of alfalfa (*Medicago sativa*), encoding proteins involved in cellular zinc (Zn) influx and efflux and Zn chelation (see **Fig. 1**). Two reference genes (i.e., *MsACT-101* and *MsEF1-□□□*) were also designed. The length of the amplicons, the primer amplification Efficiency (%) and R² of the standard curve are indicated (see **Fig. S3**). The reference sequences are indicated by the accession number of the *Medicago truncatula* sequences and by the contig number of the *M. sativa* sequences. The primers were designed using the National Center for Biotechnology Information *PRIMER Blast* online tool.

Gene*	Reference sequence (Accession number and contig number)	Forward primer (5'-3')	Reverse primer (5'-3')	Amplicon size (bp)	Efficiency (%)	R ²
<i>MsZIP1</i>	AY339054 [†] / 19855 [‡]	ATGATTAAAGCCT TCGCGGC	TCTGCTGGAACCT GTTTAGAAGG	233	99.8	0.999
<i>MsZIP2</i>	AY007281/8 2450	AGCCCAATTGGCG TAGGAAT	ACAGCAACACCAA AAAGCACA	215	99.3	0.999
<i>MsZIP3</i>	AY339055/3 3860	TGGTGTGATTTTG GCAACCG	TGACGGACCCGAA GAAACAG	325	104.9	0.999
<i>MsZIP4</i>	XM_003603 101/92651	GGAGGGTGCATTT CTCAAGC	AGCAATGCCTGTT CCAATGC	108	97.1	0.999
<i>MsZIP5</i>	XM_013605 712/66451	TGAAGGCATGGG ACTTGGA	CCAGCTGAAGCTG CATTGAA	192	99.3	0.998
<i>MsZIP6</i>	AY339058/9 668	CTTGGCGACACGT TCAATCC	CCACAAGTCCCGA AAAGGGA	188	106.0	0.998
<i>MsZIP7</i>	AY339059/6 2098	GGCTTGTGCTGGT TATTTGAT	TTTCCATGCGTCT GCTTTTGT	310	96.1	0.999
<i>MsZIF1</i>	XM_003601 836/59165	TGCCTGCATTTGG TTACCG	CTGCAGCTTCCAC ATTGTCAG	77	105.9	0.999
<i>MsHMA4</i>	XM_003626 900/19210	TGCTCAACTTGCC AAAGCAC	GGAATGAACCATC CCAGCCA	111	108.9	0.999
<i>MsYSL1</i>	XM_024781 439/4892	CAAGAAGCAAGT GCATGGGT	TCCACAGTCTTCTT TGCCTGAG	94	111.0	0.999
<i>MsMTP1</i>	FJ389717/67 347	TGCAGCATTGCG ATCTCCT	TGCATAGAAACCA AAGCACCA	114	104.5	0.999
<i>MsNAS1</i>	XM_003594 705/61146	GCTAGCTTGGCTG AAGATTGG	AGATACAAAGCAC TCGGAGACA	87	100.5	0.999
<i>MsACT-101</i>	XM_003593 074/89028	TCTCTGTATGCCA GTGGACG	TCTGTATAATCAC GCCCAGCA	140	102.4	0.999
<i>MsEF1-□</i>	XM_003618 727/56897	CCACAGACAAGC CCCTCAG	TCACAACCATAACC GGGCTTC	114	100.2	0.999

*See Fig. 1 for the full names of the genes

[†] NCBI - GenBank Accession number - <https://www.ncbi.nlm.nih.gov/genbank/>

[‡] AGED - The Alfalfa Gene Index and Expression Atlas Database - <http://plantgrn.noble.org/AGED/>

Figure legends

Figure 1. Zinc (Zn) concentrations in shoots and roots of alfalfa (*Medicago sativa*) five days after the application of Zn doses of 0, 0.01, 0.1, 0.5, 1 or 10 mg Zn plant⁻¹ to leaves. Means $\pm$ standard error of three replicates are shown. Differences among the applied Zn doses were tested separately for shoot and root by one-way analysis of variance. Different letters denote significant differences in Zn concentrations in shoots and roots independently, according to Tukey-B honestly test ($P < 0.05$).

Figure 2. Relative expression of seven transmembrane zinc (Zn) transporter genes (*MsZIP1-7*) five days after the application of Zn doses of 0, 0.1, 1 or 10 mg Zn plant⁻¹ to leaves of alfalfa (*Medicago sativa*). Means $\pm$ standard error of three replicates are shown. The expression levels were calculated relative to reference genes (*MsACT-101* for shoot and *MsEF1- α* for root) and to the control (0 mg Zn plant⁻¹). The broken line denotes the threshold between up- and down-regulation relative to the control. Differences in the expressions of each gene after different Zn doses were tested separately for shoot and root by one-way analysis of variance. Different letters denote significant differences among Zn doses, according to Tukey-B test ($P < 0.05$). The full names of the genes are reported in the legend of Figure 1.

Figure 3. Relative expression of genes related to Zn transport processes (*MsZIF1*, *MsHMA4*, *MsYSL1*, *MsMTP1* and *MsNAS1*) five days after the application of Zn doses of 0, 0.1, 1 and 10 mg Zn plant⁻¹ to leaves of alfalfa (*Medicago sativa*). Means $\pm$ standard error of three replicates are shown. The expression levels were calculated relative to reference genes (*MsACT-101* for shoot and *MsEF1- α* for root) and to the control (0 mg Zn plant⁻¹). The broken line denotes the threshold between up- and down-regulation relative to the control. Differences in the expression of each gene at the different Zn doses were tested separately for shoot and root by one-way analysis of variance.

639 Different letters denote significant differences among Zn doses, according to Tukey-B test ($P <$
640 0.05). The full names of the genes are reported in the legend of Figure 1.

641

642 **Figure 4.** Heatmaps reporting the correlations between the differences in expression of genes
643 related to zinc (Zn) transport processes in alfalfa (*Medicago sativa*) after foliar application of Zn
644 doses of 0.1, 1 and 10 mg Zn plant⁻¹ relative to a control dose of 0 mg Zn plant⁻¹. The similarity in
645 the degree of correlation in fold-change of gene expression to Zn application relative to the control
646 is based on the average linkage clustering of the Pearson correlations (r). In the clustering trees the
647 genes are indicated in brown for roots and in green for shoots, while the ranks of correlations of the
648 heatmap are indicated by color intensity (r 0 to 1: from low to strong intensity of green). Seven
649 genes encoding transmembrane Zn transporter (*MsZIP₁₋₇*) (a); four genes encoding cellular Zn
650 transporters (including vacuolar transporters) (*MsZIF1*, *MsHMA4*, *MsYSL1* and *MsMTP1*) and a
651 gene encoding a nicotianamine synthase (*MsNAS1*) (b).

652

653 **Figure 5.** Suggested model for the roles of putative genes encoding proteins involved in Zinc (Zn)
654 transport-related processes in alfalfa (*Medicago sativa*). The sites of action in the plant (i.e., root
655 cytoplasm, rc; root vacuole, rv; xylem and apoplast, X/A; phoelm, P; leaf cytoplasm, lc; leaf
656 vacuole, lv) and the element (E) fluxes (K_{1-13}) are reported. The concentration of the element is
657 indicated in each site [E]. The scheme synthesizes information across studies in various plants. Gene
658 abbreviations: *ZIP*, Zrt-/Irt-like Protein; *NAS*, Nicotianamine synthase; *ZIF*, Zinc-Induced
659 Facilitator; *MTP*, Metal Transporter Protein; *HMA*, P_{1B}-type Heavy Metal ATPase; *YSL*, Yellow
660 Stripe Like Protein; *ZIP?* indicates a generic *ZIP*; free diffusion: diffusion through leaf epidermis;
661 stomata: absorption through stomata. Plant abbreviations: Mt, *Medicago truncatula*; At,
662 *Arabidopsis thaliana*; Os, *Oryza sativa*. References: ^aLópez-Millán et al., 2004; ^bMilner et al., 2013;
663 ^cAarts, 2014; ^dClemens et al., 2013; ^eCurie et al., 2009; ^fHaydon et al., 2012; ^gDesbrosses-Fonrouge
664 et al., 2005; ^hHussain et al., 2004; ⁱPalmer and Guerinot, 2009; ^jBurleigh et al., 2003; ^kSasaki et al.,

665 2015; ¹Fageria et al., 2009.

666

667 LITERATURE CITED

668 **Aarts MG** (2014) Nicotianamine secretion for zinc excess tolerance. *Plant Physiol* **166**: 751-752.

669 **Albert IL, Nadassy K, Wodak SJ** (1998) Analysis of zinc binding sites in protein crystal
670 structures. *Protein Sci* **7**: 1700-1716

671 **Alloway BJ** (2008) Zinc in soils and crop nutrition. 2nd ed. IZA and IFA, Brussels, Belgium and
672 Paris, France

673 **Alloway BJ** (2009) Soil factors associated with zinc deficiency in crops and humans. *Environ*
674 *Geochem Hlth* **31**: 537-548

675 **Anderson MJ** (2001) A new method for non-parametric multivariate analysis of variance. *Austral*
676 *Ecol* **26**: 32-46

677 **Anderson M, Braak CT** (2003) Permutation tests for multi-factorial analysis of variance. *J Stat*
678 *Comput Sim* **73**: 85-113

679 **Anderson MJ, Ellingsen KE, McArdle BH** (2006) Multivariate dispersion as a measure of beta
680 diversity. *Ecol Lett* **9**: 683-693

681 **Andrés-Colás N, Sancenón V, Rodríguez-Navarro S, Mayo S, Thiele DJ, Ecker JR, Puig S,**
682 **Peñarrubia L** (2006) The Arabidopsis heavy metal P-type ATPase HMA5 interacts with
683 metallochaperones and functions in copper detoxification of roots. *Plant J* **45**: 225-236

684 **Baker AJ, Whiting SN** (2002) In search of the Holy Grail - a further step in understanding metal
685 hyperaccumulation? *New Phytol* **155**: 1-4

686 **Becher M, Talke IN, Krall L, Krämer U** (2004) Cross-species microarray transcript profiling
687 reveals high constitutive expression of metal homeostasis genes in shoots of the zinc
688 hyperaccumulator *Arabidopsis halleri*. *Plant J* **37**: 251-268

689 **Broadley MR, Brown P, Cakmak I, Rengel Z, Zhao F** (2012) Function of nutrients:
690 micronutrients. In: Marschner P, ed. Marschner's mineral nutrition of higher plant 3rd edn.

691 Academic Press, pp 191-248

692 **Broadley MR, White PJ, Hammond JP, Zelko I, Lux A** (2007) Zinc in plants. *New Phytol* **173**:

693 677-702

694 **Bughio N, Yamaguchi H, Nishizawa NK, Nakanishi H, Mori S** (2002) Cloning an iron-regulated

695 metal transporter from rice. *J Exp Bot* **53**: 1677-1682

696 **Burleigh SH, Kristensen BK, Bechmann IE** (2003) A plasma membrane zinc transporter from

697 *Medicago truncatula* is up-regulated in roots by Zn fertilization, yet down-regulated by

698 arbuscular mycorrhizal colonization. *Plant Mol Biol* **52**: 1077-1088

699 **Cakmak I** (2008) Enrichment of cereal grains with zinc: agronomic or genetic biofortification?

700 *Plant Soil* **302**: 1-17

701 **Cakmak I** (2012) HarvestPlus zinc fertilizer project: HarvestZinc. *Better Crops* **96**: 17-19

702 **Cakmak I, Pfeiffer WH, McClafferty B** (2010) Biofortification of durum wheat with zinc and

703 iron. *Cereal Chem* **87**: 10-20

704 **Cakmak I, McLaughlin MJ, White P** (2017) Zinc for better crop production and human health.

705 *Plant Soil* **411**: 1-4

706 **Cakmak I, Torun A, Millet E, Feldman M, Fahima T, Korol A, Nevo E, Braun HJ, Özkan H**

707 (2004) *Triticum dicoccoides*: an important genetic resource for increasing zinc and iron

708 concentration in modern cultivated wheat. *Soil Sci Plant Nutr* **50**:1047-1054

709 **Caldelas C, Weiss DJ** (2017) Zinc homeostasis and isotopic fractionation in plants: a review. *Plant*

710 *Soil* **411**: 17-46

711 **Capstaff NM, Miller AJ** (2018) Improving the yield and nutritional quality of forage crops. *Front*

712 *Plant Sci* **9**: 1-18

713 **Chaney RL** (1993) Zinc phytotoxicity. In: *Zinc in soils and plants*. Dordrecht, Netherlands:

714 Springer, pp 135-150

- 715 **Ciccolini V, Pellegrino E, Coccina A, Fiaschi AI, Cerretani D, Sgherri C, Quartacci MF,**
716 **Ercoli L** (2017) Biofortification with iron and zinc improves nutritional and nutraceutical
717 properties of common wheat flour and bread. *J Agr Food Chem* **65**: 5443-5452
- 718 **Clarke KR, Gorley RN** (2015) Getting started with PRIMER v7. Plymouth Marine Laboratory,
719 Plymouth, UK: PRIMER-E, 20
- 720 **Clemens S, Deinlein U, Ahmadi H, Höreth S, Uraguchi S** (2013) Nicotianamine is a major player
721 in plant Zn homeostasis. *Biometals* **26**: 623-632
- 722 **Curie C, Cassin G, Couch D, Divol F, Higuchi K, Le Jean M, Misson J, Shikora A, Czernic P,**
723 **Mari S** (2009) Metal movement within the plant: contribution of nicotianamine and yellow
724 stripe 1-like transporters. *Ann Bot-London* **103**: 1-11
- 725 **Deinlein U, Weber M, Schmidt H, Rensch S, Trampczynska A, Hansen TH, Husted S,**
726 **Schjoerring, JK, Talke IN, Krämer U, et al.** (2012) Elevated nicotianamine levels in
727 *Arabidopsis halleri* roots play a key role in zinc hyperaccumulation. *Plant Cell* **24**: 708-723
- 728 **Desbrosses-Fonrouge AG, Voigt K, Schröder A, Arrivault S, Thomine S, Krämer U** (2005)
729 *Arabidopsis thaliana* MTP1 is a Zn transporter in the vacuolar membrane which mediates Zn
730 detoxification and drives leaf Zn accumulation. *FEBS Lett* **579**: 4165-4174
- 731 **Deshpande P, Dapkekar A, Oak MD, Paknikar KM, Rajwade JM** (2017) Zinc complexed
732 chitosan/TPP nanoparticles: a promising micronutrient nanocarrier suited for foliar
733 application. *Carbohydr Polym* **165**: 394-401
- 734 **Desjardins P, Conklin D** (2010) NanoDrop microvolume quantitation of nucleic acids. *JOVE-J*
735 *Vis Exp* **5**: e2565
- 736 **Di Baccio D, Tognetti R, Minnocci A, Sebastiani L** (2009) Responses of the *Populus x*
737 *euramericana* clone I-214 to excess zinc: carbon assimilation, structural modifications, metal
738 distribution and cellular localization. *Environ Exp Bot* **67**: 153-163
- 739 **Eckhardt U, Marques AM, Buckhout TJ** (2001) Two iron-regulated cation transporters from
740 tomato complement metal uptake-deficient yeast mutants. *Plant Mol Biol* **45**: 437-448

741 **Eide D, Broderius M, Fett, J Guerinot ML** (1996) A novel iron-regulated metal transporter from
742 plants identified by functional expression in yeast. *P Natl Acad Sci USA* **93**: 5624-5628

743 **Erenoglu EB, Kutman UB, Ceylan Y, Yildiz B, Cakmak I** (2011) Improved nitrogen nutrition
744 enhances root uptake, root-to-shoot translocation and remobilization of zinc (⁶⁵Zn) in wheat.
745 *New Phytol* **189**: 438-448

746 **Fageria NK, Filho MB, Moreira A, Guimarães CM** (2009) Foliar fertilization of crop plants. *J*
747 *Plant Nutr* **32**: 1044-1064

748 **Foroughi S, Baker AJM, Roessner U, Johnson AAT, Bacic A, Callahan DL** (2014)
749 Hyperaccumulation of zinc by *Noccaea caerulescens* results in a cascade of stress responses
750 and changes in the elemental profile. *Metallomics* **6**: 1671-1682

751 **Foyer CH, Lam H-M, Nguyen HT, Siddique KHM, Varshney RK, Colmer TD, Cowling W,**
752 **Bramley H, Mori TA, Hodgson JM, et al** (2016) Neglecting legumes has compromised
753 human health and sustainable food production. *Nat Plants* **2**: 16112

754 **Geissler C, Powers HJ** (2017) Human nutrition. Oxford University Press, Oxford, UK

755 **Gregory PJ, Wahbi A, Adu-Gyamfi J, Heiling M, Gruber R, Joy EJM, Broadley MR** (2017)
756 Approaches to reduce zinc and iron deficits in food systems. *Global Food Secur-Agr* **15**: 1-10

757 **Grotz N, Fox T, Connolly E, Park W, Guerinot ML, Eide D** (1998) Identification of a family of
758 zinc transporter genes from *Arabidopsis* that respond to zinc deficiency. *P Natl Acad Sci USA*
759 **95**: 7220-7224

760 **Grotz N, Guerinot ML** (2006) Molecular aspects of Cu, Fe and Zn homeostasis in plants. *Biochim*
761 *Biophys Acta* **1763**: 595-608

762 **Gustin JL, Loureiro ME, Kim D, Na G, Tikhonova M, Salt DE** (2009) MTP1-dependent Zn
763 sequestration into shoot vacuoles suggests dual roles in Zn tolerance and accumulation in Zn-
764 hyperaccumulating plants. *Plant J* **57**: 1116-1127

765 **Hanikenne M, Talke IN, Haydon MJ, Lanz C, Nolte A, Motte P, Kroymann J, Weigel D,**
766 **Krämer U** (2008) Evolution of metal hyperaccumulation required cis-regulatory changes and
767 triplication of HMA4. *Nature* **453**: 391-395

768 **Haydon MJ, Cobbett CS** (2007) A novel major facilitator superfamily protein at the tonoplast
769 influences zinc tolerance and accumulation in *Arabidopsis*. *Plant Physiol* **143**: 1705-1719

770 **Haydon MJ, Kawachi M, Wirtz M, Hillmer S, Hell R, Krämer U** (2012) Vacuolar
771 nicotianamine has critical and distinct roles under iron deficiency and for zinc sequestration in
772 *Arabidopsis*. *Plant Cell* **24**: 724

773 **Hernand V, Julio E, de Borne FD, Punshon T, Ricachenevsky FK, Bellec A, Gosti F,**
774 **Berthomieu P** (2014) Inactivation of two newly identified tobacco heavy metal ATPases
775 leads to reduced Zn and Cd accumulation in shoots and reduced pollen germination.
776 *Metallomics* **6**: 1427-1440

777 **Huma ZE, Khan, ZI, Noorka IR, Ahmad K, Bayat AR, Wajid K** (2019) Bioaccumulation of
778 zinc and copper in tissues of chicken fed corn grain irrigated with different water regimes. *Int*
779 *J Environ Res* **13**: 689-703

780 **Hussain D, Haydon MJ, Wang Y, Wong E, Sherson SM, Young J, Camakaris J, Harper JF,**
781 **Cobbett CS** (2004) P-type ATPase heavy metal transporters with roles in essential zinc
782 homeostasis in *Arabidopsis*. *Plant Cell* **16**: 1327-1339

783 **Ishimaru Y, Suzuki M, Tsukamoto T, Suzuki K, Nakazono M, Kobayashi T, Wada Y,**
784 **Watanabe S., Matsushashi S, Nakanishi H, et al** (2006) Rice plants take up iron as an Fe³⁺-
785 phytosiderophore and as Fe²⁺. *Plant J* **45**: 335–346

786 **Jean ML, Schikora A, Mari S, Briat JF, Curie C** (2005) A loss-of-function mutation in AtYSL1
787 reveals its role in iron and nicotianamine seed loading. *Plant J* **44**: 769-782

788 **Keen CL, Gershwin ME** (1990) Zinc deficiency and immune function. *Annu Rev Nutr* **10**: 415-
789 431

790 **Kolaj-Robin O, Russell D, Hayes KA, Pembroke JT, Soulimane T** (2015) Cation diffusion
791 facilitator family: structure and function. *FEBS Lett* **589**: 1283-1295

792 **Kumar S, Stecher G, Li M, Knyaz C, Tamura K** (2018) MEGA X: molecular evolutionary
793 genetics analysis across computing platforms. *Mol Biol Evol* **35**: 1547-1549

794 **Kumssa DB, Joy EJ, Ander EL, Watts MJ, Young SD, Walker S, Broadley MR** (2015) Dietary
795 calcium and zinc deficiency risks are decreasing but remain prevalent. *Sci Rep* **5**: 10974

796 **Li S, Zhou X, Huang Y, Zhu L, Zhang S, Zhao Y, Chen R** (2013) Identification and
797 characterization of the zinc-regulated transporters, iron-regulated transporter-like protein
798 (ZIP) gene family in maize. *BMC Plant Biol* **13**: 114

799 **Livak K J, Schmittgen, TD** (2001) Analysis of relative gene expression data using real-time
800 quantitative PCR and the 2- $\Delta\Delta$ CT method. *Methods* **25**: 402-408

801 **López-Millán AF, Ellis DR, Grusak MA** (2004) Identification and characterization of several new
802 members of the ZIP family of metal ion transporters in *Medicago truncatula*. *Plant Mol Biol*
803 **54**: 583-596

804 **Mäser P, Thomine S, Schroeder JJ, Ward JM, Hirsch K, Sze H, Talke IN, Amtmann A,**
805 **Maathuis FJM, Sanders D, et al.** (2001) Phylogenetic relationship within cation transporter
806 families of *Arabidopsis*. *Plant Physiol* **126**: 1646-1667

807 **McDonald P, Edwards RA, Greenhalgh JFD, Morgan CA** (2002) Animal nutrition. Switzerland:
808 Pearson Education Limited, Harlow, UK

809 **Milner MJ, Seamon J, Craft E, Kochian LV** (2013) Transport properties of members of the ZIP
810 family in plants and their role in Zn and Mn homeostasis. *J Exp Bot* **64**: 369-381

811 **Nei M, Kumar S** (2000) Molecular evolution and phylogenetics. Oxford University Press, Oxford,
812 UK

813 **Nicot N, Hausman JF, Hoffmann L, Evers D** (2005) Housekeeping gene selection for real-time
814 RT-PCR normalization in potato during biotic and abiotic stress. *J Exp Bot* **56**: 2907-2914

- 815 **Nölte J** (2003) ICP Emission Spectrometry: a practical guide (Vol. 1). Wiley-VCH Weinheim,
816 Germany
- 817 **Ó Lochlainn S, Bowen HC, Fray RG, Hammond JP, King GJ, White PJ, Broadley MR** (2011)
818 Tandem quadruplication of HMA4 in the zinc (Zn) and cadmium (Cd) hyperaccumulator
819 *Noccaea caerulescens*. Plos One **6**: e17814
- 820 **Olsen LI, Palmgren MG** (2014) Many rivers to cross: the journey of zinc from soil to seed. Front
821 Plant Sci **5**: 30
- 822 **O'Rourke JA, Fu F, Bucciarelli B, Yang SS, Samac DA, Lamb JF, Li J, Dai X, Zhao PX,**
823 **Vance CP** (2015) The *Medicago sativa* gene index 1.2: a web-accessible gene expression
824 atlas for investigating expression differences between *Medicago sativa* subspecies. BMC
825 Genomics **16**: 502
- 826 **Palmer CM, Guerinot ML** (2009) Facing the challenges of Cu, Fe and Zn homeostasis in plants.
827 Nat Chem Biol **5**: 333-340
- 828 **Pedas P, Ytting CK, Fuglsang AT, Jahn TP, Schjoerring JK, Husted S** (2008) Manganese
829 efficiency in barley: identification and characterization of the metal ion transporter HvIRT1.
830 Plant Physiol **148**: 455-466
- 831 **Pita-Barbosa A, Ricachenevsky FK, Wilson M, Dottorini T, Salt DE** (2019) Transcriptional
832 plasticity buffers genetic variation in zinc homeostasis. Sci Rep **9**: 1-11
- 833 **Prasad AS** (2013) Discovery of human zinc deficiency: its impact on human health and disease.
834 Adv Nutr **4**: 176-190
- 835 **Ramesh SA, Shin R, Eide DJ, Schachtman DP** (2003) Differential metal selectivity and gene
836 expression of two zinc transporters from rice. Plant Physiol **133**: 126-134
- 837 **Rawat N, Neelam K, Tiwari VK, Dhaliwal HS** (2013) Biofortification of cereals to overcome
838 hidden hunger. Plant Breeding **132**: 437-445
- 839 **Saitou N, Nei M** (1987) The neighbor-joining method: a new method for reconstructing
840 phylogenetic trees. Mol Biol Evol **4**: 406-425

- 841 **Saltzman A, Birol E, Bouis HE, Boy E, De Moura FF, Islam Y, Pfeiffer WH** (2013)
842 Biofortification: progress toward a more nourishing future. *Glob Food Secur-Agr* **2**: 9-17
- 843 **Sankaran RP, Huguet T, Grusak MA** (2009) Identification of QTL affecting seed mineral
844 concentrations and content in the model legume *Medicago truncatula*. *Theor Appl Genet* **119**:
845 241-253
- 846 **Sasaki H, Hirose T, Watanabe Y, Ohsugi R** (1998) Carbonic anhydrase activity and CO₂-transfer
847 resistance in Zn-deficient rice leaves. *Plant Physiol* **118**: 929-934
- 848 **Sasaki A, Yamaji N, Mitani-Ueno N, Kashino M, Ma JF** (2015) A node-localized transporter
849 OsZIP3 is responsible for the preferential distribution of Zn to developing tissues in rice.
850 *Plant J* **84**: 374-384
- 851 **Sharma SS, Dietz KJ, Mimura T** (2016) Vacuolar compartmentalization as indispensable
852 component of heavy metal detoxification in plants. *Plant Cell Environ* **39**: 1112-1126
- 853 **Sinclair SA, Senger T, Talke IN, Cobbett CS, Haydon MJ, Kraemer U** (2018) Systemic
854 upregulation of MTP2-and HMA2-mediated Zn partitioning to the shoot supplements local
855 Zn deficiency responses. *Plant Cell* **30**: 2463-2479
- 856 **Sinclair SA, Krämer U** (2012) The zinc homeostasis network of land plants. *BBA-Mol Cell Res*
857 **1823**: 1553-1567
- 858 **Talke IN, Hanikenne M, Krämer U** (2006) Zinc-dependent global transcriptional control,
859 transcriptional deregulation, and higher gene copy number for genes in metal homeostasis of
860 the hyperaccumulator *Arabidopsis halleri*. *Plant Physiol* **142**: 148-167
- 861 **Tiong J, McDonald G, Genc Y, Shirley N, Langridge P, Huang CY** (2015) Increased expression
862 of six ZIP family genes by zinc (Zn) deficiency is associated with enhanced uptake and root-
863 to-shoot translocation of Zn in barley (*Hordeum vulgare*). *New Phytol* **207**: 1097-1109
- 864 **Vert G, Grotz N, Dédaldéchamp F, Gaymard F, Guerinot ML, Briat JF, Curie C** (2002) IRT1,
865 an *Arabidopsis* transporter essential for iron uptake from the soil and for plant growth. *Plant*
866 *Cell* **14**: 1223-1233

- 867 **Weber M, Harada E, Vess C, Roepenack-Lahaye EV, Clemens S** (2004) Comparative
868 microarray analysis of *Arabidopsis thaliana* and *Arabidopsis halleri* roots identifies
869 nicotianamine synthase, a ZIP transporter and other genes as potential metal
870 hyperaccumulation factors. *Plant J* **37**: 269-281
- 871 **White PJ** (2012) Long-distance transport in the xylem and phloem. In Marschner P, ed.
872 Marschner's mineral nutrition of higher plants. 3rd edn. Academic Press, pp 49-70
- 873 **White PJ** (2016) Biofortification of Edible Crops. eLS 1-8
- 874 **White PJ, Broadley MR** (2005) Biofortifying crops with essential mineral elements. *Trends Plant*
875 *Sci* **10**: 586-593
- 876 **White PJ, Broadley MR** (2009) Biofortification of crops with seven mineral elements often
877 lacking in human diets—iron, zinc, copper, calcium, magnesium, selenium and iodine. *New*
878 *Phytol* **182**: 49-84
- 879 **White PJ, Broadley MR** (2011) Physiological limits to zinc biofortification of edible crops. *Front*
880 *Plant Sci* **2**: 80
- 881 **White PJ, Pongrac P** (2017) Heavy-metal toxicity in plants. In: Shabala S, ed. *Plant Stress*
882 *Physiology*, 2nd ed., Wallingford, UK, pp 301-331.
- 883 **White PJ, Thompson JA, Wright G, Rasmussen SK** (2017) Biofortifying Scottish potatoes with
884 zinc. *Plant Soil* **411**: 151-165
- 885 **WHO** (2005) Comparative quantification of health risks: global and regional burden of diseases
886 attributable to selected major risk factors. In: Ezzati M, Lopez AD, Rodgers A., Murray CJL.
887 eds. World Health Organization, Geneva, Switzerland.
- 888 **Wickham H** (2011) ggplot2. *Wires Comput Stat* **3**: 180-185
- 889 **Wintz H, Fox T, Wu YY, Feng V, Chen W, Chang HS, Zhu T, Vulpe C** (2003) Expression
890 profiles of *Arabidopsis thaliana* in mineral deficiencies reveal novel transporters involved in
891 metal homeostasis. *J Biol Chem* **278**: 47644-47653
- 892 **Yilmaz O, Kazar GA, Cakmak I, Ozturk L** (2017) Differences in grain zinc are not correlated

893 with root uptake and grain translocation of zinc in wild emmer and durum wheat genotypes.
 894 Plant Soil **411**: 69-79
 895 **Zhao H, Eide D** (1996) The yeast ZRT1 gene encodes the zinc transporter protein of a high-affinity
 896 uptake system induced by zinc limitation. P Natl Acad Sci USA **93**: 2454-2458

Figure 1. Zinc (Zn) concentrations in shoots and roots of alfalfa (*Medicago sativa*) five days after the application of Zn doses of 0, 0.01, 0.1, 0.5, 1 or 10 mg Zn plant^{-1} to leaves. Means $\pm$ standard error of three replicates are shown. Differences among the applied Zn doses were tested separately for shoot and root by one-way analysis of variance. Different letters denote significant differences in Zn concentrations in shoots and roots independently, according to Tukey-B honestly test ($P < 0.05$).

Figure 2. Relative expression of seven transmembrane zinc (Zn) transporter genes (*MsZIP1-7*) five days after the application of Zn doses of 0, 0.1, 1 or 10 mg Zn plant⁻¹ to leaves of alfalfa (*Medicago sativa*). Means $\pm$ standard error of three replicates are shown. The expression levels were calculated relative to reference genes (*MsACT-101* for shoot and *MsEF1- α* for root) and to the control (0 mg Zn plant⁻¹). The broken line denotes the threshold between up- and down-regulation relative to the control. Differences in the expressions of each gene after different Zn doses were tested separately for shoot and root by one-way analysis of variance. Different letters denote significant differences among Zn doses, according to Tukey-B test ($P < 0.05$).

Figure 3. Relative expression of genes related to Zn transport processes (*MsZIF1*, *MsHMA4*, *MsYSL1*, *MsMTP1* and *MsNAS1*) five days after the application of Zn doses of 0, 0.1, 1 or 10 mg Zn plant⁻¹ to leaves of alfalfa (*Medicago sativa*). Means $\pm$ standard error of three replicates are shown. The expression levels were calculated relative to reference genes (*MsACT-101* for shoot and *MsEF1- α* for root) and to the control (0 mg Zn plant⁻¹). The broken line denotes the threshold between up- and down-regulation relative to the control. Differences in the expression of each gene at the different Zn doses were tested separately for shoot and root by one-way analysis of variance. Different letters denote significant differences among Zn doses, according to Tukey-B test ($P < 0.05$).

Figure 4. Heatmaps reporting the correlations between the differences in expression of genes related to zinc (Zn) transport processes in alfalfa (*Medicago sativa*) after foliar application of Zn doses of 0.1, 1 or 10 mg Zn plant⁻¹ relative to a control dose of 0 mg Zn plant⁻¹. The similarity in the degree of correlation in fold-change of gene expression to Zn application relative to the control is based on the average linkage clustering of the Pearson correlations (r). In the clustering trees the genes are indicated in brown for roots and in green for shoots, while the ranks of correlations of the heatmap are indicated by color intensity (r 0 to 1: from low to strong intensity of green). Seven genes encoding transmembrane Zn transporter (*MsZIP*₁₋₇) (a); four genes encoding cellular Zn transporters (including vacuolar transporters) (*MsZIF1*, *MsHMA4*, *MsYSL1* and *MsMTP1*) and a gene encoding a nicotianamine synthase (*MsNAS1*) (b).

Figure 5. Suggested model for the roles of putative genes encoding proteins involved in Zinc (Zn) transport-related processes in alfalfa (*Medicago sativa*). The sites of action in the plant (i.e., root cytoplasm, rc; root vacuole, rv; xylem and apoplast, X/A; phloem, P; leaf cytoplasm, lc; leaf vacuole, lv) and the element (E) fluxes (K_{1-13}) are reported. The concentration of the element is indicated in each site [E]. The scheme synthesizes information across studies in various plants. Gene abbreviations: *ZIP*, Zrt-/Irt-like Protein; *NAS*, Nicotianamine synthase; *ZIF*, Zinc-Induced Facilitator; *MTP*, Metal Transporter Protein; *HMA*, P_{1B} -type Heavy Metal ATPase; *YSL*, Yellow Stripe Like Protein; *ZIP?* indicates a generic *ZIP*; free diffusion: diffusion through leaf epidermis; stomata: absorption through stomata. Plant abbreviations: *Mt*, *Medicago truncatula*; *At*, *Arabidopsis thaliana*; *Os*, *Oryza sativa*. References: ^aLópez-Millán et al., 2004; ^bMilner et al., 2013; ^cAarts, 2014; ^dClemens et al., 2013; ^eCurie et al., 2009; ^fHaydon et al., 2012; ^gDesbrosses-Fonrouge et al., 2005; ^hHussain et al., 2004; ⁱPalmer and Guerinot, 2009; ^jBurleigh et al., 2003; ^kSasaki et al., 2015; ^lFageria et al., 2009.

Parsed Citations

Aarts MG (2014) Nicotianamine secretion for zinc excess tolerance. *Plant Physiol* 166: 751-752.

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Albert IL, Nadassy K, Wodak SJ (1998) Analysis of zinc binding sites in protein crystal structures. *Protein Sci* 7: 1700-1716

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Alloway BJ (2008) Zinc in soils and crop nutrition. 2nd ed. IZA and IFA, Brussels, Belgium and Paris, France

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Alloway BJ (2009) Soil factors associated with zinc deficiency in crops and humans. *Environ Geochem Hlth* 31: 537-548

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Anderson MJ (2001) A new method for non-parametric multivariate analysis of variance. *Austral Ecol* 26: 32-46

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Anderson M, Braak CT (2003) Permutation tests for multi-factorial analysis of variance. *J Stat Comput Sim* 73: 85-113

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Anderson MJ, Ellingsen KE, McArdle BH (2006) Multivariate dispersion as a measure of beta diversity. *Ecol Lett* 9: 683-693

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Andrés-Colás N, Sancenón V, Rodríguez-Navarro S, Mayo S, Thiele DJ, Ecker JR, Puig S, Peñarrubia L (2006) The Arabidopsis heavy metal P-type ATPase HMA5 interacts with metallochaperones and functions in copper detoxification of roots. *Plant J* 45: 225-236

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Baker AJ, Whiting SN (2002) In search of the Holy Grail - a further step in understanding metal hyperaccumulation? *New Phytol* 155: 1-4

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Becher M, Talke IN, Krall L, Krämer U (2004) Cross-species microarray transcript profiling reveals high constitutive expression of metal homeostasis genes in shoots of the zinc hyperaccumulator *Arabidopsis halleri*. *Plant J* 37: 251-268

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Broadley MR, Brown P, Cakmak I, Rengel Z, Zhao F (2012) Function of nutrients: micronutrients. In: Marschner P, ed. *Marschner's mineral nutrition of higher plant* 3rd edn. Academic Press, pp 191-248

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Broadley MR, White PJ, Hammond JP, Zelko I, Lux A (2007) Zinc in plants. *New Phytol* 173: 677-702

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Bughio N, Yamaguchi H, Nishizawa NK, Nakanishi H, Mori S (2002) Cloning an iron-regulated metal transporter from rice. *J Exp Bot* 53: 1677-1682

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Burleigh SH, Kristensen BK, Bechmann IE (2003) A plasma membrane zinc transporter from *Medicago truncatula* is up-regulated in roots by Zn fertilization, yet down-regulated by arbuscular mycorrhizal colonization. *Plant Mol Biol* 52: 1077-1088

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Cakmak I (2008) Enrichment of cereal grains with zinc: agronomic or genetic biofortification? *Plant Soil* 302: 1-17

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Cakmak I (2012) HarvestPlus zinc fertilizer project: HarvestZinc. *Better Crops* 96: 17-19

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Cakmak I, Pfeiffer WH, McClafferty B (2010) Biofortification of durum wheat with zinc and iron. *Cereal Chem* 87: 10-20

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Cakmak I, McLaughlin MJ, White P (2017) Zinc for better crop production and human health. Plant Soil 411: 1-4

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Cakmak I, Torun A, Millet E, Feldman M, Fahima T, Korol A, Nevo E, Braun HJ, Özkan H (2004) Triticum dicoccoides: an important genetic resource for increasing zinc and iron concentration in modern cultivated wheat. Soil Sci Plant Nutr 50:1047-1054

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Caldelas C, Weiss DJ (2017) Zinc homeostasis and isotopic fractionation in plants: a review. Plant Soil 411: 17-46

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Capstaff NM, Miller AJ (2018) Improving the yield and nutritional quality of forage crops. Front Plant Sci 9: 1-18

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Chaney RL (1993) Zinc phytotoxicity. In: Zinc in soils and plants. Dordrecht, Netherlands: Springer, pp 135-150

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ciccolini V, Pellegrino E, Coccina A, Fiaschi AI, Cerretani D, Sgherri C, Quartacci MF, Ercoli L (2017) Biofortification with iron and zinc improves nutritional and nutraceutical properties of common wheat flour and bread. J Agr Food Chem 65: 5443-5452

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Clarke KR, Gorley RN (2015) Getting started with PRIMER v7. Plymouth Marine Laboratory, Plymouth, UK: PRIMER-E, 20

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Clemens S, Deinlein U, Ahmadi H, Höreth S, Uraguchi S (2013) Nicotianamine is a major player in plant Zn homeostasis. Biometals 26: 623-632

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Curie C, Cassin G, Couch D, Divol F, Higuchi K, Le Jean M, Misson J, Shikora A, Czernic P, Mari S (2009) Metal movement within the plant: contribution of nicotianamine and yellow stripe 1-like transporters. Ann Bot-London 103: 1-11

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Deinlein U, Weber M, Schmidt H, Rensch S, Trampczynska A, Hansen TH, Husted S, Schjoerring, JK, Talke IN, Krämer U, et al. (2012) Elevated nicotianamine levels in Arabidopsis halleri roots play a key role in zinc hyperaccumulation. Plant Cell 24: 708-723

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Desbrosses-Fonrouge AG, Voigt K, Schröder A, Arrivault S, Thomine S, Krämer U (2005) Arabidopsis thaliana MTP1 is a Zn transporter in the vacuolar membrane which mediates Zn detoxification and drives leaf Zn accumulation. FEBS Lett 579: 4165-4174

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Deshpande P, Dapkekar A, Oak MD, Paknikar KM, Rajwade JM (2017) Zinc complexed chitosan/TPP nanoparticles: a promising micronutrient nanocarrier suited for foliar application. Carbohydr Polym 165: 394-401

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Desjardins P, Conklin D (2010) NanoDrop microvolume quantitation of nucleic acids. JOVE-J Vis Exp 5: e2565

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Di Baccio D, Tognetti R, Minnocci A, Sebastiani L (2009) Responses of the Populus x euramericana clone I-214 to excess zinc: carbon assimilation, structural modifications, metal distribution and cellular localization. Environ Exp Bot 67: 153-163

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Eckhardt U, Marques AM, Buckhout TJ (2001) Two iron-regulated cation transporters from tomato complement metal uptake-deficient yeast mutants. Plant Mol Biol 45: 437-448

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Eide D, Broderius M, Fett, J, Gueriot ML (1996) A novel iron-regulated metal transporter from plants identified by functional expression in yeast. P Natl Acad Sci USA 93: 5624-5628

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Erenoglu EB, Kutman UB, Ceylan Y, Yildiz B, Cakmak I (2011) Improved nitrogen nutrition enhances root uptake, root-to-shoot

translocation and remobilization of zinc (65Zn) in wheat. New Phytol 189: 438-448

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Fageria NK, Filho MB, Moreira A, Guimarães CM (2009) Foliar fertilization of crop plants. J Plant Nutr 32: 1044-1064

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Foroughi S, Baker AJM, Roessner U, Johnson AAT, Bacic A, Callahan DL (2014) Hyperaccumulation of zinc by Noccaea caerulea results in a cascade of stress responses and changes in the elemental profile. Metallomics 6: 1671-1682

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Foyer CH, Lam H-M, Nguyen HT, Siddique KHM, Varshney RK, Colmer TD, Cowling W, Bramley H, Mori TA, Hodgson JM, et al (2016) Neglecting legumes has compromised human health and sustainable food production. Nat Plants 2: 16112

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Geissler C, Powers HJ (2017) Human nutrition. Oxford University Press, Oxford, UK

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Gregory PJ, Wahbi A, Adu-Gyamfi J, Heiling M, Gruber R, Joy EJM, Broadley MR (2017) Approaches to reduce zinc and iron deficits in food systems. Global Food Secur-Agr 15: 1-10

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Grotz N, Fox T, Connolly E, Park W, Guerinot ML, Eide D (1998) Identification of a family of zinc transporter genes from Arabidopsis that respond to zinc deficiency. P Natl Acad Sci USA 95: 7220-7224

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Grotz N, Guerinot ML (2006) Molecular aspects of Cu, Fe and Zn homeostasis in plants. Biochim Biophys Acta 1763: 595-608

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Gustin JL, Loureiro ME, Kim D, Na G, Tikhonova M, Salt DE (2009) MTP1-dependent Zn sequestration into shoot vacuoles suggests dual roles in Zn tolerance and accumulation in Zn-hyperaccumulating plants. Plant J 57: 1116-1127

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Hanikenne M, Talke IN, Haydon MJ, Lanz C, Nolte A, Motte P, Kroymann J, Weigel D, Krämer U (2008) Evolution of metal hyperaccumulation required cis-regulatory changes and triplication of HMA4. Nature 453: 391-395

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Haydon MJ, Cobbett CS (2007) A novel major facilitator superfamily protein at the tonoplast influences zinc tolerance and accumulation in Arabidopsis. Plant Physiol 143: 1705-1719

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Haydon MJ, Kawachi M, Wirtz M, Hillmer S, Hell R, Krämer U (2012) Vacuolar nicotianamine has critical and distinct roles under iron deficiency and for zinc sequestration in Arabidopsis. Plant Cell 24: 724

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Hernand V, Julio E, de Borne FD, Punshon T, Ricachenevsky FK, Bellec A, Gosti F, Berthomieu P (2014) Inactivation of two newly identified tobacco heavy metal ATPases leads to reduced Zn and Cd accumulation in shoots and reduced pollen germination. Metallomics 6: 1427-1440

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Huma ZE, Khan, ZI, Noorka IR, Ahmad K, Bayat AR, Wajid K (2019) Bioaccumulation of zinc and copper in tissues of chicken fed corn grain irrigated with different water regimes. Int J Environ Res 13: 689-703

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Hussain D, Haydon MJ, Wang Y, Wong E, Sherson SM, Young J, Camakaris J, Harper JF, Cobbett CS (2004) P-type ATPase heavy metal transporters with roles in essential zinc homeostasis in Arabidopsis. Plant Cell 16: 1327-1339

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ishimaru Y, Suzuki M, Tsukamoto T, Suzuki K, Nakazono M, Kobayashi T, Wada Y, Watanabe S., Matsushashi S, Nakanishi H, et al (2006) Rice plants take up iron as an Fe3p-phytosiderophore and as Fe2p. Plant J 45: 335-346

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Jean ML, Schikora A, Mari S, Briat JF, Curie C (2005) A loss-of-function mutation in AtYSL1 reveals its role in iron and nicotianamine seed loading. Plant J 44: 769-782

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Keen CL, Gershwin ME (1990) Zinc deficiency and immune function. Annu Rev Nutr 10: 415-431

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Kolaj-Robin O, Russell D, Hayes KA, Pembroke JT, Soulimane T (2015) Cation diffusion facilitator family: structure and function. FEBS Lett 589: 1283-1295

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Kumar S, Stecher G, Li M, Knyaz C, Tamura K (2018) MEGA X: molecular evolutionary genetics analysis across computing platforms. Mol Biol Evol 35: 1547-1549

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Kumssa DB, Joy EJ, Ander EL, Watts MJ, Young SD, Walker S, Broadley MR (2015) Dietary calcium and zinc deficiency risks are decreasing but remain prevalent. Sci Rep 5: 10974

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Li S, Zhou X, Huang Y, Zhu L, Zhang S, Zhao Y, Chen R (2013) Identification and characterization of the zinc-regulated transporters, iron-regulated transporter-like protein (ZIP) gene family in maize. BMC Plant Biol 13: 114

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Livak K J, Schmittgen, TD (2001) Analysis of relative gene expression data using real-time quantitative PCR and the 2- $\Delta\Delta$ CT method. Methods 25: 402-408

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

López-Millán AF, Ellis DR, Grusak MA (2004) Identification and characterization of several new members of the ZIP family of metal ion transporters in *Medicago truncatula*. Plant Mol Biol 54: 583-596

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Mäser P, Thomine S, Schroeder JI, Ward JM, Hirsch K, Sze H, Talke IN, Amtmann A, Maathuis FJM, Sanders D, et al. (2001) Phylogenetic relationship within cation transporter families of Arabidopsis. Plant Physiol 126: 1646-1667

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

McDonald P, Edwards RA, Greenhalgh JFD, Morgan CA (2002) Animal nutrition. Switzerland: Pearson Education Limited, Harlow, UK

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Milner MJ, Seamon J, Craft E, Kochian LV (2013) Transport properties of members of the ZIP family in plants and their role in Zn and Mn homeostasis. J Exp Bot 64: 369-381

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Nei M, Kumar S (2000) Molecular evolution and phylogenetics. Oxford University Press, Oxford, UK

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Nicot N, Hausman JF, Hoffmann L, Evers D (2005) Housekeeping gene selection for real-time RT-PCR normalization in potato during biotic and abiotic stress. J Exp Bot 56: 2907-2914

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Nölte J (2003) ICP Emission Spectrometry: a practical guide (Vol. 1). Wiley-VCH Weinheim, Germany

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ó Lochlainn S, Bowen HC, Fray RG, Hammond JP, King GJ, White PJ, Broadley MR (2011) Tandem quadruplication of HMA4 in the zinc (Zn) and cadmium (Cd) hyperaccumulator *Noccaea caerulea*. Plos One 6: e17814

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Olsen LI, Palmgren MG (2014) Many rivers to cross: the journey of zinc from soil to seed. Front Plant Sci 5: 30

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

O'Rourke JA, Fu F, Bucciarelli B, Yang SS, Samac DA, Lamb JF, Li J, Dai X, Zhao PX, Vance CP (2015) The Medicago sativa gene index 1.2: a web-accessible gene expression atlas for investigating expression differences between Medicago sativa subspecies. BMC Genomics 16: 502

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Palmer CM, Gueriot ML (2009) Facing the challenges of Cu, Fe and Zn homeostasis in plants. Nat Chem Biol 5: 333-340

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Pedraza P, Yting CK, Fuglsang AT, Jahn TP, Schjoerring JK, Husted S (2008) Manganese efficiency in barley: identification and characterization of the metal ion transporter HvIRT1. Plant Physiol 148: 455-466

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Pita-Barbosa A, Ricachenevsky FK, Wilson M, Dottorini T, Salt DE (2019) Transcriptional plasticity buffers genetic variation in zinc homeostasis. Sci Rep 9: 1-11

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Prasad AS (2013) Discovery of human zinc deficiency: its impact on human health and disease. Adv Nutr 4: 176-190

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Ramesh SA, Shin R, Eide DJ, Schachtman DP (2003) Differential metal selectivity and gene expression of two zinc transporters from rice. Plant Physiol 133: 126-134

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Rawat N, Neelam K, Tiwari VK, Dhaliwal HS (2013) Biofortification of cereals to overcome hidden hunger. Plant Breeding 132: 437-445

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Saitou N, Nei M (1987) The neighbor-joining method: a new method for reconstructing phylogenetic trees. Mol Biol Evol 4: 406-425

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Saltzman A, Birol E, Bouis HE, Boy E, De Moura FF, Islam Y, Pfeiffer WH (2013) Biofortification: progress toward a more nourishing future. Glob Food Secur-Agr 2: 9-17

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Sankaran RP, Huguet T, Grusak MA (2009) Identification of QTL affecting seed mineral concentrations and content in the model legume Medicago truncatula. Theor Appl Genet 119: 241-253

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Sasaki H, Hirose T, Watanabe Y, Ohsugi R (1998) Carbonic anhydrase activity and CO₂-transfer resistance in Zn-deficient rice leaves. Plant Physiol 118: 929-934

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Sasaki A, Yamaji N, Mitani-Ueno N, Kashino M, Ma JF (2015) A node-localized transporter OsZIP3 is responsible for the preferential distribution of Zn to developing tissues in rice. Plant J 84: 374-384

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Sharma SS, Dietz KJ, Mimura T (2016) Vacuolar compartmentalization as indispensable component of heavy metal detoxification in plants. Plant Cell Environ 39: 1112-1126

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Sinclair SA, Senger T, Talke IN, Cobbett CS, Haydon MJ, Kraemer U (2018) Systemic upregulation of MTP2-and HMA2-mediated Zn partitioning to the shoot supplements local Zn deficiency responses. Plant Cell 30: 2463-2479

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Sinclair SA, Krämer U (2012) The zinc homeostasis network of land plants. BBA-Mol Cell Res 1823: 1553-1567

Pubmed: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Talke IN, Hanikenne M, Krämer U (2006) Zinc-dependent global transcriptional control, transcriptional deregulation, and higher gene copy number for genes in metal homeostasis of the hyperaccumulator Arabidopsis halleri. Plant Physiol 142: 148-167

Pubmed: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Tiong J, McDonald G, Genc Y, Shirley N, Langridge P, Huang CY (2015) Increased expression of six ZIP family genes by zinc (Zn) deficiency is associated with enhanced uptake and root-to-shoot translocation of Zn in barley (*Hordeum vulgare*). *New Phytol* 207: 1097-1109

Pubmed: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Vert G, Grotz N, Dédaldéchamp F, Gaymard F, Guerinot ML, Briat JF, Curie C (2002) IRT1, an Arabidopsis transporter essential for iron uptake from the soil and for plant growth. *Plant Cell* 14: 1223-1233

Pubmed: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Weber M, Harada E, Vess C, Roepenack-Lahaye EV, Clemens S (2004) Comparative microarray analysis of Arabidopsis thaliana and Arabidopsis halleri roots identifies nicotianamine synthase, a ZIP transporter and other genes as potential metal hyperaccumulation factors. *Plant J* 37: 269-281

Pubmed: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

White PJ (2012) Long-distance transport in the xylem and phloem. In Marschner P, ed. Marschner's mineral nutrition of higher plants. 3rd edn. Academic Press, pp 49-70

Pubmed: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

White PJ (2016) Biofortification of Edible Crops. eLS 1-8

Pubmed: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

White PJ, Broadley MR (2005) Biofortifying crops with essential mineral elements. *Trends Plant Sci* 10: 586-593

Pubmed: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

White PJ, Broadley MR (2009) Biofortification of crops with seven mineral elements often lacking in human diets—iron, zinc, copper, calcium, magnesium, selenium and iodine. *New Phytol* 182: 49-84

Pubmed: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

White PJ, Broadley MR (2011) Physiological limits to zinc biofortification of edible crops. *Front Plant Sci* 2: 80

Pubmed: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

White PJ, Pongrac P (2017) Heavy-metal toxicity in plants. In: Shabala S, ed. *Plant Stress Physiology*, 2nd ed., Wallingford, UK, pp 301-331.

Pubmed: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

White PJ, Thompson JA, Wright G, Rasmussen SK (2017) Biofortifying Scottish potatoes with zinc. *Plant Soil* 411: 151-165

Pubmed: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

WHO (2005) Comparative quantification of health risks: global and regional burden of diseases attributable to selected major risk factors. In: Ezzati M, Lopez AD, Rodgers A, Murray CJL. eds. *World Health Organization*, Geneva, Switzerland.

Pubmed: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wickham H (2011) ggplot2. *Wires Comput Stat* 3: 180-185

Pubmed: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wintz H, Fox T, Wu YY, Feng V, Chen W, Chang HS, Zhu T, Vulpe C (2003) Expression profiles of Arabidopsis thaliana in mineral deficiencies reveal novel transporters involved in metal homeostasis. *J Biol Chem* 278: 47644-47653

Pubmed: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Yilmaz O, Kazar GA, Cakmak I, Ozturk L (2017) Differences in grain zinc are not correlated with root uptake and grain translocation of zinc in wild emmer and durum wheat genotypes. *Plant Soil* 411: 69-79

Pubmed: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zhao H, Eide D (1996) The yeast ZRT1 gene encodes the zinc transporter protein of a high-affinity uptake system induced by zinc limitation. *P Natl Acad Sci USA* 93: 2454-2458

Pubmed: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)