A limbic circuit selectively linking active escape to food suppression

Estefania P. Azevedo¹, Bowen Tan¹, Lisa E. Pomeranz¹, Violet Ivan¹, Robert N. Fetcho², Marc Schneeberger¹, Katherine R. Doerig¹, Conor Liston², Jeffrey M. Friedman^{1,3*}, Sarah A. Stern^{1*}.

¹Laboratory of Molecular Genetics, The Rockefeller University, New York, NY, 10065.

²Department of Psychiatry, Weill Cornell Medical College-New York Presbyterian Hospital, New York, NY, 10065

³Howard Hughes Medical Institute, The Rockefeller University, New York, NY, USA.

*Correspondence should be addressed to: sstern01@rockefeller.edu or friedj@mail.rockefeller.edu

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23

Abstract

Stress and anxiety are precipitating factors for eating disorders, but the neural basis linking stress to alterations in feeding is not well understood. Here we describe a novel population of stress-responsive neurons in the lateral septum (LS) of mice that express neurotensin (LS^{NTS}) in a sexually dimorphic, estrous cycle-dependent manner. We used in vivo imaging to show that LS^{NTS} neurons are activated by stressful experiences when flight is a viable option, but not by a stressful experience associated with freezing or immobility. LS^{NTS} activation leads to a decrease of food intake and body weight in mice, without altering locomotion or other behaviors associated with anxiety. Molecular profiling of LS^{NTS} neurons showed that these neurons co-express Glp1r (glucagon-like peptide 1 receptor), and both pharmacologic and genetic manipulations of Glp1r signaling in the LS recapitulates the behavioral effects of LS^{NTS} activation. Finally, we mapped the outputs of LS^{NTS} neurons and show that activation of LS^{NTS} nerve terminals in the lateral hypothalamus (LH), a well-established feeding center, also decrease food intake. Taken together, these results show that LS^{NTS} neurons link stress and anorexia via effects on hypothalamic pathways regulating food intake.

Keywords: feeding, lateral septum, stress, circuits

1 INTRODUCTION

2 A key function of the central nervous system (CNS) is to sense external
3 conditions and activate an appropriate behavioral response (McEwen and Akil,
4 2020). The detection of stressful conditions, in particular, is of critical importance
5 by enabling animals to remove themselves from dangerous situations. These
6 responses are also associated with other adaptive behavioral effects including
7 reduced feeding until the period of danger has passed, and these stereotyped
8 responses are conserved across organisms ranging from flies to mice and humans
9 (Yau and Potenza, 2013; Surendran et al., 2017).

10 We therefore set out to map neural circuits linking stress to feeding by initially
11 studying acute restraint stress, a behavioral paradigm in rodents which leads to a
12 robust decrease in food intake for up to 48 hours (Donohoe, 1984; Shimizu et al.,
13 1989). Previous studies have shown increased expression of c-fos, an early
14 immediate gene, in many brain areas following stress, including regions of the
15 limbic system (e.g. lateral septum, Sheehan et al., 2004). However, the cell-type
16 specificity and the activation dynamics of stress-responsive neurons in specific
17 regions of the limbic system has not been determined, nor has the mechanism by
18 which these neurons regulate feeding.

19 In this study, we used an unbiased transcriptomic approach (Knight et al.,
20 2012) to map and identify neurons in the limbic system that are activated during
21 stress-induced anorexia. Initial studies revealed that neurons in the lateral septum
22 that express neurotensin are acutely activated by restraint stress and we found
23 that acute or chronic chemogenetic activation of these neurons leads to significant

1 reductions of food intake and body weight. Chronic stress is also a risk factor for
2 the development of many psychiatric diseases that are characterized by
3 maladaptive responses (Yau and Potenza, 2013), including eating disorders,
4 which have a significantly greater incidence in females (Guarda et al., 2015).
5 Accordingly, we also found that the activity of these neurons is regulated by the
6 levels of female sex hormones, raising the possibility that LS^{NTS} neurons could
7 provide an entry point for studying neural circuits that may contribute to eating
8 disorders.

9

10 **RESULTS**

11

12 **Identification of a limbic population activated by acute restraint stress**

13 In order to identify specific neuronal populations that link stress to reduced
14 food intake, we first measured food intake in wild-type C57Bl/6J mice after one
15 hour of acute restraint stress and mapped the expression pattern of c-fos
16 throughout the whole mouse brain. As previously reported, acute restraint stress
17 for a period of 1h led to a highly significant decrease in food intake for the ensuing
18 2 hours (Fig. 1A, Two-way ANOVA with Bonferroni post-hoc, *p<0.05, ***p<0.001).
19 One hour of restraint stress also led to increased numbers of c-fos+ cells in a
20 number of brain regions within the limbic system: the basolateral amygdala [BLA],
21 bed nucleus of the stria terminalis [BNST], cortex (insular cortex [IC]) (Figure 1B
22 and Table S1) and elsewhere. Consistent with previous studies (Kubo et al., 2002),
23 acute restraint stress led to a marked increase in the number of c-fos+ cells in the

1 LS compared to controls (Figure S1A, Unpaired Student's t-test, $*p < 0.05$). The LS
2 is part of the anterior limbic system and is known to regulate several emotional
3 behaviors, particularly stress, anxiety and aggression (Kubo et al., 2002; Anthony
4 et al., 2014; Wong et al., 2016). In addition, previous studies from our group and
5 another showed that the artificial activation of neurons in the LS has a negative
6 effect on feeding (Azevedo et al., 2019; Sweeney and Yang, 2016), raising the
7 possibility that specific neurons in this region might link restraint stress to a non-
8 homeostatic reduction in food intake.

9 We next set out to determine the molecular identity of the neurons in the LS
10 that are activated by acute restraint stress using PhosphoTrap, an unbiased
11 transcriptomic method to molecularly profile activated neurons (Knight et al., 2012).
12 This method takes advantage of the fact that neuronal activation results in a
13 cascade of signaling events culminating in the phosphorylation of the ribosomal
14 protein S6 protein (pS6). These phosphorylated ribosomes can be
15 immunoprecipitated from mouse brain homogenates, thereby enriching mRNAs
16 selectively expressed in the activated neuronal population (Fig. 1C, left panel).
17 After polysome immunoprecipitation and RNA extraction and sequencing, RNAs
18 enriched relative to total RNA have been shown to mark activated neurons (Knight
19 et al., 2012; Azevedo et al., 2014; Stern et al., 2019). We applied this method to
20 identify markers for neurons in the lateral septum that are activated after acute
21 restraint stress relative to control mice (naïve) (Fig. 1C). The enrichment for each
22 gene was calculated as the number of reads in the immunoprecipitated RNA (IP)
23 relative to the total input RNA (IP/INP; Fig. 1C). As expected based on data from

1 immediate early gene mapping, we found an enrichment for activity-related genes
 2 (*FosB* and *Fos*) in the samples from mice exposed to acute restraint stress, (see
 3 Fig. 1B, LS and Figure S1A). We next analyzed the data to identify other genes
 4 that were significantly enriched (>2-fold or greater and q-value <0.05) in the
 5 samples from mice exposed to acute restraint stress (RS) compared to naïve mice
 6 (q value <0.05, Fig. 1C and Table S2). We found enrichment for several markers,
 7 including the gene for the neuropeptide neurotensin (*Nts*, 2.41-fold increase
 8 compared to naïve samples). We confirmed the enrichment using qPCR and found
 9 a 3.6-fold enrichment in neurotensin mRNA in the samples from the lateral septum
 10 of mice exposed to acute restraint stress compared to samples from naïve mice
 11 (Unpaired Student's t-test, $p < 0.05$).

12

13 **LS^{NTS} neurons respond to stress in a sexual dimorphic manner**

14 We further confirmed the activation of *Nts*-expressing neurons by acute
 15 restraint stress by analyzing the colocalization of *Nts* and *Fos* mRNA expression
 16 in the LS using *in situ* hybridization (Fig. 2). Both male and female mice showed
 17 significant colocalization between *Fos* and *Nts* in the LS after restraint stress
 18 compared to their control non-stressed, naïve counterparts (~65 cells in stressed
 19 females vs. 2 cells in non-stressed females; ~37 cells in stressed males vs. 22
 20 cells in non-stressed males; ** $p < 0.01$, * $p < 0.05$, Fig. 2B and C). Because the
 21 effects of acute stress are modulated by sex hormones in particular in eating
 22 disorders, we analyzed the co-expression of *Nts* and *Fos* in stressed female mice
 23 during different phases of the estrous cycle. Female mice in estrus exposed to

1 acute restraint stress had a ~2-fold higher levels of *Nts* and *Fos* colocalization
 2 compared to male mice (Figure 2B and C). Furthermore, the colocalization of *Nts*
 3 and *Fos* was significantly higher in female mice during the estrus phase compared
 4 to those that were in diestrus phase (Figure 2B, ~ 60 cells/slice in estrus compared
 5 to ~30 cells/slice in diestrus). The extent of the colocalization of *Nts* and *Fos* after
 6 acute restraint stress (RS) was similar in females during the diestrus phase to
 7 males (Figure 2B and C, ~37 cells/slice in males and ~25 cells/slice in females in
 8 diestrus). We also found that the higher levels of *Fos* colocalization with *Nts* in
 9 stressed female and male mice was due to both an increase in the levels of *Fos* in
 10 *Nts* neurons and a concomitant induction of *Nts* mRNA during stress (Figure S1B
 11 and C). In contrast, neurotensin mRNA was almost completely absent in the LS in
 12 non-stressed female controls (Fig 1B). These results suggest that that activity of
 13 *Nts* neurons as well as the expression of the *Nts* gene in the LS is regulated by
 14 acute restraint stress in a sexually dimorphic manner and potentiated by the
 15 increased levels of female sex hormones during estrus.

16

17 **LS^{NTS} neurons regulate food intake and body weight**

18 We next tested whether chemogenetic activation of *Nts* neurons would lead
 19 to a concomitant decrease in food consumption. We injected an AAV encoding a
 20 Cre dependent excitatory DREADD, (DIO) hM3Dq, a Gq-coupled (designer
 21 receptor exclusively activated by designer drugs, Armbruster et al., 2007), into the
 22 LS of *Nts*-Cre mice. Injection of clozapine-N-oxide (CNO) into these mice
 23 increased c-fos in the LS (Figure S1D, Unpaired Student's t-test) and significantly

1 decreased food intake compared to mice treated with CNO after injection of a
 2 control Cre-dependent mCherry AAV into Nts-cre mice (Fig. 3A, red bars vs black
 3 bars, Two-way ANOVA with Bonferroni post-hoc, $*p<0.05$, $***p<0.001$) as well as
 4 saline-injected mice expressing the DREADD (Figure S1E). This robust decrease
 5 in food intake was evident after 1 hour and persisted for 4, 8 and 24 hours after a
 6 single CNO injection (Figure S1F). Chronic stimulation of Nts neurons for 3 days
 7 in the LS also significantly decreased food intake (Fig. 3B, Two-way ANOVA with
 8 Bonferroni post-hoc, $**p<0.01$, $****p<0.0001$) and significantly reduced body
 9 weight relative to the aforementioned controls (Fig. 3C, Unpaired Student's t-test,
 10 $p<0.05$).

11 We next tested whether restraint stress could override homeostatic feeding
 12 in animals that were provided with food after an overnight fast. We found that
 13 fasted mice that had also been exposed to restraint stress for 1 hour consumed
 14 significantly less food during refeeding than controls (Fig. 3D, Unpaired Student's
 15 t-test, $*p<0.05$), indicating that acute stress can override the hyperphagia that
 16 follows a fast. Consistent with this, chemogenetic activation of Nts neurons in the
 17 LS in mice also suppressed food intake in mice that were refed after that a 24h
 18 fast (Fig. 3E, Unpaired Student's t-test, $*p<0.05$).

19 To investigate whether inactivation of Nts neurons could have an opposite
 20 effect in food intake and stimulate consumption, we injected Cre conditional AAV
 21 expressing the inhibitory hM4Di receptor or a control AAV expressing mCherry into
 22 the LS of Nts-cre mice. Acute inhibition of LS^{NTS} neurons led to an increase in food
 23 intake after CNO injection compared to mCherry controls in the first 2hrs (Fig. 3F,

Two-way ANOVA with Bonferroni post-hoc, *** $p < 0.001$) and at 8 and 24h (Figure S1H). As expected, food intake was unchanged after saline-injection into mice expressing the inhibitory DREADD (Figure S1G). Chronic inhibition of Nts neurons in the LS for 3 days produced an increase in food intake that was significant in the 3rd day (Fig. 3G, Two-way ANOVA with Bonferroni post-hoc, * $p < 0.05$) but did not increase body weight (Fig. 3H, Unpaired Student's t-test, $p = 0.83$). These results suggest that inhibition of Nts neurons increases food intake.

Interestingly, LS^{NTS} chemogenetic activation did not alter behavior during open field thigmotaxis, general locomotion (Figure S2A-C), when animals were in an open-field with novelty (Figure S2D), elevated plus maze (Figure S2E-G), or during novelty-suppressed feeding (Figure S2H). Thus, the data show LS^{NTS} neurons do not appear to play a discernible role in the other anxiogenic behaviors that were tested and appear to be specific for feeding.

LS^{NTS} neurons are selectively tuned to active escape

We next set out to assay the dynamics of LS^{NTS} activation in response to stress using in vivo calcium imaging. We injected AAVs expressing a Cre-dependent GCaMP6 into the LS of NTS-Cre mice and recorded activity using *in vivo* fiber photometry in freely behaving animals while time-locking behavior with changes in calcium signals (Gunaydin et al., 2014; Fig. 4 and Figure S3A). GCaMP6s expression and optic fiber placement were confirmed by immunohistochemistry following the experiments (Figure S3B). We first assessed activity during acute restraint stress by manually restraining mice for a 1-minute

1 period. Based on the c-fos data (Fig. 1), we expected to see an increase in calcium
 2 activity while the animal was being actively restrained. Surprisingly, we did not see
 3 a change in the calcium signal when the animal was immobile. Rather, we only
 4 found increases in the calcium signal when animals were actively struggling to
 5 escape immediately before and after the manual restraint. This reduced neural
 6 activity was seen either when the animals were completely immobilized or when
 7 they had ceased struggling (Fig. 4A, one-way ANOVA, multiple comparisons with
 8 Bonferroni post-hoc, $*p < 0.05$). We then recorded LS^{NTS} calcium signals in animals
 9 that were exposed to a one-minute tail suspension test during which time the
 10 animals actively try to escape and, consistent with the previous results, we also
 11 observed activation of LS^{NTS} neurons while the animals struggled (Fig. 4B, Paired
 12 Student's t-test, $*p < 0.05$).

13 We thus considered the possibility that LS^{NTS} neurons are specifically
 14 activated by stressful situations where the animal has an ability to escape i.e;
 15 stress-induced flight. To test this further, we exposed mice to a simulated
 16 “predator” using a remote-controlled robotic spider (Fig. 4C, left panel). In this
 17 paradigm, mice are habituated to an open field arena containing a homemade
 18 opaque acrylic nest that serves as a protected “safe zone”. The remote-controlled
 19 robotic spider was placed into the arena and after 5 min of habituation the spider
 20 movement was manually controlled so that it only moved when mice left the acrylic
 21 nest. We observed that forward movement of the robotic spider toward to the
 22 mouse induced a rapid retreat leading the mice to immediately return to the inside
 23 the acrylic nest (Supplementary Video 1). We simultaneously used *in vivo* fiber

1 photometry to monitor the activity of LS^{NTS} neurons during this simulated predator
 2 attack and found that movement of the robotic spider towards the mouse led to a
 3 strong activation of LS^{NTS} neurons that persisted until the mouse returned to the
 4 nest (Fig. 4C, Paired Student's t-test, * $p < 0.05$). Similarly, we observed a rapid and
 5 sustained increase in the calcium signal when the nest was removed and mice
 6 were constantly exposed to the robotic spider (Fig. 4D, Paired Student's t-test,*
 7 $p < 0.05$). In aggregate, these results show that LS^{NTS} neurons are activated by
 8 stressful stimuli in which the animal's response includes active movement such as
 9 struggling or flight (active coping strategies) and further suggest that they are not
 10 activated by stressful stimuli associated with immobility or freezing i.e; passive
 11 coping responses.

12 To further confirm the tuning of LS^{NTS} to stress stimuli that involved active
 13 coping, we monitored the activity of LS^{NTS} neurons in mice during a contextual fear
 14 conditioning protocol (CFC), in which mice are trained to associate a novel
 15 environment with an unpredicted foot shock. This task enables a comparison of
 16 different types of stress responses because during the training phase, the foot
 17 shock leads to active movement away from the shock source, while during testing,
 18 exposure to the contextual cue results in freezing behavior (Figure S3G and H).
 19 Moreover, unlike restraint, which is imposed by the experimenter, freezing is self-
 20 initiated. Consistent with the prior results, we found that during the training session,
 21 LS^{NTS} calcium signals were increased following the foot shock, time-locked with
 22 movement, as the mice jumped away from the shock (Fig. 4E, Paired Student's t-
 23 test, $p < 0.05$). However, during the recall test the following day, when mice

exhibited freezing behavior after being placed in the chamber that they associated with a foot shock, the LS^{NTS} calcium signal was significantly decreased compared to the period before the onset of freezing or compared to the signal during the active movement exhibited during the foot shock (Fig. 4F, Paired Student's t-test, $p < 0.05$). Taken together, these data are consistent with the possibility that the activity of these neurons is tuned to active, but not passive, responses to stressful stimuli. They also indicate that these neurons remain active until the animal is in an environment that is either perceived to be "safe" or until the animal freezes as part of a passive coping strategy. In contrast, we did not find any significant temporal correlation between calcium signals in LS^{NTS} neurons and feeding (Figure S3C and D) or voluntary movement in an open field arena (Figure S3E and F).

Molecular Profiling of LS^{NTS} neurons reveal the expression of a diversity of molecules and receptors

The majority of the neurons in the LS are GABAergic and a number of neuropeptides and specific receptors are expressed in this region in addition to neurotensin (Lein et al., 2007). In order to identify possible inputs that regulate LS^{NTS} neurons, we molecularly profiled LS^{NTS} neurons by injecting a Cre-dependent adeno-associated virus expressing an GFP-tagged ribosomal protein (AAV-Introvert-EGFP10a) into the LS of NTS-Cre mice (Nectow et al., 2017). Polysomes were immunoprecipitated with an anti-GFP antibody to profile LS^{NTS} neurons using viral-based translating ribosome affinity purification followed by RNA-seq (viralTRAP; Fig. 5A). The enrichment for each gene was calculated as

1 the number of reads in the immunoprecipitated RNA (IP) relative to the total input
 2 RNA (IP/INP; Fig 5B). Several genes were significantly enriched in the
 3 immunoprecipitated fraction (IP) including Gfp, somatostatin Sst, as well as Glp1r,
 4 Cartpt and Mc3r (Fig. 5B and C, Table S3). We confirmed the co-expression of
 5 these transcripts with Nts using qPCR of the precipitated RNAs (Fig. 5C) and *in*
 6 *situ* hybridization which showed that ~80% of septal Nts neurons co-express
 7 Slc32a1 (vgat), a marker of GABAergic neurons, (Figure 5D, Unpaired Students'
 8 t-test, **p<0.01). We also verified co-expression of Nts with Glp1r, Sst, Cartpt and
 9 Mc3r with 70% of LS^{NTS} neurons expressing Glp1r, 50% expressing Sst, 40%
 10 expressing Cartpt and 20% expressing Mc3r (Fig. 5E-I).

11 Many of the genes co-expressed with Nts in LS^{NTS} neurons are known to
 12 regulate feeding behaviors in other brain areas, including Cartpt (Kristensen et al.,
 13 1998), Glp1r (Kanoski et al., 2016), Mc3r (Marks et al., 2006) and Sst (Luo et al.,
 14 2018). For this reason, we asked whether activation of neurons expressing these
 15 genes could replicate the effects of LS^{NTS} neuronal activation. Similar to
 16 chemogenetic activation of LS^{NTS} neurons, activation of Glp1r neurons within the
 17 LS with the Glp1r agonist exendin-4 (Fig.5J) resulted in a robust decrease in food
 18 intake after 2, 12 and 24h (Figure S4A-C). In addition, chemogenetic activation of
 19 Glp1r neurons by injecting the activatory hM3Dq DREADD into the LS of Glp1R-
 20 cre mice also reduced food intake (Two-way ANOVA with Bonferroni post-hoc,
 21 *p<0.05, ****p<0.0001). Neither control mice expressing the DREADD but injected
 22 with saline nor mice injected with mCherry-expressing AAVs and given CNO
 23 exhibited any change in food intake (Fig. 5K and L; Figure S4A-C). In contrast,

1 food intake was unchanged after infusion of the Mc3r agonist (gamma-melanocyte-
2 stimulating hormone; y-msh) into the septum or activation of Cartpt or Sst neurons
3 by injecting an activatory DREADD (hM3Dq) into Cartpt-cre and Sst-cre animals
4 (Figure S4D-F). Activation of these other LS^{NTS} subpopulations (Figure S4G-I)
5 also had no effect on anxiety as assessed using an elevated plus maze (Figure
6 S4J-L).

7 Glp-1 signaling has been shown to elicit an anorectic effect and blocking of
8 Glp1r signaling using specific antagonists has been shown to blunt stress-induced
9 anorexia (Terril et al., 2018). After acute restraint stress, we observed significantly
10 increased expression of *Fos* only in LS^{NTS} neurons that also co-express *Glp1r*,
11 while *Fos* expression was unchanged in *Glp1r* neurons that do not express *Nts*
12 (Fig. 5M, Two-way ANOVA with Bonferroni post-hoc, *p<0.01). Furthermore,
13 infusion of Exendin-4 (ex4), a Glp1r agonist, into the LS increased *Fos* expression
14 in ~65% of LS^{NTS} neurons co-expressing *Glp1r* (Fig. 5N, Two-way ANOVA with
15 Bonferroni post-hoc, ***p<0.0001), but not in *Nts* neurons that do not co-express
16 *Glp1r* (Fig. 5K). These results identify a stress-responsive subpopulation of LS^{NTS}
17 neurons co-expressing Glp1r that negatively regulates food intake and further
18 shows that Glp-1 (glucagon-like peptide-1) signaling to this subpopulation of LS^{NTS}
19 neurons regulates their activity.

20

21 **Mapping the downstream projections of LS^{NTS} neurons**

22 Finally, we set out to map the functional axonal projections of LS^{NTS} neurons
23 by injecting a cre-dependent AAV expressing mCherry into the LS of NTS-Cre

1 mice. We found dense axonal projections of these neurons in the hypothalamus,
2 in particular the lateral hypothalamus (LH, Fig.6A). We also confirmed this
3 projection by injecting a Cre-dependent anterograde herpes simplex virus,
4 H129ΔTK-tdTomato into the LS of NTS-Cre mice (Fig. 6B). A similar LS to LH
5 projection was identified when mapping axonal projections from Glp1r-cre mice
6 expressing a Cre-dependent mCherry from an AAV injected into the LS (Figure
7 S5A). The LH is a well-established site regulating food intake (Morrison et al.,
8 1958), and we then tested whether specific activation of an $LS^{NTS} \rightarrow LH$ projection
9 would lead to a similar decrease in food consumption as activation of LS^{NTS} cell
10 bodies. We injected a Cre-dependent AAV expressing a channelrhodopsin (Chr2)
11 into the LS of Nts-Cre mice and implanted a fiber above the LH (Figure S5B).
12 Optogenetic activation of this pathway with blue light led to a significant increase
13 in c-fos expression in the LS^{NTS} neurons (~181 c-fos+ neurons for Chr2 vs. ~8.5
14 for mCherry) and a subset of neurons in the LH (~128 c-fos+ neurons for Chr2 vs.
15 ~35 for mCherry, Figure S5C and D, Unpaired Students t-test, ** $p < 0.01$). We then
16 used an 1-h OFF-ON-OFF protocol to evaluate food intake before, during and after
17 light stimulation of LS^{NTS} terminals in the LH and observed a reversible and
18 significant decrease of food intake in mice expressing Chr2 compared to controls
19 expressing mCherry (Fig. 6C, Two-way ANOVA with post-hoc Bonferroni
20 correction, * $p < 0.05$). Furthermore, similar to the effect of activating the entire LS^{NTS}
21 population of neurons, activation of the LH-projecting LS^{NTS} neurons led to a ~50%
22 decrease in food intake compared to controls (Fig. 6C, right panel, Unpaired
23 Student's t-test). We also evaluated whether the activation of LH-projecting LS^{NTS}

1 neurons had any effect on locomotion or anxiety and found no differences during
2 open field or elevated plus maze tests (Figure S5E-G).

3

4 **DISCUSSION**

5 Stress elicits a pleiotropic set of adaptive physiologic and behavioral
6 responses enabling animals to rapidly respond to threatening situations. These
7 responses include a suppression of other behaviors such as feeding while animals
8 initiate defensive behaviors. Accordingly, the inhibition of consummatory behaviors
9 when animals are threatened decreases an animal's vulnerability to predation
10 (Petrovich et al., 2009; Kunwar et al., 2015). While acute stress is generally
11 adaptive, chronic stress can increase the risk of maladaptive behaviors and
12 psychiatric disorders including anorexia nervosa. We previously found that
13 neurons in the lateral septum, which is known to mediate the response to stress,
14 can also regulate food intake (Azevedo et al., 2019). We thus considered the
15 possibility that neurons in the LS might link stress to anorexia. We used an
16 unbiased screen to identify molecular markers of neurons that are activated by
17 stress in the LS and identified a novel population of neurons in the LS that co-
18 express of Nts and Glp1r (LS^{NTS} neurons). We further showed that these neurons
19 are activated by active coping in response to stress and that they can suppress
20 food intake and body weight via a projection to the LH. These findings thus identify
21 a molecularly defined neural circuit that responds to stressful stimuli and in turn
22 specifically regulates feeding, but not other components of the stress response.

23 These data are consistent with a number of prior studies suggesting that the

1 LS plays a role in the active coping response to stress (Singewald et al., 2011;
 2 Bondi et al., 2007) though the cell types in the LS that mediate this response had
 3 not been identified. In our study, we show a role for LS^{NTS} neurons in this response
 4 and also characterized the calcium dynamics of their response using *in vivo*
 5 recordings. Consistent with a role in active coping, we found that these neurons
 6 are specifically activated only in response to stressful stimuli associated with active
 7 movement (i.e; flight) and not with stress associated with freezing/immobility.
 8 LS^{NTS} neurons were activated during active escape in a predator-attack paradigm,
 9 during tail suspension and during the startle response to a foot shock. In contrast,
 10 the immobility associated with complete restraint or freezing after fear conditioning
 11 did not lead to activation of these neurons. This finding was initially surprising
 12 because our prior results, which mapped c-fos, had shown activation following
 13 restraint stress. However, the temporal activation of c-fos is limited and did not
 14 allow us to discern whether the LS^{NTS} neurons were activated while the animal was
 15 moving or completely immobile. Furthermore, in the initial studies using c-fos as a
 16 marker for neural activation, mice were placed in flexible restraint cones which do
 17 not completely immobilize them and allows for movement/struggle during the
 18 period of restraint. In contrast, in our *in vivo* calcium recording studies, mice were
 19 manually restrained under the experimenter's hand and we observed that mice did
 20 not actively struggle during this type of restraint, raising the possibility that manual
 21 restraint either does not cause significant stress or that the stress this elicits acts
 22 independently of LS^{NTS} neurons.

23 LS^{NTS} neurons co-express Glp1r are stress-responsive and consistent with

1 our results, a previous report showed that infusion of a Glp1r antagonist (Ex9)
 2 directly into the LS diminished the anorexigenic effects of stress, suggesting that
 3 Glp signaling in LS contributes to this response (Terril et al., 2018). The LS is
 4 innervated by pre-proglucagon (PPG) neurons in the brainstem expressing Glp-1
 5 (Trapp and Cork, 2015) and these neurons have also been shown to be activated
 6 by stress (Holt et al., 2019). In aggregate, these data suggest that LS^{NTS} neurons
 7 that expresses Glp1r provide a link between brainstem circuits conveying stress to
 8 key output pathways from brainstem and perhaps elsewhere to regulate feeding.
 9 Further studies mapping the inputs to LS^{NTS} neurons will be necessary to confirm
 10 this possibility.

11 LS^{NTS} neuronal activation decreases food intake suggesting that cells may
 12 be tuned by active coping mechanisms that dispense with voluntary food intake
 13 until appropriate countermeasures to mitigate the stressful event have been
 14 enacted. For example, this circuit would be useful for suppressing the urge to
 15 forage when predators are in the vicinity, thus serving an important evolutionary
 16 function. Consistent with this, activation of LS^{NTS} neurons suppresses food intake
 17 even in animals that have been food deprived. If true, neural plasticity and/or
 18 genetic alterations within this circuit (e.g. by mutations in the Nts gene [Lutter et
 19 al., 2017] or changes in “allostatic load” (McEwen and Akil, 2020) could contribute
 20 to persistent maladaptive eating strategies, such as those observed in eating
 21 disorders or in other cases obesity. Presumably other neural circuits mediate the
 22 effects of stressful stimuli that lead to freezing raising the possibility that different
 23 circuits have evolved to process different types of stress, and that different types

1 of chronic stress might be associated with different disorders.

2 Indeed, a strong association between stress and reduced food intake has
3 been noted in patients with eating disorders, such as Anorexia Nervosa, which has
4 the highest mortality rate of any psychiatric disorder and is more prevalent in
5 females (Treasure et al., 2020; Kaye, 2008). Moreover, Anorexia Nervosa typically
6 manifests during puberty, a critical period characterized by increased levels of
7 estrogen (Kaye, 2008; Ma et al., 2019). Consistent with an effect of sex hormones
8 on this circuit, we found that the activity of LS^{NTS} neurons is increased in stressed
9 females during estrus compared to stressed females in diestrus or compared to
10 males. This suggest that Nts expression in the LS is tightly regulated by estrogen
11 and that both stress and high levels of estrogen are necessary for maximal neural
12 activation. While Nts expression has previously been shown to be regulated by
13 estrogen in the preoptic area of the hypothalamus, a brain area responsible for
14 maternal behavior, Nts neurons in this region have not been shown to regulate
15 feeding (Alexander et al., 1994).

16 Additional evidence suggesting a possible link between LS^{NTS} neurons and
17 anorexia nervosa has recently been provided in recent WGS (whole genome
18 sequencing) and GWAS (gene-wide association studies) studies of subset of
19 female patients with AN who report restrictive eating behaviors (Lutter et al., 2017;
20 Watson et al., 2019). In particular, one WGS study (Lutter et al., 2017) reported
21 linkage of polymorphisms in Glp1r and Nts to AN. In another GWAS studies
22 (Pinheiro et al., 2010), seven genes were identified that we also find as marking
23 LS neurons that are activated by stress (see Table S1 and Table S2) including

1 *Comt*, *Rgs10*, *Pomc*, *Crh*, *Calm1*, *Gnas*, *Esrra* . Lastly, the most recent GWAS
 2 study for anorexia nervosa (Pinheiro et al., 2010) identified 8 significant loci and 4
 3 genes of particular interest. Of these, 2 genes (*Cadm1* and *Mgmt*) and another
 4 that was less strongly implicated (*Gpr27*) were significantly increased in Nts
 5 neurons according to our transcriptomic data (Table S1) and one (*Foxp1*) was
 6 significantly decreased (Table S1). While it is not yet known whether these
 7 additional genes can also regulate feeding behavior, these data add further
 8 evidence linking the activity of LS^{NTS} neurons to anorexia and possibly eating
 9 disorders.

10 If as suggested by these results, LS^{NTS} neurons play a role in the
 11 pathogenesis of anorexia nervosa, reducing their activity pharmacologically could
 12 be of therapeutic benefit. Thus, it is possible that pharmacologically targeting Glp-
 13 1 signaling in patients with eating disorders, or other means for regulating the
 14 activity of LS^{NTS} neurons, could provide important clinical insights and possibly
 15 even therapeutic effects. Further studies will be needed to assess this possibility,
 16 including analyses of the effect of modulating neurons in animal models of
 17 anorexia nervosa, though it is unclear to what extent rodent models of this disorder
 18 fully recapitulates the features of the human condition. It has also been suggested
 19 that projections from the limbic system to the hypothalamus may mediate the
 20 psychological aspects of body weight preoccupation and reduction of feeding in
 21 AN (Donohoe et al., 1994), but the identity of these potential afferents had not been
 22 characterized. Our finding that LS^{NTS} neurons reduce food intake via projections
 23 to the LH represents another potential link to eating disorders. Numerous other

1 studies have also established connections between the lateral septum and the
2 lateral hypothalamus to control food intake (Azevedo et al., 2019; Sweeney and
3 Yang, 2016) as well as other behaviors including cocaine-induced locomotion and
4 reward (McGlinchey et al., 2016; Reddy et al., 2016).

5 In summary, these data suggest that LS^{NTS} neurons play an important
6 integratory function linking stress to anorexia and serving as an anatomical bridge
7 between the limbic system and the hypothalamus. The identification of these
8 neurons thus provides an entry point for understanding how complex inputs - in
9 this case food deprivation vs. danger - compete to generate an appropriate
10 adaptive response, in some cases potentially leading to a pathologic response in
11 the case of eating disorders. These and other data further (Azevedo et al., 2019;
12 Newmeyer et al., 2019; Land et al., 2014) suggest that higher-order, limbic brain
13 regions integrate diverse sensory information to modulate subcortical feeding
14 circuits in health and possibly disease.

15

16 **MATERIALS AND METHODS**

17

18 **Animals and breeding.**

19 C57BL/6J (Stock# 000664), NTS-Cre (Stock# 017525), Sst-Cre (Stock# 028864),
20 Cartpt-Cre (Stock# 028533), and Glp1r-cre (Stock# 029283) mice were obtained
21 from Jackson Laboratories and housed according to the guidelines of the
22 Rockefeller's University Animal Facility. Males and females at age 8-20 weeks
23 were used throughout this study and all animal experiments were approved by the

1 Rockefeller University IACUC, according to NIH guidelines. Male and female mice
2 were used for NTS-cre, Sst-cre, Cartpt-Cre and Glp1r-cre experiments. In
3 experiments that used Cre⁺ and Cre⁻ mice, littermates were used. No differences
4 were determined based on gender. Mice were kept on a 12-h/12-h light/dark cycle
5 (lights on at 7:00 a.m.) and had access to food and water ad libitum, except when
6 noted otherwise. All feeding experiments used standard rodent chow pellets.

7 **Stereotaxic Injections.**

9 All AAVs used in this study were purchased from UNC Vector Core or Addgene,
10 except where noted. AAV5-DIO-hM3Dq-mCherry or AAV5-DIO-Chr2-mCherry
11 were used for activation studies. AAV5-DIO-hM4Di-mCherry was used for
12 inhibition studies. Control AAV5-DIO-mCherry virus was used for comparison and
13 for terminal projection mapping. AAV5-DIO-GCaMP6s was used for fiber
14 photometry studies. AAV5-Introvert-GFPL10a was a gift from Alexander Nectow.
15 H129ΔTK-TT virus was a gift from David Anderson and was used as described
16 previously (Lo and Anderson, 2011) Mice were anesthetized with isoflurane,
17 placed in a stereotaxic frame (Kopf Instruments) and bilaterally injected in the
18 lateral septum using the following coordinates relative to bregma: AP: +0.1; ML:
19 0.0, DV: -3.00 (Paxinos). A total of 400 nL of virus at high titer concentrations (at
20 least 10¹¹) were injected per site at a rate of 100 nL/min. For pharmacological
21 experiments, cannulae were implanted in the LS at the same coordinates as
22 above. For fiber photometry experiments, optic fiber implants (Doric) were inserted
23 at the same coordinates as above. For optogenetics experiments, optic fiber

1 implants (Thor Labs) were implanted bilaterally over the LH at the following
 2 coordinates relative to bregma: AP: -1.80 mm; ML: + 1.50 mm, DV: -5.25 mm at a
 3 10° angle (Paxinos). Implants were inserted slowly and secured to the mouse skull
 4 using two layers of Metabond (Parkell Inc). Mice were singly housed and monitored
 5 in the first weeks following optic fiber implantation. For all surgeries, mice were
 6 monitored for 72h to ensure full recovery and two weeks later mice were used in
 7 experiments. Following behavioral procedures, mice were euthanized to confirm
 8 viral expression and fiber placement using immunohistochemistry.

9

10 **Restraint Stress**

11 Except where noted for fiber photometry studies, restraint stress consisted of 1
 12 hour of immobilization in a decapicone (Braintree Scientific) in a separate room
 13 adjacent to the housing room. Mice were returned to their home cage following
 14 restraint for further food intake measurements or until euthanasia. For in vivo fiber
 15 photometry, restrain stress was given manually by the experimenter during
 16 recordings.

17

18 **Fiber photometry**

19 Mice were acclimated to patch cables for 5 min for 3 days before experiments.
 20 Analysis of the signal was done using fiber photometry system and processor
 21 (RZ5P) from TDT (Tucker-Davis Technologies), which includes the Synapse
 22 software (TDT). Post-recording analysis was performed using custom-written
 23 MATLAB codes. The bulk fluorescent signals from each channel were normalized

1 to compare across animals and experimental sessions. The 405 channel was used
2 as the control channel. GCaMP6s signals that are recorded at this wavelength are
3 not calcium-dependent; thus, changes in signal can be attributed to
4 autofluorescence, bleaching, and fiber bending. Accordingly, any fluctuations that
5 occurred in the 405 control channels were removed from the 465 channel before
6 analysis. Change in fluorescence (ΔF) was calculated as (465 nm signal – fitted
7 405 nm signal), adjusted so that a $\Delta F/F$ was calculated by dividing each point in
8 ΔF by the 405-nm curve at that time. Z-score, plot traces and colormaps were
9 calculated for all experiments using MATLAB. Behavioral variables were
10 timestamped in the signaling traces via the real-time processor as TTL signals from
11 Noldus (Ethovision) software. This allowed for precise determinations of the
12 temporal profile of signals in relation to specific behaviors. Behaviors were tested
13 as follows. First, we tested response to food. Mice were fasted overnight and then
14 placed into a clean home cage with food pellets introduced 5 minutes later. Food
15 approach and consumption were scored. We then tested restraint by manually
16 restraining mice for 5 minutes. Tail suspension was conducted by suspending mice
17 by their tails for 5 minutes. To examine the role of predator escape, mice were
18 placed into an open field that contained a designated safe zone consisting of an
19 acrylic “nest”. After 5 minutes, a remote-controlled spider (Lattburg) was
20 introduced into the open field and began to pursue the mouse at random intervals.
21 After 10-15 minutes the nest was removed. We scored times when the mouse fled
22 from the spider as well as the removal of the nest. Locomotion was scored during

1 the first 5 minutes in the open field arena. All behavior was manually scored by a
2 second investigator who did not conduct the photometry experiment.

3

4 **Chemogenetic and Pharmacological Feeding Experiments.**

5 For acute feeding, mice were single housed and injected with either saline or
6 clozapine-N-oxide (CNO; Sigma) at 1 mg/kg intraperitoneally (i.p), exendin-4
7 (Tocris, 1 mg/mL), or γ -msh (Tocris, 150 nM) directly into the lateral septum before
8 the dark phase and cumulative food intake and body weight was measured at
9 selected time periods. For chronic feeding, mice were single or double injected
10 (every 12h) with CNO or saline before dark phase and food intake and body weight
11 were measured. For experiments that require fasting, mice were single housed
12 and food was removed before dark phase (16-23h of fasting in total). Cumulative
13 food intake at each time point was calculated for analysis.

14

15 **Elevated Plus Maze**

16 Mice were placed at the center of a cross-shaped, elevated maze in which two
17 arms are closed with dark walls and two arms are open and allowed to explore for
18 10 min. Mice were injected with CNO (1mg/kg) 1h before testing. After, mice were
19 returned to their home cage and the maze floor was cleaned in between subjects.
20 All subjects were recorded using a camera and behavior (time spent in open and
21 closed arms, distance and velocity) were analyzed using Ethovision 9.0 (Noldus).

22

23 **Open Field / Open Field with Novelty**

1 Mice freely explored an open field arena (28 x 28 cm), divided into center and
2 border regions, for 5 min. The time spent (in seconds) in the center area, were
3 taken as measures of anxiety. To increase the anxiolytic nature of the task, a novel
4 object was placed into the center of the arena (open field with novelty). Mice were
5 injected with CNO (1mg/kg) 1h before testing. After, mice were returned to their
6 home cage and the maze floor was cleaned in between subjects. All subjects were
7 recorded using a camera and behaviors (time spent in center and border, distance
8 and velocity) were analyzed using Ethovision 9.0 (Noldus).

9

10 Optogenetics Feeding Experiment.

11 Mice were previously habituated to patch cables for 3 days before experiments.
12 Implanted optic fibers were attached to a patch cable using ceramic sleeves
13 (Thorlabs) and connected to 473 nm laser (OEM Lasers/OptoEngine). Laser
14 output was verified at start of each experiment. Blue light was generated by a 473
15 nm laser diode (OEM Lasers/OptoEngine) at 5-10 mW of power. Light pulse (10
16 ms) and frequency (10Hz) was controlled using a waveform generator (Keysight)
17 to activate NT+ terminals in lateral hypothalamus. Animals were sacrificed to
18 confirm viral expression and fiber placement using immunohistochemistry. Each
19 feeding session lasted 80 min and it was divided in 1 trial of 20 min to allow the
20 animal to acclimate to the cage and 3 trials of 20 min each (1h feeding session).
21 During each feeding session, light was off during the first 20 min, on for 20 min
22 and off again for the remaining 20 min. Consumed food was recorded manually
23 before and after each session. To facilitate measurement, 3 whole pellets were

1 added to cups and food crumbs were not recorded. Feeding bouts were recorded
2 using a camera and the Ethovision 9.0 software (Noldus).

3

4 **Viral-Trap (V-Trap)**

5 Affinity purification of EGFP-tagged polysomes was done 3 or 4 weeks after virus
6 injections (Nectow et al., 2017). Briefly, mice were separated into 3 biological
7 replicate groups of 4-6 mice per group and euthanized. Brains were removed, and
8 the lateral septum was dissected on ice and pooled. Tissue was homogenized in
9 buffer containing 10 mM HEPES-KOH (pH 7.4), 150 mM KCl, 5 mM MgCl₂, 0.5
10 mM DTT, 100 µg/mL cycloheximide, RNasin (Promega) and SUPERase-In™ (Life
11 Technologies) RNase inhibitors, and Complete-EDTA-free protease inhibitors
12 (Roche) and then cleared by two-step centrifugation to isolate polysome-
13 containing cytoplasmic supernatant. Polysomes were immunoprecipitated using
14 monoclonal anti-EGFP antibodies (clones 19C8 and 19F7; see Heiman et al.,
15 2008³⁰) bound to biotinylated-Protein L (Pierce; Thermo Fisher Scientific)-coated
16 streptavidin-conjugated magnetic beads (Life Technologies). A small amount of
17 tissue RNA was saved before the immunoprecipitation (Input) and both input and
18 immunoprecipitated RNA (IP) were then purified using RNAeasy Mini kit
19 (QIAGEN). RNA quality was checked using an RNA PicoChip on a bioanalyzer.
20 RIN values >7 were used. Experiments were performed in triplicates for each
21 group. cDNA was amplified using SMARTer Ultralow Input RNA for Illumina
22 Sequencing Kit and sequenced on an Illumina HiSeq2500 platform.

23

1 RNA sequencing and qPCR analysis.

2 RNA sequencing raw data was uploaded and analyzed using BaseSpace apps
3 (TopHat and Cufflinks; Illumina) using an alignment to annotated mRNAs in the
4 mouse genome (UCSC, Mus musculus assembly mm10). The average
5 immunoprecipitated (IP) and Input value of each enriched and depleted genes with
6 a q-value lower than 0.05 were plotted using GraphPad Prism (GraphPad). To
7 validate the RNA Sequencing data, qPCR using predesigned Taqman probes
8 (idtDNA) were used and cDNA was prepared using the QuantiTect Reverse
9 Transcription Kit (Life Technologies). The abundance of these genes in IP and
10 Input RNA was quantified using Taqman Gene Expression Master Mix (Applied
11 Biosystems). Transcript abundance was normalized to beta-actin. Fold of Change
12 were calculated using the $\Delta\Delta C_t$ method.

13

14 Immunohistochemistry, quantifications and imaging.

15 Mice were perfused, and brains were postfixed for 24h in 10% formalin. Brain
16 slices were taken using a vibratome (Leica), blocked for 1h with 0.3% Triton X-
17 100, 3% bovine serum albumin (BSA), and 2% normal goat serum (NGS) and
18 incubated in primary antibodies for 24h at 4°C. Then, free-floating slices were
19 washed three times for 10 min in 0.1% Triton X-100 in PBS (PBS-T), incubated for
20 1h at room temperature with secondary antibodies, washed in PBS-T and mounted
21 in Vectamount with DAPI (Southern Biotech). Antibodies used here were: anti-c-
22 fos (1:500; Cell Signaling), anti-mCherry (1:1000; Abcam) goat-anti-rabbit (Alexa
23 488 or Alexa 594, 1:1000; Molecular Probes), goat anti-chicken Alexa488 or

1 Alexa594 (1:1000; Molecular Probes). Images were taken using Axiovert 200
2 microscope (Zeiss) or LSM780 confocal (Zeiss) and images were processed using
3 ImageJ software (NIH, Schneider et al., 2012). C-fos counts were conducted for 3
4 or more sections / animal and averaged for each animal for statistical analysis.

6 **Fluorescent In Situ Hybridization**

7 For examination of gene expression and Fos experiments, tissue samples
8 underwent single molecule fluorescent in situ hybridization (smFISH). Isoflurane
9 anesthetized mice were decapitated, brain harvested and flash frozen on
10 aluminum foil on dry ice. Brains were stored at -80°C . Prior to sectioning, brains
11 were equilibrated to -16°C in a cryostat for 30 min. Brains were cryostat sectioned
12 coronally at $20\text{ }\mu\text{m}$ and thaw-mounted onto Superfrost Plus slides (25x75 mm,
13 Fisherbrand). Slides were air-dried for 60 to 90 min prior to storage at -80°C .
14 smFISH for all genes examined— *Fos*, *Nts*, *Sst*, *Glp1r*, *Cartpt*, *Mc3r* - was
15 performed using RNAscope Fluorescent Multiplex Kit (Advanced Cell Diagnostics)
16 according to the manufacturer'. Slides were counterstained for the nuclear marker
17 DAPI using ProLong Diamond Antifade mounting medium with DAPI (Thermo
18 Fisher). Sections were imaged using an LSM780 confocal (Zeiss) and processed
19 using ImageJ software. Counts were conducted for 2 or more sections/animal and
20 averaged for each animal for statistical analysis.

22 **Statistical analysis**

23 All results are presented as mean $\pm$ SEM. and were analyzed with Prism software

1 or with Matlab. No statistical methods were used to predetermine sample sizes,
 2 but our sample sizes are similar to those reported in previous publications.
 3 Normality tests and F tests for equality of variance were performed before
 4 choosing the statistical test. Unless otherwise indicated, for feeding experiments
 5 we used a two-way ANOVA with Bonferroni correction to analyze statistical
 6 differences. $P < 0.05$ was considered significant (* $p < 0.05$, ** $p < 0.01$, *** $P < 0.001$,
 7 **** $p < 0.0001$). Image quantifications were analyzed using unpaired Students' t-
 8 tests. Fiber photometry data were analyzed using paired Students' t-tests. Animals
 9 in the same litter were randomly assigned to different treatment groups and blinded
 10 to experimenters in the various experiments. Injection sites and viral expression
 11 were confirmed for all animals. Mice showing incorrect injection sites or optic fiber
 12 placement were excluded from the data analysis.

13

14 **ACKNOWLEDGEMENTS**

15 We thank Ravi Tolwani and the staff of the Comparative Biosciences Center,
 16 Connie Zhao and the staff of the Genomics Resource Center, Alison North and the
 17 staff of the Bioimaging Resource Center at Rockefeller University for technical
 18 assistance. We thank Alexander Nectow for the AAV-Introvert-GFPL10a virus and
 19 David Anderson for providing the HSV-IsI-tdTomato virus. We thank Christin Kosse
 20 for providing MATLAB codes for photometry analysis. This work was funded by a
 21 Long Term HSFP Fellowship (E.P.A.), F32DK107077 (S.A.S.), NARSAD Young
 22 Investigator Awards from the Brain and Behavior Research Foundation (E.P.A and
 23 S.A.S) the JPB Foundation and the Klarman Foundation (J.M.F).

1 **COMPETING INTEREST**

2 There is no conflict of interest.

3

4 **REFERENCES**

5 Alexander, M.J. and Leeman, S.E. (1994). Estrogen-inducible neurotensin
6 immunoreactivity in the preoptic area of the female rat. *J Comp Neurol.* 345, 496-
7 509.

8 Anthony T.E., Dee N., Bernard A., Lerchner W., Heintz N., Anderson D.J. (2014).
9 Control of Stress-Induced Persistent Anxiety by an Extra-Amygdala
10 Septohypothalamic Circuit. *Cell* 156, 522–536.

11 Armbruster BN, Li X, Pausch MH, Herlitze S, Roth BL. (2007). Evolving the lock to
12 fit the key to create a family of G protein-coupled receptors potently activated by
13 an inert ligand. *Proc Natl Acad Sci U S A.* 104, 5163-8.

14 Azevedo E. P., Pomeranz L, Cheng J, Schneeberger M, Vaughan R, Stern SA,
15 Tan B, Doerig K, Greengard P, Friedman JM. (2019). A Role of Drd2 Hippocampal
16 Neurons in Context-Dependent Food Intake. *Neuron* 102, 873–886.e5.

17 Bondi C.O., Barrera G, Lapiz MD, Bedard T, Mahan A, Morilak DA. (2007).
18 Noradrenergic facilitation of shock-probe defensive burying in lateral septum of
19 rats, and modulation by chronic treatment with desipramine. *Prog.*
20 *Neuropsychopharmacol. Biol. Psychiatry* 31, 482–495.

21 Donohoe T.P. (1984). Stress-induced anorexia: implications for anorexia nervosa.
22 *Life Sci.* 34, 203-18.

23 Guarda A.S., Schreyer C.C., Boersma G.J., Tamashiro K.L., Moran T.H. (2015).

1 Anorexia nervosa as a motivated behavior: Relevance of anxiety, stress, fear and
2 learning. *Physiol Behav.* 152, 466-72.

3 Gunaydin L., Grosenick L, Finkelstein JC, Kauvar IV, Fenno LE, Adhikari A,
4 Lammel S, Mirzabekov JJ, Airan R, Zalocusky KA, Tye KM, Anikeeva P, Malenka
5 RC, Deisseroth K. (2014). Natural neural projection dynamics underlying social
6 behavior. *Cell* 157, 1535–1551.

7 Holt, M.K., Richards, J.E., Cook, D.R., Brierley, D.I., Williams, D.L., Reimann, F.,
8 Gribble, F.M. and Trapp, S. (2019). Preproglucagon Neurons in the Nucleus of the
9 Solitary Tract Are the Main Source of Brain GLP-1, Mediate Stress-Induced
10 Hypophagia, and Limit Unusually Large Intakes of Food. *Diabetes.* 68, 21-33.

11 Kanoski, S. E., Hayes, M. R. and Skibicka, K. P. (2016) .GLP-1 and weight loss:
12 unraveling the diverse neural circuitry. *Am. J. Physiol. Regul. Integr. Comp.*
13 *Physiol.* 310, R885–95.

14 Kaye W. (2008). Neurobiology of anorexia and bulimia nervosa. *Physiol Behav.*
15 94, 121-35.

16 Knight Z.A., Tan K., Birsoy K., Schmidt S., Garrison J.L., Wysocki R.W., Emiliano
17 A., Ekstrand M.I., Friedman J.M. (2012). Molecular profiling of activated neurons
18 by phosphorylated ribosome capture. *Cell.* 151, 1126-37.

19 Kristensen P, Judge ME, Thim L, Ribel U, Christjansen KN, Wulff BS, Clausen JT,
20 Jensen PB, Madsen OD, Vrang N, Larsen PJ, Hastrup S. (1998). Hypothalamic
21 CART is a new anorectic peptide regulated by leptin. *Nature.* 393, 72-6.

22 Kubo T., Kanaya T., Numakura H., Okajima H., Hagiwara Y., Fukumori R. (2002).
23 The lateral septal area is involved in mediation of immobilization stress-induced

1 blood pressure increase in rats. *Neurosci. Lett.* 318, 25–28.

2 Kunwar P.S., Zelikowsky M., Remedios R., Cai H., Yilmaz M., Meister M.,
3 Anderson D.J. (2015). Ventromedial hypothalamic neurons control a defensive
4 emotion state. *Elife*. doi: 10.7554/eLife.06633.

5 Land, B. B., Narayanan, N.S., Liu, R.J., Gianessi, C.A., Brayton, C.E., Grimaldi,
6 D.M., Sarhan, M., Guarnieri, D.J., Deisseroth, K., Aghajanian, G.K., DiLeone, R.J.
7 (2014). Medial prefrontal D1 dopamine neurons control food intake. *Nat. Neurosci.*
8 17, 248–253.

9 Lein, E.S., Hawrylycz, M.J., Ao, N., Ayres, M., Bensinger, A., Bernard, A., Boe,
10 A.F., Boguski, M.S., Brockway, K.S., Byrnes EJ, Chen L, Chen L, Chen TM, Chin
11 MC, Chong J, Crook BE, Czaplinska A, Dang CN, Datta S, Dee NR, Desaki AL,
12 Desta T, Diep E, Dolbeare TA, Donelan MJ, Dong HW, Dougherty JG, Duncan BJ,
13 Ebbert AJ, Eichele G, Estin LK, Faber C, Facer BA, Fields R, Fischer SR, Fliss TP,
14 Frensley C, Gates SN, Glattfelder KJ, Halverson KR, Hart MR, Hohmann JG,
15 Howell MP, Jeung DP, Johnson RA, Karr PT, Kawal R, Kidney JM, Knapik RH,
16 Kuan CL, Lake JH, Laramie AR, Larsen KD, Lau C, Lemon TA, Liang AJ, Liu Y,
17 Luong LT, Michaels J, Morgan JJ, Morgan RJ, Mortrud MT, Mosqueda NF, Ng LL,
18 Ng R, Orta GJ, Overly CC, Pak TH, Parry SE, Pathak SD, Pearson OC, Puchalski
19 RB, Riley ZL, Rockett HR, Rowland SA, Royall JJ, Ruiz MJ, Sarno NR, Schaffnit
20 K, Shapovalova NV, Sivasay T, Slaughterbeck CR, Smith SC, Smith KA, Smith BI,
21 Sodt AJ, Stewart NN, Stumpf KR, Sunkin SM, Sutram M, Tam A, Teemer CD,
22 Thaller C, Thompson CL, Varnam LR, Visel A, Whitlock RM, Wohnoutka PE,
23 Wolkey CK, Wong VY, Wood M, Yaylaoglu MB, Young RC, Youngstrom BL, Yuan

- 1 XF, Zhang B, Zwingman TA, Jones AR. (2007). Genome-wide atlas of gene
2 expression in the adult mouse brain. *Nature*. 445, 168-76.
- 3 Lo, L. and Anderson, D. J. (2011). A Cre-dependent, anterograde transsynaptic
4 viral tracer for mapping output pathways of genetically marked neurons. *Neuron*
5 72, 938–950.
- 6 Luo, S.X., Huang, J., Li, Q., Mohammad, H., Lee, C.Y., Krishna, K., Kok, A.M.,
7 Tan, Y.L., Lim, J.Y., Li, H., Yeow, L.Y., Sun, J., He, M., Grandjean, J., Sajikumar,
8 S., Han, W. and Fu, Y. (2018). Regulation of feeding by somatostatin neurons in
9 the tuberal nucleus. *Science* 361, 76-81.
- 10 Lutter, M., Bahl, E., Hannah, C., Hofammann, D., Acevedo, S., Cui, H., McAdams,
11 C.J. and Michaelson, J.J. (2017). Novel and ultra-rare damaging variants in
12 neuropeptide signaling are associated with disordered eating behaviors. *PLoS*
13 *One*. 12, e0181556.
- 14 Ma, R., Mikhail, M.E., Fowler, N., Culbert, K.M., Klump, K.L. (2019). The Role of
15 Puberty and Ovarian Hormones in the Genetic Diathesis of Eating Disorders in
16 Females. *Child Adolesc Psychiatr Clin N Am*. 28, 617-628.
- 17 Marks D.L., Hruby V., Brookhart G., Cone R.D. (2006). The regulation of food
18 intake by selective stimulation of the type 3 melanocortin receptor (MC3R).
19 *Peptides*. 27, 259-64.
- 20 McEwen B.S. and Akil H. (2020). Revisiting the Stress Concept: Implications for
21 Affective Disorders. *J Neurosci*. 40, 12-21.
- 22 McGlinchey, E.M., James, M.H., Mahler, S.V., Pantazis, C., Aston-Jones, G.
23 (2016). Prelimbic to Accumbens Core Pathway Is Recruited in a Dopamine-

1 Dependent Manner to Drive Cued Reinstatement of Cocaine Seeking. *J Neurosci.*
2 36, 8700-11.

3 Morrison SD, Barnett RJ, Mayer J. Localization of lesions in the lateral
4 hypothalamus of rats with induced adipsia and aphagia. *Am J Physiol.* 193, 230-
5 4. (1958).

6 Nectow A.R., Moya MV, Ekstrand MI, Mousa A, McGuire KL, Sferrazza CE, Field
7 BC, Rabinowitz GS, Sawicka K, Liang Y, Friedman JM, Heintz N, Schmidt EF.
8 (2017). Rapid Molecular Profiling of Defined Cell Types Using Viral TRAP. *Cell*
9 *Rep.* 19, 655–667.

10 Newmyer, B. A., Whindleton, C.M., Klein, P.M., Beenhakker, M.P., Jones, M.K.,
11 Scott, M.M. (2019). VIPergic neurons of the infralimbic and prelimbic cortices
12 control palatable food intake through separate cognitive pathways. *JCI Insight* 5,
13 pii: 126283.

14 Petrovich G.D., Ross C.A., Mody P., Holland P.C. and Gallagher M. (2009).
15 Central, but not basolateral, amygdala is critical for control of feeding by aversive
16 learned cues. *J Neurosci.* 29, 15205-12.

17 Pinheiro, A.P., Bulik, C.M., Thornton, L.M., Sullivan, P.F., Root, T.L., Bloss, C.S.,
18 Berrettini, W.H., Schork, N.J., Kaye, W.H., Bergen, A.W., Magistretti, P., Brandt,
19 H., Crawford, S., Crow, S., Fichter, M.M., Goldman, D., Halmi, K.A., Johnson, C.,
20 Kaplan, A.S., Keel, P.K., Klump, K.L., La Via, M., Mitchell, J.E., Strober, M.,
21 Rotondo, A., Treasure, J. and Woodside, D.B. (2010). Association study of 182
22 candidate genes in anorexia nervosa. *Am J Med Genet B Neuropsychiatr Genet.*
23 153B, 1070-80.

1 Reddy, I.A., Pino, J.A., Weikop, P., Osses, N., Sørensen, G., Bering, T., Valle, C.,
2 Bluett, R.J., Erreger, K., Wortwein, G., Reyes, J.G., Graham, D., Stanwood, G.D.,
3 Hackett, T.A., Patel, S., Fink-Jensen, A., Torres, G.E. and Galli, A. (2016)
4 Glucagon-like peptide 1 receptor activation regulates cocaine actions and
5 dopamine homeostasis in the lateral septum by decreasing arachidonic acid levels.
6 *Transl Psychiatry*. 6, e809.

7 Schneider, C. A., Rasband, W. S. and Eliceiri, K. W. (2012). NIH Image to ImageJ:
8 25 years of image analysis. *Nat. Methods* 9, 671–675.

9 Sheehan T.P., Chambers R.A. and Russell D.S. (2004). Regulation of affect by the
10 lateral septum: implications for neuropsychiatry. *Brain Res Brain Res Rev.* 46, 71-
11 117.

12 Shimizu N., Oomura Y. and Kai Y. (1989). Stress-induced anorexia in rats
13 mediated by serotonergic mechanisms in the hypothalamus. *Physiol Behav.* 46,
14 835-41.

15 Singewald, G. M., Rjabokon, A., Singewald, N. and Ebner, K. (2011). The
16 modulatory role of the lateral septum on neuroendocrine and behavioral stress
17 responses. *Neuropsychopharmacology* 36, 793–804.

18 Stern S.A., Pomeranz L.E., Azevedo E.P., Doerig K.R., Friedman J.M. (2019). A
19 molecularly defined insular cortex --> central amygdala circuit mediates
20 conditioned overconsumption of food. Preprint at
21 <https://www.biorxiv.org/content/10.1101/684498v1>.

22 Surendran S., Hückesfeld S., Wäschle B., Pankratz M.J. (2017). Pathogen-
23 induced food evasion behavior in *Drosophila* larvae. *J Exp Biol.* 220, 1774-1780.

1 Sweeney, P. and Yang, Y. (2016). An Inhibitory Septum to Lateral Hypothalamus
2 Circuit That Suppresses Feeding. *J. Neurosci.* 36, 11185–11195.

3 Terrill, S. J., Maske, C. B. & Williams, D. L. (2018). Endogenous GLP-1 in lateral
4 septum contributes to stress-induced hypophagia. *Physiol. Behav.* 192, 17–22.

5 Trapp S. and Cork S.C. (2015). PPG neurons of the lower brain stem and their role
6 in brain GLP-1 receptor activation. *Am J Physiol Regul Integr Comp Physiol.* 309,
7 R795-804.

8 Treasure, J., Duarte, T.A., Schmidt, U. (2020). Eating disorders. *Lancet.* 395, 899-
9 911.

10 30.

11 Watson, H.J., Yilmaz, Z., Thornton, L.M., Hübel C, Coleman JRI, Gaspar HA,
12 Bryois J, Hinney A, Leppä VM, Mattheisen M, Medland SE, Ripke S, Yao S, Giusti-
13 Rodríguez P; Anorexia Nervosa Genetics Initiative, Hanscombe KB, Purves KL;
14 Eating Disorders Working Group of the Psychiatric Genomics Consortium, Adan
15 RAH, Alfredsson L, Ando T, Andreassen OA, Baker JH, Berrettini WH, Boehm I,
16 Boni C, Perica VB, Buehren K, Burghardt R, Cassina M, Cichon S, Clementi M,
17 Cone RD, Courtet P, Crow S, Crowley JJ, Danner UN, Davis OSP, de Zwaan M,
18 Dedoussis G, Degortes D, DeSocio JE, Dick DM, Dikeos D, Dina C, Dmitrzak-
19 Weglarz M, Docampo E, Duncan LE, Egberts K, Ehrlich S, Escaramís G, Esko T,
20 Estivill X, Farmer A, Favaro A, Fernández-Aranda F, Fichter MM, Fischer K,
21 Föcker M, Foretova L, Forstner AJ, Forzan M, Franklin CS, Gallinger S, Giegling
22 I, Giuranna J, Gonidakis F, Gorwood P, Mayora MG, Guillaume S, Guo Y,
23 Hakonarson H, Hatzikotoulas K, Hauser J, Hebebrand J, Helder SG, Herms S,

1 Herpertz-Dahlmann B, Herzog W, Huckins LM, Hudson JI, Imgart H, Inoko H,
2 Janout V, Jiménez-Murcia S, Julià A, Kalsi G, Kaminská D, Kaprio J, Karhunen L,
3 Karwautz A, Kas MJH, Kennedy JL, Keski-Rahkonen A, Kiezebrink K, Kim YR,
4 Klareskog L, Klump KL, Knudsen GPS, La Via MC, Le Hellard S, Levitan RD, Li
5 D, Lilienfeld L, Lin BD, Lissowska J, Luykx J, Magistretti PJ, Maj M, Mannik K,
6 Marsal S, Marshall CR, Mattingdal M, McDevitt S, McGuffin P, Metspalu A,
7 Meulenbelt I, Micali N, Mitchell K, Monteleone AM, Monteleone P, Munn-Chernoff
8 MA, Nacmias B, Navratilova M, Ntalla I, O'Toole JK, Ophoff RA, Padyukov L,
9 Palotie A, Pantel J, Papezova H, Pinto D, Rabionet R, Raevuori A, Ramoz N,
10 Reichborn-Kjennerud T, Ricca V, Ripatti S, Ritschel F, Roberts M, Rotondo A,
11 Rujescu D, Rybakowski F, Santonastaso P, Scherag A, Scherer SW, Schmidt U,
12 Schork NJ, Schosser A, Seitz J, Slachtova L, Slagboom PE, Slof-Op 't Landt MCT,
13 Slopian A, Sorbi S, Świątkowska B, Szatkiewicz JP, Tachmazidou I, Tenconi E,
14 Tortorella A, Tozzi F, Treasure J, Tsitsika A, Tyszkiewicz-Nwafor M, Tziouvas K,
15 van Elburg AA, van Furth EF, Wagner G, Walton E, Widen E, Zeggini E, Zerwas
16 S, Zipfel S, Bergen AW, Boden JM, Brandt H, Crawford S, Halmi KA, Horwood LJ,
17 Johnson C, Kaplan AS, Kaye WH, Mitchell JE, Olsen CM, Pearson JF, Pedersen
18 NL, Strober M, Werge T, Whiteman DC, Woodside DB, Stuber GD, Gordon S,
19 Grove J, Henders AK, Juréus A, Kirk KM, Larsen JT, Parker R, Petersen L, Jordan
20 J, Kennedy M, Montgomery GW, Wade TD, Birgegård A, Lichtenstein P, Norring
21 C, Landén M, Martin NG, Mortensen PB, Sullivan PF, Breen G, Bulik CM. (2019).
22 Genome-wide association study identifies eight risk loci and implicates metabo-
23 psychiatric origins for anorexia nervosa. *Nat Genet.* 51, 1207-1214.

- 1 Wong L.C., Wang L., D'Amour J.A., Yumita T., Chen G., Yamaguchi T., Chang
- 2 B.C., Bernstein H., You X., Feng J.E., Froemke R.C., Lin D. (2016). Effective
- 3 Modulation of Male Aggression through Lateral Septum to Medial Hypothalamus
- 4 Projection. *Current Biology* 26, 593–604.
- 5 Yau Y.H. and Potenza M.N. (2013). Stress and eating behaviors. *Minerva*
- 6 *Endocrinol.* 38, 255-67.

7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23

1 FIGURES

Fig 1

Figure 1. Identification of lateral septum neurons activated by acute restraint stress.
 (A) Left panel, experimental scheme. Right panel, food intake, measured in grams (g), in control (black) and RS (red) mice after 60 and 120min following RS. n=5(RS),

1 n=6(control), Two-way ANOVA with post-hoc Bonferroni correction, Time: $F(1,9)=175.9$,
2 $p<0.0001$; Subject: $F(1,9)=16.86$, $p=0.003$; Interaction: $F(1,9)=4.975$, $p=0.0527$).

3 (B) Activity mapping in the whole brain using c-fos staining in control (naïve) and restraint
4 stressed (RS) mice. Areas with the highest c-fos expression (green) difference are shown:
5 basolateral amygdala (BLA), bed nucleus of the stria terminals (BNST), insular cortex (IC)
6 and lateral septum (LS). Insets show 2x digital zoom of c-fos expression. Scale bars, 50
7 μm .

8 (C) Activity-based transcriptomics (PhosphoTrap) of the lateral septum of control (naïve)
9 and restraint stressed mice. Middle panel, plot depicting the average IP/INP value (log2)
10 of all genes analyzed in control and restraint stress (RS) samples. Enriched genes (>2-
11 fold; red) and depleted genes (<2-fold; blue) are shown, and activation or target gene
12 markers are shown: Nts (neurotensin), fos and fosb ($n = 2$). All genes depicted have q-
13 values <0.05 as calculated by Cufflinks.

14 (D) mRNA levels of Nts evaluated using Taqman qPCR in control and restraint stress (RS)
15 samples: control (black bars) and RS (red bars) are compared ($n = 6$). Unpaired Student's
16 t-test, $*p < 0.05$, $t=2.692$, $df=10$. Data are represented as mean $\pm$ SEM.

17
18
19
20
21
22
23
24
25

Fig 2

Figure 2. Nts expression is regulated by stress and the estrous cycle in females.

(A) The mouse estrous cycle: the different phases of the mouse estrous cycle (diestrus, proestrus, estrus, metestrus) are shown in different colors.

(B) In situ hybridization showing the expression of Nts (green) and fos (red) mRNA in females in estrus and diestrus exposed to acute restraint stress (RS) or naïve (control). Quantification of the number of Nts+/fos+ cells per slice/mice are shown, n=3-5, Nested one-way ANOVA, F=6.366, DFn=3, Dfd=12.

(C) In situ hybridization showing the expression of Nts (green) and fos (red) mRNA in males exposed to acute restraint stress (RS) or naïve (control). Quantification of the number of Nts+/fos+ cells per slice/mice are shown, n=6, Unpaired Student's t-test, *p<0.05, t=2.962, df=10. Data are represented as mean ± SEM.

Fig 3

Figure 3. LS^{NTS} neurons regulate food intake and body weight in mice.

(A) Left panel, experimental scheme. NTS-cre mice were injected with an AAV encoding the hM3Dq activatory DREADD receptor in the LS. Middle panel, cumulative food intake, measured in grams (g) of control mCherry (black) and hM3Dq (red) expressing NTS-cre mice after 60 and 120 min following CNO injection. n=7(mCherry), n=10(hM3Dq), Two-way ANOVA with post-hoc Bonferroni correction, Time: $F(1,15)=55.46$, $p<0.0001$; Subject: $F(1,15)=12.84$, $p=0.0027$; Interaction: $F(1,15)=5.45$, $p=0.034$).

(B) Daily food intake, measured in grams (g) daily during injections of saline (days 1-3, 7-8) or CNO (days 4-6) in control mCherry (black) or hM3Dq (red) expressing NTS-cre mice. n=8; Two-way ANOVA with post-hoc Bonferroni correction, Time: $F(7,72)=2.871$, $p=0.01$; Subject: $F(1,72)=46.92$, $p<0.0001$; Interaction: $F(7,72)=6.48$, $p<0.0001$).

1 (C) Body weight (delta), measured in grams (g) after 3 days of daily CNO injections in
2 control mCherry (black) or hM3Dq (red) expressing NTS-cre mice. n=8; Unpaired
3 Student's t-test, * $p < 0.05$, $t = 2.903$, $df = 10$.

4 (D) Left panel, experimental scheme. Right panel, cumulative food intake, measured in
5 grams (g), in control naïve mice (control, black) or mice that underwent restraint stress
6 (RS, red), measured 2h after re-feeding after an overnight fast, n=6. Unpaired Students'
7 t-test: $T(10) = 2.427$, $p = 0.0356$.

8 (E) Left panel, experimental scheme. Right panel, cumulative food intake, measured in
9 grams (g), in control mCherry (black) or hM3Dq(red) expressing NTS-cre mice, measured
10 2h following re-feeding after an overnight fast, n=8, Paired Student's t-test, * $p < 0.05$, $t =$
11 3.171 , $df = 7$.

12 (F) Left panel, experimental scheme. Right panel, cumulative food intake, measured in
13 grams (g), in control mCherry (black) and hM4Di (red) expressing NTS-cre mice after 60
14 min and 120 min following saline injection. n=7(mCherry), n=8(hM4Di). Two-way ANOVA
15 with post-hoc Bonferroni correction, Time: $F(1,19) = 415.8$, $p < 0.0001$; Subject:
16 $F(1,13) = 1.042$, $p = 0.326$; Interaction: $F(4,52) = 2.839$, $p = 0.0333$.

17 (G) Daily food intake, measured in grams (g) after daily injections of saline (days 1-3, 7-
18 8) or CNO (days 4-6) in control mCherry (black) or hM4Di (red) expressing NTS-cre mice.
19 n=7; Two-way ANOVA with post-hoc Bonferroni correction, Time: $F(7,96) = 3.949$,
20 $p = 0.0008$; Subject: $F(1,96) = 18.95$, $p < 0.0001$; Interaction: $F(7,96) = 0.876$, $p = 0.5285$.

21 (H) Body weight (delta), measured in grams (g) after 3 days of daily CNO injections in
22 control mCherry (black) or hM4Di (red) expressing NTS-cre mice. n=8; Unpaired Student's
23 t-test, $p = 0.83$, $t = 0.2168$, $df = 10$. Data are represented as mean $\pm$ SEM.

24

25

26

Fig 4

Figure 4. LS^{NTS} neurons are specifically activated by stressful situation involving active coping.

(A) Left panel, experimental scheme: mice were manually immobilized for the period of 1 min. Ca²⁺ activity was recorded before, during and after the period of manual immobilization. Middle, Average GCaMP6s z-score (black lines) and SEM (gray dotted

lines) across all recordings and time-locked to immobilization start. Right middle panel, quantification of the area under the curve of Ca^{+2} transients before, during and after immobilization. One-way ANOVA with repeated measures and post-hoc Bonferroni correction, $F(1.833, 9.165)=9.888$, $p=0.0057$. Right panel, heat maps represent the average % $\Delta F/F$ from each gCaMP6s recordings time-locked to immobilization start. $n=6$.

(B) Left panel, experimental scheme: mice were tail suspended for the period of 1 min. Ca^{+2} activity was recorded before, during and after the period of tail suspension. Middle, Average GCaMP6s z-score (black lines) and SEM (gray dotted lines) across all recordings and time-locked to tail suspension start. Right middle panel, quantification of the area under the curve of Ca^{+2} transients before, during and after tail suspension. Paired Student's t-test, $t=2.659$, $df=5$. Right panel, heat maps represent the average % $\Delta F/F$ from each gCaMP6s recordings time-locked to tail suspension start. $n=6$.

(C) Left panel, experimental scheme: mice were exposed to an open field arena containing a robotic, remote-controlled spider predator and a designated safe "nest" (black triangular hut). Ca^{+2} activity was recorded during the whole session. Middle, Average GCaMP6s z-score (black lines) and SEM (gray dotted lines) across all recordings and time-locked to the robotic spider attack. Right middle panel, quantification of the area under the curve of Ca^{+2} transients before and after the robotic spider attack. Paired Student's t-test, $t=3.507$, $df=4$. Right panel, heat maps represent the average % $\Delta F/F$ from each gCaMP6s recordings time-locked to the robotic spider attack. $n=6$.

(D) Left panel, experimental scheme: the designated safe "nest" (black triangular hut) was removed and mice that had been attacked by a robotic spider predator was exposed to the open field containing the spider. Ca^{+2} activity was recorded during the whole session. Middle, Average GCaMP6s z-score (black lines) and SEM (gray dotted lines) across all recordings and time-locked to the removal of the nest. Right middle panel, quantification

1 of the area under the curve of Ca^{+2} transients before and after the removal of the nest.
 2 Paired Student's t-test, $t=3.305$, $df=5$. Right panel, heat maps represent the average %
 3 $\Delta F/F$ from each gCaMP6s recordings time-locked to the removal of the nest. $n=6$.

4 (E) Experimental scheme for contextual fear conditioning: mice were exposed to an
 5 operant chamber where they received a mild foot shock (training) and after 24h mice were
 6 re-exposed to the operant chamber (testing). Ca^{+2} activity was recorded during training
 7 and testing.

8 (F) First panel, average GCaMP6s z-score (black lines) and SEM (gray dotted lines)
 9 across all recordings and time-locked to the foot shock received on the training day.
 10 Second panel, quantification of the area under the curve of Ca^{+2} transients before and
 11 after the foot shock. Paired Student's t-test, $t=3.230$, $df=4$. Third panel, heat maps
 12 represent the average % $\Delta F/F$ from each gCaMP6s recordings time-locked to the freezing
 13 behavior. $n=6$.

14 (G) First panel, average GCaMP6s z-score (black lines) and SEM (gray dotted lines)
 15 across all recordings and time-locked to freezing behavior on the testing day. Second
 16 panel, quantification of the area under the curve of Ca^{+2} transients before and after the
 17 freezing behavior. Paired Student's t-test, $t=2.833$, $df=8$. Third panel, heat maps represent
 18 the average % $\Delta F/F$ from each gCaMP6s recordings time-locked to the freezing behavior.
 19 $n=6$. Data are represented as mean $\pm$ SEM.

20

21

22

23

24

25

Fig 5

Figure 5. Molecular and functional profiling of LS^{NTS} neurons.

(A) Experimental scheme of the viralTrap experiment. NTS-cre mice are injected with an AAV expressing cre-dependent GFPL10 (AAV-Ef1a-FLEX-eGFPL10) into the LS and then GFP+ polysomes are immunoprecipitated.

(B) Plot depicting the average IP value and average input values (log2) of all genes analyzed. Enriched genes (>2-fold; red) and depleted genes (<2-fold; blue) are shown (q<0.05), and selected markers are shown: mc3r, Glp1r, Nts, Sst, Cartpt and Mbp (n = 3). All genes depicted have q-values <0.05 as calculated by Cufflinks.

(C) Average fold of change (IP/INP Log2) assessed by Taqman qPCR of enriched positive control markers Gfp and Nts, significantly enriched genes Mc3r, Glp1r, Sst and Cartpt (red) and depleted negative control markers Gfap and Mal (blue) (n = 3 biological replicates).

(D) Left panel, representative in situ hybridization image of Nts (green), vglut2 (red), vgat (white) and DAPI (blue). Right panel, quantification of Nts cells which express vgat (white) and vglut2 (red). $n=3$, Unpaired Students' t-test $p=0.009$; $t(4)=4.791$. Scale bars, 50 μm .

(E) Representative in situ hybridization image of Nts (green) and Glp1r (red). 2x digital zoom is presented in the inset. Scale bars for panels e-h, 50 μm .

(F) Representative in situ hybridization image of Nts (green) and Sst (red). Closer magnification is inset. 2x digital zoom is presented in the inset.

(G) Representative in situ hybridization image of Nts (green) and Cartpt (red). 2x digital zoom is presented in the inset.

(H) Representative in situ hybridization image of Nts (green) and Mc3r (red). 2x digital zoom is presented in the inset.

(I) Quantification of the percentage of Nts cells (black) expressing Nts/Glp1r (red), Sst (magenta), Cartpt (blue) and Mc3r (cyan). $n=3$ (Nts), $n=4$ (Nts/Glp1r, Nts/Mc3r), $n=6$ (Nts/Sst), $n=5$ (Nts/Cartpt)

(J) Left panel, experimental scheme. Wild-type mice are injected with exendin-4 (Ex4) directly into the LS or Glp1r-cre mice with viral expression of hM3Dq activatory DREADD receptor are tested for cumulative food intake.

(K) Cumulative food intake measured in grams (g) of mice following Vehicle (black) and Ex4 (red) injection in the LS. $n=11$ (Vehicle), $n=15$ (Ex4), Two-way ANOVA with post-hoc Bonferroni correction (Time: $F(1,24)=100.8$, $p<0.0001$; Treatment: $F(1,24)=11.54$, $p=0.0024$; Interaction: $F(1,24)=0.002$; $p=0.963$).

(L) Cumulative food intake measured in grams (g) of control mCherry (black) and hM3Dq (red) expressing Glp1r-cre mice following CNO injection. Two-way ANOVA with post-hoc Bonferroni correction. $n=10$ (mCherry), $n=11$ (hM3Dq). (Time: $F(1,19)=40.67$, $p<0.0001$; Subject: $F(1,19)=20.68$, $p=0.0002$; Interaction: $F(1,19)=9.407$, $p=0.0063$).

(M) Representative images of Nts (green), fos (red) and Glp1r (white) in control and

restraint stress (RS) mice. Right, quantification of co-localization between Nts, fos and Glp1r+ cells in control (black) and RS (red) mice. n= 4, Two-way ANOVA with post-hoc Bonferroni correction, *p<0.05, Row factor: $F(1,12)=1.460$, p=0.2502, Column Factor: $F(1,12)=5.056$, p=0.0441, Interaction: $F(1,12)=3.494$, p=0.0862. Scale bars, 50 μ m.

(N) Left, representative in situ hybridization image of Nts (green), Glp1r (blue) and Fos (red) of mice injected with vehicle or the Glp1r agonist Exendin-4 (Ex4). White arrows point to cells co-expressing Nts, Glp1r and Fos. Right, quantification of Fos+ cells expressing Nts and/or Glp1r in saline (black) or Ex4 (red) injected mice. n=3(Control), n=4 (Ex4); Two-way ANOVA with Bonferroni correction (Subject: $F(1,10)=71.7$, p<0.0001; Treatment: $F(1,10)=45.88$, p=0.0001; Interaction: $F(1,10)=37.58$, p<0.0001). Data are represented as mean $\pm$ SEM.

Fig 6

Figure 6. LS^{NTS} → LH projection regulates food intake in mice.

(A) Representative images of the LS injection site (left) and terminals in the lateral hypothalamus (left) of NTS-cre mice injected with a control mCherry reporter virus. Atlas images above the representative images depict the area of magnification.

(B) Representative image of cells labeled in the lateral hypothalamus (right) of NTS-cre mice injected with the anterograde tracer HSV-1sl-tdTomato in the LS. Middle, atlas image depicting the area of magnification to the right.

(C) Left, experimental scheme depicting mice injected with AAV expressing Chr2 or control mCherry into the LS and an optic fiber positioned over the lateral hypothalamus. Middle, food intake was measured in an Off-On-Off paradigm. Right, food intake measured

1 in grams (g) of control mCherry (black) or Chr2 (red) expressing mice before (off), during
 2 (on) and after (off) laser stimulation. n=11, Two-way ANOVA with post-hoc Bonferroni
 3 correction, Epoch: $F(2,40)=38.47$, $p<0.0001$; Subject: $F(1,20)=0.069$, $p=0.796$;
 4 Interaction: $F(2,40)=10.94$, $p=0.0002$. Data are represented as mean $\pm$ SEM.

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

1 SUPPLEMENTARY FIGURES

Supplementary Figure S1

2

3 Supplementary Figure S1. LS^{NTS} neurons regulate feeding behavior.

4 (A) Representative images of c-fos (green) in the lateral septum of control and restraint

5 stress (RS) mice. Right, quantification of c-fos+ cells in control (black) and RS (red) mice.

6 n = 4, Unpaired Student's t-test, *p < 0.05, t = 3.964, df = 4. Scale bars, 50 µm.

1 (B) Representative images of in situ hybridization experiments measuring the levels of fos
2 (red) and Nts (green) in the lateral septum of control and restraint stress (RS) male mice.
3 Left, quantification of fos+ cells in control (black) and RS (red) male mice, n= 4, Nested t-
4 test, ****p<0.0001, t=5.175, df=34. Right, quantification of nts+ cells in control (black) and
5 RS (red) male mice, n= 4, Nested t-test, **p<0.01, t=3.476, df=34. Scale bars, 50 um.

6 (C) Representative images of in situ hybridization experiments measuring the levels of fos
7 (red) and Nts (green) in the lateral septum of control and restraint stress (RS) female mice
8 in estrus or diestrus phase. Left, quantification of fos+ cells in control (black) and RS (red)
9 female mice in estrus or diestrus phase. n= 4, ****p<0.0001, F(3,34)=90.19. Right,
10 quantification of nts+ cells in control (black) and RS (red) female mice in estrus or diestrus
11 phase. n= 4, ****p<0.0001, F(3,38)=28.18. Scale bars, 50 um.

12 (D) Representative images of viral injection strategy and viral expression (mCherry, red)
13 and c-fos (green) in control virus (left) and hM3Dq (right) injected mice following CNO
14 injection. Right, quantification of c-fos+ cells in mCherry control (black) and hM3Dq (red)
15 expressing mice. n=3, Unpaired Students' t-test, p=0.003, t(4)=6.418.

16 (E) Food intake, measured in grams (g), in control mCherry (black) and hM3Dq (red)
17 expressing Nts-cre mice 1 hour (h), 2h, 4h, 8h and 24h following saline injection.
18 n=5(mCherry, n=5(hM3Dq). Mixed-model ANOVA with post-hoc Bonferroni correction
19 (Time: F(2.33,22.72)=298.8, p<0.0001; Subject: F(1,10)=6.203, p=0.032; Interaction:
20 F(4,39)=1.727, p=0.1637.

21 (F) Food intake, measured in grams (g), in control mCherry (black) and hM3Dq (red)
22 expressing Nts-cre mice 4 hour (h), 8h and 24h following CNO injection. n=7(mCherry),
23 n=10(hM3Dq). Two-way ANOVA with post-hoc Bonferroni correction (Time: F(2,18)=,
24 p<0.0001; Subject: F(1,9)=60.03, p<0.0001; Interaction: F(2,18)=12.38, p=0.0004.

25 (G) Food intake, measured in grams (g), in control mCherry (black) and hM4Di (red)

1 expressing Nts-cre mice 1 hour (h), 2h, 4h, 8h and 24h following saline injection.
2 n=7(mCherry), n=8(hM4Di). Two-way ANOVA with post-hoc Bonferroni correction (Time:
3 F(1,19)=415.8, p<0.0001; Subject: F(1,13)=1.042, p=0.326; Interaction: F(4,52)=2.839,
4 p=0.0333.
5 (H) Food intake, measured in grams (g), in control mCherry (black) and hM4Di (red)
6 expressing Nts-cre mice 4 hours (h), 8h and 24h following CNO injection. n=7(mCherry),
7 n=8(hM3Dq). Two-way ANOVA with post-hoc Bonferroni correction (Time: F(2,18)=184.5,
8 p<0.0001; Subject: F(9,18)=2.945, p=0.0245; Interaction: F(2,18)=12.38, p=0.004.

9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

Supplementary Figure S2

Supplementary Figure S2. LS^{NTS} neurons do not regulate anxiety-like behaviors.

(A) Time, measured in seconds (s), spent in the center and border of an open field in control mCherry (black) and hM3Dq (red) expressing Nts-cre mice. n=8(mCherry), n=9(hM3Dq). Two-way ANOVA with post-hoc Bonferroni correction, Subject: $F(1,15)=0.071$, $p=0.794$; Arena: $F(1,15)=135.8$, $p<0.0001$; Interaction: $F(1,15)=0.9093$, $p=0.355$.

(B) Distance, measured in centimeters (cm), traveled in the open field, in control mCherry (black) and hM3Dq (red) expressing Nts-cre mice. n=13(mCherry), n=14(hM3Dq), unpaired Students' t-test, $t(25)=0.403$, $p=0.69$.

(C) Velocity, measured as the cm per s (cm.s⁻¹), in the open field, in control mCherry (black) and hM3Dq (red) expressing Nts-cre mice. n=13(mCherry), n=14(hM3Dq), unpaired Students' t-test, $t(25)=0.915$, $p=0.369$.

(D) Time, measured in seconds (s), spent in the center and border of an open field with

1 novelty in control mCherry (black) and hM3Dq (red) expressing Nts-cre mice. n=5. Two-
2 way ANOVA with post-hoc Bonferroni correction, (Subject: $F(1,8)=0.358$, $p=0.566$; Arena:
3 $F(1,8)=21.54$, $p=0.0017$; Interaction: $F(1,8)=0.039$, $p=0.847$.

4 (E) Time, measured in seconds (s), spent in the open and closed arms of an elevated plus
5 maze in control mCherry (black) and hM3Dq (red) expressing Nts-cre mice. n=5. Two-way
6 ANOVA with post-hoc Bonferroni correction, Arm: $F(1,8)=111.2$, $p<0.0001$; Subject:
7 $F(1,8)=0.137$, $p=0.72$; Interaction: $F(1,8)=1.641$, $p=0.236$

8 (F) Distance, measured in centimeters (cm), traveled in the elevated plus maze, in control
9 mCherry (black) and hM3Dq (red) expressing Nts-cre mice. n=5, unpaired Students' t-test,
10 $t(8)=0.5347$, $p=0.6074$.

11 (G) Velocity, measured as the cm per s ($\text{cm}\cdot\text{s}^{-1}$), in the elevated plus maze, in control
12 mCherry (black) and hM3Dq (red) expressing Nts-cre mice. n=5, unpaired Students' t-test,
13 $t(8)=0.533$, $p=0.608$.

14 (H) Latency to feed, measured in seconds (s), in a novelty suppressed feeding task, in
15 control mCherry (black) and hM3Dq (red) expressing Nts-cre mice. n=5, unpaired
16 Students' t-test, $t(8)=0.027$, $p=0.979$. Data are represented as mean $\pm$ SEM.

17

18

19

20

21

22

23

24

25

Supplementary Figure S3

Supplementary Figure S3. LS^{NTS} neurons are not selectively tuned to feeding or movement.

(A) Experimental scheme. Mice were then attached to a fiber photometry setup with a 465nm LED (blue) and a 405 nm LED (purple) and a photodetector.

(B) Representative image of viral expression in the LS of Nts-cre mice injected with

1 gCaMP6s and implanted with an optic fiber in the LS.

2 (C) Average gCaMP6s z-score (black lines) and SEM (gray dotted lines) across all

3 recording sites time-locked to the start of the feeding bout. Right, Heat maps represent

4 the average % DF/F from each gCaMP6s recordings time-locked to feeding bout start.

5 (D) Area under the curve (a.u.) of fluorescent signal before (black) and after (red) the

6 feeding bout start. n=6, Paired Students' t-test, $t(5)=1.304$, $p=0.249$,

7 (E) Average gCaMP6s z-score (black lines) and SEM (gray dotted lines) across all

8 recording sites time-locked to the start of the start of locomotion.

9 (F) Area under the curve (a.u.) of fluorescent signal before (black) and after (red) the start

10 of locomotion. n=5, Paired Students' t-test, $t(4)=0.402$, $p=0.7083$.

11 (G) Experimental scheme of contextual fear conditioning.

12 (H) Percentage of freezing in Context A (CS+, red) and Context B (CS-, black). Unpaired

13 Student's t-test, $*p<0.05$, $t=3.233$, $df=4$, $n=3$. Data are represented as mean $\pm$ SEM.

Supplementary Figure S4

Supplementary Figure S4. Regulation of feeding and/or anxiety by Cartpt-, Glp1r- and Sst-expressing neurons in the lateral septum.

(A) Experimental scheme. Wild-type mice are injected with Ex4 or Mc3r agonist y-MSH directly into the LS or Glp1r-cre, Cartpt-Cre or Sst-cre mice with viral expression of hM3Dq activatory DREADD receptor are tested for cumulative food intake.

(B) Food intake, measured in grams (g) of mice 4 hours (h), 12h and 24h following Vehicle (black) and Ex4 (red) injection in the LS. $n=11$ (Vehicle), $n=15$ (Control), Two-way ANOVA with post-hoc Bonferroni correction (Time: $F(2,48)=223.6$, $p<0.0001$; Treatment: $F(1,24)=9.904$, $p=0.004$; Interaction: $F(2,48)=1.67$; $p=0.199$).

(C) Food intake, measured in grams (g) of control mCherry (black) and hM3Dq (red)

1 expressing Glp1r-cre mice 4 hours (h), 12h and 24h following CNO injection. n=xx, Two-
 2 way ANOVA with post-hoc Bonferroni correction. n=10(mCherry), n=11(hM3Dq). (Time:
 3 $F(2,38)=296.4$, $p<0.0001$; Subject: $F(1,19)=20.32$, $p=0.0002$; Interaction: $F(2,38)=6.077$,
 4 $p=0.0051$.

5 (D) Food intake, measured in grams (g) of mice 1-hour (h), 2h, 4h, 12h and 24h following
 6 Vehicle (black) and y-msh (red) injection in the LS. n=10(Vehicle), n=12(y-msh), Two-way
 7 ANOVA with post-hoc Bonferroni correction (Time: $F(4,80)=585.6$, $p<0.0001$; Treatment:
 8 $F(1,20)=0.66$, $p=0.426$; Interaction: $F(4,80)=2.67$; $p=0.038$.

9 (E) Food intake, measured in grams (g) of control mCherry (black) and hM3Dq (red)
 10 expressing Cartpt-cre mice 1 hour (h), 2h, 4h, 12h and 24h following CNO injection
 11 n=7(mCherry), n=5(hM3Dq), Two-way ANOVA with post-hoc Bonferroni correction. (Time:
 12 $F(2,17)=1,075$, $p<0.0001$; Subject: $F(1,10)=0.1767$, $p=0.6831$; Interaction: $F(4,40)=1.364$,
 13 $p=0.2637$.

14 (F) Food intake, measured in grams (g) of control mCherry (black) and hM3Dq (red)
 15 expressing Sst-cre mice 1 hour (h), 2h, 4h, 12h and 24h following CNO injection. n=xx,
 16 Two-way ANOVA with post-hoc Bonferroni correction. n=11(mCherry), n=12(hM3Dq).
 17 (Time: $F(4,68)=660.3$, $p<0.0001$; Subject: $F(1,17)=3,23$, $p=0.0903$; Interaction:
 18 $F(4,68)=1.006$, $p=0.4106$.

19 (G) Representative image of viral expression in the lateral septum of Cartpt-cre mice.

20 (H) Representative image of viral expression in the lateral septum of Glp1r-cre mice.

21 (I) Representative image of viral expression in the lateral septum of Sst-cre mice.

22 (J) Time, measured in seconds (s), spent in the open arms and closed arms of an elevated
 23 plus maze in control mCherry (black) and hM3Dq (red) expressing Cartpt-cre
 24 mice. n=7(mCherry), n=5(hM3Dq), Two-way ANOVA with post-hoc Bonferroni correction.
 25 (Arm: $F(1,10)=17.12$, $P<0.0001$; Subject: $F(1,10)=1.621$, $P=0.232$; Interaction:

1 F(1,10)=0.02, p=0.891.

2 (K) Time, measured in seconds (s), spent in the open arms and closed arms of an elevated
3 plus maze in control mCherry (black) and hM3Dq (red) expressing Glp1r-cre mice.

4 n=7(mCherry), n=8(hM3Dq), Two-way ANOVA with post-hoc Bonferroni correction. (Arm:

5 F(1,13)=79.20, P<0.0001; Subject: F(1,13)=0.3198, P=0.5814; Interaction:

6 F(1,13)=0.05114, p=0.825.

7 (L) Time, measured in seconds (s), spent in the open arms and closed arms of an elevated

8 plus maze in control mCherry (black) and hM3Dq (red) expressing Sst-cre mice. n=8, Two-

9 way ANOVA with post-hoc Bonferroni correction. (Arm: F(1,14)=39.48, p<0.0001; Subject:

10 F(1,14)=0.0007, p=0.979; Interaction: F(1,14)=0.1811, p=0.677. Data are represented as

11 mean +- SEM.

12

13

14

15

16

17

18

19

20

21

22

23

24

25

Supplementary Figure S5

Supplementary Figure S5. LS^{NTS} → LH does not regulate anxiety or locomotion.

(A) Left, experimental scheme of Glp1r-cre mice injected with AAV expressing cre-dependent mCherry in the lateral septum. Right, representative image of Glp1r fibers in the lateral hypothalamus. Scale bar are 100 μ m.

(B) Left, experimental scheme of Nts-cre mice injected with AAV expressing Chr2 in the lateral septum and optic fibers in the lateral hypothalamus. Right, Representative image of fiber optic path in the lateral hypothalamus. Scale bar are 100 μ m.

(C) Left, representative images of c-fos (green) and mCherry (red) in control mCherry (left) and Chr2 (right) expressing Nts-cre mice injected with CNO. Right, quantification of c-fos+ cells in control (black) and Chr2 (red) mice. n=3, unpaired Students' t-test, t(4)=4.999, p=0.008.

(D) Left, representative images of c-fos (red) and mCherry projections (green) in the lateral

1 hypothalamus of Nts-cre mice expressing control mCherry (left) or Chr2 (right) virus
2 injected with CNO. n=3, unpaired Students' t-test, $t(4)=5.089$, $p=0.007$.

3 (E) Distance, measured in centimeters (cm), traveled in the elevated plus maze, in control
4 mCherry (black) and Chr2 (red) expressing Nts-cre mice. n=5, unpaired Students' t-test,
5 $t(8)=0.9092$, $p=0.39$.

6 (F) Velocity, measured as the cm per s ($\text{cm}\cdot\text{s}^{-1}$), in the elevated plus maze, in control
7 mCherry (black) and Chr2 (red) expressing Nts-cre mice. n=5, unpaired Students' t-test,
8 $t(8)=1.291$, $p=0.233$.

9 (G) Time, measured in seconds (s), spent in the center and border of an open field in
10 control mCherry (black) and Chr2 (red) expressing Nts-cre mice. n=5. Two-way ANOVA
11 with post-hoc Bonferroni correction, Subject: $F(1,8)=0.051$, $p=0.827$; Border/Center:
12 $F(1,8)=18.48$, $p=0.0026$; Interaction: $F(1,8)=0.791$, $p=0.4$. Data are represented as mean
13 $\pm$ SEM.

14

15

16

17

18

Table 1. c-Fos expression in the mouse brain after restraint stress.

Brain Region	c-Fos expression
Central Amygdala (CeA)	+
Basolateral Amygdala (BLA)	++
Insular Cortex (IC)	++
Clastrum (CLA)	+++
Lateral Hypothalamic Area (LHA)	+++
Lateral Habenula (Lhb)	++
Paraventricular nucleus of the Thalamus (PVT)	++
Lateral Septum (LS)	+++
Medial Preoptic Nucleus (MPO)	+++
Periaqueductal Gray (PAG)	++
Bed Nucleus of the stria terminalis (BNST)	+++

Score: - (no expression), + (low expression), ++ (moderate expression), +++ (high expression)