1 Transcriptome analysis reveals infection strategies employed by *Fusarium* 2 *graminearum* as a root pathogen

3 Yi Ding^{a,c,*}, Donald M Gardiner^a, Kemal Kazan^{a,b,*}

4 ^aAgriculture and Food, Commonwealth Scientific and Industrial Research Organization, 306 Carmody
5 Road, St Lucia, 4067, Queensland, Australia;

6 ^bQueensland Alliance for Agriculture and Food Innovation (QAAFI), Alliance for Agriculture and Food
7 Innovation (QAAFI), The University of Queensland, Brisbane, St Lucia, 4067, Queensland, Australia;

8 ^cCurrent address: The Plant Breeding Institute, School of Life & Environmental Sciences, Faculty of
9 Science, The University of Sydney, Cobbitty, 2570, New South Wales, Australia;

10 *Corresponding authors.

11 yi.ding@sydney.edu.au (Yi Ding), Kemal.kazan@csiro.au (Kemal Kazan)

12

13 Abstract

14 The fungal pathogen *Fusarium graminearum* infect both heads and roots of cereal crops
15 causing several economically important diseases such as head blight, seedling blight, crown
16 rot and root rot. Trichothecene mycotoxins such as deoxynivalenol (DON), a well-known
17 virulence factor, produced by *F. graminearum* (*Fg*) during disease development is also an
18 important health concern. Although how *F. graminearum* infects above-ground tissues is
19 relatively well studied, very little is known about molecular processes employed by the
20 pathogen during below-ground infection. Also unknown is the role of DON during root
21 infection. In the present study, we analyzed the transcriptome of *F. graminearum* during root
22 infection of the model cereal *Brachypodium distachyon*. We also compared our *Fg*
23 transcriptome data during root infection with those reported during wheat head infection.
24 These analyses suggested that both shared and unique infection strategies employed by the
25 pathogen during colonization of different host tissues. Several metabolite biosynthesis genes

induced in *F. graminearum* during root infection could be linked to phytohormone production, implying that the pathogen likely interferes root specific defenses. In addition, to understand the role of DON in *Fg* root infection, we analyzed the transcriptome of the DON deficient *Tri5* mutant. These analyses showed that the absence of DON had a significant effect on fungal transcriptional responses. Although DON was produced in infected roots, this mycotoxin did not act as a virulence factor during root infection. Our results reveal new mechanistic insights into the below-ground strategies employed by *F. graminearum* that may benefit the development of new genetic tools to combat this important cereal pathogen.

Introduction

Fungal plant pathogens have adopted versatile strategies to colonize their hosts. While some fungal pathogens show strict host and tissue specificity, others can adjust their lifestyles to infect different hosts and tissues. Some fungal pathogens such as the rice blast *Magnaporthe oryzae* (Marcel et al., 2010; Sesma and Osbourn, 2004) and the corn smut *Ustilago maydis* (Mazaheri-Naeini et al., 2015), which commonly invade above-ground plant parts, can also undergo developmental processes resembling to root infecting fungi. Such changes in the pathogen may require sensing of host signals and we previously showed that sensing of root signals prior to root infection by *Fg* can indeed lead to developmental changes in the pathogen (Ding et al., 2020). As a member of the *Fusarium* species complex, *Fusarium graminearum* (*Fg*) causes Fusarium Head Blight (FHB) or scab, one of the most economically important diseases of cereal crops. FHB causes substantial yield losses and mycotoxin contaminations of grains, resulting in billions of dollars of economic losses worldwide and threatening our food supply and safety (Chen et al., 2019; Trail, 2009). Most studies on FHB have so far focused on wheat heads as the pathogen initially infects individual wheat florets from which it can spread to other florets through the rachis and can eventually colonize the whole spike. However, recent research has shown that *Fg* is also capable of infecting roots and young seedlings of wheat, barley and maize, causing crown rot, root rot and seedling blight (Henkes et al., 2011; Lanoue et al., 2010; Stephens et al.,

2008; Wang et al., 2015; Zhou et al., 2019). During the initiation of root infection, *Fg* forms a peg structure outside the root surface and move inter- and intra-cellularly without causing root necrosis. This early colonization stage is followed by a transition of the fungus to a necrotrophic life style where lesions develop and spread to stems and aboveground tissues (Wang et al., 2015).

The availability of the complete genome sequence of *Fg* makes investigations of global regulation of gene expression in this fungus feasible (Kazan and Gardiner, 2018a; Ma et al., 2013). Infection strategies of *Fg* evaluated by transcriptome analyses in different hosts and tissues exclusively during infection of their above-ground tissues such as heads, stems and coleoptiles revealed mostly distinct, but also common gene expression patterns (Boedi et al., 2016; Harris et al., 2016; Lysøe et al., 2011; Zhang et al., 2012, 2016). Interestingly, different cereal species can produce different defense-related metabolites (Dutartre et al., 2012) and such differences may explain why *Fg* might need to tailor its arsenal during colonization of different hosts (Harris et al., 2016). A recent comparative transcriptomic study of *Fg* also showed differential expression of fungal genes during infection of FHB resistant and susceptible wheat genotypes (Pan et al., 2018).

Although transcriptome studies have provided clues associated with host specificity of the *Fg* infection process, fungal transcriptomes of *Fg* mutants with altered virulence have rarely been tested on the same hosts. The mycotoxin deoxynivalenol (DON) is a *Fg* virulence factor during infection of wheat heads (Proctor et al., 1995). DON may also be needed during the interaction of *Fg* with its broader environment (Audenaert et al., 2013). The first step in DON biosynthesis is catalyzed by the trichodiene synthase *Tri5*, which cyclizes farnesyl pyrophosphate to trichodiene (Hohn and Beremand, 1989). In contrast to its strong expression pattern during wheat head infection, *Tri5* did not show increased *in planta* expression during wheat coleoptile infection, suggesting that DON's effect on pathogen virulence is tissue specific (Zhang et al., 2012). Other studies have suggested a crucial role for DON during the colonization of wheat stems by *Fg* and the related pathogens *F.*

80 *pseudograminearum* and *F. culmorum* (Desmond et al., 2008; Mudge et al., 2006; Powell et
81 al., 2017; Scherm et al., 2013a). The recent finding where Fhb7-mediated FHB and crown
82 rot disease resistance relies on DON detoxification also highlighted virulence function of this
83 mycotoxin in wheat (Wang et al., 2020). Furthermore, DON is known to activate defense
84 gene expression in wheat (Desmond et al., 2008). Indeed, host transcriptional changes
85 observed in *Brachypodium distachyon* (*Bd*) and wheat spikelets infected by the *Tri5* deletion
86 mutants (Δ Tri5) differed from those by wildtype *Fg* (Brauer et al., 2020; Pasquet et al., 2014).
87 DON biosynthesis in *Fg* is regulated by *Tri6* and *Tri10* transcription factors. Analyses of
88 deletion mutants for these genes by transcriptome profiling during plant infection revealed
89 significant transcriptional alterations for a large number of genes, many of which have not
90 been implicated previously in toxin production (Seong et al., 2009). Genetic analyses
91 undertaken in *Fg* have identified many genes influencing DON biosynthesis (Chen et al.,
92 2019). However, DON-non-producing mutants have not been employed for evaluating the
93 effect of this toxin on global transcriptional responses in *Fg*.

94 Phytohormones mediate immune responses in plants after pest or pathogen attack (Pieterse
95 et al., 2009). In turn, plant pathogenic fungi have evolved ways to compromise host hormone
96 pathways. This is achieved by degrading or producing phytohormones or interfering with
97 their signaling pathways (Kazan and Lyons, 2014; Patkar and Naqvi, 2017). For instance,
98 emerging evidence suggests that phytohormones such as abscisic acid (ABA), gibberellic
99 acid (GA) and ethylene (ET) produced by fungi participate in pathogenicity (Chanclud and
100 Morel, 2016). Previous studies indicated that *Fg* can likely produce auxin (IAA) and ET that
101 may be utilized for attenuating host defenses during FHB (Foroud et al., 2019; Luo et al.,
102 2016; Svoboda et al., 2019). In addition, SA hydroxylases were proposed to be involved in
103 the degradation of host SA by *Fg* (Hao et al., 2019; Qi et al., 2019; Rocheleau et al., 2019).
104 However, how hormonal compounds produced by *Fg* or the host plant are metabolized or
105 involved in host infection is poorly studied. This is at least in part due to potential co-

existence of phytohormones derived from both host and the pathogen in the infected tissue and the lack of knowledge on fungal genes involved in phytohormone biosynthesis.

Currently, potential molecular mechanisms employed by *Fg* during root infection are unknown (Kazan and Gardiner, 2018a). A global transcriptome analysis would provide a powerful way to broadly reveal previously unknown features during root infection, thus promoting development of new strategies for combating this pathogen. In addition, to what extent DON may affect global transcription in the pathogen has not been investigated. We previously reported an RNA-seq based transcriptome profiling of *Fg* prior to its physical contact with *Bd* roots. This analysis enabled us to discover novel genes that are involved in nitric oxide (NO) production in *Fg* upon sensing of root signals (e.g. metabolites found in the root exudates) and pathogen virulence (Ding et al., 2020). In this study, using *Bd* as a cereal model, we asked how *Fg* behaves as a root pathogen. To answer this question, we analyzed the transcriptome of *Fg* during infection of *Bd* roots. We analyzed phytohormone levels of infected *Bd* roots to understand potential roles played by phytohormones derived from the fungus. In addition, by comparing the transcriptomes of WT *Fg* and the DON deficient *Tri5* mutant, we uncovered novel insights into global effects of DON on fungal gene expression and metabolism during the infection of host roots.

Materials and methods

Plant and fungal materials and root infection assay

The *Fg* CS3005 WT, *Tri5* mutant (Desmond et al., 2008) and *Tri5-GFP* expressing (Gardiner et al., 2009) strains were routinely maintained on Potato Dextrose Agar (PDA, BD Difco). *Bd* (Bd21-3) seeds were surface sterilized and pre-germinated on filter paper (Whatman) placed in 150mm x 25 mm petri dishes (Corning) for 5 days. *Bd* roots inoculation with the *Fg* strains was carried out as described previously (Ding et al., 2020). Briefly, agar plugs (0.25 cm diameter) taken from fungal culture plates (Carboxymethylcellulose agar) were transferred to center of minimum media (pH 7) plates and pre-grown for 3 days. Five-day-old *Bd* seedlings

were placed above the fungal colonies and inoculated for additional 5 days. Three biological replicates for *Fg* WT and Δ Tri5 inoculated seedlings, and the *Fg* WT alone mycelia were produced by pooling materials from 10-12 plants or fungal mycelia.

RNAseq and transcriptomic analyses

Fungal and root materials were frozen in liquid nitrogen immediately after harvest. Total RNA was extracted from homogenized samples using a Qiagen RNeasy plant RNA extraction kit with on-column DNase I (Qiagen) digestion following manufacturer's instructions. RNA was quantified and quality-checked prior to sequencing. An Illumina HiSeq2500 High Output platform was used to generate 50-base pair single-end reads (Australian Genome Research Facility). Reads quality control, alignment, transcript abundance and differential expression (DE) analyses were performed according to the method described previously (Ding et al., 2020). For pairwise comparison (*Fg*WT-only vs. *Fg*WT-*Bd*, or *Fg*WT-*Bd* vs. Δ Tri5-*Bd*), different sample files were normalized and merged. Reads were measured as FPKM (Fragments Per Kilobase of gene model per Million reads mapped), and a normalization method developed by Hart et al. (2013) was used and to eliminate background noise of FPKM values where genes with a Gaussian-fit derived $\log_2$ (FPKM) value higher than -3 were considered as expressed. $|\log_2$ fold change| ≥ 1 and Benjamini and Hochberg-adjusted P value < 0.05 were applied to DE genes. RNAseq data are available at NCBI under the accession no. PRJNA631873.

Annotation and functional categorization of differentially expressed genes (DEGs)

BLAST2GO (Götz et al., 2008) was used to assign annotations for fungal DEGs. BLASTP reciprocal best hit analyses were performed in order to identify putative orthologous genes and match unique gene identifiers of the *Fg* CS3005 and PH-1 strains (Gardiner et al., 2014). Based on the PH-1 identifiers, classification ontology of DEGs from pairwise comparisons was annotated with FungiFun2 and subsequently subjected to enrichment analyses (<https://elbe.hki-jena.de/fungifun>). Functional categories were considered as enriched in the

genome if an enrichment Benjamini-Hochberg adjusted P value is smaller than 0.05. Prediction of protein cellular localizations, secretome and putative effectors, transporters, carbohydrate-active enzymes, lipases, secondary metabolism enzymes and transcription factors was according to previously described methods (Ding et al., 2020).

Identification of homologous genes between *Fg* and *Fp*

All RNAseq reads were mapped to the *Bd* reference genome first. Unmapped reads were extracted and aligned to the *Fg* CS3005 and *Fp* CS3096 reference genomes, respectively, as per previously described (Ding et al., 2020). Only reads that could be mapped to both genomes were retained. Next, read counts measured as FPKM values were log transformed, normalized and subjected to PCA analysis (Fig. S4). Homologous genes within *Fg* and *Fp* were identified using a reciprocal best BLAST hit (RBBH) approach. Gene orthologs with identity of equal or higher than 99% were kept and used for syntenic analysis using a R package shinyCircos (Yu et al., 2018).

cDNA synthesis and quantitative real-time PCR analysis

0.5-1 µg total RNA was prepared for first-strand cDNA synthesis using the superscript IV synthesis kit (Invitrogen, USA) and quantitative real-time RT-PCR (qRT-PCR) was performed using the ViiA 7 real-time PCR detection platform (Applied BioSystems). Primers were based on previous studies (Ding et al., 2020; Voigt et al., 2005). Expression levels were normalized to the fungal house-keeping gene *α-Tubulin* and were averaged over three biological replicates.

Root sectioning and fluorescence microscopy

Root dissection was carried out following a previous described method (Ursache et al., 2018). Briefly, roots were harvested at 5dpi and fixed with 4% paraformaldehyde (Sigma) in PBS buffer (pH=6.9) overnight, washed twice with PBS buffer and cleared with ClearSee solution (Ursache et al., 2018) for 3 days. After clearing, roots were hand sectioned from 3

cm above the tips and stained with 0.2% Basic Fuchsin (Sigma). To image GFP and Basic Fuchsin fluorescence, root samples were observed using 488-nm excitation / 519-nm emission, and 561-nm excitation / 625-nm emission, respectively, on a Zeiss Axio Imager M2 microscopy.

UHPLC quantification of metabolites

For metabolite extractions, mock and *Fg*-infected roots, fungal mycelia and media samples were collected at 5 dpi, immediately frozen and ground in liquid nitrogen. 100 mg of fine-ground materials were resuspended in 2 mL extraction buffer (Ethyl acetate: methanol: dichloromethane, 3:2:1 v/v). 5 µL extracts from 6 replicates of each conditioned sample were injected into a Waters ACQUITY ARC UPLC system with Photodiode Array and passed through a Phenomenex Kinetex column (C18, 1.7 µm, 100 × 2.1 mm). The mobile phases consisted of solvent A (10 mM ammonium formate in water) and solvent B (10 mM ammonium formate in acetonitrile). The gradient program was a linear gradient from 2-40% solvent B delivered over 22 min followed by 40-80% B over 1.5 min at a constant flow rate of 0.4 mL per minute. As external standards, jasmonic acid (Sigma, 10 mM), methyl-jasmonate (Sigma, 40 mM), salicylic acid (Sigma, 10 mM), deoxynivalenol (Sigma, 100 ng/µL), gibberellic acid (Sigma, 10 mM), indole-3-acetic acid (Sigma, 2 mM) were freshly prepared for serial dilutions (10x, 100x, 1000x, 10000x). Metabolites were quantified according to the concentration-gradient derived standard curves of external standards. Chromatography absorbance data were aligned using the following wavelengths: 204nm (for SA, JA and MeJA), 254nm (for IAA and DON) and 214nm (for GA), and extracted using the Empower 3 software (Waters).

Results and discussion

The *Fg* transcriptome during *Bd* root infection

Despite various studies investigating the transcriptome of *Fg* during colonization of above-ground tissues, *Fg* transcriptome during root infection has not been studied before (Kazan

and Gardiner, 2018a). Therefore, here, we first investigated the *Fg* transcriptome during root infection of the model host *Bd*. This transcriptome experiment, which was initially designed for the discovery of a novel host-sensing mechanism in *Fg* prior its physical contact with roots, included three replicates of each of 1) *Fg* grown on minimal media (MM) (*Fg*-only) 2) *Fg*WT and *Bd* roots grown together on MM without physical contact, 3) WT *Fg* colonizing *Bd* roots on MM (*Fg*WT colonization) and 4) a *Tri5* mutant colonizing *Bd* roots (Δ *Tri5* colonization). Previously, by comparing 1 with 2, we discovered new regulators involved in host sensing-mediated NO production (Ding et al., 2020). Here, we report on the comparisons between 1 and 3, 1 and 4 and 3 and 4, as detailed analyses of these have not been reported previously.

A total of 174,399,800 single-end reads, including 114,577,185 reads from *Fg* alone and *Fg*-infected *Bd* root samples (Ding et al., 2020) and 59,822,615 reads from the Δ *Tri5* root samples were generated by Illumina sequencing of mRNA libraries (Suppl. data 1). Prior to read mapping to the *Fg* CS3005 genome (Gardiner et al., 2014), reads aligned to the *Bd* genome were filtered. Of these filtered reads across all the samples, at least 91%, ranging from 5 to 19.9 million, were mapped to the *Fg* reference genome. Among these mapped reads, less than 0.3% matched to multiple genomic locations (Suppl. data 1). Out of the 12590 transcripts detected, 11942, 11291 and 11327 were found actively expressed in *Fg*-only, in WT *Fg* colonization and in *Tri5* mutant colonization conditions, respectively.

Our analysis revealed a total of 2049 genes that were differentially expressed (DE) ($\log_2$ FC ≥ 1 and adjusted p value < 0.05) (Fig. 1A, Suppl. data 2) during colonization of *Bd* roots relative to *Fg*-only. Of these, 1281 and 768 were up- and down-regulated, respectively, during root colonization. The proportion of DE genes (DEGs) (around 20% of total number of genes found in the *Fg* genome) was similar to those observed during above-ground infection by *Fg* in other studies (Brown et al., 2017; Puri et al., 2016). A previous study indicated that some defense genes are similarly regulated in wheat roots and spikes in response to *Fg* (Q. Wang et al., 2018), suggesting that common strategies might be employed by *Fg* to infect

different tissue types. Indeed, we previously reported that fungal knockouts of several DEGs identified here showed defects in both root and head infection (Ding et al., 2020).

Fungal processes employed by *Fg* during root infection

Detailed analysis of DEGs (*Fg*-only vs WT *Fg* colonization) revealed a number of enriched functional categories likely to be used by the pathogen during root infection. Below, some of these functional categories were discussed in more detail.

Genes encoding plant cell wall degrading enzymes (CWDE)

The *Fg* genome comprises a large number of genes encoding hydrolytic enzymes, transporters, secreted proteins and multiple gene clusters associated with secondary metabolite biosynthesis (Scherm et al., 2013; Sieber et al., 2014). These enzymes, non-enzymatic proteins and secondary metabolites together are generally considered fungal pathogenicity factors and their deployment during *Bd* root colonization indicates the importance of these pathway for sustaining the infection process. Indeed, over 30% of DEGs encode secondary metabolism enzymes (SMEs), secreted proteins, carbohydrate-active enzymes (CAZymes) and transporters (Fig. 1A). The most enriched categories for significantly up-regulated genes were associated with carbohydrate hydrolytic pathways (Suppl. data 3). Functional annotations showed that over half of the *in planta* activated CAZymes displayed modular structures related to CWDE acting on wall polymers such as cellulose, hemicellulose, lignin and pectin (Fig. 1B, Suppl. data 4), indicating the utilization of carbon from plant cell walls is a prominent capability of *Fg* during the colonization of *Bd* roots.

Genes encoding secreted proteins and putative effectors

Fungal pathogens produce many small secreted proteins or effectors to help facilitate host colonization. However, relatively little is known about potential *Fg* effectors. By comparing our DEGs with the previously defined *Fg* secretome (Brown et al., 2012) and using the EffectorP 2.0 software (Sperschneider et al., 2018), we identified 250 putative secreted proteins along with 65 predicted fungal effector encoding genes (Fig S1C, Suppl. data 4 and

5). Many of these secreted protein genes, including 189 induced ones (Suppl. data 5), encode putative lipases and peptidases predicted to perform hydrolytic functions (Fig. 1C). Among the 65 predicted effector-encoding genes, 52 were significantly up-regulated during *Bd* root infection (Suppl. data 5). Some of these DE genes were reported to encode effectors actively secreted by *Fg* during *in vitro* growth. For example, *FG05_04074* encodes a protein of unknown function detected in two different secretomes (Lu and Edwards, 2016; Yang et al., 2012). Other putative effectors with annotated functional domains also found in previous studies included two glycoside hydrolases (*FG05_11037* and *FG05_06466*), a putative acetylase (FG05_11280), a putative endonuclease (*FG05_03365*) and a ceratoplatanin family protein (*FG05_10212*) for which roles of protection against host defense have been proposed (Lu and Edwards, 2016; Quarantin et al., 2016; Yang et al., 2012). Interestingly, the top induced effector candidates were mostly with predicted enzymatic functions (Suppl. data 5). Function of these differentially regulated putative effectors, such as *FG05_04735* encoding a putative hypersensitive response-inducing elicitor and *FG05_02255* a LysM domain containing protein (Suppl. data 5), could be predicted in comparison with those containing similar structural domains from different fungal pathogens and generally associated with host penetration, spore dispersal, triggering plant defense responses, inhibiting chitin-induced immunity or protecting against plant lysis (De Jonge et al., 2010; Khan et al., 2016; Lo Presti et al., 2015; Marshall et al., 2011; Mentlak et al., 2012). Although exact functions of these genes up-regulated during infection are largely unknown, it can be speculated that these secreted proteins and putative effectors could benefit the fungus during the colonization of host roots.

Genes encoding secondary metabolism enzymes (SME)

Fg is known to produce many secondary metabolites (SMs) during infection (Ma et al., 2013). In line with this, a strong induction of expression could be observed for genes encoding key signature enzymes (Fig. S1A), including the longiborneol synthase *CLM1* (*FG05_10397*), butenolide synthase (*FG05_08079*), *TRI5* and the terpenoid synthase *DTC1* (*FG05_03066*)

during infection (Suppl. data 6), suggesting that the corresponding products culmorin, butenolide, trichothecene and carotenoid may be the major mycotoxins delivered by the fungus to facilitate root infection. Among them, butenolide and trichothecene pathways are known to be co-regulated *in vitro* and *in planta* (Sieber et al., 2014). In contrast, down-regulation or very low *in planta* expression of other key SME genes, such as *PKS12*, *NPS2*, *NPS1*, *PKS4* and *PKS10* (Suppl. data 6), indicates that certain types of mycotoxins such as aurofusarin, ferricrocin, malonichrome, zearalenone and fusarin C might not be highly produced by *Fg* during root colonization. Aurofusarin does not affect wheat head infection by *Fg* (Malz et al., 2005), whereas ferricrocin and malonichrome have been shown to be important for pathogenesis-related development of *Fg* (Oide et al., 2014). The tailoring enzyme genes are usually clustered and co-regulated with the corresponding signature enzyme genes in *Fg*. These genes encode cytochrome P450s, oxidoreductases, acyltransferases and methyltransferases mainly involved in the SM pathway responsible for biosynthesis and modification of SM products (Sieber et al., 2014). Therefore, up-regulation of a large portion of tailoring enzyme genes found in this study is consistent with the regulation pattern of the signature enzymes (Fig. S1B).

Genes encoding fungal transporters

Transporter encoding genes mostly induced during *Bd* root colonization comprised a large group within the DE gene list (Fig. S1C, Suppl. data 7). Indeed, the regulation of transporter genes, particularly those associated with carbohydrate and nitrogen uptake as well as the ATP-binding cassette transporters (ABC transporters), is often linked to fungal nutrient assimilation, sensing, defense and pathogenicity status in pathogenic fungi (Abou Ammar et al., 2013; Coleman and Mylonakis, 2009; Divon and Fluhr, 2007; Gardiner et al., 2013; Schuler et al., 2015; Struck, 2015; Yin et al., 2018). Interestingly, the major facilitator superfamily (MFS) transporters associated with phosphate (Pi) transport and multidrug resistance (MDR) were mostly down-regulated during root infection compared to media alone controls (Fig. S1C, Suppl. data 7). During colonization of maize stalk, *Fg* overcomes Pi

limitation by up-regulating high-affinity Pi transporter genes *FGSG_03172* and *FGSG_02426* (Zhang et al., 2016). The observation here that expressions of these genes during *Bd* root infection significantly reduced indicates a relatively rich root Pi environment under the experimental conditions and the time point examined. The down-regulation of MDR transporter genes suggested that they are possibly not essential for *Fg* to resist against root-derived anti-fungal compounds or self-derived toxins. The elevated expression of a large number of MFS-type carbohydrate transport genes together with the induction of PCWDE genes indicate that *Fg* preferentially utilizes carbon to accomplish the infection cycle and a state of glucose depletion may exist at the examined stage. The top induced ABC transporters (Suppl. data 7) exclusively belonging to the ABC-G type transporters are known to be associated with self-protection, possibly by effluxing of antifungal compounds in many pathogenic fungi (Coleman and Mylonakis, 2009). Thus, it is likely that these ABC transporters together with the MFS family multidrug resistance transporters could contribute to the virulence and fitness of *Fg* by detoxifying plant defense compounds.

We also observed that genes encoding amino acid-related transporters such as the amino acid/polyamine/organocation (APC) family and the amino acid/auxin permease (AAP) family transporters, which are the major nitrogen transporters, were differentially expressed during root infection. In contrast, no inorganic nitrogen transporter gene showed altered expression (Suppl. data 7). This suggests that *Fg* root infection requires plant-derived organic nitrogen sources, and is consistent with the finding that polyamines as well as their amino acid precursors are potent DON inducers in *Fg* and play important roles during head infection (Gardiner et al., 2010, 2009). Interestingly, the highest induced transporter (*FG05_02278*, over 10-fold logFC) gene encodes a putative APC family protein transporter involved in choline uptake. Choline was identified as one of the major fungal growth stimulators in wheat anthers and implicated in promoting *Fg* virulence (Strange et al., 1972). Thus, it is possible that choline, in addition to amino acids and their derivatives, is another major factor contributing to *Fg* root colonization.

A small set of ‘core’ genes is activated during infection of different hosts and tissues by *Fg*

To obtain additional insights into *Fg* pathogenicity, we compared the *Fg* genes found to be induced during root colonization in this study with those previously reported to be induced during the colonization of other hosts or tissues (Brown et al., 2017; Harris et al., 2016; Lysøe et al., 2011; Zhang et al., 2012, 2016). These previous studies have reported several subsets of *in-planta* expressed *Fg* genes at multiple infection time-points and different disease development stages. To make a broader comparison, *Fg* genes that were commonly induced during infection at any of the studied time-points were selected. These included 3591 *Fg* genes induced during the infection of wheat heads (Brown et al 2017), 5061 genes expressed during the infection of wheat and barley heads and maize ears (Harris et al 2016) as well as 344 and 3066 genes induced during the infection of wheat juvenile coleoptiles (Zhang et al., 2012) and maize stalks (Zhang et al., 2016), respectively (Suppl. data 8). Through these comparisons, a total of 38 *Fg* genes commonly induced across all gene lists were identified (Fig. 2A, Suppl. data 9). Some of these genes were also differentially expressed between a FHB resistant and a susceptible wheat genotype (Pan et al., 2018). Unexpectedly, no mycotoxin- or pathogenicity-related SME genes were present among these 38 genes (Fig. 2B). An ABC transporter gene, *FG05_04580* (*FgABC1*), and its flanking neighbor *FG05_04581*, which encodes a transcription factor highly inducible by the mycotoxin zearalenone (Lee et al., 2010), were found among these common genes. Deletion of *FgABC1* causes reduced virulence of *Fg* on tested wheat tissues (Abou Ammar et al., 2013; Gardiner et al., 2013). Interestingly, *FgABC1* and *FG05_04581* homologs in the closely related pathogen *F. culmorum* were both highly induced by the antifungal compound tebuconazole (Hellin et al., 2018). This indicates that *FgABC1* and *FG05_04581* could be involved in self-protection against various defensive chemicals consistent with the observation that *FgABC1* contributes to protection against the fungicide benalaxyl (Gardiner et al., 2013). Furthermore, most of these common genes encode non-SM enzymes such as

CAZymes, peptidases and putative effectors. Among them, *FG05_03624*, a gene encoding a secreted xylanase, was previously shown to promote necrosis during *Fg* head infection (Moscetti et al., 2015). Protein homologs of several of these genes were also shown to be virulence factors in other fungal pathogens. For instance, *FG05_00028* is homologous to metallopeptidases (MEP1), which were shown to be apoplastic effectors in *F. oxysporum* and *M. oryzae* (Jashni et al., 2015; Yan and Talbot, 2016). In *F. oxysporum*, the metallopeptidase FoMEP1 and the serine protease FoSEP1 act synergistically to cleave host chitinases and prevent their degradation of fungal cell walls (Jashni et al., 2015). Indeed, a serine-type proteinase inhibitor encoded by *FG05_08012* found in our gene list, shows high similarity to FoSEP1 (Suppl. data 9). Another putative hypersensitive inducing elicitor *FG05_04741* shows significant homology to the *Verticillium dahlia* effector PevD1, which was shown to be a secreted elicitor triggering host defense and cell death (Liang et al., 2018). We hypothesize that these putative *Fg* effectors may perform roles that are similar to those found in other fungal pathogens. Taken together, it can be hypothesized that a common set of *Fg* genes seems to play essential roles in *Fg* for successful colonization of different tissue types.

***Fg* genes specifically upregulated in *Bd* roots**

We have identified 257 *Fg* genes that were exclusively upregulated during *Bd* root infections (Fig. 2A). Functional category analysis showed a significant enrichment for genes involved in transmembrane transport and cellular import (FDR = 0.00334). Of 34 transporter encoding genes induced, 12 were predicted to be associated with carbon transport (Suppl. data 10). This supports the finding discussed above that carbon utilization by *Fg* plays a role during *Bd* root colonization. Three ABC-G and two MDR transporters (Suppl. data 10) found among the enriched transporters might be specifically associated with detoxification of *Bd* root defense compounds. Among the four predicted effectors induced in roots (Suppl. data 10), the putative host-necrosis inducer protein *FG05_10212* was shown to be constitutively expressed during infections of wheat heads and *in vitro* and confirmed as an extracellular

protein (Lu and Edwards, 2016). The induction of this effector might contribute to necrosis observed in the infected *Bd* roots. Notably, of the seven putative *Fg* PCWDEs whose transcripts were only induced in *Bd* roots, five use lignin as substrate (Suppl. data 10), suggesting that lignin-degradation by *Fg* in *Bd* roots. Overall, while some common infection strategies may be employed by *Fg* during infection of different hosts and tissue types, there appears to be also unique processes used based on the activation of specific *Fg* genes during root colonization.

Partially shared infection strategies may be used by *Fg* and its sister species *Fp* during above- and below-ground infection of *Bd*

Previously, above-ground responses to the infection of *Bd* seedlings by *F. pseudograminearum* (*Fp*), another fungal species that is highly similar to *Fg* at the whole genome level (Gardiner et al., 2018) and was previously considered to be the same species as *Fg* (Kazan and Gardiner, 2018b), have been investigated (Powell et al., 2017). Both *Fg* and *Fp* show highly similar infection patterns on *Bd* (Fitzgerald et al., 2015). In addition, most genes are located in similar genomic regions in both fungi, whereas only a few species-specific genes, which could not be revealed by syntenic analysis, were found in genomic locations displaying high SNP densities (Gardiner et al., 2018). To further explore organ specificity of *Fg* infection on the same host, we compared the transcriptome of *Fg* with that of *Fp* (NCBI accession no. SRR3695327), during above ground infection of *Bd* at the same time point. Only *Fg* and *Fp* orthologous genes, which could be mapped to both of the genomes and share an identity of $\geq 99\%$ were retained in this comparison. This stringent cut-off allows comparisons of only highly conserved genes that might be predicted to show similar pattern in expression and function in these two closely related fungal species.

In total, 1835 of the *Fg* DEGs were matched to *Fp* and formed a syntenic map (Fig. 3A and Suppl. data 11) consistent with the previously revealed genome structures (Gardiner et al., 2018). These genes were, in general, similarly expressed in *Fg* and *Fp* as indicated by the logarithmic transformed FPKM values (Fig. 3A). However, some variability in gene

expression between the two species could be observed, particularly for genes found on chromosome 2 where the *Fg* orthologs tend to be preferentially expressed, as reflected by the expression heat map (Fig. 3A). Parts of chromosome 2 were previously identified as regions of the *Fg* genome that are rapidly evolving (Sperschneider et al., 2015). By manual curation, we selected and annotated the top 20 variant genes (Fig. 3B) that included three MFS-type transporter genes and an acetate permease homolog as well as several putative defense associated genes encoding a cell-wall glycoprotein (FG05_03352), peptidases (FG05_08075 and FG05_08141), and glucosidases (FG05_03387 and FG05_08265). A Zn₂Cys₆ transcription factor (FG05_03727), which shares the highest similarity to the yeast multidrug and oxidative stress resistance regulator STB5 (Laroche et al., 2006), was identified. Accordingly, we found several oxidative stress responsive genes encoding a molybdopterin oxidoreductase (FG05_02880), NADH-flavin oxidoreductase (FG05_08077) and a putative flavohemoglobin (FG05_04458). Notably, *FG05_03914* encoding a putative isochorismatase (ISC) gene was only expressed in *Fg* during infection of *Bd* roots. ISC-like effectors in filamentous pathogens are conserved virulence factors that can subvert plant salicylic acid (SA) pathway and interfere with host immunity (Liu et al., 2014). Resistance to biotrophic and hemi-biotrophic pathogens is usually conferred by the host SA pathway (Pieterse et al., 2012), and therefore, ISCs might be employed by *Fg* to support its hemi-biotrophic lifestyle to attenuate SA produced by the roots. The absence of *Fp* ISC transcription is consistent with the observation that host SA levels were not elevated in *Bd* plants at the early stages of infection (Powell et al., 2017). Host plants may activate tissue-specific defense signaling in response to below- and above-ground attacks (Lyons et al., 2015). To manipulate such defense responses, a fungal pathogen must evolve to a high flexibility for successful infection progressed in different host tissues. Despite the use of two fungal species, the data provided here may suggest shared infection strategies, including the interference with host defense, employed by *Fg* to support its belowground colonization.

DON influences different fungal processes in *Fg* during *Bd* root infection

The trichothecene mycotoxin DON has been shown to significantly inhibit *Bd* root growth (Pasquet et al., 2016). However, to the best of our knowledge, there has not been any study examining the effect of DON on different fungal processes in *Fg*. Therefore, we conducted an RNA-seq analysis by infecting *Bd* roots with DON producing and deficient strains to determine *Fg* genes whose expressions are modulated by DON. We first focused on genes up-regulated in the DON deficient Δ Tri5 mutant strain relative to WT. Of 973 genes differentially expressed between Δ Tri5 and WT, 432 genes were expressed at higher levels in Δ Tri5 (Fig. 4A, Suppl. data 12). These genes were subjected to functional enrichment analysis based on the MIPS FGDB (*Fusarium graminearum* Genome Database Functional Catalogue classification) (Güldener et al., 2006). This analysis showed that these genes are enriched for transport (FDR = 0.0016) of carbon-compounds, carbohydrates, and heavy metal ions, disease, virulence and defense (FDR = 0.005) and homeostasis of phosphate (FDR = 0.02). A relatively smaller portion of fungal pathogenicity and metabolism associated genes encoding CAZymes, SMEs, transporters and secreted proteins were induced in Δ Tri5 relative to WT (Fig. 4B-C, Fig. S2 and suppl. data 12-13). Only metabolic pathway genes encoding methyl- and glycol-transferases as well as phosphate, lipid and polyamine transporters were preferentially induced (Fig. S2). Host-derived phosphates, lipids as well as polyamines may influence *Fg* infection in multiple host tissue types (Gardiner et al., 2010; Zhang et al., 2016). Their enhancement might be due to a positive feedback to the lack of DON to balance the fungal metabolism and cell structure in root proliferation. In addition, 28 CAZyme encoding genes were expressed higher in Δ Tri5 than in WT and many of these were putative PCWDEs involved in lignin degradation (Fig. 4B). Lignin is one of the major barriers against fungal pathogens (Bhuiyan et al., 2009). Defense related or unrelated lignin content at fungal penetration sites might affect *Fg* intra-cellular progression (Zhang et al., 2016). In wheat roots, *Fg* colonization could be observed in lignin-rich vascular bundles (Bhandari et al., 2018; Wang et al., 2015). Therefore, the upregulation of lignin-degrading enzymes in Δ Tri5 could be beneficial to the fungus to compensate DON deficiency and assist root colonization. In line with this, we observed reduced lignin deposition in roots

colonized by Δ Tri5 relative to the roots either mock-inoculated or colonized by WT *Fg* (Fig. 5).

Seven of the 33 secreted protein genes induced in Δ Tri5, including a pathogenesis-related protein 1 (PR1) homolog (FG05_03109) and a putative cutinase (FG05_03457), may be considered putative effectors. The most differentially regulated SME genes in Δ Tri5 were tailoring enzyme genes encoding cytochrome P450s and oxidoreductases (Fig. S2A and S2B). Surprisingly, most DEGs with elevated transcripts levels in Δ Tri5 during root colonization (405 out of 432 genes) were expressed either significantly lower in WT during *Bd* root infection than *Fg* only or remained unchanged comparing to WT *in vitro* (Suppl. data 12). Besides, we noticed that none of the above-mentioned core genes was reduced in Δ Tri5 during infection, supporting the notion that these genes may contribute to infection more than others, independently of fungal DON production.

We next looked at the 541 significantly downregulated genes in Δ Tri5 relative to WT in roots. The most enriched functional categories during root infection were C-compound and carbohydrate metabolism (80 genes, FDR = 0.00008), disease, virulence and defense (16 genes, FDR = 0.01), secondary metabolism (27 genes, FDR = 0.01), protein or peptide degradation (26 genes, FDR = 0.01) and transport facilities (42 genes, FDR = 0.03) (Suppl. data 14). In addition, we found two sets of adjacent genes *FG05_02297-FG05_02309* and *FG05_08077-FG05_08084* that showed reduced expression in Δ Tri5. Of these two sets, the latter genes, which are part of the mycotoxin butanolide cluster, shared similar regulation pattern with the *Tri* (*tricothecene*) cluster genes during infection of wheat heads (Boedi et al., 2016). In the saprophytic fungus *Trichoderma arundinaceum*, the loss of tricothecene production likely contributed to an increase of fungal secondary metabolites (Lindo et al., 2019, 2018). Therefore, DON seems to affect fungal metabolism during *Fg* infection in *Bd* roots.

Previously, *Tri5* deletion was reported to lead to observable metabolic changes in *Fg* growing in rich medium, suggesting that DON might be linked to fungal physiology and

development (Chen et al. 2011). In line with this, we also found that several differentially regulated TF genes in Δ Tri5 during root infection were putative development-associated regulatory genes (Suppl. data 15). For example, FG05_08892 (MAT1-1-1) and FG05_05151 are known to be associated with sexual development (Kim et al., 2015), FG05_01139 (FgCBF1) is a predicted chromatin remodeling regulator (Guo et al., 2016), and FG05_03597 is homologous to *Aspergillus nidulans* FlibA, which is required for the control of mycelial proliferation and activation of asexual sporulation (Yu et al., 1996). However, when grown on MM, *Tri5* was barely expressed *in vitro*, and no DON could be detected by metabolic analysis (Fig. 6D). Δ Tri5 also did not show any growth defects (Chen et al., 2011). Together, our results suggest that Δ Tri5 may colonize the roots by utilizing a small set of genes not used by the WT fungus.

DON is produced during *Bd* root colonization but does not act as a virulence factor

DON is a virulence factor during infection of wheat heads by *Fg* (Wang et al., 2020). However, it is unknown if this mycotoxin could also act as a virulence factor during infection of *Bd* roots. To determine this, the infection process was monitored using a *Fg* strain expressing Tri5-GFP fusion driving by the native *Tri5* gene promoter (Gardiner et al., 2009). Strong GFP signals could be visualized two days post-inoculation (dpi) in inoculated roots, indicating that the infection was progressing, and DON production was initiated (Fig. 7A and 7B). Consistent with this observation, *Tri5* was highly induced at 2 dpi and remained at high levels at later time points (3, 5, and 7 dpi) in the WT isolate (Fig. 7C). Previously, a temporarily similar infection pattern by *Fg* was also observed in wheat seedling roots (Wang et al., 2015). *Fg* root infection of wheat triggers induction of systemic defense responses in above-ground parts of the plant (Wang et al., 2018). In addition, DON preferentially inhibits root growth in wheat, *Bd* and *Arabidopsis* plants (Gatti et al., 2019; Masuda et al., 2007; Pasquet et al., 2016), and has been proposed to act as a major virulence factor in the early stages of *Fg* root infection (Wang et al. 2018). To assess the role of DON during root infection, a *Fg Tri5* mutant was used in inoculation experiments together with WT and Tri5-

GFP strains. *Bd* roots infected by Δ Tri5 exhibited levels of lesion development that were like those caused by WT and Tri5-GFP at 7 dpi (Fig. 7D). Therefore, while DON is a virulence factor in *Fg* during FHB of wheat and is highly induced in roots, *Bd* root infection by *Fg* seems to be independent of DON production. Indeed, various phytopathogenic phenotypes have been described for *Fg* DON deficient mutants (Boenisch and Schäfer, 2011; Cuzick et al., 2008; Jansen et al., 2005). For instance, altered levels of DON have been shown to inhibit plant apoptosis-like programmed cell death (PCD) induced by heat stress in *Arabidopsis* (Diamond et al., 2013).

Phytohormone dynamics during *Fg* colonization of *Bd* roots

It is becoming increasingly evident that plant pathogens interfere with phytohormone pathways by producing plant hormones (Kazan and Lyons, 2014). However, pathogen-produced phytohormones have rarely been examined during root infections. We therefore next examined the transcriptome of *Fg* during *Bd* root infection, coupled with metabolic analyses, to determine putative phytohormone associated genes in *Fg* and their potential involvement in *Bd* root colonization.

JA produced by *Fg* is not associated with *Bd* root colonization

The oxylipin hormone jasmonic acid and its derived metabolites collectively known as jasmonates (JAs) are derived from lipid peroxidation and can affect both host and fungal physiological processes (Tsitsigiannis and Keller, 2007). Fungal oxylipin biosynthesis is catalyzed by lipoxygenases (LOXs) (Fischer and Keller, 2016). In *F. oxysporum*, FoxLOX was found to exhibit a multifunctional activity in oxylipins production, thus proposed to possess a function in JA pathways (Brodhun et al., 2013). Interestingly, we noticed that the *Fg* homolog of FoxLOX, *FG05_05046*, was expressed during *in vitro* growth, and highly induced during root infection (4.8 fold, Table S1). JA-regulated defenses in plants can be interrupted by pathogen-derived hormone analogs (Caarls et al., 2017; Patkar and Naqvi, 2017). The *Fg* genome does not contain a homolog of the *M. oryzae* antibiotic biosynthesis

monooxygenase (*Abm*), which converts host-derived JA into 12-hydroxyjasmonic acid (12OH-JA), thus attenuating rice blast disease resistance (Patkar and Naqvi, 2017). The Arabidopsis 2OG oxygenases (JOXs) are responsible for JA hydroxylation (Caarls et al., 2017; Smirnova et al., 2017). We identified ten homologous of Arabidopsis JOXs in *Fg* (Table S1). Of these, *FG05_08081* and *FG05_02301*, whose protein products share 22-26% identity to JOXs, were induced by 8.3 and 3.9 fold, respectively, during root infection. *FG05_08081* is present in the butanolide biosynthesis gene cluster, members of which were also significantly upregulated during root infection (Suppl. data 2). However, whether *FG05_08081* and *FG05_02301* are involved in JA degradation requires further analyses.

JA-associated host defense against *Fg* has been studied during FHB development. Inhibition of JA by DON at the bottom of wheat florets promotes fungal progression through rachis nodes (Bönnighausen et al., 2019). Furthermore, late activation of JA signaling during FHB has been proposed to correlate with a necrotrophic transition of *Fg* (Ding et al., 2011). To determine if JA levels change during root infection, we quantified JA levels in *Bd* roots either mock-treated or inoculated with WT *Fg* or Δ Tri5 strains by high performance liquid chromatography (HPLC). We found that 5 ng/mg dried material of JA was produced in *Fg* mycelia grown on MM (Fig. 6A). However, JA levels found in infected and control roots were much lower than 5 ng/mg and did not display any significant difference (Fig. 6A). Interestingly, however, higher levels of JA derivative methyl-JA (MeJA) were detectable in the roots infected by Δ Tri5 than those infected by WT (Fig. 6B). Thus, DON seems to inhibit MeJA production in the infected *Bd* roots. This is consistent with the observation in wheat heads where MeJA levels in the Δ Tri5-infected tissue were less than those infected with WT *Fg* (Bönnighausen et al., 2019).

SA may synergistically interact with *Bd* root defense-related metabolic pathways

SA is a major defense hormone typically associated with plant defense against biotrophic pathogens (Glazebrook, 2005). JA and SA accumulate at different basal levels in various wheat cultivars and antagonistically fine-tune host defense responses (Powell et al., 2017).

Therefore, we next focused on *Fg* responses to SA. Although the SA-pathway may be involved in host basal resistance against *Fg* (Ding et al., 2011; Makandar et al., 2010), SA-associated systemic acquired resistance (SAR) played no role in FHB resistance (Li and Yen, 2008). During infection of *Bd* seedlings by *Fp*, SA biosynthesis was induced (Powell et al., 2017). However, the function of SA during root infection by *Fg* remains elusive.

In *Arabidopsis*, SA biosynthesis during pathogen infection mainly relies on the intermediate chorismate processed by isochorismate synthase I (ICS1), the GH3 acyl adenylase-family enzyme PBS3, and the BAHD acyltransferase-family protein EPS1 (Torrens-Spence et al., 2019). The phenylalanine ammonia lyase (PAL) pathway also mediates SA synthesis through the conversion of benzoic acid or coumaric acid, but only contributes to a small portion of total SA production (Wildermuth et al., 2002). Similarly, some bacteria can directly convert isochorismate to SA by isochorismate pyruvate lyase (IPL) (Serino et al., 1995). While the host is believed to be the source of SA production in various plant-bacteria interactions, whether there is a fungal origin of SA remains unknown. In the *Fg* genome, we found two ICS1 homologs FG05_05195 and FG05_12934. Of these, *FG05_05195* was lowly expressed (FPKM < 0.5), but *FG05_12934* exhibited constitutive and high transcript levels during both *in vitro* growth and infection of *Bd* roots (Table S1). *Fg* also has an *EPS1* homolog, *FG05_00237*, which was significantly upregulated during *Bd* root infection (Table S1). *FG05_09331* encodes a protein sharing about 50% identity with PAL. *FG05_09331* expression decreased by 2.2-fold in infected roots relative to *Fg* grown *in vitro*. No PBS3 and IPL homologs could be identified in *Fg*. While expression patterns of *ICS1* and *EPS1* homologs may coincide with observed SA production by *Fg*, the absence of a *PBS3* homolog indicates other components or pathways could be involved in the SA biosynthesis. To determine if SA is produced in *Fg-Bd* interactions, we extracted metabolites from *Bd* roots with or without *Fg* inoculation and quantified SA levels by HPLC. In *Fg* mycelia, there was also about 0.01 µg SA per mg dry material (Fig. 6C). Most SA was found in the roots inoculated with WT *Fg*, followed by uninfected roots and the lowest levels in the Δ Tri5

infected roots (Fig. 6C), indicating a potential role for DON in regulating SA levels during root infection by *Fg* although it is difficult to estimate the exact contribution of *Bd* or *Fg* to the SA levels measured.

SA has a direct effect on *Fg* growth, most likely associated with active degradation of SA by fungal hydroxylases. In recent studies (Hao et al., 2019; Qi et al., 2019; Rocheleau et al., 2019), two proteins FGSG_08116 and FGSG_03657 have been characterized with a function in SA degradation. While transcription of both FGSG_08116 and FGSG_03657 can be induced by external SA, fungal virulence in wheat heads was only influenced by deletion of the former. The function of FGSG_03657 for SA degradation was not disabled in deletion mutants. Interestingly, *FG05_03657* (a.k.a. FGSG_03657) was exclusively induced during *Bd* root infection (Suppl. data 10), suggesting that it may have a role in regulating SA levels in infected roots.

To determine if *Fg* could possess additional putative SA hydroxylase genes (Hao et al., 2019), we searched the *Fg* genome and identified 28 homologs of the SA sensor and degradation protein Shy1 from *Ustilago maydis* with FgShyC displaying at least 20% identity to Shy1 (e-value < 10⁻⁵) (Fig. S3). This similarity is much higher than the values reported previously (Hao et al., 2019; Rabe et al., 2013), where the expansion of these putative proteins in *Fg* was supported by phylogenetic analysis (Fig. 8A). In our current transcriptome, *FG05_03657* and additional ten genes encoding putative SA hydroxylases were significantly upregulated in *Fg* during root infection (Fig. 8B). Therefore, SA biosynthesis by the pathogen as well as the host can contribute to the expression and regulation of these genes.

Expression of genes involved in the biosynthesis of other phytohormones in *Fg* during *Bd* roots infection

In addition to JA and SA, other phytohormones such as gibberellins (GAs), auxins (IAAs), ethylene (ET), cytokinins (CKs) and abscisic acid (ABA) also participate in modulating host defense signaling (Pieterse et al., 2012). GAs can be synthesized by a number of *Fusarium*

species but not by *Fg* due to the lack of a corresponding biosynthesis gene cluster (Cuomo et al., 2007). As a virulence factor, GA is restricted to the necrotrophic fungal pathogen *F. fujikuroi* (Wiemann et al., 2013) and is possibly involved in attenuating host JA signaling (Navarro et al., 2008). Under our inoculation conditions, GA was detected in the roots infected by WT *Fg* but not by the Δ Tri5 mutant (Fig. 6F), indicating an endogenous GA production in *Bd* roots upon *Fg* infection and a potential positive effect of DON on root GA production. Similarly, GA accumulates in wheat heads infected by *Fg* (Bönnighausen et al., 2019), thus *Fg* seems to trigger a GA-dependent response in host roots that is similar to one observed in wheat florets (Buhrow et al., 2016). However, the association between DON and GA production requires further investigations.

Fungal genes encoding indole-3-acetaldehyde dehydrogenases (*lad*) and tryptophan aminotransferases (*laaM*) were thought to be responsible for IAA production (Reineke et al., 2008). A possible third pathway for auxin biosynthesis could be mediated by *Fg* genes homologous to *YUCCA*, a key enzyme involved in plant auxin biosynthesis (Mano and Nemoto, 2012). Interestingly, only one of the three *Fg lad* gene homologs, *FG05_02773*, showed more than 4-fold induction in *Fg* during root infection as compared to *Fg* grown *in vitro* (Table S1). In contrast, two *laaM* homologs and a *YUCCA* homolog were significantly downregulated in *Fg* inoculated roots (Table S1). Thus, the strong induction of *FG05_02773* might coincide with the production of the auxin indole-3-acetic acid (IAA) by *Fg*, which could thereafter compromise the host auxin pathway. Fungal auxin biosynthesis plays a role in pathogenicity of several pathogens (Chanclud and Morel, 2016). In *Fg*, auxin was proposed to be a virulence factor (Svoboda et al., 2019). Indeed, *Fg* is able to synthesize IAA but also sensitive to exogenous application of IAA and its biosynthetic intermediates (Qi et al., 2016). However, *Fg* infection strongly inhibited IAA levels in roots infected by *Fg* WT or Δ Tri5 (Fig. 6E). This is contradictory to the findings in wheat where auxin levels increased during FHB (Wang et al., 2018). Factors such as host tissue types and hormone antagonists (Kazan and Manners, 2009) might be responsible for such differences.

Fg can exploit host ET signaling during colonization of both dicotyledonous and monocotyledonous plants and is believed to be capable of producing ET to counteract host defense pathways (Chen et al., 2009). However, rather than ET forming enzymes (EFE), *Fg* was thought to use pathways incorporating 1-aminocyclopropane carboxylic acids (ACC) as precursors for ET biosynthesis (Svoboda et al., 2019). Although an enzymatic function for two of the five ACC enzymes encoded by the *Fg* genome could be confirmed, fungal mutants for these genes showed no defect in pathogenicity on wheat (Svoboda et al., 2019). We looked at the expression of all these five genes, including the three annotated ACC synthase genes, *FG05_05184* (*ACS1*), *FG05_07606* (*ACS2*), *FG05_13587* (*ACS3*), and two ACC deaminase (*ACD*) genes, *FG05_02678* and *FG05_12669*, but found no differential expression during *Bd* root infections (Table S1), suggesting that pathogen produced ET may not be involved in root infection.

Biosynthesis of fungal cytokinins, which can be mediated by either one or both of the fungal transfer RNA-isopentenyl transferases (tRNA-IPT) and the Lonely Guy (LOG) enzyme, is associated with host immunity and nutrient modulation and maintenance of hemi-biotrophic lifestyles during infection (Spallek et al., 2018). Unlike many other *Fusarium* species, the *Fg* genome contains only one tRNA-IPT gene homolog (*FG05_09015*) (Sørensen et al., 2018). This gene was not differentially regulated (Table S1) and only moderately expressed (FPKM < 10) during both root infections and *in vitro* growth on MM.

Exogenous ABA has no effects on disease development, *Fg* toxin production or defense hormone levels in *Fg*-challenged wheat heads, but can promote fungal hydrolase and cytoskeletal reorganization genes induced early during infection and increase wheat's susceptibility to FHB (Buhrow et al., 2016). The elucidation of fungal genes responsible for ABA production in the necrotrophic pathogen *Botrytis cinerea* and a few others has led to the hypothesis that a conserved ABA biosynthesis pathway exists in fungi (Lievens et al., 2017). In *B. cinerea*, such pathway involves four clustered genes *BcABA1-4* and a sesquiterpene cyclase gene *BcSTC5* (Izquierdo-Bueno et al., 2018). Fungal ABA was shown to act as a

virulence factor in *M. oryzae*, which also harbors a direct ABA biosynthesis pathway but lacks a BsABA3 ortholog (Spence et al., 2015). Similar to *M. oryzae*, we could identify homologs of only BcABA1, 2 and 4 by BLASTp in *Fg*. In the current root transcriptome, none of these genes was differentially regulated (Table S1). Overall, while fungal GAs and IAAs might be associated with *Fg* root infections, it is unlikely that ETs, CKs and ABAs are involved in *Bd* root infection by *Fg*.

Conclusions

The results presented here provide a detailed overview of root infection strategies employed by *Fg*, an important cereal fungal pathogen. The transcriptional regulation of pathogen metabolic pathways, virulence factors and signalling events during root infection show both unique and common features to those employed by *Fg* when infecting above-ground tissues. The mycotoxin DON, although not required for fungal virulence, produced during root infection appears to broadly affect various fungal processes and interplay with host responses. Expressions of several fungal stress and defence genes might help the pathogen to effectively deal with plant defence responses. In line with this, fungal JA, IAA and, in particular SA, seem to be used to interfere with root defenses. The findings presented in this paper will be useful for dissecting the mechanism of *Fg* belowground lifestyle and the development of novel plant protection strategies.

Acknowledgements

We thank Di Xiao and Dr. Jonathan Powell for technical assistance. Yi Ding was the recipient of a post-doctoral fellowship from the Commonwealth Scientific and Industrial Research Organization Research Office.

Conflict of interest

All authors declared no conflict of interest.

References

718 Abou Ammar, G., Tryono, R., Döll, K., Karlovsky, P., Deising, H.B., Wirsal, S.G.R., 2013.
719 Identification of ABC Transporter Genes of *Fusarium graminearum* with Roles in Azole
720 Tolerance and/or Virulence. PLoS One 8, e79042.
721 <https://doi.org/10.1371/journal.pone.0079042>

722 Audenaert, K., Vanheule, A., Höfte, M., Haesaert, G., 2013. Deoxynivalenol: A major player
723 in the multifaceted response of *Fusarium* to its environment. Toxins (Basel). 6, 1–19.
724 <https://doi.org/10.3390/toxins6010001>

725 Bhandari, D.R., Wang, Q., Li, B., Friedt, W., Römpf, A., Spengler, B., Gottwald, S., 2018.
726 Histology-guided high-resolution AP-SMALDI mass spectrometry imaging of wheat-
727 *Fusarium graminearum* interaction at the root-shoot junction. Plant Methods 14, 1–13.
728 <https://doi.org/10.1186/s13007-018-0368-6>

729 Bhuiyan, N.H., Selvaraj, G., Wei, Y., King, J., 2009. Role of lignification in plant defense.
730 Plant Signal. Behav. 4, 158–159. <https://doi.org/10.4161/psb.4.2.7688>

731 Boedi, S., Berger, H., Sieber, C., Münsterkötter, M., Maluku, I., Warth, B., Sulyok, M.,
732 Lemmens, M., Schuhmacher, R., Güldener, U., Strauss, J., 2016. Comparison of
733 *Fusarium graminearum* transcriptomes on living or dead wheat differentiates substrate-
734 responsive and defense-responsive genes. Front. Microbiol. 7, 1–24.
735 <https://doi.org/10.3389/fmicb.2016.01113>

736 Boenisch, M.J., Schäfer, W., 2011. *Fusarium graminearum* forms mycotoxin producing
737 infection structures on wheat. BMC Plant Biol. 11, 110. <https://doi.org/10.1186/1471-2229-11-110>

738 2229-11-110

739 Bönighausen, J., Schauer, N., Schäfer, W., Bormann, J., 2019. Metabolic profiling of wheat
740 rachis node infection by *Fusarium graminearum* – decoding deoxynivalenol-dependent
741 susceptibility. New Phytol. 221, 459–469. <https://doi.org/10.1111/nph.15377>

742 Brauer, E.K., Balcerzak, M., Rocheleau, H., Leung, W., Schernthaner, J., Subramaniam, R.,

743 Ouellet, T., 2020. Genome Editing of a Deoxynivalenol-Induced Transcription Factor
744 Confers Resistance to *Fusarium graminearum* in Wheat. *Mol. Plant-Microbe Interact.*
745 33, 553–560. <https://doi.org/10.1094/MPMI-11-19-0332-R>

746 Brodhun, F., Cristobal-Sarramian, A., Zabel, S., Newie, J., Hamberg, M., Feussner, I., 2013.
747 An Iron 13S-Lipoxygenase with an α -Linolenic Acid Specific Hydroperoxidase Activity
748 from *Fusarium oxysporum*. *PLoS One* 8. <https://doi.org/10.1371/journal.pone.0064919>

749 Brown, N.A., Antoniwi, J., Hammond-Kosack, K.E., 2012. The predicted secretome of the
750 plant pathogenic fungus *Fusarium graminearum*: A refined comparative analysis. *PLoS*
751 *One* 7. <https://doi.org/10.1371/journal.pone.0033731>

752 Brown, N.A., Evans, J., Mead, A., Hammond-Kosack, K.E., 2017. A spatial temporal analysis
753 of the *Fusarium graminearum* transcriptome during symptomless and symptomatic
754 wheat infection. *Mol. Plant Pathol.* 18, 1295–1312. <https://doi.org/10.1111/mpp.12564>

755 Buhrow, L.M., Cram, D., Tulpan, D., Foroud, N.A., Loewen, M.C., 2016. Exogenous abscisic
756 acid and gibberellic acid elicit opposing effects on *Fusarium graminearum* infection in
757 wheat. *Phytopathology* 106, 986–996. <https://doi.org/10.1094/PHYTO-01-16-0033-R>

758 Caarls, L., Elberse, J., Awwanah, M., Ludwig, N.R., De Vries, M., Zeilmaker, T., Van Wees,
759 S.C.M., Schuurink, R.C., Van den Ackerveken, G., 2017. Arabidopsis JASMONATE-
760 INDUCED OXYGENASES down-regulate plant immunity by hydroxylation and
761 inactivation of the hormone jasmonic acid. *Proc. Natl. Acad. Sci. U. S. A.* 114, 6388–
762 6393. <https://doi.org/10.1073/pnas.1701101114>

763 Chanclud, E., Morel, J.-B., 2016. Plant hormones: a fungal point of view. *Mol. Plant Pathol.*
764 17, 1289–1297. <https://doi.org/10.1111/mpp.12393>

765 Chen, F., Zhang, J., Song, X., Yang, J., Li, H., Tang, H., Liao, Y.C., 2011. Combined
766 metabonomic and quantitative real-time PCR analyses reveal systems metabolic
767 changes of *Fusarium graminearum* induced by Tri5 gene deletion. *J. Proteome Res.* 10,

768 2273–2285. <https://doi.org/10.1021/pr101095t>

769 Chen, X., Steed, A., Travella, S., Keller, B., Nicholson, P., 2009. *Fusarium graminearum*
770 exploits ethylene signalling to colonize dicotyledonous and monocotyledonous plants.
771 *New Phytol.* 182, 975–983. <https://doi.org/10.1111/j.1469-8137.2009.02821.x>

772 Chen, Y., Kistler, H.C., Ma, Z., 2019. *Fusarium graminearum* Trichothecene Mycotoxins:
773 Biosynthesis, Regulation, and Management . *Annu. Rev. Phytopathol.* 57, 1–25.
774 <https://doi.org/10.1146/annurev-phyto-082718-100318>

775 Coleman, J.J., Mylonakis, E., 2009. Efflux in fungi: La pièce de résistance. *PLoS Pathog.* 5.
776 <https://doi.org/10.1371/journal.ppat.1000486>

777 Cuzick, A., Urban, M., Hammond-Kosack, K., 2008. *Fusarium graminearum* gene deletion
778 mutants *map1* and *tri5* reveal similarities and differences in the pathogenicity
779 requirements to cause disease on *Arabidopsis* and wheat floral tissue. *New Phytol.* 177,
780 990–1000. <https://doi.org/10.1111/j.1469-8137.2007.02333.x>

781 De Jonge, R., Esse, H.P. Van, Kombrink, A., Shinya, T., Desaki, Y., Bours, R., Krol, S. Van
782 Der, Shibuya, N., Joosten, M.H. a J., Thomma, B.P.H.J., 2010. Immunity in Plants.
783 *Science* (80-.). 329, 953–955. <https://doi.org/10.1126/science.1190859>

784 Desmond, O.J., Manners, J.M., Stephens, A.E., Maclean, D.J., Schenk, P.M., Gardiner,
785 D.M., Munn, A.L., Kazan, K., 2008. The *Fusarium* mycotoxin deoxynivalenol elicits
786 hydrogen peroxide production, programmed cell death and defence responses in wheat.
787 *Mol. Plant Pathol.* 9, 435–445. <https://doi.org/10.1111/j.1364-3703.2008.00475.x>

788 Diamond, M., Reape, T.J., Rocha, O., Doyle, S.M., Kacprzyk, J., Doohan, F.M., McCabe,
789 P.F., 2013. The *Fusarium* Mycotoxin Deoxynivalenol Can Inhibit Plant Apoptosis-Like
790 Programmed Cell Death. *PLoS One* 8. <https://doi.org/10.1371/journal.pone.0069542>

791 Ding, L., Xu, H., Yi, H., Yang, L., Kong, Z., Zhang, L., Xue, S., Jia, H., Ma, Z., 2011.
792 Resistance to hemi-biotrophic *f. graminearum* infection is associated with coordinated

793 and ordered expression of diverse defense signaling pathways. PLoS One 6.
794 <https://doi.org/10.1371/journal.pone.0019008>

795 Ding, Y., Gardiner, D.M., Xiao, D., Kazan, K., 2020. Regulators of nitric oxide signaling
796 triggered by host perception in a plant pathogen. Proc. Natl. Acad. Sci. 201918977.
797 <https://doi.org/10.1073/pnas.1918977117>

798 Divon, H.H., Fluhr, R., 2007. Nutrition acquisition strategies during fungal infection of plants.
799 FEMS Microbiol. Lett. 266, 65–74. <https://doi.org/10.1111/j.1574-6968.2006.00504.x>

800 Dutartre, L., Hilliou, F., Feyereisen, R., 2012. Phylogenomics of the benzoxazinoid
801 biosynthetic pathway of Poaceae: Gene duplications and origin of the Bx cluster. BMC
802 Evol. Biol. 12. <https://doi.org/10.1186/1471-2148-12-64>

803 Fischer, G.J., Keller, N.P., 2016. Production of cross-kingdom oxylipins by pathogenic fungi:
804 An update on their role in development and pathogenicity. J. Microbiol. 54, 254–264.
805 <https://doi.org/10.1007/s12275-016-5620-z>

806 Fitzgerald, T.L., Powell, J.J., Schneebeli, K., Hsia, M.M., Gardiner, D.M., Bragg, J.N.,
807 McIntyre, C.L., Manners, J.M., Ayliffe, M., Watt, M., Vogel, J.P., Henry, R.J., Kazan, K.,
808 2015. Brachypodium as an emerging model for cereal–pathogen interactions. Ann. Bot.
809 115, 717–731. <https://doi.org/10.1093/aob/mcv010>

810 Foroud, N.A., Pordel, R., Goyal, R.K., Ryabova, D., Eranthodi, A., Chatterton, S., Kovalchuk,
811 I., 2019. Chemical Activation of the Ethylene Signaling Pathway Promotes Fusarium
812 graminearum Resistance in Detached Wheat Heads. Phytopathology 109, 796–803.
813 <https://doi.org/10.1094/PHYTO-08-18-0286-R>

814 Gardiner, D.M., Benfield, A.H., Stiller, J., Stephen, S., Aitken, K., Liu, C., Kazan, K., 2018. A
815 high-resolution genetic map of the cereal crown rot pathogen Fusarium
816 pseudograminearum provides a near-complete genome assembly. Mol. Plant Pathol.
817 19, 217–226. <https://doi.org/10.1111/mpp.12519>

818 Gardiner, D.M., Kazan, K., Manners, J.M., 2009. Nutrient profiling reveals potent inducers of
819 trichothecene biosynthesis in *Fusarium graminearum*. *Fungal Genet. Biol.* 46, 604–613.
820 <https://doi.org/10.1016/j.fgb.2009.04.004>

821 Gardiner, D.M., Kazan, K., Praud, S., Torney, F.J., Rusu, A., Manners, J.M., 2010. Early
822 activation of wheat polyamine biosynthesis during *Fusarium* head blight implicates
823 putrescine as an inducer of trichothecene mycotoxin production. *BMC Plant Biol.* 10,
824 289. <https://doi.org/10.1186/1471-2229-10-289>

825 Gardiner, D.M., Stephens, A.E., Munn, A.L., Manners, J.M., 2013. An ABC pleiotropic drug
826 resistance transporter of *fusarium graminearum* with a role in crown and root diseases
827 of wheat. *FEMS Microbiol. Lett.* 348, 36–45. <https://doi.org/10.1111/1574-6968.12240>

828 Gardiner, D.M., Stiller, J., Kazan, K., 2014. Genome sequence of *Fusarium graminearum*
829 isolate CS3005. *Genome Announc.* 2, e00227-14-e00227-14.
830 <https://doi.org/10.1128/genomeA.00227-14>

831 Gatti, M., Cambon, F., Tassy, C., Macadre, C., Guerard, F., Langin, T., Dufresne, M., 2019.
832 The *Brachypodium distachyon* UGT Bradi5gUGT03300 confers type II fusarium head
833 blight resistance in wheat. *Plant Pathol.* 68, 334–343. <https://doi.org/10.1111/ppa.12941>

834 Gatti, M., Cambon, F., Tassy, C., Macadre, C., Guerard, F., Langin, T., Dufresne, M., 2018.
835 The *Brachypodium distachyon* UGT Bradi5gUGT03300 confers type II Fusarium Head
836 Blight resistance in wheat. *Plant Pathol.* <https://doi.org/10.1111/ppa.12941>

837 Glazebrook, J., 2005. Contrasting Mechanisms of Defense Against Biotrophic and
838 Necrotrophic Pathogens. *Annu. Rev. Phytopathol.* 43, 205–227.
839 <https://doi.org/10.1146/annurev.phyto.43.040204.135923>

840 Götz, S., García-Gómez, J.M., Terol, J., Williams, T.D., Nagaraj, S.H., Nueda, M.J., Robles,
841 M., Talón, M., Dopazo, J., Conesa, A., 2008. High-throughput functional annotation and
842 data mining with the Blast2GO suite. *Nucleic Acids Res.*

843 <https://doi.org/10.1093/nar/gkn176>

844 Güldener, U., Mannhaupt, G., Münsterkötter, M., Haase, D., Oesterheld, M., Stümpflen, V.,
845 Mewes, H.-W., Adam, G., 2006. FGDB: a comprehensive fungal genome resource on
846 the plant pathogen *Fusarium graminearum*. *Nucleic Acids Res.* 34, D456-8.
847 <https://doi.org/10.1093/nar/gkj026>

848 Guo, L., Breakspear, A., Zhao, G., Gao, L., Kistler, H.C., Xu, J.R., Ma, L.J., 2016.
849 Conservation and divergence of the cyclic adenosine monophosphate-protein kinase A
850 (cAMP-PKA) pathway in two plant-pathogenic fungi: *Fusarium graminearum* and
851 *F.verticillioides*. *Mol. Plant Pathol.* 17, 196–209. <https://doi.org/10.1111/mpp.12272>

852 Hao, G., Naumann, T.A., Vaughan, M.M., McCormick, S., Usgaard, T., Kelly, A., Ward, T.J.,
853 2019. Characterization of a *Fusarium graminearum* salicylate hydroxylase. *Front.*
854 *Microbiol.* 10, 1–11. <https://doi.org/10.3389/fmicb.2018.03219>

855 Harris, L.J., Balcerzak, M., Johnston, A., Schneiderman, D., Ouellet, T., 2016. Host-
856 preferential *Fusarium graminearum* gene expression during infection of wheat, barley,
857 and maize. *Fungal Biol.* 120, 111–123. <https://doi.org/10.1016/j.funbio.2015.10.010>

858 Hart, T., Komori, H.K., LaMere, S., Podshivalova, K., Salomon, D.R., 2013. Finding the
859 active genes in deep RNA-seq gene expression studies. *BMC Genomics* 14.
860 <https://doi.org/10.1186/1471-2164-14-778>

861 Hellin, P., King, R., Urban, M., Hammond-Kosack, K.E., Legrève, A., 2018. The adaptation
862 of *Fusarium culmorum* to DMI fungicides is mediated by major transcriptome
863 modifications in response to azole fungicide, including the overexpression of a PDR
864 transporter (FcABC1). *Front. Microbiol.* 9, 1–15.
865 <https://doi.org/10.3389/fmicb.2018.01385>

866 Henkes, G.J., Jousset, A., Bonkowski, M., Thorpe, M.R., Scheu, S., Lanoue, A., Schurr, U.,
867 Röse, U.S.R., 2011. *Pseudomonas fluorescens* CHA0 maintains carbon delivery to

868 Fusarium graminearum-infected roots and prevents reduction in biomass of barley
869 shoots through systemic interactions. J. Exp. Bot. 62, 4337–4344.
870 <https://doi.org/10.1093/jxb/err149>

871 Hohn, T.M., Beremand, P.D., 1989. Isolation and nucleotide sequence of a sesquiterpene
872 cyclase gene from the trichothecene-producing fungus Fusarium sporotrichioides. Gene
873 79, 131–138. [https://doi.org/10.1016/0378-1119\(89\)90098-X](https://doi.org/10.1016/0378-1119(89)90098-X)

874 Izquierdo-Bueno, I., González-Rodríguez, V.E., Simon, A., Dalmais, B., Pradier, J.M., Le
875 Pêcheur, P., Mercier, A., Walker, A.S., Garrido, C., Collado, I.G., Viaud, M., 2018.
876 Biosynthesis of abscisic acid in fungi: identification of a sesquiterpene cyclase as the
877 key enzyme in Botrytis cinerea. Environ. Microbiol. 20, 2469–2482.
878 <https://doi.org/10.1111/1462-2920.14258>

879 Jansen, C., von Wettstein, D., Schäfer, W., Kogel, K.-H., Felk, A., Maier, F.J., 2005. Infection
880 patterns in barley and wheat spikes inoculated with wild-type and trichodiene synthase
881 gene disrupted Fusarium graminearum. Proc. Natl. Acad. Sci. U. S. A. 102, 16892–7.
882 <https://doi.org/10.1073/pnas.0508467102>

883 Jashni, M.K., Dols, I.H.M., Iida, Y., Boeren, S., Beenen, H.G., Mehrabi, R., 2015. Synergistic
884 Action of a Metalloprotease and a Serine Protease from Fusarium oxysporum f. sp.
885 lycopersici Cleaves Chitin-Binding Tomato Chitinases, Reduces Their Antifungal
886 Activity, and Enhances Fungal Virulence 28, 996–1008.

887 Kazan, K., Gardiner, D.M., 2018a. Transcriptomics of cereal–Fusarium graminearum
888 interactions: what we have learned so far. Mol. Plant Pathol. 19, 764–778.
889 <https://doi.org/10.1111/mpp.12561>

890 Kazan, K., Gardiner, D.M., 2018b. Fusarium crown rot caused by Fusarium
891 pseudograminearum in cereal crops: recent progress and future prospects. Mol. Plant
892 Pathol. 19, 1547–1562. <https://doi.org/10.1111/mpp.12639>

- 893 Kazan, K., Lyons, R., 2014. Intervention of Phytohormone Pathways by Pathogen Effectors.
894 Plant Cell 26, 2285–2309. <https://doi.org/10.1105/tpc.114.125419>
- 895 Kazan, K., Manners, J.M., 2009. Linking development to defense: auxin in plant-pathogen
896 interactions. Trends Plant Sci. 14, 373–382.
897 <https://doi.org/10.1016/j.tplants.2009.04.005>
- 898 Khan, N.U., Liu, M., Yang, X., Qiu, D., 2016. Fungal elicitor MoHrip2 induces disease
899 resistance in rice leaves, triggering stress-related pathways. PLoS One 11, 1–14.
900 <https://doi.org/10.1371/journal.pone.0158112>
- 901 Kim, H.-K., Jo, S.-M., Kim, G.-Y., Kim, D.-W., Kim, Y.-K., Yun, S.-H., 2015. A Large-Scale
902 Functional Analysis of Putative Target Genes of Mating-Type Loci Provides Insight into
903 the Regulation of Sexual Development of the Cereal Pathogen *Fusarium graminearum*.
904 PLoS Genet. 11, e1005486. <https://doi.org/10.1371/journal.pgen.1005486>
- 905 Lanoue, A., Burlat, V., Henkes, G.J., Koch, I., Schurr, U., Röse, U.S.R., 2010. De novo
906 biosynthesis of defense root exudates in response to *Fusarium* attack in barley. New
907 Phytol. 185, 577–588. <https://doi.org/10.1111/j.1469-8137.2009.03066.x>
- 908 Larochelle, M., Drouin, S., Robert, F., Turcotte, B., 2006. Oxidative Stress-Activated Zinc
909 Cluster Protein Stb5 Has Dual Activator/Repressor Functions Required for Pentose
910 Phosphate Pathway Regulation and NADPH Production. Mol. Cell. Biol. 26, 6690–6701.
911 <https://doi.org/10.1128/mcb.02450-05>
- 912 Lee, J., Son, H., Lee, S., Park, A.R., Lee, Y.W., 2010. Development of a conditional gene
913 expression system using a zearalenone-inducible promoter for the ascomycete fungus
914 *Gibberella zeae*. Appl. Environ. Microbiol. 76, 3089–3096.
915 <https://doi.org/10.1128/AEM.02999-09>
- 916 Li, G., Yen, Y., 2008. Jasmonate and ethylene signaling pathway may mediate *Fusarium*
917 head blight resistance in wheat. Crop Sci. 48, 1888–1896.

918 <https://doi.org/10.2135/cropsci2008.02.0097>

919 Liang, Y., Cui, S., Tang, X., Zhang, Y., Qiu, D., Zeng, H., Guo, L., Yuan, J., Yang, X., 2018.

920 An Asparagine-Rich Protein Nbnrp1 Modulate *Verticillium dahliae* Protein PevD1-

921 Induced Cell Death and Disease Resistance in *Nicotiana benthamiana*. *Front. Plant Sci.*

922 9, 303. <https://doi.org/10.3389/fpls.2018.00303>

923 Lievens, L., Pollier, J., Goossens, A., Beyaert, R., Staal, J., 2017. Abscissic acid as pathogen

924 effector and immune regulator. *Front. Plant Sci.* 8, 1–15.

925 <https://doi.org/10.3389/fpls.2017.00587>

926 Lindo, L., McCormick, S.P., Cardoza, R.E., Brown, D.W., Kim, H.S., Alexander, N.J., Proctor,

927 R.H., Gutiérrez, S., 2018. Effect of deletion of a trichothecene toxin regulatory gene on

928 the secondary metabolism transcriptome of the saprotrophic fungus *Trichoderma*

929 *arundinaceum*. *Fungal Genet. Biol.* 119, 29–46.

930 <https://doi.org/10.1016/j.fgb.2018.08.002>

931 Lindo, L., McCormick, S.P., Cardoza, R.E., Kim, H.S., Brown, D.W., Alexander, N.J., Proctor,

932 R.H., Gutiérrez, S., 2019. Role of *Trichoderma arundinaceum* tri10 in regulation of

933 terpene biosynthetic genes and in control of metabolic flux. *Fungal Genet. Biol.* 122,

934 31–46. <https://doi.org/10.1016/j.fgb.2018.11.001>

935 Liu, T., Song, T., Zhang, X., Yuan, H., Su, L., Li, W., Xu, J., Liu, S., Chen, L., Chen, T.,

936 Zhang, M., Gu, L., Zhang, B., Dou, D., 2014. Unconventionally secreted effectors of two

937 filamentous pathogens target plant salicylate biosynthesis. *Nat. Commun.* 5.

938 <https://doi.org/10.1038/ncomms5686>

939 Lo Presti, L., Lanver, D., Schweizer, G., Tanaka, S., Liang, L., Tollot, M., Zuccaro, A.,

940 Reissmann, S., Kahmann, R., 2015. Fungal Effectors and Plant Susceptibility. *Annu.*

941 *Rev. Plant Biol.* 66, 513–545. <https://doi.org/10.1146/annurev-arplant-043014-114623>

942 Lu, S., Edwards, M.C., 2016. Genome-Wide Analysis of Small Secreted Cysteine-Rich

943 Proteins Identifies Candidate Effector Proteins Potentially Involved in *Fusarium*
 944 *graminearum*-Wheat Interactions. *Phytopathology* 106, 166–176.
 945 <https://doi.org/10.1094/PHYTO-09-15-0215-R>

946 Luo, K., Rocheleau, H., Qi, P.F., Zheng, Y.L., Zhao, H.Y., Ouellet, T., 2016. Indole-3-acetic
 947 acid in *Fusarium graminearum*: Identification of biosynthetic pathways and
 948 characterization of physiological effects. *Fungal Biol.* 120, 1135–1145.
 949 <https://doi.org/10.1016/j.funbio.2016.06.002>

950 Lyons, R., Stiller, J., Powell, J., Rusu, A., Manners, J.M., Kazan, K., 2015. *Fusarium*
 951 *oxysporum* triggers tissue-specific transcriptional reprogramming in *Arabidopsis*
 952 *thaliana*. *PLoS One* 10, 1–23. <https://doi.org/10.1371/journal.pone.0121902>

953 Lysøe, E., Seong, K., Kistler, H.C., 2011. The Transcriptome of *Fusarium graminearum*
 954 During the Infection of Wheat. *Mol. Plant-Microbe Interact.* 24, 995–1000.
 955 <https://doi.org/10.1094/MPMI-02-11-0038>

956 Ma, L.-J., Geiser, D.M., Proctor, R.H., Rooney, A.P., O'Donnell, K., Trail, F., Gardiner, D.M.,
 957 Manners, J.M., Kazan, K., 2013. *Fusarium* Pathogenomics. *Annu. Rev. Microbiol.* 67,
 958 399–416. <https://doi.org/10.1146/annurev-micro-092412-155650>

959 Makandar, R., Nalam, V., Chaturvedi, R., Jeannotte, R., Sparks, A.A., Shah, J., 2010.
 960 Involvement of Salicylate and Jasmonate Signaling Pathways in *Arabidopsis* Interaction
 961 with *Fusarium graminearum*. *Mol. Plant-Microbe Interact.* 23, 861–870.
 962 <https://doi.org/10.1094/MPMI-23-7-0861>

963 Malz, S., Grell, M.N., Thrane, C., Maier, F.J., Rosager, P., Felk, A., Albertsen, K.S., Salomon,
 964 S., Bohn, L., Schäfer, W., Giese, H., 2005. Identification of a gene cluster responsible
 965 for the biosynthesis of aurofusarin in the *Fusarium graminearum* species complex.
 966 *Fungal Genet. Biol.* 42, 420–433. <https://doi.org/10.1016/j.fgb.2005.01.010>

967 Mano, Y., Nemoto, K., 2012. The pathway of auxin biosynthesis in plants. *J. Exp. Bot.* 63,

968 2853–2872. <https://doi.org/10.1093/jxb/ers091>

969 Marcel, S., Sawers, R., Oakeley, E., Angliker, H., Paszkowski, U., 2010. Tissue-Adapted
 970 Invasion Strategies of the Rice Blast Fungus *Magnaporthe oryzae*. *Plant Cell* 22, 3177–
 971 3187. <https://doi.org/10.1105/tpc.110.078048>

972 Marshall, R., Kombrink, A., Motteram, J., Loza-Reyes, E., Lucas, J., Hammond-Kosack, K.E.,
 973 Thomma, B.P.H.J., Rudd, J.J., 2011. Analysis of Two in Planta Expressed LysM
 974 Effector Homologs from the Fungus *Mycosphaerella graminicola* Reveals Novel
 975 Functional Properties and Varying Contributions to Virulence on Wheat. *Plant Physiol.*
 976 156, 756–769. <https://doi.org/10.1104/pp.111.176347>

977 Masuda, D., Ishida, M., Yamaguchi, K., Yamaguchi, I., Kimura, M., Nishiuchi, T., 2007.
 978 Phytotoxic effects of trichothecenes on the growth and morphology of *Arabidopsis*
 979 *thaliana*. *J. Exp. Bot.* 58, 1617–1626. <https://doi.org/10.1093/jxb/erl298>

980 Mazaheri-Naeini, M., Sabbagh, S.K., Martinez, Y., Séjalon-Delmas, N., Roux, C., 2015.
 981 Assessment of *Ustilago maydis* as a fungal model for root infection studies. *Fungal Biol.*
 982 119, 145–153. <https://doi.org/10.1016/j.funbio.2014.12.002>

983 Mentlak, T.A., Kombrink, A., Shinya, T., Ryder, L.S., Otomo, I., Saitoh, H., Terauchi, R.,
 984 Nishizawa, Y., Shibuya, N., Thomma, B.P.H.J., Talbot, N.J., 2012. Effector-Mediated
 985 Suppression of Chitin-Triggered Immunity by *Magnaporthe oryzae* Is Necessary for
 986 Rice Blast Disease. *Plant Cell* 24, 322–335. <https://doi.org/10.1105/tpc.111.092957>

987 Moschetti, I., Faoro, F., Moro, S., Sabbadin, D., Sella, L., Favaron, F., D'Ovidio, R., 2015. The
 988 xylanase inhibitor TAXI-III counteracts the necrotic activity of a *Fusarium graminearum*
 989 xylanase in vitro and in durum wheat transgenic plants. *Mol. Plant Pathol.* 16, 583–592.
 990 <https://doi.org/10.1111/mpp.12215>

991 Mudge, A.M., Dill-Macky, R., Dong, Y., Gardiner, D.M., White, R.G., Manners, J.M., 2006. A
 992 role for the mycotoxin deoxynivalenol in stem colonisation during crown rot disease of

993 wheat caused by *Fusarium graminearum* and *Fusarium pseudograminearum*. *Physiol.*
994 *Mol. Plant Pathol.* 69, 73–85. <https://doi.org/10.1016/j.pmpp.2007.01.003>

995 Navarro, L., Bari, R., Achard, P., Lisón, P., Nemri, A., Harberd, N.P., Jones, J.D.G., 2008.
996 DELLAs Control Plant Immune Responses by Modulating the Balance of Jasmonic Acid
997 and Salicylic Acid Signaling. *Curr. Biol.* 18, 650–655.
998 <https://doi.org/10.1016/j.cub.2008.03.060>

999 Oide, S., Berthiller, F., Wiesenberger, G., Adam, G., Turgeon, B.G., 2014. Individual and
1000 combined roles of malonichrome, ferricrocin, and TAFC siderophores in *Fusarium*
1001 *graminearum* pathogenic and sexual development. *Front. Microbiol.* 5, 1–15.
1002 <https://doi.org/10.3389/fmicb.2014.00759>

1003 Pan, Y., Liu, Z., Rocheleau, H., Fauteux, F., Wang, Y., McCartney, C., Ouellet, T., 2018.
1004 Transcriptome dynamics associated with resistance and susceptibility against fusarium
1005 head blight in four wheat genotypes. *BMC Genomics* 19, 1–26.
1006 <https://doi.org/10.1186/s12864-018-5012-3>

1007 Pasquet, J.-C., Changenet, V., Macadré, C., Boex-Fontvieille, E., Soulhat, C., Bouchabké-
1008 Coussa, O., Dalmais, M., Atanasova-Pénichon, V., Bendahmane, A., Saindrenan, P.,
1009 Dufresne, M., 2016. A Brachypodium UDP-glycosyltransferase confers root tolerance to
1010 deoxynivalenol and resistance to *Fusarium* infection. *Plant Physiol.* 172, 559–574.
1011 <https://doi.org/10.1104/pp.16.00371>

1012 Pasquet, J.-C.C., Chaouch, S., Macadré, C., Balzergue, S., Huguet, S., Martin-Magniette,
1013 M.-L.L., Bellvert, F., Deguerce, X., Thareau, V., Heintz, D., Saindrenan, P., Dufresne,
1014 M., 2014. Differential gene expression and metabolomic analyses of *Brachypodium*
1015 *distachyon* infected by deoxynivalenol producing and non-producing strains of
1016 *Fusarium graminearum*. *BMC Genomics* 15, 1–17. [https://doi.org/10.1186/1471-2164-](https://doi.org/10.1186/1471-2164-15-629)
1017 15-629

1018 Patkar, R.N., Naqvi, N.I., 2017. Fungal manipulation of hormone-regulated plant defense.

1019 PLoS Pathog. 13, 10–14. <https://doi.org/10.1371/journal.ppat.1006334>

1020 Pieterse, C.M.J., Leon-Reyes, A., Van Der Ent, S., Van Wees, S.C.M., 2009. Networking by
1021 small-molecule hormones in plant immunity. Nat. Chem. Biol. 5, 308–316.
1022 <https://doi.org/10.1038/nchembio.164>

1023 Pieterse, C.M.J., Van der Does, D., Zamioudis, C., Leon-Reyes, A., Van Wees, S.C.M.,
1024 2012. Hormonal Modulation of Plant Immunity. Annu. Rev. Cell Dev. Biol. 28, 489–521.
1025 <https://doi.org/10.1146/annurev-cellbio-092910-154055>

1026 Powell, J.J., Carere, J., Sablok, G., Fitzgerald, T.L., Stiller, J., Colgrave, M.L., Gardiner, D.M.,
1027 Manners, J.M., Vogel, J.P., Henry, R.J., Kazan, K., 2017. Transcriptome analysis of
1028 Brachypodium during fungal pathogen infection reveals both shared and distinct
1029 defense responses with wheat. Sci. Rep. 7, 1–14. [https://doi.org/10.1038/s41598-017-](https://doi.org/10.1038/s41598-017-17454-3)
1030 17454-3

1031 Proctor, R.H., Hohn, T.M., McCormick, S.P., 1995. Reduced virulence of *Gibberella zeae*
1032 caused by disruption of a trichothecene toxin biosynthetic gene. Mol. Plant. Microbe.
1033 Interact. 8, 593–601.

1034 Puri, K.D., Yan, C., Leng, Y., Zhong, S., 2016. RNA-seq revealed differences in
1035 transcriptomes between 3ADON and 15ADON populations of *Fusarium graminearum* in
1036 vitro and in planta. PLoS One 11, 1–30. <https://doi.org/10.1371/journal.pone.0163803>

1037 Qi, P.-F., Zhang, Y.-Z., Liu, C.-H., Chen, Q., Guo, Z.-R., Wang, Y., Xu, B.-J., Jiang, Y.-F.,
1038 Zheng, T., Gong, X., Luo, C.-H., Wu, W., Kong, L., Deng, M., Ma, J., Lan, X.-J., Jiang,
1039 Q.-T., Wei, Y.-M., Wang, J.-R., Zheng, Y.-L., 2019. Functional Analysis of FgNahG
1040 Clarifies the Contribution of Salicylic Acid to Wheat (*Triticum aestivum*) Resistance
1041 against *Fusarium* Head Blight. Toxins (Basel). 11, 59.
1042 <https://doi.org/10.3390/toxins11020059>

1043 Qi, P.F., Balcerzak, M., Rocheleau, H., Leung, W., Wei, Y.M., Zheng, Y.L., Ouellet, T., 2016.

1044 Jasmonic acid and abscisic acid play important roles in host-pathogen interaction
1045 between *Fusarium graminearum* and wheat during the early stages of fusarium head
1046 blight. *Physiol. Mol. Plant Pathol.* 93, 39–48.
1047 <https://doi.org/10.1016/j.pmpp.2015.12.004>

1048 Quarantin, A., Glasenapp, A., Schäfer, W., Favaron, F., Sella, L., 2016. Involvement of the
1049 *Fusarium graminearum* cerato-platanin proteins in fungal growth and plant infection.
1050 *Plant Physiol. Biochem.* 109, 220–229. <https://doi.org/10.1016/j.plaphy.2016.10.001>

1051 Rabe, F., Ajami-Rashidi, Z., Doehlemann, G., Kahmann, R., Djamei, A., 2013. Degradation
1052 of the plant defence hormone salicylic acid by the biotrophic fungus *Ustilago maydis*.
1053 *Mol. Microbiol.* 89, 179–188. <https://doi.org/10.1111/mmi.12269>

1054 Reineke, G., Heinze, B., Schirawski, J., Buettner, H., Kahmann, R., Basse, C.W., 2008.
1055 Indole-3-acetic acid (IAA) biosynthesis in the smut fungus *Ustilago maydis* and its
1056 relevance for increased IAA levels in infected tissue and host tumour formation. *Mol.*
1057 *Plant Pathol.* 9, 339–355. <https://doi.org/10.1111/j.1364-3703.2008.00470.x>

1058 Rocheleau, H., Al-harthi, R., Ouellet, T., 2019. Degradation of salicylic acid by *Fusarium*
1059 *graminearum*. *Fungal Biol.* 123, 77–86. <https://doi.org/10.1016/j.funbio.2018.11.002>

1060 Scherm, B., Balmas, V., Spanu, F., Pani, G., Delogu, G., Pasquali, M., Migheli, Q., 2013a.
1061 *Fusarium culmorum*: Causal agent of foot and root rot and head blight on wheat. *Mol.*
1062 *Plant Pathol.* 14, 323–341. <https://doi.org/10.1111/mpp.12011>

1063 Scherm, B., Balmas, V., Spanu, F., Pani, G., Delogu, G., Pasquali, M., Migheli, Q.,
1064 Johansson, P.M., Johnsson, L., Gerhardson, B., Hope, R., Aldred, D., Magan, N.,
1065 Cuomo, C.A., Guldener, U., Xu, J., Trail, F., Turgeon, B.G., Pietro, A. Di, Walton, J.D.,
1066 Ma, L., Baker, S.E., Rep, M., Adam, G., Antoniw, J., Baldwin, T., Calvo, S., Chang, Y.,
1067 Harris, L.J., Hilburn, K., Kennell, J.C., Kroken, S., Magnuson, J.K., Muehlbauer, G.,
1068 Münsterkötter, M., Nelson, D., Donnell, K.O., Ouellet, T., 2013b. The *Fusarium*
1069 *graminearum* Genome. *Mol. Plant Pathol.* 14, 295–300.

- 1070 <https://doi.org/10.1126/science.1143708>
- 1071 Schuler, D., Wahl, R., Wippel, K., Vranes, M., Münsterkötter, M., Sauer, N., Kämper, J.,
1072 2015. Hxt1, a monosaccharide transporter and sensor required for virulence of the
1073 maize pathogen *Ustilago maydis*. *New Phytol.* 206, 1086–1100.
1074 <https://doi.org/10.1111/nph.13314>
- 1075 Serino, L., Reimann, C., Baur, H., Beyeler, M., Visca, P., Haas, D., 1995. Structural genes
1076 for salicylate biosynthesis from chorismate in *Pseudomonas aeruginosa*. *Mol. Gen.*
1077 *Genet.* 249, 217–228. <https://doi.org/10.1007/BF00290369>
- 1078 Sesma, A., Osbourn, A.E., 2004. The rice leaf blast pathogen undergoes developmental
1079 processes typical of root-infecting fungi. *Nature* 431, 582–586.
1080 <https://doi.org/10.1038/nature02880>
- 1081 Sieber, C.M.K., Lee, W., Wong, P., Münsterkötter, M., Mewes, H.-W., Schmeitzl, C., Varga,
1082 E., Berthiller, F., Adam, G., Güldener, U., 2014. The *Fusarium graminearum* genome
1083 reveals more secondary metabolite gene clusters and hints of horizontal gene transfer.
1084 *PLoS One* 9, e110311. <https://doi.org/10.1371/journal.pone.0110311>
- 1085 Smirnova, E., Marquis, V., Poirier, L., Aubert, Y., Zumsteg, J., Ménard, R., Miesch, L., Heitz,
1086 T., 2017. Jasmonic Acid Oxidase 2 Hydroxylates Jasmonic Acid and Represses Basal
1087 Defense and Resistance Responses against *Botrytis cinerea* Infection. *Mol. Plant* 10,
1088 1159–1173. <https://doi.org/10.1016/j.molp.2017.07.010>
- 1089 Sørensen, J.L., Benfield, A.H., Wollenberg, R.D., Westphal, K., Wimmer, R., Nielsen, M.R.,
1090 Nielsen, K.F., Carere, J., Covarelli, L., Beccari, G., Powell, J., Yamashino, T., Kogler,
1091 H., Sondergaard, T.E., Gardiner, D.M., 2018. The cereal pathogen *Fusarium*
1092 *pseudograminearum* produces a new class of active cytokinins during infection. *Mol.*
1093 *Plant Pathol.* 19, 1140–1154. <https://doi.org/10.1111/mpp.12593>
- 1094 Spallek, T., Gan, P., Kadota, Y., Shirasu, K., 2018. Same tune, different song — cytokinins

- 1095 as virulence factors in plant–pathogen interactions? Curr. Opin. Plant Biol. 44, 82–87.
- 1096 <https://doi.org/10.1016/j.pbi.2018.03.002>
- 1097 Spence, C.A., Lakshmanan, V., Donofrio, N., Bais, H.P., 2015. Crucial roles of abscisic acid
- 1098 biogenesis in virulence of rice blast fungus *Magnaporthe oryzae*. Front. Plant Sci. 6, 1–
- 1099 13. <https://doi.org/10.3389/fpls.2015.01082>
- 1100 Sperschneider, J., Dodds, P.N., Gardiner, D.M., Singh, K.B., Taylor, J.M., 2018. Improved
- 1101 prediction of fungal effector proteins from secretomes with EffectorP 2.0. Mol. Plant
- 1102 Pathol. 19, 2094–2110. <https://doi.org/10.1111/mpp.12682>
- 1103 Sperschneider, J., Gardiner, D.M., Thatcher, L.F., Lyons, R., Singh, K.B., Manners, J.M.,
- 1104 Taylor, J.M., 2015. Genome-wide analysis in three *Fusarium* pathogens identifies
- 1105 rapidly evolving chromosomes and genes associated with pathogenicity. Genome Biol.
- 1106 Evol. 7, 1613–1627. <https://doi.org/10.1093/gbe/evv092>
- 1107 Stephens, A.E., Gardiner, D.M., White, R.G., Munn, A.L., Manners, J.M., 2008. Phases of
- 1108 infection and gene expression of *Fusarium graminearum* during crown rot disease of
- 1109 wheat. Mol. Plant-Microbe Interact. 21, 1571–1581. [https://doi.org/10.1094/MPMI-21-](https://doi.org/10.1094/MPMI-21-12-1571)
- 1110 12-1571
- 1111 Strange, R., Smith, H., Majer, J., 1972. Choline, One of Two Fungal Growth Stimulants in
- 1112 Anthers responsible for the Susceptibility of Wheat to *Fusarium graminearum*. Nature
- 1113 238, 103–104. <https://doi.org/10.1038/238103a0>
- 1114 Struck, C., 2015. Amino acid uptake in rust fungi. Front. Plant Sci. 6, 1–6.
- 1115 <https://doi.org/10.3389/fpls.2015.00040>
- 1116 Svoboda, T., Parich, A., Güldener, U., Schöffbeck, D., Twaruschek, K., Václavíková, M.,
- 1117 Hellinger, R., Wiesenberger, G., Schuhmacher, R., Adam, G., 2019. Biochemical
- 1118 Characterization of the *Fusarium graminearum* Candidate ACC-Deaminases and
- 1119 Virulence Testing of Knockout Mutant Strains. Front. Plant Sci. 10, 1–17.

1120 <https://doi.org/10.3389/fpls.2019.01072>

1121 Torrens-Spence, M.P., Bobokalonova, A., Carballo, V., Glinkerman, C.M., Pluskal, T., Shen,
1122 A., Weng, J.-K., 2019. PBS3 and EPS1 complete salicylic acid biosynthesis from
1123 isochorismate in Arabidopsis. bioRxiv 601948. <https://doi.org/10.1101/601948>

1124 Trail, F., 2009. For Blighted Waves of Grain: *Fusarium graminearum* in the Postgenomics
1125 Era: Figure 1. Plant Physiol. 149, 103–110. <https://doi.org/10.1104/pp.108.129684>

1126 Tsitsigiannis, D.I., Keller, N.P., 2007. Oxylipins as developmental and host-fungal
1127 communication signals. Trends Microbiol. 15, 109–118.
1128 <https://doi.org/10.1016/j.tim.2007.01.005>

1129 Ursache, R., Andersen, T.G., Marhavý, P., Geldner, N., 2018. A protocol for combining
1130 fluorescent proteins with histological stains for diverse cell wall components. Plant J. 93,
1131 399–412. <https://doi.org/10.1111/tpj.13784>

1132 Voigt, C.A., Schäfer, W., Salomon, S., 2005. A secreted lipase of *Fusarium graminearum* is
1133 a virulence factor required for infection of cereals. Plant J. 42, 364–375.
1134 <https://doi.org/10.1111/j.1365-313X.2005.02377.x>

1135 Wang, H., Sun, S., Ge, W., Zhao, L., Hou, B., Wang, K., Lyu, Z., Chen, L., Xu, S., Guo, J., Li,
1136 M., Su, P., Li, Xuefeng, Wang, G., Bo, C., Fang, X., Zhuang, W., Cheng, X., Wu, J.,
1137 Dong, L., Chen, W., Li, W., Xiao, G., Zhao, J., Hao, Y., Xu, Y., Gao, Y., Liu, W., Liu, Y.,
1138 Yin, H., Li, J., Li, Xiang, Zhao, Y., Wang, X., Ni, F., Ma, X., Li, A., Xu, S.S., Bai, G.,
1139 Nevo, E., Gao, C., Ohm, H., Kong, L., 2020. Horizontal gene transfer of Fhb7 from
1140 fungus underlies *Fusarium* head blight resistance in wheat. Science (80-.). 5435,
1141 eaba5435. <https://doi.org/10.1126/science.aba5435>

1142 Wang, L., Li, Q., Liu, Z., Surendra, A., Pan, Y., Li, Y., Zaharia, L.I., Ouellet, T., Fobert, P.R.,
1143 2018. Integrated transcriptome and hormone profiling highlight the role of multiple
1144 phytohormone pathways in wheat resistance against *fusarium* head blight. PLoS One

1145 13, e0207036. <https://doi.org/10.1371/journal.pone.0207036>

1146 Wang, Q., Shao, B., Shaikh, F.I., Friedt, W., Gottwald, S., 2018. Wheat Resistances to
1147 Fusarium Root Rot and Head Blight Are Both Associated with Deoxynivalenol- and
1148 Jasmonate-Related Gene Expression. *Phytopathology* 108, 602–616.
1149 <https://doi.org/10.1094/PHYTO-05-17-0172-R>

1150 Wang, Q., Vera Buxa, S., Furch, A., Friedt, W., Gottwald, S., 2015. Insights Into *Triticum*
1151 *aestivum* Seedling Root Rot Caused by *Fusarium graminearum*. *Mol. Plant-Microbe*
1152 *Interact.* 28, 1288–1303. <https://doi.org/10.1094/MPMI-07-15-0144-R>

1153 Wiemann, P., Sieber, C.M.K., von Bargaen, K.W., Studt, L., Niehaus, E.M., Espino, J.J., Huß,
1154 K., Michielse, C.B., Albermann, S., Wagner, D., Bergner, S. V., Connolly, L.R., Fischer,
1155 A., Reuter, G., Kleigrew, K., Bald, T., Wingfield, B.D., Ophir, R., Freeman, S., Hippler,
1156 M., Smith, K.M., Brown, D.W., Proctor, R.H., Münsterkötter, M., Freitag, M., Humpf,
1157 H.U., Güldener, U., Tudzynski, B., 2013. Deciphering the Cryptic Genome: Genome-
1158 wide Analyses of the Rice Pathogen *Fusarium fujikuroi* Reveal Complex Regulation of
1159 Secondary Metabolism and Novel Metabolites. *PLoS Pathog.* 9.
1160 <https://doi.org/10.1371/journal.ppat.1003475>

1161 Wildermuth, M.C., Dewdney, J., Wu, G., Ausubel, F.M., 2002. Isochorismate synthase is
1162 required to synthesize salicylic acid for plant defence. *Nature* 417, 571–571.
1163 <https://doi.org/10.1038/417571a>

1164 Yan, X., Talbot, N.J., 2016. Investigating the cell biology of plant infection by the rice blast
1165 fungus *Magnaporthe oryzae*. *Curr. Opin. Microbiol.* 34, 147–153.
1166 <https://doi.org/10.1016/j.mib.2016.10.001>

1167 Yang, F., Jensen, J.D., Svensson, B., Jørgensen, H.J.L., Collinge, D.B., Finnie, C., 2012.
1168 Secretomics identifies *Fusarium graminearum* proteins involved in the interaction with
1169 barley and wheat. *Mol. Plant Pathol.* 13, 445–453. [https://doi.org/10.1111/j.1364-](https://doi.org/10.1111/j.1364-3703.2011.00759.x)
1170 [3703.2011.00759.x](https://doi.org/10.1111/j.1364-3703.2011.00759.x)

- 1171 Yin, Y., Wang, Z., Cheng, D., Chen, X., Chen, Y., Ma, Z., 2018. The ATP-binding protein
1172 FgArb1 is essential for penetration, infectious and normal growth of *Fusarium*
1173 *graminearum*. *New Phytol.* 219, 1447–1466. <https://doi.org/10.1111/nph.15261>
- 1174 Yu, J.H., Wieser, J., Adams, T.H., 1996. The *Aspergillus* FlibA RGS domain protein
1175 antagonizes G protein signaling to block proliferation and allow development. *EMBO J.*
1176 15, 5184–5190. <https://doi.org/10.1002/j.1460-2075.1996.tb00903.x>
- 1177 Yu, Y., Ouyang, Y., Yao, W., 2018. ShinyCircos: An R/Shiny application for interactive
1178 creation of Circos plot. *Bioinformatics* 34, 1229–1231.
1179 <https://doi.org/10.1093/bioinformatics/btx763>
- 1180 Zhang, X.-W., Jia, L.-J., Zhang, Y., Jiang, G., Li, X., Zhang, D., Tang, W.-H., 2012. In *Planta*
1181 Stage-Specific Fungal Gene Profiling Elucidates the Molecular Strategies of *Fusarium*
1182 *graminearum* Growing inside Wheat Coleoptiles. *Plant Cell* 24, 5159–5176.
1183 <https://doi.org/10.1105/tpc.112.105957>
- 1184 Zhang, Y., He, J., Jia, L.J., Yuan, T.L., Zhang, D., Guo, Y., Wang, Y., Tang, W.H., 2016.
1185 Cellular Tracking and Gene Profiling of *Fusarium graminearum* during Maize Stalk Rot
1186 Disease Development Elucidates Its Strategies in Confronting Phosphorus Limitation in
1187 the Host Apoplast. *PLoS Pathog.* 12, 1–33.
1188 <https://doi.org/10.1371/journal.ppat.1005485>
- 1189 Zhou, S., Zhang, Y.K., Kremling, K.A., Ding, Y., Bennett, J.S., Bae, J.S., Kim, D.K.,
1190 Ackerman, H.H., Kolomiets, M. V., Schmelz, E.A., Schroeder, F.C., Buckler, E.S.,
1191 Jander, G., 2019. Ethylene signaling regulates natural variation in the abundance of
1192 antifungal acetylated diferuloylsucroses and *Fusarium graminearum* resistance in
1193 maize seedling roots. *New Phytol.* 221, 2096–2111. <https://doi.org/10.1111/nph.15520>

1194

1195 **Figures**

Figure 1. Global regulation of *Fg* genes during *Bd* root colonization. (A) Expression of 2049 *Fg* genes that were differentially regulated during *Bd* root infections (*Fg-Bd*) relative to *Fg* grown in culture (*Fg-only*). Proteins encoded by differentially expressed genes (DEGs) are assigned to four functional categories as shown in the right side of the figure. (B) Key features of 225 DEGs associated with carbohydrate active enzymes (CAZymes). Enzymatic functions and substrate specificities were predicted as shown in the right side of the figure. PCW: plant cell wall, FCW: fungal cell wall. (C) Key features of 251 DEGs encoding putative secreted proteins. *Bd* roots colonized by *Fg* (*Fg-Bd*) were harvested at 5 dpi and *Fg* mycelia grown without *Bd* roots were used as control (*Fg-only*). Transcript levels of fungal genes were presented as normalized FPKM (Fragments Per Kilobase of transcript per Million mapped reads) values and log2-transformed. Heatmap colour range represents high to low expression levels. The dendrogram shows distance similarity of expression of each gene.

Figure 2. The number of *Fg* genes upregulated during *Bd* root infection were also induced or actively expressed during the infection of other tissues by *Fg*, suggesting that a core number of *Fg* genes could be broadly associated with infection (A) The Venn diagram showing how many *Fg* genes that were observed to be up-regulated during *Bd* root infection in this study were common to those upregulated or differentially expressed in different tissues of wheat (*Triticum aestivum*, *Ta*), barley (*Hordeum vulgare*, *Hv*) and maize (*Zea mays*, *Zm*). (B) The heatmap showing differential expression profiles of 38 core *Fg* genes identified from (A) between *in vitro* growth and during *Bd* root infection. FPKM values from three independent biological replicates were log2-transformed. Putative proteins encoded by these genes are shown on the right.

Figure 3. Genome-wide comparison of *Fg* and *F. pseudograminearum* (*Fp*) gene expression patterns during infection of *Bd* roots (this study) and aboveground tissues (Powell et al., 2017). (A) Chromosomal locations and syntenic relationship of highly conserved *Fg* and *Fp* genes. (1-4: *Fg* chromosomes; NC_031952.1-NC_031952.4: *Fp* chromosomes). (B) The top 20 genes with the most contrasted expression values were selected from chromosome 2 of

Fg and *Fp* genomes, respectively. Heat maps in A and B display expression of genes as log2-transformed FPKM values from three biological replicates. Colour range is indicated as high (red) to low (blue).

Figure 4. Global regulation of *Fg* genes in the *Tri5* mutant during *Bd* root colonization. (A) Expression of 974 *Fg* genes that were differentially expressed WT *Fg* and Δ Tri5 during *Bd* root infection. Proteins encoded by DEGs were assigned to four functional categories as shown on the right. (B) Key features of 95 DEGs associated with carbohydrate active enzymes (CAZymes). Plant cell wall (PCW) degradation functions and substrate specificities were predicted as shown in the right-hand side. (C) Key features of 124 DEGs encoding putative secreted proteins. *Bd* roots inoculated with *Fg* WT and Δ Tri5 strains were harvested at 5 dpi for the RNA-seq analysis. Transcript levels of *Fg* genes were log2-transformed and presented as normalized FPKM values. Heatmap colour range representing high to low expression levels is shown. Dendrograms show distance similarity of expression of each gene.

Figure 5. Lignin deposition in response to *Fg* infection is reduced in *Bd* roots inoculated with Δ Tri5. *Bd* roots either mock-treated or inoculated with either WT *Fg* or Δ Tri5 were harvested at 5 dpi. Red fluorescence signals (lower panels) indicate lignin deposition stained by Basic Fuchsin. Shown are representatives of at least 6 independent roots from three biological replicates. Images were taken using 561nm excitation and detected at 600–650 nm on a Zeiss Axio Imager M2 microscopy.

Figure 6. Quantification of selected phytohormones and DON in *Bd* roots inoculated with either WT *Fg* or the Δ Tri5 mutant; jasmonic acid (JA) (A), methyl-jasmonate (MeJA) (B), salicylic acid (SA) (C), deoxynivalenol (DON) (D), indole-3-acetic acid (IAA) (E) and gibberellic acid (GA) (F) analysed by high-performance liquid chromatography. Independently grown fungal mycelium on minimal media (MM) (*Fg*-only), uninoculated *Bd* roots (*Bd*-only), as well as *Bd* roots inoculated with WT *Fg* or Δ Tri5 infected roots were collected from MM at 5 dpi and subjected to metabolite extraction. Pure MM was used as

background control. Metabolite quantifications were conducted according to the concentration-gradient derived standard curves of JA, MeJA, SA, DON, IAA and GA. Error bars indicate standard error of the mean based on six biological replicates, each comprising of 10 plants or 1 plate of fungal mycelium. Asterisks represent differences that were statistically significant (unpaired two-tailed t-test, * $P < 0.05$. **** $P < 0.0001$).

Figure 7. Deoxynivalenol (DON) produced by *F. graminearum* (*Fg*) during the infection of *B. distachyon* (*Bd*) roots does not contribute to lesion formation. Microscopic pictures were taken for *Bd* roots colonized by *Fg* strains after two days post inoculation (dpi). **(A)** DON is produced during the infection of *Bd* roots by *Fg*. Strong florescence signals could be observed in the mycelium of a *Fg* strain expressing a *Tri5-GFP* fusion construct driven by the native *Tri5* promoter. **(B)** No GFP signals could be detectable for WT *Fg* during root infection. **(C)** Transcriptional activation of *Tri5* confirms DON production in infected roots. Expressions of *Tri5*, *Flp1*, a lipase encoding gene previously identified as a virulence factor in *Fg* and a *Bd* ubiquitin-conjugating enzyme 18 gene (*UBC18*) used for fungal biomass measurements were quantified by RT-qPCR relative to the fungal α -tubulin. Root samples were harvested at 2, 3, 4, 5, and 7 dpi from three independent biological replicates each with at least 12 individual plants. **(D)** Deoxynivalenol (DON) does not contribute to lesion formation in *Bd* roots. Representative photos showing colonization of *Bd* roots with the WT *Fg* and *Tri5* deletion mutants.

Figure 8. SA hydroxylase-like genes present in the *Fg* genome and their expression profiles during growth in culture or infection of *Bd* roots. **(A)** Phylogenetic unrooted tree of SA hydroxylase protein homologs of NahG, Arabidopsis Dlo1 (AT4G10500) and Dmr6 (AT5G24530) in *Fg*. The protein evolutionary models were tested using Neighbor-Joining inference and then referred to tree building based on Maximum Likelihood method. **(B)** Expressions of 28 SA hydroxylase candidate genes during *Bd* root infections or *in vitro* growth on minimal medium (MM). Heatmap shows expression levels of each gene displayed as log-transformed FPKM values of three replicates from low (blue) to high (red). *Fg* genes

1277 significantly induced during *Bd* root infection were marked as 'yes'. An overview of
1278 concatenated alignment of all protein sequences was shown aside.

1279

1280

1281

1282

1283

1284

1285

1286

1287

1288

1289

1290

A

Expression $\log_2(\text{FPKM}+0.01)$

B

CAZymes

C

Secreted proteins

A

B

row min row max
Expression $\log_2(\text{FPKM}+0.01)$

A

Expression $\log_2(\text{FPKM}+0.01)$

B

CAzyme

C

Secreted

mock

Fg WT

Fg Δ Tri5

bioRxiv preprint doi: <https://doi.org/10.1101/2020.05.17.062288>; this version posted May 17, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

B