Molecular mapping of *qBKLZ*, a major QTL for bakanae disease resistance in rice

Sais-Beul Lee¹, Namgyu Kim², Sumin Jo¹, Yeon-Jae Hur¹, Ji-youn Lee¹, Jun-Hyeon Cho¹, Jong-Hee Lee¹, Ju-Won Kang¹, You-Chun Song¹, Maurene Bombay⁴, Sung-Ryul Kim⁴, Jungkwan Lee³, Young-Su Seo², Jong-Min Ko¹, Dong-Soo Park^{1,4*}

¹National Institute of Crop Science, Milyang 50424, Republic of Korea

²Department of Microbiology, Pusan National University, Pusan 46241, Republic of Korea

³College of Natural Resources and Life Science, Dong-A University, Pusan 49135, Republic of Korea

⁴International Rice Research Institute, Pili Drive, Los Baños, Laguna 4031, Philippines

***Correspondence:** Dong-Soo Park

Email: parkds9709@korea.kr

Tel: 82-55-350-1184

Fax: 82-55-352-3059

Funding

This work was conducted with the support of the "Cooperative Research Program for Agriculture Science & Technology Development (Project title: QTL mapping for development of functional rice with bakanae disease resistance, Project No. PJ01477401", Rural Development Administration, Republic of Korea.

Abstract

Bakanae disease is a fungal disease of rice (*Oryza sativa* L.) caused by the pathogen *Gibberella fujikuroi* (also known as *Fusarium fujikuroi*). Recently the disease incidence has increased in several Asian countries and continues to spread throughout the world. No rice varieties have been developed yet to be completely resistant to this disease. With increasing need to identify various genetic resources to impart resistance to local elite varieties, this study was carried out to identify novel quantitative trait loci (QTLs) from an *indica* variety Zenith. We performed a QTL mapping using 180 F_{2:9} recombinant inbred lines (RILs) derived from a cross between the resistant variety, Zenith, and the susceptible variety, Ilpum. A primary QTL study using the genotypes and phenotypes of the RILs indicated that the locus *qBKI^z* conferring bakanae disease resistance from the Zenith was located in a 2.8 Mb region bordered by the two SSR markers, RM1331 and RM3530 on chromosome 1. The log of odds (LOD) score of *qBKI^z* was 13.43, accounting for 30.9% of the total phenotypic variation. A finer localization of *qBKI^z* was delimited at an approximate 730 kb interval in the physical map between Chr01_1435908 (1.43 Mbp) and RM10116 (2.16 Mbp). The development of a rice variety with a higher level of resistance against bakanae disease is a major challenge in many rice growing countries. Introducing *qBKI^z* or pyramiding with other previously identified QTLs could provide effective genetic control of bakanae disease in rice.

Introduction

Bakanae disease, which means foolish seedling in Japanese, was firstly identified in 1828 in Japan [1], and has widely distributed in temperate zone as well as tropical environment and occurring throughout rice growing regions of the world [2]. Four *Fusarium* species including *F. andiyazi*, *F. fujikuroi*, *F. proliferatum* and *F. verticillioides* in the *G. fujikuroi* species complex have been associated with Bakanae disease in rice [3]. This disease typically seed-borne fungus, but may occur when the pathogen is present in plant material or soil. Infected seeds/plants result in secondary infections [4], which spreads through wind or water that carries the fungal spores from one plant to another. Bakanae disease has different symptoms starting from pre-emergence seedling death to grain infection at maturity such as tall, lanky tillers with pale green flag leaves. Infected plants have fewer tillers, and plants surviving till maturity bear only empty panicles [5], resulting in yield loss [6, 7]. Low plant survival and high spikelet sterility [5] may account for yield losses up to 50% in Japan [6, 7], 3.0-95% in India [2, 8, 9], 3.7-14.7% in Thailand, 5-23% in Spain, 40% in Nepal [10], 6.7-58.0% in Pakistan [11], 75% in Iran [12], and to 28.8% in Korea [13]. Hot water immersion and fungicide treatment are most common ways for seed

disinfection [2, 14, 15]. However, both the hot water treatment and application of fungicide are insufficient to control bakanae disease. Thermal effect does not reach the pericarp of the severely infected rice seeds. The application of fungicides is not functioning well for destroying the spores of this fungal pathogen, and some pathogen showed resistance to the fungicides [13, 16-18]. Therefore, the genetic improvement of rice using the QTLs/genes providing the bakane disease resistance would be a more effective way to control bakanae disease.

Several QTLs associated with bakanae disease resistance have been identified and those can be used for marker-assisted selection in rice breeding as well as for understanding the mechanisms of resistance. Yang et al. [19] identified two QTLs located on chromosome 1 and chromosome 10 by *in vitro* evaluation of the Chunjiang 06/TN1 doubled haploid population. Hur et al. [20] identified a major QTL, *qBK1*, on chromosome 1 from 168 BC₆F₄ near isogenic lines (NILS) generated by crossing the resistant *indica* variety Shingwang with susceptible *japonica* variety Ilpum. Lee et al. [21] delimited the location of *qBK1* to 35 kb interval between two InDel markers, InDel 18 (23.637 Mbp) and InDel 19-14 (23.672 Mbp). Fiyaz et al. [9] identified three QTLs (*qBK1.1*, *qBK1.2*, and *qBK1.3*) on chromosome 1 and one QTL (*qBK3.1*) on chromosome 3 from the highly resistant variety Pusa 1342. Physical location of *qBK1.1* is close to *qBK1* [21]. Ji et al. [22] mapped a major QTL (*qFfR1*) in 22.56–24.10 Mbp region on chromosome 1 using 180 F_{2,3} lines derived from a cross between a resistant Korean *japonica* variety Nampyeong and a susceptible Korean *japonica* line, DongjinAD. Volante et al. [23] identified *qBK1_628091* (0.6 Mbp to 1.0 Mbp on chromosome 1) and *qBK4_31750955* (31.1 Mbp to 31.7 Mbp on chromosome 4) by GWAS (Genome Wide Association Study) approach using 138 *japonica* rice germplasm. Kang et al [24] discovered the QTL *qFfR9* at 30.1 centimorgan (cM) on chromosome 9 from a *japonica* variety Samgwang. Lee et al. [15] found the QTL *qBK1^{WD}* located between markers chr01_13542347 (13.54 Mb) and chr01_15132528 (15.13 Mb) from the *japonica* variety Wonseadaesoo. They also found that resistance of gene pyramided lines harboring two QTLs, *qBK1^{WD}* and *qBK1* was significantly higher than those with only *qBK1^{WD}* or *qBK1*. This finding relates to the problem of varieties with single resistance gene losing its effect against new populations of fungal isolates [25]. Hence, identifying new resistance genes from diverse sources is important for rice breeding programs to acquire durable resistance against bakanae disease by enhancing the resistance level and/or help to overcome the breakdown of resistance genes.

In this study, we aimed to provide a new genetic source, *qBK1^z* with detailed gene locus information for developing resistant rice lines which contains single or multiple major QTLs to enhance bakanae disease resistance.

Materials and Methods

Plant materials

Zenith, a medium grain type *indica* variety of USA released in 1936, was identified as resistant to bakanae disease in a preliminary screening of rice germplasm (data not shown) using the large-scale screening method developed by Kim et al. [26]. We generated 180 F_{2:9} RILs from a cross between susceptible variety, Ilpum, and resistant variety, Zenith for QTL analysis. The population was developed in the experimental fields of the National Institute of Crop Science of the Rural Developmental Administration in Miryang, Korea.

Evaluation of bakanae resistance

The inoculation and evaluation of bakanae disease were conducted using a method that described by Lee et al. [21]. The isolate CF283 of *F. fujikuroi* (imperfect stage of *G. fujikuroi*) were obtained from National Academy Agricultural Science in Korea. Isolate was inoculated in potato dextrose broth (PDB) and cultured at 26 °C under continuous light for one week. The *F. fujikuroi* culture was washed by centrifugation with distilled water. Regarding the standardization of the density of the homogenized inoculum, the fungal spore concentration was adjusted to 1×10^6 spores/mL with a hemocytometer. Forty seeds per each line were placed in a tissue-embedding cassette (M512, Simport, Beloeil, QC, Canada). Before inoculation, the seeds in the in the tissue-embedding cassette were surface sterilized with hot water (57 °C) for 13 min, then allowed to drain and cool. Subsequently, the seeds were soaked in the spore suspensions (1×10^6 spores/mL) for 3 days for inoculation with gentle shaking four times a day for equilibration. After inoculation, thirty seeds per line were sown in commercial seedling tray, and seedlings were grown in a greenhouse (28 ± 3 °C day/ 23 ± 3 °C night, 12 h light). Bakanae disease symptom on each line was evaluated by calculating the proportion of healthy plants at 1 month after sowing. The healthy and non-healthy plants are classified as described by Kim et al. [26]. The plants exhibit elongation with thin and yellowish-green, elongation then stunted growth, stunted growth, dead seedling were classified as non-healthy plants. The plants showing same phenotype with untreated plants, slight elongation then normal growth without thin and yellowish-green was regarded as healthy plants.

Localization of *F. fujikuroi* in Zenith and Ilpum plants

Shoot base of 10 day-old seedlings derived from Zenith and Ilpum seeds inoculated by GFP-tagged *F. fujikuroi* isolate CF283 [15] was observed under confocal laser-scanning microscopy (LSM-800,

Zeiss, Germany) at GFP channel, and images were obtained using Zeiss LSM Image Browser. All experiments were conducted twice with at least three replicates.

DNA extraction and polymerase chain reaction

Genomic DNA from young leaf tissue was prepared according to the CTAB method [27] with minor modifications. Polymerase chain reaction (PCR) was performed in 25-μL reaction mixture containing 25 ng template DNA, 10 pmol of each primer, 10× e-Taq reaction buffer, 25 mM MgCl₂, 10 mM dNTP mix and 0.02 U of SolGent™ e-Taq DNA polymerase (SolGent, Daejeon, South Korea). The reaction conditions were set as follows: initial denaturation at 94°C for 2 min; 35 cycles of denaturation at 94°C for 20 s, annealing at 57°C for 40 s and extension at 72°C for 40 s; and a final extension at 72°C for 7 min. The amplification products were electrophoresed on a 3% (w/v) agarose gel and visualized by ethidium bromide staining.

QTL analysis of the F_{2,9} population and development of InDel markers for fine mapping

Polymorphic SSR markers (n = 164) that were evenly distributed on rice chromosomes were selected from the Gramene database (<http://www.gramene.org>). These markers were used to construct a linkage map and for QTL analysis of the F_{2,9} populations. The linkage map was constructed using Mapmaker/Exp v.3.0, and the genetic distance was obtained using the Kosambi map function [28]. Putative QTLs were detected using the composite interval mapping (CIM) function in WinQTLcart v.2.5 (WinQTL cartographer software [29]), with the threshold being set based on the results of 1000 permutation tests. A logarithm of the odds (LOD) ratio threshold of 3.0 was used to confirm the significance of a putative QTL. InDel markers were developed based on the fragment size differences in the sequence (in the range of 20 bp) between *japonica* (Gramene database <http://www.gramene.org>) and *indica* (BGI-RIS; <http://rice.genomics.org.cn>) in the target region on chromosome 1. The primers were designed using Primer3 software (<http://web.-bioneer.co.kr/cgi-bin/primer/primer3.cgi>).

Statistical analysis

Statistical differences between means were analyzed using Duncan's multiple range test after one-way analysis of variance (ANOVA). The level of significance was designated as P < 0.05 and was determined using the SAS Enterprise Guide 4.3 program (SAS Institute Inc., Cary, NC, USA).

Results

Bakanae disease bioassay in parents and F_{2,9} RILs

The proportion of healthy Zenith (resistant) and Ilpum (susceptible) plants was investigated after inoculation of the virulent *F. fujikuroi* isolate CF283 [26]. Most of Zenith plants did not exhibit a thin and yellowish-green phenotype, which is a typical symptom of bakanae disease as unlike Ilpum (Fig. 1A). The proportion of healthy plants was significantly different between Ilpum (14.3%) and Zenith (63.2%) (Fig. 1B).

Zenith and Ilpum were inoculated with green fluorescent protein (GFP)-tagged *F. fujikuroi* isolate CF283. Ten days after inoculation, plants with typical disease symptom of each variety were subjected to a confocal microscopy analysis. Confocal imaging of radial and longitudinal sections of the basal stem showed that the fungus penetrated and was localized easily and abundantly at vascular bundle, mesophyll tissue and hypodermis in the susceptible Ilpum variety while only background level of GFP signal was detected in the resistant Zenith (Fig. 2).

QTL analysis and mapping of *qBKI^z* using 180 F_{2,9} RILs

Based on the bakanae disease bioassay (proportion of healthy plants), the 180 F_{2,9} RIL population exhibited continuous distribution (0.0–98.0%; Fig. 3), which quantitatively confirmed the inheritance of bakanae disease resistance. The average proportion of healthy Ilpum plants was 14.3% (3.3–37.0%; $n = 10$) and that of Zenith was 63.2% (42.7–79.3%; $n = 10$).

We selected 164 SSR markers among 1,150 SSR markers from the polymorphism survey between Ilpum and Zenith from the Gramene database (<http://www.gramene.org>) covering the whole rice chromosome (S1 Fig). The genetic linkage map of Ilpum and Zenith for primary mapping was constructed with 164 polymorphic markers covering a total length of 3,140 cM with average interval of 19.14 cM as described by Lee et al. [30]. Primary QTL mapping using the 180 F_{2,9} populations showed that a significant QTL associated with bakanae disease resistance at the seedling stage was located between the SSR markers, RM1331 and RM3530 on chromosome 1, and it was designated *qBKI^z*. The LOD score of *qBKI^z* was 13.43, which accounted for 30.9% of the total phenotypic variation (Table 1).

Table 1. Putative quantitative trait locus (QTL) associated with bakanae disease resistance detected at the seedling stage by composite interval mapping of the 180 F_{2:9} populations derived from a cross between Ilpum and Zenith

QTL	Chromosome	Position (cM)	Marker interval	LOD ^a	PVE ^b (%)	Additive effect	Dominant effect
<i>qBKI^z</i>	1	4.0	RM1331 ~ RM3530	13.43	30.93	-22.10	-1.09

^a LOD: log of odds score; ^b PVE: percentage of phenotypic variation explained.

A finer localization of *qBKI^z* was determined by analyzing the chromosome segment introgression lines in the region detected from primary mapping. The *qBKI^z* region between RM1331 and RM3530 from primary mapping was narrow down with additional 55 SSR markers and designed 12 InDel markers for the insertion/deletion sites based on the differences between the *japonica* (<http://www.gramene.org>) and *indica* (<http://rice.genomics.org.cn>) sequences. Four SSR markers and six InDel markers were selected as polymorphic markers between the parents to narrow down the position of the *qBKI^z* region (S2 Table). Finally, seven homozygous recombinants were selected from the F_{2:9} lines using 14 markers in the 2.8 Mb region around the SSR markers RM1331 and RM3530 (Fig. 4 and Fig. 5).

The proportion of healthy plants of the seven homozygous recombinants was evaluated with three biological replicates according to Duncan's new multiple range test. Based on this bioassay, lines classified to Group a were regarded as resistant, and Group b as susceptible (Fig. 4). Considering the genotype and the phenotype of the recombinants, it is clear that *qSTVII^z* conferring resistance to bakanae was an approximate 730 kb interval delimited by the physical map between Chr01_1435908 (1.43 Mbp) and RM10116 (2.16 Mbp).

Discussion

Rice varieties with a single resistance gene are at an increased risk of being overcome by new pathological races [15, 30]. The development of a rice variety with a higher level of resistance against bakanae disease is a major challenge in many countries [9, 20, 31-33]. In this study, we identified *qBKI^z* locus related to bakanae disease resistance based on genotype and phenotype analyses of

homozygous recombinants on the recombinant progeny of Ilpum and Zenith, using SSR and newly developed InDel markers.

Localization of *F. fujikuroi* isolate CF283 in the susceptible Ilpum and the resistant Zenith was also examined. It was reported that successful infection of *Fusarium* species is a complex process that includes adhesion, penetration (through wounds, seeds, stomatal pores) and subsequent colonization within and between cells [34, 35]. Consistent with previous reports [36, 37], *F. fujikuroi* isolate CF283 was extensively observed on the aerenchyma, pith, cortex and vascular bundle of both leaf and stem in susceptible Ilpum, whereas this was rarely observed in resistant Zenith.

Many QTLs on bakanae disease resistance have been identified on chromosome 1. Three QTLs, *qBK1^z*, *qBK1.2* and *qBK1.3*, were found in a similar region in spite of the different source of resistant varieties (Fig 6).

Fiyaz et al. [9] mapped *qBK1.2* to a 260 kb region between markers RM10153 and RM5336, and *qBK1.3* to a 3.49 Mb region between markers RM10271 and RM578 from the Pusa 1121/Pusa1342 cross. Location of *qBK1^z* was found upstream to *qBK1.2*. Fiyaz et al. (2016) mapped *qBK1.1* to a 20 kb region between markers RM9 and RM11232 from the Pusa 1121/Pusa1342 cross. These authors hypothesized that *qBK1.1* and *qBK1* [20] might be the same QTL as they had overlapping positions. Ji et al. [22] found that QTL *qFfR1* was located in a 230 kb region of rice chromosome 1 in Korean *japonica* variety Nampyeong, and suggested that the three QTLs *qBK1*, *qBK1.1*, and *qFfR1* might indicate the same gene. Lee et al. [21] narrowed down the position of the *qBK1* locus to a 35 kb region between InDel 18 and InDel 19-14, and revealed that location of *qBK1* is close to those of *qBK1.1*, and *qFfR1*, does not overlapping each other. Two additional QTLs including *qBI* from Chunjiang 06 [19] and *qBK1^{WD}* from Wonseadaesoo [15] also were also found on chromosome 1. Gene pyramiding via phenotypic screening assays for crop breeding is considered to be difficult and often impossible due to dominance and epistatic effects of genes governing disease resistance, and the limitation of screenings being all year-round [38]. Pyramiding of multiple resistant QTLs/genes by using marker-assisted breeding (MAB) in a single plant might confer higher and/or durable resistance against bakanae disease. The effects of pyramiding resistance genes have been observed for several plant-microbe interactions. Pyramiding three bacterial blight resistance genes resulted in a high level of resistance and were expected to provide a durable pathogen resistance [39, 40]. On the other hand, pyramiding of resistant genes resulted in a level of resistance that was comparable to or even lower to than that of the line with a single gene. For example, Yasuda et al. [41] reported rice lines with pairs of blast resistance genes to be only comparable to lines with a single gene which may have a stronger suppressive effect. Similarly, pyramiding of brown planthopper (BPH) resistance genes in a

susceptible variety IR24 markedly improved BPH resistance against four BPH populations in the Philippines than the single BPH introgressed lines [42].

In our previous study of bakanae disease resistance [15] revealed that the lines harboring *qBKI^{WD}* showed a higher resistance than those with *qBKI*. Furthermore, the pyramided lines harboring *qBKI^{WD} + qBKI* had a much higher level of resistance than those possessing either *qBKI^{WD}* or *qBKI*. These results showed the utility of MAB in gene pyramiding can achieve higher resistance in many bakanae disease prone rice growing areas.

Conclusions

In this study, we identified a new major QTL *qBKI^z* conferring bakanae disease resistance from a new genetic source of *indica* variety, Zenith. Through QTL analysis and fine mapping, we narrowed down the *qBKI^z* locus into 730 kb on the short arm of chromosome 1 where is a novel locus compared with all the previously identified bakane disease resistance QTLs. Together with the previously identified QTLs of the bakane disease resistance, the new *qBKI^z* can be introduce to the elite favorable background varieties by aa marker-assisted backcrossed breeding. Furthermore, the new QTL *qBKI^z* will be a useful material for studying an interaction between the pathogen (*F. fujikuroi*) and rice host plants.

Figure legends

Fig 1. Bioassay of parents (Ilpum and Zenith) against *F. fujikuroi* isolate CF283. (A) Plant phenotype from the pathogen-inoculated seeds of Ilpum and Zenith. (B) Portion of healthy plants from the pathogen-treated seeds.

Fig 2. Confocal imaging of shoot base of 10 days-old-seedling derived from Zenith and Ilpum rice plants infected with *F. fujikuroi* (CF283 isolates) containing GFP expression cassette. (A) Radial and (B) longitudinal sections of the basal stem (Scale bar = 50 μ m).

Fig 3. Frequency distribution of the proportion of healthy $F_{2,9}$ NILs derived from a cross between Ilpum and Zenith after bakanae disease inoculation. The mean proportion of healthy plants of Ilpum and Zenith is indicated by black arrow heads.

273

274 **Fig 4.** QTL analysis of *qBKI^Z* on chromosome 1 using 180 RILs derived from a cross between Ilpum
275 and Zenith. (a) In primary mapping, *qBKI^Z* was located in 2.8 Mb regions between the RM1331 and
276 RM3530 markers on chromosome 1. (b) Location of *qBKI^Z* was narrowed down to 730 kb region
277 between the markers Chr01_1435908 and RM10116 in secondary mapping using selected seven
278 homozygous recombinants. Black bars show the homozygous regions for zenith alleles; white bars
279 indicate the homozygous regions for Ilpum alleles. R; resistant to bakanae disease, S; susceptible to
280 bakanae disease. The proportion of healthy plant was calculated from three biological replications.
281 Values (%) of the on proportion of healthy plant with different letters are significantly different by
282 Duncan's multiple range test at P = 0.05.

283

284 **Fig 5.** Phenotypic responses to bakanae disease infection of seven homozygous recombinants used in
285 secondary mapping.

286

287 **Fig 6.** Physical locations of bakanae disease resistance QTLs identified on chromosome 1.

288

289 **Supporting information**

290 S1 Fig. Linkage map constructed with 180 F_{2,9} recombinant inbred lines (RILs) derived from a cross
291 between Zenith and Ilpum. (PPT)

292

293 S2 Table. InDel and SSR markers used for the fine mapping of *qBK^Z*. (DOCX)

294

295 **Author contributions**

296 Conceptualization: SB Lee, YJ Hur, DS Park

297 Data curation: SB Lee, N Kim, YS Seo, J Lee, DS Park

Formal analysis: SB Lee, DS Park
 Funding acquisition: DS Park
 Investigation: SB Lee, N Kim
 Methodology: SB Lee, DS Park
 Project administration: JM Ko, DS Park
 Software: SM Jo, JH Cho
 Supervision: DS Park
 Validation: YS Seo, J Lee, DS Park
 Visualization: JHee Lee, J Lee, JW Kang, YC Song.
 Writing – original draft: SB Lee, DS Park
 Writing – review & editing: YS Seo, J Lee, M Bombay, SR Kim, DS Park

References

1. Ito S, Kimura J (1931) Studies on the ‘bakanae’ disease of the rice plant. Rep Hokkaido Natl Agric Exp Stat 27:1-95
2. Gupta AK, Solanki IS, Bashyal BM, Singh Y, Srivastava K (2015) Bakanae of rice - an emerging disease in Asia. J Anim Plant Sci 25:1499-1514
3. Wulff EG, Sørensen JL, Lübeck M, Nielsen KF, Thrane U et al. (2010) *Fusarium* spp. associated with rice Bakanae: ecology, genetic diversity, pathogenicity and toxigenicity. Environ Microbiol 12:649-657
4. Ora N, Faruq AN, Islam MT, Akhtar N, Rahman MM (2011) Detection and identification of seed borne pathogens from some cultivated hybrid rice varieties in Bangladesh. Middle-East J Sci Res 10:482-488
5. Singh R, Sunder S (2012) Foot rot and bakanae disease of rice: an overview. Rev Plant Pathol Vol. 5, Indian Society of Mycology and Plant Pathology, Scientific Publishers (India) 565-604
6. Mew TW, Gonzales PG (2002) A handbook of rice seedborne fungi. International Rice Research Institute, Los Baños (Philippines) and Science Publishers, Inc., Enfield
7. Ou SH (1985) Rice diseases, 2nd edn. Commonwealth Mycological Institute, Kew.
8. Sunder S, Satyavir Virk KS (1997) Studies on correlation between bakanae incidence and yield

- loss in paddy. *Indian Phytopathol* 50:99-101
9. Fiyaz RA, Yadav AK, Krishnan SG, Ellur RK, Bashyal BM et al. (2016) Mapping quantitative trait loci responsible for resistance to Bakanae disease in rice. *Rice* 9:3-10
10. Singh R, Sunder S (1997) Foot rot and bakanae of rice: Retrospects and prospects. *Intern. J Trop Pl Dis* 15:153-176
11. Yasin SI, Khan TZ, Akhtar KM, Muhammad A, Mushtaq A (2003) Economic evaluation of bakanae disease of rice. *Mycopath* 1(2):115-17
12. Saremi H, Ammarellou A, Marefat A, Okhovat SM (2008) Binam a rice cultivar, resistant for root rot disease on rice caused by *Fusarium moniliforme* in North-West, Iran. *Intern J Bot* 4:383-389
13. Park WS, Choi HW, Han SS, Shin DB, Shim HK et al. (2009) Control of bakanae disease of rice by seed soaking into the mixed solution of prochloraz and fludioxnil. *Res Plant Dis* 15:94-100
14. Hayasaka T, Ishiguro K, Shibutani K, Namai T (2001) Seed disinfection using hot water immersion to control several seed-borne diseases of rice plants. *Jpn J Phytopathol* 67:26-32
15. Lee SB, Hur YJ, Cho JH, Lee JH, Kim TH et al. (2018) Molecular mapping of *qBKL^{WD}*, a major QTL for bakanae disease resistance in rice. *Rice* 11:3. <https://doi.org/10.1186/s12284-017-0197-7>
16. Ogawa K (1988) Damage by “Bakanae” disease and its chemical control. *Jpn Pestic Inf* 52:13-15
17. Kim JM, Hong SK, Kim WG, Lee YK, Yu SH et al. (2010) Fungicide resistance of *Gibberella fujikuroi* isolates causing rice bakanae disease and their progeny isolates. *Kor J Mycol* 38:75-79
18. Lee YH, Lee MJ, Choi HW, Kim ST, Park JW et al. (2011) Development of *in vitro* seedling screening method for selection of resistant rice against bakanae disease. *Res Plant Dis* 17:288-294
19. Yang CD, Guo LB, Li XM, Ji ZJ, Ma LY et al. (2006) Analysis of QTLs for resistance to rice bakanae disease. *Chin J Rice Sci* 6:657-659
20. Hur YJ, Lee SB, Kim TH, Kwon T, Lee JH et al. (2015) Mapping of *qBKL*, a major QTL for bakanae disease resistance in rice. *Mol Breeding* 35: 78. <https://doi.org/10.1007/s11032-015-0281-x>
21. Lee SB, Kim N, Hur YJ, Cho SM, Kim TH et al. (2019) Fine mapping of *qBKL*, a major QTL for bakanae disease resistance in rice. *Rice* 12:36. <https://doi.org/10.1186/s12284-019-0295-9>
22. Ji H, TH K, Lee GS, Kang HJ, Lee SB et al. (2017) Mapping of a major quantitative trait locus for bakanae disease resistance in rice by genome resequencing. *Mol Gen Genomics*. <https://doi.org/10.1007/s00438-017-1407-0>
23. Volante A, Tondelli A, Aragona M, Valente MT, Biselli C et al. (2017) Identification of bakanae disease resistance loci in *japonica* rice through genome wide association study. *Rice* 10:29 DOI <https://doi.org/10.1186/s12284-017-0168-z>

24. Kang DY, Cheon KS, Oh J, Oh H, Kim SL et al. (2019) Rice genome resequencing reveals a major quantitative trait locus for resistance to bakanae disease caused by *Fusarium fujikuroi*. Int. J. Mol. Sci. <https://doi.org/10.3390/ijms20102598>
25. Wang GL, Valent B (2009) Advances in Genetics, Genomics and Control of Rice Blast Disease. Springer Science+Business Media B.V., Dordrecht
26. Kim MH, Hur YJ, Lee SB, Kwon TM, Hwang UH et al. (2014) Large-scale screening of rice accessions to evaluate resistance to bakanae disease. J Gen Plant Pathol 80:408-414
27. Murray MG, Thompson WF (1980) Rapid isolation of high molecular weight plant DNA. Nucleic Acids Res. 10:8(19):4321-5
28. Lander ES, Green P, Abrahamson J, Barlow A, Daly MJ et al. (1987) MAPMAKER: An interactive computer package for constructing primary genetic linkage maps of experimental and natural populations. Genomics 1: 174-181
29. Wang S, Basten CJ, Zeng ZB (2007) Windows QTL cartographer 2.5. <http://statgen.ncsu.edu/qtlcart/WQTLCart.htm>
30. Lee SB, Hur YJ, Lee JH, Kwon TM, Shin DJ et al. (2017) Molecular mapping of a quantitative trait locus *qSTV11^Z* harboring rice stripe virus resistance gene, Stv-b. Plant Breeding, 136, 61-66
31. Cumagun CJR, Arcillas E, Gergon E (2011) UP-PCR analysis of the seedborne pathogen *Fusarium fujikuroi* causing bakanae disease in rice. Int J Agric Biol 13:1029-1032
32. Bashyal BM, Aggarwal R, Banerjee S, Gupta S, Sharma S (2014) Pathogenicity, ecology and genetic diversity of the *Fusarium* spp. associated with an emerging bakanae disease of rice (*Oryza sativa* L.) in India. In: Kharwar RN, Upadhyay R, Dubey N, Raghuwanshi R (eds) Microbial diversity and biotechnology in food security, Springer India, New Delhi., p317-314
33. Hur YJ, Lee SB, Shin DJ, Kim TH, Cho JH et al. (2016) Screening of rice germplasm for Bakanae disease resistance in rice. Kor J Breed Sci 48:22-28
34. Rana A, Sahgal M, Johri BN (2017) *Fusarium oxysporum*: Genomics, Diversity and Plant–Host Interaction. In Developments in Fungal Biology and Applied Mycology; Springer: Singapore, pp. 159–199
35. Jansen C, VonWettstein D, Schäfer W, Kogel KH, Felk A et al. (2005) Infection patterns in barley and wheat spikes inoculated with wild-type and trichodiene synthase gene disrupted *Fusarium graminearum*. Proc. Natl. Acad. Sci. 102:16892–16897
36. Elshafey RAS, Tahoon AM, El-Emary FA (2018) Analysis of varietal response to bakanae infection *Fusarium fujikuroi* and gibberellic acid through morphological, anatomical and hormonal

- changes in three rice varieties. Journal of Phytopathology and Pest Management 63-87
37. Bhandari DR, Wang Q, Li B, Friedt W, Römpp A et al. (2018) Histology-guided high-resolution AP-SMALDI mass spectrometry imaging of wheat-Fusarium graminearum interaction at the root–shoot junction. Plant methods 14.1: 103. <https://doi.org/10.1186/s13007-018-0368-6>
38. Sundaram RM, Vishnupriya MR, Laha GS, Rani NS, Rao PS et al. (2009) Introduction of bacterial blight resistance into Triguna, a high yielding, mid-early duration rice variety. Biotechnol J 4:400-407
39. Singh S, Sidhu JS, Huang N, Vikal Y, Li Z et al. (2001) Pyramiding three bacterial blight resistance genes (*xa5*, *xa13* and *Xa21*) using marker-assisted selection into *indica* rice cultivar PR106. Theor Appl Genet 102:1011-1015
40. Pradhan SK, Nayak DK, Mohanty S, Behera L, Barik SR et al. (2015) Pyramiding of three bacterial blight resistance genes for broad-spectrum resistance in deepwater rice variety, Jalmagna. Rice 8:2-14
41. Yasuda N, Mitsunaga T, Hayashi K, Koizumi S, Fujita Y (2015) Effects of pyramiding quantitative resistance genes *pi21*, *Pi34*, and *Pi35* on rice leaf blast disease. Plant Dis 99:904-909
42. Jena KK, Hechanova SL, Verdeprado H, Prahalada GD, Kim SR (2017) Development of 25 near-isogenic lines (NILs) with ten BPH resistance genes in rice (*Oryza sativa* L.): production, resistance spectrum, and molecular analysis. Theor Appl Genet. 130: 2345–2360

(A)

Ilpum Zenith

(B)

Figure1

Zenith

Ilpum

Figure2

Figure3

Figure4

Ilpum Zenith 680341 680449 680453 680470 680408 680400 680384

Figure5

Figure6