Acquired fluoroquinolone resistance genes in corneal isolates of *Pseudomonas aeruginosa*

Mahjabeen Khan¹, Stephen Summers², Scott A Rice^{2,3,4}, Fiona Stapleton¹, Mark D P Willcox¹, and Dinesh Subedi^{1,5*}

¹School of Optometry and Vision Science, University of New South Wales Sydney, Australia

²The Singapore Centre for Environment Life Sciences Engineering (SCElSE), Singapore

³The School of Biological Sciences, Nanyang Technological University, Singapore

⁴The iThree Institute, The University of Technology Sydney, Sydney, New South Wales, Australia

⁵School of Biological Sciences, Monash University, Clayton, Victoria, Australia

Key words: Antimicrobial resistance, Fluoroquinolones, *Pseudomonas aeruginosa*, eye, corneal ulcer, microbial keratitis

*** Corresponding Author:** dinesh.subedi@monash.edu

Abstract

Fluoroquinolones are widely used as an empirical therapy for pseudomonal ocular infections. Based on increasing reports on acquired fluoroquinolone resistance genes in clinical isolates of *Pseudomonas aeruginosa*, we investigated 33 strains of *P. aeruginosa* isolated from the cornea of microbial keratitis patients in India and Australia between 1992 and 2018 to understand the prevalence of acquired fluoroquinolone resistance genes in ocular isolates and to assess whether the possession of those genes was associated with fluoroquinolone susceptibility. We obtained the whole genome sequence of 33 isolates using Illumina MiSeq platform and investigated the prevalence of two fluoroquinolone resistance genes *crpP* and *qnrVC1*. To examine the associated mobile genetic elements of *qnrVC1* positive strains, we obtained long read sequences using Oxford Nanopore MinION and performed hybrid assembly to combine long reads with Illumina short sequence reads. We further assessed mutations in QRDRs and antibiotic susceptibilities to ciprofloxacin, levofloxacin and moxifloxacin to examine the association between resistance genes and phenotype. Twenty strains possessed *crpP* in genetic islands characterised by possession of integrative conjugative elements. The *qnrVC1* gene was carried by four isolates on class I integrons and Tn3 transposons along with aminoglycoside and beta-lactam resistance genes. We did not observe any evidence of plasmids carrying fluoroquinolone resistance genes. Resistance to fluoroquinolones was observed in those strains which possessed *crpP*, *qnrVC1* and that had QRDRs mutations. The presence of *crpP* was not a sole cause of fluoroquinolone resistance.

Introduction

Pseudomonas aeruginosa is a highly adaptable opportunistic pathogen which is ubiquitously present in the environment. This bacterium is naturally resistant to many antimicrobials and can acquire antibiotic resistance through mutations in chromosomal genes and lateral gene transfer [1, 2]. *P. aeruginosa* is associated with different types of human infections and because of emerging multidrug-resistant strains, these infections are major global public health concerns [3].

Fluoroquinolones are broad spectrum and widely prescribed antibiotics to treat pseudomonal infections including ocular infections [4-6]. Fluoroquinolone resistance in various clinical isolates is on the rise [7]. For example, a single centre study has shown that the prevalence of fluoroquinolone resistance *P. aeruginosa* increased from 15% to 41% in ten years [8]. This increase in fluoroquinolone resistance has been linked to the excessive use of the antibiotics [9]. The rate of isolation of fluoroquinolone resistant strains also depends on the type of infections; nosocomial isolates are more resistant than isolates from non-nosocomial sources [10]. In general, fluoroquinolone resistance is relatively low in ocular isolates of *P. aeruginosa* compared to other infections [11]. However, higher resistance rates have been reported in ocular isolates in certain regions of the world and like other systemic infections this rate has been increasing over time [12]. This has raised the concern that the horizontal transfer of fluoroquinolone resistance genes can be associated with spread of fluoroquinolone resistance in ocular isolates.

Mutations that alter target sites (DNA gyrase [*gyrA/gyrB*] and topoisomerase IV [*parC/parE*]) and increased membrane permeability are common mechanisms of fluoroquinolone resistance in *P. aeruginosa* [1, 13-15]. In other Gram-negative bacteria, fluoroquinolone resistance genes such as *qnr* have been shown to be carried on plasmids [16, 17]. The gene encodes a pentapeptide repeat protein which protects DNA gyrase and topoisomerase from the action of fluoroquinolones [18]. In contrast, carriage of *qnr* on plasmid is very rare in *P. aeruginosa* [19]. Despite this low carriage, recent studies have identified fluoroquinolone resistance genes in certain mobile genetic elements. *CrpP* and *qnrVC* can be carried on the *P. aeruginosa* mega plasmids pUM505 and pBM413, respectively [20, 21]. In addition, *qnrVC* was found to be associated with a class I integron, which also carried beta-lactamase genes [20, 22]. These studies were based on a single strain and the prevalence of acquired fluoroquinolone resistance genes in *P. aeruginosa* remains unclear. This led us to examine mobile fluoroquinolone resistance genes in *P. aeruginosa*. Given the concern of increasing fluoroquinolone resistance

in ocular isolates, we have undertaken sequence analysis of 33 ocular isolates of *P. aeruginosa*, isolated from corneal ulcers in the last 25 years to assess whether the possession of fluoroquinolone resistance genes was associated with fluoroquinolone susceptibility and whether resistance genes carrying strains had any genetic commonalities.

Materials and Methods

Pseudomonas aeruginosa strains

Isolates in this study were collected from the cornea of microbial keratitis patients in India and Australia between 1992 and 2018. They comprise 20 isolates collected for this study and used for the antibiotic susceptibility testing and whole genome sequencing. Information on antibiotic susceptibility and the whole genome data of 13 isolates was obtained from our previous study [23] (Supplementary Table 1). All isolates were retrieved from the culture collection of School of Optometry and Vision Science, the University of New South Wales, Australia without identifiable patients' data.

Antibiotic susceptibility testing of *Pseudomonas aeruginosa* strains

Susceptibility of *P. aeruginosa* isolates to ciprofloxacin (Sigma-Aldrich, Inc., St. Louis Missouri, USA), levofloxacin (Sigma-Aldrich, Inc.) and moxifloxacin (European Pharmacopoeia, Strasbourg, Cedex France) was investigated using the broth microdilution method following the protocol of Clinical and Laboratory Standard Institute [24] [25]. The MIC was taken as the lowest concentration of an antibiotic in which no noticeable growth (turbidity) was observed and the break point was established according to published standards [26]. Based on MICs, the isolates were categorised into four groups, susceptible ($\leq$ resistance break point), resistant ($>$ resistance break point – 32 $\mu\text{g/mL}$), highly resistant ($> 32 - 128 \mu\text{g/mL}$) and very highly resistant ($>128 \mu\text{g/mL}$) for the analysis.

DNA extraction and Illumina sequencing

Bacterial DNA was extracted from overnight cultures using a DNeasy Blood and Tissue Kit (Qiagen, Hilden, Germany) following the manufacturer's instructions. The extracted DNA was quantified and purity-checked using Nanodrop (NanoDrop Technologies, Wilmington, DE, USA), Qubit fluorometer (Life Technologies, Carlsbad, CA, USA), and 1% agarose gel electrophoresis. The extracted DNA was dried for transport to the sequencing facility at Singapore Centre for Environmental Life Sciences Engineering, Singapore. DNA was sequenced on MiSeq (Illumina, San Diego, CA, USA) generating 300 bp paired end reads. The paired-end library was prepared using Nextera XT DNA library preparation kit (Illumina, San Diego, CA, USA). All the libraries were multiplexed on one MiSeq run.

Bioinformatics analysis of short read sequences and construction of phylogenies

The quality of raw reads was analysed using online tool FastQC v0.11.7 (<https://www.bioinformatics.babraham.ac.uk/projects/fastqc>). Adaptor sequences were removed using Trimmomatic v0.38 with quality and length filtering (SLIDINGWINDOW:4: 15 MINLEN:36) [27]. The reads were de-novo assembled using Spades v3.13.0 with the programs' default setting [28] followed by annotation using Prokka v1.12 [29]. To examine acquired resistance genes, the genomes were uploaded into online database ResFinder v3.1 of Centre for Genomic Epidemiology, DTU, Denmark [30]. The contigs carrying *crpP* and *qnrVC1* genes were selected and analysed for mobile genetic elements using BLAST, Integron finder v1.5.1 [31] and IS finder [32]. The genome was visualised and manually modified using Geneious prime v2019.2 [33]. A figure of BLAST comparison was generated using EasyFig v2.2.2 [34]. To identify mutations in the QRDRs (*gyrA*, *gyrB*, *parC* and *parE*), the genome sequences were analysed using Snippy v4.2 with program's default settings (<https://github.com/tseemann/snippy>). The genomes were also examined for the presence of type IV secretion factors (*exoU* and *exoS*) using BLAST search, and for the presence of the CRISPR cas system using the CRISPRcasFinder online tool [35]. Codon adaptation index (CAI) [36] was examined to understand the possible expression of different orthologues of *crpP* using CAIcal [37, 38]. 50s ribosomal protein L19 (*rplS*), which is a highly expressed chromosomal gene was included as the reference to show the difference in expression level between chromosomal gene and acquired genes.

Core genome single nucleotide polymorphisms (SNPs) were identified using Parsnp v from Harvest Suite [39] using settings to exclude SNPs identified in regions that had arisen by recombination. The core genome SNPs were used to construct a maximum likelihood phylogenetic tree. All genomes were examined for multi-locus sequence type (MLST) using the MLST database. Nucleotide sequences of all MLST locus were extracted and concatenated to use in Bayesian phylogenetic analyses using BEAST2 v2.4.7 with the following parameters: gamma site heterogeneity model, Hasegawa-Kishino-Yano (HKY) substitution model and relaxed-clock log-normal [40]. BEAST 2 output was summarized using TreeAnnotator with a 5% burn in. The phylogenetic trees were visualised using iTol v4 [41].

MinION sequencing and analysis

Wizard Promega DNA extraction kits (Promega, Madison, WI) was used to extract DNA from overnight broth culture. DNA was quantified and transported as mentioned above. Long reads libraries were prepared using a rapid sequencing kit (RAD-SQK004, ONT, Oxford, UK) and subsequent sequencing was conducted using the MinIon flowcell (R9.4.1) for 48 hours. Long reads were basecalled using Guppy v3.3.0 and adapters removed using Porechop v0.2.4. Assembly of both long and short reads into a hybrid genome assembly was achieved with Unicycler v0.4.3 [42], opting for default parameters and using only short reads with merging pairs. Following assembly, all assemblies were assessed for quality using Quast v5.0.2.

Results

Population structure, phylogeny and fluoroquinolone resistance of *P. aeruginosa* strains

Whole genomes were analysed from 33 corneal isolates of *P. aeruginosa*, 20 of which were sequenced as a part of this study and 13 genome that were published previously [23]. All strains were isolated between 1992 and 2018 in India (19 isolates) or Australia (14 isolates) (Supplementary Table 1). Draft genomes were mapped against the reference genome *P. aeruginosa* PAO1 and a total of 202,232 SNPs were observed among the 33 isolates, which were used to construct core genome phylogeny using Parsnp v1.2 [43]. Similar to previous reports [23, 36, 44, 45], the phylogenetic tree based on the core genome revealed two major clades (Fig 1). These clades followed a similar pattern as for other previously described strains of *P. aeruginosa*, where *exoU* carrying strains clustered together in a single clade (Phylogroup 2) [23]. Our results showed that 18 (of 33) isolates carried *exoU*, of which 16 were clustered together in phylogroup 2, which contained predominantly Indian isolates. The detection rate of *exoU* was 68% in Indian isolates and 36% in Australian isolates.

Figure 1. Maximum likelihood phylogenetic tree based on core genome SNPs analysis using *Pseudomonas aeruginosa* PAO1 as the reference, excluding SNPs identified in regions that had arisen by recombination, using the default parameters of Parsnp v1.2 [43]. Isolates from India are labelled red and Australian isolates are labelled blue. Numbers given at the nodes represent bootstrap values. The presence of *crpP*, *exoU*, *qnrVC1*, and CRISPR cas are represented by red squares. Orange squares represent the presence of mutations in the quinolone resistance determining regions (QRDRs) and fluoroquinolone (CIP = Ciprofloxacin; LEVO = Levofloxacin; and MOX = Moxifloxacin) susceptibilities are shown as a heat map with the ranges indicated in the figure. The figure was drawn using iTol v4 [41].

We further examined acquired fluoroquinolone resistance genes, mutations in quinolone resistance determining region (QRDRs), the presence of CRISPR genes and susceptibility to three different fluoroquinolones (Fig 1). None of the Australian isolates regardless of phylogenetic grouping were resistant to fluoroquinolones. In contrast, 75% of isolates of phylogroup 2 were resistant to at least one fluoroquinolone. Of the 33 strains, 73.7 % (14 out of 19) of Indian and 42.8% (6 out of 14) of Australian isolates possessed *crpP*, which has been recently shown to be on a plasmid (pUM505) and associated with ciprofloxacin resistance [20]. However, eight (40%) *crpP* carrying strains in the current study regardless of country of isolation were not resistant to the fluoroquinolones, including ciprofloxacin (Fig 1). Eleven out of 14 fluoroquinolone resistance strains had mutations in both *gyrA* and *parC* and all except one carried the *exoU* gene. In addition, four strains from the latter cohort of 11 strains carried another fluoroquinolone resistance gene, *qnrVC1*, in combination with mutations in *gyrA* and *parC*, and this was associated with a very high MIC (>128 µg/mL) to all three fluoroquinolones (Fig 1).

Given that *exoU*, *crpP* and *qnrVC1* are components of the accessory genome which is mostly shaped by the CRISPR-Cas system, a bacterial defence system against foreign DNA [46], we searched isolates for CRISPR-Cas genes using the CRISPRCasFinder database [35] following software default parameters. Only one of the *exoU*, and none of the *qnrVC1* carrying strains, were positive for CRISPR-Cas genes. However, CRISPR-Cas was observed in seven *crpP* positive strains, which did not possess *exoU* and/or *qnrVC1* (Fig 1).

Figure 2. Consensus tree of 33 *P. aeruginosa* isolates, based on Bayesian evolutionary analysis by sampling trees (BEAST) of concatenated multi-locus sequence type (MLST) under strict clock analysis [40]. The tip of the tree was constrained by date of isolation. The time scale is shown in years at the top and each internal node is labelled with posterior probability limit. Isolates from India are labelled red and Australian isolates are labelled blue. The presence of genes *crpP*, *exoU*, *qnrVC1*, and CRISPRcas are represented by red squares. Orange square represents presence of mutations in quinolone resistance determining region (QRDRs). Fluoroquinolone (CIP = Ciprofloxacin; LEVO = Levofloxacin; and MOX = Moxifloxacin) susceptibilities are shown as heat maps in the grey scale indicated in the figure. The figure was drawn using iTol v4 [41].

Bayesian phylogenetic reconstruction

To examine the evolutionary trends of isolates, a Bayesian analysis was performed based on MLST sequences. The evolutionary tree was constructed by BEAST, which uses a molecular clock model to estimate time-measured phylogenies [40]. The result suggested that the most recent common ancestor of isolates of this study appeared approximately 50 years ago and the branching of fluoroquinolone resistant strains (except PA82), which had mutations in QRDRs and/or had acquired resistance *qnrVC1* within each subclade, occurred between 1988 and 2000 (Fig 2). This coincided with the time when ciprofloxacin and ofloxacin were introduced as antibiotic treatments [47]. The analysis suggests that acquisition of *crpP* and *exoU* occurred at least 50 years ago.

Prevalence of the *crpP* gene

A recent study has shown that *crpP* encodes for a protein that is associated with increased resistance to ciprofloxacin [20]. In the current study, 63% of isolates possessed this gene, however nine out of the 20 *crpP* carrying strains were not resistant to ciprofloxacin (Fig 1). Single nucleotide polymorphisms in the *crpP* sequences from all strains were compared to the sequence of *crpP* in pUM505 using local BLASTn. The BLAST matrix (Table 1) showed that the nucleotide sequences of *crpP* genes varied between 100% and 93.43%, indicating the possibility of differences in the activity of the encoded protein. Codon usage, presented as codon adaptation index [36] (Fig 3) was examined to understand the possible expression of different orthologues of *crpP* [38]. The 50s ribosomal protein L19 (*rplS*), which is a highly expressed chromosomal gene was included in this analysis to show the difference in expression between chromosomal and acquired genes. A CAI between 0.36 and 0.39 was observed in different orthologues of *crpP*, which while low, corresponded to divergence of sequences. However, the sequence divergence was not associated with resistance to fluoroquinolones. For example, the nucleotide sequence of *crpP* of a resistant strain PA32 and a sensitive strain PA127 were 98.48% identical to that of pUM505.

This study also investigated the presence of the *crpP* homologues in NCBI databases using local BLAST with a cut off of 80% coverage and 90% sequence identity to examine the distribution of *crpP* gene in the bacterial database including plasmids as of 2019-06-15. The BLAST search against *P. aeruginosa* complete genome databases returned 52% matches and this rate of detection is broadly comparable to *crpP* prevalence observed in the current strains. No BLAST matches other than *P. aeruginosa* were observed with this search parameter. Within the *P. aeruginosa* plasmid database, two matches were observed against 45 complete plasmids using the parameters mentioned above. A CrpP-like protein, which has sequence identity of 10-46% with *P. aeruginosa crpP*, has been observed in species of the *Enterobacteriaceae* [48]. This result suggested that *crpP* can be transferred between species or strains.

Table 1: BLAST matrix showing nucleotide identity (in percentage) of *crpP* sequences of each isolate. Colour intensity corresponds to percentage identity.

	pUM505*	PA31	PA32	PA33	PA35	PA37	PA127	PA162	PA220	PA17	PA34	PA157	PA206	PA217	PA181	PA216	PA202	PA219	PA221	PA176	PA198
pUM505*		98.48	98.48	98.48	98.48	98.48	98.48	98.48	98.48	97.98	97.98	97.98	97.98	97.98	96.46	95.45	94.95	94.95	94.95	94.44	94.44
PA31	98.48		100	100	100	100	100	98.99	100	99.49	98.48	98.48	98.48	98.48	97.98	95.96	95.45	95.45	95.45	94.95	94.95
PA32	98.48	100		100	100	100	100	98.99	100	99.49	98.48	98.48	98.48	98.48	97.98	95.96	95.45	95.45	95.45	94.95	94.95
PA33	98.48	100	100		100	100	100	98.99	100	99.49	98.48	98.48	98.48	98.48	97.98	95.96	95.45	95.45	95.45	94.95	94.95
PA35	98.48	100	100	100		100	100	98.99	100	99.49	98.48	98.48	98.48	98.48	97.98	95.96	95.45	95.45	95.45	94.95	94.95
PA37	98.48	100	100	100	100		100	98.99	100	99.49	98.48	98.48	98.48	98.48	97.98	95.96	95.45	95.45	95.45	94.95	94.95
PA127	98.48	100	100	100	100	100		98.99	100	99.49	98.48	98.48	98.48	98.48	97.98	95.96	95.45	95.45	95.45	94.95	94.95
PA162	98.48	98.99	98.99	98.99	98.99	98.99	98.99		98.99	98.48	98.48	99.49	99.49	99.49	97.47	94.95	94.44	94.44	94.44	94.95	94.95
PA220	98.48	100	100	100	100	100	100	98.99		99.49	98.48	98.48	98.48	98.48	97.98	95.96	95.45	95.45	95.45	94.95	94.95
PA17	97.98	99.49	99.49	99.49	99.49	99.49	99.49	98.48	99.49		98.99	97.98	97.98	97.98	97.47	96.46	95.96	95.96	95.96	94.44	94.44
PA34	97.98	98.48	98.48	98.48	98.48	98.48	98.48	98.48	98.48	98.99		98.99	98.99	98.99	97.47	95.45	94.95	94.95	94.95	93.43	93.43
PA157	97.98	98.48	98.48	98.48	98.48	98.48	98.48	99.49	98.48	97.98	98.99		100	100	97.98	94.44	93.94	93.94	93.94	94.44	94.44
PA206	97.98	98.48	98.48	98.48	98.48	98.48	98.48	99.49	98.48	97.98	98.99	100		100	97.98	94.44	93.94	93.94	93.94	94.44	94.44
PA217	97.98	98.48	98.48	98.48	98.48	98.48	98.48	99.49	98.48	97.98	98.99	100	100		97.98	94.44	93.94	93.94	93.94	94.44	94.44
PA181	96.46	97.98	97.98	97.98	97.98	97.98	97.98	97.47	97.98	97.47	97.47	97.98	97.98	97.98		93.94	93.43	93.43	93.43	93.43	93.43
PA216	95.45	95.96	95.96	95.96	95.96	95.96	95.96	94.95	95.96	96.46	95.45	94.44	94.44	94.44	93.94		99.49	99.49	99.49	93.94	93.94
PA202	94.95	95.45	95.45	95.45	95.45	95.45	95.45	94.44	95.45	95.96	94.95	93.94	93.94	93.94	93.43	99.49		100	100	93.43	93.43
PA219	94.95	95.45	95.45	95.45	95.45	95.45	95.45	94.44	95.45	95.96	94.95	93.94	93.94	93.94	93.43	99.49	100		100	93.43	93.43
PA221	94.95	95.45	95.45	95.45	95.45	95.45	95.45	94.44	95.45	95.96	94.95	93.94	93.94	93.94	93.43	99.49	100	100		93.43	93.43
PA176	94.44	94.95	94.95	94.95	94.95	94.95	94.95	94.95	94.95	94.44	93.43	94.44	94.44	94.44	93.43	93.94	93.43	93.43	93.43		100
PA198	94.44	94.95	94.95	94.95	94.95	94.95	94.95	94.95	94.95	94.44	93.43	94.44	94.44	94.44	93.43	93.94	93.43	93.43	93.43	100

Figure 3. Codon adaptation index of *crpP* orthologues of *P. aeruginosa* strains. PA3742_rplS denotes 50s ribosomal protein L19 (rplS) of *P. aeruginosa* PAO1.

crpP associated mobile genetic elements

Although the *crpP* gene was first reported in plasmid pUM505 of a clinical strain of *P. aeruginosa* [20], BLAST analysis, as mentioned above, revealed this gene was mostly associated with the chromosome and its presence in plasmids was rare. The *crpP* containing contigs of each isolate were assessed to determine similarities between associated genomic islands. The *crpP* flanking region of pUM505 is homologous to the pathogenicity island PAPI-1 of *P. aeruginosa* PA14 [49] and pKLC102 of *P. aeruginosa* C [50, 51], which is an integrative conjugative element and is associated with carriage of virulence genes. A PAPI-1-like genomic island has also been reported in a complete genome of the ocular isolate PA34 [52]. Using PAPI-1 of PA34 as a reference, the contigs of each strain were compared and this showed that there were similarities between contigs. Those common regions were extracted and annotated manually using Geneious Prime v2019.1.3 [33]. After annotation, all of the extracted regions were compared using Easyfig [34]. The region that has been shared amongst all isolates is presented in Fig 4, where the colour code represents open reading frames. While the size of *crpP* carrying genomic islands differed between strains, the exact size of islands could not be ascertained because of the use of draft genomes for analysis. Nevertheless, three genes (integrative conjugative element, an unknown protein and single-strand binding protein) upstream and one gene (type I DNA topoisomerase) downstream of *crpP* were common in all isolates and this indicates that these genes may be involved in the transfer of *crpP* gene. A characteristic type IV pili synthesis operon was associated with 75% of the *crpP* positive strains. There was no association between size of *crpP* islands and fluoroquinolone resistance.

Figure 4. Comparison of *crpP*-associated genomic islands of *P. aeruginosa* strains. Protein-coding regions are represented by the orange arrows and common key features/associated genes among all strains are shown in various coloured arrows. The gradient pink and grey shading represent regions of nucleotide sequence identity (100% to 64%) in forward and reverse directions respectively, determined by BLASTn analysis. Figures are drawn to scale using Easyfig [34]. (parA= plasmid partition protein A, dnaB= replicative DNA helicase, yejK= nucleoid-associated protein, ICE= integrating conjugative element protein, ssb= single-stranded DNA-binding protein, crpP= ciprofloxacin resistance protein, vapC= tRNA(fMet)-specific endonuclease, topA= type I DNA topoisomerase, LysE= LysE family translocator, LysR= LysR family transcriptional regulator, toxin-anti= type II toxin-antitoxin system RelE/ParE family toxin, pilL-pilM= type IV pilus synthesis operon).

Prevalence of quinolone resistance protein (*qnr* gene)

The *qnr* is commonly carried on plasmids, especially in the *Enterobacteriaceae*. A transferable variant of the *qnr* gene, *qnrVC1* was first identified in a class I integron in *Vibrio cholerae* 01 [53]. Several *qnrVC* alleles have also been reported to be carried in various bacteria including *P. aeruginosa* [18, 54]. For example, a recent study had reported that 2.3% of clinical strains of *P. aeruginosa* carried *qnrVC* [55]. In the current study, 12% (four out of 33) of isolates possessed *qnrVC1*. All these four were isolated in 2017 and 2018. We further examined the prevalence of *qnrVC1* homologues in complete chromosomes and plasmids of other bacterial species in NCBI database using BLAST search. We observed that *qnrVC1*-like proteins were present in several bacterial species. Upon limiting the BLAST criteria to 100% coverage and 98% identity, the search returned with 19 matches from species of *Pseudomonas*, *Vibrio* and *Aeromonas*. (Supplementary Figure 1). This suggests that the *qnrVC1* gene is prevalent in various bacterial families. Curiously, the *qnrVC1* gene was not identified in the plasmid database of NCBI. Furthermore, *qnrVC1* had a G + C content of 37%, which is considerably different from that of the *P. aeruginosa* genome (66%). This evidence along with the presence of *qnrVC1* homologues in several bacterial species indicates that *qnrVC1* is a horizontally acquired gene in *P. aeruginosa*, which is potentially acquired through inter-species or inter-genera transfer.

qnrVC1 associated mobile genetic elements

We further examined the flanking nucleotide sequences of *qnrVC1* to understand the mobile genetic elements associated with transfer of the gene. Our initial analysis found that *qnrVC1* was present in small contigs (contig size <5000 bp) and this limited our aim to examine mobile genetic elements. The four *qnrVC1* positive strains were then analysed using the long-read sequencing technique ONT minION (ONT, Oxford). With hybrid assembly of long and short sequencing reads, we archived long contigs ranging in size from 400 Kbp to 2000 Kbp (PA202 N₅₀ = 680,290; PA219 N₅₀ = 2,159,014; PA221 N₅₀ = 2,110,541) in three isolates and a complete and closed genome of one strain PA198 (N₅₀ = 7,188,952).

The *qnrVC1* gene was present in two different types of mobile genetic elements. In strains PA198 and PA219, the gene was integrated into Tn3 transposons, which have 99% identity with one another (Fig 5). In addition, the Tn3 transposon carried other antibiotic resistance genes associated with aminoglycoside 3'-phosphotransferase and tetracycline resistance. BLAST comparison revealed recombination in various places within the transposon, most notably in the areas flanking the *qnrVC1-VOC* and *tetR-tetA* genes (Fig 5), which represents the site of integration of these genes. The role of VOC protein in antibiotic resistance is not well known. However, evidence suggests that proteins of VOC the family include beomycin and fosfomycin resistance proteins [56]. Integration of VOC with other antibiotic resistance

genes leads us to hypothesise that VOC can be associated with antibiotic resistance. Further research is required to investigate its role in antibiotic resistance in *P. aeruginosa*.

Figure 5. Comparison of *qnrVC1*-associated genomic islands of *P. aeruginosa* strains. Protein-coding regions are represented by the arrows and common key features/associated genes among all strains are shown in various coloured arrows. The gradient blue and red shading represent regions of nucleotide sequence identity (100% to 65%) in forward and reverse directions, respectively, determined by BLASTn analysis. The sequences are from strains top to bottom; PA198, PA219, PA202 and PA221. Figures are drawn to scale using Easyfig [34]. (Tn3-tnp = Tn3-transposase HP= hypothetical protein, ydhC= Inner membrane transport protein, tet(R)-tet(g) = tetracycline resistance genes, *qnrVC1*=quinolone resistance gene, VOC = VOC family protein, aph(6)-Id = aminoglycoside resistance protein, DNA-inv = DNA invertase, aadA10= aminoglycoside resistance protein, IS110= IS110 family transposase, qacEdealt1= quaternary ammonium compound-resistance protein, sul1= Dihydropteroate synthase, blaLcr-1=beta-lactamase gene, attC = recombination sites of gene cassette, attI = integron recombination site, IR=invert repeat).

In the other two strains, PA202 and PA221, *qnrVC1* was carried by identical class 1 integrons (Fig 5), along with two additional antibiotic resistance genes; the aminoglycoside gene *aadA10* which was within the integron and the beta-lactam gene *blaLcr-1* in the flanking region. Other studies have also shown that *qnrVC1* is a part of integrons that carry multiple resistance genes [22, 54]. The class 1 integron observed here had a typical structure containing promoter sequences and an attachment site (*attI*) required for integration and transcription of gene cassettes [57]. The *qnrVC1* gene in all four strains was associated with a perfectly matched gene recombination site (*attC*). The *attC* sequence of *qnrVC1* gene cassette had 100% identity with that of *Vibrio parahaemolyticus* (GenBank accession number EU436855), supporting the likelihood that the *Vibrionaceae* is a source of *qnr* in *P. aeruginosa* [58].

Discussion

While fluoroquinolones are the preferred empirical therapy for corneal ulcers, which are often caused by *P. aeruginosa*, increasing resistance in this bacterium raises the concern about the efficacy of these antibiotics [4-11, 59, 60]. Acquired fluoroquinolone resistance genes are starting to be reported in clinical isolates of *P. aeruginosa* [20-22, 48]. However, the rate of carriage of acquired fluoroquinolone resistance genes in ocular isolates has not been reported to date. This study examined the prevalence of acquired fluoroquinolone resistance genes, *crpP* and *qnrVC1* genes, which have recently been shown to be quinolone-associated acquired resistance genes in *P. aeruginosa*. We used whole genomes of 33 strains isolated from corneal ulcers in the last 25 years isolates from Australia and India. Fluoroquinolone resistance was common in Indian isolates and mutations in QRDRs were associated with this increased MIC to all three fluoroquinolones tested. Possession of the acquired gene *qnrVC1* substantially increased the MICs whereas *crpP* was not associated with increased fluoroquinolone resistance in those isolates that carried *crpP* but lacked QRDRs mutations or *qnrVC1*. Both quinolone resistance genes were integrated into the chromosome and their associated genomic islands were broadly common between strains.

Fluoroquinolone resistance in *P. aeruginosa* is usually associated with mutations in the genes *mexR*, *nfxB*, and *mexT* that lead to overexpression of efflux pumps and QRDRs of *gyrA*, *gyrB*, *parC* and *parE* that alter drug target sites [1, 13-15]. In addition, horizontally transferred fluoroquinolone resistance genes such as *crpP* and *qnr*-variants have been recently reported in clinical strains of *P. aeruginosa* [20-22, 55]. Our BLAST analysis against the NCBI database showed that more than half of strains in the *P. aeruginosa* complete genome database contained homologues of *crpP* and this corresponded to the detection frequency of *crpP* in the ocular strains reported in this manuscript. *CrpP* carriage rate was higher in Indian isolates (74%) compared to Australian isolates (43%) and approximately half of the *crpP* carrying strains regardless of region of isolation were not resistant to any of the fluoroquinolones tested. Furthermore, fluoroquinolone resistant *crpP*⁺ strains also had target site mutations *gyrA* and *parC* and had acquired the quinolone resistance gene *qnrVC1*. Therefore, the role of *crpP* alone on fluoroquinolone resistance could not be ascertained for these strains. Given that *crpP* is ciprofloxacin specific and associated with low-level resistance, its presence may not necessarily be responsible for MICs that exceeded the clinical breakpoint [20].

The *crpP* gene was associated with similar genomic islands in all strains. These genomic islands were characterised by the possession of integrative conjugative elements and DNA replication factors. The *crpP* gene was originally reported in the plasmid pUM505 [51] but there are no other studies showing plasmid associated *crpP* in *P. aeruginosa*. Therefore, we performed the BLAST search against the NCBI plasmid database specific to *P. aeruginosa* and observed two matches (plasmids pKLC102 and pY89) out of 45 complete plasmids. Furthermore, the most recent ancestors of *crpP* positive strains appeared at least 50 years ago, which is earlier than the appearance of resistant subclades in the phylogeny. This result, together with finding of high divergence in *crpP* orthologues between strains and lower CAI compared to a highly expressed chromosomal gene (*rlpS*) [61] suggests that acquisition of the *crpP* gene was a relatively old evolutionary event in *P. aeruginosa*. In addition, the presence of CRISPR-Cas genes in several *crpP* positive strains supports the latter hypothesis and further suggests that *crpP* was integrated into the chromosome in an early event.

The plasmid-mediated quinolone resistance *qnr* gene was first reported in 1998 [16] and this corresponds with increased usage of fluoroquinolones in the 1980s [62]. Our analysis also

revealed that resistant *P. aeruginosa* strains were phylogenetically separated from susceptible strains from the 1990s. The *qnr* genes protect DNA-gyrase and topoisomerase IV to prevent quinolone actions and confer low-level fluoroquinolones resistance, often below the clinical breakpoint [63, 64]. The *qnrVC1* gene is a mobile quinolone resistance gene that is associated with class I integrons [53]. *QnrVC1* and another variant *qnrVC6* have been reported from clinical *P. aeruginosa* strains isolated between 2007 and 2012 [21, 54]. However, in this study, the acquired fluoroquinolone resistance gene *qnrVC1* was observed in isolates sampled in 2017 and 2018, indicating that transferable fluoroquinolone resistance genes may be recently acquired or passed unnoticed in ocular isolates of *P. aeruginosa*. No strains contained *qnrVC6*. We also observed that *qnrVC1* was associated with a Tn3 transposon, which has not been previously reported in *P. aeruginosa*. Integration of several other antibiotic resistance genes in these mobile genetic elements suggests these elements may concentrate antibiotic resistance genes. *QnrVC* is reported predominantly in water bacteria including *Aeromonas* spp., and *Acinetobacter* spp [65]. In this study the recombination site (*attC*) of all four *qnrVC1* alleles had 100% identity with aquatic bacteria *Vibrio parahaemolyticus*. This corresponds with our previous study in ocular isolates of *P. aeruginosa*, where acquired resistance genes closely matched with that of environmental isolates [52, 66].

However, the current study showed that *qnrVC1* carrying strains also had QRDRs mutations and were highly resistant to all three fluoroquinolones compared to those lacking *qnrVC1*. This indicates that possession of *qnrVC1* potentially facilitates selection of strains with high-level fluoroquinolone resistance. Given that *qnrVC1* has been reported as being responsible for low-level fluoroquinolone resistance [18], the high MICs observed for these isolates is not clear but may indicate some synergistic activity between the QRDR mutations and *qnrVC1*.

Like in many other studies, fluoroquinolone resistant strains were also associated with possession of *exoU*, which is the predominate virulent genotype of *P. aeruginosa* in ocular infections [67-77]. Although the reason behind this predominance is not completely understood, the concurrent occurrence of transferable fluoroquinolone resistance genes and *exoU* may heighten the concern of selection of more virulent strains during antibiotic therapy.

In conclusion, mobile genetic elements play a key role in the rapid spread of antibiotic resistance genes. This study provides evidence for carrying of fluoroquinolone resistance genes *qnrVC1* and *crpP* in ocular isolates of *P. aeruginosa*. The *qnrVC1* gene was mobilised by a class I integron and Tn3 transposon, and associated with other antibiotics resistance genes. The *crpP* gene may have evolved prior to other transferable fluoroquinolone resistance genes. Although possession of these genes has not been shown to be associated with high level of fluoroquinolone resistance, *qnrVC1* in strains with *gyrA* and *parC* mutations could be associated with very high fluoroquinolone resistance. Further studies on a larger number of fluoroquinolone resistance strains, including molecular analysis of combined effects of mutations in resistance, should be undertaken to provide further insights into the role of acquired fluoroquinolone resistance gene in clinically significant resistance in *P. aeruginosa*.

Nucleotide accession

The nucleotide sequences are available in the GenBank under the Bio project accession number PRJNA590804 and PRJNA431326.

Acknowledgements

The authors would like to acknowledge the Singapore Centre for Environmental Life Sciences Engineering (SCELS), whose research is supported by the National Research Foundation Singapore, Ministry of Education, Nanyang Technological University and National University of Singapore, under its Research Centre of Excellence Programme. We are also thankful to

UNSW high performance computing facility KATANA for providing us cluster time for data analysis.

Conflicts of interest

No conflict of interest is declared by all authors.

References

1. Livermore DM. Multiple mechanisms of antimicrobial resistance in *Pseudomonas aeruginosa*: our worst nightmare? Clinical infectious diseases : an official publication of the Infectious Diseases Society of America. 2002;34(5):634-40. doi: 10.1086/338782. PubMed PMID: 11823954.
2. Blair JM, Webber MA, Baylay AJ, Ogbolu DO, Piddock LJ. Molecular mechanisms of antibiotic resistance. Nature reviews Microbiology. 2015;13(1):42-51. Epub 2014/12/02. doi: 10.1038/nrmicro3380. PubMed PMID: 25435309.
3. WHO priority pathogens list for R&D of new antibiotics [Internet]. 2017.
4. Linder JA, Huang ES, Steinman MA, Gonzales R, Stafford RS. Fluoroquinolone prescribing in the United States: 1995 to 2002. Am J Med. 2005;118(3):259-68. Epub 2005/03/05. doi: 10.1016/j.amjmed.2004.09.015. PubMed PMID: 15745724.
5. Blondeau JM. Fluoroquinolones: mechanism of action, classification, and development of resistance. Survey of ophthalmology. 2004;49 Suppl 2:S73-8. Epub 2004/03/19. doi: 10.1016/j.survophthal.2004.01.005. PubMed PMID: 15028482.
6. Smith A, Pennefather PM, Kaye SB, Hart CA. Fluoroquinolones: place in ocular therapy. Drugs. 2001;61(6):747-61. Epub 2001/06/12. PubMed PMID: 11398907.
7. Neuhauser MM, Weinstein RA, Rydman R, Danziger LH, Karam G, Quinn JP. Antibiotic resistance among gram-negative bacilli in US intensive care units: implications for fluoroquinolone use. Jama. 2003;289(7):885-8. Epub 2003/02/18. PubMed PMID: 12588273.
8. Gasink LB, Fishman NO, Weiner MG, Nachamkin I, Bilker WB, Lautenbach E. Fluoroquinolone-resistant *Pseudomonas aeruginosa*: assessment of risk factors and clinical impact. Am J Med. 2006;119(6):526.e19-25. Epub 2006/06/06. doi: 10.1016/j.amjmed.2005.11.029. PubMed PMID: 16750968.
9. Werner NL, Hecker MT, Sethi AK, Donskey CJ. Unnecessary use of fluoroquinolone antibiotics in hospitalized patients. BMC infectious diseases. 2011;11:187. Epub 2011/07/07. doi: 10.1186/1471-2334-11-187. PubMed PMID: 21729289; PubMed Central PMCID: PMC3145580.
10. Yayan J, Ghebremedhin B, Rasche K. Antibiotic resistance of *Pseudomonas aeruginosa* in pneumonia at a single University hospital center in Germany over a 10-year period. PloS one. 2015;10(10):e0139836. doi: 10.1371/journal.pone.0139836. PubMed PMID: 26430738; PubMed Central PMCID: PMC4592231.
11. Subedi D, Vijay AK, Willcox M. Overview of mechanisms of antibiotic resistance in *Pseudomonas aeruginosa*: an ocular perspective. Clinical & experimental optometry. 2018;101(2):162-71. Epub 2017/10/19. doi: 10.1111/cxo.12621. PubMed PMID: 29044738.
12. Smitha S, Lalitha P, Prajna VN, Srinivasan M. Susceptibility trends of *Pseudomonas* species from corneal ulcers. Indian journal of medical microbiology. 2005;23(3):168-71. PubMed PMID: 16100422.
13. Higgins PG, Fluit AC, Milatovic D, Verhoef J, Schmitz FJ. Mutations in GyrA, ParC, MexR and NfxB in clinical isolates of *Pseudomonas aeruginosa*. International journal of antimicrobial agents. 2003;21(5):409-13. PubMed PMID: 12727072.
14. Nakano M, Deguchi T, Kawamura T, Yasuda M, Kimura M, Okano Y, et al. Mutations in the gyrA and parC genes in fluoroquinolone-resistant clinical isolates of

- Pseudomonas aeruginosa*. Antimicrobial agents and chemotherapy. 1997;41(10):2289-91. Epub 1997/10/23. PubMed PMID: 9333065; PubMed Central PMCID: PMC164110.
15. Hooper DC. Mechanisms of fluoroquinolone resistance. Drug resistance updates : reviews and commentaries in antimicrobial and anticancer chemotherapy. 1999;2(1):38-55. doi: 10.1054/drup.1998.0068. PubMed PMID: 11504468.
16. Martinez-Martinez L, Pascual A, Jacoby GA. Quinolone resistance from a transferable plasmid. Lancet. 1998;351(9105):797-9. doi: 10.1016/S0140-6736(97)07322-4. PubMed PMID: 9519952.
17. Jacoby GA. Mechanisms of resistance to quinolones. Clinical infectious diseases : an official publication of the Infectious Diseases Society of America. 2005;41 Suppl 2:S120-6. Epub 2005/06/09. doi: 10.1086/428052. PubMed PMID: 15942878.
18. Robicsek A, Jacoby GA, Hooper DC. The worldwide emergence of plasmid-mediated quinolone resistance. The Lancet Infectious diseases. 2006;6(10):629-40. Epub 2006/09/30. doi: 10.1016/S1473-3099(06)70599-0. PubMed PMID: 17008172.
19. Cayci YT, Coban AY, Gunaydin M. Investigation of plasmid-mediated quinolone resistance in *Pseudomonas aeruginosa* clinical isolates. Indian journal of medical microbiology. 2014;32(3):285-9. Epub 2014/07/11. doi: 10.4103/0255-0857.136567. PubMed PMID: 25008822.
20. Chávez-Jacobo VM, Hernández-Ramírez KC, Romo-Rodríguez P, Pérez-Gallardo RV, Campos-García J, Gutiérrez-Corona JF, et al. CrpP Is a novel ciprofloxacin-modifying enzyme encoded by the *Pseudomonas aeruginosa* pUM505 plasmid. Antimicrobial agents and chemotherapy. 2018;62(6):e02629-17. doi: 10.1128/AAC.02629-17.
21. Liu J, Yang L, Chen D, Peters BM, Li L, Li B, et al. Complete sequence of pBM413, a novel multidrug resistance megaplasmid carrying qnrVC6 and blaIMP-45 from *Pseudomonas aeruginosa*. International journal of antimicrobial agents. 2018;51(1):145-50. Epub 2017/09/20. doi: 10.1016/j.ijantimicag.2017.09.008. PubMed PMID: 28923459.
22. Kocsis B, Toth A, Gulyas D, Ligeti B, Katona K, Rokusz L, et al. Acquired qnrVC1 and blaNDM-1 resistance markers in an international high-risk *Pseudomonas aeruginosa* ST773 clone. Journal of medical microbiology. 2019;68(3):336-8. Epub 2019/01/23. doi: 10.1099/jmm.0.000927. PubMed PMID: 30667355.
23. Subedi D, Vijay AK, Kohli GS, Rice SA, Willcox M. Comparative genomics of clinical strains of *Pseudomonas aeruginosa* strains isolated from different geographic sites. Scientific reports. 2018;8(1):15668. Epub 2018/10/26. doi: 10.1038/s41598-018-34020-7. PubMed PMID: 30353070; PubMed Central PMCID: PMC6199293.
24. CLSI. Clinical and Laboratory Standards Institute(CLSI). Performance standards for antimicrobial susceptibility testing; twenty-second information supplement. CLSI document M100-S22 32 (2012). 2012.
25. Clinical and Laboratory Standards Institute (CLSI). Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically; approved standard—Ninth Edition. CLSI. 2012;32(M07-A9).
26. Clinical and Laboratory Standards Institute (CLSI). Performance standards for antimicrobial susceptibility testing; twenty-second information supplement. CLSI document M100-S22. 2012;32(3).
27. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics (Oxford, England). 2014;30(15):2114-20.
28. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. J Comput Biol. 2012;19(5):455-77. doi: 10.1089/cmb.2012.0021. PubMed PMID: 22506599; PubMed Central PMCID: PMC3342519.

29. Seemann T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* (Oxford, England). 2014;30(14):2068-9. doi: 10.1093/bioinformatics/btu153. PubMed PMID: 24642063.
30. Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, et al. Identification of acquired antimicrobial resistance genes. *Journal of Antimicrobial Chemotherapy*. 2012;67(11):2640-4. doi: 10.1093/jac/dks261.
31. Cury J, Jove T, Touchon M, Neron B, Rocha EP. Identification and analysis of integrons and cassette arrays in bacterial genomes. *Nucleic acids research*. 2016;44(10):4539-50. doi: 10.1093/nar/gkw319. PubMed PMID: 27130947; PubMed Central PMCID: PMC4889954.
32. Siguier P, Perochon J, Lestrade L, Mahillon J, Chandler M. ISfinder: the reference centre for bacterial insertion sequences. *Nucleic acids research*. 2006;34(Database issue):D32-6. doi: 10.1093/nar/gkj014. PubMed PMID: 16381877; PubMed Central PMCID: PMC1347377.
33. Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, et al. Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* (Oxford, England). 2012;28(12):1647-9. Epub 2012/05/01. doi: 10.1093/bioinformatics/bts199. PubMed PMID: 22543367; PubMed Central PMCID: PMC3371832.
34. Sullivan MJ, Petty NK, Beatson SA. Easyfig: a genome comparison visualizer. *Bioinformatics* (Oxford, England). 2011;27(7):1009-10. Epub 2011/02/01. doi: 10.1093/bioinformatics/btr039. PubMed PMID: 21278367; PubMed Central PMCID: PMC3065679.
35. Couvin D, Bernheim A, Toffano-Nioche C, Touchon M, Michalik J, Néron B, et al. CRISPRCasFinder, an update of CRISPRfinder, includes a portable version, enhanced performance and integrates search for Cas proteins. *Nucleic acids research*. 2018;46(W1):W246-W51. doi: 10.1093/nar/gky425.
36. Stewart L, Ford A, Sangal V, Jeukens J, Boyle B, Kukavica-Ibrulj I, et al. Draft genomes of 12 host-adapted and environmental isolates of *Pseudomonas aeruginosa* and their positions in the core genome phylogeny. *Pathog Dis*. 2014;71(1):20-5. doi: 10.1111/2049-632X.12107. PubMed PMID: 24167005.
37. Puigbò P, Bravo IG, Garcia-Vallve S. CAIcal: a combined set of tools to assess codon usage adaptation. *Biol Direct*. 2008;3:38. Epub 2008/09/18. doi: 10.1186/1745-6150-3-38. PubMed PMID: 18796141; PubMed Central PMCID: PMC2553769.
38. Sharp PM, Li WH. The codon Adaptation Index--a measure of directional synonymous codon usage bias, and its potential applications. *Nucleic acids research*. 1987;15(3):1281-95. Epub 1987/02/11. doi: 10.1093/nar/15.3.1281. PubMed PMID: 3547335; PubMed Central PMCID: PMC340524.
39. Treangen TJ, Ondov BD, Koren S, Phillippy AM. The Harvest suite for rapid core-genome alignment and visualization of thousands of intraspecific microbial genomes. *Genome biology*. 2014;15(11):524.
40. Drummond AJ, Rambaut A. BEAST: Bayesian evolutionary analysis by sampling trees. *BMC Evol Biol*. 2007;7:214. Epub 2007/11/13. doi: 10.1186/1471-2148-7-214. PubMed PMID: 17996036; PubMed Central PMCID: PMC2247476.
41. Letunic I, Bork P. Interactive Tree Of Life (iTOL) v4: recent updates and new developments. *Nucleic acids research*. 2019. doi: 10.1093/nar/gkz239. PubMed PMID: 30931475.
42. Wick RR, Judd LM, Gorrie CL, Holt KE. Unicycler: resolving bacterial genome assemblies from short and long sequencing reads. *PLoS computational biology*. 2017;13(6):e1005595.

43. Treangen TJ, Ondov BD, Koren S, Phillippy AM. The Harvest suite for rapid core-genome alignment and visualization of thousands of intraspecific microbial genomes. *Genome Biol.* 2014;15(11):524. doi: 10.1186/PREACCEPT-2573980311437212. PubMed PMID: 25410596; PubMed Central PMCID: PMC4262987.
44. Jeukens J, Kukavica-Ibrulj I, Emond-Rheault JG, Freschi L, Levesque RC. Comparative genomics of a drug-resistant *Pseudomonas aeruginosa* panel and the challenges of antimicrobial resistance prediction from genomes. *FEMS Microbiol Lett.* 2017;364(18). Epub 2017/09/20. doi: 10.1093/femsle/fnx161. PubMed PMID: 28922838.
45. Freschi L, Jeukens J, Kukavica-Ibrulj I, Boyle B, Dupont MJ, Laroche J, et al. Clinical utilization of genomics data produced by the international *Pseudomonas aeruginosa* consortium. *Frontiers in microbiology.* 2015;6:1036. Epub 2015/10/21. doi: 10.3389/fmicb.2015.01036. PubMed PMID: 26483767; PubMed Central PMCID: PMC4586430.
46. van Belkum A, Soriaga LB, LaFave MC, Akella S, Veyrieras JB, Barbu EM, et al. Phylogenetic Distribution of CRISPR-Cas Systems in Antibiotic-Resistant *Pseudomonas aeruginosa*. *mBio.* 2015;6(6):e01796-15. Epub 2015/11/26. doi: 10.1128/mBio.01796-15. PubMed PMID: 26604259; PubMed Central PMCID: PMC4669384.
47. Emmerson AM, Jones AM. The quinolones: decades of development and use. *The Journal of antimicrobial chemotherapy.* 2003;51 Suppl 1:13-20. Epub 2003/04/19. doi: 10.1093/jac/dkg208. PubMed PMID: 12702699.
48. Chávez-Jacobo VM, Hernández-Ramírez KC, Silva-Sánchez J, Garza-Ramos U, Barrios-Camacho H, Ortiz-Alvarado R, et al. Prevalence of the *crpP* gene conferring decreased ciprofloxacin susceptibility in enterobacterial clinical isolates from Mexican hospitals. *Journal of Antimicrobial Chemotherapy.* 2019;74(5):1253-9. doi: 10.1093/jac/dky562.
49. He J, Baldini RL, Deziel E, Saucier M, Zhang Q, Liberati NT, et al. The broad host range pathogen *Pseudomonas aeruginosa* strain PA14 carries two pathogenicity islands harboring plant and animal virulence genes. *Proceedings of the National Academy of Sciences of the United States of America.* 2004;101(8):2530-5. PubMed PMID: 14983043; PubMed Central PMCID: PMC356984.
50. Kung VL, Ozer EA, Hauser AR. The accessory genome of *Pseudomonas aeruginosa*. *Microbiology and molecular biology reviews : MMBR.* 2010;74(4):621-41. doi: 10.1128/MMBR.00027-10. PubMed PMID: 21119020; PubMed Central PMCID: PMC3008168.
51. Ramirez-Diaz MI, Diaz-Magana A, Meza-Carmen V, Johnstone L, Cervantes C, Rensing C. Nucleotide sequence of *Pseudomonas aeruginosa* conjugative plasmid pUM505 containing virulence and heavy-metal resistance genes. *Plasmid.* 2011;66(1):7-18. Epub 2011/03/23. doi: 10.1016/j.plasmid.2011.03.002. PubMed PMID: 21421005.
52. Subedi D, Kohli GS, Vijay AK, Willcox M, Rice SA. Accessory genome of the multi-drug resistant ocular isolate of *Pseudomonas aeruginosa* PA34. *PloS one.* 2019;14(4):e0215038. Epub 2019/04/16. doi: 10.1371/journal.pone.0215038. PubMed PMID: 30986237; PubMed Central PMCID: PMC6464166.
53. Fonseca EL, Dos Santos Freitas F, Vieira VV, Vicente AC. New *qnr* gene cassettes associated with superintegron repeats in *Vibrio cholerae* O1. *Emerging infectious diseases.* 2008;14(7):1129-31. Epub 2008/07/05. doi: 10.3201/eid1407.080132. PubMed PMID: 18598639; PubMed Central PMCID: PMC2600354.
54. Belotti PT, Thabet L, Laffargue A, Andre C, Coulange-Mayonnove L, Arpin C, et al. Description of an original integron encompassing *blaVIM-2*, *qnrVC1* and genes encoding bacterial group II intron proteins in *Pseudomonas aeruginosa*. *The Journal of antimicrobial*

- chemotherapy. 2015;70(8):2237-40. Epub 2015/05/16. doi: 10.1093/jac/dkv103. PubMed PMID: 25977399.
55. Lin J, Chen DQ, Hong J, Huang H, Xu X. Prevalence of qnrVC Genes in *Pseudomonas aeruginosa* Clinical Isolates from Guangdong, China. *Current microbiology*. 2020. Epub 2020/04/05. doi: 10.1007/s00284-020-01974-9. PubMed PMID: 32246181.
56. He P, Moran GR. Structural and mechanistic comparisons of the metal-binding members of the vicinal oxygen chelate (VOC) superfamily. *J Inorg Biochem*. 2011;105(10):1259-72. Epub 2011/08/09. doi: 10.1016/j.jinorgbio.2011.06.006. PubMed PMID: 21820381.
57. Stokes HW, Hall RM. A novel family of potentially mobile DNA elements encoding site-specific gene-integration functions: integrons. *Molecular microbiology*. 1989;3(12):1669-83. PubMed PMID: 2560119.
58. Poirel L, Liard A, Rodriguez-Martinez JM, Nordmann P. *Vibrionaceae* as a possible source of Qnr-like quinolone resistance determinants. *The Journal of antimicrobial chemotherapy*. 2005;56(6):1118-21. Epub 2005/10/18. doi: 10.1093/jac/dki371. PubMed PMID: 16227349.
59. Keay L, Edwards K, Naduvilath T, Taylor HR, Snibson GR, Forde K, et al. Microbial keratitis predisposing factors and morbidity. *Ophthalmology*. 2006;113(1):109-16. Epub 2005/12/20. doi: 10.1016/j.ophtha.2005.08.013. PubMed PMID: 16360210.
60. Stapleton F, Keay L, Edwards K, Naduvilath T, Dart JK, Brian G, et al. The incidence of contact lens-related microbial keratitis in Australia. *Ophthalmology*. 2008;115(10):1655-62. Epub 2008/06/10. doi: 10.1016/j.ophtha.2008.04.002. PubMed PMID: 18538404.
61. Dötsch A, Klawonn F, Jarek M, Scharfe M, Blöcker H, Häussler S. Evolutionary conservation of essential and highly expressed genes in *Pseudomonas aeruginosa*. *BMC genomics*. 2010;11:234. Epub 2010/04/13. doi: 10.1186/1471-2164-11-234. PubMed PMID: 20380691; PubMed Central PMCID: PMC2864246.
62. Jacoby GA, Strahilevitz J, Hooper DC. Plasmid-mediated quinolone resistance. *Microbiology spectrum*. 2014;2(5):10.1128/microbiolspec.PLAS-0006-2013. Epub 2014/10/24. doi: 10.1128/microbiolspec.PLAS-0006-2013. PubMed PMID: 25584197.
63. Nordmann P, Poirel L. Emergence of plasmid-mediated resistance to quinolones in *Enterobacteriaceae*. *Journal of Antimicrobial Chemotherapy*. 2005;56(3):463-9. doi: 10.1093/jac/dki245.
64. Tran JH, Jacoby GA. Mechanism of plasmid-mediated quinolone resistance. *Proceedings of the National Academy of Sciences*. 2002;99(8):5638. doi: 10.1073/pnas.082092899.
65. Fonseca EL, Vicente ACP. Epidemiology of qnrVC alleles and emergence out of the *Vibrionaceae* family. *Journal of medical microbiology*. 2013;62(Pt 10):1628-30. Epub 2013/06/27. doi: 10.1099/jmm.0.062661-0. PubMed PMID: 23800600.
66. Subedi D, Vijay AK, Kohli GS, Rice SA, Willcox M. Nucleotide sequence analysis of NPS-1 beta-lactamase and a novel integron (In1427)-carrying transposon in an MDR *Pseudomonas aeruginosa* keratitis strain. *The Journal of antimicrobial chemotherapy*. 2018. Epub 2018/03/17. doi: 10.1093/jac/dky073. PubMed PMID: 29547943.
67. Heidary Z, Bandani E, Eftekhary M, Jafari AA. Virulence genes profile of multidrug resistant *Pseudomonas aeruginosa* isolated from Iranian children with UTIs. *Acta medica Iranica*. 2016;54(3):201-10. Epub 2016/04/25. PubMed PMID: 27107526.
68. Cho HH, Kwon KC, Kim S, Koo SH. Correlation between virulence genotype and fluoroquinolone resistance in carbapenem-resistant *Pseudomonas aeruginosa*. *Annals of laboratory medicine*. 2014;34(4):286-92. Epub 2014/07/02. doi: 10.3343/alm.2014.34.4.286. PubMed PMID: 24982833; PubMed Central PMCID: PMC4071185.

69. Georgescu M, Gheorghe I, Curutiu C, Lazar V, Bleotu C, Chifiriuc MC. Virulence and resistance features of *Pseudomonas aeruginosa* strains isolated from chronic leg ulcers. BMC infectious diseases. 2016;16 Suppl 1:92. Epub 2016/05/14. doi: 10.1186/s12879-016-1396-3. PubMed PMID: 27169367; PubMed Central PMCID: PMC4890939.
70. Sawa T, Shimizu M, Moriyama K, Wiener-Kronish JP. Association between *Pseudomonas aeruginosa* type III secretion, antibiotic resistance, and clinical outcome: a review. Critical care (London, England). 2014;18(6):668. Epub 2015/02/13. doi: 10.1186/s13054-014-0668-9. PubMed PMID: 25672496; PubMed Central PMCID: PMC4331484.
71. Tran QT, Nawaz MS, Deck J, Foley S, Nguyen K, Cerniglia CE. Detection of type III secretion system virulence and mutations in *gyrA* and *parC* genes among quinolone-resistant strains of *Pseudomonas aeruginosa* isolated from imported shrimp. Foodborne pathogens and disease. 2011;8(3):451-3. Epub 2010/12/02. doi: 10.1089/fpd.2010.0687. PubMed PMID: 21117986.
72. Agnello M, Finkel SE, Wong-Beringer A. Fitness cost of fluoroquinolone resistance in clinical isolates of *Pseudomonas aeruginosa* differs by type III secretion genotype. Frontiers in microbiology. 2016;7:1591. Epub 2016/10/21. doi: 10.3389/fmicb.2016.01591. PubMed PMID: 27757111; PubMed Central PMCID: PMC4890939.
73. Garey KW, Vo QP, Larocco MT, Gentry LO, Tam VH. Prevalence of type III secretion protein exoenzymes and antimicrobial susceptibility patterns from bloodstream isolates of patients with *Pseudomonas aeruginosa* bacteremia. Journal of chemotherapy (Florence, Italy). 2008;20(6):714-20. doi: 10.1179/joc.2008.20.6.714. PubMed PMID: 19129069.
74. Choy MH, Stapleton F, Willcox MD, Zhu H. Comparison of virulence factors in *Pseudomonas aeruginosa* strains isolated from contact lens- and non-contact lens-related keratitis. Journal of medical microbiology. 2008;57(Pt 12):1539-46. doi: 10.1099/jmm.0.2008/003723-0. PubMed PMID: 19018027.
75. Finck-Barbancon V, Goranson J, Zhu L, Sawa T, Wiener-Kronish JP, Fleiszig SM, et al. ExoU expression by *Pseudomonas aeruginosa* correlates with acute cytotoxicity and epithelial injury. Molecular microbiology. 1997;25(3):547-57. PubMed PMID: 9302017.
76. Agnello M, Wong-Beringer A. Differentiation in quinolone resistance by virulence genotype in *Pseudomonas aeruginosa*. PloS one. 2012;7(8):e42973. doi: 10.1371/journal.pone.0042973. PubMed PMID: 22905192; PubMed Central PMCID: PMC3414457.
77. Stewart RM, Wiehlmann L, Ashelford KE, Preston SJ, Frimmersdorf E, Campbell BJ, et al. Genetic characterization indicates that a specific subpopulation of *Pseudomonas aeruginosa* is associated with keratitis infections. Journal of clinical microbiology. 2011;49(3):993-1003. doi: 10.1128/JCM.02036-10. PubMed PMID: 21227987; PubMed Central PMCID: PMC3067716.