Protection of grapevine pruning wounds against *Phaeomoniella chlamydospora* and *Diplodia seriata* by biological and chemical methods

María del Pilar Martínez-Diz^{a,b}, Emilia Díaz-Losada^a, Ángela Díaz-Fernández^a, Yolanda Bouzas-Cid^a, David Gramaje^{c,*}

^aEstación de Viticultura e Enología de Galicia (AGACAL-EVEGA), Ponte San Clodio s/n 32428-Leiro-Ourense, Spain.

^bUniversidade da Coruña, Facultade de Ciencias, Zapateira, 15071 A Coruña, Spain.

^cInstituto de Ciencias de la Vid y del Vino (ICVV), Consejo Superior de Investigaciones Científicas - Universidad de la Rioja - Gobierno de La Rioja, Ctra. LO-20 Salida 13, Finca La Grajera, 26071 Logroño, Spain.

* Corresponding author. Instituto de Ciencias de la Vid y del Vino (ICVV), Consejo Superior de Investigaciones Científicas, Universidad de la Rioja, Gobierno de La Rioja, Ctra. LO-20 Salida 13, 26007 Logroño, Spain.

E-mail address: david.gramaje@icvv.es (D. Gramaje)

ABSTRACT

The grapevine trunk diseases (GTDs) Botryosphaeria dieback and esca threaten the sustainability of the grapevine industry worldwide. This study aimed to evaluate and compare the efficacy of various liquid (pyraclostrobin + boscalid and thiophanate methyl) and paste (paste + tebuconazole) formulation fungicide treatments, and biological control agents (*Trichoderma atroviride* SC1 and *T. atroviride* I-1237), for their potential to prevent infection of grapevine pruning wounds by *Diplodia seriata* and *Phaeomoniella chlamydospora* in two field trials over two growing seasons. Treatments were applied to freshly pruned wounds following their label dosages recommendations. After 24 hours, wounds were artificially inoculated with 400 spores of *D. seriata* or 800 spores of *P. chlamydospora*. Isolations were made from the treated pruning wounds after 12 months to evaluate the efficacy of the treatments. Fungicide formulations were superior to *Trichoderma*-based treatments for the control of both pathogens during both growing seasons, with mean percent disease control of 44 to 95% for *D. seriata* and 46 to 67% for *P. chlamydospora*. Pyraclostrobin + boscalid was the most effective treatment. *Trichoderma atroviride*-based treatments did not reduce infection by *D. seriata* or *P. chlamydospora* compared to the untreated inoculated control in both vineyards and seasons. This study represents the first vineyard assessment of several chemical and biological treatments to protect pruning wounds against GTDs fungi in Europe and provides growers with tangible preventative control practices to minimize yield losses due to GTDs.

Keywords: Botryosphaeria, chemical control, esca, *Trichoderma*, *Vitis vinifera* L.

1. Introduction

Botryosphaeria dieback and esca are two of the most harmful grapevine trunk diseases (GTDs) affecting vineyards in all major grape-producing areas worldwide. They currently are among the main biotic threats to the economic sustainability of viticulture reducing yields, productivity and longevity of vines and vineyards (Gramaje et al., 2018). Yield losses of 30-50% have been reported by Botryosphaeria dieback in highly infected vineyards of North America (Milholland, 1991). The economic impact of Botryosphaeria dieback along with another GTD such as Eutypa dieback in California was estimated to be \$USD260 million per year (Siebert, 2001). Esca incidence has reached up to 80% in several vineyards of Southern Italy (Romanazzi et al., 2009), and 12% of vineyards in France are currently no longer economically viable, due mainly to esca, with an annual estimated loss of €1 billion (Lorch, 2014).

Botryosphaeria dieback is currently associated with 26 botryosphaeriaceous taxa in the genera *Botryosphaeria*, *Diplodia*, *Dothiorella*, *Lasiodiplodia*, *Neofusicoccum*, *Neoscytalidium*, *Phaeobotryosphaeria*, and *Spencermartinsia* (Úrbez-Torres, 2011; Pitt et al., 2013a, 2013b, 2015; Rolshausen et al., 2013; Yang et al., 2017) with the species *Diplodia seriata* being one of the most frequently isolated fungi from diseased vines in several grape growing regions such as Australia (Savocchia et al., 2007), California (Úrbez-Torres et al., 2010), Chile (Auger et al., 2004), China (Yan et al., 2013), France (Larignon et al., 2001), Mexico (Úrbez-Torres et al., 2008), Portugal (Phillips, 2002), South Africa (van Niekerk et al., 2004) and Spain (Luque et al., 2014). Botryosphaeria dieback frequently shows as complete absence of spring growth from affected spurs due to necrosis formation in wood vascular tissues with bud-break failure, and shoot and trunk dieback (Úrbez-Torres, 2011). Wood symptoms are characterized by wedge-shaped perennial cankers and dark streaking in spurs, cordons and trunks vascular tissues usually beginning in pruning wounds (Úrbez-Torres et al., 2010).

Esca is mainly caused by the fungus *Phaeomoniella chlamydospora* along with *Phaeoacremonium minimum* and other *Phaeoacremonium* spp. (Gramaje et al., 2015), some *Cadophora* spp. (Travadon et al., 2015), and several basidiomycetous taxa belonging to genera *Inocutis*, *Inonotus*, *Fomitiporella*, *Fomitiporia*, *Phellinus*, and *Stereum* (Cloete et al., 2015). The most characteristic external symptoms of the chronic esca comprise multiple banding discolourations on leaves known as ‘tiger-stripe’ pattern (Surico, 2009; Gubler et al., 2015). Internal wood symptoms involve black spots in the xylem vessels, longitudinal brown to black vascular streaking, and white to light yellow soft rot that frequently develops in wood of older vines (Fischer, 2002; Lecomte et al., 2012). Apoplectic esca form is characterized by a sudden and unexpected wilting of the whole vine or one/several arms or shoots (Lecomte et al., 2012).

Infection of grapevines by GTD fungal pathogens primarily occurs through annual pruning wounds made during the dormant season (Gramaje et al., 2018). Pycnidia of Botryosphaeriaceae spp. and *P. chlamydospora* develop from dead/cankered wood, old pruning wounds, grapevine canes, crevices, cracks and on the bark of infected grapevines (Úrbez-Torres and Gubler, 2011; Baloyi et al., 2016), and in the case of *P. chlamydospora*, mycelium on infected wood can also be a source of conidia (Edwards and Pascoe, 2001; Edwards et al., 2001; Baloyi et al., 2016). Fruiting bodies of these fungi can also be found in pruning debris left in the vineyard, thus becoming a potential inoculum source for new infections (van Niekerk et al., 2010; Úrbez-Torres, 2011; Elena and Luque, 2016b).

Conidia release of Botryosphaeriaceae spp. and *P. chlamydospora* has been shown to be primarily correlated with rain events (Larignon and Dubos, 2000; Eskalen and Gubler, 2001; Kuntzmann et al., 2009; van Niekerk et al., 2010; Úrbez-Torres et al., 2010; Valencia et al., 2015). The dynamics of *P. chlamydospora* dispersal in Spain were recently described by an epidemiological equation that integrated the effects of both rain and temperature (González-Domínguez et al., 2020). Conidia of Botryosphaeriaceae spp. has been shown to be primarily

dispersed by rain splash (Úrbez-Torres et al., 2010), while inoculum of *P. chlamydospora* is predominantly aerielly dispersed (Larignon and Dubos, 2000; Eskalen and Gubler, 2001; Gubler et al., 2015; Quaglia et al., 2009). Infection occurs when conidia land on exposed and susceptible pruning wounds, germinate in xylem vessels and colonize the vine spur, cordon and trunk (Mostert et al., 2006; Epstein et al., 2008; Gubler et al., 2013; Moyo et al., 2014).

Susceptibility of pruning wounds to GTD pathogens is mainly dependent on the time of pruning, and the period between pruning and possible infection case. Several studies using artificial spore inoculations showed that susceptibility of grapevine pruning wounds is high when fungal infection occurs at the moment of pruning but decreases as the period between pruning and infection increases up to several weeks or months (Petzold et al., 1981; Munkvold and Marois, 1995; Eskalen et al., 2007; Serra et al., 2008; Úrbez-Torres and Gubler, 2011), with seasonal variation reported between grape regions caused primarily by climatic differences (Gramaje et al., 2018).

Protection of pruning wounds is essential for the management of *Botryosphaeria dieback* and *esca* in grapevine, especially if adopted early in the vineyard lifespan (Kaplan et al., 2016; Sosnowski and McCarthy, 2017). The efficacy of fungicide wound treatments against *Botryosphaeriaceae* spp. and *P. chlamydospora* has been demonstrated in Australia (Pitt et al., 2012), California (Rolshausen et al., 2010), Chile (Díaz and Latorre, 2013), New Zealand (Amponsah et al., 2012; Sosnowski and Mundi, 2019) and South Africa (Mutawila et al., 2015). The use of physical barriers such as paints and pastes formulated with or without fungicides have also shown to be effective to control infections caused by *Botryosphaeriaceae* fungi and *P. chlamydospora* (Epstein et al., 2008; Rolshausen et al., 2010; Pitt et al., 2012; Díaz and Latorre, 2013).

The high restrictions that most effective chemical active ingredients are currently facing in Europe because of environmental and human health risks (Larignon et al., 2008; Spinosi et al., 2009), make indispensable address new alternatives for controlling GTDs. Over the last years, research on biological control of GTD fungi with antagonistic microorganisms has shown promising results primarily under controlled conditions (Alfonzo et al., 2009; Mutawila et al., 2011a; Haidar et al., 2016; Rezagui et al., 2016; Álvarez-Pérez et al., 2017; Dargaignes et al., 2018; Mondello et al., 2018; Andreolli et al., 2019; Del Frari et al., 2019; Mondello et al., 2019; Trotel-Aziz et al., 2019; Niem et al., 2020). Field trials with biological control agents (BCAs) have shown variable results for preventing infection by *Botryosphaeriaceae* and *esca* fungi (Kotze et al., 2011; Mutawila et al., 2011b, 2015, 2016; Mounier et al., 2014; Reis et al., 2017; Martínez-Diz et al., 2020a).

To our knowledge, no comparative studies to evaluate the efficacy of chemical and BCA products as pruning wound protectants against GTD fungi have been performed in Europe so far. Four pruning wound treatments are currently registered in Spain for the control of GTD fungi: three *Trichoderma*-based biological products, namely Esquive, Blindar and Vintec, and Tessior, a liquid polymer containing boscalid and pyraclostrobin (MAPA, 2020). In addition, thiophanate methyl is registered in Spain against fungal trunk pathogens in almond (MAPA, 2020). The aim of this study was to evaluate and compare the efficacy of various liquid and paste formulation fungicide treatments, and BCAs, for their potential to prevent infection of grapevine pruning wounds by *D. seriata* and *P. chlamydospora* in field trials. The products assessed were those registered in Spain for control of fungal trunk pathogens or other diseases on grapevine and/or other hosts.

2. Materials and methods

2.1 Location and characteristics of the experimental vineyards

The assays were carried out at two commercial vineyards located in O Barco de Valdeorras, Galicia region (Spain), in 2018 and 2019. The vineyards were planted on 1981 (37-years-old) and 1989 (29-years-old) with ‘Godello’ cultivar grafted onto 110 Richter rootstock. Vines were spaced 120 cm from center to center, and with an interrow spacing of 225 cm, trained as bilateral cordons in a trellis system with a spur-pruning (Royat).

Vineyards were less than 500 m apart and had very similar climates. Standard cultural practices were used in both vineyards during the growing season, and the management of powdery and downy mildews was performed using only wettable sulphur and copper compounds applied at label dosages and following Integrated Pest Management (IPM) guidelines, respectively, when required. At the beginning of the study (2018), about 8% and 12% of vines had shown GTDs symptoms in each vineyard, respectively. The presence and evolution of GTDs symptoms have been inspected biannually from 2014 to present in plots of 1,500 vines at both vineyards. GTDs symptoms detected during inspection were associated mainly with esca such as tiger-pattern foliar necrosis, and shoots, arm and/or cordon death.

Both vineyards were located less than 4 km to an automatic weather station owned by Meteogalicia (Weather Service of Galician Regional Government, Xunta de Galicia) and its climatic data was considered to be representative.

2.2 Fungal isolates and inoculum preparation

Diplodia seriata isolate CJL-398 and *Phaeomoniella chlamydospora* isolate BV-130 were used. *P. chlamydospora* BV-130 was selected due to its high virulence on grapevine in previous assays (Martínez-Diz et al., 2019). This strain was isolated from a 43-year-old esca diseased vine cultivar ‘Tempranillo’ grafted onto ‘41 Berlandieri’ rootstock in 2015. *D. seriata* JL-398 was the most virulent isolate among 14 in a detached grapevine cane assay (Elena et al., 2015a). This strain was isolated from cankers and wood necrosis of grapevine.

Conidial suspensions of each pathogen were used for artificial inoculations in the field and inoculum was obtained using methods similar to those described by Elena and Luque (2016a). In the case of *D. seriata*, a mycelial plug previously plated on Potato Dextrose Agar (PDA, Conda Laboratories, Spain) at 25°C for 7 days was cultured upside down over the center of a water agar (WA, Conda Laboratories, Spain) plate. Maritime pine (*Pinus pinaster* L.) needles were cut to 1 cm long fragments and then sterilized in an autoclave following the standard protocol of 121°C for 20 min. Then, approximately 20 sterile needles fragments were placed on the WA media surface surrounding the *D. seriata* mycelial plug at about 1 to 1.5 cm and plates were incubated under warm white fluorescent and near ultraviolet light for a 12-h photoperiod regime at 25°C for 4 weeks until pycnidia formation. The day before inoculation, pine needles fragments (about n=40) with *D. seriata* pycnidia were placed along with 30 ml of sterile distilled water (SDW) in a beaker. The solution was kept overnight (about 16 h) at 4°C in permanent agitation with the aid of a magnetic stirrer to induce conidia release from the pycnidia and prevent conidia germination. The inoculation day, the resulting solution was vacuum-filtered through a 60-mm Steriflip filter (Millipore Corporation, Billerica, MA) to get a cleaner suspension. Then, conidial suspension was adjusted to 2×10^4 conidia mL⁻¹ using a hemocytometer (Brand™ Blaubrand™ Neubauer Counting Chamber, Thermo Fisher Scientific Inc., MA, USA).

Phaeomoniella chlamydospora strain was grown on PDA plates at 25°C for 3 weeks. Same day of inoculation, conidia were released from cultures by adding 10 ml of SDW and gently scraping with a sterile stick and the collected suspension adjusted to a concentration of 4×10^4 conidia mL⁻¹ based on counts from the hemocytometer. Both conidial suspensions were stored at 4°C until inoculation time to avoid early conidia germination.

Spore germination was assessed for both fungal trunk pathogens by placing four drops of the spore suspension on a PDA plate, which was then incubated at 25°C under fluorescent light for a 12-h photoperiod. After approximately 24 h, a glass cover slip was placed over each drop area on the PDA. The number of non-germinated spores over a total of 100 in each drop was counted using an optical microscope (Nikon Eclipse E400) at 100x magnification. The mean percentage of germinated spores was determined.

2.3 Pruning wound protection treatments

Wound protection treatments tested in the present assay are listed in Table 1. We evaluated the efficacy of two chemical and two BCA formulated products, and also a paste mixed with a fungicide. In general, the chemical and biological products assessed were those commercially formulated and currently registered and available in Spain for control of fungal trunk pathogens, except tebuconazole (Song), which is registered to control botrytis bunch rot (*Botrytis cinerea*) and powdery mildew (*Erysiphe necator*) in grapevine. Applications rates were selected based on the registered label dosages recommendations. Liquid formulations were prepared by the suspension of the products in tap water, which is the procedure normally used for spraying vineyard treatments in Galicia region. Pyraclostrobin + boscalid (Tessior) treatment contains a liquid polymer and it is already formulated to be directly sprayed to pruning wounds without any previous mixing. Paste treatment was prepared by mixing a liter of the paste formulation (Master) with 80 ml of tebuconazole (Song).

Regarding BCA treatments, the conidia viability of both *Trichoderma atroviride* strains (SC1 and I-1237) in the commercial products was tested to be at a minimum of 85% before the assay was set up (Pertot et al., 2016). A serial dilution of the conidia suspension was plated on PDA and the colony-forming units were counted after 24-48 h incubation at room temperature.

2.4 Field assay and experimental design

On 19 February 2018, 1-year-old canes of all vines to be treated were spur-pruned to three buds using secateurs in both vineyards, coinciding with the common pruning time in this region of Spain. Wounds treatments were applied by hand until runoff within 2 h after pruning to three wounds per vine. Liquid formulations were applied using a 500 ml hand-held spray bottle with a plastic shield on the nozzle to minimise spray drift and the paste formulation were applied with the aid of a paintbrush. Untreated controls, positive (artificially inoculated, IC) and negative (non-artificially inoculated, NC) were mock treated with sterile distilled water (SDW).

On the following day, wounds were moistened by spraying with SDW immediately prior to inoculation with the fungal trunk pathogens and a drop of Tween 20 (Sigma-Aldrich, San Luis, MO, USA) was added to each conidial suspension as a surfactant to assist spreading the spores over the pruning wound surface (Sosnowski and Mundi, 2019). Approximately 400 and 800 conidia of *D. seriata* and *P. chlamydospora*, respectively, suspended in a drop of 20 µl of SDW were then applied per wound using a micropipette. All pruning wounds were inoculated with the pathogen inoculum except NC controls, which were mock inoculated with a drop of 20 µl of SDW alone instead and being exposed to natural infection. Inoculum drops placed onto the pruning wounds were left to air dry (from some minutes to 1 h) before being wrapped with Parafilm M (Pechiney Plastic Packaging, Chicago, IL, USA) to avoid fast dehydration and favour fungal spores' penetration into xylem vessels. Due care was taken to avoid the rain for the entire duration of the trials set up, namely pruning, wound treatments application and artificial fungal inoculation (2 days).

The experiment was set up as a randomized block design with three replicates of ten plants (thirty canes) per wound protectant treatment and pathogen in each vineyard. Three replicates

of ten plants per pathogen were also used for IC in each vineyard. Additionally, three replicates of ten plants were used as NC in each vineyard. The experiment was repeated the following season (2019–20), with pruning and wound treatments applied on 12 February 2019, and artificial fungal inoculations on 13 February 2019.

2.5 Fungal recovery and identification

Canes were harvested from vines above the second bud (about 10 cm long pieces) approximately 12 months after artificial inoculation and stored in a 4°C cool room prior to laboratory assessment. Bark was first removed using a sharp knife from each cane. Then, canes were surface sterilised for 1 min in 33% sodium hypochlorite (commercial 40 g Cl/l) and rinsed twice for 1 min each in SDW. After air drying on sterile filter paper to remove moisture excess, each cane was cut into small pieces (about 12 mm²) taken from the margin between discoloured or dead and live or apparently healthy wood tissue using sterilised secateurs. Five wood fragments were plated onto each of two plates of Malt Extract Agar (MEA) amended with 0.35 g l⁻¹ of streptomycin sulphate (Sigma-Aldrich, St. Louis, MO, USA) (MEAS) giving a total of ten wood pieces per cane. Cultures were incubated at 25°C under warm fluorescent light for a 12-h photoperiod and inspected daily for 15 days. All growing fungal colonies were transferred to PDA plates and then assessed for the presence or absence fungal mycelial growth resembling *D. seriata*, *P. chlamydospora* or *Trichoderma* spp.

Identification of GTD fungal cultures was then assessed under a stereoscopic (Olympus SZX9, Olympus Corporation, Tokyo, Japan) and optical microscopes (Nikon Eclipse E400, Nikon Corporation, Tokyo, Japan) based on cultural and morphological features previously described including colony growth pattern, colour, mycelial and other characteristics such as conidial shape, size and colour (Crous and Gams, 2000; Phillips et al., 2007). Identity of GTD fungal isolates and *Trichoderma* spp. was confirmed by molecular methods. Fungal DNA was extracted from fresh mycelium after 3 weeks of incubation in PDA using the E.Z.N.A. Plant Miniprep Kit (Omega Bio-Tek, Doraville, GA, USA) following manufacturer's instructions. *D. seriata* was confirmed by sequencing part of the translation elongation factor 1- using the primer pairs EF1F-EF2R (Jacobs et al., 2004). *P. chlamydospora* was detected by PCR using the primers Pch1-Pch2 (Tegli et al., 2000). Identity of *Trichoderma* spp. was confirmed at species level by sequencing the ITS region using the universal primers ITS1F/ITS4 (Gardes and Bruns, 1993). All PCR products were visualized in 1% agarose gels (agarose D-1 Low EEO, Conda Laboratories) and sequenced in both direction by Eurofins GATC Biotech (Cologne, Germany).

2.6 Data analysis

Efficacy of each wound treatment was calculated as mean percentage recovery (MPR) of *D. seriata* and *P. chlamydospora* from each cane per treatment (Sosnowski et al., 2008, 2013). Data were checked for normality and homogeneity of variances prior to statistical analyses and transformed when required into the arcsine of the square root of the proportion (MPR/100)^{1/2}. The statistical analysis of the experimental results was carried out in a two-way ANOVA with blocks and treatments as independent variables, and MPR (%) as dependent variable. Mean percentage disease control (MPDC) was also determined as the reduction in MPR (%) as a proportion of the artificially inoculated control (IC) (MPDC=100 × [1 – (MPR treatment/MPR IC)]) (Sosnowski et al., 2008, 2013). Means were compared with ICs by the Student's *t* least significant difference (LSD) at *P* < 0.05. Data from all experiments were analysed using the Statistix 10 software (Analytical Software).

3. Results

3.1 Wound treatment evaluation against *Diplodia seriata*

During 2018-19 and 2019-20 seasons, *D. seriata* spore germination on PDA was 94% and 98.5%, respectively, and it was recovered from 58 and 68% of IC wounds, respectively (Table 2). *D. seriata* was recovered from 1% of NCs wounds at both seasons. Analysis of variance showed that there were significant differences in the relative recovery data from the different treatments between seasons ($P<0.05$). No significant differences were found in the recovery data between vineyards in each season (2018-19, $P=0.904$; 2019-20, $P=0.593$), so data from each vineyard were combined and the analysis was performed separately for each season (Table 2).

Treatment with pyraclostrobin + boscalid, thiophanate methyl, and the paste + tebuconazole significantly reduced the MPR of *D. seriata* from pruning wounds with respect to the IC at both seasons ($P<0.05$) (Table 2). During 2018-19 season, pyraclostrobin + boscalid, thiophanate methyl, and the paste + tebuconazole provided MPDC of 95, 90 and 76%, respectively, whereas these products provided MPDC of 69, 54 and 44%, respectively, during 2019-20 season. During both seasons, there was not a significant treatment effect with *Trichoderma*-based wound protectants ($P>0.05$). During 2018-19 season, *T. atroviride* SC1 and *T. atroviride* I-1237 provided MPDC of 10, and 26%, respectively, while these products provided MPDC of 22 and 32%, respectively, during 2019-20 season.

3.2 Wound treatment evaluation against *Phaeomoniella chlamydospora*

During 2018-19 and 2019-20 seasons, *P. chlamydospora* spore germination on PDA was 89% and 93%, respectively, and it was recovered from 27 and 42% of IC wounds, respectively (Table 3). *P. chlamydospora* was recovered from 0% and 3% of NCs wounds during 2018-19 and 2019-20 seasons, respectively. There were no significant differences in the relative recovery data from the different treatments between vineyards ($P=0.500$) and seasons ($P=0.080$), so data were combined for analysis.

There was a significant treatment effect ($P<0.05$) with the paste + tebuconazole and pyraclostrobin + boscalid treatments reducing MPR of *P. chlamydospora* to 12 and 18% compared with 36% from the IC wounds (MPDC of 67 and 51%; Table 3). There was not a significant treatment effect with thiophanate methyl and *Trichoderma*-based wound protectants ($P>0.05$). Thiophanate methyl provided MPDC of 46%, whereas *T. atroviride* SC1 and *T. atroviride* I-1237 provided MPDC of 0 and 17%, respectively.

3.3 *Trichoderma*-based treatments colonization

The conidia viability was on average of 97% and 95% for *T. atroviride* SC1 during 2018-19 and 2019-20 seasons, respectively. Regarding *T. atroviride* I-1237, the conidia viability was 94% during 2018-19 season and 96% during 2019-20 season. *Trichoderma* spp. were exclusively recovered from pruning wounds treated with *Trichoderma*-based formulations at varying levels. There were no significant differences in the relative recovery data between vineyards ($P=0.180$) and seasons ($P=0.075$). During 2018-19, recovery percentages were 5 and 10% for *T. atroviride* SC1 and *T. atroviride* I-1237, respectively. During 2019-20, recovery percentages were 9% and 14% for *T. atroviride* SC1 and *T. atroviride* I-1237, respectively.

3.4 Weather data

During 2018-19 season, the average of the daily mean temperature and relative humidity in the week from the day of pruning and wound treatments application (from 19 to 25 February 2018) was 6.5°C and 67.5%, respectively, with no rain events in that period. The average of daily mean temperature, daily mean relative humidity and accumulated rainfall for the whole month of February 2018 was 5.8°C, 76.6% and 84.2 mm, respectively, with nine rain events (of >1 mm) in total.

During 2019-20 season, the week from the day of pruning and application of wound treatments (from 12 to 18 February 2019) registered an average of the daily mean temperature of 8.2°C and a 71.1% on average of daily relative humidity. For the same period, there was only one rain event (18 February 2019) with a total rainfall of 10.6 mm. Regarding the whole February 2019 month, the average of the daily temperature was 8.2°C and of the daily relative humidity 73.3%. The total rainfall in the same month was 37 mm received in a total of four rain events.

4. Discussion

The present study represents the first vineyard comparison of the efficacy of paste and liquid fungicides, and BCA treatments to protect pruning wounds against GTDs fungi in Europe. Considering the high incidence of GTDs, particularly esca, and the restrictions on the use of chemicals in Europe (Mondello et al., 2018), this study provides growers with tangible preventative control practices to minimize yield losses due to GTDs. By focussing on products already registered for control of trunk diseases in almond or foliar diseases of grapevines in Spain, the lower cost of label extension compared to new product registration will increase the likelihood and success of registration for GTDs. *D. seriata* was chosen to represent *Botryosphaeria dieback*, because it is one the most common cited *Botryosphaeriaceae* species occurring on grapevines worldwide and is reported to be a virulent species in Spain (Luque et al., 2009; Elena et al., 2015b). *P. chlamydospora* was chosen to represent esca, because it is the most frequently isolated species from affected vines in most grape growing regions worldwide (Berstch et al., 2013; Gubler et al., 2015).

Our results demonstrate that paste and liquid fungicide formulations were superior to *Trichoderma*-based treatments for the control of *D. seriata* and *P. chlamydospora*. All paste and liquid fungicide treatments tested reduced recovery of both pathogens from inoculated wounds compared with the untreated inoculated control, with the exception of thiophanate methyl for *P. chlamydospora*. Similar results were observed in other studies where several fungicides and BCA treatments were compared as pruning wound protectants in the same field trial. In Australian vineyards, liquid and paste fungicide formulations were more effective than *Trichoderma*- and *Bacillus subtilis*-based formulations against *D. seriata* and *Diplodia mutila* (Pitt et al., 2012), and *Eutypa lata* (Ayres et al., 2017) infections, respectively. Halleen et al. (2010) also reported that fungicides were more effective than *Trichoderma* spp. against *E. lata* infection in field trials carried out in South Africa, in spite of the efficacy of *Trichoderma* treatments in reducing GTD fungal infection.

Application of pyraclostrobin + boscalid to pruning wounds provided high mean percentage of disease control (MPDC) for both pathogens. To date, only preliminary studies have been carried out in field trials in Germany (Kühn et al., 2017; Lengyel et al., 2019), Greece (Kühn et al., 2017; Samaras et al., 2019) and Spain (Kühn et al., 2017), where pyraclostrobin and boscalid (Tessior) was effective as pruning wound protectant reducing the grapevine wood infection caused by *Diplodia* spp. and *P. chlamydospora*. The application of a similar commercial product based on pyraclostrobin and boscalid without the liquid polymer (BASF516, BASF Australia Ltd, Sidney, New South Wales, Australia) showed a low efficacy against *E. lata* artificial pruning wound inoculations in Australian vineyards (Sosnowski et al., 2008). Wound applications of pyraclostrobin alone were effective for the control of *D. seriata*

and *P. chlamydospora* in Chile (Díaz and Latorre, 2013) and California (Rolshausen et al., 2010) vineyards. Moreover, this active ingredient significantly reduced infections caused by fungi associated with *Botryosphaeria dieback* (Rolshausen et al., 2010), *Eutypa dieback* (Sosnowski et al., 2008, 2013; Rolshausen et al., 2010; Ayres et al., 2017), and *esca* (Rolshausen et al., 2010), under field conditions.

The only treatment to provide a similar level of control than pyraclostrobin + boscalid for both pathogens was the paste with tebuconazole. Accordingly, applications of paste and liquid formulations containing tebuconazole on pruning wounds of ‘Cabernet Sauvignon’ vines significantly reduced the mean vascular discolouration length and the reisolation percentage of *D. seriata* and *P. chlamydospora* in Chilean vineyards (Díaz and Latorre, 2013). In Australia, a gel and a paint with tebuconazole applied by paintbrush to freshly pruned canes reduced *E. lata* infections to 100% and 94%, respectively (Sosnowski et al., 2013). Pitt et al. (2012) also demonstrated that a tebuconazole paste formulation provided a 38% control of *D. mutila* in a trial performed in Australia. Other physical barriers containing a paste with fungicides have resulted effective at reducing pruning wound infections by other GTD fungi (Rolshausen and Gubler, 2005; Sosnowski et al., 2008; Rolshausen et al., 2010; Pitt et al., 2012). Liquid spray applications of tebuconazole were also significantly effective reducing the recovery of *D. seriata* in Australia (Pitt et al., 2012).

Thiophanate methyl was effective in reducing infection by *D. seriata*, while no significant effect was observed against *P. chlamydospora*. Similar findings were reported by Rolshausen et al. (2010) in California, where pruning wounds applications of thiophanate methyl reached a disease control of 80% for *D. seriata* infections but did not perform as well against *P. chlamydospora* with only a 52% of disease control. In Chile, Díaz and Latorre (2013) reported the efficacy of both liquid and paste formulations of thiophanate methyl to control *D. seriata* and *P. chlamydospora* infections in pruning wounds. This chemical compound was also effective in reducing the pruning wound infections caused by *P. chlamydospora* and *Neofusicoccum luteum* in field trials carried out in South Africa (Mutawila et al., 2015) and New Zealand (Amponsah et al., 2012), respectively.

Pastes and paints are considered the most reliable protectants of pruning wounds against GTD fungi, especially when they are mixed with fungicides (Moller et al., 1977; Rolshausen and Gubler, 2005; Rolshausen et al., 2010; Sosnowski et al., 2008, 2013; Díaz and Latorre, 2013). They provide a physical barrier to protect pruning wounds from GTD fungal infection while the fungicide can also act on the pathogens if the physical barrier is compromised by rain, sap flow, or cracking when drying (Sosnowski et al., 2008). However, some other studies reported no differences in effectiveness between application of acrylic paint with or without fungicides (Sosnowski et al., 2008; Mayet and Lecomte, 2014). Pastes and paints are usually applied by hand with a paint brush, unless the product contains a liquid polymer to act as a physical barrier, which is the case of Tessior commercial product. It should be noted that application by hand is more time-consuming and can be at least two to four times the application cost with a tractor mounted sprayer (Sosnowski and McCarthy, 2017). Further research is therefore required to determine the protective mechanisms of each component and if their efficacy is also influenced by other factors such as wound size, application time, and weather variables.

Species of the fungal genus *Trichoderma* have been the most investigated BCA to act as pruning wound protectant against GTDs pathogens (John et al., 2005; Halleen et al., 2010; Kotze et al., 2011; Mutawila et al., 2015, 2016; Reis et al., 2017). Our results shown that *Trichoderma atroviride*-based treatments did not reduce infection by *D. seriata* or *P. chlamydospora* compared to the untreated inoculated control in both vineyards and seasons. This is the first report to assess the efficacy of *T. atroviride* SC1 in protecting grapevine pruning wounds from infection by GTD fungi in mature vineyards. In recent research, Berbegal et al.

(2020) applied *T. atroviride* SC1 to pruning wounds of 3-year-old vines but its efficacy as wound protectant against GTD pathogens was not tested in this specific plant part. In nurseries, *Trichoderma atroviride* SC1 showed high efficacy to reduce artificial (Pertot et al., 2016) or natural (Bebegali et al., 2020) *P. chlamydospora* infection when applied at different propagation stages. Preliminary results showed the efficacy of *T. atroviride* I-1237 to reduce the disease incidence and severity of *P. chlamydospora* and *N. parvum* on pruning wounds in Portuguese vineyards (Reis et al., 2017). Similarly, Mounier et al. (2014) demonstrated that spraying pruning wounds with *T. atroviride* I-1237 over two years significantly reduced the esca, and Botryosphaeria and Eutypa diebacks foliar symptoms expression, and the plant mortality rate due to GTDs in French vineyards. Dipping young grapevine plants in *T. atroviride* I-1237 during the nursery propagation process decreased *D. seriata* and *P. chlamydospora* DNA and necrotic lesion length compared to the untreated plants (Mounier et al., 2014). Other *Trichoderma* strains or *Trichoderma*-based commercial products have shown high efficacy in reducing the recovery of GTDs fungal pathogens from artificially inoculated pruning wounds under field conditions (John et al., 2005; Halleen et al., 2010; Kotze et al., 2011; Pitt et al., 2012; Mutawila et al., 2015).

These inconsistencies found in the *Trichoderma* products performance among our study and previous reports could be due to different reasons. Although *Trichoderma* spp. have the ability to provide long-term protection to pruning wounds and thus preventing fungal trunk pathogens infections, they firstly need to establish itself, grow and colonize wounds instead of a simply temporal establishment (Munkvold and Marois, 1993; Mutawila et al., 2011a, 2011b). In this study, pruning wound colonization by both strains of *T. atroviride* was very low at both vineyards and seasons ranging from 5 to 14%. Environmental conditions such as the temperature at the time of application might have a negative influence in the persistence and implementation of *Trichoderma* spp. (Elmer and Reglinski, 2006; Pertot et al., 2017). According to the label recommendations of each product, *T. atroviride* SC1 formulation should be applied when environmental temperature is equal or higher than 10°C for a minimum of five hours on the day of application in the field, while *T. atroviride* I-1237 formulation is supposed to be biologically active at temperatures above 5°C. The average of the daily mean temperature experienced in the week of the trials set up was 6.5°C and 8.2°C during 2018-19 and 2019-20 seasons, respectively, a fact that could explain the low colonization performed by *Trichoderma*-based formulations. In addition, a slightly higher *Trichoderma* recovery was registered during 2019-20 season (9.4 to 13.5%) than 2018-19 (4.8 to 9.8%) probably also explained by the warmer temperatures registered in this season.

Timing of pruning within the dormant season should be adjusted to periods with mild and favourable temperature values that might lead to a better implantation and development of the BCA on pruning wounds and thus increasing its effectiveness against GTD pathogens. *Trichoderma* spp. application after pruning in late winter or early spring would likely provide higher disease control than normal pruning in winter. However, late pruning is not feasible in all vineyards. In those vineyards with limited labour force, growers need to begin pruning early in the winter to ensure completion of the activity before bud break. An alternative would be to prune in late autumn or early winter. Recent research carried out in the same grape-growing region of the present study reported low abundances of GTDs pathogens infecting naturally pruning wounds after an early pruning made in November (Martínez-Diz et al., 2020b). The age and physiological state of the vine as well as the dose and product formulation have also been suggested as factors that could have an influence on effective colonization by *Trichoderma* spp. (Schubert et al., 2008; Halleen et al., 2010; Mutawila et al., 2016). However, further research is required to confirm these hypotheses.

Label instructions of most fungicide and BCA commercial formulations to protect pruning wounds recommend their application shortly after pruning to minimize the chances of GTDs

infection. In our study, we followed the official method of European countries to evaluate pruning wounds protection products against *E. lata*, which suggests carrying out fungal inoculations 24 hours after the application of preventive treatments (EPPO, 2017). Accordingly, in most of the previous pruning wound protection trials, the time elapsed between pruning wound protection and GTD fungal inoculation was 24 hours (John et al., 2005; Sosnowski et al., 2008, 2013; Halleen et al., 2010; Rolshausen et al., 2010; Kotze et al., 2011; Amponsah et al., 2012; Pitt et al., 2012; Díaz and Latorre, 2013; Ayres et al., 2017; Sosnowski and Mundi, 2019). This short time between BCA treatments application and artificial fungal inoculations could also have explained the poor performance exhibited by *Trichoderma*-based commercial formulations in our study. Previous research reported a greater biocontrol efficacy when artificial GTD pathogen infection was delayed 7 (Kotze et al., 2011; Mutawila et al., 2015) and 14 (Munkvold and Marois, 1993; John et al., 2005) days after application of *Trichoderma* spp. on pruning wounds. These findings suggest BCAs might need a period to colonise the pruning wound surface and grapevine wood to be effective, as reported by John et al. (2005).

The use of pathogen artificial inoculations is very common in the assessment of the efficacy of pruning wound protectants against GTDs to guarantee a substantial establishment of infection in untreated inoculated controls for statistical analysis (Halleen et al., 2010; Rolshausen et al., 2010; Sosnowski et al., 2008, 2013; Amponsah et al., 2012; Pitt et al., 2012; Ayres et al., 2017; Sosnowski and Mundi, 2019). In the present study, artificial inoculations with 400 (*D. seriata*) and 800 (*P. chlamydospora*) conidia were applied per wound to obtain optimal recovery percentages for robust evaluation of treatments according to doses recommendations made by Elena et al. (2015b). This fact represents a significantly higher ‘disease pressure’ than that which might be expected to occur under natural conditions. Wounds were infected naturally up to 1% by *D. seriata* and 3% by *P. chlamydospora*, in contrast with the artificially inoculated controls recovery with up to 68% and 42%, respectively. This indicates that wound protectants that showed lower efficacy rates in this study, such as BCA formulations, will most likely provide better control of both *D. seriata* and *P. chlamydospora* under ‘natural disease pressure’ in the vineyard. The efficacy of pruning wound protectants under lower artificial GTD inoculum levels or natural infections in the vineyard should be tested in future studies. Different affinities of *T. atroviride* strains for specific grapevine cultivars has been previously reported in South Africa (Mutawila et al., 2011b), and this should not be discarded as a possible cause of the low *Trichoderma* colonization rates obtained in this study.

To conclude, this study highlighted the efficacy of several fungicides with or without a physical barrier to protect grapevine pruning wounds against *D. seriata* and *P. chlamydospora* infections under field conditions. In particular pyraclostrobin + boscalid (Tessior), a registered product against GTD fungi in several countries in Europe, is recommended as pruning wound protectant to prevent infection by the most prevalent pathogens associated with Botryosphaeria dieback and esca. *Trichoderma*-based treatments showed lower efficacy against GTD fungi than that provided by fungicides and their performance seems to be related to environmental conditions and wound colonisation prior to infection by the pathogens. Good pruning practices along with strict sanitation procedures and pruning wound protection by the application of authorized products can significantly reduce the impact of GTD pathogens infections and thus increasing the lifespan of vineyards.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

This work has been funded by European funds in collaboration with the ‘Ministerio de Agricultura, Pesca y Alimentación (MAPA)’ (Spanish Government) and the ‘Xunta de Galicia’ (Regional Government) through the project ‘EVID: Innovative practices to combat grapevine diseases’ (FEADER 2017/003B). The authors acknowledge Bodegas Godeval for the provision of field sites and technical assistance for this study. We also thank Dr. J. Luque for the provision of *Diplodia seriata* isolate JL-398. M.P. Martínez-Diz was supported by the FPI-INIA program from the INIA (CPD2015-0188). D. Gramaje was supported by the Ramón y Cajal program, Spanish Government (RYC-2017-23098).

References

- Alfonzo, A., Conigliaro, G., Torta, L., Burruano, S., and Moschetti, G., 2009. Antagonism of *Bacillus subtilis* strain AG1 against vine wood fungal pathogens. *Phytopathol. Mediterr.* 48, 155-158. DOI: 10.14601/Phytopathol_Mediterr-2886
- Álvarez-Pérez, J.M., González-García, S., Cobos, R., Olego, M.A., Ibañez, A., Díez-Galán, A., Garzón-Jimeno, E., Coque, J.J.R., 2017. Use of endophytic and rhizospheric actinobacteria from grapevine plants to reduce nursery fungal infections that lead to young grapevine decline. *Appl Environ Microb.* 83, e01564-17. <https://doi.org/10.1128/AEM.01564-17>
- Amponsah, N.T., Jones, E.E., Ridgway, H.J., Jaspers, M.V., 2012. Evaluation of fungicides for the management of *Botryosphaeria* dieback diseases of grapevines. *Pest Manag. Sci.* 68, 676-683. <https://doi.org/10.1002/ps.2309>
- Andreoli, M., Zapparoli, G., Angelini, E., Lucchetta, G., Silvia Lampis, S., Vallini, G., 2019. *Pseudomonas protegens* MP12: A plant growth-promoting endophytic bacterium with broad-spectrum antifungal activity against grapevine phytopathogens. *Microbiol. Res.* 219, 123-131. <https://doi.org/10.1016/j.micres.2018.11.003>
- Auger J., Esterio, M., Ricke G., Perez, I., 2004. Black dead arm and basal canker on *Vitis vinifera* cv. Red Globe caused by *Botryosphaeria obtusa* in Chile. *Plant Dis.* 88, 1286. DOI: 10.1094/PDIS.2004.88.11.1286A
- Ayres, M.R., Wicks, T.J., Scott, E.S., Sosnowski, M.R., 2017. Developing pruning wound protection strategies for managing *Eutypa* dieback. *Aust. J. Grape Wine Res.* 23, 103-111. DOI: 10.1111/ajgw.12254
- Baloyi, M.A., Hallen, F., Mostert, L., Eskalen, A., 2016. First report of *Phaeomoniella chlamydospora* pycnidia as Petri disease inoculum sources in South African vineyards. *Plant Dis.* 100, 2528. <https://doi.org/10.1094/PDIS-04-16-0569-PDN>
- Berbegal, M., Ramón-Albalat, A., León, M., Armengol, J., 2020. Evaluation of long-term protection from nursery to vineyard provided by *Trichoderma atroviride* SC1 against fungal grapevine trunk pathogens. *Pest Manag. Sci.* 76, 967-977. DOI: 10.1002/ps.5605
- Bertsch, C., Ramirez-Suero, M., Magnin-Robert, M., Larignon, P., Chong, J., Abou- Mansour, E., Spagnolo, A., Clément, C., Fontaine, F., 2013. Grapevine trunk diseases: complex and still poorly understood. *Plant Pathol.* 62, 243-265. DOI: 10.1111/j.1365-3059.2012.02674.x
- Cloete, M., Fischer, M., Mostert, L., Halleen, F., 2015. Hymenochaetales associated with esca-related wood rots on grapevine with a special emphasis on the status of esca in South African vineyards. *Phytopathol. Mediterr.* 54, 299-312.
- Crous, P.W., Gams, W., 2000. *Phaeomoniella chlamydospora* gen. et comb. nov., a causal organism of Petri grapevine decline and esca. *Phytopathol. Mediterr.* 39, 112-118. https://doi.org/10.14601/Phytopathol_Mediterr-1530
- Daraignes, L., Gerbore, J., Yacoub, A., Dubois, L., Romand, C., Zekri, O., Roudet, J., Chambon, P., Fermaud, M., 2018. Efficacy of *P. oligandrum* affected by its association with bacterial BCAs and rootstock effect in controlling grapevine trunk diseases. *Biol. Control* 119, 59-67. <https://doi.org/10.1016/j.biocontrol.2018.01.008>

- 596 Del Frari, G., Cabral, A., Nascimento, T., Boavida Ferreira, R., Oliveira, H., 2019. *Epicoccum*
597 *layuense* a potential biological control agent of esca-associated fungi in grapevine. PLoS
598 ONE 14, e0213273. <https://doi.org/10.1371/journal.pone.0213273>
- 599 Díaz, G.A., Latorre, B.A., 2013. Efficacy of paste and liquid fungicide formulations to protect
600 pruning wounds against pathogens associated with grapevine trunk diseases in Chile. Crop
601 Prot. 46, 106-112. <http://dx.doi.org/10.1016/j.cropro.2013.01.001>
- 602 Edwards, J., Laukart, N., Pascoe, I., 2001. *In situ* sporulation of *Phaeomoniella chlamydospora*
603 in the vineyard. Phytopathol. Mediterr. 40, 61-66. DOI: 10.14601/Phytopathol_Mediterr-
604 1586
- 605 Edwards, J., Pascoe, I.G., 2001. Pycnidial state of *Phaeomoniella chlamydospora* found on
606 'Pinot Noir' grapevines in the field. Australas. Plant Pathol. 30, 67.
607 <https://doi.org/10.1071/AP00061>
- 608 Elena, G., García-Figueres, F., Reigada, S., Luque, J., 2015a. Intraspecific variation in *Diplodia*
609 *seriata* isolates occurring on grapevines in Spain. Plant Pathol. 64, 680-689. DOI: Doi:
610 10.1111/ppa.12296
- 611 Elena, G., Luque, J., 2016a. Seasonal susceptibility of pruning wounds and cane colonization
612 in Catalonia, Spain following artificial infection with *Diplodia seriata* and *Phaeomoniella*
613 *chlamydospora*. Plant Dis. 100, 1651-1659. DOI: 10.1094/PDIS-10-15-1186-RE
- 614 Elena, G., Luque, J., 2016b. Pruning debris of grapevine as a potential inoculum source of
615 *Diplodia seriata*, causal agent of Botryosphaeria dieback. Eur. J. Plant Pathol. 144, 803-
616 810. <https://doi.org/10.1007/s10658-015-0782-9>
- 617 Elena, G., Sosnowski, M.R., Ayres, M.R., Lecomte, P., Benetreau, C., Garcia-Figueres, F.,
618 Luque, J., 2015b. Effect of the inoculum dose of three grapevine trunk pathogens on the
619 infection of artificially inoculated pruning wounds. Phytopathol. Mediterr. 54, 345-354.
620 DOI: 10.14601/Phytopathol_Mediterr-16010
- 621 Elmer, P.A.G., Reglinski, T., 2006. Biosuppression of *Botrytis cinerea* in grapes. Plant Pathol.
622 55, 155e177. <https://doi.org/10.1111/j.1365-3059.2006.01348.x>
- 623 EPPO, 2017. Standard PP 1/303 (1) *Eutypa lata* on grapevine. Bulletin OEPP/EPPO Bulletin
624 47, 333-336.
- 625 Epstein, L., Sukhwinder, K., VanderGheynst, J.S., 2008. Botryosphaeria-related dieback and
626 control investigated in non-coastal California grapevines. Calif. Agric. 62, 161-166.
- 627 Eskalen, A., Feliciano, J., Gubler, W.D., 2007. Susceptibility of grapevine pruning wounds and
628 symptom development in response to infection by *Phaeoacremonium aleophilum* and
629 *Phaeomoniella chlamydospora*. Plant Dis. 91, 1100-1104. [https://doi.org/10.1094/PDIS-](https://doi.org/10.1094/PDIS-91-9-1100)
630 [91-9-1100](https://doi.org/10.1094/PDIS-91-9-1100)
- 631 Eskalen, A., Gubler, W.D., 2001. Association of spores of *Phaeomoniella chlamydospora*,
632 *Phaeoacremonium inflatipes*, and *Pm. aleophilum* with grapevine cordons in California.
633 Phytopathol. Mediterr. 40S, 429-432. DOI: 10.14601/Phytopathol_Mediterr-1613
- 634 Fischer, M., 2002. A new wood-decaying basidiomycete species associated with esca of
635 grapevine: *Fomitiporia mediterranea* (Hymenochaetales). Mycol. Prog. 1, 315-324. DOI:
636 10.1007/s11557-006-0029-4
- 637 Gardes, M., Bruns, T.D., 1993. ITS primers with enhanced specificity for Basidiomycetes:
638 application to the identification of mycorrhizae and rusts. Mol. Ecol. 2, 113-118.
- 639 González-Domínguez, E., Berlanas, C., Gramaje, D., Armengol, J., Rossi, V., Berbegal, M.,
640 2020. Temporal dispersal patterns of *Phaeomoniella chlamydospora*, causal agent of Petri
641 disease and esca, in vineyards. Phytopathology. [https://doi.org/10.1094/PHYTO-10-19-](https://doi.org/10.1094/PHYTO-10-19-0400-R)
642 [0400-R](https://doi.org/10.1094/PHYTO-10-19-0400-R)
- 643 Gramaje, D., Mostert, L., Groenewald, J.Z., Crous, P.W., 2015. *Phaeoacremonium*: From esca
644 disease to phaeohyphomycosis. Fungal Biol. 199, 759-783. DOI:
645 10.1016/j.funbio.2015.06.004

- Gramaje, D., Úrbez-Torres, J.R., Sosnowski, M.R., 2018. Managing grapevine trunk diseases with respect to etiology and epidemiology: Current strategies and future prospects. *Plant Dis.* 102, 12-39. <https://doi.org/10.1094/PDIS-04-17-0512-FE>
- Gubler, W.D., Mugnai, L., Surico, G., 2015. Esca, Petri and Grapevine leaf stripe disease, in: Wilcox, W.F., Gubler, W.D., Uyemoto, J.K. (Eds.), *Compendium of Grape Diseases, Disorders, and Pests*, 2nd Edition. APS Press, St Paul, MN, pp. 52-56.
- Gubler, W.D., Rooney-Latham, S., Vasquez, S.J., Eskalen, A., 2013. Esca (Black Measles) and Petri disease, in: Bettiga, L.J. (Ed.), *Grape Pest Management*. University of California, Agriculture and Natural Resources, Publication 3343, pp. 120-125.
- Haidar, R., Roudet, J., Bonnard, O., Dufour, M., Corio-Costet, M., Fert, M., Gautier, T., Deschamps, A., Fermaud, M., 2016. Screening and modes of action of antagonistic bacteria to control the fungal pathogen *Phaeoemoniella chlamydospora* involved in grapevine trunk diseases. *Microbiol. Res.* 192, 172-184. <https://doi.org/10.1016/j.micres.2016.07.003>
- Halleen, F., Fourie, P.H., Lombard, J., 2010. Protection of grapevine pruning wounds against *Eutypa lata* by biological and chemical methods. *S. Afr. J. Enol. Vitic.* 31, 125-132. DOI: 10.21548/31-2-1409
- Jacobs, K., Bergdahl, D.R., Wingfield, M.J., Halik, S., Seifert, K.A., Bright, D.E., Wingfield, B.D., 2004. *Leptographium wingfieldii* introduced into North America and found associated with exotic *Tomicus piniperda* and native bark beetles. *Mycol. Res.* 108, 411–418. <https://doi.org/10.1017/S0953756204009748>
- John, S., Wicks, T.J., Hunt, J.S., Lorimer, M.F., Oakey, H., Scott, E.S., 2005. Protection of grapevine pruning wounds from infection by *Eutypa lata* using *Trichoderma harzianum* and *Fusarium lateritium*. *Australas. Plant Pathol.* 34, 569-575. <https://doi.org/10.1071/AP05075>
- Kaplan, J., Travadon, R., Cooper, M., Hillis, V., Lubell, M., Baumgartner, K., 2016. Identifying economic hurdles to early adoption of preventative practices: The case of trunk diseases in California winegrape vineyards. *Wine Econ. Pol.* 5, 127-141. DOI: 10.1016/j.wep.2016.11.001
- Kotze, C., Van Niekerk, J., Mostert, L., Halleen, F., Fourie, P., 2011. Evaluation of biocontrol agents for grapevine pruning wound protection against trunk pathogen infection. *Phytopathol. Mediterr.* 50, S247-S263. DOI:10.14601/Phytopathol_Mediterr-8960
- Kühn, A., Zappata, A., Gold, R.E., Zito, R., Kortekamp, A., 2017. Susceptibility of grape pruning wounds to grapevine trunk diseases and effectiveness of a new BASF wound protectant. *Phytopathol. Mediterr.* 56, 536. DOI: 10.14601/Phytopathol_Mediterr-21865
- Kuntzmann, P., Villaume, S., Bertsch, C., 2009. Conidia dispersal of *Diplodia species* in a French vineyard. *Phytopathol. Mediterr.* 48, 150-154. DOI:10.14601/Phytopathol_Mediterr-2885
- Larignon, P., Darné, G., Ménard, E., Desaché, F., Dubos, B., 2008. Comment agissait l'arsénite de sodium sur l'esca de la vigne? *Prog. Agric. Vitic.* 125, 642-651.
- Larignon, P., Dubos, B., 2000. Preliminary studies on the biology of *Phaeoacremonium*. *Phytopathol. Mediterr.* 39, 184-189. DOI: 10.14601/Phytopathol_Mediterr-1554
- Larignon, P., Fulchic, R., Laurent, C., Dubos, B., 2001. Observation on black dead arm in French vineyards. *Phytopathol. Mediterr.* 40, S336-S342. DOI: 10.14601/Phytopathol_Mediterr-1629
- Lecomte, P., Darrieutort, G., Liminana, J.-M., Comont, G., Muruamendiaraz, A., Legorburu, F.-J., Choueiri, E., Jreijiri, F., El Amil, R., Fermaud, M., 2012. New insights into esca of grapevine: The development of foliar symptoms and their association with xylem discoloration. *Plant Dis.* 96, 924-934. <http://dx.doi.org/10.1094/PDIS-09-11-0776-RE>
- Lengyel, S., Gold, R.E., Fischer, J., Yemelin, A., Thines, E., Kühn, A., 2019. Early detection

- project - detection and quantification of *Phaeomoniella chlamydospora* and *Botryosphaeria* spp. in *Vitis vinifera* wood samples. *Phytopathol. Mediterr.* 58, 406-407. DOI: 10.14601/Phytopathol_Mediterr-10627
- Lorch, W., 2014. Fatal wood disease affects 12 percent of French vineyards. Retrieved 19 May 2020 from <https://www.wine-searcher.com/m/2014/10/fatal-wood-diseases-affect-12-percent-of-french-vineyards>
- Luque, J., Elena, G., Garcia-Figueres, F., Reyes, J., Barrios, G., Legorburu, F.J., 2014. Natural infections of pruning wounds by fungal trunk pathogens in mature grapevines in Catalonia (Northeast Spain). *Aust. J. Grape Wine Res.* 20, 134-143. DOI: 10.1111/ajgw.12046
- Luque, J., Martos, S., Aroca, A., Raposo, R., Garcia-Figueres, F., 2009. Symptoms and fungi associated with declining mature grapevine plants in northeast Spain. *J. Plant Pathol.* 91, 381-90. DOI: 10.4454/jpp.v91i2.968
- MAPA, 2020. Official Registry of Phytosanitary Products. Ministerio de Agricultura, Pesca y Alimentación, Spain. Consulted on 13 January 2020 from <https://www.mapa.gob.es/es/agricultura/temas/sanidad-vegetal/productos-fitosanitarios/registro/menu.asp>
- Martínez-Diz, M.P., Díaz-Losada, E., Barajas, E., Ruano-Rosa, D., Andrés-Sodupe, M., Gramaje, D., 2019. Screening of Spanish *Vitis vinifera* germplasm for resistance to *Phaeomoniella chlamydospora*. *Scient. Hortic.* 246, 104-109. <https://doi.org/10.1016/j.scienta.2018.10.049>
- Martínez-Diz, M.P., Díaz-Losada, E., Andrés-Sodupe, M., Bujanda, R., Maldonado-González, M.M., Ojeda, S., Yacoub, A., Rey, P., Gramaje, D., 2020a. Field evaluation of biocontrol agents against black-foot and Petri diseases of grapevine (under review. *BioRxiv* doi: <https://doi.org/10.1101/2020.05.19.101568>).
- Martínez-Diz, M.P., Eichmeier, A., Spetik, M., Bujanda, R., Díaz-Fernández, A., Díaz-Losada, E., Gramaje, D., 2020b. Grapevine pruning time affects natural wound colonization by wood-invading fungi. (under review. *BioRxiv* doi: <https://doi.org/10.1101/2020.04.20.050955>).
- Mayet, V., Lecomte, P., 2014. Un nouveau moyen de protection des blessures de taille. *Union Girondine des vins de Bordeaux*. Février 2014, 55-56.
- Milholland R.D., 1991. Muscadine grapes: Some important diseases and their control. *Plant Dis.* 75, 113-117.
- Moller, W.J., Ramos, D.E., Sanborn, R.R., 1977. *Eutypa* dieback in California apricot orchards: Chemical control studies. *Plant Dis. Rep.* 61, 600-604.
- Mondello, V., Songy, A., Battiston, E., Pinto, C., Coppin, C., Trotel-Aziz, P., Fontaine, F., 2018. Grapevine trunk diseases: a review of fifteen years of trials for their control with chemicals and biocontrol agents. *Plant Dis.* 7, 1189-1217. <https://doi.org/10.1094/PDIS-08-17-1181-FE>
- Mondello, V., Spagnolo, A., Larignon, P., Clément, C., Fontaine, F., 2019. Phytoprotection potential of *Fusarium proliferatum* for control of *Botryosphaeria* dieback pathogens in grapevine. *Phytopathol. Mediterr.* 58, 293-306. DOI: 10.14601/Phytopathol_Mediterr-10617
- Mostert, L., Groenewald, J.Z., Summerbell, R.C., Gams, W., Crous, P.W., 2006. Taxonomy and pathology of *Togninia* (*Diaporthales*) and its *Phaeoacremonium* anamorphs. *Stud. Mycol.* 54, 1-115. <https://doi.org/10.3114/sim.54.1.1>
- Mounier, E., Cortes, F., Cadious, M., Pajot, E., 2014. The benefits of *Trichoderma atroviride* I-1237 for the protection of grapevines against trunk diseases: from the nursery to the vineyard. *Phytopathol. Mediterr.* 53, 591-592. DOI: 10.14601/Phytopathol_Mediterr-15167
- Moyo, P., Allsopp, E., Roets, F., Mostert, L., Halleen, F., 2014. Arthropods vector grapevine

trunk disease pathogens. *Phytopathology* 104, 1063-1069. DOI: 10.1094/PHYTO-11-13-0303-R

Munkvold, G.P., Marois, J.J., 1993. Efficacy of natural epiphytes and colonisers of grapevine pruning wounds for biological control of *Eutypa dieback*. *Phytopathology* 83, 624-629. DOI: 10.1094/Phyto-83-624

Munkvold, G.P., Marois, J.J., 1995. Factors associated with variation in susceptibility of grapevine pruning wounds to infection by *Eutypa lata*. *Phytopathology* 85, 249-256. DOI: 10.1094/Phyto-85-249

Mutawila, C., Fourie, P.H., Halleen, F., Mostert, L., 2011a. Histo-pathology study of the growth of *Trichoderma harzianum*, *Phaeomoniella chlamydospora* and *Eutypa lata* on grapevine pruning wounds. *Phytopathol. Mediterr.* 50, S46-S60. DOI: 10.14601/Phytopathol_Mediterr-8643

Mutawila, C., Fourie, P.H., Halleen, F., Mostert, L., 2011b. Grapevine cultivar variation to pruning wound protection by *Trichoderma* species against trunk pathogens. *Phytopathol. Mediterr.* 50, S264-S276. DOI: 10.14601/Phytopathol_Mediterr-8981

Mutawila, C., Halleen, F., Mostert, L., 2015. Development of benzimidazole resistant *Trichoderma* strains for the integration of chemical and biocontrol methods of grapevine pruning wound protection. *BioControl* 60, 387-399. <https://doi.org/10.1007/s10526-014-9647-y>

Mutawila, C., Halleen, F., Mostert, L., 2016. Optimisation of time of application of *Trichoderma* biocontrol agents for protection of grapevine pruning wounds. *Aust. J. Grape Wine Res.* 22, 279-287. <https://doi.org/10.1111/ajgw.12218>

Niem, J.M., Billones-Baaijens, R., Stodart, B., Savocchia, S., 2020. Diversity Profiling of Grapevine Microbial Endosphere and Antagonistic Potential of Endophytic *Pseudomonas* Against Grapevine Trunk Diseases. *Front. Microbiol.* 11, 477. DOI: 10.3389/fmicb.2020.00477

Pertot, I., Caffi, T., Rossi, V., Mugnai, L., Hoffmann, C., Grando, M.S., Gary, C., Lafond, D., Duso, C., Thiery, D., Mazzoni, V., Anfora, G., 2017. A critical review of plant protection tools for reducing pesticide use on grapevine and new perspectives for the implementation of IPM in viticulture. *Crop Prot.* 97, 70-84. <https://doi.org/10.1016/j.cropro.2016.11.025>

Pertot, I., Prodanutti, D., Colombini, A., Pasini, L., 2016. *Trichoderma atroviride* SC1 prevents *Phaeomoniella chlamydospora* and *Phaeoacremonium aleophilum* infection of grapevine plants during the grafting process in nurseries. *Biocontrol* 61, 257-267. DOI: 10.1007/s10526-016-9723-6

Petzoldt, C.H., Moller, W.J., Sall, M.A., 1981. *Eutypa dieback* of grapevines: seasonal differences in infection and duration of susceptibility of pruning wounds. *Phytopathology* 71, 540-543.

Phillips, A.J.L., 2002. *Botryosphaeria* species associated with diseases of grapevines in Portugal. *Phytopathol. Mediterr.* 41, 3-18. DOI: 10.14601/Phytopathol_Mediterr-1655

Phillips, A.J.L., Crous, P.W., Alves, A., 2007. *Diplodia seriata*, the anamorph of “*Botryosphaeria*” *obtusata*. *Fungal Divers.* 25, 141-155.

Pitt, W.M., Sosnowski, M.R., Huang, R., Qui, Y., Steel, C.C., Savocchia, S., 2012. Evaluation of fungicides for the management of *Botryosphaeria* canker of grapevines. *Plant Dis.* 96, 1303-1308. <https://doi.org/10.1094/PDIS-11-11-0998-RE>

Pitt, W.M., Trouillas, F.P., Gubler, W.D., Savocchia, S., Sosnowski, M.R., 2013a. Pathogenicity of diatrypaceous fungi on grapevines in Australia. *Plant Dis.* 97, 749-756. <https://doi.org/10.1094/PDIS-10-12-0954-RE>

Pitt, W.M., Úrbez-Torres, J.R., Trouillas, F.P., 2013b. *Dothiorella vidmadera*, a novel species from grapevines in Australia and notes on *Spenceriartinsia*. *Fungal Divers.* 61, 209-219. <https://doi.org/10.1007/s13225-013-0244-7>

- Pitt, W.M., Úrbez-Torres, J.R., Trouillas, F.P., 2015. *Dothiorella* and *Spencermartinsia*, new species and records from grapevines in Australia. Aust. Plant Pathol. 44, 43-56. <https://doi.org/10.1007/s13313-014-0332-5>
- Quaglia, M., Covarelli, L., Zazzerini, A., 2009. Epidemiological survey on esca disease in Umbria, central Italy. Phytopathol. Mediterr. 48, 84-91. DOI: 10.14601/Phytopathol_Mediterr-2877
- Reis, P., Letousey, P., Rego, C., 2017. *Trichoderma atroviride* strain I-1237 protects pruning wounds against grapevine wood pathogens. Phytopathol. Mediterr. 56, 580. DOI: 10.14601/Phytopathol_Mediterr-21865
- Rezgui, A., Ben Ghnaya-Chakroun, A., Vallance, J., Bruez, E., Hajlaoui, M. R., Sadfi-Zouaoui, N., Rey, P., 2016. Endophytic bacteria with antagonistic traits inhabit the wood tissues of grapevines from Tunisian vineyards. Biol. Control 99, 28-37. <https://doi.org/10.1016/j.biocontrol.2016.04.005>
- Rolshausen, P.E., Akgül, D.S., Perez, R., Eskalen, A., Gispert, C., 2013. First report of wood canker caused by *Neoscytalidium dimidiatum* on grapevine in California. Plant Dis. 97, 1511. <https://doi.org/10.1094/PDIS-04-13-0451-PDN>
- Rolshausen, P.E., Gubler, W.D., 2005. Use of boron for the control of Eutypa dieback of grapevines. Plant Dis. 89, 734-738. <https://doi.org/10.1094/PD-89-0734>
- Rolshausen, P.E., Úrbez-Torres, J.R., Rooney-Latham, S., Eskalen, A., Smith, R.J., Gubler W.D., 2010. Evaluation of Pruning Wound Susceptibility and Protection Against Fungi Associated with Grapevine Trunk Diseases. Am. J. Enol. Vitic. 61, 113-119.
- Romanazzi, G., Murolo, S., Pizzichini, L., Nardi, S., 2009. Esca in young and mature vineyards, and molecular diagnosis of the associated fungi. Eur. J. Plant Pathol. 125, 277-290. <https://doi.org/10.1007/s10658-009-9481-8>
- Samaras, A., Ntasioy, P., Testempasis, S., Theocharis, S. Koundouras, S., Karaoglanidis, G., 2019. Evaluation of the fungicide Tessior (boscalid and pyraclostrobin) for control of grapevine trunk diseases in Greece. Phytopathol. Mediterr. 58, 421. DOI: 10.14601/Phytopathol_Mediterr-10627
- Savocchia, S., Steel, C.C., Stodart, B.J., Somers, A., 2007. Pathogenicity of *Botryosphaeria* species isolated from declining grapevines in sub-tropical regions of eastern Australia. Vitis 46, 27-32.
- Schubert, M., Fink, F., Schwartze, F.W.M.R., 2008. Evaluation of *Trichoderma* spp. as biocontrol agent against wood decay fungi in urban trees. Biological Control 45, 111-123. <https://doi.org/10.1016/j.biocontrol.2008.01.001>
- Serra, S., Mannoni, A.M., Ligios, V., 2008. Studies on the susceptibility of pruning wounds to infection by fungi involved in grapevine wood diseases in Italy. Phytopathol. Mediterr. 47, 234-246. DOI: 10.14601/Phytopathol_Mediterr-2727
- Siebert, J.B., 2001. *Eutypa*: the economic toll on vineyards. Wines Vines 4, 50-56.
- Sosnowski, M., McCarthy, G., 2017. Economic impact of grapevine trunk disease management in Sauvignon Blanc vineyards of New Zealand. Wine Vitic. J. 32, 42-48.
- Sosnowski, M.R., Creaser, M.L., Wicks, T.J., Lardner, R., Scott, E.S., 2008. Protection of grapevine pruning wounds from infection by *Eutypa lata*. Aust. J. Grape Wine Res. 14, 134-142. DOI: 10.1111/j.1755-0238.2008.00015.x
- Sosnowski, M.R., Loschiavo, A.P., Wicks, T.J., Scott, E.S., 2013. Evaluating treatments and spray application for the protection of grapevine pruning wounds from infection by *Eutypa lata*. Plant Dis. 97, 1599-1604. <http://dx.doi.org/10.1094/PDIS-02-13-0201-RE>
- Sosnowski, M.R., Mundi, D.C., 2019. Pruning Wound Protection Strategies for Simultaneous Control of Eutypa and Botryosphaeria dieback in New Zealand. Plant Dis. 103, 519-525. <https://doi.org/10.1094/PDIS-05-18-0728-RE>
- Spinosi, J., Févotte, J., Vial, G., 2009. Éléments techniques sur l'exposition professionnelle aux

- pesticides arsenicaux. Matrice cultures - expositions aux pesticides arsenicaux. Institut de
veille sanitaire, Saint-Maurice, France.
- Surico, G., 2009. Towards a redefinition of the diseases within the esca complex of grapevine.
Phytopathol. Mediterr. 48, 5-10. DOI: 10.14601/Phytopathol_Mediterr-2870
- Tegli, S., Bertelli, E., Surico, G., 2000. Sequence analysis of ITS ribosomal DNA in five
Phaeoacremonium species and development of a PCR-based assay for the detection of *P.*
chlamydosporum and *P. aleophilum* in grapevine tissue. Phytopathol. Mediterr. 39, 134-
149. DOI: 10.14601/Phytopathol_Mediterr-1555
- Travadon, R., Lawrence, D.P., Rooney-Latham, S., Gubler, W.D., Wilcox, W.F., Rolshausen,
P.E., Baumgartner, K., 2015. *Cadophora* species associated with wood-decay of grapevine
in North America. Fungal Biol. 119, 53-66. <https://doi.org/10.1016/j.funbio.2014.11.002>
- Trotel-Aziz, P., Abou-Mansour, E., Courteaux, B., Rabenoelina, F., Clément, C., Fontaine, F.,
Aziz, A., 2019. *Bacillus subtilis* PTA-271 Counteracts Botryosphaeria Dieback in
Grapevine, Triggering Immune Responses and Detoxification of Fungal Phytotoxins.
Front. Plant Sci. 10, 25. DOI: 10.3389/fpls.2019.00025
- Úrbez-Torres, J.R., 2011. The status of Botryosphaeriaceae species infecting grapevines.
Phytopathol. Mediterr. 50, S5-S45. DOI: 10.14601/Phytopathol_Mediterr-9316
- Úrbez-Torres, J.R., Battany, M., Bettiga, L.J., Gispert, C., McGourty, G., Roncoroni, J., Smith,
R.J., Verdegaa, P., Gubler, W.D., 2010. Botryosphaeriaceae species spore-trapping
studies in California Vineyards. Plant Dis. 94, 717-724. <https://doi.org/10.1094/PDIS-94-6-0717>
- Úrbez-Torres, J.R., Gubler, W.D., 2011. Susceptibility of grapevine pruning wounds to
infection by *Lasiodiplodia theobromae* and *Neofusicoccum parvum*. Plant Pathol. 60, 261-
270. <https://doi.org/10.1111/j.1365-3059.2010.02381.x>
- Úrbez-Torres, J.R., Leavitt, G.M., Guerrero, J.C., Guevara, J., Gubler, W.D., 2008.
Identification and pathogenicity of *Lasiodiplodia theobromae* and *Diplodia seriata*, the
causal agents of Bot canker disease of grapevines in Mexico. Plant Dis. 92, 519-529.
<https://doi.org/10.1094/PDIS-92-4-0519>
- Valencia, D., Torres, C., Camps, R., Lopez, E., Celis-Diez, J., Beosain, X., 2015. Dissemination
of Botryosphaeriaceae conidia in vineyards in the semiarid Mediterranean climate of the
Valparaíso Region of Chile. Phytopathol. Mediterr. 54, 394-402. DOI:
10.14601/Phytopathol_Mediterr-16055
- van Niekerk, J.M., Calitz, F.J., Halleen, F., Fourie, P.H., 2010. Temporal spore dispersal
patterns of grapevine trunk pathogens in South Africa. Eur. J. Plant. Pathol. 127, 375-390.
<https://doi.org/10.1007/s10658-010-9604-2>
- van Niekerk, J.M., Crous, P.W., Groenewald, J.Z., Fourie, P.H., Halleen, F., 2004. DNA
phylogeny, morphology and pathogenicity of *Botryosphaeria* species on grapevines.
Mycologia 96, 781-798. DOI: 10.2307/3762112
- Yan, J.-Y., Xie, Y., Zhang, W., Wang, Y., Liu, J.-K., Hyde, K.D., Seem, R.C., Zhang, G. Z.,
Wang, Z.-Y., Yao, S.-W., Bai, X.-J., Dissanayake, A. J., Peng, Y.-L., Li, X.-H., 2013.
Species of Botryosphaeriaceae involved in grapevine dieback in China. Fungal Divers. 61,
221-236. <https://doi.org/10.1007/s13225-013-0251-8>
- Yang, T., Groenewald, J.Z., Cheewangkoon, R., Jami, F., Abdollahzadeh, J., Lombard, L.,
Crous, P.W., 2017. Families, genera, and species of *Botryosphaeriales*. Fungal Biol. 121,
322-346. <https://doi.org/10.1016/j.funbio.2016.11.001>

Table 1. Pruning wound treatments evaluated for control of *Diplodia seriata* and *Phaeomoniella chlamydospora* under field conditions.

Trade name	Nature	Chemical/Biological group ^a	Active ingredient	Application rate	Supplier
Enovit Metil	Chemical	MBC ^b / Thiophanates	Thiophanate methyl 70%	1 g L ⁻¹	Sipcam Inagra S.L.
Tessor	Chemical	QoI ^c / methoxy-carbamates + SDHI ^d / pyridine-carboxamides	Pyraclostrobin 0.5% + boscalid 1%	n/a [*]	BASF Española S.L.U.
Master + Song	Paste + chemical	DMI ^e / Triazoles	Paste (resin 55% + vegetal oil and healing substances 45%) + tebuconazole 25%	n/a	Sipcam Jardin S.L.+ Sipcam Iberia S.L.
Vintec	BCA ^f	Microbial (fungi)	<i>Trichoderma atroviride</i> SC1 (2 x 10 ¹⁰ CFU g ⁻¹)	2 g L ⁻¹	Belchim Crop Protection España S.A.
Esquive	BCA	Microbial (fungi)	<i>T. atroviride</i> I-1237 (1 x 10 ⁸ CFU g ⁻¹)	100 g L ⁻¹	Idai Nature S.L.

^a According to Fungicide Resistance Action Committee (FRAC) Code List[®] (2020): Fungal control agents sorted by cross resistance pattern and mode of action (https://www.frac.info/docs/default-source/publications/frac-code-list/frac-code-list-2020-final.pdf?sfvrsn=8301499a_2)

^b MBC, Methyl Benzimidazole Carbamates

^c QoI, Quinone outside Inhibitors

^d SDHI, Succinate-dehydrogenase Inhibitors

^e DMI, Demethylation Inhibitors

^f BCA, Biological Control Agent

^{*} n/a, not applicable

Table 2. Efficacy of wound treatments when applied 24 h before inoculation with *Diplodia seriata* in two growing seasons.

Trade name	Active ingredient	Growing season			
		2018/2019		2019/2020	
		MPR ^a	MPDC ^b	MPR	MPDC
	Inoculated control (IC)	58 a	-	68 a	-
Vintec	<i>T. atroviride</i> SC1	52 a	10	53 ab	22
Esquive	<i>T. atroviride</i> I-1237	43 a	26	46 ab	32
Master + Song	Paste (resin 55% + vegetal oil and healing substances 45%) + tebuconazole 25%	14 b	76	38 bc	44
Enovit Metil	Thiophanate methyl 70 %	6 b	90	31 bc	54
Tessior	Pyraclostrobin 0.5% + boscalid 1%	3 b	95	21 c	69

^a Efficacy was based on the mean percent recovery (MPR) of *Diplodia seriata* from the treated canes by traditional isolation.

^b Mean percent disease control (MPDC) of treatments was calculated as the reduction in MPR as a proportion of the inoculated control.

Table 3. Efficacy of wound treatments when applied 24 h before inoculation with *Phaeomoniella chlamydospora*.

Trade name	Active ingredient	MPR ^a	MPDC ^b
	Inoculated control (IC)	36 ab	-
Vintec	<i>T. atroviride</i> SC1	45 a	0
Esquive	<i>T. atroviride</i> I-1237	30 ab	17
Enovit Metil	Thiophanate methyl 70%	19 bc	46
Tessior	Pyraclostrobin 0.5% + boscalid 1%	18 c	51
Master + Song	Paste (resin 55% + vegetal oil and healing substances 45%) + tebuconazole 25%	12 c	67

^a Efficacy was based on the mean percent recovery (MPR) of *Phaeomoniella chlamydospora* from the treated canes by traditional isolation.

^b Mean percent disease control (MPDC) of treatments was calculated as the reduction in MPR as a proportion of the inoculated control.