Snx4-mediated nucleophagy targets transcription factors controlling *ATG* gene expression

Sara E. Hanley¹, Stephen D. Willis¹ and Katrina F. Cooper^{1,2}

¹ Department of Molecular Biology, Graduate School of Biomedical Sciences, Rowan University,
Stratford, NJ, 08084, USA

^{1,2} Corresponding author: cooperka@rowan.edu, Tel: (856)-566-2887

¹ Sara E. Hanley - hanleys2@rowan.edu

¹Stephen D. Willis - willis58@rowan.edu

Characters (with spaces): 64,342

Running title: Snx4-mediated nucleophagy

ABSTRACT

Autophagy is controlled in part by the repression and activation of Autophagy-related (*ATG*) gene transcription. Here, we demonstrate that the conserved Cdk8 Kinase Module (CKM) of the mediator complex represses transcription of several *ATG* genes. To relieve this repression following nitrogen starvation, Med13 is rapidly degraded via a novel selective autophagy mechanism. This pathway requires the core autophagy machinery but is independent of known nucleophagy systems. It requires the cytosolic filament nucleoporin Gle1, the sorting nexin Snx4-Atg20 heterodimer, and the scaffold protein Atg17. This suggests a model where Med13 traverses through the nuclear pore complex, passing from Gle1 to Snx4. Snx4 then transports Med13 to autophagosomes by binding to Atg17. This previously unidentified nucleophagy pathway also mediates the autophagic degradation of two transcriptional activators of *ATG* genes (Rim15, Msn2) suggesting that this mechanism targets transcription factors that regulate *ATG* expression. This system provides a new level of selectivity, permitting the cell to fine-tune the autophagic response by controlling the turnover of both positive and negative *ATG* transcription factors.

Keywords: Gle1/Med13/Nucleophagy/Selective autophagy/Snx4

INTRODUCTION

Macro-autophagy (hereafter autophagy) is a controlled catabolic process that aids cellular survival during adverse conditions such as starvation or proteolytic stress, by degrading damaged or unnecessary proteins in the vacuole (lysosome in higher eukaryotes) (Klionsky & Codogno, 2013). In budding yeast, non-selective pathways are upregulated in response to nitrogen depletion. This triggers a cascade of events, resulting in non-specific cytosolic cargos being sequestered within autophagosomes and ultimately degraded by vacuolar proteolysis. Selective autophagy pathways use receptor proteins to recognize and deliver specific cargos such as organelles, protein aggregates, and large multi-subunit complexes to autophagosomes (Farre & Subramani, 2016).

Nuclear autophagy or nucleophagy is the least well understood of the selective autophagy mechanisms. Underscoring its importance, various pathologies namely cancer, and neurodegeneration are linked with perturbed nucleophagy (Fu *et al*, 2018). It is best characterized in yeast where macro-nucleophagy utilizes a receptor protein and involves the sequestration of a portion of the nucleus into autophagosomes (Mochida *et al*, 2015). In contrast, micro-nucleophagy (Piecemeal nucleophagy), is autophagosome independent, and forms nuclear-vacuole junctions which pinch off portions of the nucleus directly into the vacuolar lumen (Roberts *et al*, 2003). Recently, an autophagic mechanism has been described that removes defective nuclear pore complexes (NPCs) (Lee *et al*, 2020).

In *S. cerevisiae*, 41 unique autophagy-related (*ATG*) genes have been identified that control this highly coordinated and complex process (Delorme-Axford & Klionsky, 2018). Accordingly, these genes are tightly regulated at multiple levels. Recently we have shown that the cyclin C-Cdk8 kinase negatively regulates *ATG8* expression within the Ume6-Rpd3 HDAC axis (Willis *et al*, 2020). This kinase, together with Med13 and Med12 form the Cdk8 kinase module (CKM) of the mediator complex that in yeast predominantly repress transcription of a diverse set of meiosis and stress response genes (Cooper *et al*, 1997; Cooper *et al*, 1999) by interacting with DNA bound

transcription factors and RNA polymerase II (Akoulitchiev *et al*, 2000; Jeronimo *et al*, 2016; Nemet *et al*, 2014).

Activation of genes controlled by the CKM is achieved by disrupting its association with the mediator (Jeronimo & Robert, 2017). Studies from our group revealed that this is achieved by CKM disassembly. However, we observed that the mechanisms used to disassemble the CKM is dependent upon environmental cues (outlined in Fig. 1) (Cooper *et al*, 2014). In short, oxidative stress triggers cyclin C translocation to the cytoplasm (Cooper *et al*, 2012) where it mediates stress-induced mitochondrial fission and regulated cell death (RCD) in both yeast (Cooper *et al*, 2014) and mammalian cells (Ganesan *et al*, 2019; Jezek *et al*, 2019; Wang *et al*, 2015). Its nuclear release is dependent upon Med13's destruction by the UPS (Khakhina *et al*, 2014; Stieg *et al*, 2018). In contrast, following a survival cue (nitrogen starvation), cyclin C is rapidly destroyed by the UPS before its nuclear release which prevents mitochondrial fission (Willis *et al*, 2020).

This study reveals that Med13 is degraded by vacuolar proteolysis by a previously undescribed autophagic pathway which requires the cytosolic nucleoporin Gle1 and the sorting nexin heterodimer Snx4-Atg20. Moreover, two transcriptional activators that regulate *ATG* expression were also degraded upon nitrogen starvation by this mechanism. Taken together, this suggests a model in which Snx4-mediated nucleophagy of *ATG* transcriptional regulators allows fine-tuning of the autophagic response. This highly selective autophagy mechanism rapidly degrades substrates and requires the nuclear pore complex which makes this pathway distinct from previously described nucleophagy pathways.

RESULTS

Med13 is actively degraded following nitrogen starvation.

We started this investigation by addressing if Med13 was destroyed following nitrogen starvation. Wild-type cells expressing endogenous Med13-9xmyc were starved for nitrogen (SD-N), and Western blot analysis showed that Med13 protein levels decreased with a half-life of 2.6 h

(Fig 1B, D, source data Fig 1). Similarly, Med13 was rapidly degraded in replete medium containing rapamycin, a drug that mimics nitrogen starvation by inhibiting TORC1 (Li *et al*, 2014) (Fig 1B, Fig EV1A). As Med13 half-life is >6 h in unstressed cultures (Khakhina *et al.*, 2014) and *MED13* mRNA increased following 4 h in SD-N (Fig EV1B), these results indicate that Med13 is actively degraded following TORC1 inhibition.

Med13 degradation following nitrogen starvation is mediated by the vacuole.

Med13 levels were next monitored in *ump1Δ*, a mutant deficient for 20S proteasome assembly (Ramos *et al*, 1998), and no change in degradation kinetics was observed (Fig 1C, D). In contrast, in a vacuolar protease mutant (*pep4Δ prb1Δ.1*) (Takeshige *et al*, 1992; Van Den Hazel *et al*, 1996) Med13 was stable in SD-N (Fig 1C, D) (half-life >15 h) indicating that Med13 degradation requires vacuolar proteolysis. Confirming this, the same results were obtained in wild type, *ump1Δ* and *pep4Δ prb1Δ.1* cells harboring a low copy, functional Med13-3xHA plasmid (Stieg *et al.*, 2018) (Fig EV1C, D). Moreover, we found that after nitrogen starvation GFP accumulated in Med13-GFP cleavage assays. This indicates that Med13-GFP is degraded in the vacuole as the compact fold of GFP renders it resistant to vacuolar hydrolases (Shintani & Klionsky, 2004). Accordingly, repeating these cleavage assays in *pep4Δ prb1Δ.1* cells abolished the formation of free GFP and stabilized full-length Med13-GFP (Fig 1E). As anticipated from our previous studies (Khakhina *et al.*, 2014) Med13-GFP was destroyed following 0.8 mM H₂O₂ and no GFP accumulation was seen (Fig 1F). These results show that the proteolysis machinery employed to degrade Med13 is dependent upon environmental cues.

To visualize Med13 vacuolar degradation we used live-cell imaging of endogenous Med13-mNeongreen in *pep4Δ prb1Δ.1* cells. In SD media, Med13-mNeongreen is nuclear (Fig 2A) but after 4 h in SD-N, Med13 accumulated in the vacuole (Fig 2B). The deconvolved collapsed images in Fig 2C also captured Med13-mNeongreen transitioning between these organelles. After 24 h in SD-N, Med13-mNeongreen is exclusively vacuolar (Fig EV2A), and similar results were obtained

when endogenous Med13-YFP was expressed in wild-type cells treated with PMSF that blocks the activity of vacuolar serine proteases (Fig 2B) (Takeshige *et al.*, 1992). These results are consistent with a model in which nitrogen starvation triggers Med13 vacuolar proteolysis.

Med13 degradation requires core autophagy machinery.

Next, we investigated if Med13 degradation was dependent upon proteins required for autophagosome biogenesis (see Fig 3A). Med13 was significantly stabilized following nitrogen starvation in mutants defective in induction (*atg1Δ*) or phagophore fusion (*atg8Δ*) (Kamada *et al.*, 2000) (half-lives >15 h, Fig 3B, C). Likewise, GFP accumulation from Med13-GFP was abolished in various autophagy mutants (Fig 3A, D). We did observe some turnover of full-length Med13-GFP which we attribute to a combination of Med13 being expressed from the *ADH1* promotor rather than the endogenous locus used for degradation assays, and UPS activity. Consistent with this, Med13-GFP was more stable in *atg1Δ ump1Δ* than *atg1Δ* cells (Fig EV3A). Therefore, both the vacuole and core autophagy proteins are needed for Med13 degradation.

Med13 degradation does not use known selective autophagy pathways

Selective autophagy of excess or damaged cellular components in physiological conditions requires the scaffold protein Atg11 (Zientara-Rytter & Subramani, 2020). In starvation conditions, Atg17 replaces Atg11 and functions as the scaffold protein tethering the Atg1 kinase complex to the growing phagophore (Matscheko *et al.*, 2019). Analyzing the autophagic degradation of Med13-GFP (Fig 3E) or degradation of endogenous Med13-9xmyc (Fig EV3B, C) revealed that Atg17 but not Atg11 is required for Med13 destruction. Likewise, known receptors for selective autophagy pathways (Cue5, Atg19, Atg36 and Atg32, Fig EV3D) (Farre & Subramani, 2016) including those needed for micro- and macro- nucleophagy (Nvj1, Atg39 and Atg40, Fig 4A, Fig EV3B, C) (Millen *et al.*, 2009; Mochida *et al.*, 2015) are also not required. This is consistent with a model that Med13 is degraded via an undescribed nucleophagy pathway.

The cytoplasmic nucleoporin Gle1 associates with Med13 after nitrogen starvation.

To define components of this pathway, pull-down assays using Med13-3xHA followed by mass-spectroscopy was used (Fig EV4A, B and Table I for a list of candidate interactors). The conserved, essential nucleoporin Gle1, that localizes to the cytoplasmic filament of the nuclear pore complex (NPC) was identified. Gle1 is a regulator of DEAD-box proteins that are required to facilitate changes to ribonucleoprotein complexes involved in mRNA export and translation initiation (Alcazar-Roman *et al*, 2006; Aryanpur *et al*, 2017; Weirich *et al*, 2006). Co-immunoprecipitation analysis revealed that Med13-3xHA and endogenous Gle1-GFP interacted following 1 h SD-N, confirming this interaction (Fig 4B, source data Fig 2). Yeast two-hybrid assays (Y2H) showed that the large central intrinsic disordered region (IDR) of Med13, which provides a flexible interaction hub for multiple partners (Stieg *et al.*, 2018; Uversky, 2011), interacts with Gle1 (Fig 4C, Fig EV5A, B). In addition, endogenous Med13-mNeongreen and Gle1-RedStar were followed by live-cell imaging. Med13 moves from being diffuse nuclear to associating with the punctate Gle1-RedStar that surrounds the nucleus on its passage to the vacuole after nitrogen starvation (Fig 4D).

Gle1 is an essential protein (Murphy & Went, 1996), therefore, the auxin-inducible degron (AID) system was used to reduce Gle1 protein levels (Fig EV4C). The autophagic degradation of Med13-GFP was significantly decreased following auxin treatment (Fig 4E) as monitored by GFP accumulation. Moreover, after nitrogen depletion, Med13-mNeongreen was retained in the nucleus in the *GLE1*-degron (Fig 4F). However, Crm1 and Msn5, two major Ran-dependent β karyopherins that export mRNA and nuclear proteins following stress (Hutten & Kehlenbach, 2007; Mosammaparast & Pemberton, 2004; Yoshida & Blobel, 2001), are not required for Med13 autophagic degradation (Fig EV4D, E). This strongly suggests that Med13 is transported through the NPC to exit the nucleus, ending at the Gle1 exchange platform.

The sorting nexin-heterodimer Snx4-Atg20 is required for efficient autophagic degradation of Med13.

To further define components of this new nucleophagy mechanism, null alleles of candidate proteins from the mass spectroscopy were screened using Med13-GFP cleavage assays. In cells deleted for the conserved sorting nexin Snx4 (Atg24), we found that free GFP was significantly reduced (Fig 5A). Snx4 is a member of the conserved sorting nexin family of proteins. Snx4 forms distinct heterodimer complexes, with either Snx41 or Atg20, which mediate retrograde trafficking of cargo from the vacuole and endosomes to the Golgi. This role maintains homeostasis and is dispensable for non-selective autophagy (Ma & Burd, 2020; Ma *et al*, 2017; Ma *et al*, 2018; Suzuki & Emr, 2018). In contrast, Snx4-Atg20 is essential for many selective autophagy pathways including mitophagy and pexophagy (Deng *et al*, 2013; Kanki *et al*, 2009; Nice *et al*, 2002; Popelka *et al*, 2017; Shpilka *et al*, 2015)

Co-immunoprecipitation analysis confirmed the interaction between Med13 and Snx4 in SD-N (Fig 5B). Med13 degradation assays revealed that Snx4 and Atg20, but not Snx41, mediate Med13 autophagic degradation (Fig 5C, D). The half-life of Med13 in these mutants was 6.5 and 7.0 h, respectively, compared to >15 h seen in core autophagic mutants. Cytosolic and vacuolar Med13-mNeongreen levels were drastically reduced in *snx4Δ pep4Δ prb1Δ.1* mutants compared to the *pep4Δ prb1Δ.1* control (Fig 5E, F, G). These data demonstrate that the Snx4-Atg20 heterodimer is required for maximal Med13 autophagic degradation, but in its absence, limited autophagic degradation of Med13 still can occur. Taken together these results show that Snx4-Atg20 heterodimer mediates the efficient autophagic degradation of Med13

Snx4 localizes to the nuclear periphery to transport Med13 to autophagosomes.

Snx4 binds to the scaffold protein Atg17 (Nice *et al*, 2002), whose major autophagic role is binding the Atg1 kinase to the PAS (Hollenstein & Kraft, 2020). As both Snx4 and Atg17 mediate

Med13 degradation, this suggests a model in which once Med13 passes through the NPC, it is recognized by Snx4 and delivered to the growing phagophore by Snx4-Atg17 association. Consistent with this, Med13-3xHA and endogenous Atg17-GFP co-immunoprecipitated following 2 h in SD-N. In *snx4Δ*, this interaction was drastically decreased (Fig 6A). Also, endogenous Atg17-RedStar co-localized with Med13-mNeongreen after nitrogen starvation (Fig EV6A). This suggests that Snx4-Atg17 interaction promotes the efficient recruitment of Med13 to autophagosomes. Quantitative co-localization analysis with GFP-Snx4 with the nuclear marker Nab2-mCherry, showed that nitrogen depletion triggers a ~10-fold increase in perinuclear GFP-Snx4 foci (Fig 6B, Fig EV6B, C). Moreover, Med13-mNeongreen co-localized with both perinuclear and cytosolic Snx4 foci in SD-N (Fig 6C and EV6D). Together, this suggests that Snx4 localizes to the nuclear periphery to retrieve and transport Med13 to autophagosomes via Atg17 association.

The above model predicts that Snx4 and Gle1 interact whilst “handing-off” Med13 from the NPC to the sorting nexin complex. Live-cell imaging showed that perinuclear Snx4 foci are adjacent to, or co-localize with, Gle1 in unstressed cultures. After nitrogen starvation, an increased number of perinuclear Snx4 foci co-localize with Gle1 (Fig 6D and Fig EV6E), which is consistent with our model that Snx4 localizes to the NPC to retrieve Med13.

Snx4 specifically targets for Med13 autophagic degradation.

To further understand the sequential stages of this selective autophagy pathway, we next asked if Med13 nuclear localization was required for retrieval and transport by Snx4. To address this, we fused Med13-GFP to the N-terminus of Crn1, a protein associated with actin rafts, which relocalized Med13 to the plasma membrane (Humphries *et al*, 2002) (Fig 7A). Autophagic degradation assays with Crn1-Med13-GFP showed that GFP accumulated in wild-type cells, and this was mostly dependent on Snx4 (Fig 7B). This confirms that Med13 can be targeted for Snx4-mediated autophagic destruction even when located outside the nucleus. Autophagic degradation of Pgc1-GFP, an established substrate of non-selective autophagy (Welter *et al*, 2010), is Snx4

independent (Fig EV7A). This provides further evidence that Snx4 binds to Med13 after nuclear release and this interaction is highly specific.

We next addressed whether Gle1 and Snx4 remain part of the Med13 complex delivered to the vacuole. No substantial accumulation of GFP was seen from GFP-Snx4 (Fig 7C) or Gle1-GFP (Fig 7D), suggesting that neither protein is incorporated into autophagosomes or degraded by the vacuole. Importantly, by monitoring the timing of GFP accumulation from Med13-GFP, we observed that the earliest detection of GFP occurred at 2 h of nitrogen depletion when both Gle1 and Snx4 are present (Fig 7E). These data support the model that Gle1 and Snx4 mediate Med13 delivery to the autophagosome, but these proteins themselves are not vacuolar substrates.

Snx4 BAR domains interact with the C-terminal domain of Med13

To further understand the interaction between Gle1, Snx4, and Med13, we used Y2H analysis to ask if Snx4 and Gle1 associates with the same or different domains of Med13. The results show that Snx4 interacts with the structured C-terminal tail domain of Med13 (Fig 7F, EV5C). Phyre2 plot analysis of this region (Kelley *et al*, 2015) revealed two potential domains (Fig 7G), of which only one, Med13⁹⁰⁷⁻¹¹⁶³ interacted with Snx4 (Fig 7G). Therefore Snx4 interacts with a previously undescribed region of Med13 that lies adjacent to the Gle1 interaction domain. Moreover, as Snx4 is not a nuclear protein, this interaction may be direct and define a new role for Snx4 in transporting nuclear proteins.

Snx4 is a conserved member of the SNX-BAR subfamily of sorting nexin proteins. Common to all SNX family members it contains a phosphoinositide-binding phox homology (PX) domain, which binds to phosphatidylinositol 3-phosphate enriched endosomal membranes. It also contains two BAR (Bin/Amphiphysin/Rvs) domains that bind to curved membranes upon dimerization (Popelka *et al*, 2017; Stanishneva-Konovalova *et al*, 2016). Y2H interaction analyses between either Snx4 domains (PX or BAR domains) and Med13⁹⁰⁷⁻¹¹⁶³ indicated that the Med13-Snx4

interaction occurs through the BAR domain region (Fig 7H). Taken together, these results indicate that the BAR domains recognize Snx4 cargo binding as well as dimerization partners.

Med13 negatively regulates the transcription of a subset of ATG genes.

We next explored if this newly described nucleophagy pathway affects viability during nitrogen starvation. Consistent with previous studies (Nemec *et al.*, 2017), efficient survival in prolonged nitrogen starvation conditions requires Snx4 (Fig EV7B). Snx4 is dispensable for non-selective autophagy but required for many forms of non-selective autophagy (Ma *et al.*, 2018). This suggests that the role of Snx4 in these pathways is essential for cellular adaptation and survival in prolonged starvation conditions. Survival during periods of nutrient depletion requires the upregulation of ATG genes. As Med13 destruction following H₂O₂ relieves repression on SRGs (Khakhina *et al.*, 2014), here asked if a similar strategy was used for the nitrogen starvation response. RT-qPCR analysis of ATG mRNA levels in unstressed wild-type and *med13Δ* cells showed a ~fourfold increase in ATG8 mRNA levels (Fig 8A) that was mirrored by Atg8 protein levels (Fig EV7C). Furthermore, ATG1 and ATG14 mRNA levels were also elevated (Fig 8A), indicating Med13 represses transcription of a subset of ATG genes. Moreover, they are consistent with the model that the destruction of both cyclin C and Med13 following nitrogen starvation are mechanisms used to relieve this repression. These results indicate that Snx4 mediates a new specialized autophagy program able to target a transcriptional repressor that provides a positive feedback loop for the autophagic response.

Snx4-mediated nucleophagy degrades other transcription factors following nitrogen starvation.

Next, we wanted to explore the idea that Snx4-mediated nucleophagy is a mechanism used by the cell to fine-tune the autophagic response at the level of transcription. We first examined the transcriptional activator Rim15 that also regulates ATG genes. During nitrogen starvation, Rim15

enters the nucleus to directly phosphorylate and inhibit the activity of the transcriptional repressors Ume6 (Bartholomew *et al*, 2012) and Rph1 (Bernard *et al*, 2015). Surprisingly, Rim15-GFP cleavage assays revealed Rim15 degradation by Snx4-dependent vacuolar proteolysis that is independent of known nucleophagy pathways (Fig 8B).

To address if Snx4-mediated nucleophagy may have a more global role in degrading transcription factors that control *ATG* expression, the transcriptional activators Msn2 and Ccl1 were tested (Bernard *et al*, 2015; Vlahakis *et al*, 2017; Zhu *et al*, 2016). Free GFP accumulated only from Msn2-GFP in wild-type cells following nitrogen starvation, illustrating that Msn2 is degraded via the vacuole, and this degradation requires Snx4 (Fig 8C). These results support the conclusion that Snx4-mediated degradation of transcription factors targets a unique subset of nuclear proteins. These findings also provide the first evidence that the autophagic pathway directly targets regulatory proteins that control its own processes.

DISCUSSION

Here, by following Med13's fate after nitrogen starvation, we have uncovered a previously undescribed nucleophagy pathway in which both negative and positive transcriptional regulators of *ATG* genes are degraded. Our findings, together with previous studies, suggest a two-step pathway involving the NPC and Snx4 for Med13 translocation from the nucleus to the autophagic machinery (outlined in Fig 8D). First, Med13 disassociates from the CKM, shuttles through the NPC and associates with the cytoplasmic nucleoporin Gle1. In the second step, Med13 is handed-off from Gle1 to the sorting nexin heterodimer Snx4-Atg20. Once assembled, this complex localizes to the PAS via the non-selective autophagy scaffold Atg17. Lastly, Snx4-Atg20 is recycled back to the cytosol, and Med13 is degraded by vacuolar proteolysis. This new pathway is distinct from previously identified mechanisms, being the first described pathway to specifically target transcription factors for autophagic degradation. This mode of degradation does not require the canonical nucleophagy mechanisms but instead requires the NPC. Notably, the pathway uses the

non-selective scaffold Atg17, and lastly the process is rapid, occurring within 4 h making it distinct from other forms of selective autophagy which require longer starvation times.

How does Med13 find its way to the autophagosome? Med13 is a large, 160 kDa protein that must require an active process to transit from the nucleus to the cytoplasm. Its unusual structure, two folded domains separated by a ~1000 amino acid IDR region, suggests it functions as a scaffold and interaction hub. This model is supported by both physical (Nagulapalli *et al*, 2016) and genetic studies (Stieg *et al.*, 2018). Consistent with this activity, the IDR of Med13 interacts with Gle1, cyclin C, and Cdk8 (see Fig. 4) and is the target of additional regulatory protein kinases including Snf1 (Willis *et al*, 2018). The established role of the Gle1-Dbp5 complex in releasing RNPs (ribonucleoproteins) from mRNPs (messenger ribonucleoprotein particles) (Folkmann *et al*, 2011) sets a precedent for Gle1 handing off proteins once they transit through the NPC. Moreover, recent structural data places the Gle1-Dbp remodeling complex right over the NPC's central channel, allowing it to efficiently capture proteins as they reach the cytoplasmic side of the NPC (Fernandez-Martinez *et al*, 2016). Although speculative, a similar capture and release mechanism may be used in the Med13 handoff between Gle1 to Snx4. In support of this model, Gle1 and Snx4 co-localize at the NPC (Fig. 6D), but currently, it is unknown if these proteins directly interact.

The best-characterized cargo of Snx4-Atg20 is Snc1, a plasma membrane-directed v-SNARE, but it remains unknown how this protein binds to Snx4 (Bean *et al*, 2017; Hettema *et al*, 2003; Ma *et al.*, 2017). Similar to our results with Med13, Snc1 associates with the BAR domain of Snx4 using both co-immunoprecipitation analysis as well as Y2H assays (Zhang *et al*, 2009). Taken together, these results suggest that the BAR domain of Snx4 can both recognize cargos as well as be used for dimerization with Atg20. As the newly defined vital function of autophagy receptor proteins is to stably connect autophagy cargo with scaffold proteins (Hollenstein *et al*, 2019) our results place Snx4 as a receptor protein for this novel nucleophagy mechanism. Supporting this,

Snx4 co-localizes with Med13 at both perinuclear and cytoplasmic locations. Our finding that efficient Med13 and Atg17 co-immunoprecipitation requires Snx4 suggests that this sorting nexin relays Med13 from the NPC to the autophagic machinery. In addition, Snx4 mediates Med13 vacuolar degradation when this nuclear protein is relocalized to the plasma membrane, illustrating that Snx4 specifically targets and transports Med13 to growing autophagosomes. Defining the Snx4 interaction motif on Med13, as well as Msn2 and Rim15, may result in a common “Snx4 recognition motif” that could be used to identify additional transcription factors that are degraded by this mechanism. This would be useful because, despite the recent discovery of additional Snx4 cargos, (Bean *et al.*, 2017) no consensus Snx4-dependent sorting signal has been identified (Ma & Burd, 2020). This is important as Snx4 is evolutionarily conserved (Zhang *et al.*, 2018), with recent discoveries that Snx4 dysregulation is now associated with the etiology of many diseases, including, cancer, Parkinson's disease and Alzheimer's disease (Hu *et al.*, 2015).

The most unexpected finding from this research was that the complex process of autophagy is used to degrade transcription factors. More unexpectedly, was that two transcriptional activators of *ATG* genes are also regulated by this pathway. This suggests that nucleophagy can fine-tune transcription and points to a new regulatory role for autophagy beyond its canonical function of recycling damaged/unnecessary proteins or organelles. The molecular details of how these transcription factors are initially targeted and then transported through the NPC are unknown. These details will define the nuclear components required for this pathway. This is important as nucleophagy is the least well defined of all the autophagy pathways and in recent years links between deficiencies and human diseases are starting to emerge (Papandreou & Tavernarakis, 2019). Given the highly conserved nature of both Med13 and Snx4 (Tsai *et al.*, 2013; Zhang *et al.*, 2018), these studies are likely to be relevant to mammalian systems.

METHODS

Yeast strains and plasmids

Experiments were primarily performed with endogenously labeled proteins in the *S. cerevisiae* W303 background (Ronne & Rothstein, 1988) and are listed in Table S2. All strains were constructed using replacement methodology (Janke *et al.*, 2004). Other strains used were from the Research Genetics yeast knock out collection (Chu & Davis, 2008) and are derived from BY4741 strain background. The Y2H assays were performed in the Y2H Gold strain (PT4084-1, Takara 630489, Matchmaker Gold Yeast Two-Hybrid System). The *GLE1*-Auxin-inducible depletion strain (Fig 4E, RSY2456) was a gift from K. Cunningham (Snyder *et al.*, 2019). Live cells were treated with 250 μ M Auxin (Indole-3-acetic acid, Gold Bio I-110) dissolved in ethanol 30 m before nitrogen starvation. The doxycycline-inducible Crm1 N-end rule degron strain (Fig EV4D) was generated by integrating pMK632 (Gnanasundram & Kos, 2015) into the *CRM1* locus in the presence of the pCM188 TET activator plasmid to create RSY2348 (Ubi-Leu-3HA-*CRM1::NATMX*) as described in detail in (Willis *et al.*, 2020). In accordance with the Mediator nomenclature unification efforts (Bourbon *et al.*, 2004) members of the Cdk8 module, the cyclin C (*SSN8/UME3/SRB11*), Cdk8 (*SSN3/UME5/SRB10*), *MED12* (*SSN5*) and (*MED13/UME2/SRB9/SSN2*) will use *CNC1*, *CDK8*, *MED12* and *MED13* gene designations, respectively.

Plasmids used in this study are listed in Extended Data Table 3. The wild-type epitope-tagged plasmids Vph1-mCherry, Nab2-mCherry, Med13-HA and *MED13* Y2H *GAL4* activating domain plasmids have been previously described (Stieg *et al.*, 2018; Willis *et al.*, 2020). The *GAL4-BD-SNX4* fusion plasmids were constructed by amplifying *SNX4* alleles from wild-type genomic DNA with oligonucleotides containing *SaI* flanking sites and cloning into the *SaI* site of the *GAL4* binding domain plasmid pAS2 (Wang & Solomon, 2012). The Crn1-GFP-Med13 plasmid was created by amplifying the first 400 amino acids of Crn1 from genomic DNA and cloning it in-frame to the N

terminus of GFP-Med13 (pSW218). Plasmid construction details are available upon request. All constructs were verified by sequencing.

Cell growth

Yeast cells were grown in either rich, non-selective medium (YPDA: 2% (w/v) glucose, 2% (w/v) Bacto peptone, 1% (w/v) yeast extract, 0.001% (w/v) adenine sulfate) or synthetic minimal dextrose medium (SD: 0.17% (w/v) yeast nitrogen base without amino acids and ammonium sulfate, 0.5% (w/v) ammonium sulfate, 1 x supplement mixture of amino acids, 2% (w/v) glucose) allowing plasmids selection as previously described (Cooper *et al.*, 1997). For nitrogen starvation experiments, cells were grown as described (Journé *et al.*, 2009).

Cellular assays

RT-PCR analysis was executed as previously described in (Cooper *et al.*, 2012; Willis *et al.*, 2020). Oligonucleotides used during these studies are available upon request. All studies were conducted with three biological samples in technical triplicates. The standard deviation from three replicate reactions is indicated in the figures. P values were determined using the unpaired Students t-test. The six-day nitrogen starvation viability assays were executed exactly as described (Willis *et al.*, 2020) with 30,000 cells counted per timepoint using FACS and the studies were conducted in biological duplicates. Data are mean $\pm$ standard deviation

Western blot assays

Protein extracts for Western blot studies were prepared using a NaOH lysis procedure exactly as described in (Willis *et al.*, 2018). In short, protein extracts were prepared from 25 ml per timepoint with the exception of Med13-9xmyc cultures in which 50 ml was needed to visualize the protein. Proteins were separated on 6-10% SDS polyacrylamide gels depending upon their size using the Bio-RAD Mini-Trans Blot cell. To detect proteins, 1:5000 dilutions of anti-myc (UpState New York),

anti-HA (Abcam) or anti-Pgk1 (Invitrogen) antibodies were used. Western blot signals were detected using 1:5000 dilutions of either goat anti-mouse or goat anti-rabbit secondary antibodies conjugated to alkaline phosphatase (Abcam) and CDP-Star chemiluminescence kit (Invitrogen, cat.#T2307). Signals were quantified relative to Pgk1 or Tubulin (Fig EV1C) controls using CCD camera imaging (Kodak Inc.) All degradation assays were performed in triplicate. Standard deviation and significance were calculated from the mean $\pm$ standard deviation using GraphPad Prism 7.

Cleavage assays

Strains harboring GFP-fusion proteins were grown to mid-log in SD, washed and resuspended in SD-N. Protein extracts were prepared using NaOH as described above (25 mLs /TP). Proteins were separated using Invitrogen Blot™ 4-12% Bis-Tris Plus gradient gels with 1X MOPS SDS running buffer (cat.#NW04122BOX). Proteins were transferred to PVDF membranes in 1X Blot™ transfer buffer for 1h 30mins (cat.#BT00061). GFP tagged proteins were detected using 1:5000 dilution of Anti-GFP (WAKO) antibodies and goat anti-mouse secondary antibodies conjugated to alkaline phosphatase.

Co-Immunoprecipitation

For co-immunoprecipitation experiments, 1 L of cells were grown to mid-log, washed and resuspended in SD-N media (250 mLs/TP). Protein extracts were prepared using a glass bead lysis method exactly as described in (Stieg *et al.*, 2018), except Protein A beads were pre-washed with IP wash solution (500mM NaCl, 25mM Tris). 1 mg of total protein was immunoprecipitated per timepoint. Anti-GFP antibodies (Invitrogen) or Anti-HA antibodies (Abcam) were used for immunoprecipitations. Co-immunoprecipitation blot was probed with antibodies against HA (Abcam) or T7 (Novagen) epitope. Due to the drastic difference in size between Med13-3xHA and GFP tagged proteins (Snx4 and Atg17) input controls were run on separate gels. For Atg17

experiments endogenous Atg17-GFP was immunoprecipitated from whole lysates to obtain input controls. For all other input controls 50 µg of protein was resuspended in 2 x SDS-PAGE and separated on either 6% (Med13-3HA) or 10% (GFP-Snx4 and Atg17-GFP) SDS polyacrylamide gels. All co-immunoprecipitation experiments were performed in *pep4Δ prb1Δ.1* strains. (Stieg *et al.*, 2018; Willis *et al.*, 2018).

Fluorescence Microscopy

For all microscopy experiments, cells were grown to mid-log phase, washed and resuspended in SD-N for the time points indicated. Deconvolved images were obtained using a Nikon microscope (Model E800) with a 100x objective with 1.2x camera magnification (Plan Fluor Oil, NA 1.3) and a CCD camera (Hamamatsu Model C4742). Data were collected using NIS software and processed using Image Pro software. All images of individual cells were optically sectioned (0.2 µm slices at 0.3 µm spacing) and deconvolved, and slices were collapsed to visualize the entire fluorescent signal within the cell. The nuclei were visualized in live cells using Hoechst staining (Cayman Chemical 15547). Hoechst (5 µM), dissolved in water, was added to cells growing in either SD or SD-N (5 µM) 30 min before they were visualized by microscopy. The vacuole was visualized in live cells in (Fig EV2A) using FM4-64 (Invitrogen, T3166) and phenylmethane-sulfonyl-fluoride (PMSF, Sigma P7626) treatment of cells was executed exactly as described (Journé *et al.*, 2008). In order to optimize the visualization of the lowly expressed endogenous Med13-mNeongreen the Keyence microscope was used. This scope has a high sensitivity CCD and high-speed autofocus, low photobleaching mode that aids in monitoring Med13 localization in live cells. Deconvolution and processing capabilities are very limited with the compatible analyzer software. Single plane images were obtained using a Keyence BZ-X710 fluorescence microscope with a 100x objective with 1.0x camera magnification (PlanApoλ Oil, NA 1.45) and a CCD camera. Data were collected using BZ-X Analyzer software. Quantification of Med13-mNeongreen fluorescence within the vacuole was obtained using the Hybrid cell count function within the analyzer software (300 cells were counted

per sample). For analysis single extraction settings were used. Red (vacuole, Vph1-mCherry) was set as the target area and green (Med13-mNeongreen) was set as the extraction area. The percentage of cells with vacuolar Med13-mNeongreen was calculated using Area ratio (1st) (ratio of the total area of the extracted areas to the target area) and cell count values. Percentages represent a ratio of extraction area to the target area.

Statistics

All representative results included at least two independent biological experiments. P values were generated from Prism-GraphPad using unpaired Student's t-tests; NS $P \geq 0.05$; * $P \leq 0.05$, ** $P \leq 0.005$; *** $P \leq 0.001$; **** $P \leq 0.0001$. All error bars indicate mean $\pm$ SD. For quantification of Med13-9myc degradation kinetics band intensities of each time point was first divided by unstressed, T=0 band intensity. These values were then divided by Pgk1 loading band intensity values which were also normalized to their T=0 intensities. P-values shown are relative to wild-type T=4 timepoints.

ACKNOWLEDGMENTS

We thank C. De Virgilio, P. Herman and K. Cunningham for strains and plasmids. We especially thank the members of the R. Strich laboratory and the Cooper laboratory for critical reading of this manuscript.

FUNDING

This work was supported by a grant from the National Institutes of Health awarded to K.F.C. (GM113196).

COMPETING INTERESTS

The authors declare no competing or financial interests.

AUTHOR CONTRIBUTIONS

SEH, SDW and KFC conceived the study and designed experiments. SEH and SDW performed experiments. KFC performed most of the microscopy experiments executed with the Nikon microscope. SEH and KFC wrote the manuscript.

REFERENCES

- Abeliovich H, Zhang C, Dunn WA, Jr., Shokat KM, Klionsky DJ (2003) Chemical genetic analysis of Apg1 reveals a non-kinase role in the induction of autophagy. *Mol Biol Cell* 14: 477-490
- Akoulitchiev S, Chuikov S, Reinberg D (2000) TFIID is negatively regulated by cdk8-containing mediator complexes. *Nature* 407: 102-106.
- Alcazar-Roman AR, Tran EJ, Guo S, Wente SR (2006) Inositol hexakisphosphate and Gle1 activate the DEAD-box protein Dbp5 for nuclear mRNA export. *Nat Cell Biol* 8: 711-716
- Aryanpur PP, Regan CA, Collins JM, Mittelmeier TM, Renner DM, Vergara AM, Brown NP, Bolger TA (2017) Gle1 Regulates RNA Binding of the DEAD-Box Helicase Ded1 in Its Complex Role in Translation Initiation. *Mol Cell Biol* 37
- Bartholomew CR, Suzuki T, Du Z, Backues SK, Jin M, Lynch-Day MA, Umekawa M, Kamath A, Zhao M, Xie Z *et al* (2012) Ume6 transcription factor is part of a signaling cascade that regulates autophagy. *Proc Natl Acad Sci U S A* 109: 11206-11210
- Bean BD, Davey M, Conibear E (2017) Cargo selectivity of yeast sorting nexins. *Traffic* 18: 110-122
- Bernard A, Jin M, Gonzalez-Rodriguez P, Fullgrabe J, Delorme-Axford E, Backues SK, Joseph B, Klionsky DJ (2015) Rph1/KDM4 mediates nutrient-limitation signaling that leads to the transcriptional induction of autophagy. *Curr Biol* 25: 546-555
- Bourbon HM, Aguilera A, Ansari AZ, Asturias FJ, Berk AJ, Bjorklund S, Blackwell TK, Borggrefe T, Carey M, Carlson M *et al* (2004) A Unified Nomenclature for Protein Subunits of Mediator Complexes Linking Transcriptional Regulators to RNA Polymerase II. *Mol Cell* 14: 553-557

492 Chu AM, Davis RW (2008) High-throughput creation of a whole-genome collection of yeast
493 knockout strains. *Methods Mol Biol* 416: 205-220

494 Cooper KF, Khakhina S, Kim SK, Strich R (2014) Stress-induced nuclear-to-cytoplasmic
495 translocation of cyclin C promotes mitochondrial fission in yeast. *Dev Cell* 28: 161-173

496 Cooper KF, Mallory MJ, Smith JB, Strich R (1997) Stress and developmental regulation of the
497 yeast C-type cyclin Ume3p (Srb11p/Ssn8p). *EMBO J* 16: 4665-4675

498 Cooper KF, Mallory MJ, Strich R (1999) Oxidative stress-induced destruction of the yeast C-type
499 cyclin Ume3p requires phosphatidylinositol-specific phospholipase C and the 26S proteasome.
500 *Mol Cell Biol* 19: 3338-3348

501 Cooper KF, Scarnati MS, Krasley E, Mallory MJ, Jin C, Law MJ, Strich R (2012) Oxidative-stress-
502 induced nuclear to cytoplasmic relocalization is required for Not4-dependent cyclin C destruction.
503 *J Cell Sci* 125: 1015-1026

504 Delorme-Axford E, Klionsky DJ (2018) Transcriptional and post-transcriptional regulation of
505 autophagy in the yeast *Saccharomyces cerevisiae*. *J Biol Chem*

506 Deng Y, Qu Z, Naqvi NI (2013) The role of snx41-based pexophagy in magnaporthe
507 development. *PLoS One* 8: e79128

508 Farre JC, Subramani S (2016) Mechanistic insights into selective autophagy pathways: lessons
509 from yeast. *Nat Rev Mol Cell Biol* 17: 537-552

510 Fernandez-Martinez J, Kim SJ, Shi Y, Upla P, Pellarin R, Gagnon M, Chemmama IE, Wang J,
511 Nudelman I, Zhang W *et al* (2016) Structure and Function of the Nuclear Pore Complex
512 Cytoplasmic mRNA Export Platform. *Cell* 167: 1215-1228 e1225

513 Folkmann AW, Noble KN, Cole CN, Wentz SR (2011) Dbp5, Gle1-IP6 and Nup159: a working
514 model for mRNP export. *Nucleus* 2: 540-548

515 Fu N, Yang X, Chen L (2018) Nucleophagy Plays a Major Role in Human Diseases. *Curr Drug*
516 *Targets* 19: 1767-1773

517 Ganesan V, Willis SD, Chang KT, Beluch S, Cooper KF, Strich R (2019) Cyclin C directly
518 stimulates Drp1 GTP affinity to mediate stress-induced mitochondrial hyperfission. *Mol Biol Cell*
519 30: 302-311

520 Gari E, Piedrafita L, Aldea M, Herrero E (1997) A set of vectors with a tetracycline-regulatable
521 promoter system for modulated gene expression in *Saccharomyces cerevisiae*. *Yeast* 13: 837-
522 848

523 Gnanasundram SV, Kos M (2015) Fast protein-depletion system utilizing tetracycline repressible
524 promoter and N-end rule in yeast. *Mol Biol Cell* 26: 762-768

525 Hettema EH, Lewis MJ, Black MW, Pelham HR (2003) Retromer and the sorting nexins
526 Snx4/41/42 mediate distinct retrieval pathways from yeast endosomes. *EMBO J* 22: 548-557

527 Hollenstein DM, Gomez-Sanchez R, Ciftci A, Kriegenburg F, Mari M, Torggler R, Licheva M,
528 Reggiori F, Kraft C (2019) Vac8 spatially confines autophagosome formation at the vacuole in *S.*
529 *cerevisiae*. *J Cell Sci* 132

530 Hollenstein DM, Kraft C (2020) Autophagosomes are formed at a distinct cellular structure. *Curr*
531 *Opin Cell Biol* 65: 50-57

532 Hu YB, Dammer EB, Ren RJ, Wang G (2015) The endosomal-lysosomal system: from
533 acidification and cargo sorting to neurodegeneration. *Transl Neurodegener* 4: 18

534 Humphries CL, Balcer HI, D'Agostino JL, Winsor B, Drubin DG, Barnes G, Andrews BJ, Goode
535 BL (2002) Direct regulation of Arp2/3 complex activity and function by the actin binding protein
536 coronin. *J Cell Biol* 159: 993-1004

537 Hutten S, Kehlenbach RH (2007) CRM1-mediated nuclear export: to the pore and beyond.
538 *Trends Cell Biol* 17: 193-201

539 Janke C, Magiera MM, Rathfelder N, Taxis C, Reber S, Maekawa H, Moreno-Borchart A,
540 Doenges G, Schwob E, Schiebel E *et al* (2004) A versatile toolbox for PCR-based tagging of
541 yeast genes: new fluorescent proteins, more markers and promoter substitution cassettes. *Yeast*
542 21: 947-962

543 Jeronimo C, Langelier MF, Bataille AR, Pascal JM, Pugh BF, Robert F (2016) Tail and Kinase
544 Modules Differently Regulate Core Mediator Recruitment and Function In Vivo. *Mol Cell* 64: 455-
545 466

546 Jeronimo C, Robert F (2017) The Mediator Complex: At the Nexus of RNA Polymerase II
547 Transcription. *Trends Cell Biol* 27: 765-783

548 Jezek J, Chang KT, Joshi AM, Strich R (2019) Mitochondrial translocation of cyclin C stimulates
549 intrinsic apoptosis through Bax recruitment. *EMBO Rep* 20: e47425

550 Journo D, Mor A, Abeliovich H (2009) Aup1-mediated regulation of Rtg3 during mitophagy. *J Biol*
551 *Chem* 284: 35885-35895

552 Journo D, Winter G, Abeliovich H (2008) Monitoring autophagy in yeast using FM 4-64
553 fluorescence. *Methods Enzymol* 451: 79-88

554 Kamada Y, Funakoshi T, Shintani T, Nagano K, Ohsumi M, Ohsumi Y (2000) Tor-mediated
555 induction of autophagy via an Apg1 protein kinase complex. *J Cell Biol* 150: 1507-1513

556 Kanki T, Wang K, Baba M, Bartholomew CR, Lynch-Day MA, Du Z, Geng J, Mao K, Yang Z, Yen
557 WL *et al* (2009) A genomic screen for yeast mutants defective in selective mitochondria
558 autophagy. *Mol Biol Cell* 20: 4730-4738

559 Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJ (2015) The Phyre2 web portal for
560 protein modeling, prediction and analysis. *Nat Protoc* 10: 845-858

561 Khakhina S, Cooper KF, Strich R (2014) Med13p prevents mitochondrial fission and programmed
562 cell death in yeast through nuclear retention of cyclin C. *Mol Biol Cell* 25: 2807-2816

563 Klionsky DJ, Codogno P (2013) The mechanism and physiological function of macroautophagy. *J*
564 *Innate Immun* 5: 427-433

565 Lee CW, Wilfling F, Ronchi P, Allegretti M, Mosalaganti S, Jentsch S, Beck M, Pfander B (2020)
566 Selective autophagy degrades nuclear pore complexes. *Nat Cell Biol* 22: 159-166

567 Li J, Kim SG, Blenis J (2014) Rapamycin: one drug, many effects. *Cell Metab* 19: 373-379

568 Ma M, Burd CG (2020) Retrograde trafficking and plasma membrane recycling pathways of the
569 budding yeast *Saccharomyces cerevisiae*. *Traffic* 21: 45-59

570 Ma M, Burd CG, Chi RJ (2017) Distinct complexes of yeast Snx4 family SNX-BARs mediate
571 retrograde trafficking of Snc1 and Atg27. *Traffic* 18: 134-144

572 Ma M, Kumar S, Purushothaman L, Babst M, Ungermann C, Chi RJ, Burd CG (2018) Lipid
573 trafficking by yeast Snx4 family SNX-BAR proteins promotes autophagy and vacuole membrane
574 fusion. *Mol Biol Cell* 29: 2190-2200

575 Matscheko N, Mayrhofer P, Rao Y, Beier V, Wollert T (2019) Atg11 tethers Atg9 vesicles to
576 initiate selective autophagy. *PLoS Biol* 17: e3000377

577 Millen JI, Krick R, Prick T, Thumm M, Goldfarb DS (2009) Measuring piecemeal microautophagy
578 of the nucleus in *Saccharomyces cerevisiae*. *Autophagy* 5: 75-81

579 Mochida K, Oikawa Y, Kimura Y, Kirisako H, Hirano H, Ohsumi Y, Nakatogawa H (2015)
580 Receptor-mediated selective autophagy degrades the endoplasmic reticulum and the nucleus.
581 *Nature* 522: 359-362

582 Mosammaparast N, Pemberton LF (2004) Karyopherins: from nuclear-transport mediators to
583 nuclear-function regulators. *Trends Cell Biol* 14: 547-556

584 Murphy R, Wente SR (1996) An RNA-export mediator with an essential nuclear export signal.
585 *Nature* 383: 357-360

586 Nagulapalli M, Maji S, Dwivedi N, Dahiya P, Thakur JK (2016) Evolution of disorder in Mediator
587 complex and its functional relevance. *Nucleic Acids Res* 44: 1591-1612

588 Nemec AA, Howell LA, Peterson AK, Murray MA, Tomko RJ, Jr. (2017) Autophagic clearance of
589 proteasomes in yeast requires the conserved sorting nexin Snx4. *J Biol Chem* 292: 21466-21480

590 Nemet J, Jelacic B, Rubelj I, Sopta M (2014) The two faces of Cdk8, a positive/negative regulator
591 of transcription. *Biochimie* 97: 22-27

592 Nice DC, Sato TK, Stromhaug PE, Emr SD, Klionsky DJ (2002) Cooperative binding of the
593 cytoplasm to vacuole targeting pathway proteins, Cvt13 and Cvt20, to phosphatidylinositol 3-

594 phosphate at the pre-autophagosomal structure is required for selective autophagy. *J Biol Chem*
595 277: 30198-30207

596 Papandreou ME, Tavernarakis N (2019) Nucleophagy: from homeostasis to disease. *Cell Death*
597 *Differ* 26: 630-639

598 Popelka H, Damasio A, Hinshaw JE, Klionsky DJ, Ragusa MJ (2017) Structure and function of
599 yeast Atg20, a sorting nexin that facilitates autophagy induction. *Proc Natl Acad Sci U S A* 114:
600 E10112-E10121

601 Ramos PC, Hockendorff J, Johnson ES, Varshavsky A, Dohmen RJ (1998) Ump1p is required for
602 proper maturation of the 20S proteasome and becomes its substrate upon completion of the
603 assembly. *Cell* 92: 489-499

604 Roberts P, Moshitch-Moshkovitz S, Kvam E, O'Toole E, Winey M, Goldfarb DS (2003) Piecemeal
605 microautophagy of nucleus in *Saccharomyces cerevisiae*. *Mol Biol Cell* 14: 129-141

606 Ronne H, Rothstein R (1988) Mitotic sectored colonies: evidence of heteroduplex DNA formation
607 during direct repeat recombination. *Proc Natl Acad Sci U S A* 85: 2696-2700

608 Shintani T, Klionsky DJ (2004) Cargo proteins facilitate the formation of transport vesicles in the
609 cytoplasm to vacuole targeting pathway. *J Biol Chem* 279: 29889-29894

610 Shpilka T, Welter E, Borovsky N, Amar N, Shimron F, Peleg Y, Elazar Z (2015) Fatty acid
611 synthase is preferentially degraded by autophagy upon nitrogen starvation in yeast. *Proc Natl*
612 *Acad Sci U S A* 112: 1434-1439

613 Snyder NA, Kim A, Kester L, Gale AN, Studer C, Hoepfner D, Roggo S, Helliwell SB,
614 Cunningham KW (2019) Auxin-Inducible Depletion of the Essentialome Suggests Inhibition of
615 TORC1 by Auxins and Inhibition of Vrg4 by SDZ 90-215, a Natural Antifungal Cyclopeptide. *G3*
616 *(Bethesda)* 9: 829-840

617 Stanishneva-Konovalova TB, Derkacheva NI, Polevova SV, Sokolova OS (2016) The Role of
618 BAR Domain Proteins in the Regulation of Membrane Dynamics. *Acta Naturae* 8: 60-69

619 Stieg DC, Willis SD, Ganesan V, Ong KL, Scuorzo J, Song M, Grose J, Strich R, Cooper KF
620 (2018) A complex molecular switch directs stress-induced cyclin C nuclear release through
621 SCF(Grr1)-mediated degradation of Med13. *Mol Biol Cell* 29: 363-375

622 Strich R, Slater MR, Esposito RE (1989) Identification of negative regulatory genes that govern
623 the expression of early meiotic genes in yeast. *Proc Natl Acad Sci USA* 86: 10018-10022

624 Suzuki SW, Emr SD (2018) Membrane protein recycling from the vacuole/lysosome membrane. *J*
625 *Cell Biol* 217: 1623-1632

626 Takeshige K, Baba M, Tsuboi S, Noda T, Ohsumi Y (1992) Autophagy in yeast demonstrated
627 with proteinase-deficient mutants and conditions for its induction. *J Cell Biol* 119: 301-311

628 Tsai KL, Sato S, Tomomori-Sato C, Conaway RC, Conaway JW, Asturias FJ (2013) A conserved
629 Mediator-CDK8 kinase module association regulates Mediator-RNA polymerase II interaction. *Nat*
630 *Struct Mol Biol* 20: 611-619

631 Uversky VN (2011) Multitude of binding modes attainable by intrinsically disordered proteins: a
632 portrait gallery of disorder-based complexes. *Chem Soc Rev* 40: 1623-1634

633 Van Den Hazel HB, Kielland-Brandt MC, Winther JR (1996) Review: biosynthesis and function of
634 yeast vacuolar proteases. *Yeast* 12: 1-16

635 Vlahakis A, Lopez Muniozguren N, Powers T (2017) Stress-response transcription factors Msn2
636 and Msn4 couple TORC2-Ypk1 signaling and mitochondrial respiration to ATG8 gene expression
637 and autophagy. *Autophagy* 13: 1804-1812

638 Wang K, Yan R, Cooper KF, Strich R (2015) Cyclin C mediates stress-induced mitochondrial
639 fission and apoptosis. *Mol Biol Cell* 26: 1030-1043

640 Wang R, Solomon MJ (2012) Identification of She3 as an SCF(Grr1) substrate in budding yeast.
641 *PLoS One* 7: e48020

642 Wanke V, Pedruzzi I, Camerini E, Dubouloz F, De Virgilio C (2005) Regulation of G0 entry by the
643 Pho80-Pho85 cyclin-CDK complex. *EMBO J* 24: 4271-4278

644 Weirich CS, Erzberger JP, Flick JS, Berger JM, Thorner J, Weis K (2006) Activation of the
645 DExD/H-box protein Dbp5 by the nuclear-pore protein Gle1 and its coactivator InsP6 is required
646 for mRNA export. *Nat Cell Biol* 8: 668-676

647 Welter E, Thumm M, Krick R (2010) Quantification of nonselective bulk autophagy in *S.*
648 *cerevisiae* using Pgk1-GFP. *Autophagy* 6: 794-797

649 Willis SD, Hanley SE, Beishke T, Tati PD, Cooper KF (2020) Ubiquitin-proteasome-mediated
650 cyclin C degradation promotes cell survival following nitrogen starvation. *Mol Biol Cell* 31: 1015-
651 1031

652 Willis SD, Stieg DC, Ong KL, Shah R, Strich AK, Grose JH, Cooper KF (2018) Snf1 cooperates
653 with the CWI MAPK pathway to mediate the degradation of Med13 following oxidative stress.
654 *Microbial Cell* 5: 357-370

655 Yoshida K, Blobel G (2001) The karyopherin Kap142p/Msn5p mediates nuclear import and
656 nuclear export of different cargo proteins. *J Cell Biol* 152: 729-740

657 Zhang H, Chen J, Wang Y, Peng L, Dong X, Lu Y, Keating AE, Jiang T (2009) A computationally
658 guided protein-interaction screen uncovers coiled-coil interactions involved in vesicular trafficking.
659 *J Mol Biol* 392: 228-241

660 Zhang H, Huang T, Hong Y, Yang W, Zhang X, Luo H, Xu H, Wang X (2018) The Retromer
661 Complex and Sorting Nexins in Neurodegenerative Diseases. *Front Aging Neurosci* 10: 79

662 Zhu J, Deng S, Lu P, Bu W, Li T, Yu L, Xie Z (2016) The Ccl1-Kin28 kinase complex regulates
663 autophagy under nitrogen starvation. *J Cell Sci* 129: 135-144

664 Zientara-Rytter K, Subramani S (2020) Mechanistic Insights into the Role of Atg11 in Selective
665 Autophagy. *J Mol Biol* 432: 104-122

666 Zubenko GS, Park FJ, Jones EW (1983) Mutations in PEP4 locus of *Saccharomyces cerevisiae*
667 block final step in maturation of two vacuolar hydrolases. *Proc Natl Acad Sci U S A* 80: 510-514
668
669

FIGURE LEGENDS

Figure 1. Med13 is degraded via the vacuolar proteolysis following nitrogen starvation.

A Model outlining how the cyclin dependent kinase module (CKM) of the mediator complex is disassembled following stresses that mediate cell death or survival pathways. Before stress cyclin C (CC), Cdk8 (8), Med13 (13) and Med12 (12) form the CKM that predominantly represses stress response genes (SRGs) (Cooper *et al.*, 1997; Cooper *et al.*, 1999). Repression is relieved after oxidative stress and nitrogen depletion by CKM disassembly, mediated by different mechanisms. After oxidative stress, Med13 is destroyed by the UPS (Khakhina *et al.*, 2014; Stieg *et al.*, 2018), which allows cyclin C localization to the mitochondria where it triggers stress-induced mitochondrial fission and promotes cell death (Cooper *et al.*, 2014). Following nitrogen starvation, cyclin C is rapidly destroyed by the UPS before its nuclear release to prevent mitochondrial fission (Willis *et al.*, 2020) whereas the fate of Med13 is the subject of this current manuscript. N-nucleus, C-cytoplasm.

B Western blot analysis of extracts prepared from wild-type cells expressing endogenous Med13-9xMyc (RSY2211) resuspended in nitrogen starvation medium (SD-N) or treated with 200 ng/ml rapamycin for the indicated times.

C As in B except that endogenous Med13 protein levels (13Myc) were monitored in *ump1Δ* (RSY1961) and *pep4Δ prb1Δ.1* (RSY2215) strains.

D Degradation kinetics and half-life of Med13 protein levels obtained in B and C. Error bars indicate S.D., N=3 of biologically independent experiments.

E Wild-type (RSY10) or *pep4Δ prb1Δ.1* (RSY449) cells expressing Med13-GFP (pSW218) were starved for nitrogen for indicated times. For all cleavage assays, free GFP refers to the protease resistant GFP moiety that accumulates after the full-length fusion protein is degraded via the vacuole. GFP accumulation was monitored by Western blot analysis using anti-GFP antibodies. An asterisk indicates a nonspecific proteolytic fragment unrelated to autophagy.

F As in E except that wild-type cells were resuspended in SD-N or treated with 0.8 mM H₂O₂. Pgk1 protein levels were used as a loading control.

Figure 2. Med13 translocates from the nucleus to the vacuole in nitrogen starvation.

A Endogenous Med13-mNeongreen localization was monitored in *pep4Δ prb1Δ.1* cells (RSY2305) expressing Nab2-mCherry (a nuclear marker) before (growing in SD) and after 4 h in SD-N. Representative single plane images are shown.

B As in A, except that cells expressed a vacuolar marker (Vph1-mCherry).

C As in B, except that the slices were taken through the whole cell which were then collapsed and deconvolved. Representative deconvolved images are shown. Scale bar = 5 μm.

Figure 3. Med13 requires the core autophagy machinery for vacuolar degradation.

A Schematic depicting the five stages of autophagy and the major Atg protein complexes associated with each stage.

B Western blot analysis of extracts prepared from wild-type (RSY2211), *atg1Δ* (RSY2214) and *atg8Δ* (RSY2231) cells expressing endogenous Med13-9Myc resuspended in SD-N media for the indicated times.

C Degradation kinetics and half-life of Med13 protein levels obtained in B. Error bars indicate S.D., N=3 of biologically independent experiments.

D and E The indicated mutants expressing Med13-GFP (pSW218) were starved for nitrogen for the indicated times and accumulation of free GFP monitored by Western blot analysis using anti-GFP antibodies. An asterisk indicates a nonspecific proteolytic fragment unrelated to autophagy. For all experiments, Pgk1 protein levels were used as a loading control.

Figure 4. Autophagic degradation of Med13 requires the nucleoporin Gle1 and is independent of known nucleophagy pathways.

A Western blot analysis of Med13-GFP cleavage assays after 4 h nitrogen depletion in micro-nucleophagy (*nvj1Δ*, RSY2106) and macro-nucleophagy (*atg39Δ atg40Δ*, RSY2123) mutants. An asterisk indicates a nonspecific proteolytic fragment unrelated to autophagy and Pgk1 protein levels were used as a loading control.

B Co-immunoprecipitation analysis of endogenous Gle1-GFP and Med13-3HA. Whole cell lysates were immunoprecipitated with the antibodies shown from nitrogen-starved *pep4Δ prb1Δ.1* cells expressing endogenous Gle1-GFP (RSY2423) and Med13-3HA (pKC801, lanes 1 and 4) or a vector control (lane 2). *pep4Δ prb1Δ.1* cells expressing Med13-3HA alone (lane 3) was included as a control. [] represents no antibody control. Med13-HA was detected by Western blot analysis of immunoprecipitates. Western blot analysis of the proteins in the whole cell lysates for the three conditions tested is shown (input - bottom panel).

C Map of Med13 depicting different structural regions and known interacting proteins (upper panel). Different colors represent different regions of the protein and structural regions are denoted by amino acid positions. Med13-Gle1 Y2H analysis. Y2H Gold cells harboring Gal4-BD-Gle1 and the indicated Gal4-AD-Med13 subclone or empty vector control were streaked on medium selecting for plasmid maintenance (left) or induction of the *ADE2* and *HIS3* reporter genes (right) by Y2H interaction (lower panel). See Fig EV5A, B for Western blot analysis of the different constructs.

D *pep4Δ prb1Δ.1* cells expressing endogenous Med13-mNeongreen and endogenous Gle1-RedStar (RSY2450) were starved for nitrogen for 2 h and monitored by fluorescence microscopy. Deconvolved representative images are shown. The yellow arrow is pointing to Med13-GFP in the vacuole whereas the blue arrow shows co-localization. OE represents an over-exposed image to better show the colocalization of Gle1 and Med13 (pink arrows) Scale = 5μm.

E Med13-GFP cleavage assays performed in the Gle1 Auxin-inducible degron (Gle1-AID) strain (RSY2456). Cells expressing Med13-GFP (pSW320) were treated with 250 μM Auxin for 30 m

before proceeding with autophagic cleavage assays in SD-N. Free GFP accumulation was then detected using Western blot analysis.

F Fluorescence microscopy of Med13-mNeongreen localization in the Gle1 Auxin-inducible degron (Gle1-AID) strain expressing Nab2-mCherry before and after SD-N. An 8 h time point was used as the strain background is BY4741 which is not as sensitive to environmental stress as W303a. Scale = 5 μ m.

Figure 5. The sorting nexin heterodimer, Snx4-Atg20 is required for efficient autophagic degradation of Med13.

A Western blot analysis of Med13-GFP cleavage assays after 4 h nitrogen starvation in wild-type and *snx4 Δ* (RSY2272).

B Co-immunoprecipitation analysis of GFP-Snx4 and Med13-3HA. Whole cell lysates were immunoprecipitated with the antibodies shown from nitrogen-starved *pep4 Δ prb1 Δ .1* cells expressing GFP-Snx4 (RSY2299) and Med13-3HA (pKC801, lanes 1 and 2) or a vector control (lane 4). *Pep4 Δ prb1 Δ .1* cells expressing Med13-3HA alone (lane 3) was included as a control. [] represents no antibody control. Med13-HA was detected by Western blot analysis of immunoprecipitates. Western blot analysis of the proteins in the whole cell lysates for the three conditions tested is shown (input - bottom panel).

C Western blot analysis of extracts prepared from wild-type (RSY2211), *snx4 Δ* (RSY2276), *atg20 Δ* (RSY2277) and *snx41 Δ* (RSY2394) expressing endogenous Med13-9Myc resuspended in SD-N for the indicated times. Pgk1 was used as a loading control.

D Degradation kinetics and half-lives of Med13 protein levels obtained in B. Error bars indicate S.D., N=3 of biologically independent experiments.

E Fluorescence microscopy of endogenous Med13-mNeongreen localization in *snx4 Δ pep4 Δ prb1 Δ .1* (RSY2324) expressing the vacuole marker Vph1-mCherry. Cells were visualized before

(SD) and after 2 h of SD-N treatment and representative single plane images are shown. Scale = 5 μ m.

F As in D except that endogenous Med13-mNeongreen localization was followed in nitrogen-starved *pep4 Δ prb1 Δ .1* cells. Representative single plane images of the results are shown. Bar = 5 μ m.

G Quantification of accumulation of Med13-mNeongreen in vacuoles obtained from results in D and E. 100 cells counted per sample. N=3 biological samples. **** $P = >0.0001$.

Figure 6. Snx4 localizes to the nuclear periphery to retrieve Med13.

A Co-immunoprecipitation analysis of endogenous Atg17-GFP and Med13-3HA in the presence and absence of Snx4. Whole cell lysates were immunoprecipitated with the antibodies shown from nitrogen-starved *pep4 Δ prb1 Δ .1* (RSY2395) or *snx4 Δ pep4 Δ prb1 Δ .1* cells (RSY2396) expressing endogenous Atg17-GFP and Med13-3HA (pKC801, lanes 1, 2 and 6) or a vector control (lanes 4 & 5). *Pep4 Δ prb1 Δ .1* cells expressing Med13-3HA alone (lane 3) was included as a control. [] represents no antibody control. Med13-HA was detected by Western blot analysis of immunoprecipitates. Western blot analysis of the proteins present in the whole cell lysates for the conditions tested is shown (input - bottom panel).

B Representative images showing perinuclear and perivacuolar localization of GFP-Snx4 in wild-type expressing Nab2-mCherry (nuclear marker) before and after nitrogen starvation. Bar = 5 μ m. The number of perinuclear foci was counted (N=2) before and after nitrogen starvation. At least 100 cells were counted per sample. Data are the percentage of perinuclear foci among the total number of foci. Scale = 5 μ m.

C Fluorescence microscopy of *pep4 Δ prb1 Δ .1* cells expressing endogenous Med13-mNeongreen and mCherry-Snx4 (RSY2424) before and after nitrogen depletion. Hoechst staining was used to visualize the nucleus. Representative deconvolved images are shown. Bar = 5 μ m.

D GFP-Snx4 and endogenous Gle1-RedStar co-localize in wild-type cells (RSY2451) following nitrogen starvation. Representative images are shown. Bar = 5µm.

Figure 7. The BAR domain of Snx4 interacts with the C-terminal region of Med13.

A Fluorescence microscopy of Crn1-Med13-GFP (pSW288) in wild-type cells growing in SD. Hoechst staining was used to visualize the nucleus. Bar = 5µm.

B Western blot analysis of Med13-GFP (pSW218) or Crn1-Med13-GFP (pSW288) cleavage assays performed in wild-type or *snx4Δ* cells following nitrogen starvation.

C Western blot analysis of GFP-Snx4 cleavage assays performed in wild-type (RSY2283) and *pep4Δ* (RSY2299) cells following nitrogen starvation.

D Western blot analysis of Gle1-GFP cleavage assays performed in wild-type (RSY2455) cells following prolonged nitrogen starvation (24 h).

E Western blot analysis of Med13-GFP (pSW218) cleavage assays performed in wild-type cells. Pgk1 was used as a protein loading control for all experiments.

F Y2H Gold cells harboring Gal4-BD-Snx4 and the indicated Gal4-AD-Med13 construct or vector control were plated on medium selecting for plasmid maintenance (-LEU, -TRP) (top) or interaction by induction of the *ADE2* and *HIS3* reporter genes (bottom). See Fig EV5A, B for Western blot analysis of the different constructs.

G Predicted structural analysis of the Med13 tail region using Phyre2 plot analysis of this region (Kelley *et al.*, 2015). Y2H analysis of full-length Gal4-BD-Snx4 and Gal4-AD-Med13 subclones containing either the first or second domain region of the Med13 tail. Cells were streaked on medium selecting for plasmid maintenance (left) or induction of reporter genes (right) by Y2H interaction.

H Map of Snx4 depicting known domains (left panel). Y2H analysis of Snx4 PX and BAR binding domain constructs with the Med13⁹⁰⁷⁻¹¹⁶³AD construct and empty vector. Cells were streaked on

medium selecting for plasmid maintenance (left) or induction of reporter genes (right) by Y2H interaction (right panel).

Figure 8. Transcription factors regulating ATG genes are nucleophagy substrates.

A RT-qPCR analysis probing for *ATG8*, *ATG1* and *ATG14* mRNA expression in wild-type and *med13Δ* (RSY2444) cells in unstressed conditions. $\Delta\Delta C_t$ results for relative fold change ($\log_2$) values using wild-type unstressed cells as a control. Transcript levels are given relative to the internal *ACT1* mRNA control.

B Western blot analysis of Rim15-GFP (pFD846) cleavage assays in indicated mutants after nitrogen starvation. The asterisk denotes a background band.

C Western blot analysis of Ccl1-GFP (pSW230) and Msn2-GFP (pSW217) cleavage assays in wild-type or *snx4Δ* cells. Pgk1 protein levels were used as a loading control in all experiments.

D Summary model of novel Snx4-mediated nucleophagy mechanism that selectively targets transcription factors for autophagic degradation. In unstressed cells the CKM represses a subset of *ATG* genes. Following nitrogen starvation an unknown signal triggers the CKM to disassembly. Med13 is transported by unknown mechanisms through the NPC to the cytoplasmic nucleoporin Gle1. Gle1 releases Med13 to the Snx4-Atg20 heterodimer where it interacts with the Snx4 BAR domains (see insert), which in turn interacts with the Atg17 scaffold protein found on growing PAS structures.

844

Figure 1

845

Figure 2

Figure 3

846

847

848

Figure 4

Figure 5

850

Figure 6

851

Figure 7

852

853

855 **Supplementary Information**

856 **Supplemental Table I.** Candidate proteins that interact with Med13 after 90 min 200mM rapamycin
857 treatment.

Select proteins obtained from Mass Spectroscopy where: Counts were greater than 2-fold after 90 minutes of 200 ng/mL Rapamycin			Counts	
Systematic Name	Gene	Protein Function	T=0	T=90
Other				
YER155C	<i>BEM2</i>	RhoGAP	47	203
YJL005W	<i>CYR1</i>	Adenylate cyclase	19	50
YHR016C	<i>YSC84</i>	Actin-binding protein	7	18
YNL271C	<i>BNI1</i>	Formin	5	13
YBR140C	<i>IRA1</i>	GTPase activator	1	12
YDR103W	<i>STE5</i>	MAPK scaffold	4	12
YGR032W	<i>GSC2</i>	spore wall assembly	1	9
YDL198C	<i>GGC1</i>	Mitochondrial GTP/GDP transporter	3	7
Nuclear Activity				
YPR018W	<i>RLF2</i>	Chromatin assembly	21	44
YPL155C	<i>KIP2</i>	Kinesin	6	34
YDL140C	<i>RPO21</i>	RNA Polymerase II Subunit	13	33
YML065W	<i>ORC1</i>	DNA replication	14	32
YNL278W	<i>CAF120</i>	Transcriptional regulation	7	27
YDR216W	<i>ADR1</i>	Transcription factor	12	26

YER040W	<i>GLN3</i>	Transcriptional activator	10	23
YDR364C	<i>CDC40</i>	mRNA Splicing	7	15
YGL045W	<i>RIM8</i>	Activates Rim101	6	13
YER088C	<i>DOT6</i>	DNA Binding Protein	3	12
YDL089W	<i>NUR1</i>	Mitotic Exit/Nuclear Periphery	3	9
YDL207W	<i>GLE1</i>	Cytoplasmic nucleoporin	3	9
YDL220C	<i>CDC13</i>	Telomere capping	4	9
Vacuole				
YJL172W	<i>CPS1</i>	Vacuolar carboxypeptidase	4	18
YPR036W	<i>VMA13</i>	V-ATPase	4	12
YDR128W	<i>MTC5</i>	Vacuole/TORC1	3	8
YLL048C	<i>YBT1</i>	Vacuole Fusion	2	8
Kinase/Phosphatase				
YDR283C	<i>GCN2</i>	Kinase	78	160
YHR082C	<i>KSP1</i>	Kinase	25	60
YKL168C	<i>KKQ8</i>	Kinase	13	36
YDR122W	<i>KIN1</i>	Kinase	4	28
YOR039W	<i>CKB2</i>	Kinase	8	26
YDL025C	<i>RTK1</i>	Kinase	3	18
YOR061W	<i>CKA2</i>	Kinase	5	15
YGL019W	<i>CKB1</i>	Kinase	4	13
YMR165C	<i>PAH1</i>	Phosphatase	4	13
YIL035C	<i>CKA1</i>	Kinase	5	12
YAR014C	<i>BUD14</i>	Protein phosphatase regulator	3	7
Anabolism/Catabolism				

YLR044C	<i>PDC1</i>	Pyruvate decarboxylase	57	200
YKL182W	<i>FAS1</i>	Fatty acid synthetase	35	94
YKL060C	<i>FBA1</i>	Glycolysis	26	62
YEL071W	<i>DLD3</i>	Lactate Dehydrogenase	16	40
YBR196C	<i>PGI1</i>	Phosphoglucose isomerase	3	29
YLR304C	<i>ACO1</i>	Aconitase	10	25
YLR153C	<i>ACS2</i>	Acetyl-coA synthetase	8	24
YOR136W	<i>IDH2</i>	Isocitrate dehydrogenase complex	7	22
YBR208C	<i>DUR1,2</i>	Urea amidolyase	0	21
YLR355C	<i>ILV5</i>	Amino acid biosynthesis	1	19
YGL009C	<i>LEU1</i>	Leucine biosynthesis	7	17
YKL218C	<i>SRY1</i>	Amino Acid Catabolism	2	15
YGL245W	<i>GUS1</i>	tRNA synthetase	2	15
YER024W	<i>YAT2</i>	Acetyltransferase/mitochondria	1	11
YBR221C	<i>PDB1</i>	Pyruvate Dehydrogenase	3	10
YDR341C	YDR341C	Arginyl-tRNA synthetase/mitochondria	1	9
YIL128W	<i>MET18</i>	Methionine biosynthesis/CIA Complex	2	9
YGL001C	<i>ERG26</i>	Ergosterol biosynthesis	3	7
YJL060W	<i>BNA3</i>	Aminotransferase	3	7
Protein Trafficking				
YNL287W	<i>SEC21</i>	COPI vesicle coat	9	23

YBR102C	<i>EXO84</i>	ExoCytosis/Splicesome Assembly	3	19
YBL037W	<i>APL3</i>	Intracellular protein transport	5	13
YNL049C	<i>SFB2</i>	COPII vesicle coat	5	12
YNL246W	<i>VPS75</i>	Histone chaperone	4	12
YHR103W	<i>SBE22</i>	Golgi /Bud Growth	2	11
YLR114C	<i>AVL9</i>	Exocytic transport	3	9
YDR483W	<i>KRE2</i>	Golgi mannosyltransferase	2	8
YJL036W	<i>SNX4</i>	Sorting Nexin/CVT	2	8
YDR171W	<i>HSP42</i>	Chaperone	3	7
YDR186C	<i>SND1</i>	ER Targeting	3	7
Ribosome				
YPL009C	YPL009C	Ribosome quality control complex	14	63
YKL014C	<i>URB1</i>	Ribosome biogenesis	20	47
YGR162W	<i>TIF4631</i>	Translation initiation factor	12	37
YDR333C	<i>RQC1</i>	Ribosome Quality Control	13	31
YMR049C	<i>ERB1</i>	Ribosome Biogenesis	13	28
YJR041C	<i>URB2</i>	Ribosome biogenesis	1	11
YNL224C	<i>SQS1</i>	Ribosome Biogenesis	4	9
Unknown Function				
YPL032C	<i>SVL3</i>	Unknown Function/Vacuole	16	59
YER047C	<i>SAP1</i>	Unknown/AAA ATPase	21	53
YGR237C	YGR237C	Unknown Function	11	28
YLR187W	<i>SKG3</i>	Unknown Function	9	19

YLR219W	<i>MSC3</i>	Unknown Function	8	18
YBR007C	<i>DSF2</i>	Unknown/Bud Tip	5	15
YOR093C	YOR093C	Unknown Function	6	15
YBR225W	YBR225W	Unknown/Cell Wall	5	13
YDR251W	<i>PAM1</i>	Unknown Function	2	13
YGR196C	<i>FYV8</i>	Unknown Function	4	11
YDR239C	YDR239C	Unknown Function/Ribosome	2	9
Transporters/Permeases				
YDR345C	<i>HXT3</i>	glucose transporter	7	22
YKR093W	<i>PTR2</i>	Peptide Transporter	6	21
YKR039W	<i>GAP1</i>	Amino Acid Permease	4	20
YJR152W	<i>DAL5</i>	Allantoate permease	0	15
YCL025C	<i>AGP1</i>	amino acid permease	4	11
Ubiquitin				
YBL067C	<i>UBP13</i>	Ubiquitin Protease	7	19
YEL012W	<i>UBC8</i>	Ubiquitin Conjugating	7	16
YDL126C	<i>CDC48</i>	ATPase/Ubiquitin Binding	6	15
YDL122W	<i>UBP1</i>	Ubiquitin Protease	3	11
YDR328C	<i>SKP1</i>	Ubiquitin Ligase	3	9
YML088W	<i>UFO1</i>	F-box receptor protein	2	8

858

859

860 **Supplemental Table II** Yeast strains used in this study.

Strain*	Genotype	Source
RSY10*	<i>MATa ade2 ade6 can1-10 his3-11, 15 leu2-3, 112 trp1-1 ura3-1</i>	(Strich <i>et al.</i> , 1989)
RSY1812*	<i>MED13-yECitrine::KanMX6</i>	(Khakhina <i>et al.</i> , 2014)
BJ5459**/RSY449	<i>pep4::HIS3 prb1-Δ1.6R</i>	(Zubenko <i>et al.</i> , 1983)
RSY1961*	<i>ump1::HIS3 MED13-13Myc::KanMX4</i>	(Khakhina <i>et al.</i> , 2014)
RSY2000	<i>gal4Δ gal80Δ LYS2::GAL1-HIS3 GAL2-Ade2 URA3-MEL1 AUR1-C MEL1</i>	Clone Tech Y2H Gold Yeast Strain Cat. #630498
RSY2094*	<i>atg1::KanMX4</i>	This study
RSY2104*	<i>atg17::KanMX4</i>	This study
RSY2106*	<i>nvj1::KanMX4</i>	This study
RSY2160*	<i>ump1::KanMX4</i>	(Willis <i>et al.</i> , 2020)
RSY2123*	<i>atg39::KanMX4 atg40Δ::NatNT2</i>	This study
RSY2202*	<i>atg1::KanMX4 ump1::NatNT2</i>	This study
RSY2211*	<i>MED13-9Myc::NatNT2</i>	This study
RSY2213*	<i>nvj1::KanMX4 MED13-9Myc::NatNT2</i>	This study
RSY2214*	<i>atg1::KanMX4 MED13-9Myc::NatNT2</i>	This study
RSY2215**	<i>pep4::HIS3 prb1-Δ1.6R MED13-9Myc::NatNT2</i>	This study

RSY2231*	<i>atg8::KanMX4 MED13-9Myc::NatNT2</i>	This study
RSY2248*	<i>atg11::KanMX4</i>	This study
Rsy2272*	<i>snx4::HphNT1</i>	This study
RSY2276*	<i>snx4::HphNT1 MED13-9Myc::NatNT2</i>	This study
RSY2277*	<i>atg20::KanMX4 MED13-9Myc::NatNT2</i>	This study
RSY2283*	<i>ADH-GFP-SNX4::NatNT2</i>	This study
RSY2299*	<i>pep4:: KanMX4 ADH-GFP-SNX4::NatNT2</i>	This study
RSY2305**	<i>pep4::HIS3 prb1-Δ1.6R</i> <i>MED13-mNeogreen::NatNT2</i>	This study
RSY2324**	<i>pep4::HIS3 prb1-Δ1.6R snx4Δ::HphNT1</i> <i>MED13-mNeogreen::NatNT2</i>	This study
RSY2348*	<i>PTetO7-Ubi-Leu::3HA-CRM1::NatMX4</i> <i>MED13-9Myc::HphNT1</i>	This study
RSY2349*	<i>atg39::KanMX4 atg40::NatNT2 MED13-9Myc::HphNT1</i>	This study
RSY2373*	<i>atg17::KanMX4 MED13-9Myc::NatNT2</i>	This study
RSY2394*	<i>snx41::HIS3 MED13-9Myc::NatNT2</i>	This study
RSY2395**	<i>pep4::HIS3 prb1-Δ1.6R ATG17-GFP::HphNT1</i>	This study
RSY2396**	<i>pep4::HIS3 prb1-Δ1.6R snx4::KanMX4</i>	This study

	<i>ATG17-GFP::HphNT1</i>		
RSY2399*	<i>msn5::KanMX4 MED13-9Myc::NatNT2</i>		This study
RSY2400**	<i>pep4::HIS3 prb1-Δ1.6R MED13-mNeongreen::HphNT1</i>		This study
	<i>RedStar::NatNT2</i>	<i>ATG17-</i>	
RSY2423**	<i>pep4::HIS3 prb1-Δ1.6R GLE1-GFP::HphNT1</i>		This study
RSY2424**	<i>pep4::HIS3 prb1-Δ1.6R MED13-mNeongreen::HphNT1</i>		This study
	<i>mCherry-SNX4::NatNT2</i>	<i>ADH1-</i>	
RSY2444*	<i>med13::HIS3</i>		This study
RSY2450**	<i>pep4::HIS3 prb1-Δ1.6R MED13-mNeongreen::NatNT2</i>		This study
	<i>RedStar::KanMX4</i>	<i>GLE1-</i>	
RSY2451*	<i>ADH1-GFP-SNX4::NatNT2</i>		This study
	<i>RedStar::KanMX4</i>	<i>GLE1-</i>	
RSY2455*	<i>GLE1-GFP::HphNT1</i>		This study
RSY2456	<i>his3Δ1 leu2Δ met15Δ ura3Δ GLE1-TAP-AID-FLAG::Ura3</i>		(Snyder et al., 2019)

RSY2473	<i>his3Δ1 leu2Δ met15Δ ura3Δ GLE1-TAP-AID-</i>	This study
	<i>FLAG::Ura3 pep4::KanMX4 MED13-</i>	
	<i>mNeonGreen::NatNT2</i>	
BY4741	<i>MATa his3Δ1 leu2Δ met15Δ ura3Δ</i>	(Chu & Davis, 2008)
	<i>atg4::KANMX4</i>	
	<i>atg9::KANMX4</i>	
	<i>atg19:: KANMX4</i>	
	<i>atg32::KANMX4</i>	
	<i>atg36:: KANMX4</i>	
	<i>cue5::KANMX4</i>	
	<i>vps34::KANMX4</i>	

861 *Genotype of all stains is *MATa ade2 ade6 can1-10 his3-11, 15 leu2-3, 112 trp1-1 ura3-1* (W303).

862 **Genotype of all strains is *MATa his3-Δ200 can1 ura3-52 leu2Δ1 lys2-801 trp1-289* (BJ5459). The

863 *atg19, atg36, csm1, heh1, lrs4, and nur1* strains are from the yeast knock out collection (Chu and

864 Davis, 2008) and are derived from BY4741 (*MATa his3Δ1 leu2Δ met15Δ ura3Δ*).

865

866 **Extended Data Table III** Plasmids used in this study.

Plasmid Name	Gene	Epitope Tag	Marker	Promoter	2μ/ CEN/ int	Reference
pKC801	<i>MED13</i>	3HA	URA3	<i>ADH1</i>	CEN	(Stieg <i>et al.</i> , 2018)
pDS2	<i>GAL4AD-</i> <i>MED13</i> ¹⁻³⁰⁵	1HA	LEU2	<i>ADH1</i>	2μ	(Stieg <i>et al.</i> , 2018)
pDS4	<i>GAL4AD-</i> <i>MED13</i> ⁹⁰⁷⁻¹⁴²¹	1HA	LEU2	<i>ADH1</i>	2μ	(Stieg <i>et al.</i> , 2018)
pDS7	<i>GAL4AD-</i> <i>MED13</i> ³⁰⁶⁻⁵⁷⁰	1HA	LEU2	<i>ADH1</i>	2μ	(Stieg <i>et al.</i> , 2018)
pDS8	<i>GAL4AD-</i> <i>MED13</i> ⁵⁷¹⁻⁹⁰⁶	1HA	LEU2	<i>ADH1</i>	2μ	(Stieg <i>et al.</i> , 2018)
pSH3	<i>RPH1</i>	GFP	URA3	<i>ADH1</i>	CEN	This study
pSH4	<i>PGK1</i>	GFP	URA3	<i>ADH1</i>	CEN	This study
pSH8	<i>GAL4BD-SNX4</i>	1HA	TRP1	<i>ADH1</i>	2μ	This study

pSH16	<i>GAL4BD-</i> <i>SNX4</i> ¹⁻¹⁵⁷	1HA	TRP1	<i>ADH1</i>	2μ	This study
pSH17	<i>GAL4BD-</i> <i>SNX4</i> ¹⁵⁷⁻⁴²³	1HA	TRP1	<i>ADH1</i>	2μ	This study
pSH18	<i>GAL4AD-</i> <i>MED13</i> ⁹⁰⁷⁻¹¹⁶³	1HA	LEU2	<i>ADH1</i>	2μ	This study
pSH19	<i>GAL4AD-</i> <i>MED13</i> ¹¹⁶³⁻¹⁴²¹	1HA	LEU2	<i>ADH1</i>	2μ	This study
pSW217	<i>MSN2</i>	GFP	URA3	<i>ADH1</i>	CEN	This study
pSW218	<i>MED13</i>	GFP	URA3	<i>ADH1</i>	CEN	This study
pSW221	<i>VPH1</i>	mCherry	URA3	<i>ADH1</i>	CEN	(Willis <i>et al.</i> , 2020)
pSW230	<i>CCL1</i>	GFP	URA3	<i>ADH1</i>	CEN	This study
pSW288	<i>CRN1-MED13</i>	GFP	URA3	<i>ADH1</i>	CEN	This study
pSW320	<i>MED13</i>	GFP	HIS3	<i>ADH1</i>	CEN	This study
RSB2507	<i>NAB2</i>	mCherry	LEU2	<i>ADH1</i>	CEN	(Willis <i>et al.</i> , 2020)
pFD846	<i>RIM15</i>	GFP	TRP1	<i>ADH1</i>	CEN	(Wanke <i>et al.</i> , 2005)
GFP-Atg8	<i>ATG8</i>	GFP	TRP1	<i>Own</i>	CEN	(Abeliovich <i>et al.</i> , 2003)

pCM188	<i>TET</i>	-	URA3	<i>ADH1-</i>	CEN	(Gari <i>et al</i> ,
				<i>Tet02</i>		1997)

867

Supplemental Figure 1

Supplemental Figure 1. Med13 is degraded in response to rapamycin. **A** Degradation kinetics of endogenous Med13-9myc protein levels in wild-type cells treated with 200 ng/mL rapamycin. Error bars indicate S.D., N=3 biological samples. **B** RT-qPCR analysis probing for *MED13* mRNA expression in wild-type cells following nitrogen starvation. $\Delta\Delta C_t$ results for relative fold change ($\log_2$) values using wild-type unstressed cells as a control. Transcript levels are given relative to the internal *ACT1* mRNA control. **C** Western blot analysis of Med13-3HA protein levels, expressed from a single copy functional plasmid (Stieg *et al.*, 2018) (pKC801) in wild-type (RSY10), *ump1Δ* (RSY2160) and *pep4Δ prb1Δ.1* (RSY449) cells following nitrogen starvation. Pgk1 was used as a protein loading control in all experiments. **D** Quantification of Med13-3HA degradation kinetics in indicated mutants obtained from c. Error bars indicate S.D., N=3 biological samples.

Supplemental Figure 2

Supplemental Figure 2. Med13 accumulates within the vacuole following nitrogen starvation. **A** Fluorescence microscopy of *pep4Δ prb1Δ.1* cells expressing endogenous Med13-mNeongreen and Vph1-mCherry (vacuole marker) after 24 h in SD-N. Representative single plane images are shown. Bar = 5μm. **B** Fluorescence microscopy of wild-type cells expressing endogenous Med13-YFP (RSY1812) treated with 200 mM PMSF before and after 24 h in SD-N media. FM4-64 staining was used to visualize the vacuole. Representative deconvolved images are shown. Bar = 5μm.

Supplemental Figure 3

Supplemental Figure 3. The autophagic degradation of Med13 does not require known selective autophagy adapter proteins. **A** Western blot analysis of Med13-GFP cleavage assays in WY (RSY10), *atg1Δ* (RSY2094), (RSY2160) and *atg1Δ ump1Δ* (RSY2202) cells after nitrogen starvation. **B** Western blot analysis of endogenous Med13-9myc degradation kinetics in *atg17Δ* cells as well as in strains defective in known nucleophagy pathways following nitrogen starvation. **C** Degradation kinetics and half-life of Med13 protein levels obtained in a. Error bars indicate S.D., N=3 biological samples. **D** Wild-type BY4741 cells and the indicated mutants expressing Med13-GFP (pSW218) were nitrogen-starved for the indicated times and the accumulation of free GFP was monitored by Western blot analysis using anti-GFP antibodies. Pgk1 levels were used as a protein loading control for all experiments. Degradation kinetics of Med13-GFP are slower in BY4741 cells which are more resistant to stress than the W303 strain. However, all these mutants accumulated free GFP which is indicative of vacuolar degradation.

Supplemental Figure 4

Supplemental Figure 4. Mass spectrometry screen to identify proteins required for Med13 vacuolar degradation. **A** SDS-PAGE gel of immunoprecipitation assays with Med13 in unstressed cells or cells treated with 50nM rapamycin for 90 minutes. Spyro Ruby stain was used to visualize proteins immunoprecipitated with Med13-3xHA from whole cell lysate. Asterisks denote bands that were present in stressed samples and absent in unstressed samples. **B** Western blot analysis of immunoprecipitation experiments performed in wild-type and *pep4Δprb1Δ.1* cells. Cells harboring an expression plasmid for Med13-3HA were grown to mid-log and treated with 50nM rapamycin for 4 h. **C** Gle1 auxin inducible degron-degron system. The BY4741 strain harboring endogenous *GLE1-TAP-AID-FLAG* were grown to mid-log and treated with 250 μ M Auxin. Cells were then washed and resuspended in SD-N media containing 250 μ M Auxin for indicated time points. Gle1 protein depletion was monitored via Western blot analysis using antibodies against FLAG. **D** Crm1-doxycyclin inducible degron system. Cells expressing Crm1-HA under a tetracycline repressible

909 promoter and endogenous Med13-9xmyc (RSY2348) were grown to mid-log. A sample was
 910 removed prior to the addition of doxycycline for Western blot analysis to visualize Crm1-HA (far left
 911 T=0 bands). The remaining culture was treated with doxycycline for 1 hour prior to switching cells
 912 to SD-N media with doxycycline. Western blot analysis was used to monitor Crm1-HA (top panel)
 913 and Med13-9xmyc (bottom panel) degradation following doxycycline and nitrogen starvation
 914 treatment. **E** Western blot analysis of Med13-9xmyc protein levels following nitrogen starvation in
 915 *msn5Δ* cells (RSY2399).

916

Supplemental Figure 5

Supplemental Figure 5. Western blot controls for Y2H plasmids. **A**, Western blot analysis of cells expressing indicated Med13 activating domain Y2H constructs. **B** and **C** Western blot analysis of cells expressing of Gle1 and Snx4 binding domain Y2H constructs respectively.

Supplemental Figure 6

Supplemental Figure 6. Snx4 localizes to the nuclear periphery following nitrogen starvation. **A** Fluorescence microscopy monitoring colocalization of endogenous Med13-mNeon green and Atg17-Redstar after 2 h of nitrogen starvation in *pep4Δ prb1Δ.1* cells (RSY2400). Representative single plane images are shown. Bar = 5μm. **B** Fluorescence microscopy of GFP-Snx4 localization in wild-type cells following 2 h of nitrogen starvation. The number of foci increases following stress. Representative deconvolved images are shown. Bar = 5μm. **C** Fluorescence microscopy of *pep4Δ* cells expressing GFP-Snx4 and harboring Nab2-mCherry (nuclear marker) before and after nitrogen starvation. Representative deconvolved images are shown. Bar = 5μm.) **D** Fluorescence microscopy of endogenous Med13-mNeon green and mCherry-Snx4 in *pep4Δ prb1Δ.1* cells before and after nitrogen starvation. Representative deconvolved images are shown. Vacuolar Med13 population indicated by the blue arrow and perinuclear Med13-Snx4 colocalization indicated by the pink arrow. Bar = 5μm. **E** Fluorescence microscopy of colocalization experiments

935 performed in wild-type cells expressing GFP-Snx4 and endogenous Gle1-RedStar. Cells were
936 grown to mid-log, washed and resuspended in SD-N media for 2 h. Representative large field view
937 deconvolved images are shown. Bar = 5µm.

Supplemental Figure 7

Supplemental Figure 7. Snx4 is required for viability in long-term nitrogen starvation.

A Western blot analysis of Pgk1- GFP (pSH4) cleavage assays in wild-type and *snx4Δ* cells following nitrogen starvation. Endogenous Pgk1 protein levels were used as a loading control. **B** Wild-type and *snx4Δ* cells were grown to mid-log, washed and resuspended in SD-N media for the indicated number of days. The percentage of inviable cells within the population was determined using phloxine B staining and fluorescence activated cell analysis (FAC). Quantification of N=2 independent biological experiments. Data are mean ± S.D. ** $P = 0.0012$, **** $P = <0.0001$ (Willis *et al.*, 2020) **C** Quantification Atg8 protein levels in wild-type and *med13Δ* cells relative to Pgk1 loading control. Quantification of N=3 biological samples. Data are mean ± S.D. **** $P = <0.0001$

Source Data Figure 1

Source Data for figure 1. Black boxes contain cropped representative images depicted in figure 1.

Source data for figure 4B. Black boxes contain representative cropped images depicted in figure 4.