1 **Language impairment with a microduplication in 1q42.3q43**

2 Antonio Benítez-Burraco¹, Maite Fernández-Urquiza², M^a Salud Jiménez-Romero³

3

4 1. Department of Spanish, Linguistics, and Theory of Literature (Linguistics),
5 University of Seville, Seville, Spain

6 2. Department of Spanish Philology, University of Oviedo, Oviedo, Spain

7 3. Department of Education, University of Córdoba, Córdoba, Spain

8

9 Corresponding author:

10 Antonio Benítez-Burraco

11 Área de Lingüística General. Departamento de Lengua Española, Lingüística y Teoría
12 de la Literatura. Facultad de Filología. Universidad de Sevilla. C/ Palos de la Frontera
13 s/n. 41007-Sevilla (España)

14 e-mail: abenitez8@us.es

15 orcid.org/0000-0003-4574-5666

16

ABSTRACT

Deletions and duplications of the distal region of the long arm of chromosome 1 are associated with brain abnormalities and developmental delay. Because duplications are less frequent than deletions, no detailed account of the cognitive profile of the affected people is available, particularly, regarding their language (dis)abilities. In this paper we report on the cognitive and language features of a girl with one of the smallest interstitial duplications ever described in this region, affecting to 1q42.3q43 (arr[hg19] 1q42.3q43(235,963,632-236,972,276)x3). Standardized tests as well as the analysis of her language use in natural settings suggest that the proband's speech is severely impaired, exhibiting dysarthric-like features, with speech problems also resulting from a phonological deficit boiling down to a verbal auditory memory deficit. Lexical and grammatical knowledge are also impaired, impacting negatively on both expressive and receptive abilities, seemingly as a consequence of the phonological deficit. Still, her pragmatic abilities seem to be significantly spared, granting her a good command on the principles governing conversational exchanges. In silico analyses (literature mining, network analysis) and in vitro analyses (microarray) point to several genes as potential candidates for the observed deficits in the language domain. These include one gene within the duplicated region (*LYST*), one predicted functional partner (*CMIP*), and three genes outside the 1q42.3q43 region, which are all highly expressed in the cerebellum: *DDIT4* and *SLC29A1*, found strongly downregulated in the proband compared to their healthy parents, and *CNTNAP3*, found strongly upregulated.

Keywords: 1q distal duplications; language deficits; speech problems; differentially-expressed genes

INTRODUCTION

Deletions and duplications of the distal region of the long arm of chromosome 1 result in developmental anomalies, including intellectual disability and language problems. Deletions usually entail microcephaly, seizures, cognitive and psychomotor delay, growth delay, facial dysmorphisms, and cardiac defects (Juberg et al., 1981; Roberts et al., 2004; van Bon et al., 2008). That said, the exact symptoms exhibited by patients depend on the extension of the deleted region and consequently, on the number and functions of the affected genes. Duplications of this region are less frequent and patients exhibit a more variable clinical presentation, although common symptoms include intellectual disability and delayed psychomotor development, facial anomalies, and macrocephaly (Hemming et al., 2016; Morris et al. 2016).

In this paper, we report on a girl with an interstitial duplication in the 1q42.3q43 region. Rearrangements of this region are rare. Silipigni and colleagues (2017) reviewed 7 cases found in the literature, affecting to a slightly broader region (1q42.13q43), and described 2 new cases: a deletion of 1q42.13q43 and the reciprocal duplication, found in two consanguineous probands. Subjects with duplications suffer from language deficits. Specifically, the child examined by Silipigni et al. (2017) showed cognitive and social delay, and said his first words at the age of 2 only. Nonetheless, a detailed characterization of the language problems exhibited by patients with duplications of the distal region of the long arm of chromosome 1 is still pending. Not surprisingly, no conclusive genotype-phenotype link has been found for language deficits. One reason is that the role of the duplicated genes in neurodevelopment, and particularly, in language development, is mostly unknown. A second reason is that patients usually

bear complex translocations involving other chromosomes. In this paper we provide a comprehensive account of the language deficits and strengths of our proband, including a detailed psycholinguistic profile and an in-depth analysis of her language in use. We further advance a hypothesis about the molecular causes of language dysfunction. For this, we have relied on available evidence of the role played by the duplicated genes in brain development and function, but also, on predicted functional links between some of these genes and selected strong candidates for language development and language evolution in the species. Additionally, we have relied on results of our analysis of whole-genome gene expression pattern of the blood of our proband compared to her healthy parents.

MATERIAL AND METHODS

Linguistic, cognitive, and behavioral assessment

The global developmental profile of the proband was evaluated with the revised version of the Bateria de Aptitudes Diferenciales y Generales [Battery for Differential and General Abilities] (BADyG) (Yuste and Yuste, 1998) and with the Spanish version of the Inventory for Client and Agency Planning (ICAP) (Montero, 1996). The proband's linguistic skills were evaluated with the Spanish versions of the Peabody Picture Vocabulary Test (PPVT-3) (Dunn et al., 2006), the Gardner Receptive One Word Picture Vocabulary Test (ROWPVT) (Gardner, 1987), and the Illinois Test of Psycholinguistic Abilities (ITPA) (Kirk et al., 2009). The Registro Fonológico Inducido test [Induced Phonological Register] (Monfort and Juárez, 1988) was used to achieve a detailed knowledge of the proband's phonological awareness.

The Battery for Differential and General Abilities (BADyG)

The BADyG evaluates the child's abilities in 9 different domains: verbal comprehension (understanding of direct commands involving spatial concepts), verbal reasoning (finding analogical relationships between concepts), numerical comprehension (performing simple arithmetic calculations), numerical reasoning (understanding and solving simple mathematical problems), processing of rotated figures (rotating figures mentally), spatial reasoning (inducing the rules governing the assembly of geometric figures), visuo-auditive memory (recalling meanings from a story which is also displayed graphically), writing abilities (absence of dyslexic-like features), and discrimination (finding subtle differences between geometric figures). Scores are subsequently grouped to obtain 5 composite measures: verbal abilities (verbal comprehension + verbal reasoning), numerical abilities (numerical comprehension + numerical reasoning), visuospatial abilities (processing of rotated figures + spatial reasoning), logical reasoning (verbal reasoning + numerical reasoning + spatial reasoning), and general intelligence (verbal comprehension + verbal reasoning + numerical comprehension + numerical reasoning + processing of rotated figures + spatial reasoning). The version of the test employed in this case was BADyG/E1, aimed for children between 6 and 7 years old.

The Inventory for Client and Agency Planning (ICAP)

The ICAP is aimed to evaluate the subject's functional abilities and maladaptive behaviors in the following general areas: motor skills, social and communication skills, personal living skills, and community living skills. The ICAP measures the frequency and severity of 8 types of behavioral disturbances, which are organized in 3 subscales: asocial maladaptive behavior (uncooperative behavior and socially offensive behavior),

internalized maladaptive behavior (withdrawn or inattentive behavior, unusual or repetitive habits, and self-harm), and externalized maladaptive behavior (disruptive behavior, destructive to property, and hurtful to others). Behavior is rated as normal or abnormal, whereas behavioral problems are subsequently rated as marginally serious, moderately serious, serious, or very serious.

The Peabody Picture Vocabulary Test (PPVT-3)

This test is aimed to assess the correct acquisition and the level of receptive vocabulary in subject from 2.5 years to 90 years. It consists of 192 sheets with four colored drawings each. After hearing a word, the subject must indicate which illustration best represents its meaning.

Gardner Receptive One Word Picture Vocabulary Test (GROWPVT)

This test is aimed to assess receptive vocabulary in children between 2 and 11 years old, as a screening tool for detecting possible speech defects, learning disorders, auditory processing deficits, or problems for auditory-visual association. It consists of 100 picture-naming tasks: after hearing a word, the subject has to choose (either verbally or by pointing) the correct picture out of four different colored pictures.

Illinois Test of Psycholinguistic Abilities (ITPA)

This test aims to detect preserved psycholinguistic skills and specific difficulties experienced by children when trying to produce an adequate communicative contribution. In this test, psycholinguistic skills are construed as psychological processes and functions enabling the subject to convey his intentions during the communicative use of language (Kirk and McCarthy, 1961). These psychological skills

are ultimately related to the core processes involved in the reception, understanding, and transmission of a linguistic message. The ITPA comprises several subtests that are organized into two levels: the Representative Level and the Automatic Level. The Representative Level assesses the management of symbols via the evaluation of different kinds of processes: i) Receptive processes, including a) the ability to obtain meaning from a spoken word (Listening comprehension), and b) the ability to obtain meaning from a visual symbol, assessed through the capability to choose from among a group of drawings the one that is similar to the stimulus picture (Visual comprehension); ii) Organizational processes, including a) the ability to relate spoken words in a meaningful way, which is tested using a series of verbal analogies of increasing difficulty (Auditory association); and b) the aptitude to relate visual symbols, which is assessed through the matching of the stimulus picture to one drawing chosen among a set of four (Visual association); and iii) Expressive processes, including a) verbal fluency, which is estimated from the number of concepts expressed verbally (Verbal expression), and b) the ability to express concepts through gestures (Motor expression). Regarding the Automatic Level, it assesses how different habits, such as memory or remote learning, are integrated to produce an automatic chain of responses. The assessment is conducted via the evaluation of different kinds of processes: i) Integration or closure tests, including a) the competence to automatically use grammar in a task of sentence completion supported by drawings (Grammar integration or closure), b) the ability to identify a common object from its incomplete representation in a relatively complex context (Visual integration), c) the ability to produce a word from partially pronounced words (Aural integration), and d) the ability to synthesize the separate sounds of a word in order to produce the complete word (Sound gathering); ii) Sequential memory tests, including a) the ability to recognize and produce a word

from its partial pronunciation (Auditory sequential memory), and b) the ability to reproduce by heart sequences of non-significant figures after viewing the sequence for a short period of time (Visomotor Sequential Memory).

Induced Phonological Register (RFI)

This tests assesses phonological and articulatory abilities of Spanish-speaking children between 3 and 7 years old by means of elicited word production and repetition. Word production is induced by showing the child different pictures that she has to name. If the child is unable to produce the name, the speech-language therapist will provide her with the correct name and will ask the child to repeat it. The test comprises every possible sound of the Spanish language, so that it can be used to detect dyslalia as well as phonological impairment.

Language in use

In addition, the proband's language in use and communication skills were assessed through the analysis of a 12-minutes sample of her talk while chatting with her family. Three different settings were considered: 1) a meal at home with her parents, 2) a family meeting with her parents, her brother and her grandparents; and 3) a conversation with her mother and her brother after one school day. The conversations were video-recorded and then transcribed and coded using the tools provided by the CHILDES Project (MacWhinney, 2000). In brief, speech production phenomena (including relevant information about articulatory processes, prosody, and fluency) were coded in the main lines of the transcript using CHAT (*Codes for the Human Analysis of Transcripts*). Relevant information about gestural communication and the communicative use of gaze was included in the main lines within square brackets. Lastly, phonological,

morphological, syntactic, lexical and pragmatic phenomena were coded in dependent lines. Because functional approaches to language impairment demand taking into account the pragmatic consequences of structural errors, we first tagged structural errors and then evaluated their communicative effects using the “Pragmatic Evaluation Protocol of Corpora” (PREP-CORP). This protocol has been satisfactorily used to provide the pragmatic profiles of diverse neurodevelopmental disorders (Fernández-Urquiza et al., 2015, Fernández-Urquiza et al., 2017, Diez-Itza et al., 2018). Finally, a whole pragmatic profile of the proband was built up. For this, we computed the number of occurrences of each label using CLAN (*Computerized Language Analysis*) (Conti-Ramsden, 1996).

Assessment of the parents’ cognitive and language abilities

The linguistic profile of the proband’s parents was assessed with the verbal scale of the Spanish version of the WAIS-III (Wechsler, 2002), the Test de Aptitud Verbal "Buenos Aires" [Buenos Aires Verbal Aptitude Test] (BAIRES) (Cortada de Kohan, 2004), the Batería para la Evaluación de los Procesos Lectores en Secundaria y Bachillerato [Battery for the Evaluation of Reading Abilities in High School Students] (PROLEC-SE) (Ramos and Cuetos, 2011), and one specific task aimed to evaluate the comprehension of passive sentences and co-referential expressions. The verbal scale of the WAIS-III comprises 6 tasks aimed to assess the subject’s abilities in two different domains of language processing: verbal comprehension (Similarities (S), Vocabulary (V), Information (I), and Comprehension (CO)) and working memory (Digits (D) and Arithmetic (A)). The BAIRES test aims to evaluate language comprehension and production in subjects above 16 years old. It comprises two tasks which assess, respectively, the subject’s ability to understand language and her ability to provide

synonymic words and verbal definitions. Regarding the PROLEC-SE, it is mostly used to assess reading abilities of youngsters between 12 and 18 years old, but because a full reading competence is usually acquired before 18, it can be also used with adults. It comprises six tasks that evaluate three different aspects of reading: lexical processes (word reading, pseudoword reading), semantic aspects (text comprehension, text structure), and syntactic abilities (picture-sentence matching, punctuation marks). To end, the *ad-hoc* task was a sentence-picture matching task aimed to evaluate the comprehension of bound anaphora in complex sentences (that is, the ability to properly bind a noun phrase in the dependent clause to a referential element in the main clause) and of canonical long passive structures (that is, with a *by*-phrase). While the first task is highly-demanding in computational terms, the second task is usually difficult to perform by people with poor reading abilities, because passives are very infrequent in spoken Spanish.

Cytogenetic and molecular analyses

Karyotype analysis

Peripheral venous blood lymphocytes were grown following standard protocols and collected after 72 hours. A moderate resolution G-banding (550 bands) karyotyping by trypsin (Gibco 1x trypsin® and Leishmann stain) was subsequently performed. Microscopic analysis was conducted with a Nikon® eclipse 50i optical microscope and the IKAROS Karyotyping System (MetaSystem® software).

DNA from the patient and her parents was extracted from 100 µl of EDTA-anticoagulated whole blood using MagNA Pure (Roche Diagnostics, West Sussex, UK) and used for subsequent analyses.

Fragile X syndrome analysis

CGG expansions affecting the gene *FMRI* (the main determinant factor for X-fragile syndrome) were analyzed in the proband according to standard protocols. Polymerase chain reaction (PCR) of the fragile site was performed with specific primers for the fragile region of the *FMRI locus* and the trinucleotide repeat size of the resulting fragments was evaluated by electrophoresis in agarose gel.

Multiplex ligation-dependent probe amplification (MLPA)

MLPA was conducted to detect abnormal copy-number variations (CNVs) in subtelomeric regions of the chromosomes, as well as frequent interstitial CNVs. MLPA consists on the amplification of different probes using a single PCR primer pair. Each probe detects a specific subtelomeric DNA sequence. Three different kits from MRC-Holland were employed. Two of them (SALSA[®] MLPA[®] probemix P036-E3 SUBTELOMERES MIX 1 and SALSA[®] MLPA[®] probemix P070-B3 SUBTELOMERES MIX 2B) were used for examining subtelomeric regions. The third one (SALSA[®] MLPA[®] probemix P245-B1 Microdeletion Syndromes-1A) was used to detect frequent microdeletion and microduplication syndromes: 1p36 deletion syndrome, 2p16 deletion syndrome, 2q23 deletion syndrome (involving *MBD5*), 2q33 deletion syndrome (involving *SATB2*), 3q29 deletion syndrome, Wolf-Hirschhorn syndrome (resulting from a 4p16.3 deletion), Cri du Chat syndrome (resulting from a 5p15 deletion), Sotos syndrome associated to a 5q35.3 deletion, Williams syndrome (resulting from a 7q11.23 deletion), Langer-Giedion syndrome (resulting from a 8q deletion), 9q22.3 deletion syndrome, 15q24 deletion syndrome, Prader-Willi and Angelman syndromes (resulting from paternal or maternal 15q11.2 deletions), a severe variant of Rubinstein-Taybi syndrome resulting from a deletion in 16p13.3, 17q21

deletion syndrome, Miller-Dieker syndrome (resulting from a 17p deletion), Smith-Magenis syndrome (resulting from a 17p11.2 deletion), NF1 microdeletion syndrome (aka Van Asperen syndrome), linked to a 17q11.2 deletion), Phelan-McDermid syndrome (resulting from a 22q13 deletion), Di George syndrome (resulting from deletions in 22q11, distal 22q11, or 10p14), Rett syndrome (associated to *MECP2* deletion) and *MECP2* duplication syndrome. The PCR products were analyzed by capillary electrophoresis in an automatic sequencer Hitachi 3500 and further analyzed with the Coffalyser V 1.0 software from MRC-Holland.

Microarrays for whole-genome CNVs search and chromosome aberrations analysis

The DNA from the patient and her parents was hybridized on a CGH platform (Agilent Technologies). The derivative log ratio spread (DLRS) value was 0.172204. The platform included 60.000 probes. Data were analyzed with Agilent CytoGenomics 3.0.5.1 and qGenViewer, and the ADM-2 algorithm (threshold = 6.0; abs (log2ratio) = 0,25; aberrant regions had more than 4 consecutive probes).

Microarrays for DEGs search

In order to determine the genes that could be differentially expressed in the proband (DEGs) compared to her healthy parents, microarray analyses of blood samples from the three of them were performed. Total RNA was extracted with the PAXgene Blood RNA Kit IVD (Cat No./ID: 762164). RNA quality and integrity were confirmed with a Bioanalyzer RNA 6000 Nano. All samples had RNA integrity number (RIN) values above 9. An Affymetrix® Scanner 3000 7G was then used for analyzing transcriptome changes. The resulting raw data were processed with the Affymetrix® GeneChip® Command Console® 2.0 program. Next, *.CEL files were checked to certify the RNA

integrity and the suitability of the labeling and the hybridization processes. Finally, the raw data from the different arrays were normalized with the SST (Signal Space Transformation)-RMA (Robust Microarray Analysis) tool (Irizarry et al., 2003). Normalized data (*.CHP files) were subsequently used to search for DEGs in the proband compared to her parents. Statistical analyses were conducted with the LIMMA (Linear Models for Microarray Analysis) package of BioConductor, using the TAC 4.0 software.

RESULTS

Clinical History

The proband (Figure 1A) is a girl born by normal delivery after 36 weeks of low-risk gestation. The mother was a 29-year-old woman, who suffered from Crohn's disease treated with Imurel. At delivery, no signs of disease were observed in the newborn (serology was negative and there was no evidence of group B streptococcus (GBS) infection). Nonetheless, the baby exhibited a supernumerary digit in her right hand, which was subsequently removed by surgery. At birth, her weight was 3.51 kg (percentile 79th), her height was 52 cm (percentile 95th), and her occipitofrontal circumference was 36 cm (percentile 96th). Apgar scores were 9 (at 1') and 10 (at 5'). When she was 12 months old, her weight was 8,64 kg (percentile 38th), her height was 73,5 cm (percentile 42th), and her occipitofrontal circumference was 47 cm (percentile 93th). At age 1 year and 7 months, her weight was 10 kg (percentile 34th), her height was 81 cm (percentile 42th), and her occipitofrontal circumference was 48,5 cm (percentile 93th).

The proband's parents reported on hypotonia that hindered sucking during her first months of life. Hypotonia worsened with time and when she was 12 months old, a cranial magnetic resonance imaging (MRI) was performed. Main findings were a reduced white matter signal-intensity in the temporal lobe of both hemispheres, mostly impacting on the left hemisphere, as well as a mild delay in the myelination pattern of subcortical areas. At the age of 34 months, a prominent forehead ($SD < 2$) was a distinctive feature, as well as 4 hyperchromic spots and abnormally high levels of thyroid-stimulating hormone (TSH). Overall, the girl showed a significant psychomotor delay. At 12 months of age, she had not taken her first steps and showed unstable head control and sitting. Generalized hypotonia, as well as motor and cognitive delay were diagnosed at that age. She was able to stand up and wander with support only at 24 months of age. She also began to manipulate objects at that age. When she was 32 months old, she was able to move from dorsal decubitus to sitting without help. At 34 months of age, she showed typical strength, muscle tone, and spontaneous movements. At this age, she was also able to stand stably and to wander without aid. She succeeded in grabbing objects, passing them from one hand to the other, and in coordinating fine movements. When she was 4 years old, she already walked without aid, but some motor clumsiness and frequent falls were observed. She still suffered from slight hypotonia with kyphosis when sitting without support. When the proband was 6 years old, she was diagnosed with Valgus foot on both feet. She still fell down frequently and reported pain in both soles.

Early language development

Regarding language development, babbling was first observed at the age of 12 months, whereas bi-syllabic utterances were first reported when the proband was 2 years old. At that age, she already understood simple direct commands. Pointing appeared at the age of 27 months. First words were attested at the age of 32 months. When the proband was 4 years old, she was able to understand complex verbal orders. Nonetheless, expressive language was severely impaired. Accordingly, the two-word stage was reached at the age of 56 months only, with an active vocabulary of around 20 words and the use of non-inflected verbs as a hallmark of her discourse. Telegraphic speech was reported from age 6-7 only.

In spite of her marked language deficits, the proband has been attending a regular school. During the preschool education period, her teachers reported that she mostly communicated through screaming, gesturing, and bringing others to the place where objects of interest were placed, as she was unable to name them. She experienced difficulties with focused attention and she was not interested in new stimuli. She seemed to understand simple commands and sometimes she produced some understandable words. When she was 3 years old she scored significantly low in the Gardner Receptive One Word Picture Vocabulary (ROWPVT), with a verbal age of 2 years and 2 months. At the age of 6, her receptive language was still seriously delayed. She scored 16 direct points in the Peabody Picture Vocabulary Test (PPVT-3) (< 1st percentile; -3 SD), which corresponds to a verbal IQ of 55 and a verbal age of 2 years and 7 months. She communicated mostly through facial or body gestures, or touching the other with her hand, as speech and expressive language was seriously impaired. When the proband was 8 years old, her cognitive development was assessed with the

Batería de Aptitudes Diferenciales y Generales (BADyG). She scored far below typicality, being the verbal abilities the most impaired domain, whereas spatial abilities were quite preserved (Figure 1B). At the time of our evaluation, when she was 9 years and 8 months old, the proband was attending a normal classroom, although she was assisted by a support teacher and a speech therapist.

Detailed cognitive and language assessment

At the age of 9 years and 8 months, the proband's global developmental profile and adaptive behavior were evaluated with the Spanish version of the Inventory for Client and Agency Planning (ICAP). The resulting scores were suggestive of a delay of 3 years and 3 months, being motor skills and communication skills the most impaired domains, whereas personal living skills were the most preserved domain. Community living skills were moderately delayed (Figure 1C; see Supplemental file 1 for details).

Focusing on language, her psycholinguistic development was assessed in detail with the Spanish version of the Illinois Test of Psycholinguistic Abilities (ITPA). The resulting composite psycholinguistic age was 5 years and 2 months, well below her chronological age. Still, she showed a quite irregular profile, with visual and motor abilities being quite spared, but with auditory and grammatical processing abilities, as well as vocal emission, being the most impaired domains (Figure 1D; see Supplemental file 1 for details). More specifically, we found that her receptive language was seriously delayed. She scored 53 direct points in the Peabody Picture Vocabulary Test (PPVT-3), which is suggestive of a verbal age of 5 years and 1 month (see Supplemental file 1 for details). Poor vocabulary is expected to account as well for her problems for understanding language. Likewise, we found that speech production and phonological

awareness were severely delayed, exhibiting features that are typically found in children between 4-6 years old, according to the Registro Fonológico Inducido test (see Supplemental file 1 for details). Fluency problems were frequently attested. The proband experienced marked problems for correctly uttering tri-syllabic words, or even bi-syllable words containing consonant clusters. Deletions of weak syllables were frequently observed (e.g. *tuga* for *tortuga* ‘turtle’), as well as cluster reduction (e.g. *peso* for *preso* ‘prisoner’) and assimilations resulting in consonant harmony (e.g. *zaza* instead of *taza* ‘cup’ or *pambor* instead of *tambor* ‘drum’).

Figure 1. Main physical and cognitive findings in the proband. A. Facial picture of the proband. B. Developmental profile of the proband at the age of 8 years according to the Bateria de Aptitudes Diferenciales y Generales (BADyG). C. Developmental profile of the proband at the age of 9 years and 8 months according to the Spanish version of the Inventory for Client and Agency Planning (ICAP). In order to make more reliable comparisons, the resulting scores are shown as relative values referred to the expected

scores according to the chronological age of the child. Abbreviations: DA, developmental age; CA, chronological age. D. Developmental profile of the proband at the age of 9 years and 8 months according to the Spanish version of the Illinois Test of Psycholinguistic Abilities (ITPA). Abbreviations: DA, developmental age; CA, chronological age.

Lastly, we examined how the proband put her knowledge of language into use for communicating. We analyzed three different naturally-occurring interactions with her family (see Supplemental file 2 for details). At the expressive level, we confirmed the existence of widespread articulatory problems, including consonant deletions (e.g. *apatos* instead of *zapatos* ‘shoes’; *artera* instead of *cartera* ‘bag’; *e* instead of *él* ‘he’; *Carcola* instead of *Caracola* ‘The Seashell’), deletions of weak syllables (e.g. *tuche* instead of *estuche* ‘case’; *pues* instead of *después* ‘after’; *mano* instead of *hermano* ‘brother’; *jama* instead of *pijama* ‘pyjama’), and simplifications of consonant clusters (e.g. *apoco* instead of *tampoco* ‘neither’; *epués* instead of *después* ‘after’; *pima* instead of *prima* ‘cousin’). We also observed frequent sound assimilations (e.g. *Canme* instead of *Carmen* (proper noun)), non-pervasive frontalizations (e.g. *da* instead of *la* ‘the’; *dlase* instead of *clase* ‘class’), and many instances of loose articulation. Phonetic paraphasia (e.g. *felfa** instead of *lazo* ‘bow’; *tunales** instead of *naturales* ‘natural sciences’) sometimes resulted in idiosyncratic neologisms. Overall, these expressive difficulties seriously hindered the intelligibility of the proband’s discourse. The mean length of utterance in words (MLUw), a common measure of language structural complexity, was 3.9, which is typically found in children aged between 3 years and 6 months and 3 years and 11 months (Rice et al., 2010). Morphosyntactic errors included omissions and substitutions of bound morphemes (e.g. *viene* instead of *vienen*_{PL} ‘they

come'; *quién es* instead of *quiénes_{SPL} son_{PL}* 'who are they'; *el* instead of *la* 'the_{FEM}'), and omissions of different word classes such as prepositions (e.g. *veintiséis mayo* instead of *veintiséis de mayo* 'May the 26th'; *hablar* instead of *para hablar* 'to talk'), object pronouns (*pongo* instead of *me pongo* 'I wear'; *gusta* instead of *te gusta* 'do you like'), articles (*a piscina* instead of *a la piscina* 'to the swimming pool'), and auxiliary verbs (*íá* instead of *he ido* 'I have gone'). Sporadic inversions of the canonical word order in Spanish were also attested (i.e. subject-object-verb instead of subject-verb-object).

In spite of these problems with structural components of language, the proband's pragmatic skills (i.e. how language is put into use) were significantly spared. First, she mastered turn-taking during conversational exchanges (Sacks et al., 1974), including gaze selection of the interlocutor (even among six participants), or the use of paralinguistic, non-verbal, and echolalic devices for taking the floor. Second, she exhibited signals of metapragmatic awareness, as when feeling embarrassed because of her slow take of the floor during conversational exchanges due to her poor expressive abilities. Third, she seemed to master Grice's Cooperative Principle (Grice, 1975), including the ability to infer implicit meanings resulting from the flouting of Gricean maxims (Davies, 2007). Accordingly, our proband was able to detect blatant violations of the maxim of quality ('assume that an utterance will generally be true') for humorous purposes, and to flout the maxim of quantity ('give the right amount of information taking into account contextual requirements') for conveying implicit meanings.

Cytogenetic and molecular analyses

Routine cytogenetic and molecular analyses of the proband were performed when she was 8 years old. PCR analysis of the FMR1 fragile site resulted normal. MLPAs of

selected subtelomeric regions yielded normal results too. Probes for common CNVs failed to detect genomic signatures of high prevalent syndromes resulting from microdeletions or microduplications.

A comparative genomic hybridization array (array-CGH) was then performed. The array-CGH identified a duplication of 1 Mb in the 1q42.3q43 region (arr[hg19] 1q42.3q43 (235,963,632-236,972,276)x3) (figure 2A). The duplicated region includes 16 genes of which 9 are protein-coding genes (figure 2B). The proband also bears a second microduplication in chromosome 1 (arr[hg19] 1q21.2 (148,867,551-149,244,468)x3). The duplicated region contains 6 pseudogenes and 2 genes coding for snRNAs, but no protein-coding genes, and the duplication is predicted as benign.

Figure 2. Chromosomal alterations found in our proband. A. Screen capture of the array-CGH of the proband's chromosome 1 showing the microduplication at 1q42.3q43. B. Screen capture of the UCSC Genome Browser (<https://genome.ucsc.edu/>) showing the genes duplicated in the proband.

474 Our proband exhibits some of the physical, behavioral, and cognitive features of partial
 475 1q duplications (Table 1). The duplication in 1q42.3q43 was inherited from the
 476 proband's non-symptomatic father (figure 3A). The man exhibits an average verbal IQ,
 477 with no evidence of impairment in any of the assessed domains (figure 3B).
 478 Additionally, he exhibits a good command of verbal abilities as assessed with the
 479 BAIRE test, around 82nd percentile, but low reading abilities as assessed with the
 480 PROLEC-SE test, below 5th percentile, mostly resulting from poor text understanding
 481 (figure 3C). He doesn't show any of the other distinctive features of partial 1q
 482 duplications.
 483

Clinical Features	
Cognitive/behavioural features	
Intellectual disability	X
Developmental delay	X
Neurological findings	
Seizures	
Physical features	
Microcephaly	
Dysmorphic facial features	
Abnormal face shape/size	
Long and smooth philtrum	
Micrognathia	
Narrow palpebral fissures	
Wide, flat nasal bridge	
Abnormal palate (cleft or high-arched)	
Abnormal ears (low-set or posteriorly rotated, poorly formed helices)	
Abnormal fingers or toes (camptodactyly)	
Preaxial polydactyly	X
Other medical problems	
Heart disease	

Table 1. Summary table with the most relevant clinical features of our proband compared to other patients with distal 1q duplications (following Hemming et al. 2016 and Morris et al. 2016)

Figure 3. Chromosomal alterations and linguistic profile of the proband's father. A. Screen capture of the array-CGH of the chromosome 1 of the proband's father showing the microduplication at 1q42.3q43. B. Developmental profile of the proband's father according to the verbal component of the WAIS-III (for comparison, the figure includes the profile of the healthy, non-carrier mother). C. Reading abilities of the proband's father according to the PROLEC-SE test (for comparison, the figure includes the profile of the healthy, non-carrier mother).

In order to delve into the molecular causes of the speech and language problems exhibited by the proband, we conducted several in silico analyses. First, we surveyed the available literature searching for clinical cases linked or associated to the mutation, dysfunction, or dysregulation of the genes duplicated in our proband. We found that a duplication affecting *EDARADD* has been identified in a patient with autism spectrum

disorder (ASD) (Prasad et al., 2013). *GPR137B* has been associated to neurocognitive improvement of schizophrenic patients after antipsychotics intake (McClay et al., 2011). *LYST*, which is interrupted by the duplication event, is a candidate for Chediak-Higashi syndrome (OMIM # 214500), a rare autosomal recessive disorder which entails diverse neurologic problems, including cognitive decline and parkinsonism (Kaplan et al., 2008; Introne et al., 2017). Finally, two polymorphism of *MTR*, which is also interrupted by the duplication event, have been associated with bipolar disorder (BD) and schizophrenia (SZ) (Kempisty et al., 2007).

Additionally, we surveyed DECIPHER (<https://decipher.sanger.ac.uk/>) looking for similar duplication events in other individuals, but also for duplications smaller than the region duplicated in the proband, that might help establish more robust genotype-to-phenotype links regarding her language disabilities. We found that duplications encompassing *HEATR1*, *ACTN2*, and *MTR* usually entail delayed motor development, but no language problems (patient 333241). By contrast, duplications encompassing *LGALS8* and affecting *EDARADD* and *HEATR1* normally result in language delay mostly impacting on the expressive domain (patient 248308) (Figure 4).

521

522 **Figure 4.** DECIPHER patients of interest bearing chromosomal duplications at
523 1q42.3q43 with similar or smaller sizes than the one found in the proband. The patients’
524 phenotype is summarized at the bottom of the figure.

525

526 Finally, we used String 10.5 (www.string-db.org) for uncovering potential functional
527 links between the genes duplicated in our proband and genes related to language
528 function, specifically, candidate genes for i) prevalent language disorders
529 (developmental dyslexia (DD) and specific language impairment (SLI), as listed by
530 Paracchini et al. 2016, Pettigrew et al. 2016, and Chen et al. 2017); and ii) language
531 evolution, as discussed by Boeckx and Benítez-Burraco (2014a,b) and Benítez-Burraco
532 and Boeckx (2015) (many of the genes belonging to this second group are also
533 candidates for language dysfunction in broader cognitive disorders, particularly, ASD
534 and SZ, as discussed in detail in Benítez-Burraco and Murphy, 2016, Murphy and

Benítez-Burraco, 2016 and Murphy and Benítez-Burraco, 2017). The whole set of genes considered in this analysis is listed in the Supplemental file 3. String 10.5 predicts physical and functional associations between proteins relying on different sources (genomic context, high-throughput experiments, conserved coexpression, and text mining) (Szkarczyk et al. 2015). Several proteins (GRIN2A, GRIN2B, HRAS, PVALB, and ITGB4) were predicted to be direct interactors of the product of one of the genes located within the duplicated fragment, namely, ACTN2 (Figure 5). Co-expression data and experimental/biochemical data further pointed out to a functional link between HEATR1 and NOP9, as well as between LYST and GABARAP.

545

Figure 5. Interaction network of the proteins encoded by the genes duplicated in the proband. The network was drawn with String (version 10.5; Szklarczyk et al. 2015) license-free software (<http://string-db.org/>), using the molecular action visualization. It includes the products of the protein-coding genes duplicated in the subject, their potential interactors according to String, and the products of 15 strong candidates for language development and/or evolution discussed in the text. Colored nodes symbolize gene/proteins included in the query (small nodes are for proteins with unknown 3D structure, while large nodes are for those with known structures). The color of the edges represents different kind of known protein-protein associations. Green: activation, red: inhibition, dark blue: binding, light blue: phenotype, dark purple: catalysis, light purple: post-translational modification, black: reaction, yellow: transcriptional regulation. Edges ending in an arrow symbolize positive effects, edges ending in a bar symbolize negative effects, whereas edges ending in a circle symbolize unspecified effects. Grey edges symbolize predicted links based on literature search (co-mentioned in PubMed abstracts). Stronger associations between proteins are represented by thicker lines. The medium confidence value was .0400 (a 40% probability that a predicted link exists between two enzymes in the same metabolic map in the KEGG database: <http://www.genome.jp/kegg/pathway.html>). The diagram only represents the potential connectivity between the involved proteins, which has to be mapped onto particular biochemical networks, signaling pathways, cellular properties, aspects of neuronal function, or cell-types of interest.

Besides these in silico analyses, we also conducted in vitro analyses. Specifically, we performed microarray analyses of blood samples from the proband to determine whether she exhibited altered patterns of gene expression that may account for the

observed symptoms. We used her healthy parents as a control. The results of the microarrays are listed in the Supplemental file 4. First, we checked whether the genes highlighted above were differentially expressed in the proband compared to her unaffected parents (Figure 6). Genes showing an opposite expression pattern in the proband and in both healthy parents were of particular interest to us. Among the protein-coding genes located within the duplicated fragment, we found that *EDARADD* is downregulated in our proband, whereas *LYST* is upregulated. Also *ACT2* is slightly upregulated. With regards to the predicted functional partners of the duplicated genes with a known role in language development, language impairment and/or language evolution, we found that neither of them was significantly dysregulated in the girl (i.e. with $FC > 2$ compared to both healthy parents). Only *HRAS* and *CMIP* could be regarded as slightly upregulated in our subject.

Figure 6. Variation in the expression levels of genes of interest in the proband's blood compared to her healthy parents (I). The graphic shows the results of the microarray

analyses for selected genes within the duplicated fragment and for their predicted functional partners according to String 10.5, as discussed in the text.

Lastly, we searched for additional candidates for the proband's language deficits, looking for genes exhibiting strong fold changes in our subject (i.e. with FC > 5 compared to both healthy parents). We found several strongly downregulated genes (*IL5RA*, *MYOM2*, *DDIT4* and *SLC29A1*), as well as several strongly upregulated genes (*CNTNAP3*, *WLS*, and *LGALS1*) (Figure 7).

Figure 7. Variation in the expression levels of genes of interest in the proband's blood compared to her healthy parents (II). The graphic shows the results of the microarray analyses for genes exhibiting fold changes > 5 compared to both healthy parents.

EDARADD is expressed at similar levels in all brain areas, whereas *LYST* is expressed at higher levels in the cerebellum, and *ACT2* is mostly expressed in the striatum. Regarding the putative functional partners of the genes located within the duplicated fragment in our proband, *HRAS* is more expressed in the striatum, whereas *CMIP* is similarly expressed in most brain areas. Finally, with regards to the genes that are strongly downregulated in our proband, we found that *MYOM2* and *SLC29A1* are most expressed in the cortex, whereas *DDIT4* is expressed at high levels in the cerebellum and the thalamus, and *IL5RA* is preferentially expressed in the hippocampus. No data were available regarding the three genes that are strongly upregulated in our proband.

Figure 8. Expression pattern in the blood and the brain of genes of interest. These genes include candidate genes within the duplicated fragment, predicted functional partners, and genes found strongly down- or upregulated in the proband, as discussed in the text. Expression levels of the genes were retrieved from the Genotype-Tissue Expression (GTEx) project (GTEx Consortium, 2013) (<https://www.gtexportal.org/home/>). Statistical analysis and data interpretation were performed by The GTEx Consortium Analysis Working Group using the . Legend: TPM, transcripts per million.

DISCUSSION

Next generation sequencing facilities, and particularly, comparative genomic hybridization arrays, have increased exponentially the number of genes and chromosomal regions associated to clinical conditions entailing cognitive and language deficits. However, in most cases, no robust genotype-to-phenotype links have been posited. In this paper, we have characterized the cognitive profile of a girl with a microduplication in 1q42.3q43, with a focus on her problems with language. The girl exhibits some of the cognitive, behavioral, and physical features commonly found in people with duplications of the distal 1q (Table 1). She bears one of the smallest

duplications of this terminal region of chromosome 1 described to date. Around 20% of DECIPHER patients with a duplication within, overlapping, or encompassing the region duplicated in our proband (N=35) are reported to suffer from language problems. Nonetheless, the duplicated fragments usually encompass more genes than in our proband and/or patients bear other chromosomal rearrangements. Importantly too, assessment of patients usually involves standardized cognitive tests only. Overall, to the best of our knowledge, this is the first detailed report on the language disabilities of a subject bearing only a short distal duplication of the 1 chromosome. For achieving this, we examined the proband's language strengths and deficits using a battery of specific diagnostic tools, but also conducted fine-grained analyses of language use in natural settings, with a focus on both structural and functional aspects.

In the expressive domain, the proband's speech is characterized by generalized simplification processes (sound assimilation, consonant cluster reduction, weak syllable deletion), which are typically found in younger children. Because our proband exhibits a noteworthy motor delay, we hypothesize that she suffers from some form of dysarthria. Nonetheless, the results of the RFI test point as well to a phonological impairment, which seemingly contributes to her speech problems. This possibility is reinforced by the low scores obtained by our proband in the ITPA categories for verbal auditory comprehension and association, as well as for sequential auditory memory. These low scores might be pointing to a verbal auditory memory deficit, which is expected to affect the normal acquisition of phonology and ultimately, explain her speech problems, including her tendency to reduce syllabic complexity.

Regarding word morphology and sentence syntax, both are seriously impaired. The MLUw in spontaneous speech is within the range of much younger typically-developing children. Problems are also observed in the receptive domain. Accordingly, her receptive vocabulary is much poorer than expected by age. Since the acquisition of phonology is the scaffold for the acquisition of other components of language, and since comprehension usually goes ahead of expressive abilities during development (Owens, 1984), we contend that the proband's phonological deficit can be indirectly causing most of her language problems, in both the receptive and the expressive domains.

This remarkable delay in the acquisition of core structural components of language dramatically hinders the child from expressing her needs and feelings, and to a lesser extent, understanding what is happening in her social environment, thus impeding a successful communication with her family and peers. Overall, we hypothesize that this severe language impairment would underlie her learning disabilities and her cognitive deficits. The fact that the proband shows personal living skills that are typical of her chronological age and that enable her to take care of herself supports this view. Moreover, contrary to what has been reported in other developmental disorders resulting in primary intellectual disability, she behaves adequately with strangers, without exhibiting excessive signs of affection. This might also explain the preservation of her pragmatic skills, as she shows metapragmatic awareness and follows the rules governing conversational exchanges, being able to understand and generate conversational implicatures despite her limited expressive resources.

Regarding the molecular causes of the attested problems with language, in silico analyses uncovered several potential candidates for language deficits among the genes

located within the duplicated fragment, as well as functional links of interest between some of the genes within the 1q42.3q43 region and robust candidates for language development and/or evolution. Although we have not found evidence of a strong dysregulation of any of these genes, microarray analyses suggest that some of them might be (slightly) up- or downregulated in the blood of our proband. Pretty obviously, pathological changes in

Among the genes located within the duplicated region, one promising candidate is *EDARADD*, which is found significantly downregulated in our proband. This gene encodes an interactor of EDAR, involved in the development of hair, teeth and other ectodermal derivatives. Mutations in *EDARADD* usually result in ectodermal dysplasia (OMIM# 614940; OMIM# 614941) (Headon et al., 2001; Bal et al., 2007; Cluzeau et al., 2019). The gene has been recently associated to a biomarker of ageing, namely, ‘epigenetic age acceleration’, which is predictive of morbidity and mortality (Gibson et al., 2019). It is not known whether the gene plays any significant role at the brain level, where it is expressed at very low levels.

A second promising candidate is *ACTN2*, which is found slightly upregulated in the proband. This gene encodes a bundling protein that anchors actin to different intracellular structures and which contributes to regulate spine morphology and the assembly of the post-synaptic density in neurons (Hodges et al. 2014), as well as the assembly and function of N-methyl-D-aspartate (NMDA) glutamate receptors, particularly in the striatum (Dunah et al., 2000; Bouhamdan et al., 2006). *ACTN2* is predicted to interact with robust candidates for language development, language deficits and/or language evolution, particularly, *GRIN2A*, *GRIN2B*, *HRAS*, *PVALB*, and *ITGB4*. *GRIN2A* and *GRIN2B* encode two components of the subunit NR2 of the

700 NMDA receptor channel, important for long-term potentiation, and ultimately, for
 701 memory and learning, and both are candidates for SLI, SZ, and ASD (Murphy and
 702 Benítez-Burraco, 2018). Additionally, mutations in *GRIN2A* give rise to epilepsy-
 703 aphasia spectrum disorders, including rolandic epilepsies, Landau-Kleffner syndrome,
 704 and continuous spike and waves during slow-wave sleep syndrome (CSWSS), which
 705 entail speech impairment and language regression (Carvill et al., 2013; Lesca et al.,
 706 2013). Speech problems associated to mutations in the gene include imprecise
 707 articulation, alteration of pitch and prosody, and dysarthria or dyspraxia (Turner et al.,
 708 2015), which recall many of the speech problems observed in our proband. Likewise,
 709 mutations in *GRIN2B* have been also found in individuals with cognitive dysfunction
 710 and EEG anomalies (Freunscht et al., 2013; Hu et al., 2016). *HRAS* encodes a GTPase
 711 important for neural growth and differentiation, long-term potentiation, and synaptic
 712 plasticity; the gene is a candidate for Costello syndrome (OMIM# # 218040,) a
 713 condition entailing developmental delay and mild to moderate intellectual impairment,
 714 with relatively impaired language, particularly in the expressive domain (Axelrad et al.,
 715 2011; Schwartz et al. 2013). *PVALB* encodes parvalbumin, a high affinity calcium ion-
 716 binding protein with an important role in brain function. Inhibition of parvalbumin-
 717 expressing interneurons results in complex behavioral changes, including altered
 718 sensorimotor gating, reduced fear extinction, and increased novelty-seeking (Brown et
 719 al., 2015). Additionally, the parvalbumin system might represent a convergent
 720 downstream endpoint for some forms of ASD, with reduced parvalbumin-expressing
 721 neurons resulting in shifting the excitation/inhibition balance towards enhanced
 722 inhibition (Filice et al., 2016). Finally, it should be mentioned that inactivating
 723 parvalbumin-positive (Pvalb+) interneurons in the auditory cortex alters normal
 724 response to sounds (specifically, it strengthens forward suppression and alters its

frequency dependence) (Phillips et al., 2017). This is interesting in view of the auditory impairment reported in some patients with similar duplications to the one found in our proband (e.g. DECIPHER patient 279513), but particularly, in view of the verbal auditory memory deficit exhibited by our proband. Lastly, *ITGB4* interacts with *FLNA* (Travis et al, 2004), an actin-binding protein involved in actin cytoskeleton remodeling and neuronal migration (Fox et al., 1998), which in turn binds regarding predictions based on co-expression data and experimental/biochemical data, the links between *HEATR1* and *NOP9*, and particularly, between *LYST* and *GABARAP* are also worth considering. *NOP9* is a candidate for language impairment (Pettigrew et al., 2016). *LYST*, which is found slightly upregulated in our proband, has been associated to cognitive decline (Kaplan et al., 2008; Introne et al., 2017). Lastly, *GABARAP* is a candidate for DD and encodes a GABAA receptor-associated protein, which contributes to the clustering of neurotransmitter receptors and to inhibitory neural transmission (Veerappa et al., 2013).

To finish, it is worth considering the biological roles played by the genes that are found more strongly dysregulated in the blood of our proband compared to her healthy parents. Among the genes found strongly downregulated, *IL5RA* is of interest considering the psychomotor problems exhibited by the girl. This gene encodes a subunit of a cytokine receptor and it has been associated to physical activity levels (Letsinger et al., 2019). More interestingly, *DDIT4* encodes a stress-response protein that functions as negative regulator of mTOR, a kinase that regulates cell growth and proliferation, and autophagy, as well as synaptic plasticity. Increased *DDIT4* levels are involved in aspects of neuronal damage, whereas knockdown of *DDIT4* seems to inhibit neuronal apoptosis (Su et al., 2019). More specifically, in rats, overexpression of the

750 gene in the prefrontal cortex results in anxiety- and depressive-like behaviors and
 751 neuronal atrophy, whereas mutant mice with a deletion of *DDIT4* are resilient to the
 752 behavioral and neuronal consequences of chronic stress (Ota et al., 2014). Abnormally
 753 high levels of the gene have been found in the prefrontal cortex of patients with major
 754 depressive disorder (MDD) (Ota et al., 2014). The molecular mechanism underlying
 755 depressant response in the prefrontal cortex also involves *CACNA1C*. In mice,
 756 knockout of *CACNA1C* in this brain region results in antidepressant-like behaviors,
 757 whereas overexpression of *DDIT4* reverses this effect (Kabier et al., 2017). *CACNA1C*
 758 is a candidate for multiple neuropsychiatric disorders including SZ, BD, and MDD
 759 (Ferreira et al., 2008; Green et al., 2010; Curtis et al., 2011; Cross-Disorder Group of
 760 the Psychiatric Genomics Consortium, 2013; Kabir et al., 2016). Variants of *CACNA1C*
 761 have been associated to deficits in reversal learning (Sykes et al., 2019), whereas some
 762 polymorphisms have been associated to decreased semantic verbal fluency in healthy
 763 subjects (Krug et al., 2010), as well as to decreased executive function in people with
 764 BD (Soeiro-de-Souza et al., 2013). *CACNA1C* expression has been found modulated
 765 during associative learning (Sykes et al., 2018). In mice, deletion of *CACNA1C* in
 766 glutamatergic neurons gives rise to reduced synaptic plasticity, sociability, and
 767 cognition (Dedic et al., 2018). Specifically, haploinsufficiency in the gene results in
 768 deficits in prosocial ultrasonic vocalization (Kisko et al., 2018; Redecker et al., 2019).
 769 In humans, haploinsufficiency in *CACNA1C* has been related to learning difficulties,
 770 expressive language impairment, and motor-skills delay (Mio et al., 2020). By contrast,
 771 in humans, *CACNA1C* gain-of-function mutations cause Timothy syndrome (OMIM#
 772 601005), which is featured by syndactyly, autism-like behavior, and language and
 773 social delays among other features (Splawski et al., 2004; Napolitano et al., 2015).
 774 Several of these features have been attested in our proband. Finally, *SLC29A1* encodes

a nucleoside transporter that localizes to the plasma and mitochondrial membranes, and that is involved in both nucleotide synthesis and cytotoxic nucleoside uptake. The gene contributes to basal synaptic transmission, long-term potentiation, neuronal plasticity, and spatial memory (Lee et al., 2018). More specifically, it is involved in glutamatergic neurotransmission (Xu et al., 2015) and the acquisition of goal-directed behavior (Nam et al., 2013). In mice, overexpression of the gene results in changes in dependent behavior, including a greater response to ethanol and a reduced response to caffeine (Kost et al., 2011).

Among the genes found strongly upregulated in our proband compared to her healthy parents, two of them are of particular interest considering her pathological features. *WLS* contributes to regulate the secretion of Wnt signaling molecules, which are involved in different developmental and homeostatic processes (Petko et al., 2019). Considering the motor problems exhibited by our proband, it is of particular interest that *WLS* is specifically involved in the assembly of the neuromuscular junction between motoneurons and skeletal muscles to control motor behaviour, with mutations in the gene resulting in muscle weakness and neurotransmission impairment (Shen et al., 2018). Interestingly too, *WLS* is also involved in the embryonic development of the cerebellum (Yeung and Goldowitz, 2017). A second promising candidate is *CNTNAP3*, which encodes a type of neurexin involved in cell recognition within the nervous system. This gene is found upregulated in the leukocytes of patients with SZ (Okita et al., 2017). The gene is expressed abundantly in many subcortical regions of the mouse brain, including the striatum, the globus pallidus and the subthalamic nucleus, which are crucially involved in speech and language processing, but also in many other cognitive and emotional functions (Booth et al., 2007; Kotz et al., 2009; Viñas-Guasch

and Wu, 2017). In mice, the knockout of *CNTNAP3* results in a delay in motor learning (Hirata et al., 2016), as well as repetitive behaviors, deficits in social interaction, and problems for spatial learning, which recapitulate many ASD-features (Tong et al., 2019). Interestingly too, *CNTNAP3* interacts with the synaptic adhesion protein *NRLG1*, which is also a candidate for ASD and whose mutation results as well in repetitive behaviors and abnormal corticostriatal synapses (Blundell *et al.*, 2010). Finally, we wish to highlight that most of the genes reviewed above are highly expressed in the cerebellum, which is a brain area crucially involved in motor planning, but also in language processing (Vias and Dick, 2017; Mariën and Borgatti, 2018).

CONCLUSIONS

In conclusion, although the exact molecular causes of language problems (and other cognitive, behavioral and even motor deficits) observed in our proband remain to be fully elucidated, we hypothesize that her distinctive features can result from the altered expression of specific genes within the duplicated 1q42.3q43, with a potential impact on selected functional interactors with a contrasted role in language development and/or evolution. Nonetheless, some other genes outside the duplicated region, that are found strongly dysregulated in the proband, can contribute to the attested impairment in the language domain.

We hope these findings contribute to a better understanding of the behavioral and cognitive phenotype resulting from CNVs of the distal region of the long arm of chromosome 1. A better understanding of the neurobiological foundations of the symptoms found in patients is needed for implementing more efficient pedagogical approaches and intervention strategies aimed to ameliorate their deficits and reinforce

825 their strengths. This is particularly true of low-prevalent conditions, such as this type
826 of CNVs, which are generally understudied and for which good management strategies
827 and intervention tools are not commonly available.

828

829

ACKNOWLEDGMENTS

We would like to thank the proband and her family for their participation in this research. We wish to thank also Dr. Montserrat Barcos Martínez and Dr. Isabel Espejo Portero, from the “Reina Sofía” hospital in Córdoba, for giving us access to the patient and her clinical history. Preparation of this work was supported by funds from the Spanish Ministry of Economy and Competitiveness (grant number FFI2016-78034-C2-2-P [AEI/FEDER,UE] to Antonio Benítez-Burraco, with MFU and SJR as members of the project).

ETHICS

Ethics approval for this research was granted by the Comité Ético del Hospital “Reina Sofía”. Written informed consent was obtained from the proband’s parents for conducting the psycholinguistic evaluations and the molecular analyses, and for publicizing this case report and all the accompanying tables and images in scientific journals and meetings.

AUTHORS’ CONTRIBUTION

ABB conceived the paper, performed the molecular experiments, analyzed the genomic/genetic data, and wrote the paper. MFU and SJR conducted the clinical evaluation of the proband, analyzed the psycholinguistic data, and wrote the paper. The three authors approved the final version of the manuscript.

REFERENCES

- Axelrad ME, Schwartz DD, Katzenstein JM, Hopkins E, Gripp KW (2011) Neurocognitive, adaptive, and behavioral functioning of individuals with Costello syndrome: a review. *Am J Med Genet C Semin Med Genet.* 157C:115-22.
- Bal E, Baala L, Cluzeau C, El Kerch F, Ouldim K, Hadj-Rabia S, Bodemer C, Munnich A, Courtois G, Sefiani A, Smahi A (2007) Autosomal dominant anhidrotic ectodermal dysplasias at the EDARADD locus. *Hum Mutat.* 28: 703-709
- Benítez-Burraco A, Boeckx C (2015): Possible functional links among brain- and skull-related genes selected in modern humans. *Front Psychol* 6:794.
- Benítez-Burraco, A., Murphy, E. (2016) The oscillopathic nature of language deficits in autism: from genes to language evolution. *Front. Hum. Neurosci.* 10: 120.
- Boeckx C, Benítez-Burraco A (2014a) The shape of the human language-ready brain. *Front Psychol* 5:282.
- Boeckx C, Benítez-Burraco A (2014b) Globularity and language-readiness: generating new predictions by expanding the set of genes of interest. *Front Psychol* 5: 1324
- Blundell J, Blaiss CA, Etherton MR, Espinosa F, Tabuchi K, Walz C, Bolliger MF, Südhof TC, Powell CM (2010) Neuroligin-1 deletion results in impaired spatial memory and increased repetitive behavior. *J Neurosci.* 30: 2115-2129. doi:10.1523/JNEUROSCI.4517-09.2010.

877

878 Booth JR, Wood L, Lu D, Houk JC, Bitan T (2007) The role of the basal ganglia and
879 cerebellum in language processing. *Brain Res.* 1133(1):136-44.

880

881 Bouhamdan M, Yan HD, Yan XH, Bannon MJ, Andrade R (2006) Brain-specific
882 regulator of G-protein signaling 9-2 selectively interacts with alpha-actinin-2 to
883 regulate calcium-dependent inactivation of NMDA receptors. *J Neurosci.* 26: 2522-30.

884

885 Brown JA, Ramikie TS, Schmidt MJ, Báldi R, Garbett K, Everheart MG, Warren LE,
886 Gellért L, Horváth S, Patel S, Mirnics K (2015) Inhibition of parvalbumin-expressing
887 interneurons results in complex behavioral changes. *Mol Psychiatry* 20:1499-507.

888

889 Carvill GL, Regan BM, Yendle SC, O'Roak BJ, Lozovaya N, Bruneau N, Burnashev
890 N, Khan A, Cook J, Geraghty E, Sadleir LG, Turner SJ, Tsai MH, Webster R, Ouvrier
891 R, Damiano JA, Berkovic SF, Shendure J, Hildebrand MS, Szepetowski P, Scheffer IE,
892 Mefford HC (2013) GRIN2A mutations cause epilepsy-aphasia spectrum disorders.
893 *Nat. Genet.* 45: 1073-6.

894

895 Chen XS, Reader RH, Hoischen A, Veltman JA, Simpson NH, Francks C, Newbury
896 DF, Fisher SE (2017) Next-generation DNA sequencing identifies novel gene variants
897 and pathways involved in specific language impairment. *Sci Rep.* 7:46105. doi:
898 10.1038/srep46105.

899

900 Cluzeau C, Marrakchi S, Picard C, Munnich A, Smahi A, Turki H (2019) First
901 homozygous large deletion in EDARADD gene associated with a severe form of

902 anhidrotic ectodermal dysplasia. *J Eur Acad Dermatol Venereol.* 33(2):e55-e57. doi:
903 10.1111/jdv.15182.
904
905 Conti-Ramsden G (1996) CLAN (Computerized Language Analysis). *Child Lang*
906 *Teach Ther.* 12: 345–349.
907
908 Cortada de Kohan N (2004) BAIRES. Test de Aptitud Verbal “Buenos Aires”. Madrid:
909 TEA Ediciones.
910
911 Cross-Disorder Group of the Psychiatric Genomics Consortium (2013) Identification
912 of risk loci with shared effects on five major psychiatric disorders: a genome-wide
913 analysis. *Lancet* 381, 1371–1379. doi:10.1016/S0140-6736(12)62129-1.
914
915 Curtis D, Vine AE, McQuillin A, Bass NJ, Pereira A, Kandaswamy R, Lawrence J,
916 Anjorin A, Choudhury K, Datta SR, Puri V, Krasucki R, Pimm J, Thirumalai S, Quesada
917 D, Gurling HM (2011). Case-case genome wide association analysis reveals markers
918 differentially associated with schizophrenia and bipolar disorder and implicates
919 calcium channel genes. *Psychiatr Genet* 21, 1–4. doi:10.1097/YPG.0b013e3283413382
920
921 Davies BL (2007) Grice’s Cooperative Principle: Meaning and rationality. *J Pragmat.*
922 39(12): 2308–2331 doi: 10.1016/j.pragma.2007.09.002
923
924 De La Cruz López MV, González Criado M. (2011) Adaptación española del inventario
925 de Desarrollo Battelle. Madrid: TEA
926

927 Dedic N, Pöhlmann ML, Richter JS, Mehta D, Czamara D, Metzger MW, Dine J,
928 Bedenk BT, Hartmann J, Wagner KV, Jurik A, Almli LM, Lori A, Moosmang S,
929 Hofmann F, Wotjak CT, Rammes G, Eder M, Chen A, Ressler KJ, Wurst W, Schmidt
930 MV, Binder EB, Deussing JM (2018) Cross-disorder risk gene CACNA1C
931 differentially modulates susceptibility to psychiatric disorders during development and
932 adulthood. *Mol. Psychiatry* 23: 533–543. doi:10.1038/mp.2017.133.
933
934 Diez-Itza E, Martínez V, Pérez V, Fernández-Urquiza M (2018) Explicit oral narrative
935 intervention for students with Williams syndrome. *Front Psychol* 8: 2337. doi:
936 10.3389/fpsyg.2017.02337
937
938 Dunah AW, Wyszynski M, Martin DM, Sheng M, Standaert DG (2000) Alpha-actinin-
939 2 in rat striatum: localization and interaction with NMDA glutamate receptor subunits.
940 *Brain Res Mol Brain Res.* 79:77-87.
941
942 Dunn L, Dunn LM, Arribas D. (2006) Peabody, test de vocabulario en imágenes.
943 Madrid: TEA Ediciones.
944
945 Fernández-Urquiza M, Viejo A, Cortiñas S, Huelmo J, Medina B, García I, Diez-Itza
946 E. (2015) Perfiles pragmáticos comparados de síndromes genéticos neuroevolutivos (S.
947 Williams, S. Down y S. X Frágil). In Diéguez-Vide F. (Ed.), *Temas de Lingüística*
948 *Clínica* (pp. 89–90). Barcelona: Horsori Editorial.
949
950 Fernández-Urquiza M, Diez-Itza E, Cortiñas S. (2017) PREP-CORP: sistema de
951 etiquetado pragmático de corpus clínicos de lengua oral. In: Fernández López MC,

952 Martí Sánchez M, Ruiz Martínez AM (Eds), Aplicaciones de la Lingüística (pp. 167-
953 83). Alcalá de Henares: Universidad de Alcalá.
954

955 Ferreira MA, O'Donovan MC, Meng YA, Jones IR, Ruderfer DM, Jones L, Fan J, Kirov
956 G, Perlis RH, Green EK, Smoller JW, Grozeva D, Stone J, Nikolov I, Chambert K,
957 Hamshire ML, Nimgaonkar VL, Moskvina V, Thase ME, Caesar S, Sachs GS, Franklin
958 J, Gordon-Smith K, Ardlie KG, Gabriel SB, Fraser C, Blumenstiel B, Defelice M,
959 Breen G, Gill M, Morris DW, Elkin A, Muir WJ, McGhee KA, Williamson R,
960 MacIntyre DJ, MacLean AW, St CD, Robinson M, Van Beck M, Pereira AC,
961 Kandaswamy R, McQuillin A, Collier DA, Bass NJ, Young AH, Lawrence J, Ferrier
962 IN, Anjorin A, Farmer A, Curtis D, Scolnick EM, McGuffin P, Daly MJ, Corvin AP,
963 Holmans PA, Blackwood DH, Gurling HM, Owen MJ, Purcell SM, Sklar P, Craddock
964 N; Wellcome Trust Case Control Consortium (2008) Collaborative genome-wide
965 association analysis supports a role for ANK3 and CACNA1C in bipolar disorder. Nat
966 Genet 40: 1056–1058. doi:10.1038/ng.209.
967

968 Filice F, Vörckel KJ, Sungur AÖ, Wöhr M, Schwaller B (2016) Reduction in
969 parvalbumin expression not loss of the parvalbumin-expressing GABA interneuron
970 subpopulation in genetic parvalbumin and shank mouse models of autism. Mol Brain
971 9: 10.
972

973 Fox JW, Lamperti ED, Ekşioğlu YZ, Hong SE, Feng Y, Graham DA, Scheffer IE,
974 Dobyns WB, Hirsch BA, Radtke RA, Berkovic SF, Huttenlocher PR, Walsh CA. (1998)
975 Mutations in filamin 1 prevent migration of cerebral cortical neurons in human
976 periventricular heterotopia. Neuron 21: 1315-25.

977

978 Freunscht I, Popp B, Blank R, Ende S, Moog U, Petri H, Prott EC, Reis A, Rübo J,
979 Zabel B, Zenker M, Hebebrand J, Wieczorek D (2013) Behavioral phenotype in five
980 individuals with de novo mutations within the GRIN2B gene. Behav Brain Funct. 9:
981 20. doi: 10.1186/1744-9081-9-20.

982

983 Gardner M (1987) *Test figura/palabra de vocabulario receptivo*. Argentina: Editorial
984 Médica Panamericana.

985

986 Gibson J, Russ TC, Clarke TK, Howard DM, Hillary RF, Evans KL, Walker RM,
987 Bermingham ML, Morris SW, Campbell A, Hayward C, Murray AD, Porteous DJ,
988 Horvath S, Lu AT, McIntosh AM, Whalley HC, Marioni RE (2019) A meta-analysis of
989 genome-wide association studies of epigenetic age acceleration. PLoS Genet.
990 15(11):e1008104. doi: 10.1371/journal.pgen.1008104.

991

992 Green EK, Grozeva D, Jones I, Jones L, Kirov G, Caesar S, Gordon-Smith K, Fraser
993 C, Forty L, Russell E, Hamshere ML, Moskvina V, Nikolov I, Farmer A, McGuffin P;
994 Wellcome Trust Case Control Consortium, Holmans PA, Owen MJ, O'Donovan MC,
995 Craddock N (2010) The bipolar disorder risk allele at CACNA1C also confers risk of
996 recurrent major depression and of schizophrenia. Mol Psychiatry 15: 1016–1022.
997 doi:10.1038/mp.2009.49.

998

999 Grice HP (1975) Logic and Conversation. In J. Cole, P., and Morgan (Ed.), Syntax and
1000 Semantics Volume 3: Speech Acts. (pp. 41–58). New York: Academic Press.

1001

1002 Headon DJ, Emmal SA, Ferguson BM, Tucker AS, Justice MJ, Sharpe PT, Zonana J,
1003 Overbeek PA (2001) Gene defect in ectodermal dysplasia implicates a death domain
1004 adapter in development. *Nature* 414: 913-916
1005
1006 Hemming IA, Forrest AR, Shipman P, Woodward KJ, Walsh P, Ravine DG, Heng JI
1007 (2016) Reinforcing the association between distal 1q CNVs and structural brain
1008 disorder: A case of a complex 1q43-q44 CNV and a review of the literature. *Am J Med*
1009 *Genet B Neuropsychiatr Genet.* 171B(3):458-67. doi: 10.1002/ajmg.b.32427.
1010
1011 Hirata H, Takahashi A, Shimoda Y, Koide T (2016) Caspr3-deficient mice exhibit low
1012 motor learning during the early phase of the accelerated rotarod task. *PLoS One*
1013 11(1):e0147887. doi: 10.1371/journal.pone.0147887.
1014
1015 Hodges JL, Vilchez SM, Asmussen H, Whitmore LA, Horwitz AR (2014) α -Actinin-2
1016 mediates spine morphology and assembly of the post-synaptic density in hippocampal
1017 neurons. *PLoS One* 9:e101770.
1018
1019 Hu C, Chen W, Myers SJ, Yuan H, Traynelis SF (2016) Human GRIN2B variants in
1020 neurodevelopmental disorders. *J Pharmacol Sci* 132: 115-121. doi:
1021 10.1016/j.jphs.2016.10.002.
1022
1023 Introne WJ, Westbroek W, Groden CA, Bhambhani V, Golas GA, Baker EH, Lehky
1024 TJ, Snow J, Ziegler SG, Malicdan MC, Adams DR, Dorward HM, Hess RA, Huizing
1025 M, Gahl WA, Toro C (2017) Neurologic involvement in patients with atypical Chediak-
1026 Higashi disease. *Neurology* 88:e57-e65.

1027

1028 Irizarry RA, Hobbs B, Collin F, Beazer-Barclay YD, Antonellis KJ, Scherf U, Speed
1029 TP (2003) Exploration, normalization, and summaries of high density oligonucleotide
1030 array probe level data. *Biostatistics* 4(2):249-64.

1031

1032 Juberg RC, Haney NR, Stallard R (1981) New deletion syndrome: 1q43. *Am J Hum*
1033 *Genet* 33(3):455–463.

1034

1035 Kabir ZD, Lee AS, Burgdorf CE, Fischer DK, Rajadhyaksha AM, Mok E, Rizzo B,
1036 Rice RC, Singh K, Ota KT, Gerhard DM, Schierberl KC, Glass MJ, Duman RS,
1037 Rajadhyaksha AM (2017) *Cacna1c* in the prefrontal cortex regulates depression-related
1038 behaviors via REDD1. *Neuropsychopharmacology* 42(10):2032-2042. doi:
1039 10.1038/npp.2016.271.

1040

1041 Kaplan J, De Domenico I, Ward DM (2008) Chediak-Higashi syndrome. *Curr Opin*
1042 *Hematol.* 15:22-9.

1043

1044 Kaufman AS, Kaufman NL (1990) *Kaufman Brief Intelligence Test manual*. Circle
1045 Pines, MN: American Guidance Service. (Spanish versión, K-BIT: *Test Breve de*
1046 *Inteligencia de Kaufman*. Madrid: Tea Ediciones (1996))

1047

1048 Kempisty B, Sikora J, Lianeri M, Szczepankiewicz A, Czerski P, Hauser J, Jagodzinski
1049 PP. (2007) MTHFD 1958G>A and MTR 2756A>G polymorphisms are associated with
1050 bipolar disorder and schizophrenia. *Psychiatr Genet.* 17:177-81.

1051

Kirk SA, McCarthy JJ (1961) The Illinois Test of Psycholinguistic Abilities: An approach to differential diagnosis. *Am J Ment Defic.* 66, 399–412.

Kirk SA, McCarthy JJ, Kirk WD (2009) Test de Illinois de Aptitudes Psicolingüísticas (7ma edición. ed.). (S. Ballesteros, & A. Cordero, Trads.) Madrid: TEA Ediciones.

Kisko TM, Braun MD, Michels S, Witt SH, Rietschel M, Culmsee C, Schwarting RKW, Wöhr M (2018). Cacna1c haploinsufficiency leads to pro-social 50-kHz ultrasonic communication deficits in rats. *Dis Model Mech* 11. doi:10.1242/dmm.034116.

Kost S, Sun C, Xiong W, Graham K, Cass CE, Young JD, Albensi BC, Parkinson FE (2011) Behavioral effects of elevated expression of human equilibrative nucleoside transporter 1 in mice. *Behav Brain Res.* 224(1):44-9. doi: 10.1016/j.bbr.2011.05.023.

Kotz SA, Schwartze M, Schmidt-Kassow M. (2009) Non-motor basal ganglia functions: a review and proposal for a model of sensory predictability in auditory language perception. *Cortex* 45(8):982-90. doi: 10.1016/j.cortex.2009.02.010.

Krug A, Nieratschker V, Markov V, Krach S, Jansen A, Zerres K, Eggermann T, Stöcker T, Shah NJ, Treutlein J, Mühleisen TW, Kircher T (2010) Effect of CACNA1C rs1006737 on neural correlates of verbal fluency in healthy individuals. *Neuroimage* 49: 1831–1836. doi:10.1016/j.neuroimage.2009.09.028.

1076 Lee CC, Chang CP, Lin CJ, Lai HL, Kao YH, Cheng SJ., Chen HM, Liao YP, Faivre
1077 E, Buée L, Blum D, Fang JM, Chern Y (2018) Adenosine augmentation evoked by an
1078 ENT1 inhibitor improves memory impairment and neuronal plasticity in the APP/PS1
1079 mouse model of Alzheimer's disease. *Mol Neurobiol.* 55(12):8936-8952. doi:
1080 10.1007/s12035-018-1030-z.
1081
1082 Lesca G, Rudolf G, Bruneau N, Lozovaya N, Labalme A, Boutry-Kryza N, Salmi M,
1083 Tsintsadze T, Addis L, Motte J, Wright S, Tsintsadze V, Michel A, Doummar D,
1084 Lascelles K, Strug L, Waters P, de Bellescize J, Vrielynck P, de Saint Martin A, Ville
1085 D, Ryvlin P, Arzimanoglou A, Hirsch E, Vincent A, Pal D, Burnashev N, Sanlaville D,
1086 Szeppetowski P (2013) GRIN2A mutations in acquired epileptic aphasia and related
1087 childhood focal epilepsies and encephalopathies with speech and language dysfunction.
1088 *Nat. Genet.* 45: 1061-6.
1089
1090 Letsinger AC, Granados JZ, Little SE, Lightfoot JT (2019) Alleles associated with
1091 physical activity levels are estimated to be older than anatomically modern humans.
1092 *PLoS One.* 14(4):e0216155. doi: 10.1371/journal.pone.0216155.
1093
1094 MacWhinney B (2000) The CHILDES Project: Tools for Analyzing Talk. Volume I:
1095 Transcription format and programs. Mahwah, NJ: Lawrence Erlbaum.
1096
1097 Mariën P, Borgatti R. (2018) Language and the cerebellum. *Handb Clin Neurol.*
1098 154:181-202.
1099

1100 McClay JL, Adkins DE, Aberg K, Bukszár J, Khachane AN, Keefe RS, Perkins DO,
 1101 McEvoy JP, Stroup TS, Vann RE, Beardsley PM, Lieberman JA, Sullivan PF, van den
 1102 Oord EJ (2011) Genome-wide pharmacogenomic study of neurocognition as an
 1103 indicator of antipsychotic treatment response in schizophrenia.
 1104 Neuropsychopharmacology 36: 616-26.
 1105
 1106 Mio C, Passon N, Baldan F, Bregant E, Monaco E, Mancini L, Demori E, Damante G
 1107 (2020) CACNA1C haploinsufficiency accounts for the common features of interstitial
 1108 12p13.33 deletion carriers. Eur J Med Genet 103843. doi:10.1016/j.ejmg.2020.103843.
 1109
 1110 Monfort M, Juarez A (1988) Registro fonológico inducido. Rev. de Logop. Foniatr. y
 1111 Audiol. 8: 88-96.
 1112
 1113 Montero D (1996) *Evaluación de la Conducta Adaptativa en Personas con*
 1114 *Discapacidades. Adaptación y validación del ICAP*. Bilbao: Mensajero.
 1115
 1116 Montero D. (1996) Evaluación de la conducta adaptativa en personas con
 1117 discapacidades. Adaptación y validación del ICAP. Bilbao: Mensajero.
 1118
 1119 Morris ML, Baroneza JE, Teixeira P, Medina CT, Cordoba MS, Versiani BR, Roese
 1120 LL, Freitas EL, Fonseca AC, Dos Santos MC, Pic-Taylor A, Rosenberg C, Oliveira SF,
 1121 Ferrari I, Mazzeu JF (2016) Partial 1q duplications and associated phenotype. Mol
 1122 Syndromol. 6(6):297-303. doi: 10.1159/000443599.
 1123

1124 Murphy E, Benítez-Burraco A (2016) Bridging the gap between genes and language
 1125 deficits in schizophrenia: An oscillopathic approach. *Front. Hum. Neurosci.* 10: 422.
 1126

1127 Murphy E, Benítez-Burraco A (2017) Language deficits in schizophrenia and autism as
 1128 related oscillatory connectomopathies: an evolutionary account. *Neurosci Biobehav*
 1129 *Rev.* 83:742-764.
 1130

1131 Murphy E, Benítez-Burraco A (2018) Toward the language oscillogenome. *Front*
 1132 *Psychol.* 9:1999. doi: 10.3389/fpsyg.2018.01999.
 1133

1134 Nam HW, Hinton DJ, Kang NY, Kim T, Lee MR, Oliveros A, Adams C, Ruby CL,
 1135 Choi DS (2013) Adenosine transporter ENT1 regulates the acquisition of goal-directed
 1136 behavior and ethanol drinking through A2A receptor in the dorsomedial striatum. *J*
 1137 *Neurosci.* 33(10):4329-38. doi: 10.1523/JNEUROSCI.3094-12.2013.
 1138

1139 Napolitano C, Splawski I, Timothy KW, Bloise R, Priori SG (2015) Timothy
 1140 Syndrome. University of Washington, Seattle Available at:
 1141 <https://www.ncbi.nlm.nih.gov/books/NBK1403/> [Accessed July 28, 2019].
 1142

1143 Newbury DF, Winchester L, Addis L, Paracchini S, Buckingham LL, Clark A, Cohen
 1144 W, Cowie H, Dworzynski K, Everitt A, Goodyer IM, Hennessy E, Kindley AD, Miller
 1145 LL, Nasir J, O'Hare A, Shaw D, Simkin Z, Simonoff E, Slonims V, Watson J, Ragoussis
 1146 J, Fisher SE, Seckl JR, Helms PJ, Bolton PF, Pickles A, Conti-Ramsden G, Baird G,
 1147 Bishop DV, Monaco AP (2009) CMIP and ATP2C2 modulate phonological short-term

1148 memory in language impairment. *Am J Hum Genet.* 85(2):264-72. doi:
 1149 10.1016/j.ajhg.2009.07.004.
 1150
 1151 Okita M, Yoshino Y, Iga JI, Ueno SI (2017) Elevated mRNA expression of CASPR3
 1152 in patients with schizophrenia. *Nord J Psychiatry.* 71(4):312-314. doi:
 1153 10.1080/08039488.2017.1291720.
 1154
 1155 Ota KT, Liu RJ, Voleti B, Maldonado-Aviles JG, Duric V, Iwata M, Duthiel S, Duman
 1156 C, Boikess S, Lewis DA, Stockmeier CA, DiLeone RJ, Rex C, Aghajanian GK, Duman
 1157 RS (2014) REDD1 is essential for stress-induced synaptic loss and depressive behavior.
 1158 *Nat Med.* 20(5):531-5. doi: 10.1038/nm.3513.
 1159
 1160 Owens RE Jr. (1984) *Language Development: An Introduction*. London: Pearson
 1161
 1162 Paracchini S, Diaz R, Stein J (2016) Advances in dyslexia genetics—new insights into
 1163 the role of brain asymmetries. In *Advances in Genetics* 96 (Friedmann T, Dunlap JC,
 1164 Goodwin SF, eds.), Academic Press: London, pp. 53–97.
 1165
 1166 Petko J, Thileepan M, Sargen M, Canfield V, Levenson R (2019) Alternative splicing
 1167 of the Wnt trafficking protein, Wntless and its effects on protein-protein interactions.
 1168 *BMC Mol Cell Biol.* 2019 Jul 8;20(1):22. doi: 10.1186/s12860-019-0208-1.
 1169
 1170 Petko J, Thileepan M, Sargen M, Canfield V, Levenson R (2019) Alternative splicing
 1171 of the Wnt trafficking protein, Wntless and its effects on protein-protein interactions.
 1172 *BMC Mol Cell Biol.* 20(1):22. doi: 10.1186/s12860-019-0208-1.

1173

1174 Pettigrew KA, Frinton E, Nudel R, Chan MT, Thompson P, Hayiou-Thomas ME,
 1175 Talcott JB, Stein J, Monaco AP, Hulme C, Snowling MJ, Newbury DF, Paracchini S
 1176 (2016) Further evidence for a parent-of-origin effect at the NOP9 locus on language-
 1177 related phenotypes. *J Neurodev Disord.* 8: 24. doi: 10.1186/s11689-016-9157-6.

1178

1179 Phillips EAK, Schreiner CE, Hasenstaub AR (2017) Cortical interneurons differentially
 1180 regulate the effects of acoustic context. *Cell Rep* 20: 771-778.

1181

1182 Prasad A, Merico D, Thiruvahindrapuram B, Wei J, Lionel AC, Sato D, Rickaby J, Lu
 1183 C, Szatmari P, Roberts W, Fernandez BA, Marshall CR, Hatchwell E, Eis PS, Scherer
 1184 SW (2012) A discovery resource of rare copy number variations in individuals with
 1185 autism spectrum disorder. *G3 (Bethesda).* 2:1665-85.

1186

1187 Ramos J-L, Cuetos F (2011) PROLEC-SE. Evaluación de los Procesos Lectores.
 1188 Manual. Madrid: TEA Ediciones

1189

1190 Redecker TM, Kisko TM, Schwarting RKW, Wöhr M (2019) Effects of *Cacna1c*
 1191 haploinsufficiency on social interaction behavior and 50-kHz ultrasonic vocalizations
 1192 in adult female rats. *Behav Brain Res* 367: 35–52. doi:10.1016/j.bbr.2019.03.032.

1193

1194 Rice, M. L., Smolik, F., Perpich, D., Thompson, T., Rytting, N., & Blossom, M. (2010).
 1195 Mean length of utterance levels in 6-month intervals for children 3 to 9 years with and
 1196 without language impairments. *J Speech Lang Hear Res* 53(2): 333–349. doi:
 1197 10.1044/1092-4388(2009/08-0183)

1198

1199 Roberts AE, Cox GF, Kimonis V, Lamb A, Irons M (2004) Clinical presentation of 13
1200 patients with subtelomeric rearrangements and a review of the literature. *Am J Med*
1201 *Genet Part A* 128A(4):352–363.

1202

1203 Rollins B, Martin MV, Morgan L, Vawter MP (2010) Analysis of whole genome
1204 biomarker expression in blood and brain. *Am J Med Genet B Neuropsychiatr Genet.*
1205 153B: 919-36.

1206

1207 Sacks H, Schegloff E, Jefferson G (1974) A simplest systematics for the organization
1208 of turn-taking for conversation. *Language* 50(4): 696–735.

1209

1210 Schwartz DD, Katzenstein JM, Hopkins E, Stabley DL, Sol-Church K, Gripp KW,
1211 Axelrad ME (2013) Verbal memory functioning in adolescents and young adults with
1212 Costello syndrome: evidence for relative preservation in recognition memory. *Am J*
1213 *Med Genet A.* 161A:2258-65.

1214

1215 Shen C, Li L, Zhao K, Bai L, Wang A, Shu X, Xiao Y, Zhang J, Zhang K, Hui T, Chen
1216 W, Zhang B, Hsu W, Xiong WC, Mei L (2018) Motoneuron Wnts regulate
1217 neuromuscular junction development. *Elife* 7. pii: e34625. doi: 10.7554/eLife.34625.

1218

1219 Silipigni R, Monfrini E, Baccarin M, Giangiobbe S, Lalatta F, Gueneri S, Bedeschi
1220 MF (2017) Familial duplication/deletion of 1q42.13q43 as meiotic consequence of an
1221 intrachromosomal insertion in chromosome 1. *Cytogenet Genome Res.* 153(2):73-80.
1222 doi: 10.1159/000485226.

1223

1224 Soeiro-de-Souza MG, Bio DS, Dias VV, Vieta E, Machado-Vieira R, Moreno RA

1225 (2013) The CACNA1C risk allele selectively impacts on executive function in bipolar

1226 type I disorder. *Acta Psychiatr Scand* 128: 362–369. doi:10.1111/acps.12073.

1227

1228 Splawski I, Timothy KW, Sharpe LM, Decher N, Kumar P, Bloise R, Napolitano C,

1229 Schwartz PJ, Joseph RM, Condouris K, Tager-Flusberg H, Priori SG, Sanguinetti MC,

1230 Keating MT (2004) CaV1.2 calcium channel dysfunction causes a multisystem disorder

1231 including arrhythmia and autism. *Cell* 119: 19–31. doi:10.1016/j.cell.2004.09.011.

1232

1233 Su J, Wang M, Yan Y, Ju S, Chen J, Wu X (2019) Increased REDD1 facilitates neuronal

1234 damage after subarachnoid hemorrhage. *Neurochem Int.* 128:14-20. doi:

1235 10.1016/j.neuint.2019.03.019.

1236

1237 Sullivan PF, Fan C, Perou CM (2006) Evaluating the comparability of gene expression

1238 in blood and brain. *Am J Med Genet B Neuropsychiatr Genet.* 141B: 261–268.

1239

1240 Sykes L, Clifton NE, Hall J, Thomas KL (2018) Regulation of the expression of the

1241 psychiatric risk gene *Cacna1c* during associative learning. *MNP* 4: 149–157.

1242 doi:10.1159/000493917.

1243

1244 Sykes L, Haddon J, Lancaster TM, Sykes A, Azzouni K, Ihssen N, Moon AL, Lin TE,

1245 Linden DE, Owen MJ, O'Donovan MC, Humby T, Wilkinson LS, Thomas KL, Hall J

1246 (2019) Genetic Variation in the Psychiatric Risk Gene CACNA1C Modulates Reversal

1247 Learning Across Species. *Schizophr Bull* 45: 1024–1032. doi:10.1093/schbul/sby146.

1248

1249 Szklarczyk D, Franceschini A, Wyder S, Forslund K, Heller D, Huerta-Cepas J,
1250 Simonovic M, Roth A, Santos A, Tsafou KP, Kuhn M, Bork P, Jensen LJ, von Mering
1251 C (2015) STRING v10: protein-protein interaction networks, integrated over the tree of
1252 life. *Nucleic Acids Res.* 43, D447-52.

1253

1254 Tong DL, Chen RG, Lu YL, Li WK, Zhang YF, Lin JK, He LJ, Dang T, Shan SF, Xu
1255 XH, Zhang Y, Zhang C, Du YS, Zhou WH, Wang X, Qiu Z (2019) The critical role of
1256 ASD-related gene CNTNAP3 in regulating synaptic development and social behavior
1257 in mice. *Neurobiol Dis.* 130:104486. doi: 10.1016/j.nbd.2019.104486.

1258

1259 Travis MA, van der Flier A, Kammerer RA, Mould AP, Sonnenberg A, Humphries MJ
1260 (2004) Interaction of filamin A with the integrin beta 7 cytoplasmic domain: role of
1261 alternative splicing and phosphorylation. *FEBS Lett.* 569(1-3):185-90

1262

1263 Turner SJ, Mayes AK, Verhoeven A, Mandelstam SA, Morgan AT, Scheffer IE (2015)
1264 GRIN2A: an aptly named gene for speech dysfunction. *Neurology* 84: 586-93.

1265

1266 van Bon BW, Koolen DA, Borgatti R, Magee A, Garcia-Minaur S, Rooms L, Reardon
1267 W, Zollino M, Bonaglia MC, De Gregori M, Novara F, Grasso R, Ciccone R, van
1268 Duyvenvoorde HA, Aalbers AM, Guerrini R, Fazzi E, Nillesen WM, McCullough S,
1269 Kant SG, Marcelis CL, Pfundt R, de Leeuw N, Smeets D, Sistermans EA, Wit JM,
1270 Hamel BC, Brunner HG, Kooy F, Zuffardi O, de Vries BB (2008) Clinical and
1271 molecular characteristics of 1qter microdeletion syndrome: Delineating a critical region
1272 for corpus callosum agenesis/hypogenesis. *J Med Genet* 45(6): 346–354.

1273

1274 Veerappa AM, Saldanha M, Padakannaya P, Ramachandra NB (2013) Family-based
1275 genome-wide copy number scan identifies five new genes of dyslexia involved in
1276 dendritic spinal plasticity. *J Hum Genet.* 58: 539-47.

1277

1278 Vias C, Dick AS (2017) Cerebellar contributions to language in typical and atypical
1279 development: a review. *Dev Neuropsychol.* 42(6):404-421.

1280

1281 Viñas-Guasch N, Wu YJ (2017) The role of the putamen in language: a meta-analytic
1282 connectivity modeling study. *Brain Struct Funct.* 222(9):3991-4004. doi:
1283 10.1007/s00429-017-1450-y.

1284

1285 Wang J, Fu C, Zhang S, Luo J, Ouyang L, Xie B, Sun W, He S, Su J, Hu X, Fei D,
1286 Chen R, Fan X, Ou S, Chen S, Shen Y (2016) A rare occurrence of two large de novo
1287 duplications on 1q42-q44 and 9q21.12-q21.33. *Gene* 594: 59–65

1288

1289 Wechsler D (2002) WAIS – III. Escala de inteligencia para adultos de Wechsler-
1290 Tercera edición. Buenos Aires: Paidós.

1291

1292 Witt SH, Sommer WH, Hansson AC, Sticht C, Rietschel M, Witt CC (2013)
1293 Comparison of gene expression profiles in the blood, hippocampus and prefrontal
1294 cortex of rats. *In Silico Pharmacol.* 1: 15.

1295

1296 Xu Z, Xu P, Chen Y, Liu J, Zhang Y, Lv Y, Luo J, Fang M, Zhang J, Wang J, Wang
1297 K, Wang X, Chen G (2015) ENT1 inhibition attenuates epileptic seizure severity via

1298 regulation of glutamatergic neurotransmission. *Neuromolecular Med.* 17(1):1-11. doi:
1299 10.1007/s12017-014-8338-2.
1300
1301 Yeung J, Goldowitz D (2017) Wls expression in the rhombic lip orchestrates the
1302 embryonic development of the mouse cerebellum. *Neuroscience* 354:30-42. doi:
1303 10.1016/j.neuroscience.2017.04.020.
1304
1305 Yuste C, Martínez MR (2005) *La Batería de Aptitudes Diferenciales y Generales*
1306 *(BADyG)*. Madrid: CEPE.
1307
1308
1309

SUPPLEMENTAL DATA

Supplemental file 1. Tables with the scores obtained by the proband in the standardized tests used for her assessment.

Supplemental file 2. Transcriptions of the conversational exchanges in natural settings used for the analyses of the proband's language in use.

Supplemental file 3. The whole set of genes important for language used for the in silico analysis with String 10.5.

Supplemental file 4. Data resulting from the microarray analyses comparing gene expression levels in the blood of the proband and her healthy parents.

Supplemental file 5. Developmental expression profiles of genes of interest. These genes include candidate genes within the duplicated fragment, predicted functional partners, and genes found strongly down- or upregulated in the proband, as discussed in the paper. The expression data are from the Human Brain Transcriptome Database (<http://hbatlas.org/>). Six different brain regions are considered: the cerebellar cortex (CBC), the mediodorsal nucleus of the thalamus (MD), the striatum (STR), the amygdala (AMY), the hippocampus (HIP) and 11 areas of neocortex (NCX).