Title

Generation of human bronchial organoids for SARS-CoV-2 research

Authors/Affiliations

Tatsuya Suzuki¹

Yumi Itoh¹

Yusuke Sakai²

Akatsuki Saito³

Daisuke Okuzaki^{4,5,6}

Daisuke Motooka⁷

Shohei Minami⁸

Takeshi Kobayashi⁸

Takuya Yamamoto^{9,10,11,12}

Toru Okamoto^{1*}

Kazuo Takayama^{12*}

¹ Institute for Advanced Co-Creation Studies, Research Institute for Microbial Diseases, Osaka University, Suita 565-0871, Japan

² Laboratory of Veterinary Pathology, Joint Faculty of Veterinary Medicine, Yamaguchi University, Yamaguchi 753-8511, Japan

³ Department of Veterinary Science, Faculty of Agriculture, University of Miyazaki, Miyazaki 889-2192, Japan

⁴ Genome Information Research Center, Research Institute for Microbial Diseases, Osaka University, Suita 565-0871, Japan

25 ⁵ Single Cell Genomics, Human Immunology, WPI Immunology Frontier Research
 26 Center, Osaka University, Suita 565-0871, Japan
 27 ⁶ Institute for Open and Transdisciplinary Research Initiatives, Osaka University, Suita
 28 565-0871, Japan
 29 ⁷ Genome Information Research Center, Research Institute for Microbial Diseases,
 30 Osaka University, Suita 565-0871, Japan
 31 ⁸ Laboratory of Viral Replication, International Research Center for Infectious Diseases,
 32 Research Institute for Microbial Diseases, Osaka University, Suita, Osaka, 565-0871
 33 Japan
 34 ⁹ Institute for the Advanced Study of Human Biology (WPI-ASHBi), Kyoto University,
 35 Kyoto 606-8501 Japan
 36 ¹⁰ Medical-risk Avoidance based on iPS Cells Team, RIKEN Center for Advanced
 37 Intelligence Project (AIP), Kyoto 606-8507, Japan
 38 ¹¹ AMED-CREST, Japan Agency for Medical Research and Development (AMED),
 39 Tokyo 100-0004, Japan
 40 ¹² Center for iPS Cell Research and Application (CiRA), Kyoto University, Kyoto
 41 606-8507, Japan
 42
 43 ***Corresponding author**
 44 Dr. Kazuo Takayama
 45 Center for iPS Cell Research and Application, Kyoto University, Shogoin Kawaharacho
 46 53, Sakyo-ku, Kyoto 606-8397, Japan
 47 Phone: +81-75-366-7362, FAX: +81-75-366-7074
 48 E-mail: kazuo.takayama@cira.kyoto-u.ac.jp

49

50 Dr. Toru Okamoto

51 Institute for Advanced Co-Creation Studies, Research Institute for Microbial Diseases,

52 Osaka University, Yamadaoka 3-1, Suita 565-0871, Japan

53 Phone: +81-6-6879-8330, FAX: +81-6-6879-8330

54 E-mail: toru@biken.osaka-u.ac.jp

55

56 **Abbreviations**

57 2D two-dimensional

58 ACE2 angiotensin-converting enzyme 2

59 CC10 club cell protein 10

60 FGF fibroblast growth factor

61 hBEpC human bronchial epithelial cells

62 hBO human bronchial organoids

63 IFN-I type I interferon

64 IHC immunohistochemistry

65 KRT5 keratin 5

66 LDH lactate dehydrogenase

67 PSC pluripotent stem cell

68 RdRp RNA-dependent RNA polymerase

69 RNA seq RNA sequencing

70 SARS-CoV-2 severe acute respiratory syndrome coronavirus 2

71 TMPRSS2 transmembrane serine proteinase 2

72 WHO World Health Organization

73

74 **Abstract**

75 Coronavirus disease 2019 (COVID-19) is a disease that causes fatal disorders
76 including severe pneumonia. To develop a therapeutic drug for COVID-19, a model that
77 can reproduce the viral life cycle and evaluate the drug efficacy of anti-viral drugs is
78 essential. In this study, we established a method to generate human bronchial organoids
79 (hBO) from commercially available cryopreserved human bronchial epithelial cells and
80 examined whether they could be used as a model for severe acute respiratory syndrome
81 coronavirus 2 (SARS-CoV-2) research. Our hBO contain basal, club, ciliated, and
82 goblet cells. Angiotensin-converting enzyme 2 (ACE2), which is a receptor for
83 SARS-CoV-2, and transmembrane serine proteinase 2 (TMPRSS2), which is an
84 essential serine protease for priming spike (S) protein of SARS-CoV-2, were highly
85 expressed. After SARS-CoV-2 infection, not only the intracellular viral genome, but
86 also progeny virus, cytotoxicity, pyknotic cells, and moderate increases of the type I
87 interferon signal could be observed. Treatment with camostat, an inhibitor of TMPRSS2,
88 reduced the viral copy number to 2% of the control group. Furthermore, the gene
89 expression profile in SARS-CoV-2-infected hBO was obtained by performing RNA-seq
90 analysis. In conclusion, we succeeded in generating hBO that can be used for
91 SARS-CoV-2 research and COVID-19 drug discovery.

92

93

94 **Graphical abstract**

95
96
97 **Key words**

98 SARS-CoV-2, bronchial organoids, COVID-19, camostat

99

Introduction

The “2019-new coronavirus disease (COVID-19) was first reported in China in December 2019¹ and declared a pandemic by the WHO in March 2020². Severe pneumonia is most frequently observed in COVID-19 patients, and the number of COVID-19 patients and deaths are still increasing. These conditions have made it difficult for research on severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), which is the causative virus of COVID-19, to keep pace. SARS-CoV-2 is composed of four proteins: S (spike), E (envelope), M (membrane), and N (nucleocapsid) proteins. It is known that angiotensin-converting enzyme 2 (ACE2) is a SARS-CoV-2 receptor, and transmembrane serine proteinase 2 (TMPRSS2) is essential for priming S protein³. Thus, to accelerate SARS-CoV-2 research, a novel lung model that reproduces the viral life cycle with intact expression of these host factors is indispensable.

A number of animal and cell models that can be used for SARS-CoV-2 research have been reported (Takayama et al., in revision), but an *in vitro* lung model that can evaluate candidate therapeutic agents for COVID-19 is essential for conducting large-scale drug screening. Human lung organoids are excellent tools that can faithfully mimic the lung functions of living organisms⁴. The lungs consist of bronchi and alveoli. Lukassen et al. have performed single-cell analysis of alveoli and bronchi and reported that ACE2 is predominantly expressed in the transient secretory cell type, which displays a transient cell state between goblet and ciliated cells, in bronchi, but TMPRSS2 is strongly expressed in both lung tissues⁵. Therefore, a bronchial organoid containing transient secretory, goblet, and ciliated cells could be a useful model for

SARS-CoV-2 research.

Several reports have verified the usefulness of two-dimensional (2D) culturing airway epithelial cells in SARS-CoV2 studies. For example, SARS-CoV-2 infection experiments using human bronchial epithelial cells (hBEpC) showed cytopathic effects 96 hr after the infection on the layers of human airway epithelial cells ⁶. In addition, hBEpC cultured using an air-liquid interface culture system can be used to evaluate viral infection, replication, and the drug efficacy of remdesivir ⁷. 2D culture systems of hBEpC are relatively easy to use, but they cannot reproduce the cellular microenvironment in the living body and are difficult to use for long-time culture. Recently, it was shown that SARS-CoV-2 can infect and replicate in human pluripotent stem cell (PSC)-derived lung organoids containing bronchial epithelial cells and alveolar epithelial cells ⁸. However, these organoids exhibit a fetal phenotype rather than an adult type ^{9,10}. Adult-type bronchial organoids are essential because of the severe infection caused by COVID-19 in adults. Human bronchial organoids (hBO) with adult phenotype can be established from intact human lung tissue ¹¹. However, it is difficult for many researchers to obtain an intact lung biopsy sample, because the process requires the approval of an ethics committee and informed consent from the donor. Therefore, in this study, we developed a method for generating hBO from commercially available cryopreserved adult hBEpC and applied it to SARS-CoV-2 research.

Results

Generation of human bronchial organoids from cryopreserved adult bronchial epithelial cells

We searched for the conditions that could establish hBO from cryopreserved adult

hBEpC. We found that after embedding the hBEpC in Matrigel and culturing with advanced DMEM/F12 medium containing FGF2, FGF7, FGF10, Noggin, R-spondin 1, Y-27632, and SB202190 (expansion medium), hBO could be established (**Table S1**). Furthermore, we could mature the hBO by culturing them with advanced DMEM/F12 medium containing FGF2, FGF7, FGF10, Y-27632, and A83-01 (differentiation medium) (**Table S1**). Among the growth factors included in the differentiation medium, FGF2 is important for enhancing the expression levels of *ACE2* and *TMPRSS2* (**Fig. S1**). Approximately 100 hBO were present in 50 μ L of Matrigel, and the diameter of each hBO was around 100-200 μ m (**Fig. 1A**). Transmission electron microscopy (TEM) images showed the presence of cilia and goblet cells (**Fig. 1B, Fig. S2**). The *ACE2* and *TMPRSS2* expression levels in hBO were higher than in cryopreserved adult hBEpC (**Fig. 1C**). Immunohistochemical analysis showed that ACE2 was expressed in part of the outer edge of hBO, while TMPRSS2 was expressed in part of the outer edge and lumen (**Fig. 1D, Fig. S3**). Because bronchi are composed of basal, ciliated, goblet, and club cells, a gene expression analysis of markers specific to these four cell types was performed. The gene expression levels of basal, ciliated, goblet, and club cell markers in hBO were higher than in cryopreserved adult hBEpC (**Figs. 1E-1H**). Consistently, hBO were also positive for α -tubulin, CC10, mucin 5AC, and KRT5 (**Fig. 1I**). The outer edge and lumen of hBO was positive for KRT5 and acetylated α -tubulin, respectively. These results suggest that basal cells are positive for both ACE2 and TMPRSS2, but ciliated cells are positive only for TMPRSS2. Based on these observations, we succeeded in generating expandable and functional hBO from cryopreserved adult hBEpC.

RNA-seq analysis of human bronchial organoids

RNA-seq analysis was performed to further characterize hBO. A heat map, principal component analysis (PCA), and scatter plot of gene expression profiles all show hBO is closer to hBEpC than to A549 cells (**Figs. 2A-2C**). A second heat map of bronchial epithelial markers showed that hBO expressed bronchial epithelial markers more strongly than did hBEpC or A549 cells (**Fig. 2D**). These results suggest that hBO have higher bronchial functions than A549 cells or hBEpC.

SARS-CoV-2 infection experiments using human bronchial organoids

Next, we investigated whether hBO can be applied to SARS-CoV-2 research. HBO were infected with SARS-CoV-2 and then cultured in differentiation medium for 5 days (**Fig. 3A**). We observed a significant accumulation of LDH in the culture medium of infected hBO (**Fig. 3B**), suggesting that cytotoxicity was caused by the infection. At day 5 after the infection, viral gene expression in infected hBO was clearly detected (**Fig. 3C**). IHC analysis showed that S protein-positive cells were observed in part of the outer edge of hBO (**Fig. 3D**). In addition, S protein co-localized with KRT5 (**Fig. 3E**), but not with CC10 (**Fig. S4**), suggesting that SARS-CoV-2 infected and replicated in basal cells. Overall, these results indicate that SARS-CoV-2 can infect and replicate in hBO.

Currently, clinical trials using camostat, favipiravir, nafamostat, chloroquine, ritonavir/lopinavir, and remdesivir are underway around the world to develop therapeutic agents for COVID-19. However, the evaluation of these drugs using *in vitro* lung models is rare. Therefore, in this study, the effect of camostat, an inhibitor of TMPRSS2, was examined using our hBO because camostat was demonstrated as a promising candidate in cell culture models³. Upon camostat treatment, the amount of

SARS-CoV-2 viral genome was reduced to 2% of untreated infected hBO (**Fig. 3C**). In addition, LDH release from infected hBO was significantly reduced after camostat treatment (**Fig. 3B**). Finally, we examined the culture supernatants of infected hBO. We found that infectious virus was significantly observed in infected hBO, but the production of infectious virus was impaired by treatment with camostat (**Fig. 3F**). Collectively, our data indicated that hBO can secrete infectious virus into the culture medium, suggesting that our hBO system can investigate the entire life cycle of SARS-CoV-2.

Next, we examined the pathological effects of the SARS-CoV-2 infection. We observed that the number of pyknotic cells seemed to increase with the infection (**Fig. 3G**). In addition, the expression levels of type I IFN (IFN-I) and IFN-stimulated genes were moderately increased after SARS-CoV-2 infection (**Fig. S5A**). Furthermore, SARS-CoV-2 infection did not change the gene expression levels of *ACE2* or *TMPRSS2* (**Fig. S5B**). From the above, these results suggest that hBO can be used to reproduce SARS-CoV-2-induced pulmonary disorder and to evaluate the effect of therapeutic agents.

RNA-seq analysis of SARS-CoV-2-infected human bronchial organoids

RNA-seq analysis was performed to investigate the effects of SARS-CoV-2 infection and camostat treatment in detail. A heat map shows that the gene expression profile of SARS-CoV-2-infected hBO is closer to SARS-CoV-2-infected hBO treated with camostat than uninfected hBO (**Fig. 4A**). A PCA and scatter plot of the gene expressions agree with this finding (**Figs. 4B, C**). Additionally, SARS-CoV-2 infection increased the expression levels of IFN-I signaling-related genes in hBO (**Fig. 4D**).

PGSEA applied on GO biological process gene sets shows that the expression levels of genes involved in positive regulation of immune effector process, regulation of inflammatory response, interferon-gamma production, and positive regulation of acute inflammatory response were increased by SARS-CoV-2 infection and that this increase was suppressed by camostat treatment (**Fig. 4E**). These results indicated that SARS-CoV-2 infection induces IFN-I signaling-related genes and that camostat treatment reversed this phenotype.

Discussion

In this study, we succeeded in generating hBO from cryopreserved adult hBEpC and applied it to SARS-CoV-2 research. We confirmed that SARS-CoV-2 could infect and replicate in these cells and that camostat could suppress the replication. If small airway organoids and alveolar organoids can be produced from cryopreserved adult human small airway epithelial cells and alveolar epithelial cells, respectively, the infection and replication of SARS-CoV-2 in each part of the lung can be evaluated. Recently, it was shown that SARS-CoV-2 infection and replication can be observed in kidney¹², liver ductal¹³, and gut organoids¹⁴. By comparing the infection and replication ability of SARS-CoV-2 in these organoids, the sensitivity of SARS-CoV-2 in each organ could be compared.

The incorporation of mechanical stress into our organoid system could improve the accuracy of SARS-CoV-2 research. The human airway is always exposed to shear stress due to air flow. It has been reported that a functional *in vitro* lung model can be generated using a device capable of medium perfusion and expansion/contraction (organ-on-a-chip)¹⁵. Recently, it was reported that the infection and replication of

SARS-CoV-2 can be observed by culturing primary human lung airway epithelial basal stem cells and pulmonary microvascular endothelial cells on a chip device¹⁶. By applying our hBO to a similar device, we may be able to construct an *in vitro* bronchi model that more closely mimics the living body.

Nevertheless, even in its current condition, we used our hBO to evaluate the efficacy of a COVID-19 therapeutic agent and observe the cytotoxicity and innate immune responses caused by the SARS-CoV-2. Moreover, we could clarify the localization of SARS-CoV-2 in the hBO. These results could be obtained because hBO reproduce the cell populations and functions of the bronchi. Our data of RNA-seq analysis would give us a change to understand the virus affects cellular and bronchial functions.

Finally, we showed that camostat has a positive effect against COVID-19 infection in hBO, demonstrating its usefulness for COVID-19 drug discovery. Similar studies on experimental COVID-19 drugs including those currently undergoing clinical trials should be considered. Furthermore, because hBO can be generated from commercially available cryopreserved hBEpC quickly (10 days) and at large scale, we expect hBO will shorten the search for effective COVID-19 agents.

Materials and Methods

Human bronchial organoid culture

Normal human bronchial epithelial cells (hBEpC, Lonza) were suspended in 10 mg/ml cold Matrigel growth factor reduced (GFR) basement membrane matrix. 50 μ L drops of cell suspension were solidified on pre-warmed Nunc cell-culture treated multidishes (24-well plate) at 37°C for 10 min, and then 500 μ L of expansion medium (composition is shown in **Table S1**) was added to each well. The expansion medium was changed every 2 days. HBO were passaged every 10-12 days. For passaging, the hBO were suspended in 1 mL of 0.5 mM EDTA/PBS (Nacalai tesque) and mechanically sheared using a P1000 pipette tip. Then, 2 mL TrypLE Select (Thermo Fisher Scientific) was added to the suspension. After incubating for 5 min at room temperature, the hBO were again mechanically sheared using a P1000 pipette tip. 7 mL of expansion medium was added, and the organoid suspension tubes were centrifuged at 400 rpm. Organoid fragments were re-suspended in cold expansion medium and seeded as above. HBO were passaged every 10 days. To mature the hBO, the expanded hBO were cultured with differentiation medium (composition is shown in **Table S1**) for 5 days. HBO can be cryopreserved by using STEM-CELLBANKER GMP grade (TaKaRa Bio).

A549 culture

A549 cells were cultured with Ham's F12 medium (Thermo Fisher Scientific) containing 10% FBS, 1 $\times$ GlutaMAX (Thermo Fisher Scientific), and penicillin-streptomycin. A549 cells were passaged every 4 days.

SARS-CoV-2 preparation

The SARS-CoV-2 strain used in this study (SARS-CoV-2/Hu/DP/Kng/19-020) was obtained from the Kanagawa Prefectural Institute of Public Health. SARS-CoV-2 was isolated from a COVID-19 patient in Japan (GenBank: LC528232.1). The isolation and analysis of the virus will be described elsewhere (manuscript in preparation). The virus was plaque-purified and propagated in Vero E6 cells. SARS-CoV-2 was stored at -80°C. All experiments including virus infections were done in the biosafety level facility at Osaka University strictly following regulations.

SARS-CoV-2 infection and drug treatment

Approximately 100 organoids were infected with 5.0×10^4 PFU of SARS-CoV-2 in a 24-well plate containing 500 μ L differentiation medium. One-half of the differentiation medium containing SARS-CoV-2 was replaced with fresh differentiation medium every day. At 5 days after the infection, the hBO and their supernatant were collected. In the drug treatment experiments, the infected hBO were cultured with differentiation medium containing 10 μ M camostat (Sigma-Aldrich) for 5 days.

SARS-CoV-2 virus plaque assays

VeroE6/TMPRSS2 cells (JCRB1819, JCRB Cell Bank)¹⁷ were seeded on 12 well plates (1.7×10^5 cells/well) and incubated for 24 hr. The culture supernatants serially diluted by medium were inoculated and incubated for 2 hr. Culture medium was removed, fresh medium containing 1% methylcellulose (1.5mL) was added, and the culture was further incubated for 3 days. The cells were fixed with 4% Paraformaldehyde Phosphate Buffer Solution (Nacalai Tesque) and plaques were

visualized by using a Crystal violet.

Quantitative PCR

Total RNA was isolated from hBO using ISOGENE II (NIPPON GENE). cDNA was synthesized using 500 ng of total RNA with a Superscript VILO cDNA synthesis kit (Thermo Fisher Scientific). Real-time RT-PCR was performed with the SYBR Green PCR Master Mix (Applied Biosystems) using a StepOnePlus real-time PCR system (Applied Biosystems). The relative quantitation of target mRNA levels was performed by using the $2^{-\Delta\Delta CT}$ method. The values were normalized by those of the housekeeping gene, *glyceraldehyde 3-phosphate dehydrogenase (GAPDH)*. The PCR primer sequences are shown in **Table S2**.

The SARS-CoV-2 primer and probe sets were obtained from Integrated DNA Technologies (IDT, 10006606).

Ultrathin section transmission electron microscopy (TEM)

HBO fixed in phosphate buffered 2% glutaraldehyde, and subsequently post-fixed in 2% osmium tetra-oxide for 2 hr at 4°C. After fixation, they were dehydrated in a graded series of ethanol and embedded in the epoxy resin. Ultrathin sections were cut and then stained with uranyl acetate and lead staining solution and were examined using an electron microscope (HITACHI H-7600) at 100 kV.

Histopathology and immunofluorescence

Fixed bronchial organoid samples were processed and embedded in paraffin. Then they were cut into 2 μ m-thick sections. The sections were deparaffinized,

rehydrated, and stained with hematoxylin and eosin (HE). The sections were then examined using a microscope (BX53 microscope with DP73 camera, Olympus Corporation).

For the immunohistochemical stain assay, the formalin-fixed and paraffin-embedded bronchial organoid samples were treated with pH 6.0 citrate buffer for 30 sec at 125°C in a pressure cooker (Dako Japan) as antigen retrieval. Sections were incubated with each antibody (**Table S3**), followed by Histofine Simple Stain MAX-PO (Nichirei Biosciences). The sections were visualized using Peroxidase Stain DAB Kit (Nacalai Tesque) before counterstaining with Meyer's hematoxylin.

For the double immunofluorescence staining assay, the sections were deparaffinized and subjected to antigen retrieval by treating them with 0.5% trypsin for 30 min. Then the sections were blocked by 5% skim milk with albumin obtained from Bovine Serum Cohn Fraction V, pH 7.0 (Wako Pure Chemical Industries), in PBS for 30 min at room temperature to avoid non-specific reactions. The sections were then incubated with primary antibody (**Table S3**) overnight at 4°C, washed, and incubated with secondary antibody (**Table S3**) for 1 h at room temperature. After washing with PBS, the specimens were mounted with glycerol. All observations were performed using the BX53 fluorescence microscope with a DP73 camera equipped with a suitable filter set (red filter with excitation range 530-550 nm and emission range 575 nm, and green filter with excitation range 470-495 nm and emission range of 510 nm).

RNA-seq

Total RNA was prepared using the RNeasy Mini Kit (Qiagen). RNA integrity was assessed with a 2100 Bioanalyzer (Agilent Technologies). Library preparation was

performed using a NEBNext Ultra II Directional RNA Library Prep Kit for Illumina (NEB) or a TruSeq stranded mRNA sample prep kit (Illumina) according to the manufacturer's instructions. Sequencing was performed on an Illumina NextSeq500 or NovaSeq6000 platform in 152- or 101-base single-end mode, respectively. Fastq files were generated using bcl2fastq2. Adapter sequences were trimmed from the raw reads by cutadapt ver 2.7. The trimmed reads were mapped to the human reference genome sequences (hg19) using HISAT2 ver 2.1.0. The raw counts were calculated using featureCounts ver 2.0.0 and used for heatmap visualization with integrated differential expression and pathway analysis (iDEP, (<http://ge-lab.org/idep/>))¹⁸. Access to raw data concerning this study was submitted under Gene Expression Omnibus (GEO) accession number GSE150819.

LDH assay

After the SARS-CoV-2 infection, the release of LDH was monitored from an aliquot of 250 µL supernatant using the LDH-Glo cytotoxicity assay (Promega) according to the manufacturer's instructions. The absorbance was determined with a Bio-Rad microplate reader (Bio-Rad, US) at wavelength 490 nm. The release of LDH in uninfected cells was used as a control.

Statistical analyses

Statistical analysis was performed using the unpaired two-tailed Student's *t*-test. Statistical significance was evaluated by one-way analysis of variance (ANOVA) followed by Tukey's or Dunnett's post hoc tests to compare all groups.

Acknowledgements

The SARS-CoV-2 strain used in this study (SARS-CoV-2/Hu/DP/Kng/19-020) was obtained from Kanagawa Prefectural Institute of Public Health. This research was supported by the iPS Cell Research Fund. The figure 3A was created using Biorender (<https://biorender.com>). We thank Dr. Tomohiko Takasaki and Dr. Jun-Ichi Sakuragi (Kanagawa Prefectural Institute of Public Health) for providing SARS-CoV-2 strain (SARS-CoV-2/Hu/DP/Kng/19-020), Dr. Misaki Ouchida (Kyoto University) for creating graphical abstract, Dr. Peter Karagiannis (Kyoto University) for critical reading of the manuscript, Dr. Nobihito Morone (University of Cambridge) for critical discussions, Ms. Sayaka Deguchi (Osaka University) for technical assistance with the qPCR analysis, and Ms. Kazusa Okita and Ms. Eri Kawaguchi for technical assistance with the RNA-seq experiments.

Author Contributions

TS performed the SARS-CoV-2 experiments and analyses
 YI performed the SARS-CoV-2 experiments and analyses
 YS performed the immunohistochemical analysis
 AS prepared the materials for the SARS-CoV-2 experiments and analyses
 DO performed the analysis of RNA-seq data of the infected bronchial organoids
 DM collected RNA-seq data of the infected bronchial organoids
 SM performed the SARS-CoV-2 experiments and analyses
 TK performed the SARS-CoV-2 experiments and analyses
 TY collected the RNA-seq data of the bronchial organoids
 TO designed the research and performed the SARS-CoV-2 experiments and analyses

KT designed the research, generated the bronchial organoids, performed statistical analysis, and wrote the paper

Declaration of interests

The authors declare no competing financial interests.

References

- 1 Lu, H., Stratton, C. W. & Tang, Y. W. Outbreak of Pneumonia of Unknown Etiology in Wuhan China: the Mystery and the Miracle. *Journal of Medical Virology*.
- 2 Whitworth, J. COVID-19: a fast evolving pandemic. *Transactions of The Royal Society of Tropical Medicine and Hygiene* (2020).
- 3 Hoffmann, M. *et al.* SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a clinically proven protease inhibitor. *Cell* (2020).
- 4 van der Vaart, J. & Clevers, H. Airway organoids as models of human disease. *Journal of Internal Medicine* (2020).
- 5 Lukassen, S. *et al.* SARS - CoV - 2 receptor ACE2 and TMPRSS2 are primarily expressed in bronchial transient secretory cells. *The EMBO journal* (2020).
- 6 Zhu, N. *et al.* A novel coronavirus from patients with pneumonia in China, 2019. *New England Journal of Medicine* (2020).
- 7 Pizzorno, A. *et al.* Characterization and treatment of SARS-CoV-2 in nasal and bronchial human airway epithelia. *bioRxiv* (2020).
- 8 Han, Y. *et al.* Identification of Candidate COVID-19 Therapeutics using hPSC-derived Lung Organoids. *bioRxiv* (2020).
- 9 Wong, A. P. *et al.* Directed differentiation of human pluripotent stem cells into mature airway epithelia expressing functional CFTR protein. *Nature biotechnology* **30**, 876 (2012).
- 10 Firth, A. L. *et al.* Generation of multiciliated cells in functional airway epithelia from human induced pluripotent stem cells. *Proceedings of the National Academy of Sciences* **111**, E1723-E1730 (2014).
- 11 Sachs, N. *et al.* Long - term expanding human airway organoids for disease modeling. *The EMBO journal* **38** (2019).

438 12 Monteil, V. *et al.* Inhibition of SARS-CoV-2 infections in engineered human tissues
439 using clinical-grade soluble human ACE2. *Cell* (2020).

440 13 Zhao, B. *et al.* Recapitulation of SARS-CoV-2 infection and cholangiocyte damage
441 with human liver ductal organoids. *Protein & Cell*, 1-5 (2020).

442 14 Lamers, M. M. *et al.* SARS-CoV-2 productively infects human gut enterocytes.
443 *Science* (2020).

444 15 Huh, D. *et al.* Reconstituting organ-level lung functions on a chip. *Science* **328**,
445 1662-1668 (2010).

446 16 Si, L. *et al.* Human organs-on-chips as tools for repurposing approved drugs as
447 potential influenza and COVID19 therapeutics in viral pandemics. *bioRxiv* (2020).

448 17 Matsuyama, S. *et al.* Enhanced isolation of SARS-CoV-2 by TMPRSS2-expressing
449 cells. *Proceedings of the National Academy of Sciences* **117**, 7001-7003 (2020).

450 18 Ge, S. X., Son, E. W. & Yao, R. iDEP: an integrated web application for differential
451 expression and pathway analysis of RNA-Seq data. *BMC bioinformatics* **19**, 534
452 (2018).

453

454

Figure 1

Figure 1 Generation of human bronchial organoids

(A) Phase and HE staining images of human bronchial organoids (hBO). Scale bar = 20

μ m. (B) A TEM image of hBO. Larger images are shown in Fig. S2. (C) The gene

expression levels of *ACE2* and *TMPRSS2* in hBO and hBEpC were examined by qPCR.

The gene expression levels in hBO were normalized to 1.0. (D) The expressions of

ACE2 and *TMPRSS2* were examined by immunohistochemistry. Scale bar = 20 μ m.

(E-H) The gene expression levels of basal (E), ciliated (F), goblet (G), and club (H) cell

markers in hBO, hBEpC, and A549 cells were examined by qPCR. The gene expression

levels in hBO were normalized to 1.0. (I) Immunohistochemistry analysis of KRT5

(basal cell marker), acetylated α -tubulin (ciliated cell marker), Mucin 5AC (goblet cell

marker), and CC10 (club cell marker) in hBO. Scale bars = 20 μ m. All data are

represented as means $\pm$ SD ($n = 3$). Statistical significance was evaluated by one-way

ANOVA followed by Tukey's post-hoc tests. Groups that do not share the same letter

are significantly different from each other ($p < 0.05$).

Figure 2

Figure 2 Global gene expression profile of human bronchial organoids

RNA seq analysis was performed in A549 cells (A549), human bronchial organoids (hBO), and human bronchial epithelial cells (hBEpC). (A) Hierarchical clustering analysis of 2,000 variable genes was performed. (B) Principal component analysis (PCA) in “A549”, “hBO”, and “hBEpC”. (C) A scatter plot in “A549”, “hBO”, and

477 “hBEpC”. (D) A clustering analysis of bronchial markers was performed.

478

Figure 3

Figure 3 SARS-CoV-2 infection experiments in human bronchial organoids

(A) hBO were infected with SARS-CoV-2 (5.0×10^4 PFU/well) in the presence or

absence of 10 μ M camostat and then cultured with differentiation medium for 5 days.

(B) At days 1, 2, 3, 4, and 5 after the infection, an LDH assay was performed. (C) The viral genome expression levels in uninfected organoids (control), infected organoids (SARS-CoV-2), and infected organoids treated with camostat (SARS-CoV-2 + camostat) were examined by qPCR. The gene expression levels in control were normalized to 1.0. Statistical significance was evaluated by one-way ANOVA followed by Tukey's post-hoc tests. Groups that do not share the same letter are significantly different from each other ($p < 0.05$). (D) The expression of SARS-CoV-2 Spike protein was examined by immunohistochemistry. Scale bar = 20 μ m. (E) The expression of SARS-CoV-2 Spike protein and KRT5 was confirmed by immunofluorescence staining. Nuclei were counterstained with DAPI. Scale bar = 20 μ m. (F) The amount of infectious virus in the supernatant was measured. Statistical analysis was performed using the unpaired two-tailed Student's *t*-test (* $p < 0.05$). (G) HE staining images of uninfected organoids (control) and infected organoids (SARS-CoV-2) are shown. Blue circles show the existence of pyknotic cells. Scale bar = 20 μ m. All data are represented as means $\pm$ SD ($n = 3$).

Figure 4

Figure 4 Global gene expression profile of infected human bronchial organoids

RNA seq analysis of uninfected hBO (control), SARS-CoV-2-infected hBO

(SARS-CoV-2), and SARS-CoV-2-infected hBO treated with camostat (SARS-CoV-2 +

camostat). (A) A clustering analysis of 2,000 variable genes was performed. (B)

504 Principal component analysis (PCA) in “control”, “SARS-CoV-2”, “SARS-CoV2 +
505 camostat”. (C) A scatter plot in “control”, “SARS-CoV-2”, “SARS-CoV2 + camostat”.
506 (D) A heat map of IFN-I signaling-related genes in “control” and “SARS-CoV-2” is
507 shown. (E) PGSEA (Parametric Gene Set Enrichment Analysis) applied on GO
508 biological process gene sets was performed.
509

Supplemental figures

Figure S1

Figure S1 FGF2 promotes the maturation of human bronchial organoids

Expanded hBO were cultured with differentiation medium containing FGF2, EGF, HGF, R-spondin 1, or Noggin for 5 days. The gene expression levels of *ACE2* and *TMPRSS2* were examined by qPCR. The gene expression levels in “FGF2” were normalized to 1.0. All data are represented as means $\pm$ SD ($n = 3$). Statistical significance was evaluated by one-way ANOVA followed by Dunnett’s post-hoc tests (* $p < 0.05$, ** $p < 0.01$, as compared with control).

Figure S2

human bronchial organoids

goblet cells

basal cells

9 + 2 arrangement

cilia and microvilli

Figure S2 TEM image of human bronchial organoids

The larger TEM images of Fig. 1B are shown. Goblet cells, basal cells, 9+2 arrangement, cilia, and microvilli can be observed in hBO.

Figure S3

ACE2 / KRT5 / DAPI

Figure S3 ACE2 is expressed in basal cells

The expression of ACE2 and KRT5 (basal cell marker) in hBO was confirmed by immunofluorescence staining. Nuclei were counterstained with DAPI. Scale bar = 20 μ m.

Figure S4

Figure S4 SARS-CoV-2 Spike protein is not expressed in club cells

hBO were infected with SARS-CoV-2 (5.0×10^4 PFU/well) and then cultured with differentiation medium for 5 days. The expression of SARS-CoV-2 Spike protein and CC10 was confirmed by immunofluorescence staining. Nuclei were counterstained with DAPI. Scale bar = 20 μ m.

Figure S5

Figure S5 SARS-CoV-2-induced innate immune responses in human bronchial organoids

hBO were infected with SARS-CoV-2 (5.0×10^4 PFU/well) in the presence or absence of 10 μ M camostat and then cultured with differentiation medium for 5 days. (A, B) The gene expression levels of type I interferon (*IFN alpha* and *IFN beta*), interferon-stimulated genes (*ISG56*, *ISG15*, and *MxA*) (A), and SARS-CoV-2-associated genes (*ACE2* and *TMPRSS2*) (B) in uninfected organoids (control), infected organoids (SARS-CoV-2), and infected organoids treated with camostat (SARS-CoV-2 + camostat) were examined by qPCR. The gene expression levels in control were normalized to 1.0. All data are represented as means $\pm$ SD ($n = 3$).

Tables

Table S1 Composition of expansion and differentiation media for human bronchial organoids

Composition (concentration)	Expansion medium	Differentiation medium
Advanced DMEM/F12	+	+
FGF2 (5 ng/ml)	+	+
FGF7 (20 ng/ml)	+	+
FGF10 (100 ng/ml)	+	+
Noggin (100 ng/ml)	+	—
R-spondin 1 (300 ng/ml)	+	—
Y-27632 (10 μ M)	+	+
SB202190 (100 μ M)	+	—
A83-01 (1 μ M)	—	+
B27 supplement (1 $\times$)	+	+
N-Acetylcysteine (1.25 mM)	+	+
Nicotinamide (5 mM)	+	+
GlutaMAX (1 $\times$)	+	+
HEPES (10 mM)	+	+
Penicillin-Streptomycin (100 U/ml)	+	+
Primocin (50 μ g)	+	+

558 **Table S2 Primer list**

target gene	Fwd primer	Rev primer
ACE2	ACAGTCCACACTTGCCCAAAT	TGAGAGCACTGAAGACCCATT
GAPDH	GGTGGTCTCCTCTGACTTCAACA	GTGGTCGTTGAGGGCAATG
IFN alpha	GCAGATCACCCAGAAGATCG	GGCCCTTGTTATTCTCACC
IFN beta	CCTTGCTGAAGTGTGGAGGA	CCAGGCGATAGGCAGAGA
ISG15	GCAGATCACCCAGAAGATCG	GGCCCTTGTTATTCTCACC
ISG56	CCTTGCTGAAGTGTGGAGGA	CCAGGCGATAGGCAGAGA
KRT5	CCAAGGTTGATGCACTGATGG	TGTCAGAGACATGCGTCTGC
MCIDAS	ATTCCCACCAAACGGAAGCAG	CCAGGGTAGGCGACATCATAG
MUC20	ATGACAACGGACGACACAGAA	TCAGCGTTTGAGTTCCAGAG
MUC5B	GCCTACGAGGACTTCAACGTC	CCTTGATGACAACACGGGTGA
MxA	CTTATCCGTTAGCCGTGGTG	CAAGGTGGAGCGATTCTGAG
NGFR	CCTACGGCTACTACCAGGATG	CACACGGTGTCTGCTTGT
PROM1	GGCCCAGTACAACACTACCAA	ATTCCGCCTCCTAGCACTGAA
SCGB1A	TTCAGCGTGTCATCGAAACCC	ACAGTGAGCTTTGGGCTATTTT
TMPRSS2	GTCCCCACTGTCTACGAGGT	CAGACGACGGGGTTGGAAG
TUBA1A	TCGATATTGAGCGTCCAACCT	CAAAGGCACGTTTGGCATACA

559

560 **Table S3 Antibody list**

Antigen (clone number)	catalog number	company
ACE2	PGI-21115-1-AP-150	Proteintech
acetylated α tubulin (6-11B-1)	sc-23950	Santa Cruz Biotechnology
CC10 (E-11)	sc-365992	Santa Cruz Biotechnology
cytokeratin 5 (RCK103)	sc-32721	Santa Cruz Biotechnology
mucin 5AC (45M1)	sc-21701	Santa Cruz Biotechnology
SARS-CoV/SARS-CoV-2 (COVID-19) spike protein (1A9)	GTX632604	GeneTex
TMPRSS2 (H-4)	sc-515727	Santa Cruz Biotechnology

561

562