Muscle fiber type differences in nitrate and nitrite storage and nitric oxide signaling in rats

Gary M. Long^{1,2}, Derrick A. Gray³, Ashley D. Troutman^{2,3}, Amanda Fisher²,
Mary Beth Brown⁴, and Andrew R. Coggan³

¹Department of Kinesiology, Health, and Sports Science,

College of Health Sciences, University of Indianapolis, Indianapolis, IN

Departments of ²Physical Therapy and ³Kinesiology, School of Health and Human Sciences,

Indiana University Purdue University Indianapolis, Indianapolis, IN

⁴Rehabilitation Medicine, University of Washington School of Medicine, Seattle, WA

Running head: Fiber type differences in nitrate/nitrite/nitric oxide metabolism

Declarations of interest: None

Address for correspondence: Andrew R. Coggan, Ph.D., FACSM
Department of Kinesiology
Indiana University Purdue University Indianapolis
IF 101C, 250 University Boulevard
Indianapolis, IN 46202
Tel.: (317) 274-0656
Fax: (317) 278-2041
Email: acoggan@iupui.edu

Abstract

Recent studies have emphasized the importance of the nitric oxide synthase (NOS)-independent, nitrate (NO_3^-) $\rightarrow$ nitrite (NO_2^-) $\rightarrow$ nitric oxide (NO) pathway in skeletal muscle. In particular, it has been hypothesized that this pathway is especially active in type II, or fast-twitch, muscle fibers, necessitating greater NO_3^- and NO_2^- storage. We therefore measured NO_3^- and NO_2^- concentrations in the predominantly fast-twitch vastus lateralis and predominantly slow-twitch soleus muscles of rats. Contrary to the above hypothesis, we found that NO_3^- and NO_2^- concentrations were 3.4-fold and 1.8-fold higher, respectively, in the soleus. On the other hand, NO signaling (i.e., cyclic guanosine monophosphate (cGMP) level) was comparable in the two muscles. Although the physiological significance of these observations remains to be determined, we speculate that NO production via the $\text{NO}_3^- \rightarrow \text{NO}_2^- \rightarrow \text{NO}$ pathway is normally higher in slow-twitch muscles, thus helping compensate for their inherently lower NOS activity.

Keywords

fast-twitch; slow-twitch; cGMP

Introduction

The gaseous free radical nitric oxide (NO) plays an important regulatory role in a wide variety of tissues. These include skeletal muscle, in which NO availability influences blood flow, glucose uptake, mitochondrial respiration, contractile function, etc. (1). These effects are mediated via both cyclic guanosine monophosphate (cGMP)-dependent and cGMP-independent mechanisms, e.g., S-nitrosylation of the ryanodine receptor (1). Originally, NO production in muscle (and other tissues) was believed to be entirely due to the activity of the enzyme NO synthase (NOS), which catalyzes the reaction: $2 \text{ L-arginine} + 3 \text{ NADPH} + 3 \text{ H}^+ + 4 \text{ O}_2 \rightarrow 2 \text{ citrulline} + 2 \text{ NO} + 4 \text{ H}_2\text{O} + 3 \text{ NADP}^+$. The NO thus produced can be eliminated via oxidation to first nitrite (NO_2^-) and then nitrate (NO_3^-), followed by urinary excretion. More recently, however, it has been recognized that NO can also be generated via reversal of this pathway, i.e., via the reduction of NO_3^- to NO_2^- and then to NO via bacterial and/or mammalian nitroreductases (e.g., xanthine oxidoreductase (XOR)) (2,3). Compared to NOS-mediated NO production, this alternative pathway tends to be favored at low pH and low pO_2 , conditions typical of resting and especially contracting muscle (4). In fact, due in part to its large mass skeletal muscle represents the body's major NO_3^- reservoir, which potentially can be used to support NO formation locally or, after export via the circulation, in other tissues (4). In turn, muscle seems to depend upon both endogenous and exogenous sources to maintain its stores of NO_3^- and NO_2^- (4,5).

Based in part on the effects of dietary NO_3^- supplementation on muscle blood flow (6), it has been hypothesized that the $\text{NO}_3^- \rightarrow \text{NO}_2^- \rightarrow \text{NO}$ pathway is especially active in type II, or fast-twitch, skeletal muscle (7). Accordingly, Nyakayiru, van Loon, and Verdijk (8) have recently speculated in this journal that NO_3^- and NO_2^- stores may be greater in fast-twitch muscle fibers. In contrast to this hypothesis, however, as reported herein we have found that NO_3^- and NO_2^- levels are actually several-fold higher in the predominantly type I, or slow-twitch, soleus vs. the predominantly fast-twitch

superficial vastus lateralis of rats. To our knowledge, these are the first fiber type-specific measurements of muscle NO_3^- and NO_2^- concentrations in any species.

Methods

Male Sprague Dawley rats (Charles River Laboratories, Wilmington, MA) weighing ~400g were heparinized, anesthetized (5% isoflurane in 95% O_2), and 300 μL of blood withdrawn via the caudal artery. The sample was immediately centrifuged at 4°C for 10 min at 10,000 g to obtain plasma, which was stored at -80°C until analysis. Rats were then sacrificed via exsanguination and bilateral pneumectomy, after which sections of the soleus and the superficial (“white”) portion of the vastus lateralis were rapidly excised, snap-frozen in liquid N_2 , and also stored at -80°C until analysis. All study procedures were approved by the Institutional Animal Care and Use Committee at Indiana University (protocol number 11151).

Plasma and muscle NO_3^- and NO_2^- concentrations were measured using a high performance liquid chromatography (HPLC) analyzer (ENO-30, Eicom USA, San Diego, CA) as previously described in detail (9,10). Briefly, 25 μL of thawed plasma was mixed 1:1 with methanol, centrifuged at 4°C for 10 min at 10,000 g, and a 10 μL aliquot of the protein-poor supernatant injected into the HPLC. Muscle samples weighing ~10 mg were pulverized at liquid N_2 temperature in a stainless steel tissue pulverizer (Bessman Tissue Pulverizer, Thermo Fischer Scientific), transferred into pre-weighed microcentrifuge tubes, and extracted on dry ice for 30 min using 50 μL methanol containing 0.5% Triton X-100 and 0.1 mmol/L oxypurinol (to block residual XOR activity). Samples were then quickly reweighed to determine the exact amount of tissue added, centrifuged at 4°C for 10 min at 10,000 g, and stored at -80°C until analysis. A 10 μL aliquot of the tissue extract was subsequently injected into the HPLC. NO_3^- and NO_2^- levels in plasma and muscle were calculated from standard curves generated using NIST-traceable standard solutions.

Muscle cGMP content was measured using an ELISA kit (Cayman Chemicals, Ann Arbor, MI). A 25 μ L aliquot of each tissue extract was transferred to the microplate, evaporated to dryness, and resuspended in 500 μ L of assay buffer prior to analysis.

Normality of data distribution was tested using the D'Agostino-Pearson omnibus test. Differences between muscles were determined using unpaired *t* tests for unequal variance. $P < 0.05$ was considered significant. Statistical analyses were performed using GraphPad Prism version 8.4.1 (GraphPad Software, La Jolla, CA).

Results

As shown in Figure 1, muscle NO_3^- and NO_2^- concentrations were significantly higher ($P < 0.01$) in the slow-twitch soleus than in the fast-twitch vastus lateralis. On the other hand, there was no difference in the cGMP content of the two muscles (Fig. 1).

Muscle NO_3^- , NO_2^- , and/or cGMP concentrations were not significantly correlated with each other, in either muscle individually or when data from both muscles were combined (data not shown).

Plasma NO_3^- and NO_2^- concentrations averaged 29 ± 4 and 0.33 ± 0.04 $\mu\text{mol/L}$, respectively.

Discussion

It has been known for over a decade that NO_3^- and NO_2^- levels are higher in various tissues (e.g., liver, heart) than in plasma (2,11-13). This includes both mouse (14) and human (15,16) skeletal muscle. However, interest in the latter tissue has recently increased (e.g., 17), due in part to studies by Piknova and co-workers (4,5,18,19) that have highlighted the central role of muscle in whole-body $\text{NO}_3^-/\text{NO}_2^-/\text{NO}$ metabolism. Skeletal muscle, though, is a heterogeneous tissue, composed of two major fiber types that differ significantly not only in terms of their contractile properties, blood supply, energy metabolism, etc., but also in terms of NOS-mediated NO signaling (see below). In light of recent publications emphasizing the importance of NO production via the alternative $\text{NO}_3^- \rightarrow \text{NO}_2^- \rightarrow \text{NO}$ pathway in muscle, we therefore sought to determine whether there are also fiber type differences in NO_3^- and NO_2^-

storage. To do so, we used a highly-sensitive HPLC-based method (10) to measure NO_3^- and NO_2^- concentrations in the superficial vastus lateralis and soleus muscles of rats. These hindlimb muscles were chosen due to their extremes of fiber type distribution, being composed of approximately 97% fast-twitch and 87% slow-twitch fibers, respectively (20). We found that NO_3^- and NO_2^- concentrations were 3.4- and 1.8-fold higher, respectively, in the predominantly slow-twitch soleus than in the predominantly fast-twitch vastus lateralis. Extrapolating to muscles theoretically composed of 100% of just one fiber type, this implies differences in NO_3^- and NO_2^- content between slow- and fast-twitch fibers of ~4-fold and ~2-fold, respectively. Our data therefore do not support the hypothesis of Nyakayiru, van Loon, and Verdijk (8) that NO_3^- and NO_2^- storage are greater in fast-twitch fibers, at least in rats.

As indicated previously, it is well-established that, at least in rodents, NOS-dependent NO signaling is more prominent in fast-twitch than in slow-twitch muscle. For example, expression of NOS1, or nNOS, is greatest in rat fast-twitch fibers, and minimal or absent in slow-twitch muscle fibers (1). Correspondingly, total NOS activity is higher in muscles composed primarily of fast-twitch fibers (e.g., extensor digitorum longus, gastrocnemius, plantaris) than in the primarily slow-twitch soleus (1). NOS activators or inhibitors (e.g., L-NAME) also generally exhibit far greater effects on, e.g., contractile function in fast-twitch vs. slow-twitch muscle (1). Despite this, it does not necessarily follow that NOS-independent NO signaling is also greater in fast-twitch muscle. Indeed, one could argue that limited or absent NOS activity in rodent slow-twitch fibers would require more dependence upon the $\text{NO}_3^- \rightarrow \text{NO}_2^- \rightarrow \text{NO}$ pathway to augment or enable NO production. If so, this would likely require greater storage of $\text{NO}_3^-/\text{NO}_2^-$ in slow-twitch muscle, especially since such fibers are more readily recruited, and thus more NO_3^- and NO_2^- presumably regularly utilized, during ordinary contractile activity (e.g., stance, ambulation). Furthermore, despite lower microvascular pO_2 in fast-twitch vs. slow-twitch muscle at rest (i.e., ~25 vs. ~30 mmHg) and especially during electrically-stimulated contractions (i.e., ~10 vs. ~25

mmHg), which is hypothesized to enhance deoxyhemoglobin-mediated reduction of NO_2^- to NO in the capillaries (6), the rate of intracellular reduction of NO_2^- to NO may actually normally be greater, or at least equal, in slow-twitch muscle. This is because 1) the bimolecular rate constant for NO_2^- reduction by myoglobin is twice that of hemoglobin, such that in, e.g., the heart most NO_2^- reduction actually occurs in the myocytes, not in the circulation (11), and 2) the total concentration of myoglobin is several-fold higher in slow-twitch muscle (21). Due to the hyperbolic shape and low p_{50} of the oxymyoglobin dissociation curve, the latter difference means that the deoxyhemoglobin concentration will also generally be higher in slow-twitch muscle – only at very low intracellular $p\text{O}_2$, such as may occur during ischemia or intense exercise (22,23), would this not be true. Reduction of NO_2^- to NO (or NO_3^- to NO_2^-) by XOR may also very well be higher in slow-twitch muscle. This is because numerous immunohistochemical studies have demonstrated that XOR expression in muscle tissue is limited to vascular endothelial cells (24-27), and capillarization is much greater in slow-twitch muscle. Consistent with the possibility of significant NO production via the $\text{NO}_3^- \rightarrow \text{NO}_2^- \rightarrow \text{NO}$ pathway in slow-twitch fibers, we found that, at least in the basal state, the concentration of the canonical NO second messenger cGMP is just as high in the soleus as the vastus lateralis, despite limited or absent NOS activity in rodent slow-twitch muscle.

Along with the available literature (e.g., 1,3,11,21,28-33), we therefore interpret our novel observation that NO_3^- and NO_2^- concentrations are higher in the soleus than the superficial vastus lateralis muscle of rats as suggesting that although NOS-dependent NO production and signaling are more prominent in fast-twitch fibers, the NOS-independent, $\text{NO}_3^- \rightarrow \text{NO}_2^- \rightarrow \text{NO}$ pathway may be favored in slow-twitch fibers. This hypothesis is shown in Figure 2, which also illustrates potential differences in the mechanisms by which NO exerts its physiological effects in the two fiber types (i.e., by cGMP-dependent vs. cGMP-independent pathways). Although speculative, we believe that at a minimum this schema can serve as a useful “roadmap” for future research. For example, it would be of

interest to quantify XOR activity in both slow-twitch and fast-twitch muscles, which to our knowledge has not been done. Similarly, the higher NO_3^- and NO_2^- concentrations we found in the soleus implies a higher rate of import of these ions by slow-twitch fibers. It would therefore be of interest to determine whether expression of the NO_3^-/H^+ antiporter sialin, present in both pig (34) and human (35) skeletal muscle, is higher in slow-twitch vs. fast-twitch fibers.

As with any study, there are limitations to the present research. We only measured NO_3^- , NO_2^- , and cGMP levels, and did not quantify any of the other metabolites or proteins shown in Figure 2. We also made measurements only in resting muscle, and did not evaluate the effects of, e.g., increased contractile activity. Thus, although supported by the available literature our hypothesis that NO production via the $\text{NO}_3^- \rightarrow \text{NO}_2^- \rightarrow \text{NO}$ pathway is normally greater in slow-twitch than in fast-twitch muscle is clearly still untested. We also did not investigate the effects of dietary NO_3^- supplementation, which despite the differences in muscle NO_3^- and NO_2^- that we observed following a normal diet could still preferentially “target” fast-twitch muscle (7). Finally, we studied rats, and the present results need to be replicated in humans, especially given marked differences between rats and humans in other aspects of $\text{NO}_3^-/\text{NO}_2^-/\text{NO}$ metabolism, especially fiber type differences in NOS expression (1). The present method is theoretically sensitive enough to quantify the NO_3^- concentration of individual freeze-dried human muscle fibers from biopsy samples, but it would require pooling ~100 fibers of a given type to also measure NO_2^- concentration.

In summary, we have found that, contrary to previous suggestions, NO_3^- and NO_2^- concentrations are markedly higher in the predominantly slow-twitch soleus vs. the predominantly fast-twitch superficial vastus lateralis of rats. These are the first fiber type-specific measurements of these NO precursors in any species. The physiological significance of this difference remains to be determined.

References

1. Stamler JS, Meissner G. Physiology of nitric oxide in skeletal muscle. *Physiol Rev* 2001; 61:209-237.
2. Bryan NS, Fernandez BO, Bauer SM, Garcia-Saura MF, Milsom AB, Rassaf T, Maloney RE, Bharti A, Rodriguez J, Feelisch M. Nitrite is a signaling molecule and regulator of gene expression in mammalian tissues. *Nature Chemical Biology* 2005; 1:290-297.
3. Amdahl MB, DeMartino AW, Gladwin MT. Inorganic nitrite bioactivation and role in physiological signaling and therapeutics. *Biol Chem* 2019; 401:201-211.
4. Piknova B, Park JW, Swanson KM, Dey S, Noguchi CT, Schechter AN. Skeletal muscle as an endogenous nitrate reservoir. *Nitric Oxide* 2015; 47:10-16.
5. Piknova B, Won Park J, Lam KK, Schechter AN. Nitrate as a source of nitrite and nitric oxide during exercise hyperemia in rat skeletal muscle. *Nitric Oxide* 2016; 55-56:54-61.
6. Ferguson SK, Holdsworth CT, Colburn TD, Wright JL, Craig JC, Fees A, Jones AM, Allen JD, Musch TI, Poole DC. Dietary nitrate supplementation: impact on skeletal muscle vascular control in exercising rats with chronic heart failure. *J Appl Physiol* 2016; 121:661-669.
7. Jones AM, Ferguson SK, Bailey SJ, Vanhatalo A, Poole DC. Fiber type-specific effects of dietary nitrate. *Exerc Sport Sci Rev* 2016; 44:53-60.

8. Nyakayiru J, van Loon LJC, Verdijk LB. Could intramuscular storage of dietary nitrate contribute to its ergogenic effect? A mini-review. *Free Radic Biol Med* 2020 Mar 26:S0891-5849(19)31418-2. doi: 10.1016/j.freeradbiomed.2020.03.025. Online ahead of print.
9. Coggan AR, Broadstreet SR, Mahmood K, Mikhalkova D, Madigan M, Bole I, Leibowitz JL, Thomas DP, Thies D, Peterson LR. Dietary nitrate increases VO_2 peak and performance but does not alter ventilation or efficiency in patients with heart failure with reduced ejection fraction. *J Card Fail* 2018; 24:65-73.
10. Troutman AD, Gallardo EJ, Brown MB, Coggan AR. Measurement of nitrate and nitrite in biopsy-sized muscle samples using high performance liquid chromatography. *J Appl Physiol* 2018; 128:1475-1481.
11. Li H, Cui H, Kundu TK, Alzawahra W, Zweier JL. Nitric oxide production from nitrite occurs primarily in tissues not in the blood: critical role of xanthine oxidase and aldehyde oxidase. *J Biol Chem* 2008; 283:17856-17863.
12. Raat NJ, Naguchi AC, Liu VB, Raghavachari N, Liu D, Xu X, Shiva S, Munson PJ, Gladwin MT. Dietary nitrate and nitrite modulate blood and organ nitrite and the cellular ischemic stress response. *Free Rad Med Biol* 2009; 47:510-517.
13. Zweier JL, Li H, Samouilov A, Liu X. Mechanisms of nitrite reduction to nitric oxide in the heart and vessel wall. *Nitric Oxide* 2010; 22:83-90.

14. Ohtake K, Nakano G, Ehara N, Sonoda L, Ito J, Uchida H, Kobayashi J. Dietary nitrite supplementation improves insulin resistance in type 2 diabetic KKAY mice. *Nitric Oxide* 2015; 44:31-38.

15. Montes de Oca M, Torres SH, De Sanctis J, Mata A, Hernández N, Tálamo C. Skeletal muscle inflammation and nitric oxide in patients with COPD. *Eur Respir J* 2005; 26:390-397.

16. Montes de Oca M, Loed E, Torres SH, De Sanctis J, Hernández N, Tálamo C. Peripheral muscle alterations in non-COPD smokers. *Chest* 2008; 133:13-18.

17. Nyakayiru J, Kouw IWK, Cermak NM, Senden JM, van Loon LJC, Verdijk LB. Sodium nitrate ingestion increases skeletal muscle nitrate content in humans. *J Appl Physiol* 2017; 123:637-644.

18. Gilliard CN, Lam JK, Cassel KS, Park JW, Schechter AN, Piknova B. Effect of dietary nitrate levels on nitrate fluxes in rat skeletal muscle and liver. *Nitric Oxide* 2018; 75:1-7.

19. Wylie LJ, Park JW, Vanhatalo A, Kadach S, Black MI, Stoyanov Z, Schechter AN, Jones AM, Piknova B. Human skeletal muscle nitrate store: influence of dietary nitrate supplementation and exercise. *J Physiol* 2019; 597:5565-5576.

20. Armstrong RB, Phelps RO. Muscle fiber type composition of the rat hindlimb. *Am J Anat* 1984; 171:259-257.

21. Okumura N, Hashida-Okamura A, Kita K, Matsubae M, Matsubara T, Takao T, Nagai K. Proteomic analysis of slow- and fast-twitch skeletal muscle. *Proteomics* 2005; 5:2896-2906.

22. Honig CR, Connett RJ, Gayeski TE. O₂ transport and its interaction with metabolism; a systems view of aerobic capacity. *Med Sci Sports Exerc* 1992; 24:47-53.

23. Richardson RS, Noyszewski EA, Kendrick KF, Leigh JS, Wagner PD. Myoglobin O₂ desaturation during exercise. Evidence of limited O₂ transport. *J Clin Invest* 1995; 96:1916-1926.

24. Jarasch E-D, Bruder G, Heid HW. Significance of xanthine oxidase in capillary endothelial cells. *Acta Physiol. Scand Suppl* 1986; 548:39-46.

25. Hellsten-Westing Y. Immunohistochemical localization of xanthine oxidase in human cardiac and skeletal muscle. *Histochemistry* 1993; 100:215-222.

26. Moriwaki Y, Yamamoto T, Yamaguchi K, Takahashi S, Higashino K. Immunohistochemical localization of aldehyde and xanthine oxidase in rat tissues using polyclonal antibodies. *Histochem Cell Biol* 1996; 105:71-79.

27. Linder N, Rapola J, Raivio KO. Cellular expression of xanthine oxidoreductase protein in normal human tissues. *Lab Invest* 1999; 79:967-974.

28. McLane JA, Held IR. Effect of denervation on cyclic nucleotide metabolism in different types of skeletal muscle of the rat. *J Neurosci Res* 1981; 6:327-336.

29. Lau KS, Grange RW, Isotani E, Sarelius IH, Kamm KE, Huang PL, Stull JT. nNOS and eNOS modulate cGMP formation and vascular response in contracting fast-twitch skeletal muscle. *Physiol Genomics* 2000; 2:21-27.

30. Bloom TJ. Cyclic nucleotide phosphodiesterase isozymes expressed in mouse skeletal muscle. *Can J Physiol Pharmacol* 2002; 80:1132-1135.

31. Moon Y, Cao Y, Zhu J, Xu Y, Balkan W, Buys ES, Diaz F, Kerrick WG, Hare JM, Percival JM. GSNOR deficiency enhances in situ skeletal muscle strength, fatigue resistance, and RyR1 S-nitrosylation without impacting mitochondrial content and activity. *Antioxid Redox Signal* 2017; 26:165-181.

32. Moon Y, Balke JE, Madorma D, Siegel MP, Knowels G, Broukaert P, Buys ES, Marcinek DJ, Percival JM. Nitric oxide regulates skeletal muscle fatigue, fiber type, microtubule organization, and mitochondrial ATP synthesis efficiency through cGMP-dependent mechanisms. *Antioxid Redox Signal* 2017; 26:966-985.

33. Thomassen M, Hostrup M, Murphy RM, Cromer BA, Skovgaard C, Gunnarsson TP, Christensen PM, Bangsbo J. Abundance of ClC-1 chloride channel in human skeletal muscle: fiber type specific differences and effect of training. *J Appl Physiol* 2018; 125:470-478.

34. Qin L, Liu X, Sun Q, Fan Z, Xia D, Ding G, Ong HL, Adams D, Gahl WA, Zheng C, Qi S, Jin L, Zhang C, Gu L, Deng D, Ambudkar IS, Wang S. Sialin (SLC17A5) functions as a nitrate transporter in the plasma membrane. *Proc Natl Acad Sci* 2012; 109:13434-13439.

35. Srihirun S, Park JW, Teng R, Sawaengdee W, Piknova B, Schecter AN. Nitrate uptake and metabolism in human skeletal muscle cell cultures. Nitric Oxide 2020; 94:1-8.

Acknowledgements

This publication was made possible by award numbers HL121661 from the National Heart, Lung, and Blood Institute (NHLBI) and AG053606 from the National Institute on Aging (NIA) of the National Institutes of Health (NIH). Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the NHLBI, NIA, or NIH.

Figure legends

Figure 1. Nitrate (NO_3^-), nitrite (NO_2^-), and cyclic guanosine monophosphate (cGMP) concentrations in the superficial vastus lateralis and soleus muscles of male Sprague Dawley rats. *Soleus significantly higher than vastus lateralis ($P < 0.01$). Values are mean $\pm$ S.E. for $n = 9-10/\text{group}$.

Figure 2. Proposed schema of NO signaling in slow-twitch (type I) and fast-twitch (type II) muscle fibers, based on the present results along with the published literature (Refs. 1,3,11,21,28-33). Abbreviations: ClC-1, chloride channel 1. cGMP, cyclic guanosine monophosphate. GTP, guanosine triphosphate. GSNOR, S-nitrosogluthione reductase. HMb, deoxymyoglobin. NOS, nitric oxide synthase. PDE2, phosphodiesterase 2. RSH, thiol groups. RSNO, S-nitrosothiols. sGC, soluble guanyl cyclase. XOR, xanthine oxidoreductase. See Discussion for additional details.