Paleoclimate and current climate collectively shape the phylogenetic and functional diversity of trees worldwide

Wen-Yong Guo^{1,2,*}, Josep M. Serra-Diaz³, Franziska Schrod⁴, Wolf L. Eiserhardt², Brian S. Maitner⁵, Cory Merow⁶, Cyrille Violle⁷, Anne Blach-Overgaard^{1,2}, Jian Zhang⁸, Madhur Anand⁹, Michaël Belluau¹⁰, Hans Henrik Bruun¹¹, Chaeho Byun¹², Jane A. Catford¹³, Bruno E. L. Cerabolini¹⁴, Eduardo Chacón-Madrigal¹⁵, Daniela Ciccarelli¹⁶, Johannes H. C. Cornelissen¹⁷, Anh Tuan Dang-Le¹⁸, Angel de Frutos¹⁹, Arildo S. Dias²⁰, Aelton B. Giroldo²¹, Alvaro G. Gutiérrez²², Wesley Hattingh²³, Tianhua He^{24,25}, Peter Hietz²⁶, Nate Hough-Snee²⁷, Steven Jansen²⁸, Jens Kattge^{19,29}, Tamir Klein³⁰, Benjamin Komac³¹, Nathan Kraft³², Koen Kramer³³, Sandra Lavorel³⁴, Christopher H. Lusk³⁵, Adam R. Martin³⁶, Maurizio Mencuccini^{37,38}, Sean T. Michaletz³⁹, Vanessa Minden^{40,41}, Akira S. Mori⁴², Ülo Niinemets⁴³, Yusuke Onoda⁴⁴, Renske E. Onstein¹⁹, Josep Peñuelas^{38,45}, Valério D. Pillar⁴⁶, Jan Pisek⁴⁷, Matthew J. Pound⁴⁸, Bjorn J.M. Robroek⁴⁹, Brandon Schamp⁵⁰, Martijn Slot⁵¹, Ênio Sosinski⁵², Nadejda A. Soudzilovskaia⁵³, Nelson Thiffault⁵⁴, Peter van Bodegom⁵³, Fons van der Plas⁵⁵, Ian J. Wright⁵⁶, Jingming Zheng⁵⁷, Brian J. Enquist^{5,58}, Jens-Christian Svenning^{1,2}

Affiliations

¹Center for Biodiversity Dynamics in a Changing World (BIOCHANGE), Department of Biology, Aarhus University, 8000 Aarhus C, Denmark; wenyong.guo@bios.au.dk; guowuyhy@gmail.com (WYG); anne.overgaard@bios.au.dk (ABO); svenning@bios.au.dk (JCS)

²Section for Ecoinformatics & Biodiversity, Department of Biology, Aarhus University, 8000 Aarhus C, Denmark ; wolf.eiserhardt@bios.au.dk (WLE)

³Université de Lorraine, AgroParisTech, INRAE, Silva, Nancy, France; pep.serradiaz@agroparistech.fr

⁴School of Geography, University of Nottingham, Nottingham, NG7 2RD, United Kingdom; f.i.schrodt@gmail.com

⁵Department of Ecology and Evolutionary Biology, University of Arizona, Tucson AZ 85721, USA; bmaitner@gmail.com (BSM); brianjenquist@gmail.com (BJE)

⁶Ecology and Evolutionary Biology, University of Connecticut, Storrs, CT, USA; cory.merow@gmail.com

- 28 ⁷UMR 5175 CEF, Centre d'Ecologie Fonctionnelle et Evolutive, Univ Montpellier, CNRS, EPHE, IRD, Univ Paul
29 Valéry Montpellier 3, Montpellier, France; Cyrille.VIOLLE@cefe.cnrs.fr
- 30 ⁸Zhejiang Tiantong Forest Ecosystem National Observation and Research Station, School of Ecological and
31 Environmental Sciences, East China Normal University, Shanghai, 200241, P.R. China;
32 jzhang@des.ecnu.edu.cn
- 33 ⁹School of Environmental Sciences, University of Guelph, Guelph, Ontario, Canada; anand.madhur@gmail.com
- 34 ¹⁰Centre for Forest Research, Département des sciences biologiques, Université du Québec à Montréal, PO Box 8888,
35 Centre-ville station, Montréal (Qc) H3C 3P8, Canada; michael.belluau@usherbrooke.ca
- 36 ¹¹Department of Biology, University of Copenhagen, 2100 Copenhagen Ø, Denmark; HHBruun@bio.ku.dk
- 37 ¹²Department of Biological Sciences and Biotechnology, Andong National University, Andong, 36729, Korea;
38 chaehobyun@anu.ac.kr
- 39 ¹³Department of Geography, King's College London, London, WC2B 4BG, UK; jane.catford@kcl.ac.uk
- 40 ¹⁴Department of Biotechnology and Life Sciences, University of Insubria, Varese, Italy; bruno.cerabolini@uninsubria.it
- 41 ¹⁵Escuela de Biología, Universidad de Costa Rica; 11501-2060 San Jose, Costa Rica; edchacon@gmail.com
- 42 ¹⁶Department of Biology, University of Pisa, Via Luca Ghini 13, 56126 Pisa, Italy; daniela.ciccarelli@unipi.it
- 43 ¹⁷Department of Ecological Science, Faculty of Science, Vrije Universiteit, 1081 HV Amsterdam, The Netherlands;
44 j.h.c.cornelissen@vu.nl
- 45 ¹⁸University of Science, Ho Chi Minh City, Vietnam; Vietnam National University, Ho Chi Minh City, Vietnam;
46 dlatuan@gmail.com
- 47 ¹⁹German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Deutscher Platz 5e, 04103 Leipzig,
48 Germany; adefrutost@gmail.com (AdF); jkatge@bgc-jena.mpg.de (JK); onsteinre@gmail.com (REO);
- 49 ²⁰Goethe University, Institute for Physical Geography, Altenhöferallee 1, 60438 Frankfurt am Main, Germany;
50 arildodias@gmail.com
- 51 ²¹Departamento de Ensino, Instituto Federal de Educação, Ciências e Tecnologia do Ceará - IFCE campus Crateús,
52 Avenida Geraldo Barbosa Marques, 567, Crateús, Brazil. 63708-260; aeltonbg@gmail.com
- 53 ²²Departamento de Ciencias Ambientales y Recursos Naturales Renovables, Facultad de Ciencias Agronómicas,
54 Universidad de Chile, Santa Rosa 11315, La Pintana, Santiago, Chile; bosqueciencia@gmail.com
- 55 ²³Global Systems and Analytics, Nova Pioneer, Paulshof, Gauteng, South Africa; wesnhattigh@gmail.com
- 56 ²⁴School of Molecular and Life Sciences, Curtin University, PO Box U1987, Perth, WA 6845, Australia;
57 Tianhua.He@curtin.edu.au
- 58 ²⁵College of Science, Health, Engineering and Education, Murdoch University, Murdoch, WA, Australia;
- 59 ²⁶Institute of Botany, University of Natural Resources and Life Sciences Vienna, Austria; peter.hietz@boku.ac.at
- 60 ²⁷Four Peaks Environmental Science and Data Solutions, Wenatchee, WA 98801 USA;
61 nhoughsnee@fourpeaksenv.com
- 62 ²⁸Institute of Systematic Botany and Ecology, Ulm University, 89081 Ulm, Germany; steven.jansen@uni-ulm.de
- 63 ²⁹Max Planck Institute for Biogeochemistry, Hans Knöll Str. 10, 07745 Jena, Germany;
- 64 ³⁰Department of Plant & Environmental Sciences, Weizmann Institute of Science, 76100 Rehovot, Israel;
65 tamir.klein@weizmann.ac.il

- 66 ³¹Centre d'Estudis de la Neu i la Muntanya d'Andorra, Institut d'Estudis Andorrans (CENMA - IEA), Avinguda
67 Rocafort 21–23, AD600 Sant Julià de Lòria, Principality of Andorra; bkomac.cenma@iea.ad
- 68 ³²Department of Ecology and Evolutionary Biology, University of California, Los Angeles; Los Angeles, CA 90095,
69 USA; nkraft@ucla.edu
- 70 ³³Wageningen University, Forest Ecology and Management group, Droevendaalsesteeg 4 6700 AA Wageningen; and
71 Land Life Company, Mauritskade 63, 1092AD Amsterdam; koen.kramer@wur.nl
- 72 ³⁴Laboratoire d'Ecologie Alpine, LECA, UMR UGA-USMB-CNRS 5553, Université Grenoble Alpes, CS 40700,
73 38058 Grenoble Cedex 9, France; sandra.lavorel@univ-grenoble-alpes.fr
- 74 ³⁵Environmental Research Institute, University of Waikato, Hamilton, New Zealand; clusk@waikato.ac.nz
- 75 ³⁶Department of Physical and Environmental Sciences, University of Toronto Scarborough. 1265 Military Trail, M1C
76 1A4 Toronto, Ontario, Canada; adam.martin@utoronto.ca
- 77 ³⁷ICREA, Barcelona, Spain; m.mencuccini@creaf.uab.es
- 78 ³⁸CREAF, Cerdanyola del Vallès, Barcelona 08193, Catalonia, Spain; josep.penuelas@uab.cat
- 79 ³⁹Department of Botany and Biodiversity Research Centre, University of British Columbia, Vancouver, BC V6T 1Z4,
80 Canada; sean.michaletz@ubc.ca
- 81 ⁴⁰Department of Biology, Vrije Universiteit Brussel, Pleinlaan 2, 1050 Brussels, Belgium; [vanessa.minden@uni-](mailto:vanessa.minden@uni-oldenburg.de)
82 oldenburg.de
- 83 ⁴¹Institute for Biology and Environmental Sciences, University of Oldenburg, 26129 Oldenburg, Germany;
- 84 ⁴²Graduate School of Environment and Information Sciences, Yokohama National University, 79-7 Tokiwadai,
85 Hodogaya, Yokohama, 240-8501, Japan; akira.s.mori@gmail.com
- 86 ⁴³Estonian University of Life Sciences, Kreutzwaldi 1, 51006 Tartu, Estonia; ylo.niinemets@emu.ee
- 87 ⁴⁴Division of Forest and Biomaterials Science, Graduate School of Agriculture, Kyoto University, Oiwake,
88 Kitashirakawa, Kyoto, 606-8502 Japan; onoda@kais.kyoto-u.ac.jp; yusuke.onoda@gmail.com
- 89 ⁴⁵CSIC, Global Ecology Unit CREAF- CSIC-UAB, Bellaterra, Barcelona 08193, Catalonia, Spain;
- 90 ⁴⁶Department of Ecology, Universidade Federal do Rio Grande do Sul, Porto Alegre, 91501-970, Brazil;
91 vpillar@ufrgs.br
- 92 ⁴⁷Tartu Observatory, University of Tartu, Observatooriumi 1, Tõravere, 61602, Tartumaa, Estonia;
93 janpisek@gmail.com
- 94 ⁴⁸Department of Geography and Environmental Sciences, Northumbria University, Newcastle upon Tyne, NE1 8ST,
95 United Kingdom; matthew.pound@northumbria.ac.uk
- 96 ⁴⁹Aquatic Ecology & Environmental Biology Group, Faculty of Science, Institute for Water and Wetland Research,
97 Radboud University Nijmegen, Netherlands; Bjorn.Robroek@ru.nl
- 98 ⁵⁰Department of Biology, Algoma University, Sault Ste. Marie, Ontario, P6A 2G4, Canada;
99 Brandon.Schamp@algomau.ca
- 100 ⁵¹Smithsonian Tropical Research Institute, Apartado 0843-03092, Balboa, Ancón, Republic of Panama; slotm@si.edu
- 101 ⁵²Embrapa Recursos Genéticos e Biotecnologia, 70770-917, Brasília, DF, Brazil; enio.sosinski@embrapa.br
- 102 ⁵³Institute of Environmental Sciences, Leiden University, 2333 CC Leiden, the Netherlands;
103 n.a.soudzilovskaia@cml.leidenuniv.nl (NS); p.m.van.bodegom@cml.leidenuniv.nl (PMvB)

104 ⁵⁴Natural Resources Canada, Canadian Wood Fibre Centre, 1055 du P.E.P.S., P.O. Box 10380, Stn. Sainte-Foy, Quebec,
105 QC G1V 4C7, Canada; nelson.thiffault@canada.ca

106 ⁵⁵Systematic Botany and Functional Biodiversity, Life science, Leipzig University, Germany;
107 fonsvanderplas@gmail.com

108 ⁵⁶Department of Biological Sciences, Macquarie University, North Ryde, NSW, 2109, Australia; ian.wright@mq.edu.au

109 ⁵⁷Beijing Key Laboratory for Forest Resources and Ecosystem Processes, Beijing Forestry University, Beijing, 100083,
110 P. R. China; zhjmbifu@yahoo.com

111 ⁵⁸The Santa Fe Institute, 1399 Hyde Park Rd, Santa Fe, NM 87501, USA.

112 **ORCID:**

113 Wen-Yong Guo: <https://orcid.org/0000-0002-4737-2042>

114 Josep M. Serra-Diaz: <https://orcid.org/0000-0003-1988-1154>

115 Wolf L. Eiserhardt: <https://orcid.org/0000-0002-8136-5233>

116 Anne Blach-Overgaard: <https://orcid.org/0000-0002-0200-1547>

117 Jian Zhang: <https://orcid.org/0000-0003-0589-6267>

118 Hans Henrik Bruun <https://orcid.org/0000-0003-0674-2577>

119 Jane A. Catford: <https://orcid.org/0000-0003-0582-5960>

120 Bruno E. L. Cerabolini: <https://orcid.org/0000-0002-3793-0733>

121 Eduardo Chacón-Madriral: <https://orcid.org/0000-0002-8328-5456>

122 Anh Tuan Dang-Le: <https://orcid.org/0000-0002-9794-0669>

123 Arildo S. Dias: <https://orcid.org/0000-0002-5495-3435>

124 Alvaro G. Gutiérrez: <https://orcid.org/0000-0001-8928-3198>

125 Wesley Hattingh: <https://orcid.org/0000-0002-3626-5137>

126 Nate Hough-Snee: <https://orcid.org/0000-0003-4581-0931>

127 Steven Jansen: <https://orcid.org/0000-0002-4476-5334>

128 Jens Kattge: <https://orcid.org/0000-0002-1022-8469>

129 Maurizio Mencuccini: <http://orcid.org/0000-0003-0840-1477>

130 Vanessa Minden: <https://orcid.org/0000-0002-4933-5931>

- 131 Akira S Mori: <https://orcid.org/0000-0002-8422-1198>
- 132 Ülo Niinemets: <https://orcid.org/0000-0002-3078-2192>
- 133 Josep Peñuelas: <https://orcid.org/0000-0002-7215-0150>
- 134 Jan Pisek: <https://orcid.org/0000-0003-0396-2072>
- 135 Matthew Pound: <https://orcid.org/0000-0001-8029-9548>
- 136 Bjorn JM Robroek: <https://orcid.org/0000-0002-6714-0652>
- 137 Martijn Slot: <https://orcid.org/0000-0002-5558-1792>
- 138 Nelson Thiffault: <https://orcid.org/0000-0003-2017-6890>
- 139 Ian J. Wright: <https://orcid.org/0000-0001-8338-9143>
- 140 Brian J. Enquist: <https://orcid.org/0000-0002-6124-7096>
- 141 Jens-Christian Svenning: <https://orcid.org/0000-0002-3415-0862>

Abstract

Trees are of vital importance for ecosystem functioning and services at local to global scales, yet we still lack a detailed overview of the global patterns of tree diversity and the underlying drivers, particularly the imprint of paleoclimate. Here, we present the high-resolution (110 km) worldwide mapping of tree species richness, functional and phylogenetic diversities based on ~7 million quality-assessed occurrences for 46,752 tree species (80.5% of the estimated total number of tree species), and subsequent assessments of the influence of paleo-climate legacies on these patterns. All three tree diversity dimensions exhibited the expected latitudinal decline. Contemporary climate emerged as the strongest driver of all diversity patterns, with Pleistocene and deeper-time ($>10^7$ years) paleoclimate as important co-determinants, and, notably, with past cold and drought stress being linked to reduced current diversity. These findings demonstrate that tree diversity is affected by paleoclimate millions of years back in time and highlight the potential for tree diversity losses from future climate change.

Understanding the global distribution of tree diversity and its underlying drivers has been an enduring pursuit of scientists, at least as far back as Alexander von Humboldt. Achieving this aim is becoming ever more urgent due to forest degradation and land use change¹⁻⁴, and also for aiding forest restoration efforts⁵. Although our understanding of the global extent of tree cover has been greatly improved via remote sensing⁶ and large networks of forest tree plots^{7,8}, there are still gaps in our knowledge of the global patterns and drivers of tree diversity. Previous studies have often focused on tree species richness (SR) (e.g.,⁹). However, SR does not directly represent species' evolutionary history and does not provide trait-based insight into their functioning and role in ecosystems. Phylogenetic diversity (PD;^{10,11}) and functional diversity (FD;¹²) have been introduced as promising, more informative biodiversity variables than SR and have been successfully used in a wide variety of ecological applications, including conservation prioritization¹³⁻¹⁵. Indeed, FD is better coupled than SR to ecosystem functioning, e.g., productivity responses to climate change and forest multi-functionality¹⁶⁻¹⁸. In addition, PD and FD are more informative than SR in describing mechanisms of species coexistence and ecosystem functioning^{19,20}, and thus shed light on species extinction and conservation^{13,14,21-23}.

Many studies have emphasized the importance of current climate and edaphic conditions as key determinants of species diversity (e.g.,^{24,25}). However, paleoclimate could leave an influential legacy, e.g. via speciation, extinction or dispersal, on contemporary SR, and on phylogenetic and functional structures of forest ecosystems^{8,26-32}. Earth climate has experienced continuous changes during geological time³³, such as cooling or warming trends and events, and major climatic transitional periods have coincided with global ecosystem shifts³⁴⁻³⁷. Climatically-stable regions tend to have high speciation and low extinction rates, resulting in higher SR, FD, and PD^{38,39}. Contrastingly, wide climate oscillations (like glacial-interglacial cycles) can dramatically truncate species' ranges and the chances of local diversification and adaptation, increasing the likelihood of

extinction and the removal of species with suboptimal traits, thereby decreasing all three facets of diversity^{40–43}. However, rapid climate change may alternatively cause range fragmentation and further allopatric speciation as the result of isolation, potentially increasing net diversification rates²⁷.

Due to the non-equivalency between the facets of diversity^{44–47}, the responses of SR, PD and FD to different climatic conditions may vary. For example, warm and humid climates are hypothesized to increase diversification rates^{48,49}, dispersal and establishment⁵⁰, and decrease extinction^{27,51}, thus increasing SR and PD, but not necessarily FD, as comparable climates more likely predispose species towards similar functional traits^{25,52–55}. Thus, contemporary species diversity patterns can be the result of historical climate legacies and present-day environment, although the relative importance of these factors for FD and PD could be different.

Variable geological climates, i.e., warm and humid, or cold and dry in different paleo-time periods, had remarkably divergent influences on tree diversities. However, previous studies have concentrated mostly on assessing the effect of the cold and dry Last Glacial Maximum (LGM) imprints that occurred ~27 – 19 thousand years ago (kya), but deeper-time perspectives may also be important. For instance, ref.²⁶ found that palm tree diversity in Africa was affected by deep-time climate during the late Pliocene (3.3 – 3.0 million years ago [mya]) and the late Miocene (11.6 – 7.3 mya), respectively. Similarly, ref.²⁷ found that the late Miocene climate influenced global patterns of conifer phylogenetic structure. Recently, ref.³¹ reported opposite effects of LGM and Miocene tree cover on tree phylogenetic endemism. Hence, considering paleoclimate jointly across a range of time frames could be helpful in better understanding the factors shaping tree diversities. However, only a few SR studies have explicitly considered this²⁶, and even fewer in FD and PD research⁴².

Here, we go beyond global mapping of tree species richness⁹ by estimating species composition and, based thereon, functional and phylogenetic diversity. We subsequently analyze the relative roles of past and present climates in shaping global patterns of tree SR, FD, and PD. We first compiled the most updated dataset of tree species including occurrence records, functional traits, and tree phylogeny, covering 46,752 tree species or 80.5% of the species in the GlobalTreeSearch list^{47,56,57}. We subsequently mapped global tree SR, FD, and PD. To understand the potential effects of paleoclimatic change on tree diversities completely, we examined the relative importance of three paleoclimatic states in determining current SR, FD and PD patterns, with consideration of other potential contemporary covariates, such as current climate, elevation, and human activities (Table S1). Specifically, we explored the influence of paleoclimate related to important climate states of the late Cenozoic, the time frame where current species diversity to a large extent have evolved: i) the warm and humid late Miocene, *ca.* 11.63 – 7.25 mya; the mid-Pliocene Warm period, *ca.* 3.264 – 3.025 mya; the cold and dry Pleistocene glaciations (represented by the LGM, ~ 21 kya); and Pleistocene warm interglacials (IG, ~ 787 kya and ~ 130 kya) (Figs. S1 & S2). In doing so, our study addressed three main goals: (1) mapping global contemporary tree SR, FD and PD; (2) assessing the relative importance of present-day environment, Quaternary glacial-interglacial oscillations, and deeper-time effects on today's SR, FD and PD patterns, to help understand the fundamental processes determining accumulation and maintenance of tree diversity; and (3) investigating spatial divergence between FD and PD, and identifying the underlying driving factors.

Results

Global patterns of tree diversities

The global tree SR, FD, and PD distributions show classic latitudinal gradients^{58–60}, with low diversities at high latitudes and the highest diversities in the tropics (grid cell maximum value of

3261 spp. for SR and cumulative branch lengths of 641 and 61,183 Myrs for FD and PD, respectively at 110 km resolution, Fig. S3), particularly in the Neotropical lowlands (Amazonia) (Fig. 1). The latitudinal pattern is stronger in America and Asia-Australasia than in Africa-Europe, due to the interruption of the diversity gradients by deserts in northern Africa, where the diversity indices (SR, FD, and PD) are as low as at latitudes harboring the boreal climate. The similarity of the spatial patterns among the three diversity measures reflects the monotonic relationships observed between them (Fig. S4).

Drivers of global tree diversity

Due to the high associations between SR, FD, and PD, their individual relationships with the tested predictors are mostly consistent (Figs. 2 & Table S2). After controlling for spatial autocorrelation, simultaneous autoregressive models (SARs) explain more than 94% (global models) and 78% (regional models) of the variance (Table S2) in the response variables (SR, PD, and FD). Present-day annual precipitation (AP) and mean annual temperature (MAT) are the overall strongest drivers with positive effects on SR, FD, and PD globally, and for AP also regionally except for two regions where other drivers are stronger (Australasia, Nearctic). The effect of MAT varies in strength and sign among regions, showing both positive and negative effects on diversity (Fig. 2, Table S2). Elevation range and human modification index (HMc) have consistent positive effects on SR, FD and PD globally as well as regionally. Four out of the six paleoclimatic variables show significant relations to all three diversity dimensions (Fig. 2). Globally, the Miocene MAT anomaly (i.e., Miocene MAT minus present MAT), the Miocene AP anomaly, and the LGM AP anomaly have positive relations to all diversity indices, while the LGM MAT anomaly have a weak negative relation to SR ($p < 0.05$, Table S2) and no relation to FD and PD (Fig. 2). Hence, SR, PD and FD consistently increase with increasing high precipitation in the Miocene and LGM relative to the

present, while SR, but not FD or PD, is generally reduced by increasing warm during LGM at a global scale. However, although some of these global relationships are mirrored regionally, not all paleoclimatic predictors are significant nor show consistent relationships across the biogeographic regions, e.g., with LGM AP anomaly showing negative associations in Australasia and Miocene MAT anomaly in Afrotropic for all three indices (Table S2).

Taken all together, precipitation-related effects were stronger and more consistent (among regions) climatic drivers of diversity (SR, FD, PD) than were temperature-related effects, with this true both for current climate (AP) and for paleoclimates (Miocene AP anomaly; IG AP anomaly).

Spatial divergence between functional and phylogenetic diversities and its drivers

FD and PD are tightly and positively related (Fig. S4c). Deviations (FD residuals) from this linear relationship show marked spatial patterning (Fig. 3). Across North America, western and southern Europe, central Africa, eastern Asia, and eastern Australia, FD is generally higher than predicted by PD (i.e., overdispersion), whilst the opposite (i.e., FD deficit) is revealed in western Australia, much of southern and eastern Africa, west of the Andes (Peru), and central parts of northern Eurasia.

The relative importance of the factors explaining variation in FD residuals are different from those explaining their variations (Fig. 4 vs. Fig. 2b & 2c). Overall, current AP is correlated negatively with the FD residuals both globally and regionally, but is only the strongest driver at global scale (Figs. 4 & 5, Table S3). MAT and non-climatic factors show weak or no relations, except for MAT for Indo-Malay and the Neotropics. The effects of the paleoclimate are variable. At global scale, the Miocene AP anomaly and the LGM MAT anomaly are negatively related to the FD residuals, while the LGM AP anomaly is positively related (Figs. 4 & 5). However, these relationships are inconsistent across biogeographic regions (Figs. 4 & 5).

Discussion

Based on an unprecedented tree occurrence database, our study maps strong latitudinal patterns in all three diversity dimensions (SR, PD, and FD) at global scale. The SR-linked global latitudinal patterns of Faith's PD and FD matches previous empirical and modeled studies of tree species richness (e.g.,^{9,61}), tree functional diversity in the New World⁶², and tree phylogenetic diversity at a regional scale³¹. It has been reported that speciation in rainforest environments has taken place at least since the Paleocene (~58 mya)^{63,64}, probably coupled to jointly high temperatures and precipitation^{38,49,51,65}. Moreover, the relatively stable environment, compared to high latitudes, may also resulted in low extinction rates, making the tropics both “cradles (species diversifying)” and “museums (species persistence)” of species diversity^{49,66}. In addition, long speciation history and lower extinction rates in the tropics could result in both higher phylogenetic diversity and functional diversity^{38,48,49,51}, but see³⁵.

Our results provide evidence that paleoclimate complements current climate in shaping tree diversity globally and regionally, and that these effects are not only related to the recent prehistory – such as the Last Glacial period, represented by the LGM 21,000 years ago – but also much deeper time scales. These results extend previous findings for other organism groups notably for species richness and endemics^{39,42} and for trees or plant clades including trees in specific regions and biomes^{26,27,29,67–70}, to trees globally. Importantly, they go beyond species richness to the more ecologically meaningful indices, functional and phylogenetic diversity.

Notably, we found that precipitation effects were stronger and more consistent (across regions) drivers than temperature effects, especially in relation to the wet and warm middle Miocene (11.6 – 7.5 mya), and the dry, cold LGM. The middle Miocene, the warmest and wettest interval in the late

Cenozoic, was a period of forest expansion^{34,71}, due to warming coupled with elevated atmospheric CO₂ (>500 ppm)^{72–74}. This likely promoted high species diversity globally due to a higher diversification rate and lower extinction rate^{38,48,51}. The Myrtaceae family⁷⁵ and the genus *Quercus*⁷⁶ are examples that follow this pattern. As a legacy of forest expansion, the generally warmer and wetter climate in the late Miocene compared to the contemporary climate have a positive associations to tree SR, FD, and PD³⁷. We also see this in our results at the global scale and for most regions with positive effects of both Miocene AP and MAT (Fig. 2). The weak negative association between the LGM MAT anomaly and SR, but not with FD and PD, could indicate that global cold climate in LGM (Fig. S1) caused range retractions or even extinctions of certain species. Likely, the intensity of these processes were not strong enough to significantly decrease the communities' FD and PD, probably due to the high tree diversity accumulated in previous warm and humid periods^{36,75}. Indeed, both tree FD and PD showed the tendency to level-off with SR increase (Fig. S6), a similar pattern reported by⁵⁴, indicating that closely related tree species have more similar traits, i.e., the functional space tightly packed^{30,70}. The LGM precipitation anomaly was positively related to tree SR, PD and FD, likely reflecting widespread forest contractions during the generally dry LGM and tree survival in moist refugia^{77,78}. Furthermore, the diversity of drier forests itself is generally lower due to a limited number of niches and the physiological limits of species drought tolerance⁷⁹. Our results suggest that paleoclimate affects not just forest biodiversity, but also forest ecosystem functioning given the effects found here, which corroborates other studies on FD^{16,17} and PD^{20,27,67}. Notably, a recent study has found that paleoclimatic legacies in tree FD negatively affect stand productivity in Northern Hemisphere temperate forests⁸⁰.

The relationships between paleoclimate and SR, FD, and PD were partially repeated within biogeographic regions, there was also substantial inter-region variation in these relations (Fig. 2). For example, not all of the four significant relationships found globally were retained regionally,

and new relations emerged in some cases. These variable regional relations may reflect differing regional paleoclimatic histories, differences in the overall climatic and geographic setting, as well as methodological effects, e.g., different covariation among explanatory variables. For example, in Australasia, only the LGM AP anomaly showed significant, negative relationships with FD and PD, possibly because the temperature there was rather stably high during the last millions of years, with precipitation being more variable and lower (Fig. S2).

The regions representing FD surplus relative to PD, i.e., where species were found to be more functionally diverse (high FD) than expected from PD, largely coincided with high SR regions (Figs. 3 & 1a), represented by warm and humid climate today. This suggests that communities in warm and humid conditions have accumulated more FD than expected compared to dry or cold regions. This FD surplus could be caused by high competition, high heterogeneous environments, or otherwise diversifying trait evolution^{19,62,81–83}. We found that all precipitation variables were important for explaining the FD deviation from PD, even though their effects differed (Figs. 4 & 5). Surprisingly, high current precipitation tended to correspond to FD deficits, i.e., areas where species were more functionally similar than predicted by PD, both globally and in several biogeographic regions. Even though the observed FD in many wet and warm areas were higher than expected from PD, an explanation for the observed relationship could be that moist tropical forests harbor large numbers of shade-tolerant species, which have evolved along a similar evolutionary path (i.e., stabilizing selection) to adapt to the shady environment, thus showing high levels of ecological equivalence^{83,84}.

Building on recent progress in the harmonization of several databases on tree species distributions, functional traits, phylogenetic relatedness, and global paleoclimate, we have found that the tropics harbor the highest diversity across not only taxonomic, but also functional and phylogenetic

338 dimensions, while high latitudes have lower diversity values for all diversity measures.

339 Nevertheless, there are important and informative deviations between the patterns in FD and PD,

340 including a signature consistent with less ecological filtering in moist, shady tropical forest

341 environments⁸⁴. Importantly, we found evidence that current tree phylogenetic and functional

342 diversities are likely shaped not only by the contemporary environment, but also by past climate as

343 far back as the Miocene (~10 Mya). Notably, we see long-term reductions in FD and PD in relation

344 to past climatic cold or drought stress, likely affecting current forest ecosystem functioning⁸⁰.

345 These findings highlight the importance of climate for tree diversity and forest ecosystems, and that

346 losses from future climate change could have strong and very long-lasting effects.

Methods

Tree species and their range maps

In this study, we used the world tree species list⁵⁶ and species range maps compiled by^{47,57}. Briefly, the world tree species checklist (GlobalTreeSearch, GTS⁵⁶) was used to extract the global tree species list for the current study. Tree species included in the GTS is based on the definition by the IUCN's Global Tree Specialist Group (GTSG), i.e., "a woody plant with usually a single stem growing to a height of at least two meters, or if multi-stemmed, then at least one vertical stem five centimeters in diameter at breast height"⁵⁶. This list was subsequently standardized via the Taxonomic Name Resolution Service (TNRS) online tool⁸⁵ to remove synonyms. The occurrence records of the selected species were collated from five widely used and publicly accessible databases, namely: the Global Biodiversity Information Facility (GBIF; <http://www.gbif.org>), the public domain Botanical Information and Ecological Network v.3 (BIEN; <http://bien.nceas.ucsb.edu/bien/>;^{86,87}), the Latin American Seasonally Dry Tropical Forest Floristic Network (DRYFLOR; <http://www.dryflor.info/>;⁸⁸), the RAINBIO database (<http://rainbio.cesab.org/>;⁸⁹), and the Atlas of Living Australia (ALA; <http://www.ala.org.au/>). The compiled occurrence data was accessed⁵⁷ and the high-quality records were then used to generate range maps based on the alpha hull algorithm via the *Alphahull* package^{90,91} in R (ver. 3.5.1; ⁹²). We further validated the range maps using an external independent dataset⁹. The estimated range maps of the 46,752 tree species were rasterized to 110 km equal-area grid cells (~1 degree at the Equator), a resolution commonly used in global diversity studies (e.g.,⁴⁵), using the *letsR* package⁹³. For detailed information on the range map estimations and external validation, see⁴⁷.

Phylogeny

We constructed a phylogenetic tree for the tree species using the largest seed-plant phylogeny presently available (the ALLMB tree⁹⁴). This dated phylogeny combines a backbone tree⁹⁵, which was built using sequence data from public repositories (GenBank) to reflect deep relationships, with previous knowledge of phylogenetic relationships and species names from the Open Tree of Life (Open Tree of Life synthetic tree release 9.1 and taxonomy version 3, <https://tree.opentreeoflife.org/about/synthesis-release/v9.1>). This phylogeny was matched to our tree species dataset, and any species that were not in our dataset were removed from the tree. Subsequently, some species missing from the phylogeny were manually added, using the same approach as ref.⁹⁴.

Functional trait data

Eight ecologically relevant and commonly used traits⁹⁶ were selected for functional diversity analyses, i.e., leaf nitrogen content, wood density, leaf phosphorus content, leaf dry matter content, plant max height, seed dry mass, specific leaf area, and leaf area. Originally, we compiled 21 functional traits from the TRY (<https://try-db.org/TryWeb/Home.php>; ^{97,98}, TOPIC^{99–105}, and BIEN (<http://bien.nceas.ucsb.edu/bien/>; ^{86,87}) databases. As many of the species' trait were missing, we imputed missing values via an gap-filling algorithm with Bayesian Hierarchical Probabilistic Matrix Factorization (BHPMF,^{106–108}), which is mostly based on both trait-trait correlation matrix and the phylogentic signal of traits (Refer to ref.⁴⁷ for the detailed gap-filling procedure). In this process, all the 21 traits were used to maximally benefit from the correlations among them.

Environmental variables

We compiled 17 environmental variables, including current climate, paleo-climate, human effects, topographic heterogeneity and evolutionary history (Supplementary Table S1). Climate, both present-day and paleoclimate, is generally assumed to be a vital predictor of species distribution and

diversity patterns (e.g.,^{26,27,29,39,109,110}). Due to the data availability of the paleoclimates, we included two bioclimatic predictors commonly used in relevant studies: annual mean temperature (MAT) and annual precipitation (AP). Current climate variables were extracted from WorldClim (v.2, www.worldclim.org) at a resolution of 30 arc-seconds (~1 km at the equator), averaging global climate data from the period 1970 - 2000¹¹¹. We selected six paleo-time periods spanning from *ca.* 11.6 – 7.2 mya to *ca.* 21 kya, representing climatic conditions either warmer, cooler, or similar compared to the present-day climate. Specifically, each bioclimatic layer of the late Miocene climate (11.61 – 7.25 mya³⁷) and mid-Pliocene Warm period (~ 3.264 – 3.025 mya;^{112,113}) were averaged to represent the warmer climate compared to present day (hereafter Miocene). Pliocene Marine Isotope Stage M2, a glacial interval in the late Pliocene (~ 3.3 mya;^{113,114}), was used to represent the Pliocene global cooling period, while the Last Glacial Maximum (LGM, ~ 21 kya) was used to present the more recent global cooling event compared to M2^{113,115}. We further constructed a current climate (hereafter Interglacial, IG) analog using the mean value per bioclimatic layer between the Pleistocene Marine Isotope Stage 19 (MIS 19), the oldest Pleistocene interglacial (~ 787 kya¹¹³), and the Last Interglacial (LIG; ~ 130 kya¹¹⁶). The mid-Pliocene Warm Period, Pliocene M2, Pleistocene MIS19, and the LIG data were extracted from Paleoclim (www.paleoclim.org), at a resolution of 2.5 arc-minutes (~ 4.5 km at the equator)¹¹³, and the LGM data was extracted from the CHELSA database (www.chelsa-climate.org) at a resolution of 30 secs¹¹⁵.

In addition to climate, other factors, such as human activities, topographic heterogeneity, and evolutionary history, can also affect plant distributions^{9,26,117,118}. The Human Modification map (HMc¹¹⁹)¹¹⁹ was used as a proxy of human activities. Compared to the commonly used human footprint index and human influence index maps¹²⁰, HMc has been modelled with the incorporation of 13 most recent global-scale anthropogenic layers (with the median year of 2016) to account for

the spatial extent, intensity, and co-occurrence of human activities, many of which showing high direct or indirect impact on biodiversity¹²¹. HMc was extracted at a resolution of 1 km²¹¹⁹. The elevation range is the absolute difference between the maximum and minimum elevation value within a specific area. We computed the elevation/topographic range within each 110 × 110 km grid cell based on the digital elevation model at 90 m resolution (<http://srtm.csi.cgiar.org/>). Elevation range is a proxy of environmental heterogeneity, which is considered as a universal driver of biological diversity^{122,123}. To analyze the potential effects of evolutionary and biogeographic history, we also included the biogeographic regions as an additional variable. We applied the definition of biogeographic regions from ref.¹²⁴, which defines 12 regions globally using cladistic and phylogenetic analyses of plant species, and plate tectonics. However, due to the varying data size in each of the 12 regions, we combined them into six regions, i.e., Afrotropic, Australasia, Indo-Malay, Nearctic, Neotropic, Palearctic, largely similar to the biogeographic realms proposed by ref.¹²⁵. All predictors were extracted from various databases, which we describe in further detail in the supplement (Supplementary Table S1).

Except for the biogeographic regions and elevation range, mean values for all predictors were extracted at a 110 × 110 km resolution. The variable extractions and averaging were carried out in the *letsR* package. Due to the low reliability and/or missing environmental variables for many islands¹²⁶, we removed insular grid cells from small islands, and 11,950 grid cells with records were kept (Fig. S5).

Phylogenetic and functional diversity

Phylogenetic diversity (PD) was calculated for each 110 × 110 km grid cell as the sum of the branch lengths of all co-occurring species as defined by ref.¹⁰. Among the many existing, somewhat

overlapping matrices of PD, the one we selected is the most widely used due to its easy calculation and interpretation and a more robust basis for conservation^{10,13,14}.

Functional diversity (FD) was calculated in an analogous manner to PD¹²⁷. A Principal Component Analysis (PCA) was applied to the eight traits to eliminate trait redundancy. Values of all traits were log transformed to improve normality and were standardized before analysis. Then a dendrogram based on the first three PCs (explaining 84% of the total variation) was constructed using Gower's distance via the *vegan*¹²⁸ and *fastcluster*¹²⁹ packages. This dendrogram was used to calculate FD as the sum of the total branch lengths connecting a set of species in the 110 × 110 km grid cell. Both PD and FD were calculated using the *letsR* and *picante*¹³⁰ packages.

To investigate the bivariate relationships between FD and PD, an ordinary least squares model was implemented. We further plotted the residuals of model to show any deviation between FD and PD.

Statistical analyses

To test the long-term climate stability hypothesis, we calculated the anomaly for MAT and AP between the four paleo-time periods and the present-day, i.e., past minus present, to represent the amplitude of the climate changes within each time-scale (Fig. S1)^{26,27,29,39,118}. On average, compared to the present, mean annual temperature (MAT) was much higher in the Miocene, slightly higher in the Pliocene M2 period, much lower in the LGM, and similar in the IG (Fig. S2a). During Pliocene M2 and IG, annual precipitation (AP) was similar to the present-day, while the Miocene and LGM had slightly higher or lower precipitation, respectively than the contemporary precipitation (Fig. S2b). The paleo-time periods selected, thus, represent (on average) cold, warm, and similar paleo-climates compared to present-day conditions.

Pearson correlation coefficients showed a low level of correlations between MAT, AP, and their respective anomaly variables (Fig. S6). However, MAT and AP of Pliocene M2 and Pleistocene IG

anomaly showed relatively high correlations (Fig. S7) with or without accounting for the spatial autocorrelation (using the *SpatialPack* package¹³¹). Consequently, we removed the two Pliocene M2 variables from further analyses.

We used ordinary least squares models (OLSs) and simultaneous autoregressive models (SARs), if the OLS model residuals exhibited spatial autocorrelation (SAC), to evaluate the relative importance of the predictor variables in determining the variation in each of the three diversity indices and the residuals of bivariate relationships between FD and PD. We used the SAR error model because of its superior performance compared to other SAR model types¹³². The SAR error model adds a spatial weights matrix to an OLS regression to accounts for SAC in the model residuals. A series of spatial weights, i.e., k -means neighbor of each site, were tested and $k = 1.5$ was used for all SARs models as it can successfully account for the SAC (see Supplementary results of statistical analyses). Residual SAC was examined in all models (both OLS and SAR) using Moran's I test, and Moran's I correlograms were also used to visualize the spatial residuals of the models. Model explanatory power was represented by adjusted R^2 (OLSs) and Nagelkerke pseudo- R^2 (SARs)¹³³, while the Akaike Information Criterion (AIC) and Bayesian information criterion (BIC) were used to compare the models for each diversity index¹³⁴. SARs and Moran's I tests were carried out using the *spdep* package¹³⁵. Both OLS and SAR models were run by including current MAT and AP, the six anomaly variables, and the other non-climate predictors (elevation range and HMc) to investigate their relative contributions to each diversity index. In addition to the global models, we ran the same models for each biogeographic region to test whether the global relationships varied among regions. Moreover, we ran three additional global models for the FD and PD indices, selecting only one paleoclimate (both MAT and AP) from the three paleo-time periods at the time, and keeping other variables the same in each model to investigate whether the effects of the different paleoclimate predictors changed compared to the full models (including all paleo

climatic predictors). Before running the models, we inspected the normality of all predictors and log₁₀-transformed variables if needed. All response variables (three diversity indices) were log₁₀-transformed. Thereafter, we standardized all predictor variables by transforming all variables to a mean of zero and a standard deviation of one to derive more comparable estimates¹³⁶.

Supplementary results of statistical analyses

We found that for all models (both global and regional), SAR models performed better than the corresponding OLS models, regarding to AIC, BIC, and R^2 (Tables S2-S3), and all SAR models successfully accounted for SAC in model residuals ($p \gg 0.05$, Figs. S8-S11). Thus, we only represented the results from SARs models in the text, even though the significance of some predictors varied between OLS and SAR models (Fig. S12). In addition, we found that the effects of paleoclimate variables showed no change between the full models, including all paleoclimate variables and models using paleoclimate of each paleo-period (Fig. S13-S14). This clearly shows the robustness of their relationships with the tree diversity indices.

References

1. Chazdon, R. L. Beyond deforestation: Restoring forests and ecosystem services on degraded lands. *Science* **320**, 1458–1460 (2008).
2. Potapov, P. *et al.* The last frontiers of wilderness: Tracking loss of intact forest landscapes from 2000 to 2013. *Sci. Adv.* **3**, e1600821 (2017).
3. Watson, J. E. M. *et al.* The exceptional value of intact forest ecosystems. *Nature Ecology and Evolution* **2**, 599–610 (2018).
4. Hansen, M. C., Stehman, S. V & Potapov, P. V. Quantification of global gross forest cover loss. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 8650–8655 (2010).
5. Bastin, J. F. *et al.* The global tree restoration potential. *Science* (80-.). **364**, 76–79 (2019).
6. Hansen, M. C. *et al.* High-resolution global maps of 21st-century forest cover change. *Science* (80-.). **342**, 850–853 (2013).
7. Crowther, T. W. *et al.* Mapping tree density at a global scale. *Nature* **525**, 201–205 (2015).
8. Bruehlheide, H. *et al.* Global trait–environment relationships of plant communities. *Nat. Ecol. Evol.* **2**, 1906–1917 (2018).
9. Keil, P. & Chase, J. M. Global patterns and drivers of tree diversity integrated across a continuum of spatial grains. *Nat. Ecol. Evol.* **3**, 390–399 (2019).
10. Faith, D. P. Conservation evaluation and phylogenetic diversity. *Biol. Conserv.* **61**, 1–10 (1992).
11. Faith, D. P. The PD Phylogenetic Diversity Framework: Linking Evolutionary History to Feature Diversity for Biodiversity Conservation. in 39–56 (2016). doi:10.1007/978-3-319-22461-9_3
12. Petchey, O. L. & Gaston, K. J. Functional diversity: Back to basics and looking forward. *Ecology Letters* **9**, 741–758 (2006).
13. Mazel, F. *et al.* Prioritizing phylogenetic diversity captures functional diversity unreliably. *Nat. Commun.* **9**, 2888 (2018).
14. Tucker, C. M. *et al.* A guide to phylogenetic metrics for conservation, community ecology and macroecology. *Biol. Rev.* **92**, 698–715 (2017).
15. Jarzyna, M. A. & Jetz, W. Detecting the Multiple Facets of Biodiversity. *Trends in Ecology and Evolution* **31**, 527–538 (2016).
16. Huang, X., Su, J., Li, S., Liu, W. & Lang, X. Functional diversity drives ecosystem multifunctionality in a *Pinus yunnanensis* natural secondary forest. *Sci. Rep.* **9**, 1–8 (2019).
17. Díaz, S. *et al.* Functional Diversity — at the Crossroads between Ecosystem Functioning and Environmental Filters. in *Terrestrial Ecosystems in a Changing World* 81–91 (Springer, Berlin, Heidelberg, 2007). doi:10.1007/978-3-540-32730-1_7
18. Díaz, S. & Cabido, M. Vive la différence: Plant functional diversity matters to ecosystem processes. *Trends in Ecology and Evolution* **16**, 646–655 (2001).
19. Safi, K. *et al.* Understanding global patterns of mammalian functional and phylogenetic diversity. *Philos. Trans. R. Soc. B Biol. Sci.* **366**, 2536–2544 (2011).

- 535 20. Donoghue, M. J. A phylogenetic perspective on the distribution of plant diversity. *Proc. Natl. Acad. Sci. U. S. A.* **105**, 11549–11555 (2008).
- 536
- 537 21. Oliveira, B. F., Sheffers, B. R. & Costa, G. C. Decoupled erosion of amphibians' phylogenetic and
- 538 functional diversity due to extinction. *Glob. Ecol. Biogeogr.* **29**, 309–319 (2020).
- 539 22. Faith, D. P. Phylogenetic diversity, functional trait diversity and extinction: Avoiding tipping points
- 540 and worst-case losses. *Philos. Trans. R. Soc. B Biol. Sci.* **370**, 1–10 (2015).
- 541 23. Park, D. S., Feng, X. & Maitner, B. S. Darwin's naturalization conundrum can be explained by spatial
- 542 scale. *Proceedings of the National Academy of Sciences* 201918100 (2020).
- 543 doi:10.5281/zenodo.3710499
- 544 24. Whittaker, R. J. & Field, R. Tree Species Richness Modelling: An Approach of Global Applicability?
- 545 *Oikos* **89**, 399–402 (2000).
- 546 25. Wright, I. J. *et al.* Modulation of leaf economic traits and trait relationships by climate. *Glob. Ecol.*
- 547 *Biogeogr.* **14**, 411–421 (2005).
- 548 26. Blach-Overgaard, A., Kissling, W. D., Dransfield, J., Balslev, H. & Svenning, J. C. Multimillion-year
- 549 climatic effects on palm species diversity in Africa. *Ecology* **94**, 2426–2435 (2013).
- 550 27. Eiserhardt, W. L., Borchsenius, F., Sandel, B., Kissling, W. D. & Svenning, J. C. Late Cenozoic climate
- 551 and the phylogenetic structure of regional conifer floras world-wide. *Glob. Ecol. Biogeogr.* **24**, 1136–
- 552 1148 (2015).
- 553 28. Ordonez, A. & Svenning, J.-C. Consistent role of Quaternary climate change in shaping current plant
- 554 functional diversity patterns across European plant orders. *Sci. Rep.* **7**, 42988 (2017).
- 555 29. Svenning, J. C. & Skov, F. Could the tree diversity pattern in Europe be generated by postglacial
- 556 dispersal limitation? *Ecology Letters* **10**, 453–460 (2007).
- 557 30. Ordonez, A. & Svenning, J. C. Geographic patterns in functional diversity deficits are linked to glacial-
- 558 interglacial climate stability and accessibility. *Glob. Ecol. Biogeogr.* **24**, 826–837 (2015).
- 559 31. Sandel, B. *et al.* Current climate, isolation and history drive global patterns of tree phylogenetic
- 560 endemism. *Glob. Ecol. Biogeogr.* **29**, 4–15 (2020).
- 561 32. Blonder, B. *et al.* Late Quaternary climate legacies in contemporary plant functional composition.
- 562 *Glob. Chang. Biol.* **24**, 4827–4840 (2018).
- 563 33. Zachos, J., Pagani, H., Sloan, L., Thomas, E. & Billups, K. Trends, rhythms, and aberrations in global
- 564 climate 65 Ma to present. *Science* **292**, 686–693 (2001).
- 565 34. Dunn, R. E., Strömberg, C. A. E., Madden, R. H., Kohn, M. J. & Carlini, A. A. Linked canopy, climate,
- 566 and faunal change in the Cenozoic of Patagonia. *Science (80-.).* **347**, 258–261 (2015).
- 567 35. Meseguer, A. S. & Condamine, F. L. Ancient tropical extinctions at high latitudes contributed to the
- 568 latitudinal diversity gradient. *Evolution (N. Y.)*. evo.13967 (2020). doi:10.1111/evo.13967
- 569 36. Herbert, T. D. *et al.* Late Miocene global cooling and the rise of modern ecosystems. *Nat. Geosci.* **9**,
- 570 843–847 (2016).
- 571 37. Pound, M. J. *et al.* A Tortonian (Late Miocene, 11.61–7.25Ma) global vegetation reconstruction.
- 572 *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **300**, 29–45 (2011).
- 573 38. Svenning, J. C., Borchsenius, F., Bjorholm, S. & Balslev, H. High tropical net diversification drives the

- 574 New World latitudinal gradient in palm (Arecaceae) species richness. in *Journal of Biogeography* **35**,
575 394–406 (John Wiley & Sons, Ltd (10.1111), 2008).
- 576 39. Sandel, B. *et al.* The influence of late quaternary climate-change velocity on species endemism.
577 *Science* (80-.). **334**, 660–664 (2011).
- 578 40. McGlone, M. S., Buitenwerf, R. & Richardson, S. J. The formation of the oceanic temperate forests of
579 New Zealand. *New Zealand Journal of Botany* **54**, 128–155 (2016).
- 580 41. Liebergesell, M. *et al.* Functional resilience against climate-driven extinctions - Comparing the
581 functional diversity of European and north American tree floras. *PLoS One* **11**, e0148607 (2016).
- 582 42. Svenning, J.-C., Eiserhardt, W. L., Normand, S., Ordonez, A. & Sandel, B. The Influence of
583 Paleoclimate on Present-Day Patterns in Biodiversity and Ecosystems. *Annu. Rev. Ecol. Evol. Syst.* **46**,
584 551–572 (2015).
- 585 43. Dynesius, M. & Jansson, R. Evolutionary consequences of changes in species' geographical
586 distributions driven by Milankovitch climate oscillations. *Proc. Natl. Acad. Sci. U. S. A.* **97**, 9115–9120
587 (2000).
- 588 44. Devictor, V. *et al.* Spatial mismatch and congruence between taxonomic, phylogenetic and
589 functional diversity: the need for integrative conservation strategies in a changing world. *Ecol. Lett.*
590 **13**, no-no (2010).
- 591 45. Pollock, L. J., Thuiller, W. & Jetz, W. Large conservation gains possible for global biodiversity facets.
592 *Nature* **546**, 141–144 (2017).
- 593 46. Mazel, F., Mooers, A. O., Riva, G. V. D. & Pennell, M. W. Conserving Phylogenetic Diversity Can Be a
594 Poor Strategy for Conserving Functional Diversity. *Syst. Biol.* **66**, 1019–1027 (2017).
- 595 47. Guo, W.-Y. *et al.* Half of the world's tree biodiversity is unprotected and is increasingly threatened
596 by human activities. *bioRxiv* 2020.04.21.052464 (2020). doi:10.1101/2020.04.21.052464
- 597 48. Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M. & West, G. B. Toward a metabolic theory of
598 ecology. in *Ecology* **85**, 1771–1789 (2004).
- 599 49. Brown, J. H. Why are there so many species in the tropics? *J. Biogeogr.* **41**, 8–22 (2014).
- 600 50. Steinbauer, M. J. *et al.* Accelerated increase in plant species richness on mountain summits is linked
601 to warming. *Nature* **556**, 231–234 (2018).
- 602 51. Gillooly, J. F. & Allen, A. P. Linking global patterns in biodiversity to evolutionary dynamics using
603 metabolic theory. *Ecology* **88**, 1890–4 (2007).
- 604 52. Wright, I. J. *et al.* The worldwide leaf economics spectrum. *Nature* **428**, 821–827 (2004).
- 605 53. Wright, I. J. *et al.* Global climatic drivers of leaf size. *Science* (80-.). **357**, 917–921 (2017).
- 606 54. Swenson, N. G. *et al.* Constancy in functional space across a species richness anomaly. *Am. Nat.* **187**,
607 E83–E92 (2016).
- 608 55. Ordonez, A. & Svenning, J. C. Greater tree species richness in eastern North America compared to
609 Europe is coupled to denser, more clustered functional trait space filling, not to trait space
610 expansion. *Glob. Ecol. Biogeogr.* **27**, 1288–1299 (2018).
- 611 56. Beech, E., Rivers, M., Oldfield, S. & Smith, P. P. GlobalTreeSearch: The first complete global database
612 of tree species and country distributions. *J. Sustain. For.* **36**, 454–489 (2017).

- 613 57. Serra-Diaz, J. M., Enquist, B. J., Maitner, B., Merow, C. & Svenning, J. C. Big data of tree species
614 distributions: how big and how good? *For. Ecosyst.* **4**, 30 (2017).
- 615 58. Hillebrand, H. On the Generality of the Latitudinal Diversity Gradient. *Am. Nat.* **163**, 192–211 (2004).
- 616 59. Gaston, K. J. Global patterns in biodiversity. *Nature* **405**, 220–227 (2000).
- 617 60. Pianka, E. R. Latitudinal Gradients in Species Diversity: A Review of Concepts. *Am. Nat.* **100**, 33–46
618 (2002).
- 619 61. Barthlott, W. *et al.* Geographic Patterns of Vascular Plant Diversity at Continental to Global Scales
620 (Geographische Muster der Gefäßpflanzenvielfalt im kontinentalen und globalen Maßstab).
621 *Erdkunde* **61**, 305–315 (2007).
- 622 62. Lamanna, C. *et al.* Functional trait space and the latitudinal diversity gradient. *Proc. Natl. Acad. Sci. U.*
623 *S. A.* **111**, 13745–13750 (2014).
- 624 63. Wing, S. L. *et al.* Late Paleocene fossils from the Cerrejon Formation, Colombia, are the earliest
625 record of Neotropical rainforest. *Proc. Natl. Acad. Sci.* **106**, 18627–18632 (2009).
- 626 64. Antonelli, A. *et al.* Amazonia is the primary source of Neotropical biodiversity. *Proc. Natl. Acad. Sci.*
627 **115**, 6034–6039 (2018).
- 628 65. Chu, C. *et al.* Direct and indirect effects of climate on richness drive the latitudinal diversity gradient
629 in forest trees. *Ecology Letters* **22**, 245–255 (2019).
- 630 66. Fine, P. V. A. Ecological and Evolutionary Drivers of Geographic Variation in Species Diversity. *Annu.*
631 *Rev. Ecol. Evol. Syst.* **46**, 369–392 (2015).
- 632 67. Feng, G. *et al.* Species and phylogenetic endemism in angiosperm trees across the Northern
633 Hemisphere are jointly shaped by modern climate and glacial–interglacial climate change. *Glob. Ecol.*
634 *Biogeogr.* **28**, 1393–1402 (2019).
- 635 68. Kissling, W. D. *et al.* Quaternary and pre-Quaternary historical legacies in the global distribution of a
636 major tropical plant lineage. *Glob. Ecol. Biogeogr.* **21**, 909–921 (2012).
- 637 69. Blonder, B. *et al.* Linking environmental filtering and disequilibrium to biogeography with a
638 community climate framework. *Ecology* **96**, 972–985 (2015).
- 639 70. Ordonez, A. & Svenning, J. C. Functional diversity of North American broad-leaved trees is
640 codetermined by past and current environmental factors. *Ecosphere* **7**, (2016).
- 641 71. Pound, M. J., Haywood, A. M., Salzmann, U. & Ridg, J. B. Global vegetation dynamics and
642 latitudinal temperature gradients during the Mid to Late Miocene (15.97–5.33Ma). *Earth-Science*
643 *Rev.* **112**, 1–22 (2012).
- 644 72. Osborne, C. P. Atmosphere, ecology and evolution: What drove the Miocene expansion of C4
645 grasslands? *Journal of Ecology* **96**, 35–45 (2008).
- 646 73. Greenop, R., Foster, G. L., Wilson, P. A. & Lear, C. H. Middle Miocene climate instability associated
647 with high-amplitude CO₂ variability. *Paleoceanography* **29**, 845–853 (2014).
- 648 74. Steinhorsdottir, M., Jardine, P. E. & Rember, W. C. Near-Future p CO₂ during the hot Mid Miocene
649 Climatic Optimum. *Paleoceanogr. Paleoclimatology* (2020). doi:10.1029/2020PA003900
- 650 75. Lucas, E. J. & Bünge, M. O. Myrtaceae in the Atlantic forest: their role as a ‘model’ group. *Biodivers.*
651 *Conserv.* **24**, 2165–2180 (2015).

- 652 76. Deng, M., Jiang, X. L., Hipp, A. L., Manos, P. S. & Hahn, M. Phylogeny and biogeography of East Asian
653 evergreen oaks (*Quercus* section *Cyclobalanopsis*; Fagaceae): Insights into the Cenozoic history of
654 evergreen broad-leaved forests in subtropical Asia. *Mol. Phylogenet. Evol.* **119**, 170–181 (2018).
- 655 77. Piñeiro, R., Dauby, G., Kaymak, E. & Hardy, O. J. Pleistocene population expansions of shade-tolerant
656 trees indicate fragmentation of the African rainforest during the ice ages. *Proc. R. Soc. B Biol. Sci.*
657 **284**, 20171800 (2017).
- 658 78. Zozomová-Lihová, J., Melichárková, A., Svitok, M. & Španiel, S. Pleistocene range disruption and
659 postglacial expansion with secondary contacts explain the genetic and cytotype structure in the
660 western Balkan endemic *Alyssum austrodalmaticum* (Brassicaceae). *Plant Syst. Evol.* **306**, 1–25
661 (2020).
- 662 79. Niinemets, Ü. & Valladares, F. Tolerance to shade, drought, and waterlogging of temperate northern
663 hemisphere trees and shrubs. *Ecol. Monogr.* **76**, 521–547 (2006).
- 664 80. Conradi, T., Van Meerbeek, K., Ordonez, A. & Svenning, J. Biogeographic historical legacies in the net
665 primary productivity of Northern Hemisphere forests. *Ecol. Lett.* **23**, 800–810 (2020).
- 666 81. Muscarella, R. *et al.* Functional convergence and phylogenetic divergence during secondary
667 succession of subtropical wet forests in Puerto Rico. *J. Veg. Sci.* **27**, 283–294 (2016).
- 668 82. Swenson, N. G. & Enquist, B. J. Opposing assembly mechanisms in a Neotropical dry forest:
669 implications for phylogenetic and functional community ecology. *Ecology* **90**, 2161–2170 (2009).
- 670 83. Kraft, N. J. B., Valencia, R. & Ackerly, D. D. Functional traits and niche-based tree community
671 assembly in an Amazonian forest. *Science (80-.).* **322**, 580–582 (2008).
- 672 84. Hubbell, S. P. Neutral theory and the evolution of ecological equivalence. *Ecology* **87**, 1387–1398
673 (2006).
- 674 85. Boyle, B. *et al.* The taxonomic name resolution service: An online tool for automated standardization
675 of plant names. *BMC Bioinformatics* **14**, 16 (2013).
- 676 86. Enquist, B. J., Condit, R. R., Peet, R. K., Schildhauer, M. & Thiers, B. M. The Botanical Information and
677 Ecology Network (BIEN): Cyberinfrastructure for an integrated botanical information network to
678 investigate the ecological impacts of global climate change on plant biodiversity. *PeerJ* **4**, e2615v2
679 (2016).
- 680 87. Enquist, B. J. *et al.* The commonness of rarity: Global and future distribution of rarity across land
681 plants. *Sci. Adv.* **5**, eaaz0414 (2019).
- 682 88. Banda, K. R. *et al.* Plant diversity patterns in neotropical dry forests and their conservation
683 implications. *Science (80-.).* **353**, 1383–1387 (2016).
- 684 89. Dauby, G. *et al.* RAINBIO: a mega-database of tropical African vascular plants distributions.
685 *PhytoKeys* **74**, 1–18 (2016).
- 686 90. Pateiro-López, B. & Rodríguez-Casal, A. Generalizing the Convex Hull of a Sample: The R Package
687 alphahull. *J. Stat. Softw.* **34**, (2015).
- 688 91. Kirkpatrick, D. G. & Seidel, R. On the Shape of a Set of Points in the Plane. *IEEE Trans. Inf. Theory* **29**,
689 551–559 (1983).
- 690 92. R Core Team. R: A Language and Environment for Statistical Computing. (2019).

93. Vilela, B. & Villalobos, F. LetsR: A new R package for data handling and analysis in macroecology. *Methods Ecol. Evol.* **6**, 1229–1234 (2015).
94. Smith, S. A. & Brown, J. W. Constructing a broadly inclusive seed plant phylogeny. *Am. J. Bot.* **105**, 302–314 (2018).
95. Magallón, S., Gómez-Acevedo, S., Sánchez-Reyes, L. L. & Hernández-Hernández, T. A metacalibrated time-tree documents the early rise of flowering plant phylogenetic diversity. *New Phytol.* **207**, 437–453 (2015).
96. Díaz, S. *et al.* The global spectrum of plant form and function. *Nature* **529**, 167–171 (2016).
97. Kattge, J. *et al.* TRY - a global database of plant traits. *Glob. Chang. Biol.* **17**, 2905–2935 (2011).
98. Kattge, J. *et al.* TRY plant trait database – enhanced coverage and open access. *Glob. Chang. Biol.* **26**, 119–188 (2020).
99. Aubin, I., Beaudet, M. & Messier, C. Light extinction coefficients specific to the understory vegetation of the southern boreal forest, Quebec. *Can. J. For. Res.* **30**, 168–177 (2000).
100. Aubin, I., Messier, C. & Kneeshaw, D. Population structure and growth acclimation of mountain maple along a successional gradient in the southern boreal forest. *Écoscience* **12**, 540–548 (2005).
101. Aubin, I. *et al.* Traits to stay, traits to move: A review of functional traits to assess sensitivity and adaptive capacity of temperate and boreal trees to climate change. *Environmental Reviews* **24**, 164–186 (2016).
102. Thiffault, N., Titus, B. D. & Munson, A. D. Silvicultural options to promote seedling establishment on Kalmia-Vaccinium-dominated sites. *Scand. J. For. Res.* **20**, 110–121 (2005).
103. Titus, B. D. *et al.* Post-harvest nitrogen cycling in clearcut and alternative silvicultural systems in a montane forest in coastal British Columbia. *For. Chron.* **82**, 844–859 (2006).
104. Wiebe, S., Morris, D., Luckai, N. & Reid, D. Coarse Woody Debris Dynamics Following Biomass Harvesting: Tracking Carbon and Nitrogen Patterns During Early Stand Development in Upland Black Spruce Ecosystems. *Int. J. For. Eng.* **23**, 25–32 (2012).
105. Masse, J., Prescott, C. E., Müller, C. & Grayston, S. J. Gross nitrogen transformation rates differ in reconstructed oil-sand soils from natural boreal-forest soils as revealed using a ¹⁵N tracing method. *Geoderma* **282**, 37–48 (2016).
106. Shan, H. *et al.* Gap Filling in the Plant Kingdom---Trait Prediction Using Hierarchical Probabilistic Matrix Factorization. in *Proceedings of the 29th International Conference on Machine Learning* 1303–1310 (2012).
107. Fazayeli, F., Banerjee, A., Kattge, J., Schrod, F. & Reich, P. B. Uncertainty quantified matrix completion using bayesian hierarchical matrix factorization. in *Proceedings - 2014 13th International Conference on Machine Learning and Applications, ICMLA 2014* 312–317 (IEEE, 2014). doi:10.1109/ICMLA.2014.56
108. Schrod, F. *et al.* BHPMF - a hierarchical Bayesian approach to gap-filling and trait prediction for macroecology and functional biogeography. *Glob. Ecol. Biogeogr.* **24**, 1510–1521 (2015).
109. Dynesius, M. & Jansson, R. Evolutionary consequences of changes in species' geographical distributions driven by Milankovitch climate oscillations. *Proc. Natl. Acad. Sci.* **97**, 9115–9120 (2000).

- 730 110. Jetz, W. & Rahbek, C. Geographic range size and determinants of avian species richness. *Science* (80-
731 .). **297**, 1548–1551 (2002).
- 732 111. Fick, S. E. & Hijmans, R. J. WorldClim 2: new 1-km spatial resolution climate surfaces for global land
733 areas. *Int. J. Climatol.* **37**, 4302–4315 (2017).
- 734 112. Hill, D. J. The non-analogue nature of Pliocene temperature gradients. *Earth Planet. Sci. Lett.* **425**,
735 232–241 (2015).
- 736 113. Brown, J. L., Hill, D. J., Dolan, A. M., Carnaval, A. C. & Haywood, A. M. Paleoclim, high spatial
737 resolution paleoclimate surfaces for global land areas. *Sci. Data* **5**, 180254 (2018).
- 738 114. Dolan, A. M. *et al.* Modelling the enigmatic Late Pliocene Glacial Event - Marine Isotope Stage M2.
739 *Glob. Planet. Change* **128**, 47–60 (2015).
- 740 115. Karger, D. N. *et al.* Climatologies at high resolution for the earth's land surface areas. *Sci. Data* **4**,
741 170122 (2017).
- 742 116. Otto-Bliesner, B. L., Marshall, S. J., Overpeck, J. T., Miller, G. H. & Hu, A. Simulating arctic climate
743 warmth and icefield retreat in the last interglaciation. *Science* (80-.). **311**, 1751–1753 (2006).
- 744 117. Kreft, H. & Jetz, W. Global patterns and determinants of vascular plant diversity. *Proc. Natl. Acad. Sci.*
745 **104**, 5925–5930 (2007).
- 746 118. Zhang, J., Nielsen, S. E., Mao, L., Chen, S. & Svenning, J. C. Regional and historical factors supplement
747 current climate in shaping global forest canopy height. *J. Ecol.* **104**, 469–478 (2016).
- 748 119. Kennedy, C. M., Oakleaf, J. R., Theobald, D. M., Baruch-Mordo, S. & Kiesecker, J. Managing the
749 middle: A shift in conservation priorities based on the global human modification gradient. *Glob.*
750 *Chang. Biol.* **25**, 811–826 (2019).
- 751 120. Sanderson, E. W. *et al.* The Human Footprint and the Last of the Wild. *Bioscience* **52**, 891–904 (2002).
- 752 121. Schulze, K. *et al.* An assessment of threats to terrestrial protected areas. *Conservation Letters* **11**,
753 e12435 (2018).
- 754 122. Slavich, E., Warton, D. I., Ashcroft, M. B., Gollan, J. R. & Ramp, D. Topoclimate versus macroclimate:
755 How does climate mapping methodology affect species distribution models and climate change
756 projections? *Divers. Distrib.* **20**, 952–963 (2014).
- 757 123. Stein, A., Gerstner, K. & Kreft, H. Environmental heterogeneity as a universal driver of species
758 richness across taxa, biomes and spatial scales. *Ecology Letters* **17**, 866–880 (2014).
- 759 124. Morrone, J. J. Biogeographical regions under track and cladistic scrutiny. *Journal of Biogeography* **29**,
760 149–152 (2002).
- 761 125. Olson, D. M. *et al.* Terrestrial Ecoregions of the World: A New Map of Life on Earth. *Bioscience* **51**,
762 933 (2001).
- 763 126. Harter, D. E. V. *et al.* Impacts of global climate change on the floras of oceanic islands - Projections,
764 implications and current knowledge. *Perspectives in Plant Ecology, Evolution and Systematics* **17**,
765 160–183 (2015).
- 766 127. Mouchet, M. *et al.* Towards a consensus for calculating dendrogram-based functional diversity
767 indices. *Oikos* **117**, 794–800 (2008).
- 768 128. Oksanen, J. *et al.* vegan: Community Ecology Package. (2019).

769 129. Müllner, D. fastcluster²: Fast Hierarchical, Agglomerative Clustering Routines for R and Python. *J.*
770 *Stat. Softw.* **53**, 1–18 (2015).

771 130. Kembel, S. W. *et al.* Picante: R tools for integrating phylogenies and ecology. *Bioinformatics* **26**,
772 1463–1464 (2010).

773 131. Osorio, F., Vallejos, R. & Cuevas, F. SpatialPack: Computing the Association Between Two Spatial
774 Processes. (2016).

775 132. Kissling, W. D. & Carl, G. Spatial autocorrelation and the selection of simultaneous autoregressive
776 models. *Glob. Ecol. Biogeogr.* **17**, 59–71 (2008).

777 133. Nagelkerke, N. J. D. A note on a general definition of the coefficient of determination. *Biometrika* **78**,
778 691–692 (1991).

779 134. Burnham, K. P. & Anderson, D. R. Multimodel inference: Understanding AIC and BIC in model
780 selection. *Sociological Methods and Research* **33**, 261–304 (2004).

781 135. Bivand, R. S. & Wong, D. W. S. Comparing implementations of global and local indicators of spatial
782 association. *Test* **27**, 716–748 (2018).

783 136. Schielzeth, H. Simple means to improve the interpretability of regression coefficients. *Methods Ecol.*
784 *Evol.* **1**, 103–113 (2010).

Acknowledgments

We thank Brad Boyle for valuable database and informatics assistance and advice, and TRY contributors for sharing their data. This work was conducted as a part of the BIEN Working Group, 2008–2012. We thank all the data contributors and numerous herbaria who have contributed their data to various data compiling organizations (see the herbarium list below) for the invaluable data and support provided to BIEN. We thank the New York Botanical Garden; Missouri Botanical Garden; Utrecht Herbarium; the UNC Herbarium; and GBIF, REMIB, and SpeciesLink. The staff at CyVerse provided critical computational assistance.

We acknowledge the herbaria that contributed data to this work: A, AAH, AAS, AAU, ABH, ACAD, ACOR, AD, AFS, AK, AKPM, ALCB, ALTA, ALU, AMD, AMES, AMNH, AMO, ANGU, ANSM, ANSP, AQP, ARAN, ARIZ, AS, ASDM, ASU, AUT, AV, AWH, B, BA, BAA, BAB, BABY, BACP, BAF, BAFC, BAI, BAJ, BAL, BARC, BAS, BBB, BBS, BC, BCMEX, BCN, BCRU, BERE, BESA, BG, BH, BHCB, BIO, BISH, BLA, BM, BOCH, BOL, BOLV, BONN, BOON, BOTU, BOUM, BPI, BR, BREM, BRI, BRIT, BRLU, BRM, BSB, BUT, C, CALI, CAN, CANB, CANU, CAS, CATA, CATIE, CAY, CBM, CDA, CDBI, CEN, CEPEC, CESJ, CGE, CGMS, CHAM, CHAPA, CHAS, CHR, CHSC, CIB, CICY, CIIDIR, CIMI, CINC, CLEMS, CLF, CMM, CMMEX, CNPO, CNS, COA, COAH, COCA, CODAGEM, COFC, COL, COLO, CONC, CORD, CP, CPAP, CPUN, CR, CRAI, CRP, CS, CSU, CSUSB, CTES, CTESN, CU, CUVC, CUZ, CVRD, DAO, DAV, DBG, DBN, DES, DLF, DNA, DPU, DR, DS, DSM, DUKE, DUSS, E, EA, EAC, EAN, EBUM, ECON, EIF, EIU, EMMA, ENCB, ER, ERA, ESA, ETH, F, FAA, FAU, FAUC, FB, FCME, FCO, FCQ, FEN, FHO, FI, FLAS, FLOR, FM, FR, FRU, FSU, FTG, FUEL, FULD, FURB, G, GAT, GB, GDA, GENT, GES, GH, GI, GLM, GMDRC, GMNHJ, GOET, GRA, GUA, GZU, H, HA, HAC, HAL, HAM, HAMAB, HAO, HAS, HASU, HB, HBG, HBR, HCIB, HEID, HGM, HIB, HIP, HNT, HO, HPL, HRCB, HRP, HSC, HSS, HU, HUA, HUAA, HUAL, HUAZ, HUCP, HUEFS, HUEM, HUFU, HUI, HUSA, HUT, HXBH, HYO, IAA, IAC, IAN, IB, IBGE, IBK, IBSC, IBUG, ICEL, ICESI, ICN, IEA, IEB, ILL, ILLS, IMSSM, INB, INEGI, INIF, INM, INPA, IPA, IPRN, IRVC, ISC, ISKW, ISL, ISTC, ISU, IZAC, IZTA, JACA, JBAG, JBGP, JCT, JE, JEPS, JOTR, JROH, JUA, JYV, K, KIEL, KMN, KMNH, KOELN, KOR, KPM, KSC, KSTC, KSU, KTU, KU, KUN, KYO, L, LA, LAGU, LBG, LD, LE, LEB, LIL, LINC, LINN, LISE, LISI, LISU, LL, LMS, LOJA, LOMA, LP, LPAG, LPB, LPD, LPS, LSU, LSUM, LTB, LTR, LW, LYJB, LZ, M, MA, MACF, MAF, MAK, MARS, MARY, MASS, MB, MBK, MBM, MBML,

MCNS, MEL, MELU, MEN, MERL, MEXU, MFA, MFU, MG, MGC, MICH, MIL, MIN, MISSA, MJG, MMMN, MNHM, MNHN, MO, MOL, MOR, MPN, MPU, MPUC, MSB, MSC, MSUN, MT, MTMG, MU, MUB, MUR, MVFA, MVFQ, MVJB, MVM, MW, MY, N, NA, NAC, NAS, NCU, NE, NH, NHM, NHMC, NHT, NLH, NM, NMB, NMNL, NMR, NMSU, NSPM, NSW, NT, NU, NUM, NY, NZFRI, O, OBI, ODU, OS, OSA, OSC, OSH, OULU, OWU, OXF, P, PACA, PAMP, PAR, PASA, PDD, PE, PEL, PERTH, PEUFR, PFC, PGM, PH, PKDC, PLAT, PMA, POM, PORT, PR, PRC, PRE, PSU, PY, QCA, QCNE, QFA, QM, QRS, QUE, R, RAS, RB, RBR, REG, RELC, RFA, RIOC, RM, RNG, RSA, RYU, S, SACT, SALA, SAM, SAN, SANT, SAPS, SASK, SAV, SBBG, SBT, SCFS, SD, SDSU, SEL, SEV, SF, SFV, SGO, SI, SIU, SJRP, SJSU, SLPM, SMDB, SMF, SNM, SOM, SP, SPF, SPSF, SQF, SRFA, STL, STU, SUU, SVG, TAES, TAI, TAIF, TALL, TAM, TAMU, TAN, TASH, TEF, TENN, TEPB, TEX, TFC, TI, TKPM, TNS, TO, TOYA, TRA, TRH, TROM, TRT, TRTE, TU, TUB, U, UADY, UAM, UAMIZ, UB, UBC, UC, UCMM, UCR, UCS, UCSB, UCSC, UEC, UESC, UFG, UFMA, UFMT, UFP, UFRJ, UFRN, UFS, UGDA, UH, UI, UJAT, ULM, ULS, UME, UMO, UNA, UNB, UNCC, UNEX, UNITEC, UNL, UNM, UNR, UNSL, UPGB, UPEI, UPNA, UPS, US, USAS, USF, USJ, USM, USNC, USP, USZ, UT, UTC, UTEP, UU, UVIC, UWO, V, VAL, VALD, VDB, VEN, VIT, VMSL, VT, W, WAG, WAT, WELT, WFU, WII, WIN, WIS, WMNH, WOLL, WS, WTU, WU, XAL, YAMA, Z, ZMT, ZSS, and ZT.

W.-Y.G., J.M.S.-D., and J.-C.S. acknowledge support from the Danish Council for Independent Research | Natural Sciences (Grant 6108-00078B) to the TREECHANGE project. J.-C.S. also considers this work a contribution to his VILLUM Investigator project “Biodiversity Dynamics in a Changing World” funded by VILLUM FONDEN. C.B. was supported by the National Research Foundation of Korea (NRF) grant funded by the Korean government (MSIT) (2018R1C1B6005351). A.S.M. was supported by the Environment Research and Technology Development Fund (S-14) of the Ministry of the Environment, Japan. J.P. (Jan Pisek) was supported by the Estonian Research Council grant PUT1355. J.P. (Josep Peñuelas) was funded by the European Research Council Synergy grant ERC-2013-SyG-610028 IMBALANCE-P. A.G.G. (Alvaro G. Gutiérrez) was funded by FONDECYT 11150835 and 1200468. The BIEN working group was supported by the National Center for Ecological Analysis and Synthesis, a center funded by NSF EF-0553768 at the University of California, Santa Barbara and the State of California. Additional support for the BIEN working group was provided by iPlant/CyVerse via NSF DBI-0735191. B.J.E. and C.M. were supported by NSF ABI-1565118 and NSF HDR-1934790. B.J.E.

was also supported by the Global Environment Facility SPARC project grant (GEF-5810). B.J.E., C.V., and B.S.M. acknowledge the FREE group funded by the synthesis center CESAB of the French Foundation for Research on Biodiversity (FRB) and EDF. J.-C.S. and B.J.E. acknowledge support from the Center for Informatics Research on Complexity in Ecology (CIRCE), funded by the Aarhus University Research Foundation under the AU Ideas program.

Author contributions

W.-Y.G., J.M.S.-D., and J.-C.S. conceived the project; J.M.S.-D., W.-Y.G., and all others collected the data; W.-Y.G. analyzed the data; W.-Y.G. interpreted the data; W.-Y.G., J.M.S.-D., and J.-C.S. wrote the manuscript. All authors contributed data, discussed the results, revised manuscript drafts, and contributed to writing and approved the final manuscript.

Competing interests

The authors declare no competing interests.

Data and materials availability:

All the occurrences are deposited in BIEN (<https://bien.nceas.ucsb.edu/bien/>), and the phylogeny and imputed functional trait data are available via ref. ⁴⁷.

Figures

Fig. 1 Global patterns of tree (a) species richness, (c) functional diversity, and (e) phylogenetic diversity. In (b), (d) and (f), the fitted line is the lowess regression. Maps use the Behrmann projection at 110 km × 110 km spatial resolution. Myr: Million years.

Fig. 2 Effects of the tested environmental variables on tree (a) specie richness (SR), (b) functional diversity (FD) and (c) phylogenetic diversity (PD). Estimates (± 1 standard error) of effects were obtained from simultaneous autoregressive (SAR) models. Different colors and shapes indicate biogeographic regions. Non-significant variables ($p > 0.05$) are indicated in grey. Results from OLS models are shown in Table S2. HMc: human modification index; MAT: mean annual temperature; AP: Annual precipitation; IG: Pleistocene Interglacial; LGM: Last Glacial Maximum. Anomaly was calculated as the past minus the present state.

874

875 **Fig. 3** Global patterns of the residuals (deviation) from the ordinary least regression between functional
876 diversity (FD) and phylogenetic diversity (PD) ($FD = 0.90PD$, $R^2 = 0.987$, $p < 0.0001$). Brown (positive)
877 areas are areas of higher FD than expected based on PD, whereas blue (negative) areas depict areas with
878 lower FD than expected from the observed PD. Map uses the Behrmann projection at $110 \text{ km} \times 110 \text{ km}$
879 spatial resolution.

Fig. 4 Effects of the tested environmental variables on the residuals from the regression between functional diversity (FD) and phylogenetic diversity (PD) (Fig. 3). Estimate (± 1 standard error) of effects were obtained from simultaneous autoregressive (SAR) models. Different colors and shapes indicate biogeographic regions. Non-significant variables ($p > 0.05$) are indicated in grey. Results from OLS models are shown in Table S3. HMc: human modification index; MAT: mean annual temperature; AP: Annual precipitation; IG: Pleistocene Interglacial; LGM: Last Glacial Maximum. Anomaly was calculated as the past minus the present state.

Fig. 5 Bivariate regressions between the residuals from the regression of functional diversity (FD) on phylogenetic diversity (PD) and significant explanatory climate variables. Anomaly was calculated as the past minus the present state. For each subplot, only significant relationships are shown (Fig. 4, $p < 0.05$).