Spatio-temporal surveys of the brown meagre *Sciaena umbra* using passive acoustics for management and conservation

Lucia Di Iorio^{1*}, Patrick Bonhomme², Noémie Michez³, Bruno Ferrari³, Alexandra Gigou⁴, Pieraugusto Panzalis⁵, Elena Desiderà⁵, Augusto Navone⁵, Pierre Boissery⁶, Julie Lossent¹, Benjamin Cadville⁷, Marie Bravo-Monin⁷, Eric Charbonnel⁷, Cédric Gervaise¹

1 CHORUS Institute, INP Phelma Minatec, 38016 Grenoble, France

2 Parc national des Calanques, 13008 Marseille, France

3 Parc naturel marin du golfe du Lion / Office français de la biodiversité, 66700 Argelès-sur-Mer, France

4 Délégation de façade maritime Méditerranée, Office français de la biodiversité, 13001 Marseille, France

5 Area Marina Protetta di Tavolara – Punta Coda Cavallo, 07026, Olbia, Italy

6 Agence de l'Eau Rhône Méditerranée Corse, Imm Le Noailles, 13001 Marseille, France

7 Parc marin de la Côte Bleue, 13620 Carry-le-Rouet, France

* Corresponding author: lucia.diiorio@chorusacoustics.com

Abstract

Conservation of exploited fish populations is a priority for environmental managers. Spatio-temporal knowledge on reproductive sites is mandatory for species and habitat conservation but is often difficult to assess, particularly over vast geographic areas. Regular and long-term standardized surveys are necessary to identify reproductive sites, assess population trends and their distribution. Here we emphasize the utility of Passive Acoustic Monitoring (PAM) for the survey and management of a depleted vulnerable Mediterranean fish species, the brown meagre, *Sciaena umbra*. Acoustic surveys of reproductive calls were conducted combining 1) spatial data from standardized surveys within three MPAs and from 49 unprotected sites throughout the Northwestern Mediterranean basin, as well as 2) temporal data from a two-year-long survey at a presumed spawning location. The MPA surveys, which rapidly scanned ~30-50 km of the rocky coastlines per MPA, unveiled maps of distribution and reproductive activity of the brown meagre, including potential spawning sites. They were also effective in emphasizing effects linked to management actions: Full-protection zones had a higher number of vocalizations (70% of the listening sites) compared to less protected zones (30% of the sites) or sites outside MPAs (45% of the sites). This was also reflected in the number of singers that was generally low (< 3 individuals) in less protected zones and outside MPAs, implying lower fish densities. Highest calling aggregations were observed in potential spawning areas that represented only 0.04% of all listening sites, and were almost all in older, fully protected MPAs, which thus play a key role for fish stock recovery. The two-year survey revealed a 5-month reproductive season (from May to October) with a strong positive correlation between calling activity and temperature. Overall this study confirms the role of PAM as an efficient, replicable and standardized non-invasive method for population management that can identify functional sites and key protection zones, provide valuable information on reproduction, spatial and temporal occurrence, but also on population trends and climate-driven changes.

45

46

47 **Keywords:** vulnerable fish, reproductive sites, fish sounds, passive acoustic monitoring, marine
48 protected areas, Mediterranean Sea

49

50 **Highlights**

- 51 • Monitoring of threatened species and their key habitats is critical for environmental
52 managers.
- 53 • Management requires methods to assess population trends at large spatial and temporal
54 scales.
- 55 • Passive acoustics (PA) is efficient in mapping and monitoring vulnerable fish species.
- 56 • Distribution, reproductive sites and population dynamics can be assessed over vast
57 geographical areas.
- 58 • We show the utility of PA to identify key conservation zones and assess effects of
59 management actions.

60

1 INTRODUCTION

Knowledge of spawning habitats and breeding sites is essential for the conservation of exploited fish stocks. Recording when and where a fish species is reproducing provides pivotal data on key habitats and allows to better manage both vulnerable species and their critical habitats (Luczkovich et al., 2008b). Identification, mapping and monitoring of functional fish habitats is therefore a priority for environmental managers of Marine Protected Areas (MPA), in particular with regard to endangered and over-exploited species (Harmelin and Ruitton, 2006; Luczkovich et al., 2008a). Regular and long-term surveys of target species at large spatial scales are necessary to evaluate population trends, distribution, recovery, and the efficiency of protection measures (Guidetti et al., 2014a; Harmelin-Vivien et al., 2015). However, although very effective, traditional methods based on visual surveys are logistically difficult to implement at high-frequency intervals, over wide areas and for nocturnally active organisms (Harmelin-Vivien et al. 2015). Management programs would therefore benefit from complementary standardized and replicable methods allowing to cover long distances over short time periods and to monitor key sites and key species continuously, over the long-term with limited human investment.

Passive acoustic monitoring (PAM) has proven to be a powerful tool for both conservation and fishery science (e.g., Gannon, 2008; Luczkovich et al., 2008a; Mann et al., 2008). It is a non-invasive flexible listening technique that involves the use of hydrophones to record the collection of sounds emanated from an environment (i.e., the soundscape). PAM exploits the sounds emitted by fish mainly for communication (e.g., Amorim, 2006; Ladich, 2015) to help identifying species occurrence, spawning areas, cryptic species, follow reproductive activity and population turnover, and to appraise diversity (Desiderà et al., 2019; Lobel, 2002; Luczkovich et al., 2008b; Picciulin et al., 2018). Despite this, and the fact that a variety of

commercial and vulnerable fish species produce sounds, there is a limited number of studies adopting PAM for large-scale mapping of protected fish species and their critical habitats in the Mediterranean Sea, and they all concern the brown meagre *Sciaena umbra* Linnaeus 1758 (Bonacito et al., 2002; Parmentier et al., 2018; Picciulin et al., 2012).

The brown meagre is an iconic demersal fish species of Mediterranean coastal habitats listed in the Annexe III (Protected Fauna Species) of the Barcelona and Berne Conventions. Globally, this species is classified as Near Threatened (Chao, 2015) and as vulnerable in the Mediterranean Sea by the IUCN (Abdul Malak et al., 2011; Bizsel et al., 2011). *S. umbra* is highly vulnerable to fishing due to its aggregative behaviour and easily accessible shallow-water habitat (Lloret et al., 2008). With a population decline in the Mediterranean Sea of approximately 70% between 1980 and 2005 (Bizsel et al., 2011), the brown meagre falls within the categories of seriously depleted and collapsed fish species. This is why in 2013, France adopted a moratorium banning spear and recreational fishing of the brown meagre, which was renewed in December 2018. *S. umbra* is also of high ecological and conservation value as it is considered a useful bioindicator (Mouillot et al., 2002). Its presence is indicative of high environmental quality and community richness (García-Rubies et al., 2013; Mouillot et al., 2002; Picciulin et al., 2013), and it is very sensitive to the effect of protection measures (Guidetti et al., 2014b; Harmelin-Vivien et al., 2015b; Harmelin et al., 1995). Although there is conspicuous increase in its abundance in no-take integral reserves (Francour, 1991; García-Rubies et al., 2013; Harmelin, 2013), outside most protected areas, the brown meagre is frequently targeted (Lloret et al., 2008; Morales-Nin et al., 2005) and found to be rare or absent (Harmelin-Vivien et al., 2015b; Letourneur et al., 2003). Appropriate monitoring of *S. umbra* populations and their trends should involve the combination of extensive large-scale observations with regular long-term monitoring of key sites within and outside MPAs

(Harmelin & Ruitton, 2007; Harmelin, 2013). This would allow appropriate assessment of species distribution as well as its spatial extension and to identify and follow functional sites (e.g., feeding, reproduction, spawning).

S. umbra shows strong site-fidelity particularly during the autumn and winter months (Alós and Cabanellas-Reboredo, 2012; Harmelin and Ruitton, 2006), but there is a lack of knowledge on the reproductive movements occurring during the summer. As many other sciaenid species, the brown meagre forms breeding aggregations (Bono et al., 2001; Grau et al., 2009; Ragonese et al., 2002). Locations of aggregations and spawning sites need to be better documented in order to be protected and managed (Harmelin-Vivien et al., 2015). During the spawning period, i.e., May until August (Chauvet, 1991, Grau et al., 2009), *S. umbra* males produce sounds (Parmentier et al., 2017, Picciulin et al., 2012). In other Sciaenids, these types of vocalizations have been shown to serve to aggregate individuals, attract mates to leks (Gilmore, 2002; Parsons et al., 2009), advertise male readiness to spawn, and synchronise gamete release (Connaughton & Taylor 1995, Lobel, 1992). *S. umbra* vocalizations are loud percussive sounds produced by a sonic muscles associated to the males' swim bladder (Dijkgraaf, 1947, Parmentier et al., 2017). These percussive sounds can either be produced as irregular sounds (I-calls), or in long regular stereotyped sequences of calls (R-calls) (Picciulin et al. 2012) emitted for several hours at nighttime (Fig. 1). In Sciaenids these R-calls represent reproductive displays. Their occurrence in high densities forms so-called choruses, which are indicative of spawning aggregations (Guest and J., 1978; Picciulin et al. 2012, Parsons et al., 2009, Luczkovich et al., 2008b; Mok and Gilmore, 1983). Therefore, sound production has the potential to serve 1) as a proxy for *S. umbra*'s reproductive activity, 2) to identify nocturnal individual occurrence as well as breeding areas and 3) to locate and monitor aggregations and spawning sites over extended time periods (Ramcharitar et al., 2006). Despite the potential of PAM to fill some of

the knowledge gaps about the brown meagre's ecology and reproductive behaviour and a few pioneering studies in the Gulf of Venice (Italy) using PAM for spatial monitoring (Bonacito et al., 2002; Colla et al., 2018; Picciulin et al., 2013b), there are no studies reporting the application and validation of this technique by environmental managers. The aims of this study were to underline the efficiency of PAM for the survey and management of *S. umbra* and its potential spawning grounds within and outside MPAs by mapping its acoustic activity 1) along the coastline of three wide-ranging MPAs, 2) at large spatial scales (Northwestern Mediterranean), and 3) over two years in a no-take reserve.

2 METHODS

2.1 Spatial sampling within three MPAs

2.1.1 Study sites

The study sites were three vast MPAs in the Northwestern Mediterranean Sea: The MPA of the Calanques National Park (CNP), the Natural Marine Park of the Gulf of Lion (NMPGL), both in France, and the MPA of Tavolara-Punta Coda Cavallo (TPCC) in Sardinia, Italy (Fig. 2). The CNP is the youngest French national park established in 2012 and situated between Marseille and La Ciotat (Fig. 2). The AMP comprises a 435 km² area that includes seven no-take zones (NTZ, no fishing) and one reinforced protection zone (RPZ) that allows yet regulates artisanal and recreational fishing activity. The NMPGL is also a young MPA, established in 2011 with a vast surface of 4000 km² along 100 km coastline. The NMPGL includes the Natural Marine Reserve of Cerbère-Banyuls (MRCB), which is the oldest marine reserve in France created in 1974, measuring 6.50 km² and consisting of a RPZ where only recreational fishing and diving are allowed and regulated and a fully protected no-take and no-access zone (FPZ) (Fig. 2). Finally, the TPCC MPA in Northeastern Sardinia, Italy, established in 1997 but

effectively managed around 2003-04, comprises 76 km of coastline and a surface area of 153.57 km². Three types of zones with different levels of protection have been established: (1) A zones, corresponding to FPZs (i.e., no-take and no-access); (2) B zones, corresponding to RPZ, where only licensed local artisanal fishing and diving are allowed; (3) C zones, corresponding to general protection zones (PZ), where both artisanal and recreational fishing are authorized as well as diving (Fig. 2).

2.1.2 Acoustic data acquisition and recording protocol

Spatial mapping of *S. umbra* based on their vocalisations has been proposed and successfully adopted in the Gulf of Venice (Italy) (Bonacito et al., 2002; Colla et al., 2018; Picciulin et al., 2013). In this study sound recordings were acquired using two or three simultaneously deployed drifting devices equipped with an EA-SDA14 compact autonomous recorder (RTSys®, France) and a HTI-92-WB (High Tech Inc.®) hydrophone with a flat sensitivity equal to -155 ± 3 dB re 1 V μ Pa⁻¹ between 5 and 50 kHz. Acoustic recordings were sampled continuously at a rate of 78 kHz and a 24-bit resolution. The hydrophone was located between 5 m and 9 m below the sea surface. Positions were defined based on the mean propagation range of *S. umbra* calls estimated using reported source levels (Codarin et al. 2009, Parmentier et al. 2017) and assuming cylindrical spreading loss. Consequently, maximal inter-recording position-distance was set at 600m. Priority and secondary zones for the survey were selected by the MPA's agents based on habitat, protection level and existing knowledge. Deployments were performed using a standardized protocol: Along rocky coastlines, the devices were deployed within maximum 200m from the coast, one after the other at 200 to 600m distance, with water depths ranging between 8 and 20m. After deployment of the drifting recorders at a target positions, the boat moved 200 m away and stopped the engine in order to avoid noise interference. After 7 min of recording, each device was recovered. At each deployment and recovery, GPS time and position

were noted. All drifts were conducted during the summer, from 7 p.m. to 12 p.m., which includes the period of maximal vocalizing activity of the brown meagre ([Parmentier et al. 2017](#); [Picciulin et al. 2013](#)) (Table 1).

2.2 Spatial sampling across the Northwestern Mediterranean basin

Sound recordings from 49 stations of the CALME acoustic network along the French Western Mediterranean coast established by the RMC Water Agency and the CHORUS Institute ([Gervaise et al. 2018](#)) were used; 22 from *Posidonia oceanica* meadows, and 25 from coralligenous reefs, and two from rocky reefs obtained in summer 2016 and 2017 (Table 1). The CALME network was not designed for *S. umbra* monitoring but is dedicated to key habitat and noise surveillance within the scope of EU Water Framework Directive (2000/60/EC) and the Marine Strategy Framework Directive (2008/56/EC). The same recording equipment (hydrophones and autonomous recorders) was deployed as for the drifting devices. The recorders were bottom moored with the hydrophone at 1 m from the seafloor (Fig. 3). Recorders acquired sounds continuously at a 78 kHz sampling rate and 24-bit resolution. For 46 of the 50 recording stations, at each recording date, the recorder was submerged in the afternoon and recovered the next morning for a duration of at least 14 hours. Recordings were made during night-time because the brown meagre, as most temperate fishes, vocalises at night ([Parmentier et al., 2017](#); [Parsons et al., 2016](#); [Picciulin et al., 2012](#)). In four sites, recordings were made over more than 24h.

2.3 Long-term data collection

One long-term recording from the CALME network situated in typical *S. umbra* habitat was used to assess annual trends in acoustic presence. The recording was obtained from the Réserve de Couronne in the Côte Bleue Marine Park (CBMP), near Marseille, France (43° 19' 32", 5° 10' 11"). This reserve was created in 1996 and comprises 210 ha. It is a FPZ, where all kind of fishing activities, mooring and scuba diving are prohibited. An EA-SDA14 compact autonomous recorder (RTSys®, France) and an HTI-92-WB (High Tech Inc.®) with the same characteristics described for the drifting device was used but equipped with an external battery pack for long-term deployments. Data were recorded continuously and year-round in 2017 and 2018. The recorder was placed on a spawning site of seabreams (*Sparus aurata*) and seabass (*Dicentrarchus labrax*) known to regularly host brown meagres. The recording site was situated near an artificial reef and a natural coralligenous reef, next to a patchy *Posidonia oceanica* meadow, at a depth of 25 m. Bottom temperature profiles and trends were obtained from a temperature probe installed nearby by the park's agents since 1998.

2.4 Data processing

All data was first down-sampled to 4kHz. On the one hand, data acquired during the acoustic small-scale spatial surveys within the MPAs and those obtained from the large-scale CALME network were visually inspected using Raven PRO 1.5 (The Cornell Lab of Ornithology; Hanning window, LFFT = 512, overlap: 50%). All recordings from the MPA surveys were inspected for presence/absence of I-calls, R-calls and choruses, as the presence of these different call types may indicate functional differences of sites. I-calls are characterized by variable numbers of pulses with no fixed call repetition rate. In contrast, R calls are composed of 5-6 pulses repeated in long regular stereotyped sequences (Picciulin et al. 2012) (Fig. 1). They are mainly produced between 5 p.m. and 12 p.m. with a peak between 8 p.m. and 10 p.m. and show spatial as well as temporal consistency (at least over 10 years) in their acoustic

features (Picciulin et al. 2012, Parmentier et al. 2017). If possible, based on signal-to-noise differences of simultaneously R-calling individuals, an approximation was made on the number of singers. In the presence of choruses, the abundance of calls is so high that individual call sequences cannot be discriminated any more (Fig. 1).

The large-scale sampling sites across the Northwestern Mediterranean were visually inspected for the presence of R-calls and choruses only, which are representative of courtship and spawning behaviour. On the other hand, long-term data was processed using a custom-made sound detection algorithm based on the rhythmic properties of calls and capable of discerning vocalizing individuals (Le Bot et al., 2015). This rhythmic analysis algorithm is based on time of arrivals of single pulses of a call within a frequency band of interest and uses a complex autocorrelation function with a sliding window to build a time-rhythm representation (Le Bot et al. 2015). The algorithm parameters were set to detect good-quality *S. umbra* calls that are generally temporally stereotyped (i.e., R-calls with mean pulse periods of 240ms; Parmentier et al. 2017) and with a minimum signal-to-noise-ratio of 6dB. Time series of the detected *S. umbra* calls were plotted over time. When available, temperature curves were provided and correlated with the acoustic activity.

3 RESULTS

3.1 Spatial distribution within MPAs

3.1.1 Calanques National Park

The three recording sessions comprised a total of 72 recording stations covering over 40km of coastline and representing a surface of 20.4 km² (considering a detection range of 300 m). 40 recording drifts were performed within NTZs (no-take zones) of the MPA and 32 in areas with

a lower protection regime (RPZ) (Fig. 4). *S. umbra* R- & I-calls were recorded at 27.7% of the stations, both within (55% of call detections) and outside (45%) NTZs. R-calls (reproductive calls) equally occurred at 4 stations inside and 4 outside NTZ, indicating that reproductive activity also takes place in less protected areas. Most I-calls were detected inside NTZs (9 out of 12 stations). No choruses were detected and overall, calling activity was not abundant (only one, maximum two R-calls per stations and a few single I-calls), indicating a small number of vocalizing individuals.

3.1.2 Marine Natural Park of the Gulf of Lion

96 sites were acoustically scanned during the four night-time surveys, allowing to cover the entire rocky coastline of the park (i.e., 30 km). *S. umbra* calls were present at 46 recording stations (i.e., 48% of the stations); 27 were within the marine reserve (MRCB) and 19 outside the reserve along the rocky cost of the marine park (Fig. 4). As the recording effort was greater within the reserve than in the rest of the park, there may be a bias towards the higher number of detections within the reserve. Nevertheless, the results show an effect of type and duration of the protection measures. In fact, the reserve was dense in detections, with *S. umbra* calls present in 77% compared to 33% of the stations in the rest of the park. Furthermore, in the reserve, acoustic reproductive behaviour was intense: 21 (of 27) stations with R-calls, three with choruses, and 50% of all stations with at least two singers. In the rest of the park outside the reserve, R-calls were recorded at 12 stations (66%), I-calls at 6 stations and simultaneous singing was detected at one station only. Overall vocal activity was therefore generally smaller compared to the reserve. From 4 sites recorded both in June and August in the NMPGL, two showed acoustic activity in both months, which may be indicative of site fidelity.

3.1.2 MPA of Tavolara-Punta Coda Cavallo

54 stations were acoustically sampled during three recording sessions covering 20 km coastline: 16 in the full-protection zones (no-take & no-access) A, 24 in the RPZ zones B, and 14 in the less protected zone C (Fig. 4). At almost 70% of all stations, *S. umbra* calls were present. Zonation, i.e., the degree of protection and enforcement level, strongly influenced the observed calling behaviour: All stations in fully protected A zones showed sustained calling activity, including 11 stations with R-calls, one with I-calls and 2 with choruses. 12 out of the 16 stations showed elevated simultaneous singing of multiple individuals (2-4 or choruses). The B zone also showed a strong vocal activity (70% of the stations) with 10 stations in which R-calls were recorded, three in which I-calls occurred, three in which both R-and I-calls were present and one with a chorus. Simultaneous singing activity of at least 2 individuals occurred at 5 stations. The C-zone was the one with the fewest detections of *S. umbra* calls. Out of 14 stations, only 3 presented calling activity; one with I-calls, one with R-calls and one showing chorusing activity. Moreover, the Northern part of the MPA is more exposed to anthropogenic noise of ferries and other boats leaving the harbour of Olbia. Signal-to-noise ratio was overall small in N-Tavolara and Capo Ceraso because of increased background noise due to vessel traffic. Masking of communication signals may therefore be a cause for reduced signal production and/or detection.

3.2 Spatial sampling across the Northwestern Mediterranean basin

S. umbra was also present outside MPAs. On 23 out of 49 stations inspected, R-calls and choruses were detected (Fig. 5). This corresponds to an encounter probability of 47% (Fig 4) despite the fact that the recording stations were not specifically designed for *S. umbra* monitoring. The comparison between sites in seagrass meadows and coralligenous reefs indicates that *S. umbra* is acoustically present at 11 sites within seagrass meadows (50% of the

sites) and at 8 sites on coralligenous reefs (30% of the sites). Although not quantified here, the number of simultaneously singing males was generally low (< 3 individuals), suggesting overall low fish densities. Potential spawning sites indicated by the presence of choruses only occurred at four sites; three in Corsica, and one in Sardinia. Two sites were within and two outside no-take zones.

3.3 Long-term seasonal monitoring within an MPA no-take zone

The annual recordings in the Reserve of Couronne in CBMP highlighted an acoustic activity of *S. umbra* spanning over a period of 5 months, with a strong consistency in sound production from one year to the other. In 2017, the first *S. umbra* calls were present in April but singing started on May 21 and stopped on October 11. In 2018 brown meagres started singing on May 22 and stopped on October 17 (Fig. 6). The singing period was almost identical and comprising part of spring and autumn. Singing activity was most intense during the summer months (June-September), and present 72% of the time (Fig. 6). Singing was also highly positively correlated with water temperature ($\rho = 0.8$) (Fig. 6).

DISCUSSION

This study highlights the efficiency of PAM in studying and following *S. umbra* populations over different spatial and temporal scales for species management. The standardized method adopted to map the acoustic distribution of *S. umbra* using multiple drifting devices allowed to establish maps of the acoustic presence and reproductive activity of almost the totality of the MPAs' rocky coastlines (up to 50km) in only 3-4 evenings. This confirms the efficiency of the method in rapidly assessing *S. umbra* distribution over vast areas (Picciulin et al., 2013). Distribution maps of this kind, obtained with limited human and logistic effort, are difficult to

establish with traditionally used techniques. The results from these nocturnal surveys, showed that in the CNP and the NMPGL (excluding the reserve) *S. umbra* was present in approximately 30% of the recorded sites. In the Reserve of Cerbère-Banyuls (within the NMPGL) and TPCC, the probability of detecting *S. umbra* calls was between 70 and 80%, thus more than two times higher than in the NMPGL and the CNP. These differences may be linked to different management measures and particularly enforcement against anthropogenic pressures (i.e., fishing and diving), to the surveillance programs adopted by the different MPAs, to habitat characteristics, but also to the onset of the protection measures. In fact, the CNP and the NMPGL are young MPAs established approximately 7 years ago, while the MRCB and TPCC have been managed for over 20 years. Moreover, within the TPCC MPA, *S. umbra* calls were present in 100% of the sites within the fully protected zones A, 70% within reinforced B-zones and only 21% in the partially protected C-zones. Together these results indicate an overall effect of “age” of the MPA as well as a reserve-effect given by the level of protection and enforcement. This is consistent with the findings of long-term visual observations in Mediterranean MPAs indicating conspicuous increases in brown meagre abundance in fully protected or no-take zones over the last decade and an overall slow recovery of *S. umbra* populations (Francour, 1991; García-Rubies et al., 2013; Guidetti et al., 2014; Harmelin-Vivien et al., 2008; 2015; Harmelin and Ruitton, 2006). The role of enforced protection is further supported by the fact that almost all (8 out of 11) sites with chorusing activity, which are indicative of potential spawning aggregations, occurred within fully protected zones of older MPAs. Furthermore, in younger MPAs, calling activity was generally lower and the number of simultaneously singing individuals per site was smaller (< 3) compared to older more enforced MPA zones.

Overall, the R-call was the most abundant call type and it occurred within MPAs as well as in non-protected areas of the Northwestern Mediterranean. Outside MPAs, *S. umbra* reproductive calls were present in 47% of the sites, both in *Posidonia oceanica* seagrass meadows (50% of the seagrass sites) and in coralligenous reefs (30% of the coralligenous reefs). However, except for two chorusing sites in Corsica, the number of singers at one site was generally low, possibly indicating low animal densities. Corsica showed the highest number of sites with singing brown meagres (11 out of 23). On the mainland, they were mainly detected in the Eastern part of the region Provence-Alpes-Côte d'Azur (PACA, stations 8-6). In Western PACA only the CBMP reserve recorded brown meagre acoustic activity, while none was observed in the Languedoc-Roussillon region, which was however only represented by three stations (Fig. 5). These differences may be linked to the presence of suitable habitats for the species or fishing and may also be due to the fact that the recording sites were not specifically chosen to monitor the brown meagre. The coast of the gulf of Lion is dominated by sandy substrate with limited rocky and *P. oceanica* habitats. Overall, the *P. oceanica* sites with *S. umbra* detections were characterized by higher percentage of rocky substrate compared to the sites without detections (14% vs 6% rock). This may partly explain the distributions observed and why this species, typically associated to rocky habitats, was present in *P. oceanica* meadows. A more detailed analysis on habitat characteristics may provide insights to elucidate the observed distributions. The acoustic presence of *S. umbra* also outside MPAs, although generally in low densities, might be due to 1) the extended spatial coverage of the CALME network, 2) the presence of habitats suitable for the occurrence of the study species but also 3) the long-term management measures to increase water and habitat quality (EU Water Framework Directive) and 4) the ongoing moratorium in France that bans spearfishing and hook-and-line recreational fishing until December 2023 and thus partially protects *S. umbra* populations (Harmelin-Vivien et al., 2015; Rocklin et al., 2011).

377

378 Overall, these results show that, although strongly linked to habitat characteristics and
 379 protection level, *S. umbra* courtship activity occurs both within and outside MPAs (at least in
 380 France), and that PAM can be used to identify distribution and potential reproductive sites.
 381 Within the scope of spatial planning, PAM can contribute to define protection areas around
 382 functionally relevant zones (Erisman et al., 2012; Luczkovich et al., 2008b). This is particularly
 383 true for sites with chorusing activity that is associated with spawning aggregations, with chorus
 384 intensity being related to fish abundance and biomass in other Sciaenids (Gannon and Gannon,
 385 2010; Rowell et al., 2017). Large aggregations of the brown meagre have been reported in the
 386 Mediterranean Sea (Fiorentino et al., 2001; Ragonese et al., 2002). Surveys combining passive
 387 and active acoustics will help confirming the link between acoustic activity and fish abundance
 388 (Erisman and Rowell, 2017; Mann et al., 2008). Furthermore, in *S. umbra* it remains unknown
 389 if choruses are related to egg production and spawning as in the weakfish (*Cynoscion regalis*),
 390 spotted seatrout (*Cynoscion nebulosus*), red drum (*Sciaenops ocellatus*), silver perch
 391 (*Bairdiella chrysoura*), meagre (*Argyrosomus regius*) or the mullet (*Argyrosomus*
 392 *japonicus*) (Lagardère and Mariani, 2006; Lowerre-Barbieri et al., 2008a; Luczkovich et al.,
 393 1999; Parsons et al., 2006). Ichthyoplankton and adult surveys at chorusing sites are required
 394 to assess the link between sound production and spawning (Luczkovich et al. 2008a). However,
 395 the results of the acoustic activity of *S. umbra* reported here clearly show that chorusing is rare
 396 (11 out of 271 sites) and prevails within fully protected or no-take zones in well-enforced
 397 MPAs. This is particularly relevant and further underlines the importance of marine reserves in
 398 protecting spawning sites and ultimately populations.

399

400 The vocal activity patterns reported here also allow to better describe reproductive movements
 401 of the brown meagre and the displacement of individuals for breeding aggregations that are still

poorly documented (*c.f.*, [Harmelin-Vivien et al. 2015](#)). PAM offers a means to identify the sites of vocal males and help assess the extent of their poorly known spatial movements and home ranges. It also shows its utility to fill some of the knowledge gaps in the nocturnal behaviour of this species ([Harmelin-Vivien et al. 2015](#)). *S. umbra* males in the Northwestern Mediterranean predominantly vocalize during the first half of the night, with a peak about one hour after sunset. This is therefore a time period at which, during the reproductive season, animals are engaged in reproductive behaviours. The diel pattern described here coincides with the one reported in Corsica (France), Sardinia (Italy) and in the Adriatic Sea (Italy) ([Parmentier et al. 2017](#), [Picciulin et al. 2013](#)) and is commonly observed in many sciaenid species ([Fish and Cummings, 1972](#); [Mann and Grothues, 2009](#); [Saucier and Baltz, 1993](#)).

Two-year recordings in the Reserve of CBMP showed that the brown meagre is vocally active from May to October. Seasonal calling activity is common in Sciaenids (e.g., [Connaughton and Taylor, 1994](#); [Montie et al., 2015](#)), but this is the first description of annual calling activity and natural variability in the brown meagre that may help better assess the spawning period of this vulnerable specie ([Monczak et al., 2017](#)). The spawning season of *S. umbra* is suggested to be from May to August, as estimated from histological examination and gonadosomatic index ([Chauvet, 1991](#); [Grau et al., 2009](#)). Here we show that singing activity was sustained (i.e., 72% of occurrence) from late spring to autumn, indicating that the reproductive season of *S. umbra* is likely protracted, and occurring over a five-months period. Additional long-term recordings at different potential spawning sites will allow to confirm this extended reproductive season. The duration of the singing activity was equivalent in both years, starting on May 21 in 2017 and May 22 in 2018 and ending on October 11 in 2017 and October 17 in 2018. This relatively abrupt onset of calling activity suggests the presence of an environmental trigger. Temperature has been shown to initiate reproductive behaviour and chorusing in other sciaenid species

(Mann and Grothues, 2009; Monczak et al., 2017; Rice et al., 2016). In this study, there was a strong positive correlation between water temperature and calling activity ($\rho = 0.8$), supporting the temperature-related hypothesis. Furthermore, although singing was sustained throughout the summer, it showed fluctuations in calling intensity (i.e., number of calls per hour). Peaks of vocal activity occurred in June, July, August and September and coincided with peaks in water temperature at the recording site. Similar temperature-related fluctuations in calling during the spawning season have been described in other Sciaenids (Connaughton and Taylor, 1994; Monczak et al., 2017; Montie et al., 2015) and have been related to seasonal cycles in the sonic muscle of males (Connaughton and Taylor, 1994). The smaller pulse-period observed during intense calling activity may be related to faster muscle twitch and contraction velocity at higher temperatures (Connaughton et al., 2002, Bolgan et al. 2020). An alternative explanation for the observed variations in calling intensity may be the influence of the lunar cycle (McCauley, 2012, Monczak et al., 2017). However, in this study intervals between peaks varied between 14 and 36 days (mean 25 days), thus indicating no clear lunar cycle pattern. In other sciaenid species, changes in calling behaviour have been related to spawning events (Lowerre-Barbieri et al., 2008; Mok and Gilmore, 1983; Montie et al., 2017). Although this remains to be elucidated for the brown meagre, the species is a multiple-spawning fish and within a population, females do not show synchronous ovarian maturity, which results in an extended spawning season (Grau et al. 2009).

CONCLUSIONS

This study emphasizes the utility of PAM for the survey and conservation of exploited, vulnerable Mediterranean fish species. It represents a complementary, non-invasive, low-cost, standardized, replicable and efficient monitoring technique for environmental managers (Luczkovich et al., 2008a; Mann et al., 2008; Rountree et al., 2011). Here we focussed on the

brown meagre, but the method can be extended to other vocal species (e.g., groupers). Harmelin-Vivien and co-authors (2015) underlined the need to combine extensive observations along the coast and regular long-term monitoring of favourable sites to document spatial distribution and the effect of increased protection of *S. umbra*. Here we show that PAM will allow to fill some of the knowledge gaps on species distribution, behaviour and reproduction needed for the conservation of the species using a combination of spatial and long-term acoustic recordings. Information of this kind is essential for the conservation of exploited or vulnerable fish species (Erisman et al., 2017).

The importance of recurrent monitoring surveys of threatened species inside and outside MPAs is critical to estimate population trends. Repeated standardized evening surveys over the season and across years will also allow, with limited human and logistic effort, to study site-fidelity and identify preferential sites of brown meagre populations over wide areas and help decision-makers to adopt protection acts and establish management zones (Luczkovich et al., 2008b). Here we showed that PAM increases our knowledge on the distribution and reproductive activity of this vulnerable species in rarely surveyed areas (i.e. outside MPAs) and over large geographical scales. PAM is also effective in emphasizing effects linked to management actions (e.g., no-take zones). Combining the obtained findings with existing visual observations, abundance estimations (e.g., using underwater visual census data or active acoustics), individual telemetry tracking data (Parsons et al., 2009; Taylor et al., 2006) as well as ichthyoplankton surveys on chorusing sites will provide critical information on the reproductive behaviour of the species, their distribution and population trends and allow to further consolidate the use of PAM for monitoring and managing this vulnerable species. Furthermore, simultaneous annual recordings on chorusing or known aggregation sites will help identify spawning events (e.g; Luczkovich et al., 2008b; Rice et al. 2015), trends in fish abundance via

the calling activity (Erisman and Rowell, 2017; Rowell et al., 2017), and shifts in reproductive timelines associated with climate change (Monczak et al. 2017). Altogether, the findings reported here highlight the use of PAM as a means to support managers in their efforts to monitor, recover, and ultimately protect key ecological, commercial and emblematic species such as the Mediterranean brown meagre.

CRediT authorship contribution statement

Lucia Di Iorio: Conceptualization, Methodology, Formal analysis, Writing – initial, review & editing, Investigation, Visualization, Supervision, Project administration. Patrick Bonhomme: Conceptualization, Investigation, Resources, Visualization, Funding acquisition. Noémie Michez: Investigation, Resources, Writing - Review & Editing. Bruno Ferrari: Investigation, Resources. Alexandra Gigou: Conceptualization, Funding acquisition. Pieraugusto Panzalis: Investigation, Resources, Elena Desiderà: Investigation, Writing - Review & Editing. Augusto Navone: Resources, Pierre Boissery: Resources, Writing - Review & Editing, Funding acquisition. Julie Lossent: Methodology, Investigation. Benjamin Cadville: Investigation, Resources. Marie Bravo-Monin: Investigation, Resources. Eric Charbonnel: Resources, Writing - Review & Editing, Cédric Gervaise: Software, Methodology, Formal analysis, Conceptualization, Writing – review & editing, Investigation, Supervision, Project administration.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

500

501 **ACKNOWLEDGEMENTS**

502 The authors would like to thank the teams of the four MPAs as well as F. Cadène, V. Hartmann
 503 the team of the Marine Reserve of Cerbère-Banyuls for their technical support and their
 504 implication in the survey. Many thanks also to L. Bramanti, CNRS Observatoire Océanologique
 505 de Banyuls and the team of Andromède Océanologie for their support in the field. Thanks to J.
 506 Decelle for constructive comments on a previous version of the manuscript. This work has been
 507 possible thanks of the financial support of the Calanques National Park, the RMC Water
 508 Agency (convention 2016 1858), the French Agency for Biodiversity (AFB/2019 140),
 509 CHORUS Institute, and by the *Fonds Français pour l'Environnement Mondial* (FFEM) and the
 510 *Fondation du Prince Albert II de Monaco* via the MedPAN Calls for Small Projects 2019.

511

512

513 **REFERENCES**

- 514 Abdul Malak, D., Livingstone, S., Pollard, D., Polidoro, B., Cuttelod, A., Bariche, M.,
 515 Bilecenoglu, M., KE, C., Collette, B., Francour, P., Goren, M., Kara, H., Massutí, E.,
 516 Papaconstantinou, C., Tunesi, L., 2011. Overview of the Conservation Status of the
 517 Marine Fishes of the Mediterranean Sea.
- 518 Alós, J., Cabanellas-Reboredo, M., 2012. Experimental acoustic telemetry experiment reveals
 519 strong site fidelity during the sexual resting period of wild brown meagre, *Sciaena*
 520 *umbra*. J. Appl. Ichthyol. 28, 606–611. [https://doi.org/10.1111/j.1439-](https://doi.org/10.1111/j.1439-0426.2012.01955.x)
 521 [0426.2012.01955.x](https://doi.org/10.1111/j.1439-0426.2012.01955.x)
- 522 Amorim, M.C.P., 2006. Diversity of Sound Production in Fish. Diversity 1, 71–105.

- 523 Bizsel, C., Yokes, B., Pollard, D., Kara, M.H., Bariche, M., Quignard, J.P., 2011. *Sciaena*
524 *umbra*. IUCN Red List Threat. Species 2011.
- 525 Bonacito, C., Costantini, M., Picciulin, M., Ferrero, E.A., Hawkins, A.D., 2002. Passive
526 Hydrophone Census of *Sciaena Umbra* (Sciaenidae) in the Gulf of Trieste (Northern
527 Adriatic Sea, Italy). Bioacoustics 12, 292–294.
528 <https://doi.org/10.1080/09524622.2002.9753726>
- 529 Bono, G., Fiorentino, F., Camilleri, M., Gancitano, S., Giusto, G., Ragonese, S., Rizzo, P.,
530 Rosso, B., 2001. On a spawning aggregation of the brown meagre *Sciaena umbra* L.,
531 1758 (Sciaenidae, Osteichthyes) in the Maltese waters (Sicilian Channel-Central
532 Mediterranean). Rapp. Comm. Int. Mer Médit. 36, 266.
- 533 Chao, L., 2015. *Sciaena umbra*. IUCN Red List Threat. Species.
534 [https://doi.org/http://dx.doi.org/10.2305/IUCN.UK.2015-](https://doi.org/http://dx.doi.org/10.2305/IUCN.UK.2015-4.RLTS.T198707A83232286.en)
535 [4.RLTS.T198707A83232286.en](https://doi.org/http://dx.doi.org/10.2305/IUCN.UK.2015-4.RLTS.T198707A83232286.en).
- 536 Chauvet, C., 1991. Le corb ou brown meagre (*Sciaena umbra* - Linnaeus, 1758) quelques
537 éléments de sa biologie, in: Bouderesque, C.F., Avon, M., Gravez, V. (Eds.), Es Espèces
538 Marines à Protéger En Méditerranée. GIS Posidonie, France, pp. 229–235.
- 539 Colla, S., Pranovi, F., Fiorin, R., Malavasi, S., Picciulin, M., 2018. Using passive acoustics to
540 assess habitat selection by the brown meagre *Sciaena umbra* in a northern Adriatic Sea
541 mussel farm. J. Fish Biol. 92, 1627–1634. <https://doi.org/10.1111/jfb.13589>
- 542 Connaughton, M.A., Fine, M.L., Taylor, M.H., 2002. Use of sounds for localisation of
543 spawning weakfish in Delaware Bay (USA) and effects of fish size, temperature and
544 season on sound parameters. Bioacoustics 12, 294–296.
545 <https://doi.org/10.1080/09524622.2002.9753727>

- 546 Connaughton, M.A., Taylor, M.H., 1994. Seasonal cycles in the sonic muscles of the
547 weakfish , *Cynoscion regalis*. U.S. Fish.Bull 92, 697–703.
- 548 Desiderà, E., Guidetti, P., Panzalis, P., Navone, A., Valentini-Poirrier, C.-A., Boissery, P.,
549 Gervaise, C., Di Iorio, L., 2019. Acoustic fish communities: sound diversity of rocky
550 habitats reflects fish species diversity and beyond? Mar. Ecol. Prog. Ser. 608, 183–197.
551 <https://doi.org/10.3354/meps12812>
- 552 Dijkgraaf, S., 1947. Ein Töne erzeugender Fisch im Neapler Aquarium. Experientia 3, 493–
553 494. <https://doi.org/10.1007/BF02154563>
- 554 Erisman, B., Aburto-Oropeza, O., Gonzalez-Abraham, C., Mascareñas-Osorio, I., Moreno-
555 Báez, M., Hastings, P.A., 2012. Spatio-temporal dynamics of a fish spawning
556 aggregation and its fishery in the Gulf of California. Sci. Rep. 2, 284.
557 <https://doi.org/10.1038/srep00284>
- 558 Erisman, B., Heyman, W., Kobara, S., Ezer, T., Pittman, S., Aburto-Oropeza, O., Nemeth,
559 R.S., 2017. Fish spawning aggregations: where well-placed management actions can
560 yield big benefits for fisheries and conservation. Fish Fish. 18.
561 <https://doi.org/10.1111/faf.12132>
- 562 Erisman, Brad E, Rowell, T.J., 2017. A sound worth saving: acoustic characteristics of a
563 massive fish spawning aggregation. Biol. Lett. 13, 20170656.
564 <https://doi.org/10.1098/rsbl.2017.0656>
- 565 Erisman, Brad E., Rowell, T.J., 2017. A sound worth saving: acoustic characteristics of a
566 massive fish spawning aggregation. Biol. Lett. 13, 20170656.
567 <https://doi.org/10.1098/rsbl.2017.0656>
- 568 Fiorentino, F., Camilleri, M., Bono, G., Gancitano, S., Giusto, G.B., Ragonese, S., Rizzo, P.,

569 Rosso, B., 2001. On a spawning aggregation of the brown meagre *Sciaena umbra* L.
570 1758 (Sciaenidae, Osteichthyes) in the Maltese waters (Sicilian Channel—Central
571 Mediterranean). Rapp. Comm. Int. Mer Médit. 36, 266.

572 Fish, J.F., Cummings, W.C., 1972. A 50-dB Increase in Sustained Ambient Noise from Fish
573 (*Cynoscion xanthulus*). J. Acoust. Soc. Am. 52, 1266–1270.
574 <https://doi.org/10.1121/1.1913243>

575 Francour, P., 1991. The effect of protection level on a coastal fish community at Scandola,
576 Corsica. Rev. d'écologie 46, 65–81.

577 Gannon, D.P., 2008. Passive Acoustic Techniques in Fisheries Science: A Review and
578 Prospectus. Trans. Am. Fish. Soc. 137, 638–656. <https://doi.org/10.1577/T04-142.1>

579 Gannon, D.P., Gannon, J.G., 2010. Assessing trends in the density of Atlantic croaker
580 (*Micropogonias undulatus*): A comparison of passive acoustic and trawl methods. Fish.
581 Bull. 108, 106–116.

582 García-Rubies, A., Hereu, B., Zabala, M., 2013. Long-Term Recovery Patterns and Limited
583 Spillover of Large Predatory Fish in a Mediterranean MPA. PLoS One 8, e73922.
584 <https://doi.org/10.1371/journal.pone.0073922>

585 Gilmore, R.G., 2002. Passive acoustic transects: mating calls and spawning ecology in East
586 Florida Sciaenids, in: Massachusetts Institute of Technology (Ed.), An International
587 Workshop on the Applications of Passive Acoustics in Fisheries. Cambridge, MA, pp.
588 33–42.

589 Grau, A., Linde, M., Grau, A.M., 2009a. Reproductive biology of the vulnerable species
590 *Sciaena umbra* Linnaeus, 1758 (Pisces: Sciaenidae). Sci. Mar. 73.
591 <https://doi.org/10.3989/scimar.2009.73n1067>

- 592 Grau, A., Linde, M., Grau, A.M., 2009b. Reproductive biology of the vulnerable species
593 *Sciaena umbra* Linnaeus, 1758 (Pisces: Sciaenidae). *Sci. Mar.* 73, 67–81.
594 <https://doi.org/10.3989/scimar.2009.73n1067>
- 595 Guest, W.C., J., L., 1978. A note on courtship behavior and sound production of red drum.
596 *Copeia* 2, 337–338.
- 597 Guidetti, P., Baiata, P., Ballesteros, E., Di Franco, A., Hereu, B., Macpherson, E., Micheli, F.,
598 Pais, A., Panzalis, P., Rosenberg, A.A., Zabala, M., Sala, E., 2014a. Large-scale
599 assessment of mediterranean marine protected areas effects on fish assemblages. *PLoS*
600 *One* 9. <https://doi.org/10.1371/journal.pone.0091841>
- 601 Guidetti, P., Baiata, P., Ballesteros, E., Di Franco, A., Hereu, B., Macpherson, E., Micheli, F.,
602 Pais, A., Panzalis, P., Rosenberg, A.A., Zabala, M., Sala, E., 2014b. Large-Scale
603 Assessment of Mediterranean Marine Protected Areas Effects on Fish Assemblages.
604 *PLoS One* 9, e91841. <https://doi.org/10.1371/journal.pone.0091841>
- 605 Harmelin-Vivien, M., Cottalorda, J.-M., Dominici, J.-M., Harmelin, J.-G., Le Diréach, L.,
606 Ruitton, S., 2015. Effects of reserve protection level on the vulnerable fish species
607 *Sciaena umbra* and implications for fishing management and policy. *Glob. Ecol.*
608 *Conserv.* 3, 279–287. <https://doi.org/10.1016/J.GECCO.2014.12.005>
- 609 Harmelin-Vivien, M., Le Diréach, L., Bayle-Sempere, J., Charbonnel, E., García-Charton,
610 J.A., Ody, D., Pérez-Ruzafa, A., Reñones, O., Sánchez-Jerez, P., Valle, C., 2008.
611 Gradients of abundance and biomass across reserve boundaries in six Mediterranean
612 marine protected areas: Evidence of fish spillover? *Biol. Conserv.* 141, 1829–1839.
613 <https://doi.org/10.1016/j.biocon.2008.04.029>
- 614 Harmelin, J.-G., Bachet, F., Garcia, F., 1995. Mediterranean Marine Reserves: Fish Indices as
615 Tests of Protection Efficiency. *Mar. Ecol.* 16, 233–250. <https://doi.org/10.1111/j.1439->

0485.1995.tb00408.x

Harmelin, J., 2013. Le mérou brun et le corb : deux Grands Témoins de 50 ans de protection du milieu marin dans le Parc national de Port-Cros (France, Méditerranée). *Sci. Rep. Port-Cros Natl. Park. Fr.* 27, 263–277.

Harmelin, J.G., Ruitton, S., 2006. La population de corb (*Sciaena umbra* : Pisces) du parc national de Port-Cros (France), état en 2005 et évolution depuis 1990 : un indicateur halieutique et biogéographique pertinent. *Sci. Rep. Port-Cros Natl. Park* 22 pp.

Ladich, F., 2015. *Sound communication in fishes*. Springer.

Lagardère, J.P., Mariani, A., 2006. Spawning sounds in meagre *Argyrosomus regius* recorded in the Gironde estuary, France. *J. Fish Biol.* 69, 1697–1708.
<https://doi.org/10.1111/j.1095-8649.2006.01237.x>

Le Bot, O., Mars, J.I., Gervaise, C., Simard, Y., 2015. Rhythmic analysis for click train detection and source separation with examples on beluga whales. *Appl. Acoust.* 95, 37–49. <https://doi.org/10.1016/j.apacoust.2015.02.005>

Letourneur, Y., Ruitton, S., Sartoretto, S., 2003. Environmental and benthic habitat factors structuring the spatial distribution of a summer infralittoral fish assemblage in the north-western Mediterranean Sea. *J. Mar. Biol. Assoc. United Kingdom* 83, 193–204.
<https://doi.org/10.1017/S0025315403006970h>

Lloret, J., Zaragoza, N., Caballero, D., Font, T., Casadevall, M., Riera, V., 2008. Spearfishing pressure on fish communities in rocky coastal habitats in a Mediterranean marine protected area. *Fish. Res.* 94, 84–91. <https://doi.org/10.1016/J.FISHRES.2008.07.002>

Lobel, P.S., 2002. Diversity of fish spawning sounds and the application of passive acoustic monitoring. *Bioacoustics* 12, 286–289. <https://doi.org/10.1080/09524622.2002.9753724>

639 Lowerre-Barbieri, S.K., Barbieri, L.R., Flanders, J.R., Woodward, A.G., Cotton, C.F.,
640 Knowlton, M.K., 2008. Use of Passive Acoustics to Determine Red Drum Spawning in
641 Georgia Waters. *Trans. Am. Fish. Soc.* 137, 562–575. <https://doi.org/10.1577/T04-226.1>

642 Luczkovich, J.J., Mann, D. a., Rountree, R. a., 2008a. Passive Acoustics as a Tool in Fisheries
643 Science. *Trans. Am. Fish. Soc.* 137, 533–541. <https://doi.org/10.1577/T06-258.1>

644 Luczkovich, J.J., Pullinger, R.C., Johnson, S.E., Sprague, M.W., 2008b. Identifying Sciaenid
645 Critical Spawning Habitats by the Use of Passive Acoustics. *Trans. Am. Fish. Soc.* 137,
646 576–605. <https://doi.org/10.1577/T05-290.1>

647 Luczkovich, J.J., Pullinger, R.C., Johnson, S.E., Sprague, M.W., 2008c. Identifying Sciaenid
648 Critical Spawning Habitats by the Use of Passive Acoustics. *Trans. Am. Fish. Soc.* 137,
649 576–605. <https://doi.org/10.1577/t05-290.1>

650 Luczkovich, J.J., Sprague, M.W., Johnson, S.E., Pullinger, R.C. 1999. Delimiting spawning
651 areas of weakfish *Cynoscion regalis* (famili Sciaenidae) in Pamlico sound, North
652 Carolina using passive hydroacoustic surveys. *Bioacoustics* 10, 143–160.
653 <https://doi.org/10.1080/09524622.1999.9753427>

654 Mann, D., Grothues, T., 2009. Short-term upwelling events modulate fish sound production at
655 a mid-Atlantic Ocean observatory. *Mar. Ecol. Prog. Ser.* 375, 65–71.
656 <https://doi.org/10.3354/meps07720>

657 Mann, D.A., Hawkins, A.D., Jech, J.M., 2008. Active and Passive Acoustics to Locate and
658 Study Fish, in: *Fish Bioacoustics*. Springer New York, pp. 279–309.
659 https://doi.org/10.1007/978-0-387-73029-5_9

660 McCauley, R.D., 2012. Fish choruses from the Kimberley, seasonal and lunar links as
661 determined by long term sea noise monitoring. *Aust. Acoust. Soc. Conf.* 2012, Acoust.

2012 Acoust. Dev. Environ. 441–446.

Mok, H.K., Gilmore, R.G., 1983. Analysis of sound production in estuarine aggregations of *Pogonias cromis*, *Bairdiella chrysoura*, and *cynoscion nebulosus* (Sciaenidae) [Florida]. Bull. Inst. Zool. Acad. Sin.

Monczak, A., Berry, A., Kehrer, C., Montie, E., 2017. Long-term acoustic monitoring of fish calling provides baseline estimates of reproductive timelines in the May River estuary, southeastern USA. Mar. Ecol. Prog. Ser. 581, 1–19. <https://doi.org/10.3354/meps12322>

Montie, E.W., Hoover, M., Kehrer, C., Yost, J., Brenkert, K., O'Donnell, T., Denson, M.R., 2017. Acoustic monitoring indicates a correlation between calling and spawning in captive spotted seatrout (*Cynoscion nebulosus*). PeerJ 5. <https://doi.org/10.7717/peerj.2944>

Montie, E.W., Vega, S., Powell, M., 2015. Seasonal and Spatial Patterns of Fish Sound Production in the May River, South Carolina. Trans. Am. Fish. Soc. 144, 705–716. <https://doi.org/10.1080/00028487.2015.1037014>

Morales-Nin, B., Moranta, J., García, C., Tugores, M.P., Grau, A.M., Riera, F., Cerdà, M., 2005. The recreational fishery off Majorca Island (western Mediterranean): some implications for coastal resource management. ICES J. Mar. Sci. 62, 727–739. <https://doi.org/10.1016/j.icesjms.2005.01.022>

Mouillot, D., Culioli, J.-M., Chi, T. Do, 2002. Indicator species analysis as a test of non-random distribution of species in the context of marine protected areas. Environ. Conserv. 29, 385–390. <https://doi.org/10.1017/S0376892902000267>

Parmentier, E., Di Iorio, L., Picciulin, M., Malavasi, S., Lagardère, J.P., Bertucci, F., 2018. Consistency of spatiotemporal sound features supports the use of passive acoustics for

long-term monitoring. *Anim. Conserv.* 21, 211–220. <https://doi.org/10.1111/acv.12362>

Parsons, M.J., McCauley, R.D., Mackie, M.C., Siwabessy, P., Duncan, A.J., 2009.

Localization of individual mulloay (*Argyrosomus japonicus*) within a spawning aggregation and their behaviour throughout a diel spawning period. *ICES J. Mar. Sci.* 66, 1007–1014. <https://doi.org/10.1093/icesjms/fsp016>

Parsons, M.J.G., Mccauley, R.D., Mackie, M., 2006. Spawning sounds of the mulloay (*Argyrosomus japonicus*). *Proc. Acoust.* 499–505.

Parsons, M.J.G., Salgado Kent, C.P., Recalde-Salas, A., McCauley, R.D., 2016. Fish choruses off Port Hedland, Western Australia. *Bioacoustics* 1–18. <https://doi.org/10.1080/09524622.2016.1227940>

Picciulin, M., Bolgan, M., Codarin, A., Fiorin, R., Zucchetta, M., Malavasi, S., 2013. Passive acoustic monitoring of *Sciaena umbra* on rocky habitats in the Venetian littoral zone. *Fish. Res.* 145, 76–81. <https://doi.org/10.1016/j.fishres.2013.02.008>

Picciulin, M., Calcagno, G., Sebastianutto, L., Bonacito, C., Codarin, A., Costantini, M., Ferrero, E. a., 2012. Diagnostics of nocturnal calls of *Sciena umbra* (L., fam. Sciaenidae) in a nearshore Mediterranean marine reserve. *Bioacoustics* 22, 1–12. <https://doi.org/10.1080/09524622.2012.727277>

Picciulin, M., Kéver, L., Parmentier, E., Bolgan, M., 2018. Listening to the unseen: Passive acoustic monitoring reveals the presence of a cryptic fish species. *Aquat. Conserv. Mar. Freshw. Ecosyst.* <https://doi.org/10.1002/aqc.2973>

Ragonese, S., Camilleri, M., Gancitano, S., Rizzo, P., Bono, G., Fiorentino, F., 2002.

Evaluating age at sexual maturity in *Sciaena umbra* L. 1758 (Osteichthyes, Sciaenidae) on the basis of otolith microstructure 9, 789–791.

- 708 Ramcharitar, J., Gannon, D.P., Popper, A.N., 2006. Bioacoustics of Fishes of the Family
709 Sciaenidae (Croakers and Drums). Trans. Am. Fish. Soc. 135, 1409–1431.
710 <https://doi.org/10.1577/t05-207.1>
- 711 Rice, A.N., Morano, J.L., Hodge, K.B., Muirhead, C.A., 2016. Spatial and temporal patterns
712 of toadfish and black drum chorusing activity in the South Atlantic Bight. Environ. Biol.
713 Fishes. <https://doi.org/10.1007/s10641-016-0511-z>
- 714 Rocklin, D., Tomasini, J.-A., Culioli, J.-M., Pelletier, D., Mouillot, D., 2011. Spearfishing
715 Regulation Benefits Artisanal Fisheries: The ReGS Indicator and Its Application to a
716 Multiple-Use Mediterranean Marine Protected Area. PLoS One 6, e23820.
717 <https://doi.org/10.1371/journal.pone.0023820>
- 718 Rountree, R.A., Gilmore, R.G., Goudey, C.A., Hawkins, A.D., Luczkovich, J.J., Mann, D.A.,
719 2011. Listening to Fish. Chang. Publ. Wiley. <https://doi.org/10.1577/1548->
720 [8446\(2006\)31\[433:LTF\]2.0.CO;2](https://doi.org/10.1577/1548-8446(2006)31[433:LTF]2.0.CO;2)
- 721 Rowell, T.J., Demer, D.A., Aburto-Oropeza, O., Cota-Nieto, J.J., Hyde, J.R., Erisman, B.E.,
722 2017. Estimating fish abundance at spawning aggregations from courtship sound levels.
723 Sci. Rep. 7, 3340. <https://doi.org/10.1038/s41598-017-03383-8>
- 724 Saucier, M.H., Baltz, D.M., 1993. Spawning site selection by spotted seatrout, *Cynoscion*
725 *nebulosus*, and black drum, *Pogonias cromis*, in Louisiana. Environ. Biol. Fishes 36,
726 257–272. <https://doi.org/10.1007/BF00001722>
- 727 Taylor, M., Laffan, S., Fielder, S., Suthers, I., 2006. Key habitat and home range of mullet
728 *Argyrosomus japonicus* in a south-east Australian estuary: finding the estuarine niche to
729 optimise stocking. Mar. Ecol. Prog. Ser. 328, 237–247.
730 <https://doi.org/10.3354/meps328237>

TABLES

Table 1. Summary table of the different survey types and the recording sites. For precautionary reasons with regard to illegal fishing, the coordinates of the positions are not provided.

ID	Site	Habitat	Survey	Survey dates	Year	<i>S. umbra</i>	Recorders	Recorder support	Recording stations
	CNP	rock	MPA	26-28/06/2018	2018	1	3	floating device	72
	NMPGL	rock	MPA	24-27/06 & 29/08/2019	2019	1	3	floating device	96
	TPCC	rock	MPA	22-24/07/2019	2019	1	2	floating device	54
1	Banuyls	coralligenous	NW Med Sea	2016/06/21	2016	0	1	bottom mooring	1
1	Banuyls	seagrass	NW Med Sea	2016/06/21	2016	1	1	bottom mooring	1
2	Agde	coralligenous	NW Med Sea	2016/06/20	2016	1	1	bottom mooring	1
3	Grand Travers	coralligenous	NW Med Sea	2016/06/17	2016	0	1	bottom mooring	1
4	Ciotat	seagrass	NW Med Sea	2016/06/02	2016	0	1	bottom mooring	1
4	Ciotat	coralligenous	NW Med Sea	2016/06/02	2016	1	1	bottom mooring	1
5	Sicie	seagrass	NW Med Sea	2016/06/03	2016	0	1	bottom mooring	1
5	Sicie	coralligenous	NW Med Sea	2016/06/03	2016	1	1	bottom mooring	1
6	Giens	coralligenous	NW Med Sea	2016/06/04	2016	1	1	bottom mooring	1
7	Ribaud	seagrass	NW Med Sea	2016/06/04	2016	1	1	bottom mooring	1
8	Bormes	coralligenous	NW Med Sea	2016/06/05	2016	0	1	bottom mooring	1
8	Bormes	seagrass	NW Med Sea	2016/06/05	2016	1	1	bottom mooring	1
9	Cap Lardier	coralligenous	NW Med Sea	2016/06/06	2016	0	1	bottom mooring	1
9	Cap Lardier	seagrass	NW Med Sea	2016/06/06	2016	1	1	bottom mooring	1
10	Bonneau	seagrass	NW Med Sea	2016/06/08	2016	1	1	bottom mooring	1
10	Bonneau	coralligenous	NW Med Sea	2016/06/08	2016	1	1	bottom mooring	1
11	Cap Roux	seagrass	NW Med Sea	2016/06/09	2016	0	1	bottom mooring	1
11	Cap Roux	coralligenous	NW Med Sea	2016/06/09	2016	0	1	bottom mooring	1
12	Golfe Juan	seagrass	NW Med Sea	2016/06/10	2016	0	1	bottom mooring	1
12	Golfe Juan	coralligenous	NW Med Sea	2016/06/10	2016	0	1	bottom mooring	1
13	Bacon	coralligenous	NW Med Sea	2016/06/11	2016	0	1	bottom mooring	1
13	Bacon	seagrass	NW Med Sea	2016/06/11	2016	0	1	bottom mooring	1
14	Nice	coralligenous	NW Med Sea	2016/06/12	2016	0	1	bottom mooring	1
14	Nice	seagrass	NW Med Sea	2016/06/13	2016	1	1	bottom mooring	1
15	Eze	coralligenous	NW Med Sea	2016/06/13	2016	0	1	bottom mooring	1
16	Cap Martin	coralligenous	NW Med Sea	2016/06/15	2016	0	1	bottom mooring	1
17	Maccinagio	seagrass	NW Med Sea	2017/06/02	2017	1	1	bottom mooring	1
17	Maccinagio	coralligenous	NW Med Sea	2017/06/02	2017	1	1	bottom mooring	1
18	Bastia	coralligenous	NW Med Sea	2017/05/31	2017	0	1	bottom mooring	1
18	Bastia	seagrass	NW Med Sea	2017/05/31	2017	0	1	bottom mooring	1
19	Tarco	coralligenous	NW Med Sea	2017/05/30	2017	0	1	bottom mooring	1
19	Tarco	seagrass	NW Med Sea	2017/05/30	2017	1	1	bottom mooring	1
20	Rondinara	coralligenous	NW Med Sea	2017/05/29	2017	0	1	bottom mooring	1
20	Rondinara	seagrass	NW Med Sea	2017/05/29	2017	1	1	bottom mooring	1
21	Murtoli	seagrass	NW Med Sea	2017/05/27	2017	1	1	bottom mooring	1
22	Parata	seagrass	NW Med Sea	2017/06/12	2017	1	1	bottom mooring	1
22	Parata	coralligenous	NW Med Sea	2017/06/12	2017	1	1	bottom mooring	1
23	Cappo Rossu	seagrass	NW Med Sea	2017/06/10	2017	1	1	bottom mooring	1
23	Cappo Rossu	coralligenous	NW Med Sea	2017/06/10	2017	0	1	bottom mooring	1
24	Focolara	coralligenous	NW Med Sea	2017/06/09	2017	0	1	bottom mooring	1
24	Focolara	seagrass	NW Med Sea	2017/06/09	2017	1	1	bottom mooring	1
25	Calvi	seagrass	NW Med Sea	2017/06/05	2017	1	1	bottom mooring	1
25	Calvi	coralligenous	NW Med Sea	2017/06/05	2017	1	1	bottom mooring	1
26	Agriates	coralligenous	NW Med Sea	2017/06/03	2017	0	1	bottom mooring	1
26	Agriates	seagrass	NW Med Sea	2017/06/03	2017	1	1	bottom mooring	1
CBMP	Côte Bleue	seagrass	NW Med Sea	2016/05/30	2016	0	1	bottom mooring	1
27	Cala Gonone	rock	NW Med Sea	several months	2017	1	1	bottom mooring	1
CBMP	Côte Bleue	rock	NW Med Sea	yearround	2017/2018	1	1	bottom mooring	1
TPCC	Tavolara	coralligenous	NW Med Sea	several months	2017	1	1	bottom mooring	1

739 **FIGURES**

741 **Figure 1.** Oscillograms (top) and spectrograms (bottom) of *S. umbra* call categories. a) Irregular
742 I-calls, b) series of stereotyped R-calls, and c) choruses, as described by Picciulin et al. 2012.

743

Figure 2. Maps of the MPAs where drifting surveys were conducted. Top left panel: France and Italy with the locations of the four MPAs. Other panels: Maps of the three MPAs in which *S. umbra* distribution was assessed using drifting devices. The different protection levels are indicated by different colours described in the legend: FPZ = Full Protection Zone (=Reserve = A zone), NTZ = No-Take-Zone (only in PNC), RPZ = Reinforced Protection Zone (= B zone), PZ = Protection Zone (= C zone). CNP: Calanques National Park, NMPGL: Natural Marine Park of the Gulf of Lion with the Marine Reserve of Cerbère-Banyuls (dark grey), TPCC: MPA of Tavolara-Punta Coda Cavallo.

Figure 3. Map of the 49 recording sites of the large-scale monitoring program CALME and the long-term recording site in the Côte Bleue Marine Park (CBMP). Dark grey circles indicate coralligenous reef recording sites, bright grey circle indicate rocky long-term recording stations, white circles are seagrass meadow recording sites. The picture illustrates the mooring system with the recording device. The locations corresponding to the numbers are listed in table 1. *PACA*: Provence Alpes Côte d’Azur region, *LR*: Languedoc-Roussillon region.

Figure 4. Maps of the drifting surveys and *S. umbra* detections within the MPAs. Top left panel: France and Italy with the locations of the four MPAs. Other panels: Maps of the three MPAs illustrating all the recording positions (white circles). Zones of *S. umbra* call detections are highlighted in red. Because of management and precautionary reasons with regard to illegal fishing, more detailed detection positions are not shown here. The different protection levels are indicated by different colours described in the legend: FPZ = Full Protection Zone (=Reserve = A zone), NTZ = No-Take-Zone (only in PNC), RPZ = Reinforced Protection Zone (= B zone), PZ = Protection Zone (= C zone). CNP: Calanques National Park, NMPGL: Natural Marine Park of the Gulf of Lion, TPCC: MPA of Tavolara-Punta Coda Cavallo.

Figure 5. Map of the *S. umbra* detections within the acoustic monitoring network CALME. Green circles indicate *S. umbra* acoustic presence in the seagrass meadow and long-term recording sites, red circles indicate in the coralligenous reef sites. Dark grey circles indicate coralligenous reef recording sites without detections, and white circles are seagrass meadow recording sites without detections. CBMP : Côte Bleue Marine Park. PACA: Provence Alpes Côte d'Azur region, LR: Languedoc-Roussillon region.

Figure 6. Biannual series of *S. umbra* calls in the Côte Bleue Marine Park, no-take Reserve of Couronne (top panel) and temperature profile at the recording site (bottom panel). Call production and temperature show strong correlations ($\rho = 0.8$).