1 **Molecular characterization of *Fasciola gigantica* in Punjab, Pakistan to infer the**
2 **dispersal route among the neighbouring countries of the Indian subcontinent**

3
4 Zia Ur Rehman^a, Atsushi Tashibu^b, Michiyo Tashiro^b, Imran Rashid^{a*}, Qasim Ali^d, Osama
5 Zahid^c, Kamran Ashraf^a, Wasim Shehzad^a, Umer Chaudhry^{c*}, Madoka Ichikawa-Seki^{b*}

6
7 ^a *Department of Parasitology, University of Veterinary and Animal Sciences, Lahore, 54200,*
8 *Pakistan*

9
10 ^b *Laboratory of Veterinary Parasitology, Faculty of Agriculture, Iwate University, 3-18-8*
11 *Ueda, Morioka 020-8550, Japan*

12
13 ^c *Roslin Institute, University of Edinburgh, Midlothian, Scotland, EH25 9RG, United*
14 *Kingdom*

15
16 ^d *Department of Parasitology, Gomal University Dera Ismail Khan, Khyber Pakhtoon Khaw,*
17 *Pakistan*

18
19
20 *Corresponding authors:

21 Imran Rashid

22 Tel : +92 42 99 213 697

23 E-mail: imran.rashid@uvas.edu.pk

24
25 Madoka Ichikawa-Seki

26 Tel.: +81 19 621 6218

27 E-mail: madoka@iwate-u.ac.jp

28
29 Umer Chaudhry

30 Email: uchaudhr@exseed.ed.ac.uk

31

Abstract

Fasciola gigantica is considered to be a major pathogen causing fasciolosis in the Indian subcontinent, resulting in millions of dollars production losses to the livestock industry. To understand the dispersal origin and the spread patterns of *F. gigantica* is important for preventing the disease. A total of 53 *Fasciola* flukes collected from buffalo and goat in the Punjab province of Pakistan, were identified as *F. gigantica* based on the multiplex PCR for the phosphoenolpyruvate carboxykinase (*pepck*) and the PCR-restriction fragment length polymorphism (RFLP) for DNA polymerase delta (*pold*). A significant genetic difference between *F. gigantica* from buffalo and goats in Pakistan was indicated by the genetic analysis of two distinct mitochondrial markers [NADH dehydrogenase subunit 1 (*nad1*) and cytochrome C oxidase subunit 1 (*cox1*)]. Phylogenetic analysis of the seventeen *nad1* haplotypes of *F. gigantica* from Pakistan with those in neighbouring countries of the Indian subcontinent revealed that all the haplotypes were clustered in haplogroup A. *Fasciola gigantica* with the eight haplotypes might be expanded in Pakistan from Indian origin, along with the migration of the domestic animals, since they were related to Indian haplotypes. In contrast, the remaining nine haplotypes were not shared with any neighbouring countries, suggesting independent origin, or possibly come from neighbouring Middle East countries. Our study provides a proof of concept for a method that could be used to investigate the epidemiology of *F. gigantica* regarding the development of sustainable parasite control strategies.

Key words: *Fasciola gigantica*, *pepck*, *pold*, *nad1*, *cox1*.

1. Introduction

Fasciolosis, caused by the liver fluke of genus *Fasciola*, is a neglected zoonotic disease that results in a severe economic losses in the livestock industry (Aghayan et al., 2019). Fasciolosis is one of the most widely spread diseases reported from over 50 countries mostly in Asia, Africa and America (Mas-Coma, 2003; Mas-Coma et al., 2005; Toledo and Fried, 2014). The incidents of fasciolosis have increased over the past two decades, possibly because of the changes in farming practices, development of anthelmintic resistance and climatic changes (Sabourin et al., 2018). The genus *Fasciola* comprises of two important species. *Fasciola hepatica* is found in temperate zones, whereas *F. gigantica* is generally considered to be a parasite of tropical areas (Mas-Coma et al., 2009). Both species co-exist in subtropics, which can result in the formation of intermediate or hybrid forms mainly in Asian countries (Ichikawa and Itagaki, 2010b; Rokni et al., 2010). *Fasciola* infects the livers and bile ducts of ruminants and other mammals, while the snails of Lymnaeidae family act as their intermediate hosts (Toledo and Fried, 2014; Usip et al., 2014).

Mitochondrial markers have been used for the phylogenetic characterizations of *Fasciola* species to examine the propagation route of this group of parasites in many countries (Ai et al., 2011; Elliott et al., 2014; Ichikawa-Seki et al., 2017; Ichikawa and Itagaki, 2012; Semyanova et al., 2006; Thang et al., 2019). In the Asian subcontinent, *F. gigantica* has been divided into three haplogroups. Haplogroups B and C have been predominant mainly in Southeast Asian countries including Thailand (Chaichanasak et al., 2012), Myanmar (Ichikawa et al., 2011) and Indonesia (Hayashi et al., 2016). Haplogroup A has been distributed in the Indian subcontinent including East India (Hayashi et al., 2015), Bangladesh (Mohanta et al., 2014), Nepal (Shoriki et al., 2014) and Myanmar (Ichikawa et al., 2011). The zebu cattle (*Bos indicus*) and water buffalo (*Bubalus bubalis*) have been considered to be the definitive host of *F. gigantica* (Kikkawa et al., 2003). Hayashi et al. (2015) demonstrated that the haplogroup A of *F. gigantica* originated in the Indus River vally, and the unregulated animal movement might be involved in the spread of this parasite species. However, limited information is available on *F. gigantica* fluke in Pakistan to reveal the expansion history of this parasite through Indus River vally of the Indian subcontinent.

In this paper, we describe a study using liver flukes collected from buffalo and goat slaughtered in abattoirs in the Punjab province of Pakistan, with the following aims: i) to perform the species identification using the most reliable nuclear markers: phosphoenolpyruvate carboxykinase (*pepck*) and DNA polymerase delta (*pold*) genes (Shoriki et al., 2016); ii) to determine the propagation route of *F. gigantica* in the Indian

subcontinent using mitochondrial NADH dehydrogenase subunit 1 (*nad1*) and cytochrome C oxidase subunit 1 (*cox1*) genes.

2. Materials and Methods

2.1. Sample collection and gDNA extraction

A total of 14 *Fasciola* infected livers (7 buffalo and 7 goats) were collected from the animals slaughtered at the Punjab Agriculture & Meat Company (PAMCO) Lahore (31.4330° N, 74.1945° E). The livers were transported from abattoir to the laboratory on ice, where flukes were recovered from the biliary ducts by dissection. A total of 53 flukes (25 from 7 buffalo and 28 from 7 goats) (Table 1) were thoroughly washed with phosphate-buffered saline (PBS), and preserved in 70% ethanol until use. A small portion of the vitelline glands from the posterior part of each fluke was used for DNA extraction using the High Pure DNA Extraction Kit (Roche, Mannheim, Germany) following the manufacturer's protocols, and stored at -20°C until further use.

2.2. Multiplex PCR and PCR-RFLP of *pepck* and *pold* genes of *Fasciola* species

The fragments of *pepck* were amplified through the multiplex PCR assay previously described by Shoriki et al. (2016). The PCR amplicons were run on 1.8% agarose gels for 30 min to detect the *F. hepatica* (approximately 500bp), *F. gigantica* (approximately 240bp) or hybrid fragment patterns (both the fragments). The fragments of *pold* were analysed through the PCR-RFLP assay previously described by Shoriki et al. (2016). The PCR products were subsequently digested with *AluI* enzyme (Toyobo, Osaka, Japan) at 37 °C for three hours. The resultant products were run on 1.8% agarose gels for 30 min to detect the *F. hepatica* (approximately 700bp), *F. gigantica* (approximately 500bp) or hybrid fragment patterns (both the fragments).

2.3. PCR amplification and sequencing of *nad1* and *cox1* genes

The fragments of *nad1* (535bp) and *cox1* (430bp) genes were amplified through a PCR assay previously described by Itagaki et al. (2005). The PCR amplicons were purified using NucleoSpin Gel and PCR Clean-up Kit (Macherey-Nagel, Düren, Germany) and sequenced by Eurofin Genomics K.K. (Tokyo, Japan). The resultant DNA sequences were assembled using ATGC v. 6.0.3 (Genetyx Co., Tokyo, Japan) and the haplotypes were identified by GENETYX v. 10.0.2 (Genetyx Co.).

2.3. Median-joining network and diversity indices of *nad1* and *cox1* genes

Median-joining (MJ) network was constructed using Network v. 5.0.1.1 software (Tajima, 1989) to determine the phylogenetic relationships among the *nad1* and *cox1* haplotypes. Median-joining (MJ) network has been further used to compare the *nad1* haplotypes of the present study with the reference *nad1* haplotypes of *F. gigantica* from India (Ichikawa and Itagaki, 2010a), Bangladesh (Mohanta et al., 2014), Nepal (Shoriki et al., 2014), Myanmar (Ichikawa et al., 2011), Thailand (Chaichanasak et al., 2012), Vietnam (Itagaki et al., 2009), Indonesia (Hayashi et al., 2016), China (Peng et al., 2009), Korea (Ichikawa and Itagaki, 2012) and Japan (Itagaki et al., 2005). The reference *nad1* haplotypes were retrieved from the GenBank and their frequencies were referred from the previous reports.

The diversity indices, including the number of variable sites (S), the number of haplotypes (h), and the nucleotide diversity (π) were calculated using DnaSP software v. 5.1 (Librado and Rozas, 2009). Tukey's test was performed by GraphPad Prism v. 7.04 (GraphPad Software Inc., San Diego, CA, USA) to detect the significant differences in π values among the populations. First, the samples of the present study were compared to each other to find the differences between buffalo and goat. In the next step, the indices of *nad1* were compared with those of the reference populations of *F. gigantica* from India (Hayashi et al., 2015), Bangladesh (Mohanta et al., 2014), Nepal (Shoriki et al., 2014) and Myanmar (Ichikawa et al., 2011) to find relationships between the neighbouring countries.

The pairwise fixation index (F_{ST}) values between the *F. gigantica* samples derived from buffalo and goat were calculated using Arlequin program v. 3.5.2.2 (Loftus et al., 1994) to find genetic differences. If the F_{ST} values approaching 1 indicate extreme genetic differentiation between the two populations.

Results

3.1. Species identification

The fragment analysis by the multiplex PCR and the PCR-RFLP for *pepck* and *pold* showed that a total of 53 flukes collected from buffalo and goat were *F. gigantica* (Table 1). These results are complemented with the previous reports which suggest that *F. gigantica* is the predominant species in Punjab province of Pakistan (Chaudhry et al., 2016; Rehman et al., 2020).

3.2. Mitochondrial haplotype distribution of *F. gigantica* between buffalo and goat

The haplotype distribution was analysed separately for 53 individual *F. gigantica* flukes. A total of seventeen *nad1* haplotypes were detected in *F. gigantica* (Table 1). The MJ network revealed that PAK-nad1Fg1 and PAK-nad1Fg2 were the two predominant haplotypes present in both buffalo and goat (Fig. 1A) with four nucleotide substitutions between them. A total of seventeen *cox1* haplotypes were detected in *F. gigantica* (Table 1). The MJ network revealed that PAK-cox1Fg1 and PAK-cox1Fg2 were the predominant haplotypes found in both buffalo and goat (Fig. 1B), with three nucleotide substitutions between them.

The π for *nad1* and *cox1* were compared between *F. gigantica* samples acquired from both hosts. For the *nad1* gene, a higher π value was observed in the *F. gigantica* populations obtained from goat, but the result was opposite for the *cox1* gene (Table 2), and therefore, more diverse i.e. older population could not be determined. The F_{ST} values between the two hosts for both the genes (*nad1*: 0.12931, *cox1*: 0.16914) were statistically significant ($P < 0.05$), indicating the existence of genetic differentiation.

3.3. Comparative analysis of *F. gigantica* with that from neighboring countries

The MJ network analysis of seventeen *nad1* haplotypes revealed that PAK-nad1Fg2 haplotype had an identical sequence with the *F. gigantica* haplotypes from India (ND1-E6), Nepal (Fg-ND1-N1), Myanmar (Fg-M15), Bangladesh (Fg-NDI-Bd9) and Thailand (Fg-ND1-Thai13) (Fig. 2). The PAK-nad1Fg9, 10, 11, 12, 14 and 15 haplotypes had a single or double nucleotide substitutions present nearly to the PAK-nad1Fg2 haplotype. The PAK-nad1Fg13 haplotype had an identical sequence with the *F. gigantica* haplotypes from India (ND1-E7) and Myanmar (Fg-M16), and PAK-nad1Fg10 was identical to ND1-IN14 haplotype from India (Fig. 2). In contrast, the remaining nine haplotypes including PAK-nad1Fg1 (a most dominated haplotype) were not related to any reference haplotypes from neighbouring countries (Fig. 2).

The π values for *nad1* were compared between *F. gigantica* samples from Pakistan and neighbouring countries. The data suggest that the *F. gigantica* of the present study had the highest π value among the populations (Table 3), implying a higher nucleotide diversity in Pakistani populations as compared to the neighbouring countries.

Discussion

Historically, it has been suggested that *F. gigantea* might originate and spread by zebu cattle (*Bos indicus*) and water buffalo (*Bubalus bubalis*) in the Indian subcontinent (Peng et al., 2009). The water buffalo was domesticated in the Indus River valley (modern-day Pakistan) at around 7,000 to 8,000 BC, whereas zebu cattle was domesticated in the Indus Valley, Near and Middle East and Eastern Europe around 5,000 years ago (Bradley et al., 1996; Loftus et al., 1994; Tanaka et al., 1996). Since Pakistan is a part of the Indian subcontinent, free movement of zebu cattle and water buffalo probably play a significant role in the spread of *F. gigantea* in the region. Over the past few decades, high levels of animal movement have been reported in domestic ruminants in the Indian subcontinent (Kelley et al., 2016; Vilas et al., 2012). The farmers rear multiple species of animals to meet their livelihood in this region (Devendra, 2007). The mixed farming system might play an important role in the spread of *F. gigantea*. The animal movement patterns differ between farms, and *F. gigantea* infects a wide range of hosts including domestic and wild animals. Therefore, human activities potentially enable the spread of this parasite (Rojo-Vázquez et al., 2012). Hence genetic analysis are needed to understand the corresponding origin and spread of *F. gigantea* infections, which aid in the development of parasite control strategies (Hayashi et al., 2016).

In the present study, a significant genetic differentiation between *F. gigantea* samples from buffalo and goat was suggested by the F_{ST} value, which might be due to the difference in host immunity (Haroun and Hillyer, 1986; Piedrafita et al., 2004; Roberts et al., 1997) or due to the variances in the geographical position of these two hosts in the Punjab province of Pakistan. Generally, the lowlands of Punjab are more prone to flooding and reported to be highly populated with buffalo, in comparison, goat are resided in higher areas or keep moving to different areas due to human travelling (Afshan et al., 2014). However, the expansion history of *F. gigantea* between the hosts could not be inferred in the present study; the older host in Punjab could not be determined since the opposite statistical differences were observed in the π values of *nad1* and *cox1*.

The current study revealed that at least two origins of *F. gigantea* in Pakistan with reference to the neighbouring countries. In haplogroup A, the eight *F. gigantea* haplotypes of the present study had a close relationship with the haplotypes from India, Bangladesh, Nepal, Myanmar and Thailand. The MJ network indicates that the haplotypes detected in India were apparently more diverse than Pakistan, which indicates the hypothesis of the expansion of

these haplotypes from India to Pakistan. In contrast, the nine Pakistani haplotypes were not shared with any neighbouring countries, suggesting an independent origin, or possibly come from neighbouring Middle East countries where genetic analysis of *F. gigantica* has never been conducted. Further studies from different areas of Pakistan, as well as neighbouring Middle East countries, will be required to reveal the origin and dispersal direction of these haplotypes.

In summary, the present study provides preliminary insights into the origin and spread of *F. gigantica* in Pakistan and the neighbouring countries of the Indian subcontinent. We have also described the genetic difference between *F. gigantica* populations derived from buffalo and goat. Overall, the study provides a benchmark and opens a new avenue for more detailed analysis in this region. It might be helpful to involve higher samples size, more host species from different areas to get more conclusive results of the genetic diversity of *F. gigantica* among buffalo, goat, sheep and cattle as well as their possible spread patterns in the country and the subcontinent.

Acknowledgements

The authors acknowledge Mr Zain Ali khan, Mr Obaid Mohammad Abdullah and Mr Muhammad Omer Gulzar for their support and participation during sample collection. This work was supported by a Grant-in-Aid for Scientific Research (B) from MEXT KAKENHI (Grant Number16H05806).

Conflicts of interest

The authors declare no conflicts of interest.

Reference

- Afshan, K., Fortes-Lima, C.A., Artigas, P., Valero, M.A., Qayyum, M., Mas-Coma, S., 2014. Impact of climate change and man-made irrigation systems on the transmission risk, long-term trend and seasonality of human and animal fascioliasis in Pakistan. *Geospatial Health*, 317-334.
- Aghayan, S., Gevorgian, H., Ebi, D., Atoyan, H.A., Addy, F., Mackenstedt, U., Romig, T., Wassermann, M., 2019. *Fasciola* spp. in Armenia: Genetic diversity in a global context. *Veterinary parasitology* 268, 21-31.
- Ai, L., Weng, Y., Elsheikha, H., Zhao, G., Alasaad, S., Chen, J., Li, J., Li, H., Wang, C., Chen, M., 2011. Genetic diversity and relatedness of *Fasciola* spp. isolates from different hosts and geographic regions revealed by analysis of mitochondrial DNA sequences. *Veterinary parasitology* 181, 329-334.

- Bradley, D.G., MacHugh, D.E., Cunningham, P., Loftus, R.T., 1996. Mitochondrial diversity and the origins of African and European cattle. *Proceedings of the National Academy of Sciences* 93, 5131-5135.
- Chaichanasak, P., Ichikawa, M., Sobhon, P., Itagaki, T., 2012. Identification of *Fasciola* flukes in Thailand based on their spermatogenesis and nuclear ribosomal DNA, and their intraspecific relationships based on mitochondrial DNA. *Parasitol Int* 61, 545-549.
- Chaudhry, U., Van Paridon, B., Shabbir, M., Shafee, M., Ashraf, K., Yaqub, T., Gilleard, J., 2016. Molecular evidence shows that the liver fluke *Fasciola gigantica* is the predominant *Fasciola* species in ruminants from Pakistan. *Journal of helminthology* 90, 206-213.
- Devendra, C., 2007. Small farm systems to feed hungry Asia. *Outlook on Agriculture* 36, 7-20.
- Elliott, T., Muller, A., Brockwell, Y., Murphy, N., Grillo, V., Toet, H., Anderson, G., Sangster, N., Spithill, T., 2014. Evidence for high genetic diversity of NAD1 and COX1 mitochondrial haplotypes among triclabendazole resistant and susceptible populations and field isolates of *Fasciola hepatica* (liver fluke) in Australia. *Veterinary parasitology* 200, 90-96.
- Haroun, E., Hillyer, G.V., 1986. Resistance to fascioliasis--a review. *Veterinary parasitology* 20, 63-93.
- Hayashi, K., Ichikawa-Seki, M., Allamanda, P., Wibowo, P.E., Mohanta, U.K., Sodirun, Guswanto, A., Nishikawa, Y., 2016. Molecular characterization and phylogenetic analysis of *Fasciola gigantica* from western Java, Indonesia. *Parasitol Int* 65, 424-427.
- Hayashi, K., Ichikawa-Seki, M., Mohanta, U.K., Singh, T.S., Shoriki, T., Sugiyama, H., Itagaki, T., 2015. Molecular phylogenetic analysis of *Fasciola* flukes from eastern India. *Parasitology international* 64, 334-338.
- Ichikawa-Seki, M., Tokashiki, M., Opara, M.N., Iroh, G., Hayashi, K., Kumar, U.M., Itagaki, T., 2017. Molecular characterization and phylogenetic analysis of *Fasciola gigantica* from Nigeria. *Parasitology international* 66, 893-897.
- Ichikawa, M., Bawn, S., Maw, N.N., Htun, L.L., Thein, M., Gyi, A., Sunn, K., Katakura, K., Itagaki, T., 2011. Characterization of *Fasciola* spp. in Myanmar on the basis of spermatogenesis status and nuclear and mitochondrial DNA markers. *Parasitology international* 60, 474-479.
- Ichikawa, M., Itagaki, T., 2010a. Discrimination of the ITS1 types of *Fasciola* spp. based on a PCR-RFLP method. *Parasitology research* 106, 757-761.
- Ichikawa, M., Itagaki, T., 2010b. Discrimination of the ITS1 types of *Fasciola* spp. based on a PCR-RFLP method. *Parasitology research* 106, 757-761.
- Ichikawa, M., Itagaki, T., 2012. Molecular analysis of aspermic *Fasciola* flukes from Korea on the basis of the nuclear ITS1 region and mitochondrial DNA markers and comparison with Japanese aspermic *Fasciola* flukes. *Journal of Veterinary Medical Science*, 1203050785-1203050785.
- Itagaki, T., Kikawa, M., Sakaguchi, K., Shimo, J., Terasaki, K., Shibahara, T., Fukuda, K., 2005. Genetic characterization of parthenogenic *Fasciola* sp. in Japan on the basis of the sequences of ribosomal and mitochondrial DNA. *Parasitology* 131, 679-685.
- Itagaki, T., Sakaguchi, K., Terasaki, K., Sasaki, O., Yoshihara, S., Van Dung, T., 2009. Occurrence of spermic diploid and aspermic triploid forms of *Fasciola* in Vietnam and their molecular characterization based on nuclear and mitochondrial DNA. *Parasitol Int* 58, 81-85.

- Kelley, J.M., Elliott, T.P., Beddoe, T., Anderson, G., Skuce, P., Spithill, T.W., 2016. Current threat of triclabendazole resistance in *Fasciola hepatica*. *Trends in parasitology* 32, 458-469.
- Kikkawa, Y., Takada, T., Sutopo, Nomura, K., Namikawa, T., Yonekawa, H., Amano, T., 2003. Phylogenies using mtDNA and SRY provide evidence for male-mediated introgression in Asian domestic cattle. *Animal genetics* 34, 96-101.
- Librado, P., Rozas, J., 2009. DnaSP v5: a software for comprehensive analysis of DNA polymorphism data. *Bioinformatics* 25, 1451-1452.
- Loftus, R.T., MacHugh, D.E., Bradley, D.G., Sharp, P.M., Cunningham, P., 1994. Evidence for two independent domestications of cattle. *Proceedings of the National Academy of Sciences* 91, 2757-2761.
- Mas-Coma, S., 2003. Adaptation capacities of *Fasciola hepatica* and their relationships with human fascioliasis: from below sea level up to the very high altitude. *Taxonomy, ecology and evolution of metazoan parasites* 2, 81-123.
- Mas-Coma, S., Bargues, M.D., Valero, M., 2005. Fascioliasis and other plant-borne trematode zoonoses. *International journal for parasitology* 35, 1255-1278.
- Mas-Coma, S., Valero, M.A., Bargues, M.D., 2009. Chapter 2. *Fasciola*, lymnaeids and human fascioliasis, with a global overview on disease transmission, epidemiology, evolutionary genetics, molecular epidemiology and control. *Adv Parasitol* 69, 41-146.
- Mohanta, U.K., Ichikawa-Seki, M., Shoriki, T., Katakura, K., Itagaki, T., 2014. Characteristics and molecular phylogeny of *Fasciola* flukes from Bangladesh, determined based on spermatogenesis and nuclear and mitochondrial DNA analyses. *Parasitology research* 113, 2493-2501.
- Peng, M., Ichinomiya, M., Ohtori, M., Ichikawa, M., Shibahara, T., Itagaki, T., 2009. Molecular characterization of *Fasciola hepatica*, *Fasciola gigantica*, and aspermic *Fasciola* sp. in China based on nuclear and mitochondrial DNA. *Parasitology research* 105, 809-815.
- Piedrafita, D., Raadsma, H., Prowse, R., Spithill, T.W., 2004. Immunology of the host-parasite relationship in fasciolosis (*Fasciola hepatica* and *Fasciola gigantica*). *Canadian Journal of Zoology* 82, 233-250.
- Rehman, Z.U., Zahid, O., Rashid, I., Ali, Q., Akbar, M.H., Oneeb, M., Shehzad, W., Ashraf, K., Sargison, N.D., Chaudhry, U., 2020. Genetic diversity and multiplicity of infection in *Fasciola gigantica* isolates of Pakistani livestock. *Parasitol Int* 76, 102071.
- Roberts, J., Estuningsih, E., Widjayanti, S., Wiedosari, E., Partoutomo, S., Spithill, T., 1997. Resistance of Indonesian thin tail sheep against *Fasciola gigantica* and *F. hepatica*. *Veterinary Parasitology* 68, 69-78.
- Rojo-Vázquez, F.A., Meana, A., Valcárcel, F., Martínez-Valladares, M., 2012. Update on trematode infections in sheep. *Veterinary parasitology* 189, 15-38.
- Rokni, M.B., Mirhendi, H., Mizani, A., Mohebbi, M., Sharbatkhori, M., Kia, E.B., Abdoli, H., Izadi, S., 2010. Identification and differentiation of *Fasciola hepatica* and *Fasciola gigantica* using a simple PCR-restriction enzyme method. *Experimental parasitology* 124, 209-213.
- Sabourin, E., Alda, P., Vázquez, A., Hurtrez-Boussès, S., Vittecoq, M., 2018. Impact of human activities on fasciolosis transmission. *Trends in parasitology*.
- Semyenova, S.K., Morozova, E.V., Chrisanova, G.G., Gorokhov, V.V., Arkhipov, I.A., Moskvina, A.S., Movsessyan, S.O., Ryskov, A.P., 2006. Genetic differentiation in eastern European and western Asian populations of the liver fluke, *Fasciola hepatica*, as revealed by mitochondrial nad1 and cox1 genes. *Journal of Parasitology* 92, 525-531.

- Shoriki, T., Ichikawa-Seki, M., Devkota, B., Rana, H.B., Devkota, S.P., Humagain, S.K., Itagaki, T., 2014. Molecular phylogenetic identification of *Fasciola* flukes in Nepal. *Parasitol Int* 63, 758-762.
- Shoriki, T., Ichikawa-Seki, M., Suganuma, K., Naito, I., Hayashi, K., Nakao, M., Aita, J., Mohanta, U.K., Inoue, N., Murakami, K., Itagaki, T., 2016. Novel methods for the molecular discrimination of *Fasciola* spp. on the basis of nuclear protein-coding genes. *Parasitology international* 65, 180-183.
- Tajima, F., 1989. Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics* 123, 585-595.
- Tanaka, K., Solis, C.D., Masangkay, J.S., Maeda, K., Kawamoto, Y., Namikawa, T., 1996. Phylogenetic relationship among all living species of the genus *Bubalus* based on DNA sequences of the cytochrome b gene. *Biochemical genetics* 34, 443-452.
- Thang, T.N., Hakim, H., Rahimi, R.R., Ichikawa-Seki, M., 2019. Molecular analysis reveals expansion of *Fasciola hepatica* distribution from Afghanistan to China. *Parasitology international* 72, 101930.
- Toledo, R., Fried, B., 2014. *Digenetic trematodes*, Vol 480. Springer.
- Usip, L., Ibanga, E., Edoho, H., Amadi, E., Utah, E., 2014. Prevalence of Fascioliasis and the economic loss of condemned liver due to *Fasciola* infection in Cattle slaughtered at three abattoirs in Eket Urban, Akwa Ibom State of Nigeria. *Global Advanced Research Journal of Food Science and Technology* 3, 54-75.
- Vilas, R., Vázquez-Prieto, S., Paniagua, E., 2012. Contrasting patterns of population genetic structure of *Fasciola hepatica* from cattle and sheep: implications for the evolution of anthelmintic resistance. *Infection, Genetics and Evolution* 12, 45-52.

Figure Legend

Fig. 1: Median-joining network based on the mitochondrial (A) *nadI* and (B) *coxI* haplotypes of *F. gigantica* from Pakistani origin. Each circle indicates a single haplotype. Black colour indicates the haplotypes from Buffalo and white colour haplotypes from the goat. Small circles are the median vectors needed to connect the haplotypes.

Fig. 2: Median-joining network based on the mitochondrial *nadI* haplotypes of *F. gigantica* in Pakistan and other countries. The *Fasciola* flukes from Pakistan are shown in black colour. Each circle indicates a single haplotype. Small circles are the median vectors which are needed to connect the haplotypes. The haplotype codes are shown within or adjacent to the circles. Numbers on each circle and node indicate the number of flukes and the number of substitution sites, respectively.

F. gigantica
haplogroup B

F. gigantica
haplogroup A

hybrid *Fasciola* flukes

Table 1: Profiles of *Fasciola* flukes used in this study.

Host ID	Host species	Number of flukes	Nuclear DNA		Mitochondrial DNA				species
			<i>pepck</i>	<i>pold</i>	<i>nad1</i>	Accession No.	<i>cox1</i>	Accession No.	
B8	Buffalo	4	Fg	Fg	PAK-nad1Fg2	LC517886	PAK-cox1Fg17	LC520088	<i>F. gigantica</i>
			Fg	Fg	PAK-nad1Fg11	LC517895	PAK-cox1Fg12	LC517913	<i>F. gigantica</i>
			Fg	Fg	PAK-nad1Fg12	LC517896	PAK-cox1Fg13	LC517914	<i>F. gigantica</i>
			Fg	Fg	PAK-nad1Fg2	LC517886	PAK-cox1Fg2	LC517903	<i>F. gigantica</i>
B10	Buffalo	4	Fg	Fg	PAK-nad1Fg5	LC517889	PAK-cox1Fg5	LC517906	<i>F. gigantica</i>
			Fg	Fg	PAK-nad1Fg10	LC517894	PAK-cox1Fg2	LC517903	<i>F. gigantica</i>
			Fg	Fg	PAK-nad1Fg1	LC517885	PAK-cox1Fg1	LC517902	<i>F. gigantica</i>
			Fg	Fg	PAK-nad1Fg1	LC517885	PAK-cox1Fg6	LC517907	<i>F. gigantica</i>
B11	Buffalo	4	Fg	Fg	PAK-nad1Fg8	LC517892	PAK-cox1Fg4	LC517905	<i>F. gigantica</i>
			Fg	Fg	PAK-nad1Fg14	LC517898	PAK-cox1Fg12	LC517913	<i>F. gigantica</i>
			Fg	Fg	PAK-nad1Fg1	LC517885	PAK-cox1Fg1	LC517902	<i>F. gigantica</i>
			Fg	Fg	PAK-nad1Fg4	LC517888	PAK-cox1Fg1	LC517902	<i>F. gigantica</i>
B15	Buffalo	4	Fg	Fg	PAK-nad1Fg2	LC517886	PAK-cox1Fg12	LC517913	<i>F. gigantica</i>
			Fg	Fg	PAK-nad1Fg10	LC517894	PAK-cox1Fg2	LC517903	<i>F. gigantica</i>
			Fg	Fg	PAK-nad1Fg2	LC517886	PAK-cox1Fg11	LC517912	<i>F. gigantica</i>
			Fg	Fg	PAK-nad1Fg2	LC517886	PAK-cox1Fg10	LC517911	<i>F. gigantica</i>
B16	Buffalo	4	Fg	Fg	PAK-nad1Fg1	LC517885	PAK-cox1Fg1	LC517902	<i>F. gigantica</i>
			Fg	Fg	PAK-nad1Fg1	LC517885	PAK-cox1Fg1	LC517902	<i>F. gigantica</i>
			Fg	Fg	PAK-nad1Fg1	LC517885	PAK-cox1Fg1	LC517902	<i>F. gigantica</i>
			Fg	Fg	PAK-nad1Fg1	LC517885	PAK-cox1Fg1	LC517902	<i>F. gigantica</i>
B24	Buffalo	2	Fg	Fg	PAK-nad1Fg2	LC517886	PAK-cox1Fg10	LC517911	<i>F. gigantica</i>
			Fg	Fg	PAK-nad1Fg2	LC517886	PAK-cox1Fg10	LC517911	<i>F. gigantica</i>
B29	Buffalo	3	Fg	Fg	PAK-nad1Fg2	LC517886	PAK-cox1Fg14	LC517915	<i>F. gigantica</i>
			Fg	Fg	PAK-nad1Fg13	LC517897	PAK-cox1Fg2	LC517903	<i>F. gigantica</i>
			Fg	Fg	PAK-nad1Fg9	LC517894	PAK-cox1Fg11	LC517912	<i>F. gigantica</i>
Subtotal	7	25							
G1	Goat	4	Fg	Fg	PAK-nad1Fg3	LC515887	PAK-cox1Fg1	LC517902	<i>F. gigantica</i>
			Fg	Fg	PAK-nad1Fg1	LC517885	PAK-cox1Fg1	LC517902	<i>F. gigantica</i>
			Fg	Fg	PAK-nad1Fg4	LC517888	PAK-cox1Fg1	LC517902	<i>F. gigantica</i>
			Fg	Fg	PAK-nad1Fg1	LC517885	PAK-cox1Fg1	LC517902	<i>F. gigantica</i>
G2	Goat	4	Fg	Fg	PAK-nad1Fg2	LC517886	PAK-cox1Fg1	LC517902	<i>F. gigantica</i>
			Fg	Fg	PAK-nad1Fg1	LC517885	PAK-cox1Fg1	LC517902	<i>F. gigantica</i>
			Fg	Fg	PAK-nad1Fg2	LC517886	PAK-cox1Fg2	LC517903	<i>F. gigantica</i>
			Fg	Fg	PAK-nad1Fg2	LC517886	PAK-cox1Fg9	LC517010	<i>F. gigantica</i>
G3	Goat	4	Fg	Fg	PAK-nad1Fg1	LC517885	PAK-cox1Fg1	LC517902	<i>F. gigantica</i>
			Fg	Fg	PAK-nad1Fg6	LC517890	PAK-cox1Fg1	LC517902	<i>F. gigantica</i>
			Fg	Fg	PAK-nad1Fg2	LC517886	PAK-cox1Fg2	LC517903	<i>F. gigantica</i>
			Fg	Fg	PAK-nad1Fg1	LC517885	PAK-cox1Fg1	LC517902	<i>F. gigantica</i>
G4	Goat	4	Fg	Fg	PAK-nad1Fg1	LC517885	PAK-cox1Fg1	LC517902	<i>F. gigantica</i>
			Fg	Fg	PAK-nad1Fg15	LC517899	PAK-cox1Fg1	LC517902	<i>F. gigantica</i>
			Fg	Fg	PAK-nad1Fg1	LC517885	PAK-cox1Fg1	LC517902	<i>F. gigantica</i>
			Fg	Fg	PAK-nad1Fg6	LC517890	PAK-cox1Fg1	LC517902	<i>F. gigantica</i>
G5	Goat	4	Fg	Fg	PAK-nad1Fg1	LC517885	PAK-cox1Fg3	LC517904	<i>F. gigantica</i>
			Fg	Fg	PAK-nad1Fg1	LC517885	PAK-cox1Fg7	LC517908	<i>F. gigantica</i>

			Fg	Fg	PAK-nad1Fg1	LC517885	PAK-cox1Fg3	LC517904	<i>F. gigantea</i>
			Fg	Fg	PAK-nad1Fg1	LC517885	PAK-cox1Fg3	LC517904	<i>F. gigantea</i>
G6	Goat	4	Fg	Fg	PAK-nad1Fg7	LC517891	PAK-cox1Fg1	LC517902	<i>F. gigantea</i>
			Fg	Fg	PAK-nad1Fg3	LC515887	PAK-cox1Fg15	LC517916	<i>F. gigantea</i>
			Fg	Fg	PAK-nad1Fg1	LC517885	PAK-cox1Fg8	LC517909	<i>F. gigantea</i>
			Fg	Fg	PAK-nad1Fg1	LC517885	PAK-cox1Fg1	LC517902	<i>F. gigantea</i>
G7	Goat	4	Fg	Fg	PAK-nad1Fg16	LC517900	PAK-cox1Fg1	LC517902	<i>F. gigantea</i>
			Fg	Fg	PAK-nad1Fg16	LC517900	PAK-cox1Fg15	LC517916	<i>F. gigantea</i>
			Fg	Fg	PAK-nad1Fg1	LC517885	PAK-cox1Fg1	LC517902	<i>F. gigantea</i>
			Fg	Fg	PAK-nad1Fg17	LC517901	PAK-cox1Fg16	LC517917	<i>F. gigantea</i>
Subtotal			7	28					
Total			14	53					

Table 2: Diversity indices of *F. gigantica* populations based on the nucleotide sequences of mitochondrial markers.

Gene	Host	<i>N</i>	<i>S</i>	<i>h</i>	π	<i>SD</i>
<i>nad1</i>	Buffalo	25	14	11	0.00533	0.00049
	Goat	28	17	9	0.00668	0.00135
<i>cox1</i>	Buffalo	25	12	11	0.00612	0.00047
	Goat	28	10	8	0.00493	0.00125

N: number of flukes accessed; *S*: number of variable sites; *h*: number of haplotypes; π : nucleotide diversity. All values were statistically significant ($P < 0.05$).

Table 3 Diversity indices of *F. gigantica* within haplogroup A based on the nucleotide sequences of *nad1* gene.

Countries	<i>N</i>	<i>S</i>	<i>h</i>	π	<i>SD</i>
Pakistan	53	26	17	0.00650	0.00071
India	132	41	43	0.00252*	0.00023
Bangladesh	21	14	10	0.00312	0.00075
Nepal	20	16	10	0.00366	0.00088
Myanmar	13	7	6	0.00225*	0.00072

N: number of flukes accessed; *S*: number of variable sites; *h*: number of haplotypes; π : nucleotide diversity; *SD*: standard deviation; *: Statistically non-significant between the identical letter ($P > 0.05$), others were significant ($P < 0.05$).

F. gigantica from Thailand ($n = 1$) and Indonesia ($n = 1$) in haplogroupA was excluded from the calculation.