The framework of lncRNAs and genes at early pollen developmental stage in a PTGMS wheat line

Short title: Cold-induced pollen sterile in a wheat PTGMS line

Author for contact details: Li-ping Zhang

Jian-fang Bai^{1,2†} JB, Zi-han Liu^{1,2†} ZL, Yu-kun Wang^{1,2,4†} YW, Hao-yu Guo^{1,2,4} HG, Li-Ping Guo^{1,2} LG, Zhao-guo Tan^{1,2} ZT, Shao-hua Yuan^{1,2} SY, Yan-mei Li^{1,2} YL, Ting-ting Li^{1,2,4} TL, Wen-jing Duan^{1,2,4} WD, Jie-ru Yue^{1,2} JY, Feng-ting Zhang^{1,2} FZ, Chang-ping Zhao^{1,2*} CZ and Li-ping Zhang^{1,2*}

ORCID IDs: 0000-0001-8384-7001 (J.-F. B.); 0000-0002-3438-7130 (Z.-H. L.); 0000-0003-1574-6254 (Y.-K. W.); 0000-0003-3933-2478 (L.-P. Z.)

¹ Beijing Engineering Research Center for Hybrid Wheat, Beijing Academy of Agriculture and Forestry, Beijing 100097, China;

² The Municipal Key Laboratory of Molecular Genetic of Hybrid Wheat, Beijing 10097, China;

³ College of Life Science, Capital Normal University, Beijing 100048, China;

⁴ Division of Biological Science, Graduate School of Science and Technology, Nara Institute of Science and Technology, Ikoma, Japan

17

†shared first authorship.

* to whom correspondence should be addressed:

Dr. Liping Zhang, Beijing Engineering and Technique Research Center for Hybrid Wheat, Beijing Academy of Agriculture and Forestry, No. 9 Shuguang Park Middle Rd., Haidian District, Beijing 100097, China. Tel. +8601051503765, E-mail: lpzhang8@126.com

One sentence summary: Differential transcriptome analysis was utilized to identify the key factors for fertility transformation in a wheat photo-thermosensitive genic male sterile line.

Authors' contributions

JB supervised the experiments, wrote manuscript. ZL: observed cytological events and wrote manuscript. YW: edited the manuscript and completed the writing. LG: analyzed the RAN-seq data. HG: performed the RT-PCR and qPCR. ZT and WD: performed ultrathin sections. SY: managed the plants. YL, JY and TL: collected the samples. FZ: supervised the experiments. CZ: conceived the original screening and research plans. LZ: conceived the research plan, serves as the author responsible for contact and ensures communication.

33 All authors carefully checked and approved this version of the manuscript.

34 **Funding**

35 This study was supported by the National Key Research Project (2016YFD0101601);
 36 National Natural Science Foundation of China (31872881); Outstanding Scientist
 37 Cultivation Program of BAAFS (JKZX201907); Science and Technology Plan of Beijing
 38 (Z191100004019014); National Science Foundation for Young Scientists of China
 39 (31901490).

40

41

42 **Abstract:**

43 Wheat photo-thermosensitive genic male sterile (PTGMS) line is a vital material in the
44 two-line hybrid wheat breeding system in which functional pollen production is highly
45 associated with temperature during early developmental stage. Understanding the potential
46 mechanism of pollen infertility induced by low temperature in PTGMS wheat is crucial for
47 the effective utilization of genetic resources to guide wheat breeding. Herein, we combined
48 full-length single-molecular sequencing and Illumina short reads sequencing data to obtain
49 the high-resolution spatio-temporal transcriptome map of pollen under low temperature
50 stress at mother cell, dyad and tetrad stages in PTGMS line BS366. Cytological descriptions
51 and whole transcriptome analysis revealed a global landscape of low temperature altered
52 pollen fertility transformation via regulating the transcriptional patterns of cytoskeleton-
53 related lncRNAs and their target genes, which involved in the calcium signaling and vesicle
54 trafficking pathways on cytoskeleton homeostasis at different stages of meiosis. Overall,
55 our results provided the transcriptional and cytological evidences for understanding the low
56 temperature-induced pollen sterility deficiency in PTGMS wheat line.

57 **Keywords:** Wheat, photo-thermosensitive genic male sterile (PTGMS), cytoskeleton,
58 alternative splicing, lncRNA

59 **Introduction**

60 Pollen abortion contains multiple physiological, biochemical, and molecular changes such as the
61 abnormal degradation of the tapetum (Zheng et al., 2019), defective pollen wall (Marianne et al., 2002;
62 Wu et al., 2015), the level of kinetic of ATPases (Sane et al., 1997), the distribution and concentration of
63 Ca^{2+} in anther (Tian et al., 1998), the regulation of the cytoskeleton (Tang et al., 2012; Wang et al., 2018),
64 accumulation ROS in tapetum (Liu et al., 2018a), and the abnormalities of cell signaling transductions (Liu
65 et al., 2018b). Based on mentions above, it is well known that cytological biological events and related
66 genes may play important roles in regulation of plants male fertility. In CMS lines wheat, premature or
67 delayed PCD by tapetal cells disorganized the supply of the nutrients to microspores, thereby resulting
68 in pollen abortion (Meng et al., 2016). Previous studies suggested this irregular tapetal PCD was tightly
69 controlled by evolutionarily conserved transcriptional cascades (Liu et al., 2020). In recent years, hybrid
70 breeding has a remarkable success in several allogamous species such as maize, sunflower, sorghum,
71 sugar beet, and rye, but not be fully exploited in autogamous crops (Longin et al., 2012). Wheat is an
72 autogamous crop, however, hybrid seed production requires cross-pollination of the female parent by
73 pollen from the male parent. Photoperiod and/or thermo-sensitive genic male sterility (P/TGMS) is an
74 important material in two-line breeding system to explore the potential of heterosis. Previous studies
75 showed that the male sterility of P/TGMS lines are contributed to abnormal pollen development (Bai et
76 al., 2017). Further cytological studies showed that P/TGMS wheat, it exhibited disordered distribution

of the cytoskeleton, including microfilaments and microtubules when exposed to a sterile environment during the fertility-sensitive stage (Tang et al., 2011; Wang et al., 2018). Additionally, previous RNA-seq studies also have mostly focused on the transcripts of male sterility and thousands of differentially expressed genes have been reported (Liu et al., 2016). As we all know, in male sterile wheat, the process of pollen abortion reflects extremely complex reprogramming of gene expression involving chromatin modification, transcription, posttranscriptional processing, posttranslational modification, and protein turnover. However, the complex posttranscriptional and translational levels molecular mechanisms of the male sterility, especially the wheat P/TGMS line induced by low temperature, are currently still not clear.

As a hexaploid, wheat has a large and complex genome, estimated to reach approximately 17G, which composes three closely-related and independently maintained genomes that are the result of a series of naturally occurring hybridisation events. With the continuous advancement of technology, second-generation sequencing technology (e.g., Illumina sequencing), could feature high-throughput capability and provide high-quality reads. However, the short-read length potentially introduces errors in inaccurately identify the transcript results. Third-generation sequencing is a single-molecule real-time sequencing technology, (e.g., PacBio sequencing), could provide full-length transcript information, detect single-molecule structure, provide complete mRNA structure and is, therefore, well suited for transcript recovery and isoform detection in species with well sequenced and/or incomplete genome sequences (Abdel-Ghany et al., 2016; Wang et al., 2016). In some crop plants, many scientists have been studied the isoform change when plant response to stress at gene transcriptional levels, such AS level and long non coding RNA (lncRNA) (Abdel-Ghany et al., 2016; Wang et al., 2016) using combination of TGS and NGS technology. AS is a widely recognized RNA processing mechanism in eukaryotic species, playing a major role in the molecular biology of the cell, and within humans it has been implicated in multiple genetic disorders (Wang et al., 2016). AS is a critical posttranscriptional event which comes from alternate splice site choices in a single gene locus, including intron retention (IR), exon skipping (ES), alternative 5' splicing site (Alt5'SS) and alternative 3' splicing site (Alt3'SS) (Wang et al., 2019). In higher plant, it has been reported that the AS events are involved in a wide range of developmental and physiological processes including responses to stress. For example, about 60% of Arabidopsis intron-containing genes are generated by AS (Marquez et al., 2012). It has been demonstrated that AS is important for cold response when SFs mis-expressed during cold sensitivity or tolerance treatment (Laloum et al., 2018). In wheat, Liu et al. (2018) performed genome-wide analysis of alternative splicing (AS) responses to drought stress (DS), heat stress (HS) and their combination (HD) in wheat seedling to investigate the regulation of AS during these stress processes (Liu et al., 2018c). More recently, Wang et al. (2019) also investigated the spatio-temporal landscape of heat adaptations in wheat filling grain and flag leaves at transcriptional and AS levels by hybrid sequencing (second- and third-generation sequencing). These studies strongly suggest that AS networks are central co-ordinators of the stress response. Although AS plays an important role in stress response, it is not clear in studies related to pollen abortion. In addition to AS the lncRNA were also generated by hybrid sequencing (Wang et al., 2019). Many studies have shown that lncRNA could regulate genes at the transcriptional and post-

transcriptional levels by acting as signals, decoys, scaffolds, and guides (Heo and Sung, 2011). However, virtually nothing is known about the extent and timing of the contribution of AS and lncRNA or how AS and lncRNA determine the dynamic changes of transcriptome required for regulating male sterility for P/TGMS line. Here, to illustrate the regulation of these factors and genes on fertility transformation, we used hybrid sequencing strategy and used a conventional fertile wheat variety as a control and differential genes background to analyze the role of AS and lncRNA in regulating male sterility during fertility transformation stages in wheat PTGMS line BS366.

Results

Morphological characteristics of mature pollen under different conditions

In order to identify the influence of environment on the fertility of PTGMS line BS366, we observed the phenotype of BS366 at the trinucleate stage under different fertility conditions and used conventional wheat Jing411 as control (Fig. 1). Scanning electron microscopy (SEM) examination at trinuclear stage revealed that epidermis cells of the Jing411 in two conditions and BS366 anthers in fertile conditions were arranged closely, on the contrary, the epidermis cells of BS366 anthers in sterile conditions were incomplete and occurred loss. Moreover, the inner epidermal ubisch bodies of anther for BS366 in sterile conditions were abnormal and accumulated more sparsely distributed. Based on observations of microspores at the trinucleate stage, the microspores were uniformly spheroid and had finely reticulate ornamentation on their surface in anther of Jing411 and BS366 in fertile conditions, while the sterile microspores were extremely atrophied. According to I₂-KI staining, BS366 in fertile conditions pollen grains were 60% stained black, whereas pollen grains were all wrinkled and inadequately stained in sterile condition, and exhibited completely aborted characteristics, indicating that low temperature could induce the male fertility conversion in PTGMS line BS366.

Overview of sequencing data

Previous study showed that the period from the pollen mother cell stage to the tetrad stage is the most sensitive to low temperature for the pollen of BS366 (Bai et al., 2017). And previous cytological studies also showed that there were abnormalities of film-forming body and cell plate in this process (Tang et al., 2011). To comprehensively investigate the wheat transcriptomes present during the fertility transition, hybrid sequencing were performed on the anther of BS366 and Jing411 from different fertility condition with three pollen development stages (S1: pollen mother cell stage, S2: dyad stage, and S3: tetrad stage) (Fig. S1). Totally, 247,486 and 240,993 circular consensus sequence reads for BS366 and Jing411 were yielded, respectively (Table S2). In total, 209,967 and 190,280 reads of full-length non-chimeric (FLNC) (84.84% and 78.96%) from these circular consensus sequence reads, were identified based on the inclusion of 5' and 3' primer, and 3' poly(A) tails, followed by error correction. Then, 209,967 and 190,280 high-quality isoforms were uniquely mapped to the IWGSC RefSeq v1.0 (Table S3). Finally, 16,000 and 14,864 transcripts were obtained

from BS366 and Jing411, followed by error correction using Illumina short reads (Table S4). The length of transcripts was mainly concentrated in the range of 10^3 - 10^4 for BS366, Jing411 and IWGSC RefSeq v1.0, suggesting the generated data are accurate and reliable, could be used as a reference in this study (Fig. 2 and Fig. S2). These transcripts were derived from 13,577 and 13,010 gene loci. Of which, 1,185 and 902 are new gene loci and 6,248 and 4,918 are new transcripts (Table S4). By comparison with the IWGSC RefSeq v1.0 annotation, PacBio transcripts could be classified into seven groups including PacBio data, DEG, DAS, lncRNA, fusion gene analysis (Fig. 3).

Identification of alternative splicing events during fertility transition

AS is an important biological event of post-transcriptional regulation in organisms (Wang et al., 2019). A few studies have explored the relationship between AS and fertility transition which regulated by external environment in plants (Capovilla et al., 2015). Therefore, we analyzed the patterns of fertility transition-induced AS events from the RNA sequencing data. In this study, four main AS events including Alternative 3' splice site (Alt3'SS), Exon skipping (ES), Intron retention (IR) and Mutually exclusive exon (MEX) were identified from three pollen developments stages of BS366 and Jing411. As shown in Table 1, total 35,248 and 33,099 AS events with 20,312 and 19,490 AS genes were identified from three pollen developments in BS366 and Jing411, respectively. Of which 11,619, 11,840 and 11,789 AS events in BS366 and 10,980, 11,084 and 11,035 AS events were determined on subgenomes A, B and D, respectively (Table S5). It was found that ES was the most abundant (91.08%-91.98%) AS events in both BS366 and Jing411, followed by MEX (4.84%-5.24%), IR (3.03%-3.67%) in BS366 (Table 1), and some orders in Jing411 and for AS genes (Table S5).

Identification of sterility -related AS events and genes in BS366

The high-resolution temporal transcriptomes allowed us to determine the specific stage at which differentially spliced genes (DSGs) and differentially expressed genes (DEGs, fold change ≥ 2.0 , FDR-adjusted P -value < 0.05) showed a significant change, along with the magnitude and trend of that change. To identify the specific time of significant changes of DSGs and DEGs as well as analysis the sterility-related DSGs and DEGs in BS366, DSGs and DEGs from three key pollen development stages of BS366 during fertility transition BS366 were screened. By comparing the DSGs of three stages, in total, 108, 130 and 141 genes with 118, 141 and 160 AS events were identified from pollen mother cell stage, dyad stage and tetrad stage respectively after critically filtering process. For DEGs, in total, 39, 73 and 412 DEGs were identified from three stages (Figure 4A). The change trend of DSGs and DEGs in the three stages was synergistically rising, in which the DSGs and DEGs contained in tetrad stage have significant changes, which may be the key stage of fertility transformation.

Here four DSGs were found in shared DSGs and DEGs in the three stages (Fig. 4B), in which the genes encoding zinc finger domain protein 1A (*TraesCS5D02G371100*) and UTP--glucose-1-

phosphate uridylyltransferase (*TraesCS5A02G353700*) involved in pollen formation (Chivasa et al., 2013), and the gene encoding coatomer subunit gamma-2 (*TraesCS1D02G156000*) was associated with vesicle transport (Hamlin et al., 2014). Moreover, cytoskeleton-related gene kinesin-4 (*TraesCS3B02G196600*), vesicle transport-related gene encoding vacuolar protein sorting-associated protein (*TraesCS4B02G382900*), and pollen formation-related genes encoding zinc finger BED domain-containing protein (*TraesCS3B02G126800*) and MYB transcription factors (*TraesCS3B02G243600*) were found in co-DEGs of the three stages (Figure 4C). Interestingly, the gene encoding kinesin-4 (*TraesCS3B02G196600*) was differentially expressed in three stages at the same time and occurred differentially AS in stage 3 (Fig. 4D and E), and the kinesin-4 has been proved to be involved in vesicle transport and cytoskeleton formation (van Riel et al., 2017), thereby inferring that AS of this gene could work together with transcriptional regulation to involve in fertility transformation of PTGMS line BS366. qPCR analysis verified the correctness of RNA-seq results (Fig. 5F). The primers used in this study were listed in Table S1.

Comparative analysis of the biological functions regulated at AS and transcription levels

GO analysis showed that the most of DSGs were correlated with some cytological and molecular events in the process of pollen development including cytoskeleton (such as “cytoskeleton” and “cytoplasmic microtubule organization”), calcium regulation (such as “calcium ion binding” and “calcium-dependent phospholipid binding”), vesicle transport (such as “vesicle-mediated transport” and “vacuole”), and pollen formation (such as “cell wall organization” and “glucose-1-phosphate uridylyltransferase activity”) during the three key stages of fertility conversion (Table S6). It has been reported these GO terms identified are involved in the male sterility, for example, the DSGs encoding the protein MOR1 (*TraesCS3B02G371500* in S1, *TraesCS3A02G339900* in S2, and *TraesCS3B02G371500* in S3) was critical for the orderly assembly of microtubules (Kawamura et al., 2006). Microtubules and microfilaments control the whole double fertilization process of pollen, is necessary for the normal development of pollen. Therefore, low temperature induced DSGs was a crucial further layer of regulation for pollen development, thereby possibly leading to fertility conversion.

To further identify the sterility-related genes, weighted gene co-expression network analysis (WGCNA) were performed with all DEGs from BS366 based on the seed setting rate (Table S7). The analysis of module-trait relationships analysis showed that the module ‘Red’ ($r = -0.98$, $p = 3e-27$) was highly correlated with male sterility in the six samples (Fig. 5B and C). Interestingly, GO analysis suggested that these DEGs in this module were also mainly concentrated in the GO terms associated with cytoskeleton, calcium regulation, vesicle transport, and pollen formation, which were consistent with that of DSGs. Furthermore, cytoskeletal-regulatory complex EF hand

(*TraesCS7A02G553100*), actin-related protein 9 (*TraesCS2D02G029100*), and microtubule-associated protein RP/EB family member 3 (*TraesCS2D02G523000*) were greater connectivity in these DEGs, suggesting they may be strongly associated with male sterility (Fig. 5D). Thus, transcriptional regulation may play a major role in cytoskeleton, calcium regulation, vesicle transport, and pollen formation, which coordinates with AS regulation to induce pollen abortion in wheat.

Differences of diverse transcription factors (TFs) during fertility transition

Transcription factors (TF) that perceive environmental signals and activate the expression of related genes play master roles in gene regulatory networks in the processes of growth and development including pollen development in plants (Wang et al., 2018). In this work, total 522 TFs with 633 transcripts from Pacbio data were annotated which belonging to major 48 families (Table S8). Furthermore, 34 wheat TFs were differentially expressed (DE-TFs), and 6 genes were differentially spliced (DS-TFs) during fertility transition (S1, S2 and S3) in BS366 after removing background, respectively (Fig. 6 and Tables S9). In addition, enrichment analysis was performed on these TFs at each pollen development stage to reveal the fertility transition-related signaling. As shown in Fig. 6, bZIP family was significantly enriched in DE-TFs at dyad stage, MYB and bHLH families were enriched in DE-TFs at tetrad stage. More, there were several members of the HMG, ARF, NAC and bHLH families were differentially expressed at the tetrad stage providing evidence that these families participated in fertility transition signaling transduction.

Function analysis of predicted lncRNAs and their targets

After filtering by CPC, CNCI, CPAT and Pfam, total 53 lncRNAs (>200bp) were obtained from Pacbio data (Fig. 7A). Generally, lncRNAs regulate gene expression via *cis* (regulation of neighboring loci) or *trans*-acting mechanisms. It has been proposed that lncRNAs that are synthesized at a low level are likely to act in *cis*, whereas those accumulate at a higher level are able to act in *trans* (Kornienko et al., 2013). Identification and analysis of candidate target genes could provide insight into the functions of lncRNAs in fertility transition of PTGMS line. In this study, 38 ncRNA-mRNA pairs were identified as *cis*-regulation and five were *trans*-regulation lncRNA-mRNA pairs from 53 lncRNAs after filtering (Table S10). In this five *trans*-regulation lncRNA-mRNA pairs, only PB.18919.1 have *trans*-regulation targets and other four lncRNAs have both *cis* and *trans*-regulation targets. There were 14 lncRNAs which no targets were found (Table S10). In addition, like other non-coding RNA, such as miRNA (Bai et al., 2017), the same lncRNA can regulate multiple mRNA genes, different lncRNA molecules can also be synergistic regulation of the same mRNA gene (Table S10), indicating that lncRNAs with their targets may participate in multiple regulation pathway.

To investigate lncRNA functions in regulation of fertility transformation, GO analysis on predicted targets was performed. As shown in Fig. 7B, the most frequent "molecular function" term was "peptide alpha-N-acetyltransferase activity", followed by "voltage-gated chloride channel

activity", and "microtubule motor activity" for targets. The most frequent "biological process" term was "microtubule organizing center organization", followed by "microtubule nucleation" and "spindle assembly". The results indicated that these lncRNA with their targets are involved in process of cell division and important for male sterility in wheat.

qPCR and heatmap analysis were also performed in lncRNAs and their corresponding targets to verified their expression patterns during different pollen development stages (Fig. 7C and D). Coordinated expression was found between most lncRNAs and their targets (Figure 7D and Table S11). PB.51.4, PB.235.1, PB.18919.1 and PB5272.1 with targets (*TraesCS5A02G351600*, *TraesCS7D02G239700*, *TraesCS5A02G390000* and *TraesCS2D02G553800* down regulated during pollen development in both fertile condition and sterile condition. However, the other target gene of PB.18919.1, *TraesCS3D02G288100* (Copper transport protein ATX1) showed up-regulated in sterile condition and down -regulated in fertile condition (Fig. 7D).

RT-PCR validation of the DSGs

In this study, four DSGs were selected randomly from pollen mother cell stage, dyad and tetrad stage in different condition to validate the accuracy of AS events using reverse transcription polymerase chain reaction (RT)-PCR (Fig. 8). The isoforms of each DSG were designed primers to amplify all predicted transcripts and cloned using Sanger sequencing (Table S1). The results as shown by a gel banding pattern in Figure 8, the size of each amplified fragment was consistent with that of predicted fragment (Fig. 8). It was also found that expression of transcript isoforms exhibits a stage-preferential pattern. For example, *TraesCS3B02G326100* (encoding a Lipid-A-disaccharide synthetase protein), which was produced by an A5'SS event, was preferentially expressed at dyad stage and tetrad stage in fertile condition and at pollen mother cell stage in sterile condition. It therefore is example of stage specific RNA isoforms.

Effects of altered alternative splicing and gene expression on important cellular events

Hybrid sequencing showed that most of DSGs, DESs and lncRNA regulated targets could be related to chromosomal movement, process of cell division, cytoskeleton activity, cell plate formation. To further verify the results of the above analysis, the changes of microtubules, microfilaments, cell plate, chromosomes and calcium throughout critical periods of fertility transformation in pollen cells of BS366 under fertile and sterile conditions were observed (Fig. 9A). During pollen mother cell stage, in the fertile condition, the pollen mother cell took on a normal oval shape, the cytoskeleton was evenly distributed and the polar perinuclear microtubules were initially formed. However, in the sterile condition, the overall cellular morphology of pollen mother cell was wrinkled, chromatin was abnormally condensed and arranged scattered in the nuclear region, and microtubules and microfilaments appear as a radial and disordered filament and the polar microtubules were not obvious. Up to dyad stage, the fertile pollen mother cell proceeded normal cytokinesis and formed a distinct cell plate, while the division of sterile cell underwent disruption, and it's worth noting that the sterile dyad occurred absence of the cell plate. Especially during the

tetrad stage, the sterile tetrad happened severe malformation and the cytoskeleton was sparse and disordered compared with BS366 of fertile condition. Thus, under low temperature stress, cytoskeleton related genes and lncRNA regulated targets underwent differential alternative splicing and differential expression, which led to the abnormal meiosis of pollen mother cells, including the concentration of chromatin, the scattered distribution of cytoskeleton and the absence of cell plates.

It was also found that some DEGs such as *TraesCS5B02G336100* (phospholipase D delta), *TraesCS3B02G456100* (60S ribosomal protein L38) and *TraesCS5B02G160500* (calmodulin-binding receptor-like cytoplasmic kinase 3) were all up-regulated and associated with Ca^{2+} distribution (Fig. 5E). Meanwhile, six DSGs encoding calmodulin related protein also was screened (Fig. 5A). Many studies have shown that the dynamic distribution of Ca^{2+} was related to pollen sterility in anthers. In order to further verify whether fertility conversion is related to Ca^{2+} distribution, we used potassium antimonate to observe the distribution of Ca^{2+} in pollen and tapetum of BS366 under fertile and sterile conditions during pollen development (Fig. 9B). During the process of pollen mother cell division to tetrad, abundant Ca^{2+} precipitates were gradually accumulated on the cell surface, and the distribution of Ca^{2+} in the cytoplasm was very little, which maintained a low concentration of Ca^{2+} distribution environment. However, in the sterile condition, Ca^{2+} was less on the surface of sterile pollen cells, and excessive accumulation of Ca^{2+} precipitated in cytoplasm. Therefore, we concluded that the up-regulation of calcium-related DEGs and the regulation of DSGs may lead to the abnormal function of calcium pump or calcium channel in the cells and the inability to discharge the excess Ca^{2+} out of the cells, thereby leading to the increase of Ca^{2+} concentration in the cytoplasm of sterile pollen, and ultimately leading to pollen abortion.

According to the above DE-TFs, DEGs and DSGs analysis, we found many DEGs annotated as polygalacturonase, glycosyl transferase family 8, ABC transporter were all significantly down-regulated. In addition, DSGs encoding MYB-related protein, ABC transporter B family member 29, pectinesterase 31 and UTP--glucose-1-phosphate uridylyltransferase were identified, which may participate in plant cell wall synthesis (Schubert et al., 2019). In plants, MYB transcription factors play a very important role in the pollen development process, involving various key steps in the pollen formation process, including the programmed cell death (PCD) of tapetum, the deposition of callose, the formation of pollen wall and the accumulation of sporopollenin. Once one of the above abnormalities occurs, it will lead to pollen abortion (Schubert et al., 2019). Previous studies have shown that the large superfamily ABC transporter proteins are involved in translocation of a broad range of substances across membranes using energy from ATP hydrolysis such as transport of sporopollen (the main component of pollen exine) so they are also required for pollen exine formation (Chang et al., 2018). The tapetum provides sucrose, proteins, lipids, and sporopollenin to support the growth and development of the pollen via its degradation and secretion, and the progress of secreting nutrients to pollen is considered to be vesicular trafficking (Liu et al., 2020). Here, vesicular trafficking related DEGs encoding Sec23/Sec24 trunk domain protein, coatomer WD

associated region were significantly down-regulated, and identified 15 related DSGs (Fig. 5A and E). To verify the correctness of the above analysis, TEM was used to observe the degradation of tapetum and the formation of pollen wall of BS366 in fertile and sterile conditions (Fig. 9C). During pollen mother cell stage and dyad stage, there was no obvious difference between tapetum of BS366 under two conditions. Up to the tetrad stage, the fertile tapetum structure was complete and connected closely with the middle layer, whereas the tapetum of sterile condition separated from the middle layer. From the release of microspores from tetrapods to the trinuclear stage, the degradation of sterile tapetum was significantly faster than that of fertile tapetum. During this process, the tapetosome of sterile condition were also deformed and missing. Because of the degradation of tapetum in advance, the callose around the tetrad deposited abnormally, subsequently, the pollen wall of microspore released from the tetrad also deformed, which was manifested as the abnormal accumulation and uneven distribution of sporopollenin in the outer wall of pollen. Thus, we conclude that low temperature stress may induce the change of genes encoding MYB transcription factor, ABC transporter and glucose metabolism related enzymes, which may lead to the advanced degradation of tapetum and the deformity of pollen wall.

Discussion

Disorder of cytoskeleton is an essential factor for the pollen abortion regulation

In the past decades, many studies showed that cytoskeleton was involved in male sterility in plants. In the PTGMS rice line Peiai 64S displayed abnormal distribution in microtubules at the meiosis stage, no polar microtubules in pollen cells at the zygotene stage and rarefied perinuclear microtubules in diakinesis (Xu et al., 2001). Although Wang et al. (2018) found that disordered and asymmetrical distribution of microfilaments and microtubules in sterility pollens, nevertheless, the relationship of cytoskeleton homeostasis with pollen abortion are still unclear in PTGMS wheat line BS366 (Wang et al., 2018). Some studies showed that microtubules display dynamic instability, bouts of rapid growth followed by catastrophic shrinking, and the balance between these phases can be modulated by MAPs which generally increase polymerisation (Kawamura and Wasteney, 2008). In our study, a MAPs-related hub gene (*TraesCS2D02G523000*), encoding the microtubule-associated protein RP/EB family member 3, was significantly up-regulated during the three stages in the sterile condition in BS366 (Fig. 5E), suggesting that the abnormal expression of MAPs-related gene may lead to the aggregation disorder of cytoskeleton. For our DSGs analysis, 17 DSGs including two genes annotated as *MOR1* were found to be related to cytoskeleton (Table S6). In *Arabidopsis thaliana*, *MOR1*, the homologue of *Xenopus MAP215*, promoted rapid growth and shrinkage, and suppressed the pausing of microtubules in vivo (Kawamura and Wasteney, 2008). In this study, *MOR1* with differential alternative splicing, may hinder the rapid growth of cytoskeleton abnormally in the critical stage of abortion, thereby affect the development of pollen. In addition, lncRNAs with their targets were also involved in cytoskeleton. Cell cycle regulation related gene *NSA2* (Nop seven-associated 2) could blocked the cell cycle in G1/S transition in

Arabidopsis thaliana. Here, *NSA2* (*TraesCS2B02G131900*) (target of lncRNA PB.3805.1) was up regulated in SS3 and down regulated in FS3. At the same time, our cytological observation also verified the above analysis (Fig. 9).

The abnormal calcium messenger system might be related to pollen abortion

At present, it is known that the calcium messenger system is often in the center of signal cascade. Tian et al. (1998) described the anomalies in the distribution of calcium in anthers of PGMS rice, which displayed the failure of pollen development and pollen abortion. It was found that calcium precipitates were abundant in the middle layer and endothecium in sterile anthers, but not in the tapetum (Tian et al., 1998). In this study, calcium messenger system-related genes were excavated from DSG and DEG sets, for instance, *TraesCS6D02G176800*, which was annotated as *BON 1*; *TraesCS4A02G407100*, which was annotated as Calmodulin-binding transcription activator 2 (*CAMTA*); *TraesCS6B02G230900*, which was annotated as Calmodulin binding protein-like (*CBPL*); *TraesCS5B02G160500*, which was annotated as calmodulin binding receptor like cytoplasmic kinase 3 (*CRCK3*), and *TraesCS7A02G553100*, which was annotated as cytoskeletal-regulatory complex EF hand from DEGs, suggesting that calcium messenger system related genes might play essential roles in pollen fertility transformation after encountering low temperature during meiosis (Table S6). Ca^{2+} , Calmodulin (CaM) and CaM binding propeins (CaMBP) are involved in the process of dynamic distribution of cytoskeleton that regulated by microtubule/microfilament associated proteins (MAPs). Microfilament assembly will be promoted when the concentration of $[\text{Ca}^{2+}]_{\text{cyt}}$ is decreased, and that will be inhibited when the concentration of $[\text{Ca}^{2+}]_{\text{cyt}}$ is increased (Helper and Callaham, 1987). In resistant cowpea, elevation of $[\text{Ca}^{2+}]_{\text{cyt}}$ lead to deassembly of microtubule during rust fungal infection (Xu and Heath, 1998). To future investigated the mechanism of calcium in pollen fertility transformation, the potassium antimonate was used to locate Ca^{2+} in fertile and sterile anthers of BS366. It was found that pollen cell of BS366 with high concentration of $[\text{Ca}^{2+}]_{\text{cyt}}$ and low Ca^{2+} on the cell surface showed pollen abortion, suggested that the reason for pollen abortion may be due to the calcium pump or calcium channel dysfunction, which could not discharge excess Ca^{2+} outside the cell, so that the middle layer of anther was abundant in calcium precipitation (Fig. 9). The up-regulation of calcium-related DEGs and the regulation of DAS may be the reason of calcium message system dysfunction (Fig. 5A and E).

The abnormal vesicle trafficking is related to pollen abortion

In plants, vesicle trafficking is the main way of material and information exchange between organelles and is also important for maintaining homeostasis, which is involved in many biological processes such as cell wall formation, cell secretion and environmental response (Singh et al., 2018). The membrane vesicle transport machinery includes phospholipids and integral membrane proteins, such as vesicle-associated membrane proteins (VAMPs), the latter being the major constituent of

soluble NSF (N-ethylmaleimide-sensitive factor) attachment protein receptors (SNARE) complexes (Han et al., 2017). The vesicle-associated proteins (VAPs) are type II integral ER membrane-bound proteins tethered to the membranes and have been implicated in different processes such as membrane trafficking, lipid transport and metabolism, and unfolded protein response. SNARE complexes are responsible for fusion of vesicles with the target membranes, which is involved in many processes, such as cell plate formation, ion channel regulation, plant growth and development, plant tropism response (Jena, 2011). Sec23/Sec24, the core component of the coat protein complex II (COPII), functions to transport newly synthesized proteins and lipids from the endoplasmic reticulum (ER) to the Golgi apparatus in cells for secretion (Jing et al., 2019). However, information on the role of vesicle trafficking related proteins in wheat pollen development is scanty. In the present study, the gene encoding putative vesicle-associated protein 4-2 (VAP4-2) exhibited significant AS pattern change in PTGMS line BS366. Moreover, the WGCNA analysis results showed the hub DEGs include a soluble NSF attachment protein (SNAP), a conserved oligomeric Golgi complex component (COG2), a dynamin-related protein 5A and three Sec23/Sec24 trunk domain proteins, and it may affect the process of pollen obtaining nutrients through abnormal vesicle transport. Thus, combined with cytological observation and sequencing analysis, we thought that the changes of these genes might result in the abnormal development of pollen. It was also found some DSGs encoding MYB-related protein, UTP--glucose-1-phosphate uridylyltransferase, and ABC transporter B family member 29, as well as some DEGs encoding cytochrome P450 and ABC transporter F family member 3, and these abnormal changes may affect sporopollenin synthesis and formation of the pollen wall (Chang et al., 2016).

Putative cytoskeleton-related and AS mediated pollen sterile network in PTGMS wheat

According to the putative functions and changes in the DSGs, DEGs, DE-TFs, and lncRNA and their experimental verification in the present study, we propose an intriguing cytoskeleton related transcriptome and AS response mediated pollen sterile network for PTGMS wheat, as shown in Fig. 10. This network has several functional components comprising the calcium regulation, vesicle trafficking, distribution of cytoskeleton and pollen development. The abnormal alternative splicing of these genes encoding kinesin-related protein and myosin as well as the down-regulated genes encoding dynein may hinder the post-translational modification of microtubules, which may further lead to the disordered distribution of microtubules in pollen. Therefore, the low temperature environment, as a signal, may activate or repress the transcription factors, lncRNA or splicing factors of Ca^{2+} and vesicle trafficking, and these in turn regulate the transcription or AS of downstream genes, which in turn disrupted the distribution of the cytoskeleton, thereby hindering pollen development, and ultimately leading to male sterility in BS366.

Conclusions

In conclusion, through the mechanistic study of the pollen sterility phenotypic change in wheat

PTGMS line BS366 by analysis combining second- and third-generation sequencing and investigations of ultrastructural, we demonstrated that Ca^{2+} and vesicle trafficking related DEGs, DSGs and lncRNA affected the assembly and deassembly as well as post-translational modification of cytoskeleton, thereby causing disorder of cytoskeleton, eventually led to pollen sterility (Fig. 10). Our study sheds new light on the underlying mechanism of how cytoskeleton contributes to male sterility in plants and the data could be used as a benchmark for future studies of the molecular mechanisms of PTGMS in other crops.

Materials and methods

Plant materials, growth conditions, and sample collection

In this study, the wheat (*Triticum aestivum* L.) PTGMS line BS366 (Bai et al. 2017) and the conventional wheat line Jing411 were used as plant materials. All plants were planted in experimental fields in Beijing (China, N 39°54', E 116°18') in plastic pots in early October and managed conventionally. The treatments for wheats and anther sample collection were performed according to Bai et al. (2017). The overall anther development period was divided into six stages: S1: pollen mother cell stage; S2: dyad stage; S3: tetrad stage; S4: uninucleate stage; S5: binucleate stage and S6: trinucleate stage as defined in Browne et al. 2018 (Browne et al., 2018). Samples from fertile and sterile conditions, were collected and named FS1, FS2, FS3, FS4, FS5, FS6 and SS1, SS2, SS3, SS4, SS5, SS6 for each developmental stage, respectively. Of which S1, S2 and S3 were for sequencing, S4, S5 and S6 were for phenotypic characterization assistant.

Phenotypic characterization at the trinucleate stage

Anthers at the trinucleate stage from fertile and sterile conditions were photographed with a Nikon E995 digital camera (Nikon, Japan) mounted on a Motic K400 dissecting microscope (Preiser Scientific, Louisville, KY, USA). To further analyze pollen fertility, the mature pollen grains were stained using I₂-KI staining and photographed with a microscope (Zeiss stemi 305). For SEM analysis, anthers in the trinucleate stage were collected, fixed in 2.5% glutaraldehyde, dehydrated, air dried in silica, coated with gold-platinum in a sputter coater, and finally examined by SEM (Hitachi S-3400N) (Yang et al., 2018).

RNA Sequencing

After confirming the anther development periods, three critical pollen fertility transformation stages (S1, S2 and S3) were selected for further sequencing analysis (Figure 1). Illumina RNA seq and Iso-Seq library were constructed using the method of Wang et al. (2019). The RNAs of 36 samples (three stages of two cultivars in two different conditions, three biological replicates per stage of two cultivars) were subjected to 150bp paired-end sequencing using HiSeq X Ten platform (Illumina), and then the RNAs of 18 samples from each cultivars were mixed in equal concentration and sequenced on the PacBio RS II platform. Sequencing were performed according to the

479 manufacturer's standard protocol.

480 **Identification of full length transcripts**

481 Raw data obtained from Illumina sequencing were processed and filtered by Illumina pipeline
 482 (<https://www.illumina.com/>) to generated FastQ files. Raw data obtained from PacBio sequencing
 483 were processed using SMRT Pipe analysis workflow of the PacBio SMRT Analysis software suite
 484 (<https://www.pacb.com/products-and-services/analytical-software/smrt-analysis/>). Raw
 485 polymerase reads were filtered and trimmed to generate the ik, subreads and read of inserts (ROIs),
 486 requiring a minimum polymerase read length of 50 bp, a minimal read score of 0.65, a minimum
 487 subread length of 50 bp, Iso-seq pipeline with minFullPass of 0 and a minimum predicted accuracy
 488 of 0.8. Next, full-length, non-chemiric (FLNC) transcripts were determined by searching for the
 489 polyA tail signal and the 5' and 3' cDNA primers in ROIs. ICE (Iterative Clustering for Error
 490 Correction) was used to obtain consensus isoforms and full-length (FL) consensus sequences from
 491 ICE was polished using Quiver. High quality FL transcripts were classified with the criteria post-
 492 correction accuracy above 99%. Then, FL consensus sequences were mapped to reference genome
 493 using Genomic Mapping and Alignment Program (GMAP) using parameters 'cross-species -allow-
 494 close-indels0' and filtered for 99% alignment coverage and 85% alignment identity (Wu and
 495 Watanabe, 2005). Here, 5' difference was not considered when collapsing redundant transcripts.
 496 Integrity assessment for transcripts with no redundant using BUSCO (Simao et al., 2015).

497 **Fusion transcript delectation**

498 The fusion candidates were detected using criteria that: a single transcript must: 1) map to 2 or more
 499 loci, 2) minimum coverage for each loci is 5% and minimum coverage in bp is at least 1 bp, and
 500 total coverage is at least 95% and 3) distance between the loci is at least 10kb.

501 **AS detection and fertility related AS event identification**

502 Transcripts were validated against known reference transcript annotations with the python library
 503 MatchAnnot. In this study, AS events including ES, IR, A5'SS, A3'SS and MEX were detected and
 504 quantified using rMATs and AStalavista tool (version 3.0) (<http://astalavista.sammeth.net/>) (Foissac
 505 and Sammeth, 2007). Candidate splicing event was calculated using reads mapped to splicing
 506 junctions. Differential splicing genes under FS1, FS2, and FS3 compared with SS1, SS2, and SS3,
 507 respectively, were selected with FDR ≤ 0.05 .

508 **Identification of lncRNA**

509 The full-length transcripts were aligned to genome of *Triticum aestivum* L. from *Ensembl Plants*
 510 database (<http://plant.ensembl.org/index.html>). Those which could not be aligned were considered
 511 as novel transcripts. The novel transcripts (>200bp) were processed to identify lncRNAs based on
 512 four computational approaches include Coding Potential Calculator (CPC) (Kong et al., 2007),
 513 Coding-Non-Coding Index (CNCI) (Sun et al., 2013), Coding Potential Assessment Tool (CPAT)

(Wang et al., 2013) and Pfam. The lncRNA from intersection of these four computational approaches were used further analysis.

Identification of the transcription factors

The TFs were identified based on the domains of known TFs in the plant transcription factor database PlnTFDB 3.0 (<http://plntfdb.bio.uni-potsdam.de/v3.0/>). The domains of the protein corresponding to the newly identified transcripts in our analysis and the annotated transcripts in the IWGSC RefSeq v1.0 (generated from both high-confidence genes and low confidence genes) were searched against the included domains and excluded domains of each TF in the PlnTFDB database using the hmm search function of the HMMER software, and only proteins with exactly the same included domains and not with the excluded domains were regarded as TFs. All TFs were against DEGs to confirm the fertility-related TFs.

Cytological observation

For the transmission electron microscopy (TEM) observation, anthers were fixed, embedded, and stained as described by Zhang et al. (2014). The ultrathin sections were observed and obtained with transmission electron microscope (Hitachi, H-7650, Tokyo, Japan) and an 832 charge-coupled device camera (Gatan, Abingdon, VA, USA). In pollen cells, microfilaments and microtubules were marked by tetramethylrhodamine isothiocyanate (TRITC)- phalloidin (Sigma, St. Louis, MO, USA) and anti- α -tubulin (mouse IgG monoclonal anti- α -tubulin, T-9026; Sigma), respectively. The staining procedures were the same as those described by Wang et al. (2018). For DNA staining, 4', 6-diamidino-2-phenylindole (DAPI) was used for counterstaining. The DAPI staining procedures were the same as those described by Li et al. (2019). Preparations were observed and images were captured using a laser scanning confocal microscope (Nikon A1R, Tokyo, Japan).

Data access

The data reported in this article have been deposited in the National Genomics Data Center (NGDC) Genome Sequence Archive (GSA) database under the BioProject accession no PRJCA002516. (<https://bigd.big.ac.cn/>).Dr

Acknowledgments

We are grateful to Dr Feng Xu for his helpful suggestions and bioinformatics analysis assistance.

Figure and table Legends

Figure 1: Phenotypes of mature anthers and pollen of Jing411 (A, B) and BS366 (C, D) at the trinucleate stage under fertile and sterile conditions. Scale bars in anther are equivalent to 1 mm, in epidermis, ubisch bodies and pollen are equivalent to 50µm. Abbreviations: epidermis (E), ubisch bodies (Uby).

Figure 2: Summary of the direct RNA sequencing data of BS366. A-C: The bubble scatter plots show the relationship between the fraction of detected transcripts by the direct RNA sequencing with the transcript length and the level transcript expression. The violin-boxplots on the right show the overall distribution of the expression of transcripts. D: The histogram plot shows the distribution of read length of high quality reads obtained from BS366 (red), Jing411 (green) and IWGSC, respectively.

Figure 3: CIRCOS visualization of different data at the genome-wide level. The density was calculated in a 10-Mb sliding window

A: Karyotype of the wheat genome.

B: Comparison of transcript density between the IWGSC RefSeq v1.0 annotation and the PacBio data. From the upper to lower tracks: transcripts in IWGSC RefSeq v1.0, transcripts in BS366 and 411, transcripts in BS366 and 411 in pollen, respectively.

C-E: Distribution of differentially expressed genes (DEGs) and differentially spliced genes (DSG) for S3, S2 and S1 of BS366 and Jing411 in between fertile and sterile conditions. From the upper to lower tracks in each part: DEGs for BS366 in between fertile and sterile conditions, DSGs for BS366 in between fertile and sterile conditions, DEGs for Jing411 in between fertile and sterile conditions, DSGs for Jing411 in between fertile and sterile conditions.

F: Distribution of transcription factors in BS366 (upper track) and Jing411 (lower track).

G-K: Identified of lncRNAs from pfam (G), CPC (H), CPAT (I), CNCI (J) and overlap of them (K).

L: Linkage of fusion transcripts in BS366(red) and Jing411(blue).

Figure 4: Identification and comparison analysis of sterility-related AS genes and sterility-related genes during fertility transition. A: The changes of gene number of DSGs and DEGs during anther development stages.

B-E: Venn diagram of DSGs (B) and DEGs (C) in three stages, genes in DSGs and DEGs (D), and genes in common DSGs and common DSGs &DEGs (E)

F: qPCR analysis of seven genes of common DSGs &DEGs

Figure 5: Analysis male sterility-related DSGs and DEGs. **A:** Heat map for pollen sterility-related DSGs. **B:** Hierarchical cluster tree showing the modules of co-expressed genes, where the lower panel shows the Modules in different colors. **C:** Module-trait correlations and corresponding *p*-values (in parentheses), where the left panel shows the module eigen genes and the right panel shows a color scale for the module trait correlations ranging from -1 to 1. **D:** Cytoscape representation of the co-expressed genes in important pathways in the red module. **E:** Heat map for male sterility-related DEGs.

Figure 6: Differentially expressed TFs (DE-TFs) and differentially spliced TFs (DS-TFs) in different anther development stages

Figure 7: Analysis of identified lncRNAs. **A:** Identified lncRNAs from Pacbio data by using CPC, CNCI, CPAT and Pfam. **B:** Go enrichment analysis of targets of lncRNAs. **C:** Heat map for male sterility-related targets of lncRNA. **D:** qPCR analysis the expression of randomly selected lncRNAs and targets.

Figure 8: Validation of full-length isoforms using Semi-quantitative RT-PCR. RT-PCR validation of AS events for three genes. Gel bands in each figure show DNA makers and PCR results in three stages under two condition. Transcript structure of each isoform is shown in right panel. Yellow boxes show exons and lines with arrows show introns. PCR primers (F, forward and R, reverse) are shown on the first isoform of each gene. The length of each full-length isoform is shown after the transcript structure.

Figure 9: Cytological observation of BS366 under different conditions. **A:** the distribution of cytoskeleton of different conditions from pollen mother stage to terad stage. **B:** the distribution of Ca^{2+} of BS366 under different conditions from pollen mother stage to terad stage. **C:** the ultrastructural observation of anther, tapletum and pollen cell of BS366 under different conditions from pollen mother stage to trinucleate stage. FS1: pollen mother cell stage of fertile condition, FS2: dyad stage of fertile condition, FS3: tetrad stage of fertile condition, FS4: uninucleate stage of fertile condition, FS5: binucleate stage of fertile condition, FS6: trinucleate stage of fertile condition, SS1: pollen mother cell stage of sterile condition, SS2: dyad stage of sterile condition, SS3: tetrad stage of sterile condition, SS4: uninucleate stage of sterile condition, SS5: binucleate stage of sterile condition, SS6: trinucleate stage of sterile condition. Dd: dyad, E: epidermis, En: endothecium, PMC: pollen mother cell, T: tapetum, Td: tetrads. Bars are 4 μ m in A and 1 μ m in B and C.

Figure 10: Proposed a cytoskeleton related transcriptome and AS response mediated regulation networks and the signaling pathway involved in male sterility of PTGMS wheat line BS366.

Low temperature activates or repress transcription factors, lncRNA or splicing factors of Ca^{2+} and vesicle trafficking and these in turn regulate the transcription or AS of downstream genes, which in turn disrupted the distribution of the cytoskeleton, thereby hindering pollen development, and ultimately leading to male sterility in wheat PTGMS line BS366. lncRNA, long non coding

611 RNA; TF, transcription factor; SF, splicing factors; DEG, differentially expressed gene; DSG,
612 differentially spliced gene.

613 **Figure S1:** Flowchart of sample collection and RNA-sequencing analysis. The anthers with three
614 stages including pollen mother cell stage, dyad and tetrad stage, were taken from middle of spikes
615 from two condition in both BS366 and Jing411. In total, 36 samples (three stages for each of the
616 two varieties in two conditions, three biological replicates per stage) were sequenced using
617 second-generation sequencing, and two mixed samples (the RNAs of 18 samples from each
618 variety mixed in equal volume) were sequenced using third-generation sequencing

619 **Figure S2:** Summary of the direct RNA sequencing data of BS366 and Jing411. The bubble scatter
620 plots show the relationship between the fraction of detected transcripts by the direct RNA
621 sequencing with the transcript length and the level transcript expression. The violin-boxplots on
622 the right show the overall distribution of the expression of transcripts.

623 **Table S1:** Primers used in this study

624 **Table S2:** Summary information of circular consensus sequence reads

625 **Table S3:** Summary Statistics of Full-length non-chimeric and isform

626 **Table S4:** Summary Statistics of transcript and gene loci

627 **Table S5:** Statistics of alternative splicing events in different anther development stages during
628 fertility transition in BS366 and Jing411

629 **Table S6:** GO analysis of pollen sterile related DAS, DSGs and targets of lncRNA

630 **Table S7:** The seed setting rate of BS366 and Jing 411 in different conditions

631 **Table S8:** Identificatied TFs from Pacbio data

632 **Table S9:** Statistics of differentially expressed TFs (DE-TFs) and differentially spliced TFs (DS-
633 TFs) in different anther development stages

634 **Table S10:** lncRNA and their corresponding targets

635 **Table S11:** Annotation of targets of lncRNA

636

LITERATURE CITED

- Abdel-Ghany SE, Hamilton M, Jacobi JL, Ngam P, Devitt N, Schilkey F, Ben-Hur A, Reddy ASN** (2016) A survey of the sorghum transcriptome using single-molecule long reads. *Nat Commun* **7**: 11
- Bai JF, Wang YK, Wang P, Duan WJ, Yuan SH, Sun H, Yuan GL, Ma JX, Wang N, Zhang FT, Zhang LP, Zhao CP** (2017) Uncovering male fertility transition responsive mirna in a wheat photo-thermosensitive genic male sterile line by deep sequencing and degradome analysis. *Front Plant Sci* **8**: 1370
- Browne RG, Iacuone S, Li SF, Dolferus R, Parish RW** (2018) Anther Morphological Development and Stage Determination in *Triticum aestivum* L. *Front Plant Sci* **9**: 13
- Capovilla G, Pajoro A, Immink RGH, Schmid M** (2015) Role of alternative pre-mRNA splicing in temperature signaling. *Curr Opin Plant Biol* **27**: 97-103
- Chang Z, Chen Z, Wang, N, Xie G, Lu J, Yan W, Zhou J, Tang X, Deng X** (2016) Construction of a male sterility system for hybrid rice breeding and seed production using a nuclear male sterility gene. *Proc Natl Acad Sci USA* **113**: 14145-14150
- Chang ZY, Jin MN, Yan W, Chen H, Qiu SJ, Fu S, Xia JX, Liu YC, Chen ZF, Wu JX, Tang XY** (2018) The ATP-binding cassette (ABC) transporter OsABCG3 is essential for pollen development in rice. *Rice* **11**: 15
- Chivasa S, Tome DFA, Slabas AR** (2013) UDP-Glucose pyrophosphorylase is a novel plant cell death regulator. *J Proteome Res* **12**: 1743-1753
- Foissac S, Sammeth M** (2007) ASTALAVISTA: dynamic and flexible analysis of alternative splicing events in custom gene datasets. *Nucleic Acids Res* **35**: W297-W299
- Hamlin JNR, Schroeder LK, Fotouhi M, Dokainish H, Ioannou MS, Girard M, Summerfeldt N, Melancon P, McPherson PS** (2014) Scyl1 scaffolds class II Arfs to specific subcomplexes of coatomer through the gamma-COP appendage domain. *J Cell Sci* **127**: 1454-1463
- Han J, Pluhackova K, Boeckmann RA** (2017) The Multifaceted Role of SNARE Proteins in Membrane Fusion. *Front Physiol* **8**: 00005
- Helper PK, Callaham DA** (1987) Free Calcium Increases during Anaphase in Stamen Hair Cells of *Tradescantia*. *J Cell Biol* **105**: 2137-2143
- Heo JB, Sung S** (2011) Vernalization-Mediated Epigenetic Silencing by a Long Intronic Noncoding RNA. *Science* **331**: 76-79
- Jena BP** (2011) Role of SNAREs in Membrane Fusion. *In* T Dittmar, KS Zanker, eds, *Cell Fusion in Health and Disease: I: Cell Fusion in Health*, Vol 713, pp 13-32
- Jing JC, Wang B, Liu PJ** (2019) The Functional Role of SEC23 in Vesicle Transportation, Autophagy and Cancer. *Int J Biol Sci* **15**: 2419-2426
- Kawamura E, Himmelsbach R, Rashbrooke MC, Whittington AT, Gale KR, Collings DA, Wasteney GO** (2006) MICROTUBULE ORGANIZATION 1 regulates structure and function of microtubule arrays during mitosis and cytokinesis in the *Arabidopsis* root. *Plant Physiol* **140**: 102-114
- Kawamura E, Wasteney GO** (2008) MOR1, the *Arabidopsis thaliana* homologue of *Xenopus* MAP215, promotes rapid growth and shrinkage, and suppresses the pausing of microtubules in vivo. *J Cell Sci* **121**: 4114-4123

679 **Kong L, Zhang Y, Ye ZQ, Liu XQ, Zhao SQ, Wei L, Gao G** (2007) CPC: assess the protein-coding
680 potential of transcripts using sequence features and support vector machine. *Nucleic Acids Res*
681 **35**: W345-W349

682 **Kornienko AE, Guenzl PM, Barlow DP, Pauler FM** (2013) Gene regulation by the act of long non-
683 coding RNA transcription. *BMC Biology* **11**: 59

684 **Laloum T, Martin G, Duque P** (2018) Alternative splicing control of abiotic stress responses. *Trends*
685 *Plant Sci* **23**: 140-150

686 **Li W, Liu Z, Meng C, Jia Y, Zhang L, Song X** (2019) Identification of genes related to the regulation
687 of anther and pollen development in Mu-type cytoplasmic male sterile wheat (*Triticum aestivum*
688 L.) by transcriptome analysis. *Crop and Pasture Sci* **70**: 306

689 **Liu H, Tan M, Yu H, Li L, Zhou F, Yang M, Zhou T, Zhao Y** (2016) Comparative transcriptome
690 profiling of the fertile and sterile flower buds of a dominant genic male sterile line in sesame
691 (*Sesamum indicum* L.). *BMC Plant Biol* **16**

692 **Liu Z, Li S, Li W, Liu Q, Zhang L, Song X** (2020) Comparative transcriptome analysis indicates that
693 a core transcriptional network mediates isonuclear alloplasmic male sterility in wheat (*Triticum*
694 *aestivum* L.). *BMC Plant Biol* **20**: 10

695 **Liu Z, Shi X, Li S, Hu G, Zhang L, Song X** (2018a) Tapetal-delayed programmed cell Death (PCD)
696 and oxidative stress-induced male sterility of aegilops uniaristata cytoplasm in wheat. *Int J Mol*
697 *Sci* **19**: 1708

698 **Liu Z, Shi X, Li S, Zhang L, Song X** (2018b) Oxidative stress and aberrant programmed cell death are
699 associated with pollen abortion in isonuclear alloplasmic male-sterile wheat. *Front plant sci* Vol
700 **9**, p 595

701 **Liu ZS, Qin JX, Tian XJ, Xu SB, Wang Y, Li HX, Wang XM, Peng HR, Yao YY, Hu ZR, Ni ZF, Xin**
702 **MM, Sun QX** (2018c) Global profiling of alternative splicing landscape responsive to drought,
703 heat and their combination in wheat (*Triticum aestivum* L.). *Plant Biotechnol J* **16**: 714-726

704 **Longin C, Mühleisen J, Maurer H, Zhang H, Gowda M, Reif J** (2012) Hybrid breeding in autogamous
705 cereals. *Theor Appl Genet* **125**: 1087-1096

706 **Marianne BS, Reid GP, Harry TH** (2002) Microscopy of a cytoplasmic male-sterile soybean from an
707 interspecific cross between Glycine max and G. soja (*Leguminosae*). *Am J bot* **89**: 417-426

708 **Marquez Y, Brown JWS, Simpson C, Barta A, Kalyna M** (2012) Transcriptome survey reveals
709 increased complexity of the alternative splicing landscape in *Arabidopsis*. *Genome Res* **22**:
710 1184-1195

711 **Meng L, Liu Z, Zhang L, Hu G, Song X** (2016) Cytological characterization of a thermo-sensitive
712 cytoplasmic male-sterile wheat line having K-type cytoplasm of *Aegilops kotschyi*. *Breeding*
713 *Sci* **66**: 752-761

714 **Sane AP, Nath P, Sane PV** (1997) Differences in kinetics of F1-ATPases of cytoplasmic male sterile,
715 maintainer and fertility restored lines of sorghum. *Plant Sci* **130**: 19-25

716 **Schubert R, Dobritsch S, Gruber C, Hause G, Athmer B, Schreiber T, Marillonnet S, Okabe Y,**
717 **Ezura H, Acosta IF, Tarkowska D, Hause B** (2019) Tomato MYB21 acts in ovules to mediate
718 jasmonate-regulated fertility. *Plant Cell* **31**: 1043-1062

719 **Simao FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM** (2015) BUSCO: assessing
720 genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* **31**:
721 3210-3212

722 **Singh B, Khurana P, Khurana JP, Singh P** (2018) Gene encoding vesicle-associated membrane
723 protein-associated protein from *Triticum aestivum* (TaVAP) confers tolerance to drought stress.
724 *Cell Stress Chaperones* **23**: 411-428

725 **Sun L, Luo HT, Bu DC, Zhao GG, Yu KT, Zhang CH, Liu YN, Chen RS, Zhao Y** (2013) Utilizing
726 sequence intrinsic composition to classify protein-coding and long non-coding transcripts.
727 *Nucleic Acids Res* **41**: 8

728 **Syed NH, Kalyna M, Marquez Y, Barta A, Brown JWS** (2012) Alternative splicing in plants - coming
729 of age. *Trends in Plant Sci* **17**: 616-623

730 **Tang ZH, Zhang LP, Xu CG, Yuan SH, Zhang FT, Zheng YL, Zhao CP** (2012) Uncovering small
731 rna-mediated responses to cold stress in a wheat thermosensitive genic male-sterile line by deep
732 sequencing. *Plant Physiol* **159**: 721-738

733 **Tang ZH, Zhang LP, Yang D, Zhao CP, Zheng YL** (2011) Cold stress contributes to aberrant
734 cytokinesis during male meiosis I in a wheat thermosensitive genic male sterile line. *Plant Cell*
735 *and Environ* **34**: 389-405

736 **Tian HQ, Kuang A, Musgrave ME, Russell SD** (1998) Calcium distribution in fertile and sterile anthers
737 of a photoperiod-sensitive genic male-sterile rice. *Planta* **204**: 183-192

738 **van Riel WE, Rai A, Bianchi S, Katrukha EA, Liu Q, Heck AJR, Hoogenraad CC, Steinmetz MO,**
739 **Kapitein LC, Akhmanova A** (2017) Kinesin-4 KIF21B is a potent microtubule pausing factor.
740 *Elife* **6**

741 **Wang B, Tseng E, Regulski M, Clark TA, Hon T, Jiao YP, Lu ZY, Olson A, Stein JC, Ware D** (2016)
742 Unveiling the complexity of the maize transcriptome by single-molecule long-read sequencing.
743 *Nat Commun* **7**: 13

744 **Wang L, Park HJ, Dasari S, Wang SQ, Kocher JP, Li W** (2013) CPAT: Coding-Potential Assessment
745 Tool using an alignment-free logistic regression model. *Nucleic Acids Res* **41**: 7

746 **Wang X, Chen S, Shi X, Liu D, Zhao P, Lu Y, Cheng Y, Liu Z, Nie X, Song W, Sun Q, Xu S, Ma C**
747 (2019) Hybrid sequencing reveals insight into heat sensing and signaling of bread wheat. *Plant*
748 *J* **98**: 1015-1032

749 **Wang YK, Bai JF, Wang P, Duan WJ, Yuan SH, Zhang FT, Gao SQ, Liu LH, Pang BS, Zhang LP,**
750 **Zhao CP** (2018) Comparative transcriptome analysis identifies genes involved in the regulation
751 of the pollen cytoskeleton in a genic male sterile wheat line. *Plant Growth Regul* **86**: 133-147

752 **Wu TD, Watanabe CK** (2005) GMAP: a genomic mapping and alignment program for mRNA and EST
753 sequences. *Bioinformatics* **21**: 1859-1875

754 **Wu Y, Min L, Wu Z, Yang L, Zhu L, Yang X, Yuan D, Guo X, Zhang X** (2015) Defective pollen wall
755 contributes to male sterility in the male sterile line 1355A of cotton. *Sci Rep* **5**: 9608

756 **Xu H, Heath MC** (1998) Role of calcium in signal transduction during the hypersensitive response
757 caused by basidiospore-derived infection of the cowpea rust fungus. *Plant Cell* **10**: 585-598

758 **Xu SX, Liu XD, Feng JH, Lu YG** (2001) Comparative studies on the changes of microtubule
759 distribution and reorganization during the meiotic stages of development in normal (IR36) and
760 a temperature/photoperiod sensitive male sterile line (Peiai 64S) of rice (*Oryza sativa* L.). *Acta*
761 *Botanica Sinica* **43**: 221-226

762 **Yang X, Geng X, Liu Z, Ye J, Xu M, Zang L, Song X** (2018) A Sterility Induction Trait in the Genic
763 Male Sterility Wheat Line 4110S Induced by High Temperature and its Cytological Response.
764 *Crop Sci* **58**: 1866-1876

765 **Zhang DD, Liu D, Lv XM, Wang Y, Xun ZL, Liu ZX, Li FL, Lu H** (2014) The cysteine protease
766 CEP1, a key executor involved in tapetal programmed cell death, regulates pollen development
767 in *Arabidopsis*. *Plant Cell* **26**: 2939-2961
768 **Zheng S, Li J, Ma L, Wang H, Zhou H, Ni E, Jiang D, Liu Z, Zhuang C** (2019) *OsAGO2* controls
769 ROS production and the initiation of tapetal PCD by epigenetically regulating *OsHXX1*
770 expression in rice anthers. *Proc Natl Acad Sci USA* **116**:7549-7558
771

Figure 1: Phenotypes of mature anthers and pollen of Jing411 (A, B) and BS366 (C, D) at the trinucleate stage under fertile and sterile conditions. Scale bars in anther are equivalent to 1 mm, in epidermis, ubisch bodies and pollen are equivalent to 50µm. Abbreviations: epidermis (E), ubisch bodies (Uby).

Figure 2: Summary of the direct RNA sequencing data of BS366. **A-C:** The bubble scatter plots show the relationship between the fraction of detected transcripts by the direct RNA sequencing with the transcript length and the level transcript expression. The violin-boxplots on the right show the overall distribution of the expression of transcripts. **D:** The histogram plot shows the distribution of read length of high quality reads obtained from BS366 (red), Jing411 (green) and IWGSC, respectively.

Figure 3: CIRCOS visualization of different data at the genome-wide level. The density was calculated in a 10-Mb sliding window

A: Karyotype of the wheat genome.

B: Comparison of transcript density between the IWGSC RefSeq v1.0 annotation and the PacBio data. From the upper to lower tracks: transcripts in IWGSC RefSeq v1.0, transcripts in BS366 and 411, transcripts in BS366 and 411 in pollen, respectively.

C–E: Distribution of differentially expressed genes (DEGs) and differentially spliced genes (DSG) for S3, S2 and S1 of BS366 and Jing411 in between fertile and sterile conditions. From the upper to lower tracks in each part: DEGs for BS366 in between fertile and sterile conditions, DSGs for BS366 in between fertile and sterile conditions, DEGs for Jing411 in between fertile and sterile conditions, DSGs for Jing411 in between fertile and sterile conditions.

800 **F:** Distribution of transcription factors in BS366 (uper track) and Jing411 (lower track).

801 **G-K:** Identificated of lncRNAs from pfam (**G**), CPC (**H**), CPAT (**I**), CNCI (**J**) and overlap of them
802 (**K**).

803 **L:** Linkage of fusion transcripts in BS366(red) and Jing411(blue).

804

805

Figure 4: Identification and comparison analysis of sterility-related AS genes and sterility-related genes during fertility transition. **A:** The changes of gene number of DSGs and DEGs during anther development stages.

B-E: Venn diagram of DSGs (**B**) and DEGs (**C**) in three stages, genes in DSGs and DEGs (**D**), and genes in common DSGs and common DSGs & DEGs (**E**)

F: qPCR analysis of seven genes of common DSGs & DEGs

Figure 5: Analysis male sterility-related DSGs and DEGs. **A:** Heat map for pollen sterility-related DSGs. **B:** Hierarchical cluster tree showing the modules of co-expressed genes, where the lower panel shows the Modules in different colors. **C:** Module-trait correlations and corresponding *p*-values (inparentheses), where the left panel shows the module eigen genes and the right panel shows a color scale for the module trait correlations ranging from -1 to 1. **D:** Cytoscaper representation of the co-expressed genes in important pathways in the red module. **E:** Heat map for male sterility-related DEGs.

Figure 6: Differentially expressed TFs (DE-TFs) and differentially spliced TFs (DS-TFs) in different anther development stages

Figure 7: Analysis of identified lncRNAs. **A:** Identified lncRNAs from Pacbio data by using CPC, CNCI, CPAT and Pfam. **B:** Go enrichment analysis of targets of lncRNAs. **C:** Heat map for male sterility-related targets of lncRNA. **D:** qPCR analysis the expression of randomly selected lncRNAs and targets.

Figure 8: Validation of full-length isoforms using Semi-quantitative RT-PCR. RT-PCR validation of AS events for three genes. Gel bands in each figure show DNA makers and PCR results in three stages under two condition. Transcript structure of each isoform is shown in right panel. Yellow boxes show exons and lines with arrows show introns. PCR primers (F, forward and R, reverse) are shown on the first isoform of each gene. The length of each full-length isoform is shown after the transcript structure.

Figure 9: Cytological observation of BS366 under different conditions. **A:** the distribution of cytoskeleton of different conditions from pollen mother stage to terad stage. **B:** the distribution of Ca^{2+} of BS366 under different conditions from pollen mother stage to terad stage. **C:** the ultrastructural observation of anther, tapletum and pollen cell of BS366 under different conditions from pollen mother stage to trinucleate stage. FS1: pollen mother cell stage of fertile condition, FS2: dyad stage of fertile condition, FS3: tetrad stage of fertile condition, FS4: uninucleate stage of fertile condition, FS5: binucleate stage of fertile condition, FS6: trinucleate stage of fertile condition, SS1: pollen mother cell stage of sterile condition, SS2: dyad stage of sterile condition, SS3: tetrad stage of sterile condition, SS4: uninucleate stage of sterile condition, SS5: binucleate stage of sterile condition, SS6: trinucleate stage of sterile condition. Dd: dyad, E: epidermis, En: endothecium, PMC: pollen mother cell, T: tapetum, Td: tetrads. Bars are 4 μm in A and 1 μm in B and C.

Figure 10: Proposed a cytoskeleton related transcriptome and AS response mediated regulation networks and the signaling pathway involved in male sterility of PTGMS wheat line BS366.

Low temperature activates or repress transcription factors, lncRNA or splicing factors of Ca²⁺ and vesicle trafficking and these in turn regulate the transcription or AS of downstream genes, which in turn disrupted the distribution of the cytoskeleton, thereby hindering pollen development, and ultimately leading to male sterility in wheat PTGMS line BS366. lncRNA, long non coding RNA; TF, transcription factor; SF, splicing factors; DEG, differentially expressed gene; DSG, differentially spliced gene.