Signalome-wide assessment of erythrocyte response to *Plasmodium* reveals novel targets for host-directed antimalarial intervention

Jack D. Adderley¹, Simona John von Freyend², Sarah A. Jackson², Megan J. Bird², Amy L. Burns³, Burcu Anar⁴, Tom Metcalf⁴, Jean-Philippe Semblat⁵, Oliver Billker^{4,6}, Danny W. Wilson^{3,7}, Christian Doerig¹

¹Centre for Chronic Infectious and Inflammation Disease, Biomedical Sciences Cluster, School of Health and Biomedical Sciences, RMIT University, Bundoora VIC 3083, Australia

²Infection and Immunity Program, Biomedicine Discovery Institute and Department of Microbiology, Monash University, Clayton VIC 3800, Australia

³Research Centre for Infectious Diseases, School of Biological Sciences, University of Adelaide, Adelaide SA 5005, Australia

⁴Wellcome Trust Sanger Institute, Hinxton, Cambridgeshire, CB10 1SA, United Kingdom.

⁵Institut National de la Transfusion Sanguine, Inserm UMR S1134, 75015 Paris, France

⁶Molecular Infection Medicine Sweden (MIMS), Department of Molecular Biology, Umeå University, Umeå, SE-901 87, Sweden.

⁷Burnet Institute, Melbourne, Victoria, Australia 3004

Abstract

Intracellular pathogens are known to mobilise host signaling pathways to manipulate gene expression in their host cell to promote their own survival. Surprisingly, evidence is emerging that specific signal transduction elements are activated in a-nucleated erythrocytes in response to infection with malaria parasites, but the extent of this phenomenon remains unknown. Here, we fill this knowledge gap by providing a comprehensive and dynamic assessment of host erythrocyte signaling during the course of infection with *Plasmodium falciparum*. We used an antibody microarray that comprises 878 antibodies directed against human signaling proteins, >600 of which are phospho-specific, to interrogate the status of host erythrocyte signaling pathways at three stages of parasite development during the asexual cycle. This confirmed the pre-existing fragmentary data on the activation of a PAK-MEK pathway, and revealed the modulation of a large number of additional signaling elements during infection.

We focussed on the receptor tyrosine kinase c-MET, also known as the hepatocyte growth factor receptor, and the MAP kinase pathway component B-Raf that is reported to lie downstream of c-MET in a number of cell types. Array data validated by Western blotting revealed that activation sites of c-MET are phosphorylated in trophozoite-infected erythrocytes, and we show that treatment of parasite cultures with c-MET or B-Raf selective inhibitors have nanomolar potency against *in vitro* proliferation of *P. falciparum* and the phylogenetically distant species *P. knowlesi*. Furthermore, we demonstrate that a c-MET inhibitor impairs *in vivo* proliferation of the rodent malaria parasite *P. berghei* in mice.

Overall, we provide a comprehensive dataset on the modulation of host erythrocyte signaling during infection with malaria parasites, as well as a proof of concept that human signaling kinases identified as activated by malaria infection represent attractive targets for antimalarial intervention.

Introduction

Malaria, caused by infection of mosquito-borne apicomplexan parasites in the genus *Plasmodium*, remains one of the most devastating infectious diseases globally. Among the six species that infect humans, *P. falciparum* is the most virulent and is responsible for the majority of malaria-related deaths. The remaining human malaria species *P. vivax*, *P. ovale*, *P. malariae* and two zoonotic species, *P. knowlesi* and *P. simium*^{1,2}, are major contributors to global malaria morbidity and must be considered in the context of new treatment strategies for malaria. Recent years have seen a 50% drop in malaria-related mortality, yet the disease still kills an estimated 445,000 people every year, mostly young children in sub-Saharan Africa³. Progress towards malaria elimination and ultimately eradication has stagnated, and the emergence of parasites resistant to the most recently deployed global front-line treatment, artemisinin-combination therapies (ACT)^{3,4}, is a major concern. It remains critical to develop novel antimalarial drugs with untapped modes of action.

Protein kinases (PKs) are core components of signaling pathways in eukaryotic cells. Phosphorylation of the target protein can lead to conformational changes and generate or mask binding motifs, and thus can affect its activity, binding properties, stability or subcellular localisation. PK activity can be regulated positively or negatively by phosphorylation of specific amino acids, or by binding to specific activator or inhibitor proteins. Activation leads to a conformational change making the active site accessible to ATP and the substrate protein. PKs are eminently druggable targets, as illustrated by the fact that > 48 small molecule kinase inhibitors have reached the market, mostly in the context of cancer chemotherapy^{5,6}. The kinomes of apicomplexan parasites and humans are quite divergent owing to the large phylogenetic distance between these organisms⁷, and parasite-encoded PKs have been proposed as attractive potential targets for selective intervention⁸. Host-directed therapy (HDT) is another avenue that is currently gaining traction to combat infectious diseases generally⁹. We and others^{10,11} have proposed that host erythrocyte (and hepatocyte) PKs represent excellent potential targets for antimalarial intervention. Many anti-infective drugs are rendered ineffective by the selection of mutations in their pathogen-encoded targets, which explains the rapid emergence of resistance across essentially all pathogen taxons. Targeting host enzymes would deprive the pathogen of this most direct pathway to resistance. Furthermore, repurposing anti-cancer

kinase inhibitors as agents against infectious diseases would greatly alleviate the lack of resources that severely affects anti-infective drug development.

Malaria pathology is caused by the blood stage of the parasite lifecycle. Blood stage malaria begins with the infection of erythrocytes by *Plasmodium* merozoites, followed by rapid growth and asexual multiplication. Newly formed merozoites are then released into the bloodstream and invade erythrocytes to begin the next replication cycle (~48 hrs for *P. falciparum*). Several human signaling molecules of the host erythrocyte have been implicated in parasite survival during infection. These include G-coupled protein receptors¹²⁻¹⁴, the protein kinases MEK (MAP/ERK kinase)¹⁵, PAK (p21-activated kinase)¹⁵, Protein kinase C (PKC)¹⁶, and peroxiredoxin¹⁷⁻¹⁹. In our study of the erythrocyte PAK-MEK pathway¹⁵, we showed that host MEK1 is phosphorylated on residue Ser298 in infected erythrocytes. pSer298 is known to cause activation of the enzyme through stimulating trans-auto-phosphorylation on its activation loop. The only PK known to phosphorylate MEK1 on Ser298 is PAK1²⁰, and indeed we showed that the latter enzyme is also activated in infected erythrocytes¹⁵. PAK isoforms are notorious for having a multitude of substrates²¹ and serve as a node to integrate signals from a number of upstream receptors and transmit these signals to several downstream effector pathways (including the mitogen-activated protein kinase pathway [MAPK] that includes the aforementioned MEK). We, therefore, hypothesised that our description of the PAK-MEK pathway addresses but a minute fraction of the signaling events that may be modulated by infection. Here, in order to comprehensively assess the host erythrocyte signaling response to infection, we use a microarray of antibodies designed to recognise human signaling proteins and their phosphorylation status. Comparison of the signals obtained from extracts of uninfected erythrocytes with those obtained from erythrocytes infected with *P. falciparum* at three main developmental stages of the replication cycle (rings, trophozoites and schizonts) allowed us to generate a comprehensive and dynamic picture of the modulation of host erythrocyte signaling by infection. This identified several host kinases as potential targets for HDT; on this basis, we further demonstrate that selective inhibitors against human c-MET and B-Raf display high potency against *P. falciparum* and *P. knowlesi* *in vitro*, and show that a c-MET inhibitor has *in vivo* activity against *P. berghei* in a murine model of malaria.

Results

Kinexus antibody microarray analysis. To investigate dynamic changes in host erythrocyte signaling during *P. falciparum* asexual proliferation, we employed an antibody microarray developed by Kinexus (Vancouver, Canada). The array consists of 878 unique antibodies, 265 of which are pan-specific, i.e. recognise both the phosphorylated and unphosphorylated forms of the target protein; the remaining 613 antibodies are phosphorylation-specific, recognising signaling molecules only if their activating or inhibitory phosphorylation sites are modified by the addition of a phosphate group. Some of the most important and well-known signaling molecules, such as members of the mitogen-activated protein kinase (MAPK) pathways or the PKC isoforms, are detected by multiple antibodies aimed at various phosphorylation sites within the same protein. The array thus provides a comprehensive picture as to how signaling mediated by these molecules changes during infection. Each array device comprises two identical chambers, each carrying two spots for each of the 878 antibodies, thus delivering each read-out in duplicate. Two sample lysates are labelled with protein-binding fluorescent dye [e.g. from *Plasmodium*-infected (iRBC) and non-infected (uRBC) red blood cells] and compared on each device, one sample per chamber. Binding of proteins to the antibodies on the array is quantitated through signal intensity for each spot.

Robustness of the approach. To evaluate the reproducibility and robustness of the approach, we compared four uninfected erythrocyte (uRBC) samples from different blood donors that were cultured for ~24 hrs under the same conditions as the iRBCs. The mean signal intensity and standard deviation for each phospho-specific antibody on the array were determined for these four samples (Fig. 1; the entire datasets are available in Supplemental dataset 1). Despite some variation, individual data points for each antibody remained within two standard deviations of their mean, illustrating that sample-to-sample variation was minimal.

Overall changes in phosphosignaling in response to infection. To assess the impact of infection on host cell signaling, whole-cell protein extracts were obtained from synchronised *P. falciparum* cultures at three time windows during the erythrocytic cycle: 4-12 hrs post-invasion (hpi) (rings, n=3), 24-28 hpi (trophozoites, n=3) and 44-48 hpi (segmented schizonts, n=2). Each of these extracts were loaded onto a separate antibody array and

compared to the mean of the uRBC lysates harvested in parallel and containing RBCs from the same donor as the iRBC sample (see above). In stark contrast to the minimal variation between the uRBC samples (see above), analysis of the iRBC extracts showed that 6% (ring-stage), 23% (trophozoite-stage) and 12% (schizont stage) of the phospho-specific antibody intensities differed by more than two standard deviations from uRBC mean. The datasets for each of the three developmental stages showed no internal variation greater than two standard deviations between the biological replicates (Supplemental Figure 1), further confirming the robustness and reliability of the approach (Fig. 1; the complete datasets are available in Supplemental Dataset 1). Overall, this implies (i) that there is little variation between biological replicates of uRBCs, attesting to the reliability of the system, and (ii) that infection with *P. falciparum* causes significant variation in the signals yielded by phospho-specific antibodies, suggesting that *P. falciparum* strongly impacts host erythrocyte phosphosignalling during infection.

Figure 1 – Distribution of Z-scores for each sample. All antibody microarray signals from each sample (four from uRBCs and the mean signal for each iRBC harboring parasites at the ring (n=3), trophozoite (n=3) and schizont stage (n=2)) were compared to the uninfected erythrocyte mean value for each signal, and the Z-score determined for each phosphorylation-specific signal. The Z-score is defined as the number of standard deviations from the uRBC mean for each sample; a positive or negative value indicates the direction of the

signal (increase or decrease, respectively) from the uRBC mean. No signals from the uRBC samples (n=4) had associated Z-scores above 2 or below -2, while the three iRBC time points showed clear variations from uRBCs.

Data filtering- removal of cross-reacting antibodies. To address possible cross-reactivity of the antibodies with parasite-derived proteins, we compared signals from purified unsynchronised parasites (pellet obtained by saponin lysis followed by centrifugation) with those of the purified erythrocyte cytoplasm (saponin supernatant). Saponin disrupts the erythrocyte membrane, thereby releasing erythrocyte cytoplasmic proteins and exported parasite proteins, while the parasite and insoluble erythrocytic material can be pelleted²². The array was loaded with 20x more (protein mass) pellet material than supernatant material, to confer high stringency to the cross-reactivity filter. A heatmap of the results is shown in Supplemental Figure 2a (full data available in Supplemental Dataset 2). Antibodies showing a fold change > 1.5 between the erythrocyte cytoplasm (saponin supernatant) and the parasite extract (saponin pellet), amounting to 224 signals (37%) of the phospho-specific antibodies, were withdrawn from further analysis (Supplemental Figure 2 and Supplemental Dataset 2). This level of cross-reactivity is not surprising, as many signaling proteins display conservation between *Homo sapiens* and *P. falciparum*, especially in their regulatory sites⁷. Some of these cross-reactive antibodies may recognise parasite-encoded orthologues of the human target proteins and thus prove to be useful tools to study *P. falciparum* signal transduction; however, this lies outside of the scope of the present study.

Data filtering - removal of low-signal antibodies. Some antibodies displayed a weak fluorescence signal, likely due to the low abundance of the target protein. The antibodies yielding a signal intensity below a fluorescence reading of 1000 units in both the erythrocyte control and parasite-infected samples were removed from further analysis, as recommended by Kinexus (Supplemental Dataset 1). This included 26 signals from the ring array, and 41 and 48 signals from the trophozoite and schizont arrays, respectively.

Broad analysis of the post-filtering iRBC dataset

Following low and cross-reactive signal removal, 1 of the ring stage signals, 29 of the trophozoite stage and 17 of the schizont stage signals had fold changes greater than 2 or

below 0.5 compared to their uninfected counterpart, revealing dynamic changes in the phosphorylation of host signaling proteins during *P. falciparum* asexual development (Fig. 2a: heatmap of all retained signals; Fig 2b: distribution of increased and decreased signals at the three development stages). The small number of changes in ring-infected cells may in part be due to the fact that these samples contained only 33% infected cells (as magnetic purification of infected cells, which allows close to 100% iRBCs for the trophozoite and schizont stages, cannot be implemented for ring stages). The observed effect on host cell phosphosignaling was considerably larger at later stages of infection (trophozoite/schizont), with the majority of these changes being attributed to increases in phosphorylation (80% and 65%, respectively). A dot plot for each of the three life stages (Fig. 2c) further illustrates this trend and highlights the strongest individual changes.

Figure 2 – Microarray data for the ring, trophozoite and schizont stages. **a)** Heat map of phosphospecific antibody array data for ring- trophozoite- and schizont-infected cells after removal of cross-reactive and weak signals. **b)** Phosphorylation-specific signals (post-filtering) ranked according to fold change. Yellow represents changes with a fold change greater than 2; blue represents changes with a fold change less than 0.5. Full datasets are presented in Supplemental Dataset 1 and 2. **c)** Dot plot highlighting differences in phosphosite signals represented as fold change. Strong changes are listed with their associated phosphorylation site.

Shortlisting of high-confidence changes in phosphosignaling. Of the reliable antibodies remaining after removal of low and cross-reactive signals, 16/345 (ring stage), 102/351 (trophozoite stage) and 35/346 (schizont stage) were significantly different ($p < 0.5$, unpaired t-test) from the uninfected erythrocyte signals. Figure 3 lists the most noteworthy changes

observed from this analysis (the full list of the significant changes is available in Supplemental Dataset 3).

Some of the most notable changes implicated the PKC isoforms δ and θ . Several PKC isoforms have previously been reported to be activated during *P. falciparum* infection of erythrocytes, but the phosphorylation status of these kinases was uncharacterised^{23,24}. Several phosphorylation sites of PKC δ / θ were significantly phosphorylated in trophozoite- and schizont-infected cells (Fig. 3; a list of the phosphorylation specific signals for PKC δ / θ is available in Supplemental Figure 3). The most striking changes were observed for Thr507 (PKC δ) and Ser695 (PKC θ). Thr507 is a regulatory residue located within the activation loop of PKC δ ²⁵ while, Ser695 is within the hydrophobic motif of PKC θ , and its phosphorylation is required for optimal kinase activity²⁶. Additionally, phosphorylation of Ser695 may also be involved in the localisation of PKC θ to the plasma membrane²⁷. These data are consistent with previous reports that host cell PKC activity is detectable during schizont development (30-40 hrs post-invasion) with limited activity before 25 hrs^{23,24}, and considerably refine these previous findings, notably with respect to dynamic phosphorylation changes on specific residues of the various PKC isoforms during parasite development.

Rings

Trophozoites

Schizonts

Protein	Phospho Site (Human)	Fold change	p value
FAK	Y397	1.80	0.043
DDR1	Y796+Y797	1.69	0.027
B-Raf	S729	1.69	0.029
SCYL1	S754	1.63	0.045
PTEN	S380+T382+T383	1.43	0.042
HSP90AB1	Y484	1.38	0.044
DDR2	Y736	1.34	0.044
CHK1	S317	1.30	0.019
JAK2	Y570	1.20	0.044
Rac1/cdc42	S71	0.65	0.010
RSK1/2	S380/S386	0.64	0.006
Tau	S713	0.58	0.050
Paxillin 1	Y31	0.56	0.012
Tau	S713	0.53	0.011
p38a MAPK	T180+Y182	0.52	0.045
Rb	S780	0.52	0.029

Protein	Phospho Site (Human)	Fold change	p value
Smad1	S465	4.56	0.0003
IKK ϵ	S172	2.88	0.007
InsR	Y1189	2.65	0.005
SCYL1	S754	2.65	0.0002
B-Raf	S729	2.53	0.020
PTEN	S380+T382+T383	2.17	0.001
RET	Y905	2.00	0.013
PKC δ	T507	1.96	0.004
PKC ζ	S695	1.90	0.006
ERK2	T185+Y187	1.85	0.001
PKC η	Y545	1.80	0.013
MET	T1241	1.72	0.023
DDR2	Y736	1.65	0.008
PKC ζ	S695	1.59	0.040
DDR2	Y740	1.55	0.002
MET	Y1234+Y1235	1.52	0.025
DDR1	Y796+Y797	1.51	0.022
MET	S1236	1.38	0.035
Paxillin 1	Y31	0.62	0.021
PTEN	S380+T382+S385	0.53	0.008
Tau	S713	0.50	0.006
p38a MAPK	T180+Y182	0.39	0.014

Protein	Phospho Site (Human)	Fold change	p value
PRKD2	S197+S198	3.56	0.005
Smad1	S465	2.66	0.008
B-Raf	S729	2.63	0.002
ERK1	Y204	2.11	0.037
EIF2AK3	T982	2.07	0.005
PKC δ	T507	2.05	0.005
DDR2	Y736	2.01	0.001
IKK ϵ	S172	2.00	0.034
InsR	Y1189	1.92	0.032
SCYL1	S754	1.84	0.040
PTEN	S380+T382+T383	1.80	0.012
DDR2	Y740	1.79	0.001
PKC ζ	S695	1.43	0.044
ERK2	T185+Y187	1.42	0.033
SMG1	T3550	0.67	0.006
ABL1	Y469	0.64	0.025
CLK1	S337	0.63	0.042
Paxillin 1	Y31	0.57	0.026
Tau	S713	0.50	0.012
ErbB3	Y1328	0.50	0.046

Figure 3 – Shortlist of the highest confidence signals (p<0.05) for rings, trophozoites and schizonts. Increases in phosphorylation compared to uninfected erythrocytes are marked in yellow; decreases are marked in blue. All values listed are the fold change from the uRBC control mean (n=4); the corresponding p-value for each signal is also listed. Each signal has a p-value <0.05, as determined through an unpaired t-test. All listed signals passed stringent parasite cross-reactivity and signal intensity filters (see text for details and Figure 2).

MAP-Kinase signaling in infected erythrocytes – B-Raf, MEK and ERK. The canonical MAPK pathway is a highly conserved, three-tiered cascade, in which a MAP kinase kinase kinase (MAP3K, or MEKK for MAP/ERK kinase kinase) phosphorylates and thereby activates a MAP2K (or MEK), which in turn phosphorylates and activates a MAP kinase. A well-understood MAPK pathway involves Raf, MEK and ERK. Both Raf isoforms are activated by Ras-GTPase binding, which enables phosphorylation of T598 and S601 (on B-Raf) and S338 and Y340 (on C-Raf) ²⁸⁻³⁰. This results in the full activation of Raf, enabling downstream phosphorylation of MEK and consequently ERK1/2 (pathway depicted in Fig. 4b). *P. falciparum* has no homologues for either Ras, Raf or MEK, but expresses two MAPKs, neither of which is a close homologue of ERK1/2 ³¹. Figure 4 lists the phosphorylation sites detected on the array that are relevant to canonical MAPK pathways (Fig. 4a) and indicates the activating phosphorylation sites for each of the three kinases involved (Fig. 4b).

Proteins in the MAPK signaling pathway showed notable changes in phosphorylation in iRBCs compared to uRBCs. B-Raf and C-Raf as the ‘gatekeeper’ proteins of this pathway showed changes in phosphorylation patterns with *P. falciparum* RBC infection. Unfortunately, the activating Ras-induced phosphorylation sites on B-Raf (T598 and S601) and C-Raf (S338 and Y340) are not represented in the array and could not be assessed. Antibodies against C-Raf S296 yielded a consistent fold change of >1.2 for all three parasite stages. Phosphorylation of C-Raf on S296 is thought to be mediated by ERK1/2 in the context of a negative feedback loop ³², which would be consistent with activation of the MAPK pathway detected here (see below).

B-Raf phosphorylation at S446 and S447, sites associated with priming B-Raf for activation by Ras ^{30,34}, increased by 1.79-fold at the ring stages, but decreased back to uRBC levels in trophozoites and schizonts. B-Raf S446 is constitutively phosphorylated in some cell lines, but this remains to be shown in cells of the erythropoietic lineage ^{30,34-36}. In contrast, phosphorylation of B-Raf at S729, a Ras-independent phosphorylation event carried out by

AMP-activated protein kinase (AMPK) in response to energy stress, increased at all three life cycle stages with a fold change ranging from 1.69 to 2.63 across parasite development (Fig 3, Fig. 4a). Of the two AMPK antibodies on the array, one was discounted due to cross-reactivity with a parasite protein, while the other (AMPKa2 S377) shows a slight increase (1.17 fold change) in phosphorylation at the trophozoite time point supporting that this kinase may also be activated in iRBCs (Supplemental Dataset 1).

We next determined the total amount of B-Raf and phosphorylated B-Raf S729 at each lifecycle stage using Western blot. Detection of total B-Raf protein using a pan-antibody independent of phosphorylation showed that the amount of full-length B-Raf was greatly reduced at trophozoite and schizont stages compared to uRBCs and rings (Fig. 4c). This fits with the reduced phosphorylation identified for S446 and S447 at trophozoites and schizonts stages. Paradoxically, while we observed increasing S729 and total B-Raf signals from rings to schizont stages on the array, this was not reflected in increased signal for full-length B-Raf on the Western Blot. Instead, the increasing signal on the array corresponded to increasing bands on the Western blot that have lower molecular mass (see Supplemental Figure 4 for additional data). Together this indicates that B-Raf is indeed phosphorylated at the ring stage of *P. falciparum* development, before degradation in the later stages of parasite development (see Discussion).

Overall, the microarray and Western blot data points to activation of B-Raf at ring-stages, and degradation at later stages. To assess if parasite requirements for host B-Raf activation were also ring-stage-specific, as suggested by the apparent degradation of B-Raf in trophozoite- and schizont-infected cells (Fig. 4c), we treated highly synchronized parasites in 4-hour windows with a potent B-Raf inhibitor (SB-590885, see below). Only when the treatment was applied at ring-stage did it display parasitocidal activity, whereas there was no effect when the cultures were treated at the trophozoite or schizont stage, in stark contrast to artemisinin, which displayed (as expected) a strong effect at the trophozoite stage (Fig. 4d).

Further supporting activation of Rafs in iRBCs, the B-Raf/C-Raf substrates MEK1/2 showed increased phosphorylation on several residues at both the ring and schizont time points (Fig. 4a). MEK1/2 phosphorylation of S298 (resulting from p21-activated kinase [PAK] activity) during the late stages of *P. falciparum* infection has previously been established by Western

blot analysis ¹⁵, aligning with the 1.35-fold increase of pS298 at the schizont stage observed in the array experiment. Other phospho-sites such as T292 and S218+S222 (MEK1/2) show a signal increase in schizont-infected red cells. MEK1 T292 is involved in negative feedback of the kinase ³⁷, while S218+S222 and S222+S226 are the activation loop phosphosites in MEK1 and MEK2, respectively ^{38,39}. All the MEK1/2 phospho-specific antibodies, as well as the ERK1 (Y204) antibody, showed some cross-reactivity with parasite protein (underlined in Fig. 4a), which makes the overall MEK1/2 data difficult to interpret (Fig. 4a).

MEK1/2 phosphorylate ERK1 and ERK2 (on T202- 204 for ERK1, on T185 -Y187 for ERK2), and these residues are the only known targets of MEK1 activity ^{40,41}. ERK2 is slightly more phosphorylated in ring-infected cells than in uninfected erythrocytes (fold change = 1.09); at the trophozoite stage it has increased further (fold change = 1.85), and at the schizont stage ERK2 phosphorylation is still elevated (fold change = 1.42). The array results for the ERK1 activation sites (T202 and Y204) at the ring-stage of *P. falciparum* development are however difficult to interpret as these signals indicate increases as well as decreases. ERK1 T207 is a known autophosphorylation regulatory site and showed a fold change value of 1.25 in trophozoite-infected red cells-, supporting that regulation of ERK1 activity occurs during infection ⁴². These data support that *P. falciparum* iRBCs modify the host cell Raf/Mek/ERK pathway during growth and that the activity of RBC B-Raf is required for parasite maturation beyond ring stages.

Figure 4 – Activation of a MAPK pathway in infected erythrocytes. **a)** Summary table of the MAPK pathway components and phosphorylation sites covered by the antibody microarray, including RAF, MEK1/2 and ERK1/2 with the fold change from the uninfected erythrocyte control signal for each of the three time points (erythrocytes infected with rings, trophozoites or schizonts) shown. Signals which were significantly different during infection are marked by * ($p < 0.05$, unpaired t-test). Phosphosites underlined in the table were flagged as cross-reactive with parasite material in Fig. 2, but are included here for completeness. **b)** Schematic representation of the MAPK pathway with the relevant phosphorylation sites present in the array data set covered in **a**. Green 'P' represents phosphorylation sites associated with the activity of the respective kinase, while the red 'P's represent negative feedback phosphorylation sites. The grey 'P' represents a known priming site on B-Raf. **c)** Western blot validation of B-Raf phosphorylation. A phospho-specific antibody to B-Raf (S729) detected phosphorylated B-Raf in ring-stage infected erythrocytes (upper panel). A pan-B-Raf antibody detected B-Raf in the uninfected and ring-stage infected erythrocyte samples, with a strong decrease at the trophozoite and schizont stages (middle panel). An antibody against Glycophorin-C was used as a loading control (lower panel). **d)** Synchronised *P. falciparum* cultures were treated in 4-hr windows with 2.45 μ M SB590885 (5x IC_{50} , see Fig 6), 0.091 μ M artemisinin or DMSO (vehicle control), starting at 0 (rings), 22 (trophozoites) and 38 (schizonts) hours post-invasion (the black arrows indicate the end of treatment, where the medium was replaced with fresh medium without drug). The parasitemia was then monitored until the next cycle of parasite development to assess parasite survival. Starting parasitemia was normalised to 5% across the biological replicates ($n=3$) error bars represent the standard error of the mean.

c-MET signaling in infected erythrocytes. Following the identification of the MAPK pathway activation, we focused our attention on signaling components that lay upstream of the MAPK pathway and found that the receptor tyrosine kinase c-MET, which is known to signal to MAPK pathways in other systems⁴³, was clearly activated by infection (Fig. 3). C-MET is expressed in many cells of mesenchymal origins^{44,45}, and its activation occurs through the binding of hepatocyte growth factor (HGF) to the extracellular domain, prompting homodimerization and trans-autophosphorylation and activation⁴⁶. C-MET controls several key downstream pathways including (but not limited to) the MAPK pathway (proliferation, motility and cell cycle progression) and the phosphatidylinositol 3-kinase (PI3K)/AKT pathway (cell survival)⁴³. There is no homologue of c-MET (or any member of the Tyrosine kinase group) in *P. falciparum*⁷.

The array contained a total of nine phospho-specific antibodies targeting c-MET, 7 of which passed cross-reactivity and low-intensity signal thresholds (Fig. 5a). There was no noteworthy change in the phosphorylation of any of these residues at the ring stage. In contrast, five out of the seven sites show an increase in phosphorylation in trophozoite-

infected cells (fold change >1.38, Fig. 5a). The signals from two antibodies against c-Met phosphorylation sites decreased at the trophozoite stage. One of these is directed against Y1230+Y1234+Y1235; this antibody is affected by high error between duplicates (848% at the trophozoite stage) and was deemed unreliable. The other antibody recognises pY1230 and is the only reliable c-MET antibody to show a decrease of phosphorylation at the trophozoite stage. At the schizont stage, most sites were slightly higher than the uninfected control. Two unique pan-specific c-MET antibodies were also present on the array. Both pan-specific antibodies showed minimal signal variation between samples, indicating that the level of c-MET remained relatively constant as expected (Fig 5a).

Antibodies against c-MET S975 and Y1349+Y1356 were not present on the array or removed from analysis because of low-intensity signal, respectively. Reliable antibodies were available for pY1230 and pY1234+pY1235, which lie within the c-MET catalytic domain and are phosphorylated in response to binding of HGF, activating the kinase⁴⁷. While increased phosphorylation was evident for Y1234+Y1235 across lifecycle stages, with a peak signal seen at trophozoites, this occurred only at schizont stage for Y1230. Increased Y1234+Y1235 phosphorylation compared to uRBCs was confirmed by Western blot analysis using independently-sourced Y1234+Y1235 and pan-MET antibodies (not used on the microarray), with peak signal again seen in trophozoites (Fig. 5c). This confirms that c-MET is indeed phosphorylated in the catalytic domain during trophozoite infection. As observed for B-Raf, we saw a decrease in apparent size of c-MET during the ring-to trophozoite transition, which may be due to differences in phosphorylation status, or to partial degradation of the protein (see Discussion).

Phosphorylation of c-MET Y1003 peaks at the trophozoite stage. In nucleated cells, phosphorylated Y1003 recruits the c-CBL (Casitas B-lineage lymphoma) protein, which promotes ubiquitination and degradation of c-MET via the 26S proteasome^{48,49}. The activity of the 20S proteasome has been previously reported in mature erythrocytes; in contrast, no 26S activity was observed in these cells⁵⁰. The absence of 26S proteasomal activity in erythrocytes could explain why c-MET protein maintains a steady-state level throughout blood-stage parasite development (Fig. 5a), despite Y1003 phosphorylation.

Figure 5 – Activation of c-MET in infected erythrocytes. **a)** c-MET phosphorylation sites covered by the antibody microarray, with the fold change from uRBC control values for the three-time points (ring, trophozoite and schizont). Signals which were significantly different during infection are marked by * ($p < 0.05$, unpaired t-test). **b)** Schematic of the c-MET protein. The juxtamembrane sequence contains both S975 and Y1003, which represent c-CBL binding sites that target the receptor for degradation (red circles)^{51,52}. The catalytic domain contains Y1230, Y1234, Y1235, S1236 and T1241. All activating tyrosine phosphorylations (green circles) occur through receptor homodimerisation^{43,46,53}. The function of S1236 and T1241 (black circles) is uncharacterised. The multifunctional docking site contains two phosphosites, Y1349 and Y1356, which aid in the binding of adapter molecules such as grb2 and GAB1⁴⁶. **c)** Western blot validation of c-MET phosphorylation across asexual development. A phospho-specific antibody to c-MET (Y1234+Y1235) detected phosphorylated c-MET at the ring and trophozoite stages (upper panel). A pan c-MET antibody detected c-MET

across all four samples, with a double band in the uninfected, ring and trophozoite samples (middle panel). An antibody against Glycophorin-C was used as a loading control (lower panel).

Small-molecule inhibitors of host kinases identified with the array impair parasite

proliferation. We previously published that U0126, a highly selective allosteric inhibitor of MEK1/2, as well as other structurally distinct selective inhibitors of these kinases, display low μM potency against *P. falciparum* *in vitro*¹⁵. The findings reported here that several additional human kinases such as c-MET and B-Raf are activated by infection raise the question of whether their inhibition would likewise impair parasite proliferation. Several selective inhibitors of c-MET and B-Raf have emerged from the cancer drug discovery pipeline, allowing us to measure the IC_{50} values in parasite proliferation assays of the following inhibitors: Crizotinib (an inhibitor of c-Met and the related receptor tyrosine kinase ALK [anaplastic lymphoma kinase]); PHA-665752 (a c-MET-selective inhibitor), SB-590885 (a B-Raf selective inhibitor) and Sorafenib (a B-Raf inhibitor which also targets C-Raf, albeit with a 10-fold lower potency)⁵⁴ (Fig. 6b&c). Despite the absence of B-Raf, c-MET and ALK homologues in the *Plasmodium* kinome, all these inhibitors displayed low nano- to micromolar IC_{50} values against both *P. falciparum* and *P. knowlesi* in parasite proliferation assays (Fig. 6a). We cannot at this stage exclude off-target effects against *Plasmodium* proteins. However, our observation that the B-Raf inhibitor SB-590885 is effective only against ring stages, i.e. during the period where B-Raf is present and before its apparent degradation during the ring to trophozoite transition (Fig 4a& d), together with the absence of c-MET or B-Raf homologues in the parasite's kinome, strongly suggests that inhibition of these host erythrocyte kinases impairs parasite proliferation. This in turn indicates that activation of these human kinases by malaria parasites may play an important role in their survival inside erythrocytes.

To determine at which point in the asexual cycle the parasite die if treated with c-MET inhibition at the start of the cycle, and to expand the aforementioned data pertaining to B-Raf inhibition (Fig. 4c), cultures of *P. falciparum* were highly synchronised to a 0 – 4 hrs post-invasion window of development. Cultures were treated with 5x IC_{50} of SB-590885, Crizotinib, PHA-665752, the frontline antimalarial Artemisinin, or with the merozoite invasion inhibitor heparin. Parasites were monitored via flow cytometry every 12 hr over 84 hrs. The normalised live parasitemia detected for each time point was determined through

co-staining with Hoechst 33342 (DNA stain) and Mito-tracker Orange (mitochondrial potential stain). The average values of the biological replicates are represented graphically in Fig. 6d. Treatment with Artemisinin resulted in a reduction in live parasitemia at 12-16hpi, indicative of ring-stage parasite death. Treatment with heparin caused a drop in live parasitemia post 48hpi, as expected for an invasion inhibitor. In contrast, the first notable reduction in parasitemia observed after treatment with Crizotinib, PHA-665752 or SB-590885 occurs prior to the schizont stage. This indicates that all three human kinase inhibitors exert their largest effect on parasite survival at a time that is consistent with the activation time of their target kinases (and, in the case of B-Raf, with the time where the full-length protein is present, i.e. ring stage), as revealed by the microarray and Western blot analysis. Notably, all compounds except heparin resulted in a live parasitemia below 0.2% by 84-88hpi, with SB-590885 and Crizotinib reducing live parasitemia to <0.1%. For an initial assessment of the *in vivo* efficacy of the c-MET inhibitor PHA-665752, we treated mice infected with the rodent malaria parasite *Plasmodium berghei*. A treatment regime and formulation that was previously used to control tumour growth in mice with the same inhibitor⁵⁶ led to a 3.3-fold reduction in parasite load after two doses of treatment *in vivo* (Fig. 6e, day 4).

a	Met Inhibitor	Met/Alk Inhibitor	Raf inhibitor	
	PHA-665752	Crizotinib	SB-590885	Sorafenib
IC_{50} (nM)				
<i>P. falciparum</i>	173	858	490	3688
<i>P. knowlesi</i>	87	312	260	-
Structure

b *P. falciparum* IC_{50} determination

c *P. knowlesi* IC_{50} determination

d Effect on Synchronous *P. falciparum* growth

e Effect on *P. berghei* growth in mice

Figure 6 – IC₅₀ and phenotypic effect of human kinase inhibitors on *P. falciparum* and *P. knowlesi* blood-stage development **a)** IC₅₀ values of the human kinase inhibitors Crizotinib (MET/ALK inhibitor), PHA-665752 (Met inhibitor), SB-590885 (B-Raf inhibitor) and Sorafenib (B-Raf inhibitor, *P. falciparum* only) on asynchronous *P. falciparum* and ring-stage *P. knowlesi* asexual growth. **b)** Growth inhibition curve for asynchronous *P. falciparum*, IC₅₀ measured after 72 hrs of compound exposure, n=3, n=2 for sorafenib. **c)** Growth inhibition curve for *P. knowlesi*, IC₅₀s measured after 50 hrs of compound exposure, n=3. **d)** Effect of treatment with 5x the IC₅₀ value of the compounds Crizotinib, PHA-665752, SB-590885, Artemisinin, DMSO (equivalent volume of DMSO in Crizotinib sample) and heparin (15 µl/ml) on highly synchronous (4-hr window) *P. falciparum* cultures over 88 hrs (n=5). The Y-axis represents the live parasitemia values for each time point normalised to a starting total parasitemia of 2%. All error bars represent the standard error of the mean. **e)** Effect of treating mice with PHA-665752 on the growth of *P. berghei*. The graph shows relative luminescence units (RLU) in the blood of BALB/c mice infected with parasites constitutively expressing firefly luciferase and treated or not with PHA-665752 at 25 mg/kg at times indicated by arrows. * = p < 0.001 by two-tailed T-test; n = 7 mice per treatment group.

Discussion

The antibody microarray analysis generated a large number of testable hypotheses on the involvement of host erythrocyte signaling proteins during infection by the human malaria parasite *P. falciparum*. We achieved Western blot validation for three of the host kinases (MEK1 [15], c-Met [this study] and B-Raf [this study]) that the array data suggest are activated by infection. In the other infection systems that we characterised through this approach (hepatocyte/hepatitis C virus ⁵⁷ and mosquito cells/*Wolbachia* endosymbiont ⁵⁸) the functional relevance of the most interesting array hits was established through siRNA campaigns. In the erythrocyte/*Plasmodium* system where siRNA-based knock-down of host genes is not feasible, we propose that similar Western blot validation should be undertaken as the first step in further work on any of the numerous additional enzyme/pathways identified as hits in the array.

The need for systematic validation notwithstanding, the extent to which host erythrocyte signaling is modulated by *Plasmodium* infection is surprising. The array data are consistent with earlier observations implicating some of the host cell signaling pathways in the infection process, and allowing an in-depth, highly specific understanding of the mobilisation of such pathways (for example, see above the section pertaining to PKC isoforms). The observation (Fig. 4c) that B-Raf is present at the ring-stage but appears to be

degraded in trophozoite- and schizont-infected RBCs is consistent with a thorough modulation of host erythrocyte signaling pathways during infection, not only through activation/inactivation of enzymes by phosphorylation, but also through affecting the abundance of specific signaling elements. In line with this observation, the host erythrocyte cAMP-dependent kinase PKA has been reported to similarly be degraded in infected cells⁵⁹. This is reminiscent of the strategy of other intracellular pathogen to modify the composition of signaling pathways in their host cell. A well-characterised example is that of the Lethal Factor, a component of the anthrax toxin that is secreted by *Bacillus anthracis* and that possesses highly selective endopeptidase activity directed against host cell MEKs, thereby inactivating MAPK pathways (reviewed in ⁶⁰). Down-regulation of MAPK pathways through degradation of specific components is not restricted to host-pathogen interactions. For example, during hematopoietic cell apoptosis, IL-3-deprivation results in the cleavage of Raf-1 by caspase 9, causing separation of the N-terminal regulatory domain from the C-terminal kinase domain⁶¹. Likewise, c-MET and several receptor tyrosine kinases⁶² can undergo specific proteolytic cleavage events, which mediate various biological outcomes (reviewed in ⁶³). It is therefore interesting to note that in our Western blot analysis, c-MET appears as a band of lower mass in trophozoite/schizont RBCs than at the ring stage (Fig. 5). It is tempting to speculate that intracellular pathogens hijack such cellular processes to suppress host pathways. Our observation that host erythrocyte B-Raf is activated early in infection (ring-stage), and apparently subsequently degraded, warrants further studies that will yield fascinating insights into host-parasite interactions. In particular, it will be of great interest to investigate if some host pathways identified as activated in the present study are dependent on the replacement of specific host cell kinase by parasite-encoded kinases, several of which are known to be exported to the host erythrocyte (e.g. ^{64,65}).

A major long-term objective of this research is the identification of antimalarial compounds with untapped modes of action. It is a clear possibility that part of the erythrocyte signaling response to infection may be directed towards activating innate immune defence mechanisms, and thus downregulating such responses would be undesirable. However, based on the hypothesis that at least some of the signaling events triggered in the host cell by infection are required for parasite survival, we tested highly selective inhibitors against some of the activated kinases and showed that these compounds have high potency (in the

low nM range) in *in vitro* culture systems. The nanomolar activity against both *P. falciparum* and the phylogenetically distant *P. knowlesi* for c-Met and B-Raf inhibitors indicate that the reliance on the activation of host erythrocyte pathways spans various *Plasmodium* species. Importantly, the markedly impaired proliferation of *P. berghei* in a murine malaria model treated with the c-Met inhibitor PHA-665752 shows that targeting host kinases to control malaria infection *in vivo* is achievable, strengthening the case for kinase-focussed host-targeted intervention.

Major attractive features of targeting host-cell kinases as a strategy to develop novel antimalarials are threefold: First is the effectiveness of specific inhibitors across phylogenetically distant *Plasmodium* species, illustrated by the facts (i) that inhibitors against three host cell kinases activated by infection (MEK1, c-Met and B-Raf) display similar *in vitro* potency on *P. falciparum* and *P. knowlesi*, and (ii) that MEK and c-Met inhibitors also have activity against the *P. berghei* rodent malaria parasite *ex vivo* (MEK¹⁵) and *in vivo* (c-MET). A second tremendous advantage of this strategy is that the most parsimonious pathway to resistance, i.e. the selection under drug pressure of genotypes encoding a mutated target with decreased susceptibility, is not possible if the target is encoded by the host. Third, large kinase-directed compound libraries are available from the intensive efforts made over the past two decades in the context of cancer chemotherapy (including inhibitors of many of the kinases identified in this study as being activated in infected erythrocytes), providing a unique repository of potential antimalarials. These can now be exploited as a basis for fundamental research into host-parasite interactions, and as leads for the development of novel antimalarials with untapped modes of actions.

Methods

Plasmodium spp. culture and life stage synchronisation

Human erythrocytes were supplied by the Australian Red Cross. *P. falciparum* (clone 3D7) and *P. knowlesi* (YH1) were grown in human erythrocytes as previously described⁶⁶⁻⁶⁹. *P. falciparum* synchronisation (4-hr window) was achieved using 5% w/v Sorbitol as well as 15 µl/ml culture of heparin⁷⁰. Both techniques were utilised for the trophozoite and schizont stage samples used on the antibody microarray, whereas an 8-hr synchronisation window

was achieved through sorbitol alone for the ring stage. Purification of *P. falciparum* trophozoite and schizont-infected cells was achieved using a super-MACS column ⁷¹. *P. knowlesi* cultures for *in vitro* assays were partially synchronised (~12-hr age range) using heparin ⁶⁷.

Kinexus antibody microarray preparation

The Kinex 900P array kits were purchased from Kinexus. Protein extracts were prepared as per the manufacturer's instruction. Each array was, unless otherwise specified, loaded with protein extracts normalized to equivalent cell number using a hemocytometer. Array scanning was conducted by the manufacturer at their facility. Arrays were loaded with protein extracts of rings 4-12 hrs (approximately 33% parasitemia, n=3), trophozoites 24-28 hrs (magnet purified, n=3) and schizonts 44-48 hrs (magnet purified, n=2). Parallel uRBC control extracts containing RBCs from the same donor as the iRBC treatment were loaded onto the arrays to measure baseline RBC phosphorylation (these cells were cultured in the same conditions as the parasite-infected samples for approximately 24 hours before harvesting). To identify phosphosites of parasite proteins cross-reactive to the array, saponin pellets and supernatants of magnet purified, saponin lysed trophozoites/schizonts were compared. Saponin pellet samples were loaded at 2 mg/ml on the array, 20x more pellet material than is present in 2 mg/ml of equivalent saponin supernatant material.

Western blotting

Western blot analysis of infected and uninfected erythrocytic material performed on protein extracts prepared by resuspending cells in M-PER mammalian protein extraction reagent (Pierce) supplemented with 1x protease inhibitory cocktail (EDTA free) (Roche), 300 mM benzamidine, 200 mM phenylmethylsulfonyl fluoride, 500 mM sodium fluoride, 100 mM sodium orthovanadate and 500 mM β -glycerophosphate. Lysates were cleared by centrifugation (10,000 g for 15 mins at 4°C); samples were heated (70°C) in Laemmli reducing sample buffer before separation on 4-12% Tris/Bis gradient SDS-PAGE gels (Invitrogen). After electrophoresis, proteins were transferred to nitrocellulose membrane (Amersham) and blocked (1-hr at room temperature) with 5% Skim milk (Diploma) in Tris-buffered saline containing 0.05% Tween 20 (TBST). Immunoblotting was performed using

the following antibodies; phosphorylated c-MET (Y1234/Y1235; D26, Cell Signaling Technologies), pan c-MET (D1C2, Cell Signaling Technologies), pan B-Raf (OTI4B2, Bio-Rad), phosphorylated B-Raf (S729, AB-PK535, Kinexus), Glycophorin-C (Abcam). All primary antibodies used at 1:500 dilution and incubated overnight at 4°C. Anti-Rabbit/Mouse-horseradish peroxidase-conjugated secondary antibody was used for all experiments at 1:2500 dilution (Monoclonal antibody facility Monash University Clayton, Vic, Australia).

IC₅₀ determination

P. falciparum – The IC₅₀ values of Crizotinib, PHA-665752, SB590885, Sorafenib and Artemisinin were determined with asynchronous cultures with a starting parasitemia of 0.25% in 2% haematocrit. Cultures were incubated with the compounds for 72 hrs before adding SYBR-gold nucleic acid stain (1:10000 dilution) (ThermoFisher Scientific) modified from ⁷² substituting SYBR-green for SYBR-gold. Plates were incubated with SYBR-gold for 1 hr in the dark prior to reading fluorescence on a Tecan plate reader (I-control software), and the inhibitory IC₅₀ determined using GraphPad PRISM (GraphPad software).

P. knowlesi – *P. knowlesi* growth inhibition assays using ring-stage parasites were set up at 1% parasitaemia and 1% haematocrit in 96-well round-bottom plates at a final volume of 45 µL as described ^{73,74}. A 10x final concentration of drug and controls was added to make the final volume to 50 µL and the drug assay cultured for 50 hrs until parasites reached late trophozoites in the next growth cycle. Assays were stained with 10 µg/mL ethidium bromide (EtBr, Bio-Rad) for 1hr and washed prior to flow cytometry (Becton Dickinson LSR) assessment of parasitaemia with gating as per established protocols ⁷⁵. Parasitaemia counts were quantitated using FlowJo software (Tree Star) and the inhibitory IC₅₀ determined using GraphPad PRISM (GraphPad Software).

Time-dependent inhibitor treatment of parasites monitored via flow cytometry

Highly synchronous cultures (0 - 4 hrs post-invasion) at 2% haematocrit treated with DMSO (vehicle), Crizotinib, PHA-665752, SB-590885 or artemisinin at 5x the IC₅₀ values were monitored at 12-hr time intervals over two full asexual intra-erythrocytic cycles (84 hrs) by

flow cytometry. Live parasitaemia was quantitated by a dual-colour flow cytometry staining protocol using 2 μ M Hoechst-33342 staining for 8 minutes and 75 nM MitoTracker Orange for 25 minutes. Staining was completed in v-bottom plates with two washes in complete RPMI. Stained cells were transferred to polypropylene tubes and diluted one in four in complete RPMI. Cells were then immediately analysed on an LSR BDFortessaTM with the laser UV379 and filter 450/50 for Hoechst-33242 staining, and laser YG585 with filter 585/15 for MitoTracker Orange staining. Biological replicates (n=5) were normalised to a 2% total parasitemia value at the first-time point (0 – 4 hrs post-invasion).

***P. falciparum* stage-dependent inhibitor treatment and washout out assay**

P. falciparum cultures were synchronised into a 6-hour development window at 2% haematocrit and treated with 5x the 72hour IC₅₀ of SB590885, Artemisinin or DMSO at 0-6hpi (ring stage treatment) 22-28hpi (trophozoite stage treatment) and 38-44hpi (schizont stage treatment). Treatments were washed from the respective cultures after 4 hours (denoted by the black arrow) and monitored until the next cycle of parasite development to assess parasite survival using flow cytometry (2 μ M Hoechst-33342 staining for 8 minutes). Starting parasitemia was normalised to 5% across the biological replicates (n=3) error bars represent the standard error of the mean.

***P. berghei* in vivo drug treatment assay**

Animal research was conducted under licenses from the UK Home Office, and protocols were approved by the Animal Welfare and Ethical Review Body of the Wellcome Sanger Institute. Mice were purchased from Envigo, kept in specific-pathogen-free conditions and subjected to regular pathogen monitoring by sentinel screening. They were housed in individually ventilated cages furnished with autoclaved aspen woodchip, fun tunnel and Nestlets at 21°C +/- 2°C under a 12:12 hr light-dark cycle at a relative humidity of 55 +/- 10%. They were fed a commercially prepared autoclaved dry rodent diet and water, both available *ad libitum*. The health of animals was monitored by routine daily visual health checks. Cohorts of seven female, 10-week-old BALB/c mice per treatment group were infected by i.v. inoculation with 10⁶ infected erythrocytes. Assays to determine parasite growth were performed as described ⁷⁶, using the reporter parasite line PbGFPLuc_{con} (RMgm-29 in the rodent malaria genetic modification database,

<http://www.pberghei.eu/index.php?rmgm=29>), which expresses a GFP-firefly luciferase fusion protein under the control of the constitutive *eef1a* promoter ⁷⁷. Treatment with PHA-665752 or solvent control was 3hr thereafter and on day two post infection by i.p. injection of 25 mg/kg/day of PHA-665752. The drug was prepared fresh by diluting a stock of 50 mg PHA-665752 in 0.5 ml DMSO (w/v) in water (1:40, v/v). The same drug formulation and treatment regime had proved effective in controlling carcinoma growth ⁵⁶.

Data Availability Statement

All raw data pertaining to this study are available in Supplemental Datasets 1 to 3.

Acknowledgements

We thank Dr Kylie Quinn for lending her Flow Cytometry expertise and the staff of Monash FlowCore Facility, in particular Andrew Fryga and Adam Dinsdale, for their continued experimental assistance. We also thank Prof Roger Daly, Dr. Teresa Carvalho, A/Prof Jose Garcia-Bustos and Prof Brian Cooke for fruitful discussions. This work was supported by the Australian Government National Health and Medical Research Council (Project Grants APP1082619 [CD] and APP1143974 [DW]), Australian Research Council PhD Scholarship (AB), Hospital Research Foundation Fellowship (DW) and internal support from Monash University and the University of Adelaide (Beacon Fellowship (DW)).

References

- 1 Singh, B. *et al.* A large focus of naturally acquired Plasmodium knowlesi infections in human beings. *Lancet* **363**, 1017-1024, doi:10.1016/S0140-6736(04)15836-4 (2004).
- 2 Brasil, P. *et al.* Outbreak of human malaria caused by Plasmodium simium in the Atlantic Forest in Rio de Janeiro: a molecular epidemiological investigation. *Lancet Glob Health* **5**, e1038-e1046, doi:10.1016/s2214-109x(17)30333-9 (2017).
- 3 (WHO), W. H. O. World malaria report. (2017).

653 4 Haldar, K., Bhattacharjee, S. & Safeukui, I. Drug resistance in Plasmodium. *Nat Rev Microbiol*
654 **16**, 156-170, doi:10.1038/nrmicro.2017.161 (2018).

655 5 Bhullar, K. S. *et al.* Kinase-targeted cancer therapies: progress, challenges and future
656 directions. *Mol Cancer* **17**, 48, doi:10.1186/s12943-018-0804-2 (2018).

657 6 Roskoski, R., Jr. Properties of FDA-approved small molecule protein kinase inhibitors.
658 *Pharmacol Res* **144**, 19-50, doi:10.1016/j.phrs.2019.03.006 (2019).

659 7 Ward, P., Equinet, L., Packer, J. & Doerig, C. Protein kinases of the human malaria parasite
660 Plasmodium falciparum: the kinome of a divergent eukaryote. *BMC Genomics* **5**, 79,
661 doi:10.1186/1471-2164-5-79 (2004).

662 8 Lucet, I. S., Tobin, A., Drewry, D., Wilks, A. F. & Doerig, C. Plasmodium kinases as targets for
663 new-generation antimalarials. *Future Med Chem* **4**, 2295-2310, doi:10.4155/fmc.12.183
664 (2013).

665 9 Schor, S. & Einav, S. Combating Intracellular Pathogens with Repurposed Host-Targeted
666 Drugs. *ACS Infect Dis* **4**, 88-92, doi:10.1021/acsinfecdis.7b00268 (2018).

667 10 Deitsch, K. W. Self-targeting blocks malaria parasite invasion. *J Exp Med* **212**, 1143,
668 doi:10.1084/jem.2128insight5 (2015).

669 11 Zenonos, Z. A. *et al.* Basigin is a druggable target for host-oriented antimalarial
670 interventions. *J Exp Med* **212**, 1145-1151, doi:10.1084/jem.20150032 (2015).

671 12 Murphy, S. C. *et al.* Erythrocyte G protein as a novel target for malarial chemotherapy. *PLoS*
672 *Med* **3**, e528, doi:06-PLME-RA-0363R3 [pii]
673 10.1371/journal.pmed.0030528 (2006).

674 13 Harrison, T. *et al.* Erythrocyte G protein-coupled receptor signaling in malarial infection.
675 *Science* **301**, 1734-1736, doi:10.1126/science.1089324
676 301/5640/1734 [pii] (2003).

677 14 Ntumngia, F. B., Thomson-Luque, R., Pires, C. V. & Adams, J. H. The role of the human Duffy
678 antigen receptor for chemokines in malaria susceptibility: current opinions and future
679 treatment prospects. *J Receptor Ligand Channel Res* **9**, 1-11, doi:10.2147/JRLCR.S99725
680 (2016).

681 15 Sicard, A. *et al.* Activation of a PAK-MEK signalling pathway in malaria parasite-infected
682 erythrocytes. *Cell Microbiol* **13**, 836-845, doi:10.1111/j.1462-5822.2011.01582.x (2011).

683 16 Hall, B. S., Daramola, O. O., Barden, G. & Targett, G. A. Modulation of protein kinase C
684 activity in Plasmodium falciparum-infected erythrocytes. *Blood* **89**, 1770-1778 (1997).

- 17 Brizuela, M. *et al.* Treatment of erythrocytes with the 2-cys peroxiredoxin inhibitor, Conoidin A, prevents the growth of *Plasmodium falciparum* and enhances parasite sensitivity to chloroquine. *PLoS One* **9**, e92411, doi:10.1371/journal.pone.0092411 (2014).
- 18 Koncarevic, S. *et al.* The malarial parasite *Plasmodium falciparum* imports the human protein peroxiredoxin 2 for peroxide detoxification. *Proceedings of the National Academy of Sciences of the United States of America* **106**, 13323-13328, doi:10.1073/pnas.0905387106 (2009).
- 19 Carvalho, T. G. *et al.* The ins and outs of phosphosignalling in *Plasmodium*: Parasite regulation and host cell manipulation. *Mol Biochem Parasitol* **208**, 2-15, doi:10.1016/j.molbiopara.2016.05.006 (2016).
- 20 Park, E. R., Eblen, S. T. & Catling, A. D. MEK1 activation by PAK: a novel mechanism. *Cell Signal* **19**, 1488-1496, doi:10.1016/j.cellsig.2007.01.018 (2007).
- 21 Radu, M., Semenova, G., Kosoff, R. & Chernoff, J. PAK signalling during the development and progression of cancer. *Nat Rev Cancer* **14**, 13-25 (2014).
- 22 Elandalloussi, L. M. & Smith, P. J. Preparation of pure and intact *Plasmodium falciparum* plasma membrane vesicles and partial characterisation of the plasma membrane ATPase. *Malar J* **1**, 6 (2002).
- 23 Millholland, M. G. *et al.* A host GPCR signaling network required for the cytolysis of infected cells facilitates release of apicomplexan parasites. *Cell Host Microbe* **13**, 15-28, doi:10.1016/j.chom.2012.12.001 (2013).
- 24 Bouyer, G. *et al.* *Plasmodium falciparum* infection induces dynamic changes in the erythrocyte phospho-proteome. *Blood Cells Mol Dis* **58**, 35-44, doi:10.1016/j.bcmd.2016.02.001 (2016).
- 25 Welman, A., Griffiths, J. R., Whetton, A. D. & Dive, C. Protein kinase C delta is phosphorylated on five novel Ser/Thr sites following inducible overexpression in human colorectal cancer cells. *Protein Sci* **16**, 2711-2715, doi:10.1110/ps.072874607 (2007).
- 26 Liu, Y., Graham, C., Li, A., Fisher, R. J. & Shaw, S. Phosphorylation of the protein kinase C-theta activation loop and hydrophobic motif regulates its kinase activity, but only activation loop phosphorylation is critical to in vivo nuclear-factor-kappaB induction. *Biochem J* **361**, 255-265, doi:10.1042/bj3610255 (2002).
- 27 Wang, X., Chuang, H. C., Li, J. P. & Tan, T. H. Regulation of PKC-theta function by phosphorylation in T cell receptor signaling. *Front Immunol* **3**, 197, doi:10.3389/fimmu.2012.00197 (2012).

- 28 Zhang, B. H. & Guan, K. L. Activation of B-Raf kinase requires phosphorylation of the conserved residues Thr598 and Ser601. *EMBO J* **19**, 5429-5439, doi:10.1093/emboj/19.20.5429 (2000).
- 29 Chong, H., Lee, J. & Guan, K. L. Positive and negative regulation of Raf kinase activity and function by phosphorylation. *EMBO J* **20**, 3716-3727, doi:10.1093/emboj/20.14.3716 (2001).
- 30 Leicht, D. T. *et al.* Raf kinases: function, regulation and role in human cancer. *Biochim Biophys Acta* **1773**, 1196-1212, doi:10.1016/j.bbamcr.2007.05.001 (2007).
- 31 Dorin-Semblat, D. *et al.* Functional characterization of both MAP kinases of the human malaria parasite *Plasmodium falciparum* by reverse genetics. *Mol Microbiol* **65**, 1170-1180, doi:10.1111/j.1365-2958.2007.05859.x (2007).
- 32 Lake, D., Correa, S. A. & Muller, J. Negative feedback regulation of the ERK1/2 MAPK pathway. *Cell Mol Life Sci* **73**, 4397-4413, doi:10.1007/s00018-016-2297-8 (2016).
- 33 Hekman, M. *et al.* Novel C-Raf phosphorylation sites: serine 296 and 301 participate in Raf regulation. *FEBS Lett* **579**, 464-468, doi:10.1016/j.febslet.2004.11.105 (2005).
- 34 Brummer, T. *et al.* Functional analysis of the regulatory requirements of B-Raf and the B-Raf(V600E) oncoprotein. *Oncogene* **25**, 6262-6276, doi:10.1038/sj.onc.1209640 (2006).
- 35 Mason, C. S. *et al.* Serine and tyrosine phosphorylations cooperate in Raf-1, but not B-Raf activation. *EMBO J* **18**, 2137-2148, doi:10.1093/emboj/18.8.2137 (1999).
- 36 Brummer, T., Shaw, P. E., Reth, M. & Misawa, Y. Inducible gene deletion reveals different roles for B-Raf and Raf-1 in B-cell antigen receptor signalling. *EMBO J* **21**, 5611-5622 (2002).
- 37 Eblen, S. T. *et al.* Mitogen-activated protein kinase feedback phosphorylation regulates MEK1 complex formation and activation during cellular adhesion. *Mol Cell Biol* **24**, 2308-2317, doi:10.1128/mcb.24.6.2308-2317.2004 (2004).
- 38 Roskoski, R., Jr. MEK1/2 dual-specificity protein kinases: structure and regulation. *Biochem Biophys Res Commun* **417**, 5-10, doi:10.1016/j.bbrc.2011.11.145 (2012).
- 39 Bromberg-White, J. L., Andersen, N. J. & Duesbery, N. S. MEK genomics in development and disease. *Brief Funct Genomics* **11**, 300-310, doi:10.1093/bfpg/els022 (2012).
- 40 Payne, D. M. *et al.* Identification of the regulatory phosphorylation sites in pp42/mitogen-activated protein kinase (MAP kinase). *EMBO J* **10**, 885-892 (1991).
- 41 Goetz, E. M., Ghandi, M., Treacy, D. J., Wagle, N. & Garraway, L. A. ERK mutations confer resistance to mitogen-activated protein kinase pathway inhibitors. *Cancer Res* **74**, 7079-7089, doi:10.1158/0008-5472.CAN-14-2073 (2014).

- 42 Lai, S. & Pelech, S. Regulatory roles of conserved phosphorylation sites in the activation T-loop of the MAP kinase ERK1. *Mol Biol Cell* **27**, 1040-1050, doi:10.1091/mbc.E15-07-0527 (2016).
- 43 Organ, S. L. & Tsao, M. S. An overview of the c-MET signaling pathway. *Ther Adv Med Oncol* **3**, S7-S19, doi:10.1177/1758834011422556 (2011).
- 44 Zarnegar, R. & Michalopoulos, G. K. The many faces of hepatocyte growth factor: from hepatopoiesis to hematopoiesis. *J Cell Biol* **129**, 1177-1180 (1995).
- 45 Teofili, L. *et al.* Expression of the c-met proto-oncogene and its ligand, hepatocyte growth factor, in Hodgkin disease. *Blood* **97**, 1063-1069 (2001).
- 46 Rodrigues, G. A. & Park, M. Autophosphorylation modulates the kinase activity and oncogenic potential of the Met receptor tyrosine kinase. *Oncogene* **9**, 2019-2027 (1994).
- 47 Emaduddin, M., Bicknell, D. C., Bodmer, W. F. & Feller, S. M. Cell growth, global phosphotyrosine elevation, and c-Met phosphorylation through Src family kinases in colorectal cancer cells. *Proc Natl Acad Sci U S A* **105**, 2358-2362, doi:10.1073/pnas.0712176105 (2008).
- 48 Peschard, P. *et al.* Mutation of the c-Cbl TKB domain binding site on the Met receptor tyrosine kinase converts it into a transforming protein. *Mol Cell* **8**, 995-1004 (2001).
- 49 Jeffers, M., Taylor, G. A., Weidner, K. M., Omura, S. & Vande Woude, G. F. Degradation of the Met tyrosine kinase receptor by the ubiquitin-proteasome pathway. *Mol Cell Biol* **17**, 799-808 (1997).
- 50 Neelam, S., Kakhniashvili, D. G., Wilkens, S., Levene, S. D. & Goodman, S. R. Functional 20S proteasomes in mature human red blood cells. *Exp Biol Med (Maywood)* **236**, 580-591, doi:10.1258/ebm.2011.010394 (2011).
- 51 Abella, J. V. *et al.* Met/Hepatocyte growth factor receptor ubiquitination suppresses transformation and is required for Hrs phosphorylation. *Mol Cell Biol* **25**, 9632-9645, doi:10.1128/MCB.25.21.9632-9645.2005 (2005).
- 52 Trusolino, L., Bertotti, A. & Comoglio, P. M. MET signalling: principles and functions in development, organ regeneration and cancer. *Nat Rev Mol Cell Biol* **11**, 834-848, doi:10.1038/nrm3012 (2010).
- 53 Carrolo, M. *et al.* Hepatocyte growth factor and its receptor are required for malaria infection. *Nat Med* **9**, 1363-1369, doi:10.1038/nm947 (2003).
- 54 King, A. J. *et al.* Demonstration of a genetic therapeutic index for tumors expressing oncogenic BRAF by the kinase inhibitor SB-590885. *Cancer Res* **66**, 11100-11105, doi:10.1158/0008-5472.CAN-06-2554 (2006).

783 55 Miranda-Saavedra, D., Gabaldon, T., Barton, G. J., Langsley, G. & Doerig, C. The kinomes of
784 apicomplexan parasites. *Microbes Infect* **14**, 796-810, doi:10.1016/j.micinf.2012.04.007
785 (2012).

786 56 Zhi, J. *et al.* Effects of PHA-665752 and vemurafenib combination treatment on in vitro and
787 murine xenograft growth of human colorectal cancer cells with BRAF(V600E) mutations.
788 *Oncol Lett* **15**, 3904-3910, doi:10.3892/ol.2018.7770 (2018).

789 57 Haqshenas, G. *et al.* Signalome-wide assessment of host cell response to hepatitis C virus.
790 *Nat Commun* **8**, 15158, doi:10.1038/ncomms15158 (2017).

791 58 Haqshenas, G. *et al.* A Role for the Insulin Receptor in the Suppression of Dengue Virus and
792 Zika Virus in Wolbachia-Infected Mosquito Cells. *Cell Rep* **26**, 529-535 e523,
793 doi:10.1016/j.celrep.2018.12.068 (2019).

794 59 Syin, C. *et al.* The H89 cAMP-dependent protein kinase inhibitor blocks Plasmodium
795 falciparum development in infected erythrocytes. *Eur J Biochem* **268**, 4842-4849,
796 doi:10.1046/j.1432-1327.2001.02403.x (2001).

797 60 Turk, B. E. Exceptionally Selective Substrate Targeting by the Metalloprotease Anthrax Lethal
798 Factor. *Adv Exp Med Biol* **1111**, 189-203, doi:10.1007/5584_2018_273 (2019).

799 61 Cornelis, S. *et al.* Apoptosis of hematopoietic cells induced by growth factor withdrawal is
800 associated with caspase-9 mediated cleavage of Raf-1. *Oncogene* **24**, 1552-1562,
801 doi:10.1038/sj.onc.1208401 (2005).

802 62 Merilahti, J. A. M. & Elenius, K. Gamma-secretase-dependent signaling of receptor tyrosine
803 kinases. *Oncogene* **38**, 151-163, doi:10.1038/s41388-018-0465-z (2019).

804 63 Fernandes, M., Duplaquet, L. & Tulasne, D. Proteolytic cleavages of MET: the divide-and-
805 conquer strategy of a receptor tyrosine kinase. *BMB Rep* **52**, 239-249 (2019).

806 64 Dorin-Semblat, D. *et al.* Malaria Parasite-Infected Erythrocytes Secrete PfCK1, the
807 Plasmodium Homologue of the Pleiotropic Protein Kinase Casein Kinase 1. *PLoS One* **10**,
808 e0139591, doi:10.1371/journal.pone.0139591 (2015).

809 65 Nunes, M. C., Goldring, J. P., Doerig, C. & Scherf, A. A novel protein kinase family in
810 Plasmodium falciparum is differentially transcribed and secreted to various cellular
811 compartments of the host cell. *Mol Microbiol* **63**, 391-403, doi:10.1111/j.1365-
812 2958.2006.05521.x (2007).

813 66 Trager, W. & Jensen, J. B. Human malaria parasites in continuous culture. *Science* **193**, 673-
814 675 (1976).

- 67 Lyth, O. *et al.* Cellular dissection of malaria parasite invasion of human erythrocytes using
viable Plasmodium knowlesi merozoites. *Sci Rep* **8**, 10165, doi:10.1038/s41598-018-28457-z
(2018).
- 68 Walliker, D. *et al.* Genetic analysis of the human malaria parasite Plasmodium falciparum.
Science **236**, 1661-1666 (1987).
- 69 Dankwa, S. *et al.* Ancient human sialic acid variant restricts an emerging zoonotic malaria
parasite. *Nat Commun* **7**, 11187, doi:10.1038/ncomms11187 (2016).
- 70 Hill, D. L., Eriksson, E. M. & Schofield, L. High yield purification of Plasmodium falciparum
merozoites for use in opsonizing antibody assays. *J Vis Exp*, doi:10.3791/51590 (2014).
- 71 Ribaut, C. *et al.* Concentration and purification by magnetic separation of the erythrocytic
stages of all human Plasmodium species. *Malar J* **7**, 45, doi:10.1186/1475-2875-7-45 (2008).
- 72 Johnson, J. D. *et al.* Assessment and continued validation of the malaria SYBR green I-based
fluorescence assay for use in malaria drug screening. *Antimicrob Agents Chemother* **51**,
1926-1933, doi:10.1128/AAC.01607-06 (2007).
- 73 Wilson, D. W., Crabb, B. S. & Beeson, J. G. Development of fluorescent Plasmodium
falciparum for in vitro growth inhibition assays. *Malar J* **9**, 152, doi:10.1186/1475-2875-9-
152 (2010).
- 74 Nguyen, N. *et al.* Broad activity of diphenyleneiodonium analogues against Mycobacterium
tuberculosis, malaria parasites and bacterial pathogens. *Eur J Med Chem* **148**, 507-518,
doi:10.1016/j.ejmech.2017.10.010 (2018).
- 75 Wilson, D. W., Langer, C., Goodman, C. D., McFadden, G. I. & Beeson, J. G. Defining the
timing of action of antimalarial drugs against Plasmodium falciparum. *Antimicrob Agents
Chemother* **57**, 1455-1467, doi:10.1128/AAC.01881-12 (2013).
- 76 Lin, J. W. *et al.* Screening inhibitors of P. berghei blood stages using bioluminescent reporter
parasites. *Methods Mol Biol* **923**, 507-522, doi:10.1007/978-1-62703-026-7_35 (2013).
- 77 Franke-Fayard, B. *et al.* A Plasmodium berghei reference line that constitutively expresses
GFP at a high level throughout the complete life cycle. *Mol Biochem Parasitol* **137**, 23-33,
doi:10.1016/j.molbiopara.2004.04.007 (2004).