First report of the Eocene bivalve *Schedocardia* (Mollusca, Cardiidae) from Cuba

JOHANSET ORIHUELA^{1*}

¹ Department of Earth and Environment (Geosciences), Florida International University,
Miami, Florida 33199, USA, Email: Jorih003@fiu.edu * corresponding author

YASMANI CEBALLOS IZQUIERDO²

² Instituto de Geofísica y Astronomía, calle 212 No.2906 entre 29 y 31, La Coronela, La Lisa,
La Habana, Cuba, Email: yasmaniceballos@gmail.com

ROGER W. PORTELL³

³ Florida Museum of Natural History, 1659 Museum Road, University of Florida, Gainesville,
Florida 32611, USA, Email: portell@flmnh.ufl.edu

ABSTRACT

Herein we provide the first report of the cardiid bivalve *Schedocardia* from Cuba. The single, partial, valve external mold was derived from the Madruga Formation which is characterized by a richly diverse marine fauna including echinoderms, brachiopods, benthic and planktonic foraminiferans, but from which bivalves were not previously reported. The unit is considered late Paleocene in age (Thanetian), but the presence of *Schedocardia* supports a possible age extension of the formation into the early Eocene (Ypresian). Moreover, we provide a reconsideration of the historical factors that affected the accepted type locality of the outcrop, which allows for an alternative interpretation of the fauna found therein.

Keywords: Mollusca; Cardiidae; Fossils; Eocene; Madruga Formation; Cuba

RESUMEN

Se reporta por primera vez el género bivalvo *Schedocardia* de la Formación Madruga, para el registro fósil de Cuba. De esta formación se ha reportado una rica fauna de invertebrados incluyendo equinodermos, braquiópodos y, especialmente de foraminíferos, pero no moluscos bivalvos. Se ha considerado que la Formación Madruga se depositó durante el Paleoceno tardío (Thanetiano), no obstante, la presencia de *Schedocardia* apoya la extensión de edad de la formación hasta el Eoceno temprano (Ypresiano). Además, se consideran factores antropogénicos que han afectado la localidad tipo de la formación.

Palabras clave: Molusco; Cardiido; Fósil; Eoceno; Madruga; Cuba

INTRODUCTION

The Paleocene of Cuba is represented in localized outcrops throughout the island, but a characteristic one is located near the town of Madruga, in the central-eastern part of the Mayabeque Province, where a complex of ophiolites, Cretaceous volcanic rocks, and latest Cretaceous, Paleogene and Neogene age sedimentary formations crop out (Furrazola-Bermúdez et al., 1964; Albear et al., 1985).

This region has been the subject of paleontological investigations since the early-middle 20th century, as part of oil and mineral prospecting by North American companies (De Golyer, 1918; Lewis, 1932a, b; Palmer, 1932; Wright and Sweet, 1924). Based on field trips made between 1929 and 1946 around Cuba, Palmer (1948) compiled a list of fossil-bearing localities, including more than 100 sites for the town of Madruga. In his publication, Palmer mentioned a locality previously identified by Lewis (1932a) as “*Cut under railroad bridge two kilometers west of Madruga at Central San Antonio*” and cataloged it as a notable locality (Loc. 757 in Palmer, 1948). Since Lewis (1932a) had not selected a holotype (type section) nor did he record its faunal content, Bermúdez (1950) erected the roadcut under the railroad bridge as the type locality for the Madruga Formation (Loc. Bermúdez Sta. 76b) and assigned it a Paleocene age.

This locality was originally described as “Madruga marls” by Lewis (1932a), who erroneously considered it of Late Cretaceous age. Its fossil fauna was reported later by Palmer (1934), Palmer (1932, 1948), Cushman and Bermúdez (1948a, b, 1949), Bermúdez (1938, 1950), Cooper (1955, 1979), Sachs (1959), and Kier (1984). They had found mostly Cretaceous and Paleocene large foraminifers, but also assemblages consisting of small benthic and planktonic species including several index taxa (Palmer, 1934; Cushman and Bermúdez, 1948a, b, 1949; Bermúdez, 1950; Furrazola-Bermúdez et al., 1964; Franco-Álvarez et al.,

1992). Interestingly, brachiopods, scaphopods, and echinoid fragments were also reported. The brachiopods collected by Palmer, deposited in the USNM (United States National Museum, Washington D.C.), were later described and figured by Cooper (1955, 1979). Although these works provide a detailed account of the foraminifer faunule from the Madruga outcrop, bivalves had not been reported, and the echinoids and scaphopods still need investigation.

Recent fieldwork at this locality yielded several interesting, well-preserved fossil bivalve molds, representing a new addition to the fauna of the Madruga Formation undetected in over 80 years of collecting history at the site. Furthermore, our new observations contribute to the reinterpretation of the age and environment of deposition of the Madruga Formation.

MATERIALS AND METHODS

Locality

The Madruga Formation type section and collecting site of the bivalves herein reported is located at a highway cut near the Boris Luis Santa Coloma Sugar Plantation, outskirts of the town of Madruga, Mayabeque Province: GPS coordinates Lat. 22.906787°, Long. -81.872793°, datum WGS 84, altitude 159 m above modern sea level (taken from Google Earth). This is the same location, below the railroad bridge at the Carretera Central west of Madruga town, reported by Palmer (1948) as Loc. 757, and Loc. Sta. 76b in Bermúdez (1950). This section is the current lectostratotype of the Madruga Formation (Albear et al., 1985; Franco-Álvarez et al., 1992). The outcrop section discussed is ~4 m above the highway level, on each side. The bivalves herein described were collected at a +1 m

and +2.5 m (Fig. 1). This is nearby the original localities visited by Palmer and Bermúdez, but the exact locality of their collecting stations is unknown because they did not indicate them.

The specimens were collected from loose fragments found within the conglomerates that make up the stratigraphic column of the formation at the referenced locality (Fig. 1). Their lithology, coloration, and position within the column indicate they are loose clastics of the Madruga Formation, and not the Late Cretaceous bed that lies discordantly below.

Geological setting: Lithology and Deposition environment

The Madruga section has been described as brown or reddish-brown calcareous well-stratified, cemented, greywacke sandstones, sandy-shales, and shales with polymictic conglomerate intercalations. This formation lies discordantly over Late Cretaceous deposits, which are often included within the Madruga Formation. Clastic material derived from older Cretaceous volcanic-sedimentary rocks are conglomerates that reach from 30 to 40 cm up to 1 to 2 m in diameter. Radiolarian sandstones have been reported (Cushman and Bermúdez, 1948a:68), but the origin of the silica (biogenic or volcanic) is unknown. The age of the unit was originally considered Late Cretaceous by Lewis (1932a:539, 1932b) but later defined as late Paleocene based on several index Foraminifera (Bermúdez, 1950, 1961).

Stratigraphically, the Madruga Formation lies concordant over the Mercedes Formation and discordant over the Peñalver (Late Cretaceous/early Paleocene) and Via Blanca Formations (Late Cretaceous) (Fig. 2). It is covered concordantly by the Capdevilla Formation (early Eocene) and discordantly by the Cojimar Formation (early – middle Miocene) and Nazareno Formation (late middle Eocene – late Eocene). The Cretaceous rocks

(K₂^{cp-m?}) can be confused with the overlaying Paleocene layers (Albear et al., 1985; Franco-Álvarez et al., 1992). At the type locality, contacts with other formations are not exposed within the outcrop but are partially exposed at nearby locations. Fossil specimens are sparse and rare throughout the outcrop.

The environment of deposition has been interpreted as a marine synorogenic piggyback basin of medium to deep depth (bathyal), formed under unstable tectonic environment, transportation, and turbidity currents (Bralower and Iturralde-Vinent, 1997; Iturralde-Vinent and MacPhee, 1999; Iturralde-Vinent et al., 2016; Núñez-Cambra and Iturralde-Vinent, 2016).

Abbreviations

Institutional— Specimen repository: Paleontology collection of the Madruga Museum (MPAL), Mayabeque Province, Cuba. LBF-large benthic foraminifera; PKF, planktonic foraminifera; SBF, small benthic Foraminifera. United States National Museum (USNM), Washington, D.C.

RESULTS

Systematic Paleontology

Class: Bivalvia Linnaeus, 1758

Order: Cardiida Férussac, 1822

Family: Cardiidae Lamarck, 1809

Cardiinae Schneider, 2002

Genus: *Schedocardia* Stewart, 1930

Schedocardia cf. hatchetigbeensis (Aldrich, 1886)

Specimens- MPAL-1001 (Fig. 3), deposited at the paleontology collection of the Museum of Madrugá city. Collected by Yasmani Ceballos Izquierdo on June 2017, at 3-4 m west of the railroad bridge, on the north margin of the roadcut (Figs. 1-2).

Description- The fossil is an incomplete external mold of the right valve in a calcareous matrix. The external mold is obliquely quadrate, elliptic-ovoid, or schediform in shape (D in Schneider, 2002); radial ribs are narrow and nearly symmetrical throughout. There are small-simple foramina on the interstices between the radial ribs, valve margins crenulated. The interstices between radial ribs are wider than the width of the rib (Fig. 4). External ribs strongly expressed, but not alternating (i.e., wide to narrow) on a central slope. There are >19 ribs present on our specimen, but the exact number is unknown due to incompleteness. The spines or their foramina are more pronounced in the lower valve margins. Only the upper part of the valve external mold is visible. No spines or nobs present. Flanking ribs closely packed together. Cardinals, muscle scars, or hinge not visible (Fig. 4). The umbo is generally incomplete, but likely opisthogyrate and wide. Marked, ovoid-shaped and incised foramina occur almost symmetrically along the curve-space between the valve ribs (Fig. 4). In cross-section, these either extend straight down or curved at the deepest tip. These appear to represent scars of spines that have broken off.

A smaller cardiid specimen is embedded in the same sample opposite this cast (not illustrated here).

DISCUSSION

Identification

The specimen (MPAL-1001) is comparable to *Schedocardia juncea* (Olsson, 1930), and *Schedocardia gatunense* (Dall, 1900), in their elliptic-ovate shape of the valve and higher, less oblique shell anteriorly (Figs. 3-5). However, it differs from these three species in having prominent ribs near the umbo, many more costae, interstices that are flat but wide, not narrow, and ribs that do not bear beads (as in *S. gatunense*). It also differs from *Schedocardia brewerii* Gabb, 1864 in having large spines (Stewart, 1930: 257) more closely packed costae and interstices, and more ovate-elliptic shape of the valve (more elongated in *S. brewerii*, see pl. 3, fig. 11 in Moore, 2002).

Compared to images of the syntypes of *S. hatchetigbeensis* (Aldrich, 1886) (USNM 638802 and 645087, both from Hatchetigbee Bluff, Alabama, USA) the specimen from Madruga Formation agrees closely with this species in having several radial rows of tiny scars alternating with larger scars, narrower interstices, lacking rib beads, low spines inside these spaces, and ovate shape of the valve. However, the species is not fully identical, and may represent a new form. Due to their overall similarity, we tentatively refer it here to *S. cf. hatchetigbeensis* until further specimens are available (Figs. 5-6).

In addition to the *Schedocardia*, another cardiid bivalve, *Acanthocardia* (MPAL-1002), was discovered in the same deposit (Fig. 7). This specimen has an obliquely quadrate shell, with 21 visible ribs, crenulate valve margins, and spinose/nodose ribs. Probable double ribs can be seen on the imbricated valve side. Although the specimen is incomplete, it resembles in overall morphology *Acanthocardia tuberculata* (Linnaeus, 1758) and *Acanthocardia echinata* (Linnaeus, 1758). However, the specimen is too incomplete to assign to species.

Chronology

The Madruga Formation has been formerly assigned to the late Paleocene (upper) by the researchers of the Cuban Geological Lexicon and others (Sánchez-Roig, 1949; Sachs, 1957; Bermúdez, 1950, 1961; Furrázola-Bermúdez et al., 1964; Franco-Álvarez et al., 1992) based on the presence of several index taxa, such as the planktic foram *Morozovella velascoensis* (Cushman, 1925), index of PKF zone P5, and *Pseudophragmina* (= *Athecocyclina*) *stephensoni* (Vaughan, 1929), a large benthic foraminifer of carbonate banks (Özcan et al., 2019). More recently, researchers have extended the age of the formation to the early Eocene (Ypresian) due to the presence of the LBF *Euconuloides wellsi* (Cole and Bermúdez, 1944), which is absent in Paleogene assemblages (Blanco-Bustamante et al., 1999).

The genus *Schedocardia* appears in the fossil record in the late Paleocene (Keen, 1980; Schneider, 2002), but did not have its acme until the Eocene. *Schedocardia hatchetigbeensis* has been reported from the early Eocene (Ypresian) of Alabama and Texas, USA (Toulmin, 1977; Garvie, 2013; Sessa et al., 2012). Other *Schedocardia* species, such as *S. juncea* and *S. gatunense* are reported from the late Eocene of Panama, Venezuela, Colombia, and Peru (Woodring, 1982). Therefore, the genus is generally considered to be an Eocene indicator (Woodring, 1982:542). The occurrence of *Schedocardia* in the Madruga Formation seems to support the extension of the Cuban formation at least to the early Eocene.

Cushman and Bermúdez (1948a, b) correlated the Madruga Formation with late Paleocene formations from Alabama and Texas, so it is not so surprising that *S. cf. hatchetigbeensis* appears here. These North American formations are now considered early Eocene (Toulmin, 1977; Sessa et al., 2012).

Environment

Schedocardia and *Acanthocardia* are considered facultative mobile infaunal (benthic) suspension feeders (Clarkson, 1986). Their occurrence, along with multiple large benthic

forams (LBF's), echinoids, and brachiopods suggests that after the late Paleocene, during the early Eocene, the area of deposition became a shallower marine environment, or were redeposited from shallower environments into deeper by more turbid environments (Blanco-Bustamante et al., 1999). The presence of the conglomerates and polymictic elements may suggest closer proximity to a deltaic environment. Moreover, these suggest that at least some of the fauna reported for the Madruga Formation may have originated in shallower waters of a nearby shelf-forereef environment, transported or reworked by currents or other agents into deeper depositional environments interpreted in the current literature (Albear et al., 1985; Franco-Álvarez et al., 1992; Iturralde-Vinent, 2011a). This interpretation agrees with the overall trend in the orogenic activity of the Cuban terrain during the Paleocene-early/middle Eocene (Bralower and Iturralde-Vinent, 1997; Iturralde-Vinent and MacPhee, 1999; Iturralde-Vinent, 1995, 2011a, b; Iturralde-Vinent et al., 2016).

Limitations and final considerations

Our observations call attention to several limitations that must be considered for our current interpretation of this outcrop and its fauna. The construction of the highway highly disturbed the area of the outcrop. At the time of the Palmer and Bermúdez visits (the 1930's-1940's), those outcrops were well-exposed due to the recent highway construction but are currently severely deteriorated or disturbed by urbanization. Windows into the fossil-bearing strata are rare due to vegetation and soil cover (YCI unp. observations). Former research, however, did not take into consideration the anthropogenic alterations inflicted on the local stratigraphy by the construction of the roads (Figs. 1, 8, and 9). These modifications seem to have severely altered or mixed the external bedding at our collecting locality (Fig. 9), and the horizontality of several beds at several locations. The use of dynamite and heavy machinery likely affected the local stratigraphy and occurrence of its fossil fauna, which up to now has not been

considered in any of the published accounts of the outcrop since the 1920's (Figs. 8 and 9). These human-caused alterations could explain some of the intermixing of the pale Late Cretaceous clastics within some parts of the brown Madruga Formation. Furthermore, the historic photographs taken during the construction of the highway and overpass show the presence of a more lithified outcrop-base than the more friable, marly, polymictic sandstones covered by soil and vegetation of today. It is possible that large boulders now present in the outcrop were positioned there then. Even with all the external modifications caused by the building of the central highway on the Madruga Formation type locality, the mollusk specimens here described are referable to that formation. All these factors, and the presence of these hitherto unreported faunal components prompt a reassessment of the whole outcrop that could help clarify the stratigraphy, fauna, and age of the Madruga Formation and its formational history. Much more data is needed to test some of these initial observations and hypotheses. However, we are confident that such an investigation will reveal a far richer Madruga Formation than currently characterized.

CONCLUSIONS

The Paleocene bivalve fauna of Cuba has been scantily investigated, although outcrops of marine sedimentary rocks of this age are widespread throughout the island. One of the foremost Paleocene Cuban geologic units is the Madruga Formation, which occurs near the town of Madruga, in Mayabeque Province. This formation has produced diverse foraminifera, brachiopods, echinoids, and bivalves. This last group which up to now had not been reported. Thus, by recording the cardiid bivalve genera *Schedocardia* and *Acanthocardia*, we provide a first report for the Cuban fossil record. Historical insight gleaned from photographs taken during the construction of the highway call attention to the high level of disturbance and

anthropogenization which the outcrop has been subjected since the late 1920's and cautions as to the depositional and chronologic interpretation of the samples taken there since. Moreover, our observations could call for a reassessment of the stratotype outcrop of the Madruga Formation.

ACKNOWLEDGMENTS

We thank Manuel Iturralde-Vinent and Lazaro Viñola López (Florida Museum of Natural History) for their observations and comments that greatly improved earlier versions of the manuscript. Jan Johan ter Poorten (Naturalis Biodiversity Center) provided further corrections and suggestions, aided in the identification of *Schedocardia*, and kindly allowed us to use his photographs of USNM. Madruga city historian Carlos Miguel Suárez Sardiñas kindly provided important historical insight, historical photographs, and aid during the fieldwork. Sandy Ebersole and Lynn Harrell (both Geological Survey of Alabama) kindly provided images of the *S. hatchetigbeensis* valve in Figure 6.

REFERENCES

- Albear, J.F., M. Iturralde-Vinent, G. Furrázola-Bermúdez, and J.R. Sánchez-Arango. 1985. Contribución a la geología de las provincias de La Habana y Ciudad de la Habana. Editorial Científico-Técnica, La Habana, 155 pp.
- Aldrich, T.H. 1886. Preliminary report upon the Tertiary fossils of Alabama and Mississippi. Bulletin of the Geological Survey of Alabama 1 (I): 15–60.
- Bermúdez, P.J. 1938. Nueva especie de *Bulimina* del Cretácico Superior cubano. Memorias de la Sociedad Cubana de Historia Natural “Felipe Poey” 12(2): 80–90.

- 307 Bermúdez, P.J. 1950. Contribución al estudio del Cenozoico cubano. Memorias de la
- 308 Sociedad Cubana de Historia Natural “Felipe Poey” 19(3): 204–375.
- 309 Bermúdez, P.J. 1961. Las Formaciones Geológicas de Cuba. Ministerio de Industrias,
- 310 Instituto Cubano de Recursos Minerales, Geología Cubana, 177 pp.
- 311 Bralower, T. and M. Iturralde-Vinent 1997. Micropaleontological dating of the collision
- 312 between the North American continental margin and the Greater Antilles Volcanic Arc
- 313 in Western Cuba. *Palaios* 12: 133–150. doi: 10.2307/3515303.
- 314 Blanco-Bustamante, S., G. Fernández-Rodríguez, and R.H. Fluegeman. 1999. A note on the
- 315 Biostratigraphy of Paleocene-Eocene Larger Foraminifera from Western Cuba.
- 316 *Micropaleontology* 45(2): 19–26.
- 317 Cooper, G.A. 1955. New brachiopods from Cuba. *Journal of Paleontology*, 29(1): 64–70.
- 318 Cooper, G.A. 1979. Tertiary and Cretaceous brachiopods from Cuba and the Caribbean.
- 319 *Smithsonian Contributions to Paleobiology* 37: 1–56.
- 320 Cushman, J.A. and P.J. Bermúdez. 1948a. Some Paleocene foraminifera from the Madruga
- 321 Formation of Cuba. *Contributions for the Cushman Laboratory for Foraminiferal*
- 322 *Research* 24: 68–75.
- 323 Cushman, J.A., and P.J. Bermúdez. 1948b. Additional species of Paleocene foraminifera from
- 324 the Madruga Formation of Cuba. *Contributions for the Cushman Laboratory for*
- 325 *Foraminiferal Research* 24: 85–89.
- 326 Cushman, J. A. and P.J. Bermúdez. 1949. Some Cuban species of *Globorotalia*.
- 327 *Contributions for the Cushman Laboratory for Foraminiferal Research* 25:26-45.
- 328 De Golyer, E. 1918. The geology of Cuban petroleum deposits. *American Association of*
- 329 *Petroleum Geologists Bulletin* 2: 133–167.
- 330 Férussac, A. E. J. P. J. F. d'Audebard de. 1821-1822. Tableaux systématiques des animaux
- 331 mollusques classés en familles naturelles, dans lesquels on a établi la concordance de

332 tous les systèmes; suivis d'un prodrome général pour tous les mollusques terrestres ou
333 fluviatiles, vivants ou fossiles. - pp. j-xlvij [= 1-47], [1], 1-27, 1-110, [1]. Paris,
334 Londres. (Bertrand, Sowerby).

335 Fluegeman, R.H. and M.P. Aubry. 1999. Lower Paleogene biostratigraphy of Cuba.
336 Micropaleontology 45(2): 1–91.

337 Franco-Álvarez, G.L., M. Acevedo-González, H. Álvarez-Sánchez, C. Artime Peñeñori, A.
338 Barriento-Duarte, S. Blanco-Bustamante, M. Cabrera, et al. 1992. Léxico
339 Estratigráfico de Cuba. Centro de Nacional de Información Geológica, La Habana, pp.
340 658.

341 Furrázola-Bermúdez, G., C.M. Judoley, M.S. Mijailovskaya, Y.S. Miroljubov, I.P.
342 Novojatsky, A. Nuñez Jiménez, and J.B. Solsona. 1964. Geología de Cuba. Editorial
343 Nacional de Cuba, La Habana, 239 pp.

344 Garvie, C.L. 2013. Studies on the molluscan paleomacrofauna of the Texas Paleogene.
345 Bulletins of American Paleontology 384-386: 1–222.

346 Keen, A.M. 1980. The pelecypod family Cardiidae: A taxonomic summary. Tulane Studies in
347 Geology and Paleontology 16: 1–40.

348 Kier, P. 1984. Fossil spatangoid echinoids of Cuba. Smithsonian Contributions to
349 Paleobiology 55: 1–336.

350 Iturralde-Vinent, M. 1995. Cuencas sedimentarias del Paleoceno-Eoceno de Cuba. Boletín de
351 la Sociedad Venezolana de Geología 20 (1-2): 75–80.

352 Iturralde-Vinent, M. 2011a. Geología de las rocas del Arco Volcánico del Paleógeno. In:
353 Compendio de Geología de Cuba y del Caribe. Segunda Edición (published December
354 2012), DVD-ROM. Editorial CITMATEL, ISBN: 9-789592-572863.

355 Iturralde-Vinent, M. 2011b. Cuencas sinorogénicas del Campaniano tardío al Eoceno
356 Superior, desarrolladas sobre rocas del arco volcánico y máfico-ultramáficas. In:

357 Compendio de Geología de Cuba y del Caribe. Segunda Edición, DVD-ROM.

358 Editorial CITMATEL, ISBN: 9-789592-572863. Premio Academia de Ciencias de

359 Cuba.

360 Iturralde-Vinent, M. 2015. Estratigrafía de las cuencas sinorogénicas del Campaniano tardío

361 al Daniano desarrolladas sobre rocas de arco volcánico y retroarco en Cuba. Anuario

362 de la Sociedad Cubana de Geología 2: 27–50.

363 Iturralde-Vinent, M., A. García-Casco, Y. Rojas-Agramonte, J.A. Proenza-Fernández, J.B.

364 Murphy, and R.J. Stern. 2016. The geology of Cuba: A brief overview and synthesis.

365 GSA Today 26(10): 4–10, doi: 10.1130/GSATG296A.1.

366 Iturralde-Vinent, M. and R.D.E. MacPhee. 1999. Paleogeography of the Caribbean region,

367 implications for Cenozoic biogeography. Bulletin of American Museum Natural

368 History 238: 1–95.

369 Lamarck, J.B. 1809. *Philosophie Zoologique*. Musée d'Histoire Naturelle, Paris.

370 Lewis, J.W. 1932a. Geology of Cuba (with discussion by R. J. Metcalf). American

371 Association of Petroleum Geologists Bulletin 16(6): 533–555.

372 Lewis, J.W. 1932b. Occurrence of oil in igneous rocks in Cuba. American Association of

373 Petroleum Geologists Bulletin 16(8): 809–818.

374 Linnaeus, C. 1758. *Systema Naturae*. Laurentii Salvii, Homiae.

375 Moore, E.J. 2002. Family Cardiidae, 21 p. In: Tertiary marine pelecypods of California and

376 Baja California, Chapter F: <http://www.cmug.com/~chintimp/Cardiidae.htm>

377 Núñez-Cambra, K.E., and M. Iturralde-Vinent. 2016. Mapa geológico de la República de

378 Cuba. Ministerio de Energía y Minas. Instituto de Geología y Paleontología-Servicio

379 Geológico.

380 Özcan, E., S.F. Mitchell, G. Less, E. Robinson, J.R. Bryan, J. Pignatti, and A.O. Yücel. 2019.

381 A revised suprageneric classification of American orthophragminids with emphasis on

late Paleocene representatives from Jamaica and Alabama. *Journal of Systematic Palaeontology* 17: 1551-1579. doi.org/10.1080/14772019.2018.1539778.

Palmer, D.K. 1934. Some large Foraminifera from Cuba. *Memorias de la Sociedad Cubana de Historia Natural "Felipe Poey"* 8(4): 235–269.

Palmer, K.V. and D.C. Brann. 1965. Catalogue of the Paleocene and Eocene Mollusca of the southern and eastern United States. Part 1. Pelecypoda, Amphineura, Pteropoda, Scaphopoda, and Cephalopoda. *Bulletins of American Paleontology* 48: 1–471.

Palmer, R.H. 1932. Informe geológico del área de Madruga. Oficina Nacional de Recursos Minerales, Ministerio de Energía y Minas, La Habana (unpublished), 27 pp.

Palmer, R.H. 1948. List of Palmer Cuban fossil localities. *Bulletins of American Paleontology* 31(128): 1–178.

Sánchez-Roig, M. 1949. Paleontología Cubana I: Los Equinodermos Fósiles de Cuba. Compañía Editora de Libros y Folletos, La Habana 1: 1–330.

Sachs, K.N. 1957. Restudy of some Cuban larger Foraminifera. *Contributions from the Cushman Foundation for Foraminiferal Research* 8(3): 106–120.

Sessa, J.A., T.J. Bralower, M.E. Patzkowsky, J.C. Handley, and L.C. Ivany. 2012. Environmental and biological controls on the diversity and ecology of Late Cretaceous through early Paleogene marine ecosystems in the U.S. Gulf Coastal Plain. *Paleobiology* 38(2): 218–239.

Schneider, J. 2002. Phylogeny of cardiid bivalves (cockles and giant clams): revision of the Cardiinae and the importance of fossils in explaining disjunct biogeographical distributions. *Zoological Journal of the Linnean Society* 136: 321–369.

Toulmin, L.D. 1977. Stratigraphic Distribution of Paleocene and Eocene Fossils in the Eastern Gulf Coast Region. Geological Survey of Alabama, Monograph 13(1): 1–602.

Wright, A. and P.W.K. Sweet. 1924. The Jurassic as a source of oil in western Cuba. Bulletin of the American Association of Petroleum Geologists 8(4): 516–519.

Woodring, W.P. 1982. Geology and paleontology of Canal Zone and adjoining parts of Panama: Description of Tertiary mollusks (pelecypods: Propeamussiidae to Cuspidariidae). United States Geological Survey Professional Paper 306(F): 541–759.

CAPTIONS

Figures

Figure 1: Collecting locality of the Madruga Formation, near the city of Madruga, Province of Mayabeque, Cuba. 1.1 Mayabeque province, 1.2 Madruga municipality, and 1.3 Madruga Formation type locality indicated by the red arrow near the railroad bridge. Historical photographs of the construction of the road under the railroad bridge, where the exposed outcrops of the Madruga Formation can be seen during construction (1.4) and after (1.5).

Figure 2: Geological map indicating the outcrops associated with the Madruga Formation at the collection locality discussed (marked with an asterisk *). Madruga Formation (md), Paleocene (P₁₋₃), Nazareno Formation (nz) lower Eocene (P₄₋₅), Peñalver Formation (pñ) Upper Cretaceous (KTB/K₁₋₅), Via Blanca Formation (vb), Upper Cretaceous (Campanian-Maastrichtian, K₅₋₆), Cojimar Formation (cjr) lower Miocene (N₁₋₂), and Güines Formation (gn) middle-upper Miocene (N₃₋₄). UB indicates ultrabasic ophiolite outcrops.

Figure 3: *Schedocardia cf. hatchetigbeensis* (MPAL-1001) right valve, incomplete external mold in the matrix, collected from the Madruga Formation.

Figure 4: Close-ups of the *Schedocardia* cf. *hatchetigbeensis* (MPAL-1001). Note the ovoid foramen on the valve, and the smaller foramina on the rib tops, suggesting alternating rows of spines.

Figure 5: *Schedocardia hatchetigbeensis* syntypes (USNM 638802 =*Cardium hatchetigbeense* and USNM 645087; type =*Cardium hatchetigbeense*) from the Hatchetigbee Formation at Hatchetigbee Bluff, Tombigbee River, Washington County, Alabama. Two specimens in the center row are stereopairs. Photographs used with permission of Jan Johan ter Poorten.

Figure 6. Left valve of *Schedocardia hatchetigbeensis*. Specimen image courtesy of Geological Survey of Alabama. Specimen catalog number GSA-I8608 labelled as *Acanthocardia hatchetigbeensis* from the Hatchetigbee Formation at Hatchetigbee Bluff, Tombigbee River, Washington County, Alabama,

Figure 7: *Acanthocardia* sp. (MPAL-1002) external mold in matrix, collected from the indicated outcrop of the Madruga Formation.

Figure 8: Historical photograph of the road under the railroad bridge during construction in 1928.

Figure 9: Modern photograph of a section of the Madruga Formation outcrop, at the type locality, in which the exposed stratigraphic column is visible, along with the effect of the use of explosives on the natural stratigraphy. The specimens discussed here were not collected at this precise location.

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6.

475

476 Figure 7.

477

478 Figure 8.

479

480

481 Figure 9.

482