1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23

Title: The intrinsic neonatal hippocampal network: rsfMRI findings

Abbreviated Title: The intrinsic neonatal hippocampal network

Authors:

Athena L. Howell (contribution: running registration and fMRI analyses, statistical analyses and writing the manuscript),
David E. Osher (preprocessing fMRI data and editing the manuscript),
Jin Li (preprocessing fMRI data and editing the manuscript),
and Zeynep M. Saygin (PI/advisor, guiding the research and analyses, checking results and editing the manuscript)

The Ohio State University, 43210

Please address correspondence to:

Athena L. Howell, The Ohio State University, 225 Psychology Building, 1835 Neil Avenue,
Columbus, OH, 43210. E-mail: howell.551@osu.edu

# of pages: 19 (in body of manuscript)
# of Figures: 5 Figures
# of words in abstract: 201

- 24 # of words in introduction: 575
- 25 # of words in discussion: 1077
- 26 Acknowledgements; The authors declare no competing financial interests

Abstract

Many adults cannot voluntarily recall memories before the ages of 3-5, a phenomenon referred to as “infantile amnesia”. The development of the hippocampal network likely plays a significant part in the emergence of the ability to form long-lasting memories. In adults, the hippocampus has specialized and privileged connections with certain cortical networks, which presumably facilitate its involvement in memory encoding, consolidation, and retrieval. Is the hippocampus already specialized in these cortical connections at birth? And are the topographical principles of connectivity (e.g. long-axis specialization) present at birth? We analyzed resting-state hippocampal connectivity in neonates scanned within one week of birth (Developmental Human Connectome Project) and compared them to adults (Human Connectome Project). We explored the connections of the whole hippocampus and its long-axis specialization to seven canonical cortical networks. We found that the neonatal hippocampal networks show clear immaturity at birth: adults showed hippocampal connectivity that was unique for each cortical network, whereas neonates showed no differentiation in hippocampal connectivity across these networks. Further, neonates lacked long-axis specialization (i.e., along anterior-posterior axis) of the hippocampus in its differential connectivity patterns to the cortical networks. This immaturity in connectivity may contribute to immaturity in memory formation in the first years of life.

50 “New and Noteworthy”:

51 While animal data, and anatomical and behavioral human data from young children
52 suggest that the hippocampus is immature at birth, to date, there are no direct assessments of
53 human hippocampal functional connectivity (FC) very early in life. Our study explores the FC of
54 the hippocampus to the cortex at birth, allowing insight into the development of human memory
55 systems.

56

Introduction

Many adults cannot voluntarily recall memories before the ages of 3-5, a phenomenon referred to as “infantile amnesia” (Alberini & Travaglia, 2017). One potential reason for this is that the hippocampus (the primary brain structure responsible for episodic memory formation in adults) and its connections with the rest of the brain may be immature at birth. Indeed, the hippocampus does appear to be immature at birth; evidence in macaques suggests it continues to mature after one year of age (roughly age 3-5 in humans) (Jabés et al., 2011) and human data indicates that volumetric and structural changes in the hippocampus continue through childhood (DeMaster et al., 2014; Gilmore et al., 2012; Seress 2007). Further, episodic memory performance may be influenced by changes in the patterns of hippocampal connectivity from middle childhood to adulthood, including along the long-axis of the hippocampus (Blankenship et al., 2017; DeMaster et al., 2014; Ghetti et al., 2010; Gogtay et al., 2006; Poppenk & Moscovitch, 2011; Riggins et al., 2016). At younger ages, hippocampal gray matter volume has been linked to early language ability (Can et al., 2013) and one recent study showed potential hippocampal activation for learned items in 2-year old toddlers (Prabhakar et al., 2018). However, the intrinsic connectivity of hippocampus very early in life is less well understood. Therefore, an understanding of the hippocampal network at birth and its development may lead to greater understanding of memory development.

Recently, Wael and colleagues (2018) showed the hippocampus has a clear intrinsic pattern of functional connectivity (FC) to a set of cortical networks in adults. Specifically, they showed higher (i.e. most positive) connectivity from the hippocampus to the Default Mode and Limbic networks and lowest (i.e. least positive) connectivity to the Frontoparietal and Ventral Attention networks (from Yeo et al., 2011). Further, this connectivity pattern differed between the anterior

and posterior portions of the hippocampus, with anterior hippocampus showing larger differences in connectivity to the networks than posterior hippocampus. This so-called long-axis specialization of the hippocampus is consistent with previous research showing that the anterior and posterior hippocampus display different patterns of structural and functional connectivity and may be uniquely activated in response to cognitive, memory and spatial demands (for reviews see Poppenk et al., 2013, Strange et al., 2014). The development of the hippocampal network and the long-axis gradient likely plays a significant part in the emergence of the ability to form long-lasting memories. For instance, the work of Riggins et al. (2016) examines the relationship of anterior/posterior connectivity and episodic memory in 4- and 6-year old children and finds developmental differences even between these two ages. Although this work in young children is notable, the fact remains that we know very little about the hippocampus, its connections, and its relationship to memory-formation during the earliest stages of life.

To this end, we compared the resting-state hippocampal connectivity patterns to a set of cortical networks in neonates and adults. Resting state connectivity, determined by spontaneously correlated activity of disparate brain regions, is used as a reliable marker of intrinsic functional connectivity (FC) between brain regions (Biswal et al., 1995; Raichle, 2009; Smith, 2013; Sporns, 2013); further, FC at rest is predictive of task-based activity (Cole et al., 2014; Osher et al., 2019; Smith et al., 2009; Tobyn et al., 2018).

More recently, developmental studies using FC have shown the FC of some networks is mature at birth while others take months or longer to become adultlike (for reviews see Gao et al., 2017 and Grayson & Fair, 2017). In particular, multiple studies indicate the connectivity of visual and somatomotor networks is not only functional but highly adult-like at birth (Gao et al., 2015b; Lin et al., 2008; Liu et al., 2008). Other areas, such as the default mode network, dorsal attention

network, frontoparietal network, and some perceptual regions show relatively immature functional and structural characteristics at birth and experience large modifications postnatally (Gao et al., 2015b; Natu et al., 2019), although the frontoparietal network may have important functional roles even within the first year of life (e.g. Linke et al., 2018).

To assess hippocampal maturity at birth, we analyzed FC between seven intrinsic networks and the hippocampus as a whole as well as along the hippocampal long-axis in both neonates and adults. We also compared neonatal vs. adult hippocampal connectivity to the cortex at a finer, voxelwise scale. Based on previous literature suggesting the immaturity of the hippocampus at birth, we hypothesized that neonates would differ from adults in their hippocampal connectivity to the cortex, particularly to the more immature networks (e.g. default mode and frontoparietal).

Materials and Methods

Participants

Neonates:

Neonatal data comes from the initial release of the Developing Human Connectome Project (dHCP) (<http://www.developingconnectome.org>, Markopoulos et al., 2018). Neonates were recruited and imaged in London at the Evelina Neonatal Imaging Centre after gathering informed parental consent to image and release the data. The study was approved by the UK Health Research Authority. 40 neonates were included in our analyses (15 female, 36-44 weeks old at scan).

Adults:

Adult data comes from the Human Connectome Project (HCP), WU-Minn HCP 1200 Subject Data Release (<https://www.humanconnectome.org/study/hcp-young-adult>, Van Essen et al., 2013). Participants were scanned at Washington University in St. Louis (WashU). We included

40 participants in our analyses (15 female; 20-35 years old). These adult participants were motion-matched to the neonates. Specifically, we matched each neonatal participant with an adult from the HCP dataset with the same gender who showed the most similar motion parameter (i.e., framewise displacement, FD). While the resulting group of adults were motion-matched to the neonates, we found that the groups were significantly different in the average gray-matter tSNR, with neonates exhibiting higher tSNR values ($t(78)=-6.8774, p=1.3469 \times 10^{-9}$). To ensure that any results were not driven by tSNR differences between groups, we identified an additional group of HCP adults whose tSNR was matched to the tSNR of the neonates ($t(78)=-1.5237, p=.132$) and replicated our results (see Extended Data, Figure 1-1).

MRI Acquisition

Neonates:

All acquisition information comes from the dHCP data release documentation. Imaging was carried out on a 3T Philips Achieva (running modified R3.2.2 software) using an imaging system specifically designed for neonates with a 32 channel phased array head coil (Hughes, E.J., et al.). Neonates were scanned during natural sleep; resting-state FC patterns have been shown to stay largely consistent while awake, asleep, or under anesthesia (Liu et al., 2015; Larson-Prior et al., 2009).

Resting-state fMRI

High temporal resolution fMRI developed specifically for neonates was collected using multiband (MB) 9x accelerated echo-planar imaging (TE/TR=38/392ms, voxel size = 2.15 x 2.15 x 2.15mm³). The resting state scan lasted approximately 15 minutes and consisted of 2300 volumes

for each run. No in-plane acceleration or partial Fourier transform was used. Single-band reference scans with bandwidth matched readout and additional spin-echo acquisitions were also acquired with both AP/PA fold-over encoded directions.

Anatomical MRI

High-resolution T2-weighted and inversion recovery T1-weighted multi-slice fast spin-echo images were acquired with in-plane resolution $0.8 \times 0.8\text{mm}^2$ and 1.6mm slices overlapped by 0.8mm (T2-weighted: TE/TR= 156/12000ms; T1 weighted: TE/TR/TI = 8.7/4795/1740ms)

Adults:

All acquisition information comes from the HCP data release documentation. Scanning for the 1200 WU-Minn HCP subject was carried out on a customized 3T Connectome Scanner adapted from a Siemens Skyra (Siemens AG, Erlanger, Germany), equipped with a 32-channel Siemens receiver head coil and a “body” transmission coil specifically designed by Siemens to accommodate the smaller space (due to special gradients) of the WU-Minn and MGH-UCLA Connectome scanners.

Resting-State fMRI

Participants were scanned using the Gradient-echo EPI sequence (TE/TR = 33.1/720ms, flip angle = 52° , 72 slices, voxel size = $2 \times 2 \times 2\text{mm}^3$). Scanning lasted approximately 15 minutes consisting of 1200 volumes for each run. Each participant finished two resting-state fMRI sessions. For each session, two phases were encoded: one right-to-left (RL) and the other left-to-right (LR). For our analyses, we used the LR phase encoding from the first session. Participants were

instructed to relax and keep their eyes open and fixated on a bright, projected cross-hair against a dark background.

Anatomical MRI

High-resolution T2-weighted and T1-weighted images were acquired with an isotropic voxel resolution of 0.7mm³ (T2-weighted 3D T2-SPACE scan: TE/TR=565/3200ms; T1-weighted 3D MPRAGE: TE/TR/TI = 2.14/2400/1000ms).

MRI Preprocessing

Neonates:

The dHCP data was preprocessed using the dHCP minimal preprocessing pipelines (Makropoulos et al., 2018). Anatomical MRI preprocessing included bias correction, brain extraction using BET from FSL (FMRIB Software Library) and segmentation of the T2w volume using their DRAW-EM algorithm (Makropoulos et al., 2014). The resulted gray and white matter segmentations were used as anatomical masks in further analyses; these masks were manually checked for accuracy.

Minimal preprocessing for the resting-state fMRI included (Fitzgibbon et al., 2016) distortion correction, motion correction, 2-stage registration of the MB-EPI functional image to the T2 structural image, temporal high-pass filtering (150s high-pass cutoff), and ICA denoising using FSL's FIX (Salimi-Khorshidi, et al., 2014). In addition to this minimal preprocessing, we smoothed the data (Gaussian filter, FWHM = 3mm) across the gray matter, and applied a band-pass filter at 0.009-0.08 Hz. To further denoise the data, we used aCompCor (Behzadi et al., 2007)

to regress out physiological noise (heartbeat, respiration, etc.) from the white matter and cerebrospinal fluid (CSF).

Adults:

HCP data was preprocessed using the HCP minimal preprocessing pipelines (Glasser et al., 2013). For the anatomical data, a Pre-FreeSurfer pipeline was applied to correct gradient distortion, produce an undistorted “native” structural volume space for each adult participant by ACPC registration (hereafter referred to as “acpc space”), extract the brain, perform a bias field correction, and register the T2-weighted image to the T1-weighted image. Additionally, each participant’s brain was aligned to a common MNI152 template brain (with 0.7mm isotropic resolution). Then, the FreeSurfer pipeline (based on FreeSurfer 5.3.0-HCP) was performed with a number of enhancements specifically designed to capitalize on HCP data (Glasser et al., 2013). The goal of this pipeline was to segment the volume into predefined structures, to reconstruct the white and pial cortical surfaces, and to perform FreeSurfer’s standard folding-based surface registration to their surface atlas (fsaverage).

For the resting-state fMRI data, minimal functional analysis pipelines included: removing spatial distortions, motion correction, registering the fMRI data to structural and MNI152 templates, reducing the bias field, normalizing the 4D image to a global mean, and masking the data with the final brain mask. After completing these steps, the data were further denoised using the ICA-FIX method (Salimi-Khorshidi, et al., 2014). To mirror the adult and neonatal preprocessing pipelines, we unwarped the data from MNI152 to acpc space, allowing both groups to be analyzed in “native” space. We then applied spatial smoothing (Gaussian filter, FWHM = 3mm) within the gray matter, band-pass filtered at 0.009-0.08 Hz and implemented aCompCor to regress out physiological noise, just as we did with the neonates.

All subsequent analyses in neonates and adults were performed in each subject's native space, except for the whole-brain voxelwise analysis.

Connectivity analyses

We used the 7-network cortical parcellation identified by Yeo et al. (2011). For the whole-hippocampus and long-axis analyses, the hippocampal label was binarized from FreeSurfer's (surfer.nmr.mgh.harvard.edu) `aparc+aseg` parcellation and visually inspected for accuracy in each subject. For the first long-axis gradient analysis this label was further sectioned into anterior and posterior portions via manual segmentation using FreeSurfer, with the uncus apex as the dividing marker (Poppenk & Moscovitch., 2011). All labels (cortical networks, hippocampal labels) were originally in CVS average-35 MNI152 space and then registered to each individual subject's anatomical data using ANTs (Advanced Normalization Tool) 3dWarpMultiTransform (ANTs version 2.1.0; <http://stnava.github.io/ANTs>; Avants et al., 2011). ANTs is routinely used for developmental dataset registrations (Alexander et al., 2019; Dean et al., 2018). The resulting registrations were checked for accuracy. Similarly, for the long-axis gradient analysis, the hippocampal label in CVS was split into nine equally spaced "slices" along the anterior-posterior axis. Using the same ANTs registration technique for all ROIs provided an extra measure of consistency between groups and between analyses; however, as an added quality check we ran our whole-hippocampus to network analysis using the binarized hippocampal label provided by the dHCP and HCP for each individual. These second results are nearly identical (see Extended Data, Figure 2-1) to the first (Figure 2) thus increasing confidence that our results are not due to registration error.

After registration to the anatomical data, we registered the labels onto the functional data in neonates using an inverse warp of the `func2anat` matrix provided by the dHCP. In adults, the

labels in acpc space after ANTs registration were then resampled to 2mm cubic voxels to align with the functional data. We manually checked individuals from each sample to ensure the accuracy and fit of the labels to the individual functional data. We extracted the BOLD activation in each label over the time course, averaged within each label, and correlated the hippocampal activity—first whole hippocampus, then along the long-axis (for both anterior-posterior and gradient slices)—with activity in each of the 7 networks to create a Fisher’s Z-scored correlation matrix using Matlab 2018b (The MathWorks, Inc., Natick, Massachusetts, United States). We also explored differences in the hippocampal connectivity to the whole cortex at a voxelwise scale between adults and neonates to determine whether specific regions within the networks were driving adult-neonate differences. Hippocampal connectivity to the cortex was calculated by correlating the average hippocampal signal and the signal of each voxel within the cortical gray matter mask during the time course for each individual in functional space. To compare the connectivity between adults and neonates, images from both groups were registered to the template space (i.e., CVS average-35 MNI152) before running a between-group analysis. Although this is the only template-space analysis we performed, template-space analyses have been routinely performed to compare infants to adults using similar registration methods (e.g. Gao et al., 2009; Gao et al., 2015a).

Experimental Design and Statistical Analyses

Where t-tests were performed between regions we corrected for multiple comparisons using the Holm-Bonferroni correction (Holm, 1979); all connectivity values were Fisher’s Z transformed (Fisher, 1915) to normalize the data.

Before doing any of the planned analyses, we first performed data quality checks. To make sure there was no significant motion difference between groups, we calculated the framewise

displacement (FD) (Power et al., 2012) based on the six motion parameters estimated from a rigid-body transformation provided by dHCP and HCP. We manually checked the registration of the gray and white matter masks as well as the network and hippocampal labels in the adults and neonates to the registration was accurate. Because we are performing comparisons of correlations between groups, we next wanted to ensure that the correlation distributions were similar and were normally distributed in both neonates and adults; we did this by assessing the correlation of each voxel to every other voxel in the brain and plotting the distribution of those correlations. We also performed between-subject reliability of correlation matrices within and across the adult and neonate groups. We calculated the connectivity of each region (i.e. each of the seven networks and the hippocampus) to every other region for each subject. This connectivity matrix was then correlated with every other subject's value either between- or within-groups to assess inter-subject reliability; in other words, we correlated the connectivity of every adult to every other adult (within-group) and neonate to neonate, as well as comparing every adult to every neonate (between-group).

Our first analysis examined the relationship of the whole hippocampus to the seven cortical networks. After running a one-way ANOVA with network as the independent variable and connectivity as the dependent variable for both groups, we computed pairwise comparisons between each unique combination of connectivity values to the networks (e.g. Hipp-Lim vs Hipp-VA) to determine networks with significantly different FC to the hippocampus (Snedecor and Cochran, 1989). Rose plots comparing the connectivity pattern of adults and neonates were created by subtracting the mean connectivity across all networks from each individual network (for adults and neonates separately) and plotting the resulting magnitude to show the relative connectivity

patterns of the hippocampus to the networks for each group and to compare these patterns between groups.

For our hippocampal-cortical voxelwise analysis, we used FSL's randomise function to compare between groups and perform permutation testing (to correct for multiple comparisons) in order to determine areas of greater connectivity in adults vs neonates and visa versa. After mapping the individual correlation matrices from subject space into a common template space, we used randomise with default 5000 permutations and clustered the results using FSL's threshold-free cluster enhancement (TFCE), which corrects for family-wise error (FWE). This produced a list of potential clusters with each cluster's associated p-value; the p-values were then thresholded at a $p < 0.0005$, and only those clusters that remained significant after that point are reported in this paper.

For the first long-axis hippocampus analysis, we first computed a two-way ANOVA in each group (separately) using location (i.e. anterior or posterior hippocampus) and network as independent variables and FC as the dependent variable. Pairwise comparisons were then made between the anterior and posterior FC values to each network for each group (e.g. adult antHipp-Lim vs adult postHipp-Lim). For the second long-axis analysis, we conducted a two-way ANOVA at each slice using group and network as independent variables and connectivity as the dependent variable. We also computed a one-way ANOVA at each slice for each group with network as the independent variable. As in the whole-hippocampal analysis, rose plots were created by subtracting out the mean connectivity to all networks (e.g. mean connectivity of adult anterior hippocampus to all networks) from each network and group in the anterior and posterior labels individually to demonstrate comparative connectivity differences between the anterior and posterior regions in each group.

Results

(FIGURE 1 HERE)

Preliminary data-checks

Comparison of the framewise displacement in adults and neonates showed no significant difference of FD between adults and neonates ($t(78)=-0.48$, $p=0.63$). Visual inspection of the gray and white matter masks (which are critical for resting-state preprocessing) in Figure 1a shows they are accurately delineating gray/white matter in both neonates and adults; the cortical networks and hippocampal labels also appear to be correctly localized, suggesting that the regions are accurately identified in both neonates and adults (Figure 1a). Figure 1b demonstrates that both neonates and adults have normally-distributed correlation values that are centered around 0. Between-subject reliability of correlation matrices within and across the adult and neonate groups showed the connectivity matrices (i.e. region-to-region connectivity of each of the seven networks and the hippocampus to each other) of each adult subject to each other adult subject were highly correlated, as were the matrices of each neonate subject to each other neonate subject, and a pairwise comparison of subject variability within groups (e.g. adult-adult correlations compared to neonate-neonate correlations) was not significant ($t(78)=0.76$, $p=0.45$). But subject-to-subject correlations across the two groups were significantly lower than the within-group correlations (adult-adult vs adult-neo $t(78) = 14.09$, $p=3.87 \times 10^{-23}$, neonate-neonate vs adult-neonate ($t(78)=11.95$, $p=2.63 \times 10^{-19}$) suggesting that while the connectivity data are reliable, neonates have different connectivity patterns than adults.

Whole Hippocampus

We first explored the connectivity of the whole hippocampus to the cortical networks. In adults, there was a main effect of network suggesting that some networks are more strongly

connected with the hippocampus than others (Figure 2; one-way ANOVA, $F(6,273)=47.11$, $p=1.84 \times 10^{-39}$). Subsequent pairwise comparisons showed a clear hierarchy of connectivity, such that hippocampal connectivity was highest to the Limbic (Lim) network (vs hippocampal connectivity to: Ventral Attention or VA ($t(78)=12.95$, $p_{HB}=8.42 \times 10^{-20}$); FrontoParietal or FP ($t(78)=11.76$, $p_{HB}=1.20 \times 10^{-17}$), Dorsal Attention or DA ($t(78)=10.09$, $p_{HB}=1.50 \times 10^{-14}$); Visual or Vis ($t(78)=7.20$, $p_{HB}=4.52 \times 10^{-9}$); and SomatoMotor or SM ($t(78)=5.97$, $p_{HB}=7.91 \times 10^{-7}$)). Hippocampal connectivity to the Default Mode Network (DM) was higher than hippocampal connectivity to: VA ($t(78)=10.32$, $p_{HB}=5.80 \times 10^{-15}$); FP ($t(78)=9.07$, $p_{HB}=1.33 \times 10^{-12}$); DA ($t(78)=7.24$, $p_{HB}=4.04 \times 10^{-9}$); Vis ($t(78)=4.63$, $p_{HB}=1.45 \times 10^{-4}$); and SM ($t(78)=3.16$, $p_{HB}=0.014$)). Hippocampal-SM connectivity was 3rd highest, and higher than hippocampal connectivity to: VA ($t(78)=7.83$, $p_{HB}=3.19 \times 10^{-10}$); FP ($t(78)=6.46$, $p_{HB}=1.09 \times 10^{-7}$); and DA ($t(78)=4.35$, $p_{HB}=3.61 \times 10^{-4}$)). Hippocampal-Vis connectivity was the next highest (vs VA ($t(78)=5.49$, $p_{HB}=5.31 \times 10^{-6}$); FP ($t(78)=4.16$, $p_{HB}=6.38 \times 10^{-4}$), and connectivity with DA was higher than with VA ($t(78)=3.89$, $p_{HB}=1.47 \times 10^{-3}$). In summary, hippocampal connectivity was highest to Lim, followed by DM, then SM, Vis, and DA; hippocampal connectivity was lowest (i.e. negatively correlated) with the FP and VA networks. In previous literature, the hippocampus is occasionally included as a part of the DM network; our finding of high Hippocampal-DM correlation and anti-correlation between the hippocampus and attention (i.e. FP and VA) networks falls in line with earlier work on the connectivity of the DM network (e.g. Buckner, Andrews-Hanna & Schacter, 2008) and is a good sign of the reliability of our results.

In contrast to the adult pattern, although neonates did show a main effect of network ($F(6,273)=5.12$, $p=2.27 \times 10^{-5}$), pairwise comparisons indicated that only the Lim and SM networks significantly differ from the rest, with significantly greater connectivity from the

hippocampus to Lim vs DA ($t(78)=5.31$, $p_{HB}=2.15 \times 10^{-5}$) and Lim vs FP ($t(78)=4.22$, $p_{HB}=1.33 \times 10^{-3}$) and significantly greater connectivity to SM vs DA ($t(78)=3.35$, $p_{HB}=0.023$).

Pairwise comparisons between adults and neonates showed significant differences between the groups, with significantly less connectivity in adults to Vis ($t(78)=-2.64$, $p_{HB}=0.040$), DA ($t(78)=-2.77$, $p_{HB}=0.035$), FP ($t(78)=-5.33$, $p_{HB}=5.49 \times 10^{-6}$) and VA ($t(78)=-8.62$, $p_{HB}=5.86 \times 10^{-13}$) networks compared to neonates.

(FIGURE 2 HERE)**Hippocampus to Cortex voxelwise analysis**

We next explored the connectivity of the hippocampus to the entire cortex at a voxelwise scale; because our previous analysis only focused on 7 canonical networks, we may have missed differences between neonates and adults at a finer grain than that seen on a network level. Thresholding the unpaired t-test results of the whole-brain clusters at $p < 0.0005$ produced 26 significant FWE-corrected (Smith & Nichols, 2009) clusters in the neonates > adults comparison (i.e. 26 clusters where neonatal hippocampal FC significantly exceeds adult hippocampal FC) and 14 significant clusters in the adults > neonates comparison (Figure 3). Specifically, neonates show greater hippocampal FC to frontal and parietal areas, bilateral lingual and pericalcarine cortex and cuneus when compared to adults; frontoparietal differences were particularly prevalent within the right hemisphere. Adults, on the other hand, displayed greater hippocampal FC than the neonates primarily to bilateral isthmus cingulate and precuneus. Cluster sizes and indices for clusters greater than 200 voxels along with peak voxel location and associated brain regions are reported in Figures 3-1 and 3-2 and largely follow the results from the 7-network analysis—the neonatal hippocampus shows greater FC to frontoparietal and attention-relevant areas, whereas the adult hippocampus shows greater FC with regions associated with the default mode and limbic networks.

(FIGURE 3 HERE)

Anterior-Posterior Hippocampus

We next explored the anterior vs. posterior hippocampal connectivity patterns in neonates and adults; previous literature in both humans and other animals suggest functional differentiation of the anterior and posterior hippocampal segments, and thus we may expect these segments to have differences in FC to the 7 cortical networks. In adults, a two-way ANOVA indicated a main effect of network ($F(6,546)=60.04$, $p=5.04 \times 10^{-57}$) and an interaction between network and anterior/posterior hippocampus ($F(6,546)=14.31$, $p=3.54 \times 10^{-15}$) (Figure 4). In neonates, the two-way ANOVA showed only a significant main effect for network ($f(6,546)=7.67$, $p=6.30 \times 10^{-8}$) (Figure 4). Pairwise comparisons between the anterior and posterior portions of the hippocampus in adults show greater anterior vs posterior connectivity to the Lim ($t(78)=3.53$, $p_{HB}=0.0035$), DMN ($t(78)=2.38$, $p_{HB}=0.03$) and SM ($t(78)=3.19$, $p_{HB}=0.0082$) networks, and decreased anterior vs posterior connectivity to the DA ($t(78)=-3.07$, $p_{HB}=0.0087$), Frontoparietal ($t(78)=-5.79$, $p_{HB}=9.99 \times 10^{-7}$) and VA ($t(78)=-3.92$, $p_{HB}=0.0011$) networks. These results suggest the anterior hippocampus was primarily driving the negative correlations with VA & FP seen at the level of the whole hippocampus in adults. Neonates, however, show no significant differences between the anterior and posterior portions of the hippocampus to any of the networks, suggesting no differentiation/specialization of the hippocampal segments in their connections to the rest of the brain.

(FIGURE 4 HERE)

Long-Axis Gradient

Finally, we investigated the long-axis gradient, which has been demonstrated to map onto a differential functional gradient of the hippocampus. We broke up the hippocampus in each

subject into 9 different segments along the anterior-posterior axis and compared the 7-network connectivity to these segments in neonates and adults. Adults showed clear differentiation of network connectivity along the long-axis while neonates showed no clear differentiation (Figure 5). The Lim and DM in adults appeared to have an initial rise and fall of FC along the anterior-posterior gradient of the hippocampus which differentiated them from the Vis, SM, and DA, and the FP and VA showed a similar rise and fall of negative FC along the gradient. One-way ANOVAs for adults and neonates at each slice indicated a main effect of network in adults in all but the most posterior slice (Slice 1 ($F(6,273)=23.61$, $p=1.89 \times 10^{-22}$), Slice 2 ($F(6,273)=40.45$, $p=4.19 \times 10^{-35}$), Slice 3 ($F(6,273)=50.09$, $p=2.61 \times 10^{-41}$), Slice 4 ($F(6,273)=49.56$, $p=5.48 \times 10^{-41}$), Slice 5 ($F(6,273)=49.07$, $p=1.11 \times 10^{-40}$), Slice 6 ($F(6,273)=25.10$, $p=1.12 \times 10^{-23}$), Slice 7 ($F(6,273)=13.40$, $p=2.57 \times 10^{-13}$), Slice 8 ($F(6,273)=5.51$, $p=2.12 \times 10^{-5}$)). In the neonates, there was no main effect of network in any of the slices (at $p < 0.001$). To compare between the two groups, we performed two-way ANOVAs (with network and group as independent variables and FC value as the dependent variable) for each of the 9 slices. There was a significant interaction between network and group for the anterior 7 slices (Slice 1 ($F(6,546)=7.30$, $p=1.64 \times 10^{-7}$), Slice 2 ($F(6,546)=16.25$, $p=2.95 \times 10^{-17}$), Slice 3 ($F(6,546)=18.98$, $p=3.99 \times 10^{-20}$), Slice 4 ($F(6,546)=17.22$, $p=2.83 \times 10^{-18}$), Slice 5 ($F(6,546)=17.93$, $p=5.05 \times 10^{-19}$), Slice 6 ($F(6,546)=5.79$, $p=7.26 \times 10^{-6}$), Slice 7 ($F(6,546)=4.87$, $p=7.27 \times 10^{-5}$) and Slice 8 ($F(6,546)=3.89$, $p=8.26 \times 10^{-4}$), but no group differences for the most posterior slice. These results show that the biggest differentiation of hippocampal connectivity to the 7 networks occurs in the anterior 2/3s of the hippocampus in adults and that neonates do not show this differentiation.

(FIGURE 5 HERE)

Discussion

Our results show that the intrinsic connectivity of the hippocampal network is not fully mature at birth. Previous functional and volumetric evidence in both non-human primates and humans suggests that the hippocampus continues to develop beyond one year of age, even into middle childhood (Blankenship et al., 2017; Jabes et al., 2011; Keresztes et al., 2018; Lavenex and Banta Lavenex, 2013; Riggins et al., 2016). Although there is some evidence to suggest that the hippocampus is playing a key role in memory formation even early on in rodents (Alberini & Travaglia, 2017; Travaglia et al., 2018), it has been suggested that the long-lasting memories of very young children may be created in a fundamentally different way from adult long-term memories and may rely on cortical mechanisms rather than the traditional hippocampal method (Ellis & Turke-Browne, 2018; Gómez & Edgkin, 2016). Interestingly, multiple studies comparing preterm to term infants show no differences in gray matter volume in the hippocampus with decreased gestational age, implying the better part of hippocampal growth is accomplished prior to birth (Alexander et al., 2019; Ge et al., 2015; Thompson et al., 2008); our work suggests that although the physical bulk of the hippocampus may exist at birth, its connections do not. Specifically, the hippocampus does not have preferential connectivity to any particular network at birth and lacks any long-axis gradient of connectivity, suggesting that the hippocampus, the cortical networks it interacts with, or some combination of both, are immature at birth and may therefore be unable to form long-term memories using adult-like mechanisms. Indeed, the cortex itself is still maturing early on (e.g. Gao et al., 2015b; Ofen et al., 2007; Salzwedel et al., 2019) and it is likely this cortical immaturity, in addition to hippocampal immaturity, is contributing to the differences in memory formation between adults and neonates.

Adults showed a clear hierarchy of FC to the seven networks (consistent with Wael et al. (2018)), whereas neonates lacked this hierarchy. Further, the comparison between adults and

neonates shows significant differences between groups to all networks except the SM network, and only marginally significant differences between groups in the Vis and DA networks. The similarity between adults and neonates in connectivity to the SM and Vis networks may be due to the relative maturity of these areas at birth (Arcaro & Livingstone, 2017; Deen et al., 2017; Hurk et al., 2017; Gao et al. 2017, Dall’Orso et al., 2018).

To more specifically determine which regions in the networks were responsible for the differences seen between adults and neonates, we conducted a voxel-wise cortical analysis. Our results indicate that neonates have higher connectivity to much of the cortex as compared to adults with the exception of areas of bilateral medial orbitofrontal, isthmus cingulate and precuneus. This is consistent with Riggins et al.’s (2016) conclusion that 4-year old children rely more on regions “outside” the canonical hippocampal network to complete episodic memory tasks, and other research suggesting the infant cortex is more broadly tuned than in adults (Ellis & Turk-Browne, 2018). The few regions where adults display higher FC than neonates reside mainly within DM network and highlight the immaturity of this network: adults show significantly greater DM-Hippocampal connectivity than neonates, consistent with Gao et al.’s (2015) finding that this network is one of the last to develop in the first year of life.

Our anterior-posterior analysis and long-axis gradient analyses again suggest that the FC differentiation of the hippocampus is lacking at birth. Consistent with previous literature, adults display changes along the long-axis such that the anterior hippocampus shows greater connectivity to the Lim network than the posterior hippocampus but greater posterior vs anterior FC to the attention (i.e. FP and VA) networks; in fact, the anterior hippocampus is especially anti-correlated with these networks, as is consistent with previous literature (Buckner et al., 2009; Wael et al., 2018). The greatest differentiation in FC to the networks in adults occurred

within the anterior two-thirds of the hippocampus. In contrast, neonates showed no specificity to any of the networks along the long-axis or the anterior-posterior analysis. Blankenship et al. (2017), Langnes et al. (2018), and Riggins et al. (2016) show evidence of specialization along the longitudinal axis in 4- and 6-year old children but no such evidence is seen in our results, suggesting that maturational changes within the hippocampus may occur before age 4 to produce the preferential connectivity seen in children and adults. Future studies of infants and toddlers can better elucidate when after birth this change in specialization of the long-axis occurs.

Several limitations warrant discussion. A major problem in imaging children is motion artifact. We used the motion-corrected data that were released by the dHCP, took steps in preprocessing to ensure that physiological artifacts were removed from the data in both neonates and adults (Power et al., 2014; Yan et al., 2013), and further motion-matched the neonatal and adult groups. Given that motion-related artifacts are a major confound in FC analyses (Power et al., 2012; Satterthwaite et al., 2013), our approach should minimize the risk of spurious correlations. Other steps we took to minimize potential confounds included visual inspection of spatial registration results (and using established registration procedures that have been previously performed on infants (Alexander et al., 2019; Dean et al., 2018; Gao et al., 2009; Gao et al., 2015a)), performing the analyses in the native-space of each individual, and checking the reliability of the correlation values across participants in each group to ensure they were not particularly noisy in the neonatal group. A result of particular note is that neonates showed primarily positive FC from the hippocampus to the networks, while adults showed slightly negative FC for some networks. Blankenship et al., (2017) similarly fail to show any negative hippocampal FC in their sample of 4- and 6-year old children (but this may be due to their preprocessing steps, see Murphy & Fox, 2017 for discussion). Here, we used the same

preprocessing steps in both neonates and adults and used aCompCor and other preprocessing steps that should not necessarily remove negative correlations if they were there. Indeed, we found a normal distribution of correlation values in both neonates and adults (Figure 1) suggesting that negative correlations do exist in neonates, but not between the hippocampus and the cortex. Further, regardless of the negative vs. positive correlation differences we observe a difference in the pattern of FC in adults (demonstrated in the rose plots) primarily in the anterior portion of the hippocampus; this is missing in neonates.

Differences in arousal states between the groups present another challenge. Mitra et al., (2017) showed differences in resting-state connectivity between sleeping infants and waking adults. However, observation of Mitra et al.'s data suggests although the magnitude of connectivity may differ between arousal states, the overall pattern of connectivity remains similar (i.e. the same clusters of connectivity are observed in sleep and in rest and their relative comparison to other clusters remains similar across sleep states and age groups). Further, although notable differences are seen between the 24-mo sleeping infants and waking adults in Mitra et al., this difference is far less pronounced in the younger 6-mo infants. Based on previous EEG studies (e.g. Roffwarg, 1966), it is possible that younger infants experience less slow-wave sleep and more REM sleep and thus, younger infants (vs. older infants) during sleep would be expected to look more like awake adults due to the high similarity of REM and wakefulness activity patterns in EEG, particularly in infants. Because we would expect more awake-like REM sleep and less slow-wave sleep in young infants, we believe that the neonates in the current study are unlikely to show major wake/sleep confounds in their connectivity patterns. Finally, analysis of the same dataset but specifically of visual network connectivity showed striking *similarities* in connectivity patterns (<https://www.biorxiv.org/content/10.1101/712455v1>) and therefore further

suggest that any differences in data acquisition and/or sleep states between adults and neonates are unlikely to systematically lead to the differences in network connectivity that we find here.

Finally, we found the motion- and gender-matched HCP adults used in this manuscript tended to have lower tSNR than their respective dHCP counterparts. To ascertain this discrepancy was not the cause of observed group differences, we identified a separate group of 40 HCP adults whose tSNR matched that of the 40 neonates used here and performed our whole hippocampus to network analysis on this group. The resulting pattern matches the pattern observed from the previous analyses (i.e. using the original 40 HCP adults), reiterating that identified differences in connectivity patterns between adults and neonates are likely not spurious byproducts of discrepant data quality (Extended Data, Figure 1-1).

In conclusion, our results suggest that the resting-state FC patterns of the human hippocampus are immature at birth. This immaturity may play a key role in infantile amnesia and the vast differences between adults and neonates shown here suggests a fundamentally different memory and learning system from that of adults may be present at this point in development.

References

Alberini, C. M., & Travaglia, A. (2017). Infantile Amnesia: A Critical Period of Learning to Learn and Remember. *Journal of Neuroscience*, 37(24), 5783–5795.

Alexander, B., Kelly, C. E., Adamson, C., Beare, R., Zannino, D., Chen, J., ... Thompson, D. K. (2019). Changes in neonatal regional brain volume associated with preterm birth and perinatal factors. *NeuroImage*, 185, 654–663.

Arcaro, M. J., & Livingstone, M. S. (2017). A hierarchical, retinotopic proto-organization of the primate visual system at birth. *ELife*, 6.

537 Avants, B. B., Tustison, N. J., Song, G., Cook, P. A., Klein, A., & Gee, J. C. (2011). A
538 Reproducible Evaluation of ANTs Similarity Metric Performance in Brain Image
539 Registration. *NeuroImage*, 54(3), 2033–2044.

540 Behzadi, Y., Restom, K., Liau, J. & Liu, T.T.J.N. A component based noise correction method
541 (CompCor) for BOLD and perfusion based fMRI. *Neuroimage*, 37, 90-101 (2007)

542 Biswal, B., Yetkin, F. Z., Haughton, V. M., & Hyde, J. S. (1995). Functional connectivity in the
543 motor cortex of resting human brain using echo-planar mri. *Magnetic Resonance in*
544 *Medicine*, 34(4), 537–541.

545 Blankenship, S. L., Redcay, E., Dougherty, L. R., & Riggins, T. (2017). Development of
546 hippocampal functional connectivity during childhood. *Human Brain Mapping*, 38(1),
547 182–201.

548 Buckner, R. L., Andrews - Hanna, J. R., & Schacter, D. L. (2008). The Brain’ s Default
549 Network. *Annals of the New York Academy of Sciences*, 1124(1), 1–38.

550 Buckner, R. L., Sepulcre, J., Talukdar, T., Krienen, F., Liu, H., Hedden, T., ... Johnson, K. A.
551 (2009). Cortical Hubs Revealed by Intrinsic Functional Connectivity: Mapping, Assessment
552 of Stability, and Relation to Alzheimer’s Disease. *The Journal of Neuroscience : The*
553 *Official Journal of the Society for Neuroscience*, 29(6), 1860–1873.

554 Can, D. D., Richards, T., & Kuhl, P. (2013). Early gray-matter and white-matter concentration in
555 infancy predict later language skills: A whole brain voxel-based morphometry study.
556 *Brain and Language*, 124(1), 34–44. <https://doi.org/10.1016/j.bandl.2012.10.007>

557 Cole, M. W., Bassett, D. S., Power, J. D., Braver, T. S., & Petersen, S. E. (2014). Intrinsic and
558 task-evoked network architectures of the human brain. *Neuron*, 83(1), 238–251.

559 Dall’Orso, S., Steinweg, J., Allievi, A. G., Edwards, A. D., Burdet, E., & Arichi, T. (2018).
560 Somatotopic Mapping of the Developing Sensorimotor Cortex in the Preterm Human Brain.
561 *Cerebral Cortex*, 28(7), 2507–2515.

562 Dean, D. C., Planalp, E. M., Wooten, W., Schmidt, C. K., Kecskemeti, S. R., Frye, C., ...
563 Davidson, R. J. (2018). Investigation of brain structure in the 1-month infant. *Brain*
564 *Structure and Function*, 223(4), 1953–1970.

565 Deen, B., Richardson, H., Dilks, D. D., Takahashi, A., Keil, B., Wald, L. L., ... Saxe, R. (2017).
566 Organization of high-level visual cortex in human infants. *Nature Communications*, 8,
567 13995.

568 DeMaster, D., Pathman, T., Lee, J. K., & Ghetti, S. (2014). Structural development of the
569 hippocampus and episodic memory: Developmental differences along the
570 anterior/posterior axis. *Cerebral Cortex (New York, N.Y.: 1991)*, 24(11), 3036–3045.

571 Ellis, C. T., & Turk-Browne, N. B. (2018). Infant fMRI: A Model System for Cognitive
572 Neuroscience. *Trends in Cognitive Sciences*, 22(5), 375–387.

573 Fisher, R. A. (1915). Frequency Distribution of the Values of the Correlation Coefficient in
574 Samples from an Indefinitely Large Population. *Biometrika*, 10(4), 507.

575 Fitzgibbon, S.P., et al. The developing Human Connectome Project (dHCP): minimal 406
576 functional pre-processing pipeline for neonates. in *Fifth Biennial Conference on Resting*
577 *State and Brain Connectivity* (2016)

578 Gao, W., Alcauter, S., Elton, A., Hernandez-Castillo, C. R., Smith, J. K., Ramirez, J., & Lin, W.
579 (2015). Functional Network Development During the First Year: Relative Sequence and
580 Socioeconomic Correlations. *Cerebral Cortex (New York, NY)*, 25(9), 2919–2928.

581 Gao, W., Alcauter, S., Smith, J. K., Gilmore, J., & Lin, W. (2015). Development of Human Brain
582 Cortical Network Architecture during Infancy. *Brain Structure & Function*, 220(2),
583 1173–1186.

584 Gao, W., Lin, W., Grewen, K., & Gilmore, J. H. (2017). Functional Connectivity of the Infant
585 Human Brain: Plastic and Modifiable. *The Neuroscientist*, 23(2), 169–184.

586 Gao, W., Zhu, H., Giovanello, K. S., Smith, J. K., Shen, D., Gilmore, J. H., & Lin, W. (2009).
587 Evidence on the emergence of the brain’s default network from 2-week-old to 2-year-old
588 healthy pediatric subjects. *Proceedings of the National Academy of Sciences*, 106(16),
589 6790–6795.

590 Ge, X., Shi, Y., Li, J., Zhang, Z., Lin, X., Zhan, J., Ge, H., Xu, J., Yu, Q., Leng, Y., Teng, G.,
591 Feng, L., Meng, H., Tang, Y., Zang, F., Toga, A. W., & Liu, S. (2015). Development of
592 the human fetal hippocampal formation during early second trimester. *NeuroImage*, 119,
593 33–43.

594 Ghetti, S., DeMaster, D. M., Yonelinas, A. P., & Bunge, S. A. (2010). Developmental
595 Differences in Medial Temporal Lobe Function during Memory Encoding. *Journal of*
596 *Neuroscience*, 30(28), 9548–9556.

597 Gilmore, J. H., Shi, F., Woolson, S. L., Knickmeyer, R. C., Short, S. J., Lin, W., ... Shen, D.
598 (2012). Longitudinal Development of Cortical and Subcortical Gray Matter from Birth to
599 2 Years. *Cerebral Cortex*, 22(11), 2478–2485.

600 Glasser, M.F., et al. The minimal preprocessing pipelines for the Human Connectome Project.
601 *Neuroimage*, 80, 105-124 (2013)

602 Gómez, R. L., & Edgin, J. O. (2016). The extended trajectory of hippocampal development:
603 Implications for early memory development and disorder. *Developmental Cognitive*
604 *Neuroscience*, 18, 57–69.

605 Gogtay, N., Nugent, T. F., Herman, D. H., Ordonez, A., Greenstein, D., Hayashi, K. M., ...
606 Thompson, P. M. (2006). Dynamic mapping of normal human hippocampal
607 development. *Hippocampus*, 16(8), 664–672.

608 Grayson, D. S., & Fair, D. A. (2017). Development of large-scale functional networks from birth
609 to adulthood: A guide to the neuroimaging literature. *NeuroImage*, 160, 15–31.

610 Holm, S. (1979). A simple sequentially rejective multiple test procedure. *Scandinavian Journal*
611 *of Statistics*, 6, 65–70

612 Hughes, E.J., et al. A dedicated neonatal brain imaging system. *Magn Reson Med.* 78, 794-804
613 (2017).

614 Hurk, J. van den, Baelen, M. V., & Beeck, H. P. O. de. (2017). Development of visual category
615 selectivity in ventral visual cortex does not require visual experience. *Proceedings of the*
616 *National Academy of Sciences*, 114(22), E4501–E4510.

617 Jabès, A., Lavenex, P. B., Amaral, D. G., & Lavenex, P. (2011). Postnatal development of the
618 hippocampal formation: A stereological study in macaque monkeys. *The Journal of*
619 *Comparative Neurology*, 519(6), 1051–1070.

620 Jenkinson, M., Beckmann, C. F., Behrens, T. E. J., Woolrich, M. W., & Smith, S. M. (2012).
621 FSL. *NeuroImage*, 62(2), 782-790.

622 Keresztes, A., Ngo, C. T., Lindenberger, U., Werkle-Bergner, M., & Newcombe, N. S. (2018).
623 Hippocampal Maturation Drives Memory from Generalization to Specificity. *Trends in*
624 *Cognitive Sciences*, 22(8), 676–686.

Langnes, E., Vidal-Piñeiro, D., Sneve, M. H., Amlien, I. K., Walhovd, K. B., & Fjell, A. M. (2018). Development and Decline of the Hippocampal Long-Axis Specialization and Differentiation During Encoding and Retrieval of Episodic Memories. *Cerebral Cortex*, 29(8), 3398–3414.

Larson-Prior, L. J., Zempel, J. M., Nolan, T. S., Prior, F. W., Snyder, A. Z., & Raichle, M. E. (2009). Cortical network functional connectivity in the descent to sleep. *Proceedings of the National Academy of Sciences*, 106(11), 4489–4494.

Lavenex, P., & Banta Lavenex, P. (2013). Building hippocampal circuits to learn and remember: Insights into the development of human memory. *Behavioural Brain Research*, 254, 8–21.

Lin, W., Zhu, Q., Gao, W., Chen, Y., Toh, C.-H., Styner, M., ... Gilmore, J. H. (2008). Functional Connectivity MR Imaging Reveals Cortical Functional Connectivity in the Developing Brain. *American Journal of Neuroradiology*, 29(10), 1883–1889.

Linke, A. C., Wild, C., Zubiaurre-Elorza, L., Herzmann, C., Duffy, H., Han, V. K., Lee, D. S. C., & Cusack, R. (2018). Disruption to functional networks in neonates with perinatal brain injury predicts motor skills at 8 months. *NeuroImage. Clinical*, 18, 399–406.

Liu, W.-C., Flax, J. F., Guise, K. G., Sukul, V., & Benasich, A. A. (2008). Functional connectivity of the sensorimotor area in naturally sleeping infants. *Brain Research*, 1223, 42–49.

Liu, X., Yanagawa, T., Leopold, D. A., Fujii, N., & Duyn, J. H. (2015). Robust Long-Range Coordination of Spontaneous Neural Activity in Waking, Sleep and Anesthesia. *Cerebral Cortex*, 25(9), 2929–2938.

Makropoulos, A., et al. Automatic whole brain MRI segmentation of the developing neonatal brain. *IEEE Transactions on Medical Imaging*, 33, 1818-1831 (2014).

648 Makropoulos, A., et al. The developing human connectome project: A minimal processing
649 pipeline for neonatal cortical surface reconstruction. *Neuroimage*, 173, 88-112 (2018).

650 Murphy, K., & Fox, M. D. (2017). Towards a consensus regarding global signal regression for
651 resting state functional connectivity MRI. *NeuroImage*, 154, 169–173.

652 Mitra, A., Snyder, A. Z., Tagliazucchi, E., Laufs, H., Elison, J., Emerson, R. W., Shen, M. D.,
653 Wolff, J. J., Botteron, K. N., Dager, S., Estes, A. M., Evans, A., Gerig, G., Hazlett, H. C.,
654 Paterson, S. J., Schultz, R. T., Styner, M. A., Zwaigenbaum, L., IBIS Network, ...
655 Raichle, M. (2017). Resting-state fMRI in sleeping infants more closely resembles adult
656 sleep than adult wakefulness. *PloS One*, 12(11),

657 Natu, V. S., Gomez, J., Barnett, M., Jeska, B., Kirilina, E., Jaeger, C., Zhen, Z., Cox, S., Weiner,
658 K. S., Weiskopf, N., & Grill-Spector, K. (2019). Apparent thinning of human visual
659 cortex during childhood is associated with myelination. *Proceedings of the National*
660 *Academy of Sciences of the United States of America*, 116(41), 20750–20759.

661 Ofen, N., Kao, Y.-C., Sokol-Hessner, P., Kim, H., Whitfield-Gabrieli, S., & Gabrieli, J. D. E.
662 (2007). Development of the declarative memory system in the human brain. *Nature*
663 *Neuroscience*, 10, 1198.

664 Osher, D. E., Brissenden, J. A., & Somers, D. C. (2019). Predicting an individual's dorsal
665 attention network activity from functional connectivity fingerprints. *Journal of*
666 *Neurophysiology*, 122(1), 232–240.

667 Poppenk, J., Evensmoen, H. R., Moscovitch, M., & Nadel, L. (2013). Long-axis specialization of
668 the human hippocampus. *Trends in Cognitive Sciences*, 17(5), 230–240.

669 Poppenk, J., & Moscovitch, M. (2011). A Hippocampal Marker of Recollection Memory Ability
670 among Healthy Young Adults: Contributions of Posterior and Anterior Segments.
671 *Neuron*, 72(6), 931–937.

672 Power, J. D., Barnes, K. A., Snyder, A. Z., Schlaggar, B. L., & Petersen, S. E. (2012). Spurious
673 but systematic correlations in functional connectivity MRI networks arise from subject
674 motion. *NeuroImage*, 59(3), 2142–2154.

675 Power, J. D., Mitra, A., Laumann, T. O., Snyder, A. Z., Schlaggar, B. L., & Petersen, S. E.
676 (2014). Methods to detect, characterize, and remove motion artifact in resting state fMRI.
677 *NeuroImage*, 84, 320–341.

678 Prabhakar, J., Johnson, E. G., Nordahl, C. W., & Ghetti, S. (2018). Memory-related hippocampal
679 activation in the sleeping toddler. *Proceedings of the National Academy of Sciences of the*
680 *United States of America*, 115(25), 6500–6505. <https://doi.org/10.1073/pnas.1805572115>

681 Raichle, M. E. (2009). A Paradigm Shift in Functional Brain Imaging. *Journal of Neuroscience*,
682 29(41), 12729–12734.

683 Riggins, T., Geng, F., Blankenship, S. L., & Redcay, E. (2016). Hippocampal functional
684 connectivity and episodic memory in early childhood. *Developmental Cognitive*
685 *Neuroscience*, 19, 58–69

686 Roffwarg, H. P., Muzio, J. N., & Dement, W. C. (1966). Ontogenetic development of the human
687 sleep-dream cycle. *Science (New York, N.Y.)*, 152(3722), 604–619.

688 Salimi-Khorshidi, G., et al. Automatic denoising of functional MRI data: combining 409
689 independent component analysis and hierarchical fusion of classifiers. *Neuroimage*, 90,
690 449-468 (2014)

691 Salzwedel, A. P., Stephens, R. L., Goldman, B. D., Lin, W., Gilmore, J. H., & Gao, W. (2019).
692 Development of Amygdala Functional Connectivity During Infancy and Its Relationship
693 With 4-Year Behavioral Outcomes. *Biological Psychiatry: Cognitive Neuroscience and*
694 *Neuroimaging*, 4(1), 62–71.

695 Satterthwaite, T. D., Elliott, M. A., Gerraty, R. T., Ruparel, K., Loughead, J., Calkins, M. E., ...
696 Wolf, D. H. (2013). An improved framework for confound regression and filtering for
697 control of motion artifact in the preprocessing of resting-state functional connectivity data.
698 *NeuroImage*, 64, 240–256.

699 Seress, L. (2007). Comparative anatomy of the hippocampal dentate gyrus in adult and
700 developing rodents, non-human primates and humans. In H. E. Scharfman (Ed.),
701 *Progress in Brain Research* (pp. 23–798).

702 Smith, S. M., Fox, P. T., Miller, K. L., Glahn, D. C., Fox, P. M., Mackay, C. E., ... Beckmann,
703 C. F. (2009). Correspondence of the brain's functional architecture during activation and
704 rest. *Proceedings of the National Academy of Sciences of the United States of America*,
705 106(31), 13040–13045.

706 Smith, S. M., & Nichols, T. E. (2009). Threshold-free cluster enhancement: Addressing
707 problems of smoothing, threshold dependence and localisation in cluster inference.
708 *NeuroImage*, 44(1), 83–98.

709 Smith, S. M., Vidaurre, D., Beckmann, C. F., Glasser, M. F., Jenkinson, M., Miller, K. L., ...
710 Van Essen, D. C. (2013). Functional connectomics from resting-state fMRI. *Trends in*
711 *Cognitive Sciences*, 17(12), 666–682.

712 Snedecor, GW, Cochran, WG - Ames: Iowa State Univ. Press Iowa, 1989

713 Sporns, O. (2013). Structure and function of complex brain networks. *Dialogues in Clinical*
714 *Neuroscience*, 15(3), 247–262.

715 Strange, B. A., Witter, M. P., Lein, E. S., & Moser, E. I. (2014). Functional organization of the
716 hippocampal longitudinal axis. *Nature Reviews Neuroscience*, 15(10), 655–669.

717 Thompson, D. K., Wood, S. J., Doyle, L. W., Warfield, S. K., Lodygensky, G. A., Anderson, P.
718 J., Egan, G. F., & Inder, T. E. (2008). Neonate hippocampal volumes: Prematurity,
719 perinatal predictors, and 2-year outcome. *Annals of Neurology*, 63(5), 642–651.

720 Tobyne, S. M., Somers, D. C., Brissenden, J. A., Michalka, S. W., Noyce, A. L., & Osher, D. E.
721 (2018). Prediction of individualized task activation in sensory modality-selective frontal
722 cortex with ‘connectome fingerprinting.’ *NeuroImage*, 183, 173–185.

723 Travaglia, A., Steinmetz, A. B., Miranda, J. M., & Alberini, C. M. (2018). Mechanisms of
724 critical period in the hippocampus underlie object location learning and memory in infant
725 rats. *Learning & Memory*, 25(4), 176–182.

726 Wael, R. V. de, Larivière, S., Caldairou, B., Hong, S.-J., Margulies, D. S., Jefferies, E., ...
727 Bernhardt, B. C. (2018). Anatomical and microstructural determinants of hippocampal
728 subfield functional connectome embedding. *Proceedings of the National Academy of*
729 *Sciences*, 115(40), 10154–10159.

730 Wheelock, M. D., Hect, J. L., Hernandez-Andrade, E., Hassan, S. S., Romero, R., Eggebrecht, A.
731 T., & Thomason, M. E. (2019). Sex differences in functional connectivity during fetal brain
732 development. *Developmental Cognitive Neuroscience*, 36, 100632.

733 Yan, C.-G., Cheung, B., Kelly, C., Colcombe, S., Craddock, R. C., Di Martino, A., ... Milham,
734 M. P. (2013). A comprehensive assessment of regional variation in the impact of head
735 micromovements on functional connectomics. *NeuroImage*, 76, 183–201.

Yeo, B. T., Krienen, F. M., Sepulcre, J., Sabuncu, M. R., Lashkari, D., Hollinshead, M., ...
Buckner, R. L. (2011). The organization of the human cerebral cortex estimated by
intrinsic functional connectivity. *Journal of Neurophysiology*, 106(3), 1125–1165.

Legends

Figure 1: Preliminary Data Checks a) Gray matter (red), white matter (yellow) and network
registrations on the anatomical images of a representative adult and neonate subject; registration
image: blue=hippocampus, white = Vis, red = SM, purple (dark) = DA, pink = VA, green = Lim,
yellow = FP, orange = DM b) voxelwise correlations distributions of a representative adult and
neonate c) between-subject and between-group correlations demonstrate high within-group
reliability of connectivity but low between-group reliability between adults and neonates. (*)
indicates significance at $p < 0.05$; ns denotes non-significance

Figure 1-1: tSNR-Matched Adult Hippocampus to Networks. Hippocampal-network
connectivity of 40 tSNR-matched HCP adults again shows very similar results to the motion-
matched and binarized-hippocampal analyses. Hippocampal connectivity in adults shows a clear
hierarchy, with strong positive connectivity to Lim and DMN and negative connectivity to FP
and VA (*) indicates significance at $p_{HB} < 0.05$; (***) indicates significance at $p_{HB} < 0.005$.

Figure 2: Hippocampal Connectivity to Cortical Networks a) Comparison of hippocampal
connectivity to the seven cortical networks in adults showed a hierarchy of hippocampal
connectivity, whereby the highest FC was with Lim, followed by DM, SM, and Vis, almost no
FC with DA, and negative FC with FP and VA. b) In contrast, neonates show the same level of
FC to almost all of the 7 networks. Rose plot to the right shows adult connectivity compared to
neonates to highlight the differences between groups in the pattern of hippocampal FC to these

networks. Brain images on the top right depict connectivity between the hippocampus (left) and the seven cortical networks (right). (*) indicates significance at $p_{HB} < 0.05$; (***) indicates significance at $p_{HB} < 0.005$. Lim=Limbic; DM=Default Mode; SM=Somatomotor; Vis=visual; DA=Dorsal Attention; FP=FrontoParietal; VA=Ventral Attention

Figure 2-1: Binarized Whole Hippocampus to Networks. Hippocampal connectivity to the networks using a binarized HCP/dHCP hippocampal ROI yields very similar results to the ANTs registered hippocampal ROI (see figure 2). As with the initial analysis, hippocampal connectivity in adults shows a clear hierarchy whereas neonates display very few differences in hippocampal connectivity strength to the networks. (*) indicates significance at $p_{HB} < 0.05$; (***) indicates significance at $p_{HB} < 0.005$.

Figure 3: Hippocampal Connectivity to Cortex. Comparison of adult and neonate hippocampal connectivity to the cortex at a voxelwise grain. FWE-corrected results for the contrast of neonate > adult connectivity is shown in warm colors and the contrast of adult > neonate is denoted by cool colors.

Figure 3-1: Hippocampal Connectivity to Cortex, Adult>Neo Clusters are listed from largest to smallest. Peak coordinates (MAX) are listed in MNI space as well as center of gravity (COG) for each cluster

Figure 3-2: Hippocampal Connectivity to Cortex, Neo>Adult Clusters are listed from largest to smallest. Peak coordinates (MAX) are listed in MNI space as well as center of gravity (COG) for each cluster

Figure 4: Anterior/Posterior Hippocampal Connectivity to Networks a) Anterior vs posterior hippocampal-network connectivity in adults b) Anterior vs posterior hippocampal-network connectivity in neonates c) Rose plot comparing anterior vs posterior hippocampal-network connectivity pattern in adults d) Rose plot comparing anterior vs posterior hippocampal-network connectivity pattern in neonates. Brain image to the right shows the anterior (red) and posterior (yellow) hippocampal labels (*) indicates significance at $p_{HB} < 0.05$; (***) indicates significance at $p_{HB} < 0.005$.

Figure 5: Connectivity along the Long-Axis Gradient to Networks Comparison of the connectivity of the long axis gradient of the hippocampus to the 7 networks in a) adults and b) neonates. The slices are arranged anterior-to-posterior. Lighter coloring surrounding each line represents the standard error. Brain image on the right demonstrates the hippocampus (blue) segmented into slices (white lines). (*) are slices where the ANOVA shows an interaction between network and group at $p < 0.001$.

Figures

(Figure 1)

804 (Extended Data for Figure 1, Figure 1-1)

805

(Figure 2)

809

810 (Extended data for Figure 2; Figure 2-1)

811

812 (Figure 3)

<i>Cluster</i>	<i>Regions</i>	<i>Voxels</i>	<i>MAX</i>	<i>MAX</i>	<i>MAX</i>	<i>COG</i>	<i>COG</i>	<i>COG</i>	
				<i>(X)</i>	<i>(Y)</i>	<i>(Z)</i>	<i>(X)</i>	<i>(Y)</i>	<i>(Z)</i>
1	(L) Posterior Cingulate; Isthmus Cingulate, Precuneus	2370	8.08	-10	-57	17	-6.23	-55.2	20.8
2	(R) Isthmus Cingulate; Precuneus	1255	8.27	15	-54	19	9.3	-56.1	19.1
3	(L) Inferior Parietal	574	6.85	-42	-77	43	-44.3	-74.8	39.2
4	(L) Middle Temporal Cortex	403	5.72	-62	-1	-20	-63.2	-7.54	-18.2
5	(L) Medial Orbital Frontal	303	7.45	-10	39	-11	-7.75	41.9	-11.7
6	(L) Middle Temporal Cortex; Superior Temporal Cortex	235	6.88	-52	-13	-14	-53.1	-11.6	-13.7

813 (Extended Data for Figure 3; Figure 3-1)

814

815

816

817

818

819

820

<i>Cluster</i>	<i>Regions</i>	<i>Voxels</i>	<i>MAX</i> <i>(X)</i>	<i>MAX</i> <i>(Y)</i>	<i>MAX</i> <i>(Z)</i>	<i>COG</i> <i>(X)</i>	<i>COG</i> ⁴⁴ <i>(Y)</i>	<i>COG</i> <i>(Z)</i>	
1	(R) Rostral Middle Frontal; Pars Triangularis; Pars Orbitalis; Lateral Orbitofrontal; Pars Opercularis; Insula; Caudal Middle Frontal; Precentral; Postcentral	16290	8.72	57	14	4	44.9	26.2	21.8
2	(R) Superior Frontal; Paracentral	4702	6.88	4	26	61	9.85	12.7	59.4
3	(L) Supramarginal	3278	8.54	-65	-42	34	-59.6	-38.9	30.7
4	(R) Supramarginal; Inferior Parietal	3226	7.93	62	-36	48	61.8	-35.9	37.1
5	(R) Lingual; Pericalcarine (L) Lingual; Pericalcarine	2794	6.22	-19	-66	2	1.81	-77.1	5.9
6	(L) Rostral Middle Frontal	796	6.85	-34	51	29	-36.3	46.6	28.9
7	(R) Lateral Orbitofrontal; Pars Orbitalis	458	5.59	46	22	-7	39.4	24.6	-7.38
8	(L) Superior Frontal	350	5.74	-17	7	66	-13.6	8.52	69.4
9	(R) Insula	238	5.25	42	3	-6	40.4	6.44	-3.65

821 (Extended Data for Figure 3; Figure 3-2)

(Figure 4)

825

826 (Figure 5)

827

828

829