Brain orchestration of pregnancy and maternal behavior in mice

David André Barrière^{1,2#}, Arsène Ella^{1,3}, Frédéric Szeremeta⁴, Hans Adriaensen¹, William Mème⁴, Elodie Chaillou¹, Martine Migaud¹, Sandra Mème⁴, Frédéric Lévy¹,
Matthieu Keller^{1,#}

¹ *Physiologie de la Reproduction et des Comportements, UMR INRAE/CNRS/Université de Tours/IFCE, Nouzilly, France.*

² *Université Paris-Saclay, CEA, CNRS, BAOBAB, NeuroSpin, 91191, Gif-Sur-Yvette, France.*

³ *MRC Cognition and Brain Sciences Unit, University of Cambridge, UK.*

⁴ *Complexes Métalliques et IRM, Centre de Biophysique Moléculaire, UPR44301 CNRS, Orléans, France.*

[#]Corresponding authors:

matthieu.keller@inrae.fr and david.a.barriere@gmail.com

17 **Abstract**

18 Reproduction induces changes within the brain to prepare for gestation and motherhood.
 19 However, the dynamic of these central changes and their relationships with the development
 20 of maternal behavior remain poorly understood. Here, we describe a longitudinal
 21 morphometric neuroimaging study in female mice between pre-gestation and weaning, using
 22 new magnetic resonance imaging (MRI) resources comprising a high-resolution brain
 23 template, its associated tissue priors (60- μ m isotropic resolution) and a corresponding mouse
 24 brain atlas (1320 regions of interest). Using these tools, we observed transient hypertrophies
 25 not only within key regions controlling gestation and maternal behavior (medial preoptic area,
 26 bed nucleus of the *stria terminalis*), but also in the amygdala, caudate nucleus and
 27 hippocampus. Additionally, unlike females exhibiting lower levels of maternal care, highly
 28 maternal females developed transient hypertrophies in somatosensory, entorhinal and
 29 retrosplenial cortices among other regions. Therefore, coordinated and transient brain
 30 modifications associated with maternal performance occurred during gestation and lactation.

31

32 **Key words:** Gestation, lactation, maternal brain, MRI, atlas, voxel-based morphometry.

33 **Abbreviations:**

- 34 AC-PC: anterior commissure-posterior commissure
- 35 AMBMC: Australian Mouse Brain Mapping Consortium
- 36 AOB: accessory olfactory bulb
- 37 BNST: bed nucleus of the *stria terminalis*
- 38 CNS: central nervous system
- 39 CSF: cerebrospinal fluid
- 40 DARTEL: diffeomorphic anatomical registration using exponentiated lie algebra
- 41 df: degree of freedom
- 42 FA: flip angle
- 43 FLASE: fast large-angle spin-echo
- 44 FoV: field of view
- 45 MRI: magnetic resonance imaging
- 46 MOB: main olfactory bulb
- 47 mPOA: medial preoptic area
- 48 GM: gray matter
- 49 GMC: gray matter concentration
- 50 PVN: paraventricular nucleus of the hypothalamus
- 51 RARE: rapid acquisition with relaxation enhancement
- 52 ROC: receiver operating characteristic
- 53 ROI: region of interest
- 54 TE: echo time
- 55 TR: repetition time
- 56 VBM: voxel-based morphometry
- 57 WM: white matter

58 **Introduction**

59 Motherhood is among the most transformative experiences in the lives of female
60 mammals. While virgin females tend to avoid neonates, the end of the gestation period and
61 the birth process lead to a behavioral switch characterized by an attraction towards infant
62 cues, the expression of nurturing behavior and ultimately the establishment of infant
63 bonding^{1,2}. Decades of scientific research dedicated to the maternal brain have revealed a core
64 neural circuitry that includes the medial preoptic area (mPOA) and the adjoining ventral part
65 of the bed nucleus of the *stria terminalis* (BNSTv), and that is highly critical for the onset of
66 maternal behavior^{1,3-5}. Functional modulations of the mPOA/BNSTv consistently disrupt
67 maternal motivation and expression in numerous species^{1,2,6}. This core maternal circuitry
68 regulates maternal behavior through its direct projections to the ventral tegmental area, which
69 promotes reward system activation^{1,7}, as well as through its connections with cortical regions,
70 including the prefrontal cortex⁸⁻¹⁰. This crucial central circuitry is finely regulated by multiple
71 neural networks that integrate both internal and external stimulations. The proper expression
72 of maternal care towards offspring is prepared through the neuroendocrine action of sex
73 steroids and neuropeptides such as oxytocin among others during the gestation period¹¹. These
74 internal factors induce rewiring of the maternal brain, including through structural plasticity
75 through increasing neuronal soma size or astrocytic complexity within the mPOA¹², and
76 changes in neurogenesis mainly in the main olfactory bulb (MOB)¹³ but also in the
77 mPOA/BNST in rodents¹⁴. In human, regional morphological changes of gray matter (GM)
78 within the parahippocampal gyrus, precuneus, cingulate, insula and frontal cortex have been
79 observed in primiparous women using magnetic resonance imaging (MRI)¹⁵. Additionally,
80 olfactory cues coming from the neonate are integrated by the MOB and the accessory
81 olfactory bulb (AOB) through an amygdalo-hypothalamic pathway, which is responsible for
82 attraction/repulsion behavioral outcomes¹⁶. Hence, the development of the maternal brain is

83 dependent on both integration of external and internal cues acting through multiple brain
84 pathways and regions to prepare the brain to gestation and motherhood. Nevertheless, the
85 relationship between the brain rewiring over the gestation and lactation periods and the
86 establishment of the maternal behavior is poorly documented.

87 To assess the dynamics of the maternal brain, a longitudinal MRI morphometric study
88 over a complete reproductive experience was performed in mouse to investigate changes in
89 the gray matter concentration (GMC) using voxel-based morphometry (VBM). VBM is a
90 well-established and well-validated image analysis technique that provides an unbiased and
91 comprehensive assessment of anatomical differences throughout the brain, and has been
92 successfully used to study GM changes within the mouse brain^{17–21}. However, available
93 mouse brain MRI resources are often partial or provided in different spatial orientations or
94 spatial resolutions (**Table 1**). As an example, the Australian Mouse Brain Mapping
95 Consortium (AMBMC), offers a high resolute template and detailed atlases of the mouse
96 brain including the cerebellum²², hippocampus²³, diencephalon²⁴ and cortices²⁵.
97 Unfortunately, segmentation of the MOB and the AOB and hindbrain is lacking, and this
98 resource does not provide associated tissues probabilistic maps necessary for VBM. In other
99 hand, the Allen Mouse Brain Common Coordinate Framework, is the most advanced mouse
100 brain atlas²⁶. This new atlas delimitates discrete structures within the thalamus, hindbrain,
101 olfactory system and diencephalon and provides a full segmentation of cortical layers
102 however, MRI template and brain tissues priors are still lacking for VBM investigations.

103 Thus, we combined these resources and associated probabilistic maps to emulate a
104 complete resource dedicated to the mouse brain. Using these resources and a longitudinal
105 VBM approach, we were able to assess the dynamic morphological changes of the brain
106 during the whole reproductive period and demonstrate how these changes predict the quality
107 of maternal behavior.

Results

Mouse MRI atlas

VBM strategies require a template image and its associated priors of gray matter (GM), white matter (WM) and cerebrospinal fluid (CSF) for brain segmentation and normalization. In addition, a complete atlas of the mouse brain is mandatory for the identification of regions of interest (ROIs) highlighted by the VBM analysis. Given the limitations of available tools to thoroughly study GMC changes during the gestation and lactation periods in mice, we developed first a new set of resources using the AMBMC, an ultra-high-resolution template built from *ex vivo* brain images finely normalized within the same space²⁵ and the Allen Mouse Brain Common Coordinate Framework²⁶ (**Figure 1**). Our resources comprise the following four components: 1) a complete mouse brain template with a spatial resolution suitable for mouse brain analysis (60- μ m isotropic resolution); 2) the corresponding GM, WM and CSF probabilistic maps for brain normalization together with a VBM analysis built from 138 T₂-weighted images; 3) a complete mouse brain atlas derived from Paxinos and Franklin's mouse brain atlas²⁷ and composed of a mosaic of 1320 ROIs (**Figure 2A**); 4) a brain mesh permitting brain plot generation and data visualization (**Figure 2B** and **Supplemental Video 1**). We visually inspected and carefully checked the results of the normalization process against the original coregistered atlas. Then, the labeled structures were reclassified and aggregated according to the brain regions to which they belonged (auditory, insular, temporal cortices, etc.), with respect to their anatomical topography (cortex, basal ganglia, etc.), tissue type (GM, WM and CSF) and hemisphere (left or right). Cortical structures were subdivided into functional (*e.g.*, primary and secondary motor cortices) or structural (agranular, dysgranular, agranular/dysgranular, granular and posterior agranular insular cortices) areas (**Figure 2C**), as well as into different cortical layers (**Figure 2D**).

Subcortical structures, as for example, the hypothalamus (**Figure 2E**) and the hippocampus (**Figure 2F**), were fully segmented according to Paxinos and Franklin's atlas²⁷.

Morphometric changes occurred during the gestation and lactation periods

Next, we used our new resources to study the variations of GMC in mouse brain from the beginning of gestation until weaning. Using MRI T₂-weighted anatomical acquisitions, we estimated the GMC maps which offer for each animal a global estimation of the GM. Longitudinal comparison of GMC between virgin mice (control group, n= 11) and mice who became pregnant and raised their young until weaning (parous group, n=12) permits to highlight local modifications of GM during the whole reproductive cycle. A comparison of baseline and early gestation GMC maps between the control and parous groups did not reveal significant differences. However, at the end of the gestation period, significant increases in GMC were observed within several brain regions in the parous group compared to the control group (**Table S1** and **Figure 3A**). Time course analysis revealed differences in GMC profiles between control and parous groups at the end of the gestation period, early in lactation and at the end of the lactation period. Specifically, GMCs within the mPOA and the BNST were consistently significantly higher in the parous group at that times. In addition, within the agranular insular cortex in the late gestation period and the early lactation period, GMC was significantly higher in parous group compare to control group (**Figure 4A**). During the early lactation period, we also found specific and significant increases in GMCs of the parous group within numerous brain regions (**Table S2** and **Figure 3B**). Among these structures, the hippocampus (CA1 layer), amygdalar area and piriform area showed a transient increase in GMC at the early lactation time point that returned to baseline values at the end of the lactation period in parous group compared to control group (**Figure 4B**). In contrast, the caudate putamen, arcuate nucleus and paraventricular nucleus of the hypothalamus (PVN)

showed significantly higher GMCs in the parous mice than in the control mice during the lactation period (**Table S3** and **Figures 3C and 4C**). Together, our data demonstrate that the late gestation period is associated with a pronounced increase in GMC in the mPOA/BNST, the core neural system of maternal motivation, lasting up to the late lactation period. Furthermore, the early lactation period is associated with increased GMC in other key maternal motivation areas in midbrain regions, including the hypothalamus, caudate putamen and amygdala. These GMC differences between both groups were no longer observed after weaning.

Morphometric changes during gestation predict the quality of maternal behavior

In the last part of this work, we evaluated whether these morphometric changes might reflect differences in maternal performance. Based on 15 min of behavioral observation during the pup retrieval test performed one week after birth, we evaluated the maternal performance of each mother by measuring the first, second and third pup retrieval times, pup-licking duration, crouching time, rearing time, digging time and self-grooming time (**Figure 5A**). Interestingly, we observed a large distribution of values for both crouching (284.2 s, SD $\pm$ 268.7 s) and digging (98.22 s, SD $\pm$ 159.3 s) durations within the parous group. Whereas crouching duration relates to maternal behavior, digging duration is widely recognized as a discriminative marker of stress-related behavior in rodents²⁸. Hence, we used crouching and digging durations to cluster animals using the k-means clustering procedure, thereby clustering parous animals into a high maternal behavior group (those with a high crouching time and low digging time) and a low maternal behavior group (those with low crouching time and high digging time).

A comparison of maternal performance parameters between the two clustered groups revealed significantly lower crouching and pup-licking times and a significantly higher

digging time (**Figure 5B**) in the low maternal behavior group (n=6) than in the high maternal behavior group (n=6). Moreover, a comparison of GMC maps revealed both cortical and subcortical differences between the two clustered groups, mainly in the late gestation period (**Table S4** and **Figure 5C**) but also during the early lactation period (**Table S5** and **Figure 5D**). Indeed, transient increases in GMCs within the entorhinal area, lateral part of the orbital area, AOB, and medial preoptic area were observed at the end of pregnancy in the high maternal behavior group were absent in the low maternal behavior group (**Figure 6A**). In addition, the high maternal behavior group showed consistent higher GMCs in the hippocampus, retrosplenial area and barrel field of the primary somatosensory cortex (**Figure 6B**) from the end of the gestation until the end of the lactation period.

Interestingly, using a receiver operating characteristic (ROC) analysis, we found that GMCs values within the entorhinal area and AOB at late gestation are reliable predictors for mouse maternal performance after birth. These GMC values significantly distinguished low maternal performance from high maternal performance postpartum (entorhinal area: sensitivity = 100, confidence interval (CI) = 61% to 100%; specificity = 83, CI = 44% to 99%; likelihood ratio = 6; **Figure 7A**; AOB: sensitivity = 100, CI = 61% to 100%; specificity = 83, CI = 44% to 99%; likelihood ratio = 6, **Figure 7B**). The GMC values of both the entorhinal area and AOB observed at late gestation were also significantly correlated with maternal behavior (crouching and digging times). These results reveal that the GMC differences in olfactory (AOB and entorhinal cortex) and mnemonic (entorhinal area)-related brain regions occurring during the late gestation period significantly predicted the quality of maternal behavior.

Discussion

Using a new comprehensive neuroimaging resource dedicated to mouse brain, this longitudinal study reveals that pregnancy and lactation coincide with pronounced and transient cerebral changes. Transient increases in GMCs were observed in key regions controlling maternal behavior (mPOA, BNST, and PVN), as well as regions involved in emotions (amygdala), in motivation and reward (caudate nucleus, orbitofrontal cortex) and in mnemonic functions (hippocampus). Interestingly, increase in GMC was also revealed in the insular cortex thought to link social and emotional skills. Moreover, we showed that females expressing high levels of maternal behavior had developed specific increases in GMCs in structures involved in olfactory (MOB and AOB) and somatosensory (somatosensory cortex) information processing, in memory (hippocampus, entorhinal cortex, retrosplenial cortex) and in reward and reinforcement (striatum). Interestingly, these hypertrophies were already significant at the end of the gestation period thus being predictive of the quality of maternal care (**Supplemental Video 2**).

Implementation of new resources to support the analysis of mouse brain MRI data

The use of preclinical MRI is a target of growing interest for the study of brain structure and function in both healthy and pathological conditions. The use of advanced MRI techniques, coupled with the development of advanced animal models, is a powerful way to push new breakthroughs in the understanding of brain functioning and pathology. Herein, in the first step in our study, from the recent major advances in the development of brain mouse atlas we generated a new set of neuroinformatic tools offering for the first time a complete resource dedicated to MRI studies of the mouse brain, namely, an accurate brain atlas (1320 ROIs), a high-resolution brain template and the associated GM, WM and CSF priors (60- μ m isotropic resolution). The GM, WM and CSF probabilistic maps built and used in this study

were calculated from 138 T₂-weighted anatomical images, resulting in robust tissue class priors not only for VBM analysis but also for functional MRI and diffusion tensor imaging analysis in mice.

This comprehensive set of MRI compatible template and atlas for the mouse brain, allows a unified and standardized analysis of multimodal mouse brain MRI data and paves the way for the development of multicentric preclinical studies. Indeed, animal models deliver crucial information for the understanding of brain structure and function both in healthy and pathological conditions. Our template and mouse brain atlas were conceived to bridge the gap between basic and clinical neurosciences by providing to the preclinical neuroimaging community specific resources designed to be used in conjunction with the neuroinformatic tools and methodologies commonly used in human MRI studies. We anticipate that these resources will help neuroscientists to conduct their analyses of anatomical and functional datasets in a more standardized way, with the final goal of reaching more reproducible conclusions (https://www.nitrc.org/projects/tmbta_2019).

Gestation and lactation periods induce strong but transient GMC hypertrophy

The establishment of accurate mouse MRI resources permit to study the variations of GMC longitudinally, *in vivo* and during the gestation and lactation periods in female mice. We observed that several brain regions became transiently hypertrophic during pregnancy or in the lactation period until weaning. A set of structures comprising the core of the maternal circuit – especially the mPOA and BNST – displayed long-lasting hypertrophy that started at late gestation, culminated during the first week of lactation, and then disappeared at weaning. The initial GMC increase observed during the gestation period probably reflects changes induced by hormonal priming¹¹. Indeed, both the mPOA and BNST express a high number of steroid hormone and neuropeptide receptors^{29,30}. These factors are well known to trigger

significant plasticity changes within the core maternal circuitry that are necessary for the preparation and adaptation of the brain to motherhood^{3,11}. From parturition, the GMC differences observed during the whole lactation period highlight that mPOA/BNST receives a variety of sensory inputs from the pups, integrates that information with the females' endocrine status, and then projects to brain sites involved in socially-relevant motivation, affective state, and cognition^{2,31}. Pup stimulation, electrolytic and neurotoxic lesions and local steroid hormone injections³²⁻³⁵ have been shown to modify the intrinsic activity of these nuclei and consequently responsible for motivation and expression of maternal behavior. The mPOA is engaged throughout the postpartum period but differentially according to the needs of the developing pups. It has been shown that neurons of the mPOA in late postpartum inhibits maternal responses allowing the changing expression and waning of maternal behavior across postpartum¹⁰. The sustained increase in GMC reported in late lactation could reflect the involvement of the mPOA to appropriately influence maternal behavior.

Our study highlights another set of brain structure that became hypertrophic only during the period from parturition to weaning – specifically, the PVN and arcuate nucleus of the hypothalamus. These changes illustrate the structural plasticity occurring in these regions. For instance, at parturition and during lactation oxytocin neurons of the PVN undergo dramatic neuronal, glial and synaptic changes such as an increase in size of the oxytocin neurons and an amplification of their synaptic input³⁶. Oxytocin release at parturition facilitates the onset of maternal behavior by acting on the mPOA and is also important for maternal memory³⁷. Finally, lesions of the PVN disrupt the onset of maternal behavior³⁸⁻⁴⁰. In the arcuate nucleus, dopaminergic cells are responsible for suckling induced prolactin release⁴¹ and neurons projecting to the arcuate nucleus are involved in the maintenance of maternal motivation⁴².

Additionally, we report changes in GMCs found in olfactory related structures (main olfactory bulb, piriform cortex), somatosensory areas and auditory areas which reflect the multisensory control of maternal behavior. This finding is in accordance with a functional MRI study performed in rats which demonstrates that pup suckling is associated with increased neuronal activity within the midbrain, striatum and cortical sensory areas (somatosensory, olfactory and auditory cortices)⁹.

GMC variations within the hippocampus and entorhinal cortex, highlight the role of two essential structures involved in learning and memory processing during the reproductive period. Our data support evidence that the hippocampus undergoes profound neural changes during lactation. Indeed, lactating females have elevated spine densities in the hippocampus⁴³ and show significant dendritic remodeling in pyramidal neurons⁴⁴. Changes in hippocampal neurogenesis occurs during lactation and may support the enhancement of spatial memory necessary to foraging behavior in lactating females^{43,45-47}.

Finally, our study also revealed a hypertrophy of the agranular insular cortex, which has never been reported in this context. The agranular insular cortex is a laminar part of the insular cortex and can be considered as a hub structure linking large-scale brain systems⁴⁸. Indeed, the insula receives direct thalamic and somatosensory afferents carrying sensitive stimulations. In addition to its sensory afferents, the insula displays structural connectivity with the limbic system (basolateral, lateral and central amygdalar nuclei) as well as with the BNST, mediodorsal nucleus of the thalamus, lateral hypothalamus and perirhinal and lateral entorhinal cortices⁴⁸. The insula also connects brain regions implicated in motivation and reward, such as the nucleus accumbens and caudate putamen⁴⁸. Hence, our findings and the current literature suggest that before and after birth, the insular cortex may integrate and combine information from both external and internal stimulation and act as a relay between higher cortical and subcortical structures. Together, our results describe the dynamics of

neurophysiological adaptation occurring in the brain from the early gestation period to weaning, thereby ensuring efficient maternal behavior and, by extension, the development of the offspring.

GMC modifications in the olfactory system at the end of the gestation predict the level of maternal behavior post-partum

All these transient modifications of GMC in parous animals indicate that various brain regions undergo considerable plasticity from birth to weaning. Then, we sought to determine whether inter-individual variations in maternal behavior were associated with similar variations of GMC. Based on their behavioral performance in the pup retrieval test, we used a k-means clustering strategy to divide maternal female mice into two groups displaying high *versus* low levels of maternal behavior. This analysis revealed that several transient and several long-lasting increases in GMCs were observed in the high maternal behavior group that were absent in the low maternal behavior group. Brain regions showing significant differences included the olfactory bulbs, somatosensory system, limbic system, especially the orbitofrontal area, and mnesic system, including the retrosplenial cortex, hippocampus and entorhinal area. Some of these structures are directly responsive to pup stimulation, and the observed dynamics may have been induced by mother-offspring interactions. For example, higher GMC values in the somatosensory cortex and olfactory bulbs in the high maternal behavior group potentially reflected increased suckling duration and proximity between the mother and pups, respectively.

Strikingly, differences in GMCs were detected in the entorhinal area, orbitofrontal area, olfactory bulb, hippocampus, retrosplenial area and primary somatosensory area before parturition. These findings suggest that the maturation of these structures, probably through hormone-dependent plasticity mechanisms, is a key determinant of the intensity of maternal behavior expressed during the lactation period. Using a ROC procedure, we found that GMC

values of the entorhinal area and AOB at the end of gestation were significantly predictive of the maternal behavior postpartum. Interestingly, previous studies in mice reported an increase in cell proliferation during gestation in the subventricular zone, the neurogenic niche which provides newly generated neurons within the olfactory bulb (for review see ¹³). These adult-born olfactory neurons are fully responsive to pup odor exposure⁴⁹ and are in part involved in some components of maternal behavior^{50,51}.

Hence, the correlations observed between GMC values in AOB at the late gestation period and maternal behavior performances may suggest that impairments of neural plasticity of this olfactory region would induce maladaptive neuroendocrine processing of the maternal brain at the end of the gestation period impacting maternal behavior performance. Taken together, our data provide the first potential imaging-based predictive biomarkers of the quality of maternal behavior and suggest the key role of the maturation of the olfactory system at the end of the pregnancy in the development of adaptative maternal behavior in mice.

Conclusion

Our study provides a new generation of neuroinformatic tools which will help basic neuroscientists to conduct structural and functional MRI investigations. Using these resources, we found that the development of the maternal brain is associated with substantial mesoscopic changes in critical regions. These modifications can be interpreted as cell size changes, neural or glial cell genesis/apoptosis, spine density or blood flow modifications⁵²⁻⁵⁴. As cellular and molecular plasticity events are key for the adaptation of the brain to motherhood, molecular, cellular and behavioral investigations must be performed to obtain a more precise view of the physiological mechanisms responsible for GMC variations.

Methods

Animals. Twenty-three female RjOrl:SWISS virgin mice (8 weeks old; 20-25 g; Janvier Laboratory, Le Genest-Saint-Isle, France) were maintained on a 12-h light/dark cycle with access to food (standard chow) and water *ad libitum*. Animals were acclimatized 6 per cage to the housing facility for 7 days prior to manipulation. Females were randomly divided into two groups: a parous group (n = 12), in which each female was exposed to a male (RjOrl:SWISS, 8 weeks old; 20-25 g; Janvier Laboratory, Le Genest-Saint-Isle, France) for 5 days, became pregnant, and raised their offspring (litter size: 6 to 14 pups) until weaning at 21 days postpartum, and a control group (n = 11), in which virgin females were not exposed to male mice. Each parous female was individually housed after male exposure. Control females were housed together in a separate room from parous females.

The MRI protocol was optimized to keep mice anesthetized for 2 h during each of the six acquisitions. During lactation MRI acquisitions, pups were kept under a heat lamp. One week after birth, maternal behavior was assessed as described in the behavioral section. All experiments were conducted in accordance with the local research ethics committee (APAFIS #6626-201002281145814V1) and are reported in accordance with the ARRIVE guidelines.

MRI acquisition. *In vivo* 3D MRI of the entire brain was performed three days before male exposure (baseline), at one week of gestation (early gestation), two days before the expected day of birth (late gestation), one week postpartum (early lactation), three weeks postpartum (late lactation) and two weeks after weaning (weaning). One female in the parous group and one female in the control group were scanned under similar conditions on the same day.

Mice were anesthetized using isoflurane (2.5%; induction in O₂/air mixture 1:1) (TEM-SEGA, F-33600 Pessac, France) and then transferred and placed head first *prone* within an MRI-compatible cradle that incorporated a stereotaxic system dedicated for mouse

head MRI, connected to a heater with circulating water to maintain body temperature and supplied with 1-2% isoflurane *via* a fitted mask. Respiration rate was recorded during all the experiments using an MRI-compatible monitoring system (PC-SAM model #1025; SA Instruments Inc., Stony Brook, NY, USA) and used to adjust the isoflurane rate to maintain a rate between 20 and 40 respirations per minute. After a recovery period of one hour, mouse returned to her pups. MRI studies were conducted at the Centre de Biophysique Moléculaire d'Orléans and were performed on a 7T/160 mm PharmaScan spectrometer (Bruker Biospin, Wissembourg, France) equipped with an actively shielded B-GA09 gradient set, with 90-mm inner diameter and 300-mT/m gradient intensity. A 23-mm inner diameter Bruker birdcage coil with a cradle dedicated to a mouse head was used. Data acquisitions were performed on an Advance III console running ParaVision 5.1 software. T₂-weighted images were acquired using a 3D fast large-angle spin-echo (FLASE) sequence which allows 3D brain mapping with a high resolution in a suitable time for *in vivo* acquisition^{55,56}. Thus, the sequence with echo time (TE) = 20 ms, 1 repetition, acquisition matrix = 160 × 140 × 95, and a field of view (FoV) of 19.2 × 16.8 × 11.4 mm³, resulting in a final resolution of 120 μm isotropic voxels^{55,56}. To obtain the FLASE sequence⁵⁵, which is a specific sequence that is not included in the sequence package provided with ParaVision, the usual rapid acquisition with relaxation enhancement (RARE) spin-echo sequence was modified; in particular, the RARE-factor was fixed to 1 allowing a flip angle (FA) higher than 90° for the excitation pulse, while maintaining a 180° refocusing pulse⁵⁵. Thus, T₂-weighted images were obtained with a repetition time (TR) as short as 300 ms, 10 times lower than that needed for a classical T₂-weighted spin-echo sequence. The sequence was optimized for acquisition in 1 h 28 min, with an isotropic resolution of 120 μm, a TR of 300 ms, an TE of 20 ms, an excitation pulse (FA) of 115°, with 1 repetition and with a matrix of 160 x 140 x 95 corresponding to a FoV of 19.2 x 16.8 x 11.4 mm³ and contained the whole mouse brain.

Maternal behavioral test. One week after birth, the maternal behavior of each female was evaluated using the pup retrieval test⁵⁷. Briefly, three pups were removed from the nest and placed at three different corners within the home cage. The latency to retrieve each pup and the time spent licking the pups, crouching in the nest over the pups and performing nonmaternal behaviors such as self-grooming and digging were recorded over 15 min. Retrieval was defined as the animal picking up a pup and transporting it to the nest. Crouching was defined as the animal assuming the nursing posture. Nursing and licking were permitted whether they took place in the nest. All videos were analyzed using BORIS software version 4.1.4.

K-means clustering. Clustering analysis of the behavioral data was performed using MATLAB Simulink 10b (The Mathworks, Inc., USA). Normality was verified and no outlier subjects were detected. To classify animals according to their maternal performance, a k-means clustering algorithm was used with crouching and digging times as behavioral markers. Digging was chosen because it is indicative of high maternal stress^{28,58}. This algorithm iteratively grouped the animals by creating k initial centroids, assigning each animal to the closest centroid, iteratively re-calculating the centroids from the mean of its assigned animals and re-assigning the animals to each centroid until there were no more changes across iterations⁵⁹. This clustering divided parous animals into a high maternal behavior group (with high crouching and low digging time) and low maternal behavior group (with low crouching and high digging time).

Mouse brain template and atlas building. For the MRI protocol, we developed a brain template and an atlas from the AMBMC brain template and the Allen Mouse Brain Common Coordinate Framework, respectively (**Figure 1**). First, we down-sampled the AMBMC

template and its associated atlases and the Allen Mouse Brain Atlas and its associated Nissl images to a suitable resolution for MRI analysis (60- μ m isotropic resolution; **Figure 1**, step 1). Then, all images were manually aligned to the anterior commissure/posterior commissure (AC/PC) axis, and the center of the images was defined relative to the AC (**Figure 1**, step 2). The resulting template was then segmented into GM, WM and CSF probability maps using the unified segmentation approach⁶⁰ of Statistical Parametric Mapping 8 (**SPM8**) and the mouse brain priors provided by the **SPMMouse toolbox** (**Figure 1**, step 3). In parallel, our T₂-weighted anatomical images were realigned, coregistered, bias-corrected and normalized to our template. Using the **SPMMouse toolbox**, we also segmented the images as described above, and from these preprocessed images, we obtained a large set of 138 images for each tissue class (**Figure 1**, steps 4-7). From these images, we built population-specific GM, WM and CSF priors. To build these priors, for each tissue class, we applied a diffeomorphic anatomical registration using an exponentiated lie algebra (DARTEL) approach, which is an automated, unbiased, and nonlinear template-building algorithm⁶¹ (**Figure 1**, step 8). This new set of population-specific tissue priors was used for both atlas building and final VBM preprocessing.

To normalize the Allen Mouse Brain Atlas to our brain template, we used the associated Nissl-stained images because (1) Nissl staining corresponds to the GM prior in terms of histology, and (2) this image was already coregistered to the atlas (**Figure 1**, step 4). Therefore, we applied the segmentation function provided by **SPM8** using the GM prior previously calculated from the Nissl image to generate the “Nissl2template” normalization matrix. We used this matrix to normalize the atlas to the template, while avoiding interpolation to maintain the label indices as integers (**Figure 1**, step 7). Then, a visual inspection of each normalized label was carried out to assess whether the normalization process modified the position and volume of the structure too much. When necessary, holes

were filled and labels were redrawn according to Paxinos and Franklin's atlas and using the FreeSurfer package. Finally, the olfactory bulbs and hind brain regions were completed, the corpus collosum and ventricles were drawn from the WM and CSF priors, and the cerebellum labels were replaced by the AMBMC cerebellum labels, which are more accurate. Finally, the atlas image was symmetrized (left-right). Our mouse brain template, priors and atlas were normalized within the same space and with the same final resolution (60- μ m isotropic resolution), resulting in our final mouse brain atlas composed of a mosaic of 1320 ROIs covering the entire brain (**Supplemental Video 1**).

VBM data preprocessing. Previously preprocessed normalized T₂-weighted data were segmented into GM, WM, and CSF within SPM8 using the images of the population-specific priors (**Figure 1**, step 9). Then, to produce a more accurate registration within each mouse as well as across all mice, a longitudinal VBM analysis was applied using the strategy described by Asami *et al*⁶². First, a *subject-specific* template was created by the DARTEL algorithm using the previous tissue class images (*i.e.*, GM, WM, and CSF maps) obtained from each mouse for the six time points. The DARTEL procedure releases *individual-specific* flow field maps, permitting the application of diffeomorphic normalization on images of each tissue class to spatially normalize each time point on a *subject-specific* template space. Each normalized tissue class image was modulated by the Jacobian determinant to account for the expansion and/or contraction of brain regions over time. Then, a *population-specific* template was created by the DARTEL algorithm using all *subject-specific* templates of the tissue class images. Here, the DARTEL procedure releases *population-specific* flow field maps, permitting the application of diffeomorphic normalization of each animal onto the images of each tissue class. Finally, tissue class images were modulated by the Jacobian determinant, and the final modulated GM images were spatially smoothed with an isotropic Gaussian

kernel with a 3-mm full-width at half-maximum and convolved with GMC images to create GMC maps (**Figure 1**, steps 10-12).

VBM statistics and analysis. SPM8 was used to reveal the temporal and regional changes in the GMC maps. A second-level SPM analysis comprising a flexible factorial model, which is equivalent to a 2×2 mixed-model ANOVA with group as the between-subject factor and time point as the within-subject factor, was used to compare the control *versus* the parous groups and the low *versus* high maternal behavior groups⁶³. The factors included in the analysis were subjects, group (control *versus* parous, or low maternal behavior *versus* high maternal behavior), and time points (baseline, early gestation, late gestation, early lactation, late gestation, and weaning). A brain mask was used to constrain the analysis within the brain. For each cluster, the significance of the peak voxel was set as $p < 0.01$ ($t_{(126)} = 2.356$, control *versus* parous; $t_{(60)} = 2.39$, low *versus* high maternal behavior), and the minimum cluster extent was set at 25 voxels. The results are presented on axial brain slice series generated by the Xjview SPM plugin. Corresponding surfacing results were produced with BrainNet viewer 1.6⁶⁴, allowing the generation of both brain meshes and brain plots to visualize data and create supplemental videos.

Postprocessing statistical analysis. Cluster peaks revealed by the flexible factorial analyses were identified using our atlas and an *in situ* procedure developed with MATLAB Simulink 10b (The Mathworks, USA). For each comparison, clusters were binarized, and the obtained masks were used to extract GMC values of corresponding regions from the GMC map using the REX plugin. GMC data and behavioral measurements were then compiled and analyzed using GraphPad Prism 6.02 software. GMCs were compared between groups and for each time point using a two-way ANOVA with repeated measures followed by a two-stage setup method of Benjamini, Krieger and Yekutieli as recommended by the software. Maternal and

502 nonmaternal behaviors of the low maternal and high maternal groups were compared using a
 503 multicomparison t-test with a false discovery rate (FDR) approach ($Q = 1\%$). Correlation
 504 analyses were performed using a parametric two-tailed Pearson test. Specificity and
 505 selectivity analyses were performed using the ROC curve method. Statistical significance was
 506 defined as $p < 0.05$ (*) for these analyses.

Acknowledgments

The authors acknowledge the regional council of Centre Val-de-Loire for funding this research through the IMACERVOREPRO grant (convention 201500104011, 2015-2018) awarded to Matthieu Keller.

Author Contributions

D.A.B. contributed to the atlas and template building, data analysis, drafted and revised the manuscript. **A.E.** contributed to the imaging sequence troubleshooting, data analysis and contributed to the critical revisions. **F.S.** contributed to MRI sequence programming, imaging protocol setting, data acquisitions and contributed to the analysis and critical revisions. **H.A.**, **W.M.**, **E.C.**, **M.M.**, **S.M.** contributed to the study conception and design and contributed to the critical revisions of the manuscript. **F.L.** contributed to raise the funding, study conception and design and contributed to the critical revisions of the manuscript. **M.K.**, is the principal investigator of the study, raised the funding, coordinated the project, revised and validated the manuscript.

References

1. Numan, M. & Insel, T. R. *The neurobiology of parental behavior*. (Springer, 2003).
2. Lonstein, J. S., Lévy, F. & Fleming, A. S. Common and divergent psychobiological mechanisms underlying maternal behaviors in non-human and human mammals. *Horm. Behav.* **73**, 156–185 (2015).
3. Bridges, R. S. Neuroendocrine regulation of maternal behavior. *Front. Neuroendocrinol.* **36**, 178–196 (2015).
4. Kohl, J. & Dulac, C. Neural control of parental behaviors. *Curr. Opin. Neurobiol.* **49**, 116–122 (2018).
5. Kohl, J., Autry, A. E. & Dulac, C. The neurobiology of parenting: A neural circuit perspective. *BioEssays* **39**, e201600159 (2017).
6. Lévy, F. & Keller, M. Olfactory mediation of maternal behavior in selected mammalian species. *Behav. Brain Res.* **200**, 336–345 (2009).
7. Numan, M. & Stolzenberg, D. S. Medial preoptic area interactions with dopamine neural systems in the control of the onset and maintenance of maternal behavior in rats. *Front. Neuroendocrinol.* **30**, 46–64 (2009).
8. Afonso, V. M., Sison, M., Lovic, V. & Fleming, A. S. Medial prefrontal cortex lesions in the female rat affect sexual and maternal behavior and their sequential organization. *Behav. Neurosci.* **121**, 515–526 (2007).
9. Febo, M. A bold view of the lactating brain: functional magnetic resonance imaging studies of suckling in awake dams. *J. Neuroendocrinol.* **23**, 1009–1019 (2011).
10. Pereira, M. & Morrell, J. I. Functional Mapping of the Neural Circuitry of Rat Maternal Motivation: Effects of Site-Specific Transient Neural Inactivation: Mapping maternal motivation circuitry. *J. Neuroendocrinol.* **23**, 1020–1035 (2011).
11. Brunton, P. J. & Russell, J. A. The expectant brain: adapting for motherhood. *Nat. Rev. Neurosci.* **9**, 11–25 (2008).
12. Kinsley, C. H. & Lambert, K. G. Reproduction-Induced Neuroplasticity: Natural Behavioural and Neuronal Alterations Associated with the Production and Care of Offspring. *J. Neuroendocrinol.* **20**, 515–525 (2008).
13. Lévy, F., Gheusi, G. & Keller, M. Plasticity of the Parental Brain: A Case for Neurogenesis: Parental behaviour and neurogenesis. *J. Neuroendocrinol.* **23**, 984–993 (2011).
14. Akbari, E. M., Chatterjee, D., Lévy, F. & Fleming, A. S. Experience-dependent cell survival in the maternal rat brain. *Behav. Neurosci.* **121**, 1001–1011 (2007).
15. Hoekzema, E. *et al.* Pregnancy leads to long-lasting changes in human brain structure. *Nat. Neurosci.* **20**, 287 (2017).
16. Kohl, J. *et al.* Functional circuit architecture underlying parental behaviour. *Nature* **556**, 326–331 (2018).
17. Ashburner, J. & Friston, K. J. Unified segmentation. *NeuroImage* **26**, 839–851 (2005).
18. Sawiak, S. J., Wood, N. I., Williams, G. B., Morton, A. J. & Carpenter, T. A. Voxel-based morphometry with templates and validation in a mouse model of Huntington's disease. *Magn. Reson. Imaging* **31**, 1522–1531 (2013).
19. Sawiak, S. J., Picq, J.-L. & Dhenain, M. Voxel-based morphometry analyses of in vivo MRI in the aging mouse lemur primate. *Front. Aging Neurosci.* **6**, (2014).
20. Keifer Jr, O. P. *et al.* Voxel-based morphometry predicts shifts in dendritic spine density and morphology with auditory fear conditioning. *Nat. Commun.* **6**, (2015).
21. Lein, E. S. *et al.* Genome-wide atlas of gene expression in the adult mouse brain. *Nature* **445**, 168–176 (2007).
22. Ullmann, J. F. P. *et al.* Segmentation of the C57BL/6J mouse cerebellum in magnetic resonance images. *NeuroImage* **62**, 1408–1414 (2012).
23. Richards, K. *et al.* Segmentation of the mouse hippocampal formation in magnetic resonance images. *NeuroImage* **58**, 732–740 (2011).
24. Watson, C. *et al.* An ontologically consistent MRI-based atlas of the mouse diencephalon. *NeuroImage* **157**, 275–287 (2017).
25. Ullmann, J. F. P., Watson, C., Janke, A. L., Kurniawan, N. D. & Reutens, D. C. A segmentation protocol and MRI atlas of the C57BL/6J mouse neocortex. *NeuroImage* **78**, 196–203 (2013).
26. Wang, Q. *et al.* The Allen Mouse Brain Common Coordinate Framework: A 3D Reference Atlas. *Cell* **181**, 936–953.e20 (2020).
27. Paxinos and Franklin's the Mouse Brain in Stereotaxic Coordinates, Compact - 5th Edition. <https://www.elsevier.com/books/paxinos-and-franklins-the-mouse-brain-in-stereotaxic-coordinates-compact/franklin/978-0-12-816159-3>.

28. Deacon, R. M. J. Digging and marble burying in mice: simple methods for *in vivo* identification of biological impacts. *Nat. Protoc.* **1**, 122 (2006).

29. Tsuneoka, Y. *et al.* Neurotransmitters and neuropeptides in gonadal steroid receptor-expressing cells in medial preoptic area subregions of the male mouse. *Sci. Rep.* **7**, 1–16 (2017).

30. McHenry, J. A., Rubinow, D. R. & Stuber, G. D. Maternally responsive neurons in the bed nucleus of the stria terminalis and medial preoptic area: Putative circuits for regulating anxiety and reward. *Front. Neuroendocrinol.* **38**, 65–72 (2015).

31. Olazábal, D. E. *et al.* Flexibility and adaptation of the neural substrate that supports maternal behavior in mammals. *Neurosci. Biobehav. Rev.* **37**, 1875–1892 (2013).

32. Bridges, R. S. & Hays, L. E. Steroid-induced alterations in mRNA expression of the long form of the prolactin receptor in the medial preoptic area of female rats: Effects of exposure to a pregnancy-like regimen of progesterone and estradiol. *Brain Res. Mol. Brain Res.* **140**, 10–16 (2005).

33. Lonstein, J. S. & De Vries, G. J. Maternal behaviour in lactating rats stimulates c-fos in glutamate decarboxylase-synthesizing neurons of the medial preoptic area, ventral bed nucleus of the stria terminalis, and ventrocaudal periaqueductal gray. *Neuroscience* **100**, 557–568 (2000).

34. Miceli, M. O., Fleming, A. S. & Malsbury, C. W. Disruption of maternal behaviour in virgin and postparturient rats following sagittal plane knife cuts in the preoptic area-hypothalamus. *Behav. Brain Res.* **9**, 337–360 (1983).

35. Lee, A., Clancy, S. & Fleming, A. S. Mother rats bar-press for pups: effects of lesions of the mpoa and limbic sites on maternal behavior and operant responding for pup-reinforcement. *Behav. Brain Res.* **100**, 15–31 (1999).

36. El Majdoubi, M., Poulain, D. A. & Theodosis, D. T. Activity-dependent morphological synaptic plasticity in an adult neurosecretory system: magnocellular oxytocin neurons of the hypothalamus. *Biochim. Cell Biol.* **78**, 317–327 (2000).

37. D’Cunha, T. M., King, S. J., Fleming, A. S. & Lévy, F. Oxytocin receptors in the nucleus accumbens shell are involved in the consolidation of maternal memory in postpartum rats. *Horm. Behav.* **59**, 14–21 (2011).

38. Insel, T. R. & Harbaugh, C. R. Lesions of the hypothalamic paraventricular nucleus disrupt the initiation of maternal behavior. *Physiol. Behav.* **45**, 1033–1041 (1989).

39. Numan, M. & Corodimas, K. P. The effects of paraventricular hypothalamic lesions on maternal behavior in rats. *Physiol. Behav.* **35**, 417–425 (1985).

40. Olazábal, D. E., Kalinichev, M., Morrell, J. I. & Rosenblatt, J. S. MPOA cytotoxic lesions and maternal behavior in the rat: effects of midpubertal lesions on maternal behavior and the role of ovarian hormones in maturation of MPOA control of maternal behavior. *Horm. Behav.* **41**, 126–138 (2002).

41. Freeman, M. E., Kanyicska, B., Lerant, A. & Nagy, G. Prolactin: structure, function, and regulation of secretion. *Physiol. Rev.* **80**, 1523–1631 (2000).

42. Csérvénák, M. *et al.* Thalamic neuropeptide mediating the effects of nursing on lactation and maternal motivation. *Psychoneuroendocrinology* **38**, 3070–3084 (2013).

43. Kinsley, C. H. *et al.* Motherhood and the hormones of pregnancy modify concentrations of hippocampal neuronal dendritic spines. *Horm. Behav.* **49**, 131–142 (2006).

44. Pawluski, J. L. & Galea, L. A. M. Hippocampal morphology is differentially affected by reproductive experience in the mother. *J. Neurobiol.* **66**, 71–81 (2006).

45. Pawluski, J. L., Walker, S. K. & Galea, L. A. M. Reproductive experience differentially affects spatial reference and working memory performance in the mother. *Horm. Behav.* **49**, 143–149 (2006).

46. Eid, R. S. *et al.* Early and late effects of maternal experience on hippocampal neurogenesis, microglia, and the circulating cytokine milieu. *Neurobiol. Aging* **78**, 1–17 (2019).

47. Duarte-Guterman, P., Leuner, B. & Galea, L. A. M. The long and short term effects of motherhood on the brain. *Front. Neuroendocrinol.* **53**, 100740 (2019).

48. Gogolla, N. The insular cortex. *Curr. Biol.* **27**, R580–R586 (2017).

49. Mak, G. K. & Weiss, S. Paternal recognition of adult offspring mediated by newly generated CNS neurons. *Nat. Neurosci.* **13**, 753–758 (2010).

50. Larsen, C. M. & Grattan, D. R. Prolactin-induced mitogenesis in the subventricular zone of the maternal brain during early pregnancy is essential for normal postpartum behavioral responses in the mother. *Endocrinology* **151**, 3805–3814 (2010).

51. Sakamoto, M. *et al.* Continuous neurogenesis in the adult forebrain is required for innate olfactory responses. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 8479–8484 (2011).

629 52. Kelly, J. R. *et al.* Breaking down the barriers: the gut microbiome, intestinal permeability and stress-related
630 psychiatric disorders. *Front. Cell. Neurosci.* **9**, 392 (2015).

631 53. Arentsen, T. *et al.* The bacterial peptidoglycan-sensing molecule Pglyrp2 modulates brain development and
632 behavior. *Mol. Psychiatry* **22**, 257–266 (2017).

633 54. Barrière, D. A. *et al.* Structural and functional alterations in the retrosplenial cortex following neuropathic
634 pain. *Pain* (2019) doi:10.1097/j.pain.0000000000001610.

635 55. DiIorio, G., Brown, J. J., Borrello, J. A., Perman, W. H. & Shu, H. H. Large angle spin-echo imaging. *Magn.*
636 *Reson. Imaging* **13**, 39–44 (1995).

637 56. Ma, Y. *et al.* In vivo 3D digital atlas database of the adult C57BL/6J mouse brain by magnetic resonance
638 microscopy. *Front. Neuroanat.* **2**, (2008).

639 57. Keller, M., Pawluski, J. L., Brock, O., Douhard, Q. & Bakker, J. The α -fetoprotein knock-out mouse model
640 suggests that parental behavior is sexually differentiated under the influence of prenatal estradiol. *Horm. Behav.*
641 **57**, 434–440 (2010).

642 58. van den Brom, C. E. *et al.* Altered myocardial substrate metabolism is associated with myocardial
643 dysfunction in early diabetic cardiomyopathy in rats: studies using positron emission tomography. *Cardiovasc.*
644 *Diabetol.* **8**, 39 (2009).

645 59. Magalhães, R. *et al.* The dynamics of stress: a longitudinal MRI study of rat brain structure and connectome.
646 *Mol. Psychiatry* (2017) doi:10.1038/mp.2017.244.

647 60. Ashburner, J. & Friston, K. J. Voxel-based morphometry--the methods. *NeuroImage* **11**, 805–821 (2000).

648 61. Ashburner, J. A fast diffeomorphic image registration algorithm. *NeuroImage* **38**, 95–113 (2007).

649 62. Asami, T. *et al.* Longitudinal loss of gray matter volume in patients with first-episode schizophrenia:
650 DARTEL automated analysis and ROI validation. *NeuroImage* **59**, 986–996 (2012).

651 63. Barrière, D. A. *et al.* In vivo magnetic resonance imaging reveals the effect of gonadal hormones on
652 morphological and functional brain sexual dimorphisms in adult sheep. *Psychoneuroendocrinology* **109**, 104387
653 (2019).

654 64. Xia, M., Wang, J. & He, Y. BrainNet Viewer: a network visualization tool for human brain connectomics.
655 *PloS One* **8**, e68910 (2013).

656 **Tables**

Ex vivo / In vivo	Strain	Sex	Age (days)	Number of animals	Magnetic field intensity	Anatomical contrast	Spatial resolution	MRI template	Atlas	Regions of interest	Tissue Probability maps	References
Ex vivo	C57BL/6J	Male	63	6	9.4 Tesla	T ₁ PD T ₂ W	90 x 90 x 90 µm ³	Yes	Yes	21	No	Ali et al. 2005
In vivo	C57BL/6J	Male	100	6	11.7 Tesla	T ₁	60 x 60 x 60 µm ³	Yes	Yes	NA	No	MacKenzie-Graham et al. 2011
Ex vivo	129S1/SvImJ	Male	56	9	7 Tesla	T ₂	60 x 60 x 60 µm ³	Yes	Yes	9	No	Kovachev et al. 2005
Ex vivo	C57BL/6J	NA	PO	8	11.7 Tesla	T ₁	40 x 40 x 40 µm ³	Yes	Yes	12	Yes	Lee et al. 2005
In vivo	C57BL/6J	Male	84-98	12	9.4 Tesla	T ₂	100 x 100 x 100 µm ³	Yes	Yes	20	No	Ma et al. 2008
In vivo	C3H/HeSnJ	NA	11	15	7 Tesla	T ₁	156 x 156 x 156 µm ³	Yes	Yes	6	No	Brack et al. 2006
Ex vivo	129S1/SvImJ C57BL/6J CTD1	Male	126	27	7 Tesla	T ₂	60 x 60 x 60 µm ³	Yes	Yes	42	No	Chen et al. 2008
Ex vivo	C57BL/6J	NA	63	6	9.4 Tesla	T ₁ T ₂	21.5 x 21.5 x 21.5 µm ³ 43 x 43 x 43 µm ³	Yes	Yes	33	No	Badea et al. 2008
Ex vivo	C57BL/6J	20 males 20 females	84	40	7 Tesla	T ₁	32 x 32 x 32 µm ³	Yes	Yes	62	No	Dorr et al. 2008
Ex vivo	C57BL/6J and BXD	Male	63	12	9.4 Tesla	T ₁ T ₂	21.5 x 21.5 x 21.5 µm ³ 43 x 43 x 43 µm ³	Yes	Yes	20	No	Sharief et al. 2008
Ex vivo	C57BL/6J	Male	66-78	14	9.4 Tesla	T ₁ T ₂ T ₂ *	21.5 x 21.5 x 21.5 µm ³	Yes	Yes	37	No	Johnson et al. 2010
Ex vivo	C57BL/6J	Male	84	18	16.4 Tesla	T ₁ /T ₂ *	30 x 30 x 30 µm ³	Yes	Yes (Partial: hippocampus)	40	No	Richards et al. 2011
Ex vivo	C57BL/6J	Male	84	18	16.4 Tesla	T ₁ /T ₂ *	30 x 30 x 30 µm ³	Yes	Yes (Partial: cerebellum)	38	No	Ullman et al. 2013
Ex vivo	C57BL/6J	Male	84	18	16.4 Tesla	T ₁ /T ₂ *	30 x 30 x 30 µm ³	Yes	Yes (Partial: basal ganglia)	35	No	Ullman et al. 2013
Ex vivo	C57BL/6J	Male	84	18	16.4 Tesla	T ₁ /T ₂ *	30 x 30 x 30 µm ³	Yes	Yes (Partial: neocortex)	74	No	Ullman et al. 2013
In vivo	C57BL/6J	Male	126	82	4.7 Tesla	T ₂	70 x 70 x 70 µm ³	No	NA	NA	Yes	Sawak et al. 2013
Ex vivo	C57BL/6J	Male	84	18	16.4 Tesla	T ₁ /T ₂ *	30 x 30 x 30 µm ³	Yes	Yes (Partial: diencephalon)	89	No	Watson et al. 2017
Ex vivo	C57BL/6J	1,051 males 621 female	77	1675	NA	NA	10 x 10 x 10 µm ³	No	Yes	1,327	No	Wang et al. 2020

657

658 **Table 1.** Comparison of mouse brain resources currently available in the literature (NA = not
659 available).

Figures

Figure 1. Processing of mouse brain templates and building an atlas from the AMBMC template and Allen Brain Atlas for data analysis and visualization. To create our resources, we used both AMBMC mouse brain template and the mouse Allen Brain Atlas and its associated Nissl images. (1) We down-sampled to a suitable resolution for MRI analysis (60-μm isotropic resolution) and (2) realign in the AC/PC axis. The resulting template was then segmented into GM, WM and CSF probability maps (3). These probability maps were used to segment all the images which have been previously normalized to the template (5,6,7). We obtained a large set of 138 images for each tissue class which have been used to build a population-specific GM, WM and CSF priors. using an exponentiated lie algebra (DARTEL) approach (8). This new set of population-specific tissue priors was used to segment again normalized T₂ images (9) for the final VBM preprocessing (10, 11, 12). To normalize the Allen Brain Atlas, we manually realign (13) and normalized the associated Nissl-stain mouse brain using the GM priors generated previously (14, 15). Both linear and nonlinear transformations have been applied to the Allen mouse brain atlas. Then, a visual inspection of

676 each normalized label was carried out and, when necessary, redrawn according to Paxinos and
 677 Franklin's atlas. Finally, the olfactory bulb and hind brain regions were completed, the *corpus*
 678 *callosum* and ventricles were drawn from the WM and CSF priors, and the cerebellum labels
 679 were replaced by the AMBMC cerebellum labels.
 680 AMBMC = Australian Mouse Brain Mapping Consortium mouse brain template, AC/PC =
 681 anterior commissure/posterior commissure, CSF = cerebrospinal fluid, GM = gray matter,
 682 WM = white matter.

cerebrospinal fluid probabilistic maps and the associated anatomical atlas (60- μ m isotropic resolution). **(B)** Dorsal (left panel) and lateral (right panel) 3D representations of the anatomical mouse brain atlas. **(C)** Lateral views of the cortical areas after normalization of the Allen Mouse Brain Atlas to the AMBMC anatomical template. The cortex was segmented into cortical areas such as the cingulate, visual, auditory, frontal, visceral, insular, olfactory, retrosplenial, motor, somatosensory, temporal, entorhinal and gustatory systems. Each area was subdivided into secondary areas (*e.g.*, primary and secondary motor cortices) or structural areas (*i.e.*, agranular, dysgranular, agranular/dysgranular, granular and posterior agranular insular cortices). **(D)** The four images depict the different cortical layers (I, II/III, IV/V and VIa/VIb). **(E and F)** 3D rendering and axial sections of subcortical structures (hypothalamus and hippocampus).

Legend for labeled regions: Hypothalamus: *ARh* = *arcuate hypothalamic nucleus*; *LHA* = *lateral hypothalamic area*; *LPOA* = *lateral preoptic area*; *MPOA* = *medial preoptic area*; *MPOn* = *medial preoptic nucleus*; *PH* = *posterior hypothalamic nucleus*. *AHN* = *anterior hypothalamic nucleus*.

Hippocampus: *Apr* = *area prostriata*; *CA1*, *CA2*, *CA3* = *cornu ammonis areas 1, 2 and 3*; *DG* = *dentate gyrus*; *FC* = *fasciola cinerea*; *PARA* = *parasubiculum*; *POST* = *postsubiculum*; *PRO* = *prosubiculum*; *PRE* = *presubiculum*. For further details, see https://www.nitrc.org/projects/tmbta_2019.

Figure 3. Longitudinal effects of the reproductive cycle on brain morphometry. Coronal slices (at left) and brain plots (at right) showing gray matter concentration (GMC) differences between control and parous animals at the end of gestation (**A**) and during early lactation (**B**) and late lactation (**C**). SPM flexible factorial analysis revealed an interaction between control mice and parous mice in the late gestation period (**A**), early lactation period (**B**) and late lactation period (**C**); voxel-

712 level threshold $p < 0.01$, $t_{(126)} = 2.356$, cluster threshold = 25 voxels. *BNST* = *bed nucleus of*
 713 *the stria terminalis*; *Hyp* = *hypothalamus*; *mPOA* = *medial preoptic area*; *dHippo* = *dorsal*
 714 *hippocampus*; *ObF/PrL* = *orbitofrontal/prelimbic area*; *aOb/Ob* = *accessory olfactory*
 715 *nucleus/olfactory bulb*; *Pir* = *piriform cortex*; *Amyg/Ent* = *amygdala/entorhinal cortex*.

Figure 4. Longitudinal time course analysis of gray matter concentration (GMC) over the reproductive cycle. Time course comparisons in GMC between the control (black dots and lines) and parous (red dots and lines) groups showing 3 different time profiles. (A) GMC values within the medial preoptic area, the bed nucleus of *stria terminalis* (BNST) and the agranular insular cortex reveal a significant increase in GMC during the late gestation (LG) period maintained until weaning (W). (B) Specific and transient increases in GMCs are observed in the amygdala, the piriform area and the hippocampus during early lactation (EL). (C) The arcuate nucleus, PVN and caudate putamen display an increase in GMC through both EL and late lactation (LL) periods. Orange and blue areas represent the gestation and lactation periods, respectively. Data are expressed as the mean $\pm$ standard error of the mean (SEM); two-way ANOVA followed by Holm-Sidak multiple comparisons test; * $p < 0.05$, ** $p < 0.01$ and *** $p < 0.001$, compared with control mice.

Figure 5. Distribution of animals according to the quality of their maternal behavior assessed with the pup retrieval test and brain morphometric. K-means clustering of parous animals to classify mice into low and high maternal behavior groups based on behavior during the pup retrieval test (**A**). Comparisons between the low and high maternal behavior groups revealed significant differences in crouching, pup-licking and digging times (**B**). Brain slices (left panel) and brain plots (right panel) comparing gray matter concentration (GMC) modifications and surface maps of GMC differences between females exhibiting low and

those exhibiting high maternal behavior at the end of the gestation period (C) and early lactation period (D).

Low and high maternal behavioral data were compared using a Student's t-test with post hoc corrections for multiple comparisons using an FDR approach ($Q = 1\%$) and are expressed as the mean $\pm$ SEM; ** $p < 0.01$ and *** $p < 0.001$. SPM flexible factorial analysis revealed an interaction between low and high maternal behavior parous mice in the late gestation period (A) and early lactation period (B); voxel-level threshold $p < 0.01$, $t_{(60)} = 2.39$, cluster threshold = 25 voxels. *RS* = retrosplenial cortex; *vHippo* = ventral hippocampus; *S1* = primary somatosensory cortex; *aOB* = accessory olfactory bulb; *OB* = olfactory bulb; *Pir* = piriform cortex; *EntH* = entorhinal cortex; *Str* = striatum; *OrbF* = orbitofrontal cortex.

Figure 6. Longitudinal analysis in gray matter concentration (GMC) during the reproductive cycle in control females and females exhibiting low or high maternal behavior. Time-course comparison of GMC between the control (black dots and lines), low maternal behavior (pink dots and lines) and high maternal behavior (red dots and lines) groups revealed two types of

time profile. GMC analysis in the entorhinal area, orbitofrontal area, the accessory olfactory bulb and the medial preoptic nucleus revealed an acute and specific increase in GMC values in the high maternal behavior group at the late gestation period (**A**). In contrast, GMC analysis in the ventral hippocampus, the retrosplenial area and the primary somatosensory area revealed an increase in GMC in the high maternal behavior group at the late gestation period, and this increase was maintained until weaning (**B**).

Orange and blue areas represent the gestation and lactation periods, respectively. Data were compared using a two-way ANOVA followed by Holm-Sidak multiple comparisons test and expressed as the mean $\pm$ SEM; * $p < 0.05$, ** $p < 0.01$ when high maternal mice were compared with control mice and $^{**} p < 0.01$, $^{***} p < 0.001$ when high maternal mice were compared with low maternal mice.

764

765 **Figure 7.** Estimation of the sensitivity and specificity of late-gestation GMC measures in the
766 entorhinal area (A) and accessory olfactory bulb (B) to predict postpartum maternal
767 performance.

768 Receiver operating characteristic (ROC) curves were estimated using a Wilson/Brown test
769 with a 95% confidence interval. Correlations were estimated using a Pearson correlation test.

770 ROC and correlation analyses were considered significant at $p < 0.05$.

Supplementary Data

# Label	Region of Interest	Hemisphere	T Value	p Value	x (mm)	y (mm)	z (mm)
712	Visceral area, layer 2/3	Left	2.3789	0.0094	-41.5	-3.5	-3.5
1842	Agranular insular area, dorsal part, layer 2/3	Left	3.9264	< 0.0001	-34.5	16.5	-8.5
652	Gustatory areas, layer 2/3	Left	3.4262	0.0004	-30.5	25.5	-4.5
2562	Cortical amygdalar area, posterior part, medial zone	Left	2.6591	0.0044	-23.5	-16.5	-18.5
3002	Olfactory tubercle	Left	4.2298	< 0.0001	-22.5	13.5	-23.5
2502	Piriform area	Left	3.6483	0.0002	-21.5	26.5	-14.5
5122	Medial amygdalar nucleus	Left	2.5052	0.0067	-21.5	-10.5	-11.5
2972	Caudoputamen	Left	2.5003	0.0068	-20.5	7.5	-6.5
3522	Lateral dorsal nucleus of thalamus	Left	2.3933	0.0090	-17.5	-4.5	4.5
292	Primary somatosensory area, lower limb, layer 2/3	Left	2.4134	0.0086	16.5	4.5	23.5
1602	Orbital area, lateral part, layer 2/3	Left	2.4677	0.0074	-15.5	37.5	-0.5
2462	Anterior olfactory nucleus	Left	2.3608	0.0098	-11.5	33.5	-5.5
4132	Lateral hypothalamic area	Left	2.5217	0.0064	-10.5	-10.5	-18.5
2982	Nucleus accumbens	Left	2.3953	0.0090	-9.5	18.5	-15.5
3192	Diagonal band nucleus	Left	2.5835	0.0054	-5.5	14.5	-22.5
1731	Orbital area, medial part, layer 1	Right	2.5912	0.0053	1.5	37.5	0.5
2421	Main olfactory bulb	Right	3.7602	0.0001	2.5	35.5	-17.5
3861	Medial preoptic area	Right	2.6983	0.0040	6.5	11.5	-21.5
3971	Anterior hypothalamic nucleus	Right	2.4173	0.0085	8.5	-0.5	-20.5
4201	Tuberal nucleus	Right	2.4250	0.0083	8.5	-3.5	-20.5
4131	Lateral hypothalamic area	Right	3.3026	0.0006	11.5	-10.5	-16.5
11	Frontal pole, layer 1	Right	2.5614	0.0058	12.5	41.5	5.5
4141	Lateral preoptic area	Right	2.5685	0.0057	12.5	5.5	-18.5
2461	Anterior olfactory nucleus	Right	2.4141	0.0086	14.5	35.5	7.5
2631	Dentate gyrus, molecular layer	Right	2.4871	0.0071	14.5	-16.5	12.5
3121	Medial amygdalar nucleus	Right	2.6129	0.0050	21.5	-10.5	-11.5
3001	Olfactory tubercle	Right	2.3745	0.0095	22.5	10.5	-24.5
1841	Agranular insular area, dorsal part, layer 2/3	Right	3.0066	0.0016	23.5	32.5	-2.5
2501	Piriform area	Right	2.9740	0.0018	26.5	24.5	-12.5
2971	Caudoputamen	Right	2.7880	0.0031	27.5	-7.5	-5.5
621	Supplemental somatosensory area, layer 6a	Right	2.4045	0.0088	29.5	8.5	2.5
591	Supplemental somatosensory area, layer 2/3	Right	2.5642	0.0057	36.5	16.5	-0.5

Table S1. Local variation in gray matter concentration between control and parous mice at the end of the gestation period. SPM flexible factorial analysis revealed an interaction between the control and parous groups at the late gestation time point.

# Label	Region of Interest	Hemisphere	T Value	p Value	x {mm}	y {mm}	z {mm}
712	Visceral area, layer 2/3	Left	4.0471	< 0.0001	-41.5	3.5	3.5
592	Supplemental somatosensory area, layer 2/3	Left	4.2243	< 0.0001	-40.5	6.5	-2.5
1842	Agranular insular area, dorsal part, layer 2/3	Left	3.9710	< 0.0001	-34.5	16.5	-8.5
352	Primary somatosensory area, mouth, layer 2/3	Left	1.9021	0.0293	-31.5	22.5	5.5
2972	Caudoputamen	Left	1.8455	0.0331	-31.5	-8.5	-0.5
342	Primary somatosensory area, mouth, layer 1	Left	1.7124	0.0437	-30.5	23.5	6.5
72	Primary motor area, layer 2/3	Left	1.7423	0.0411	-29.5	27.5	-0.5
232	Primary somatosensory area, barrel field, layer 2/3	Left	2.0289	0.0220	-27.5	6.5	18.5
532	Primary somatosensory area, unassigned, layer 2/3	Left	1.7753	0.0384	-27.5	11.5	16.5
2812	Subiculum	Left	3.0917	0.0012	-25.5	-24.5	10.5
2642	Dentate gyrus, polymorph layer	Left	1.7318	0.0420	-20.5	-23.5	-9.5
4092	Ventral premammillary nucleus	Left	2.3287	0.0107	-8.5	-14.5	-18.5
5522	Medial vestibular nucleus	Left	1.9545	0.0261	-5.5	-47.5	1.5
4442	Superior colliculus, motor related, intermediate gray layer	Left	2.9857	0.0017	2.5	14.5	13.5
2411	Olfactory areas	Right	2.0350	0.0217	0.5	23.5	-11.5
4031	Medial preoptic area	Right	1.6858	0.0461	5.5	-10.5	-19.5
2421	Main olfactory bulb	Right	2.4342	0.0081	2.5	34.5	-17.5
4541	Olivary pretectal nucleus	Right	2.6632	0.0044	6.5	-12.5	11.5
3211	Red nuclei of the stria terminalis	Right	1.7760	0.0383	10.5	8.5	-6.5
4131	Lateral hypothalamic area	Right	2.0659	0.0202	11.5	-9.5	-17.5
4211	Zona incerta	Right	1.9015	0.0293	11.5	8.5	11.5
2791	Presubiculum	Right	3.0783	0.0013	15.5	-18.5	12.5
3121	Medial amygdalar nucleus	Right	1.8183	0.0351	16.5	-10.5	-16.5
2561	Cortical amygdalar area, posterior part, medial zone	Right	3.5177	0.0003	22.5	-15.5	-19.5
401	Primary somatosensory area, upper limb, layer 1	Right	1.7139	0.0435	24.5	9.5	19.5
531	Primary somatosensory area, unassigned, layer 2/3	Right	1.7406	0.0412	26.5	9.5	18.5
61	Primary motor area, layer 1	Right	1.7713	0.0385	27.5	27.5	7.5
221	Primary somatosensory area, barrel field, layer 1	Right	2.2903	0.0118	27.5	5.5	19.5
2731	Entorhinal area, medial part, dorsal zone, layer 1	Right	1.9890	0.0241	27.5	-32.5	-1.5
231	Primary somatosensory area, barrel field, layer 2/3	Right	1.8671	0.0316	29.5	7.5	17.5
2571	Piriform-amygdalar area	Right	3.3855	0.0005	29.5	-6.5	-24.5
351	Primary somatosensory area, mouth, layer 2/3	Right	1.9950	0.0238	32.5	14.5	11.5
171	Primary somatosensory area, nose, layer 2/3	Right	1.8466	0.0330	33.5	12.5	11.5
2501	Piriform area	Right	3.4453	0.0004	36.5	15.5	17.5

Table S2. Local variation in gray matter concentration between control and parous mice at the beginning of the lactation period. SPM flexible factorial analysis revealed an interaction between the control and parous groups at the early lactation time point.

# Label	Region of Interest	Hemisphere	T Value	p Value	x {mm}	y {mm}	z {mm}
712	Visceral area, layer 2/3	Left	2.1711	0.01566	-11.5	0.5	2.5
2812	Subiculum	Left	2.0845	0.01938	-28.5	-24.5	9.5
2972	Caudoputamen	Left	2.8735	0.00238	-28.5	-7.5	-3.5
2502	Piriform area	Left	2.3017	0.01143	-27.5	16.5	-16.5
2562	Cortical amygdalar area, posterior part, medial zone	Left	2.2071	0.01445	-23.5	-17.5	-17.5
2992	Fundus of striatum	Left	2.0065	0.0232	-20.5	8.5	-17.5
4442	Superior colliculus, motor related, intermediate gray layer	Left	1.9022	0.02927	-16.5	-21.5	10.5
2142	Anterior visual area, layer 2/3	Left	1.7105	0.04382	-13.5	-0.5	25.5
4082	Dorsal premammillary nucleus	Left	2.6229	0.00488	-5.5	-14.5	-15.5
3011	Lateral septal nucleus, caudal (caudodorsal) part	Right	2.2155	0.01416	-1.5	8.5	7.5
3861	Medial preoptic area	Right	2.8305	0.0027	6.5	11.5	-21.5
3211	Bed nuclei of the stria terminalis	Right	1.7190	0.04052	7.5	14.5	12.5
4131	Lateral hypothalamic area	Right	1.7901	0.03721	11.5	-6.5	-17.5
2621	Field CA3	Right	2.6408	0.00465	21.5	-18.5	-6.5
5161	Dorsal cochlear nucleus	Right	1.7477	0.04063	23.5	-43.5	-4.5
2561	Cortical amygdalar area, posterior part, medial zone	Right	1.8753	0.03104	24.5	-18.5	-17.5
2731	Entorhinal area, medial part, dorsal zone, layer 1	Right	1.9285	0.02763	27.5	-32.5	-1.5
2971	Caudoputamen	Right	2.8706	0.0024	30.5	8.5	1.5
701	Visceral area, layer 1	Right	1.8073	0.03589	41.5	3.5	-3.5

Table S3. Local variation in gray matter concentration between control and parous mice at the end of the lactation period. SPM flexible factorial analysis revealed an interaction between the control and parous groups at the late lactation time point.

# Label	Region of Interest	Hemisphere	T Value	p Value	x {mm}	y {mm}	z {mm}
992	Ventral auditory area, layer 6b	Left	2.4578	0.0084	-36.5	-15.5	5.5
2292	Temporal association areas, layer 6a	Left	2.7134	0.0043	-36.5	-24.5	8.5
242	Primary somatosensory area, barrel field, layer 4	Left	2.4559	0.0084	-33.5	0.5	12.5
2752	Entorhinal area, medial part, dorsal zone, layer 3	Left	3.7707	0.0002	-30.5	-29.5	-9.5
2812	Subiculum	Left	2.6370	0.0053	-29.5	-23.5	10.5
2212	Rostrolateral visual area, layer 4	Left	2.7190	0.0043	-23.5	-7.5	22.5
2622	Field CA3	Left	4.2394	0.0000	21.5	18.5	6.5
4362	Substantia nigra, reticular part	Left	2.4952	0.0076	-17.5	-23.5	-5.5
72	Primary motor area, layer 2/3	Left	2.5921	0.0060	-12.5	13.5	20.5
2162	Retrosplenial area, dorsal part, layer 6a	Left	3.1820	0.0012	-11.5	-0.5	23.5
4132	Lateral hypothalamic area	Left	3.0784	0.0016	-11.5	-6.5	-17.5
2462	Anterior olfactory nucleus	Left	4.0594	0.0001	10.5	31.5	12.5
4212	Zona incerta	Left	3.0013	0.0020	-9.5	-10.5	-10.5
3862	Medial preoptic area	Left	2.6620	0.0050	-8.5	4.5	-20.5
4412	Superior colliculus, motor related, deep gray layer	Left	2.8740	0.0028	-8.5	-23.5	16.5
4012	Medial mammillary nucleus, medial part	Left	2.4376	0.0088	-4.5	-16.5	-16.5
4112	Ventromedial hypothalamic nucleus	Left	2.5173	0.0072	-3.5	-6.5	-22.5
4272	Superior colliculus, zonal layer	Left	2.8049	0.0034	2.5	14.5	17.5
4052	Tubermammillary nucleus, dorsal part	Left	2.4993	0.0076	-1.5	-15.5	-19.5
4441	Superior colliculus, motor related, intermediate gray layer	Right	2.8749	0.0062	0.5	-26.5	19.5
4451	Periaqueductal gray	Right	2.4815	0.0079	3.5	-32.5	13.5
2651	Dentate gyrus, granule cell layer	Right	2.7718	0.0037	7.5	7.5	10.5
4111	Ventromedial hypothalamic nucleus	Right	2.6915	0.0046	7.5	-6.5	-21.5
4431	Superior colliculus, motor related, intermediate white layer	Right	2.4866	0.0078	9.5	-28.5	18.5
2421	Main olfactory bulb	Right	3.9879	0.0001	10.5	38.5	-8.5
1071	Anteromedial visual area, layer 2/3	Right	4.0275	0.0001	12.5	-4.5	25.5
2971	Caudoputamen	Right	2.4763	0.0080	16.5	10.5	-7.5
2411	Olfactory areas	Right	3.9513	0.0001	17.5	30.5	-7.5
2621	Field CA3	Right	3.7783	0.0002	25.5	-17.5	-8.5
2811	Subiculum	Right	2.4734	0.0081	25.5	23.5	11.5
1241	Posterolateral visual area, layer 1	Right	2.5546	0.0066	27.5	-31.5	19.5
2741	Entorhinal area, medial part, dorsal zone, layer 2	Right	4.0416	0.0001	29.5	-31.5	-8.5

Table S4. Local variation in gray matter concentration between low maternal behavior parous mice and high maternal behavior parous mice at the end of the gestation period. SPM flexible factorial analysis revealed an interaction between the low *versus* high maternal behavior groups at the late gestation time point.

# Label	Region of Interest	Hemisphere	T Value	p Value	x {mm}	y {mm}	z {mm}
2732	Entorhinal area, medial part, dorsal zone, layer 1	Left	2.5882	0.0060	28.5	33.5	1.5
2632	Dentate gyrus, molecular layer	Left	2.5173	0.0072	-20.5	-16.5	6.5
412	Primary somatosensory area, upper limb, layer 2/3	Left	3.1161	0.0014	-19.5	14.5	18.5
2602	Field CA1	Left	2.5276	0.0070	-19.5	-11.5	17.5
122	Secondary motor area, layer 2/3	Left	2.8890	0.0027	-15.5	24.5	15.5
2422	Main olfactory bulb	Left	3.5084	0.0004	-8.5	45.5	6.5
3652	Parafascicular nucleus	Left	3.0525	0.0017	-3.5	-12.5	-1.5
2672	Induseum griseum	Left	2.4662	0.0082	-1.5	19.5	5.5
1482	Anterior cingulate area, dorsal part, layer 1	Left	2.4675	0.0082	-0.5	21.5	14.5
2082	Retrosplenial area, ventral part, layer 1	Left	2.3913	0.0099	-0.5	-3.5	23.5
2421	Main olfactory bulb	Right	2.7201	0.0043	2.5	43.5	-14.5
3651	Parafascicular nucleus	Right	2.9370	0.0023	6.5	12.5	0.5
2411	Olfactory areas	Right	3.2967	0.0008	8.5	42.5	4.5
2601	Field CA1	Right	3.6131	0.0003	18.5	-9.5	17.5
381	Primary somatosensory area, mouth, layer 6a	Right	3.1451	0.0013	24.5	14.5	5.5
2821	Prosubiculum	Right	2.9853	0.0020	24.5	-19.5	14.5
2691	Entorhinal area, lateral part, layer 2	Right	3.0830	0.0015	38.5	29.5	5.5

Table S5. Local variation in gray matter concentration between low maternal behavior parous mice and high maternal behavior parous mice at the beginning of the lactation period. SPM flexible factorial analysis revealed an interaction between the low *versus* high maternal behavior groups at the early lactation time point.