Whole-genome sequence analysis reveals evolution of antimicrobial resistance in a Ugandan colistin resistant *Acinetobacter baumannii*

Dickson Aruhomukama^{1,2}, Ivan Sserwadda² and Gerald Mboowa^{2, 3*}

1. **Dickson Aruhomukama:** ¹Department of Medical Microbiology, College of Health Sciences, School of Biomedical Sciences, Makerere University, Kampala, 7072, Uganda, ²Department of Immunology and Molecular Biology, College of Health Sciences, School of Biomedical Sciences, Makerere University, Kampala, 7072, Uganda. Email: dickson.aruhomukama@chs.mak.ac.ug; Mobile number: +256 706 511287
2. **Ivan Sserwadda:** ²Department of Immunology and Molecular Biology, College of Health Sciences, School of Biomedical Sciences, Makerere University, Kampala, 7072, Uganda. Email: ivangunz23@gmail.com; Mobile number: +256 702 208225
3. Correspondence: **Gerald Mboowa;** ²Department of Immunology and Molecular Biology, College of Health Sciences, School of Biomedical Sciences, Makerere University, Kampala, 7072, Uganda, ³The African Center of Excellence in Bioinformatics and Data Intensive Sciences, the Infectious Disease Institute, Makerere University P.O. Box 22418, Kampala, Uganda. Email: gerald.mboowa@chs.mak.ac.ug; Mobile number: +256 775 766249

Abstract

In recent times, pan-drug resistant *Acinetobacter baumannii* have emerged and continue to spread among critically ill patients, this poses an urgent risk to global and local human health. This study sought to provide the first genomic analysis of a pan-drug resistant *Acinetobacter baumannii* from Uganda and Africa, and to tell a story of mobile genetic element-mediated antibiotic resistance evolution in the isolate. It was an in-silico study in which intrinsic and acquired antibiotic resistance genes, and/or chromosomal resistance mutations were identified using PATRIC, CARD, NDARO and ResFinder. Screening for insertion sequences was done using ISfinder. Also, plasmid screening, phylogenetic analysis and sequence typing were performed using PlasmidFinder, PATRIC and Gubbin, and MLST respectively.

The isolate belonged to the Sequence type 136, belonging to Clonal complex 208 and Global complex 2. This isolate shared close homology with strains from Tanzania. Resistance in the isolate was chromosomally and mobile genetic element-mediated by *Acinetobacter*-derived cephalosporinases and carbapenem hydrolyzing class D β -lactamses, *bla*_{OXA-2}, 51, 58, 88, 317, *bla*_{ADC-2}, 25. Colistin resistance was associated with previously documented mutants, *lpxA* and *lpxC*. Other key resistance genes identified were: *aph(3'')-Ib*, *aph(6)-Id*, *aph(3')-Ia*, *aac(3)-Ild*, *aac(3)-IIa*, *aph(3')-I*, *aph(3'')-I*, *aph(6)-Ic*, *aph(6)-Id*, *aac(3)-II*, III, IV, VI, VIII, IX, X, *macA*, *macB*, *tetA*, *tetB*, *tetR*, *dfrA*, and those of the *floR* family. RSF1010 like IncQ broad-host-range plasmids and features of pACICU1, pACICU2, and p3ABAYE *Acinetobacter baumannii* plasmids namely partitioning proteins ParA and B were present. Insertion sequences present included IS3, IS5, IS66 and those of the ISLre2 families.

The study described for the first time a pan-drug resistant *Acinetobacter baumannii* from Uganda, and told a story of mobile genetic element-mediated antibiotic resistance evolution in

the isolate despite being limited by pan-drug resistance phenotypic data. It provides a basis to track trends in antibiotic resistance and identification of emerging resistance patterns in *Acinetobacter baumannii* in Uganda.

Keywords: *Acinetobacter baumannii*, Whole-genome sequencing, Pan-drug resistance, Uganda

Background

Current times have witnessed the emergence and continuous spread of pan-drug resistant *Acinetobacter baumannii* among critically ill patients, this poses an urgent risk to global human health¹. The World Health Organisation (WHO) in its 2017 report on "priority pathogens – a catalog of 12 families of bacteria that pose the greatest threat to human health" categorized *Acinetobacter baumannii* in priority 1 or the critical category¹. *Acinetobacter baumannii* is a Gram-negative opportunist pathogen with a high incidence in immunocompromised individuals, particularly those attending intensive-care units². This organism exhibits an extraordinary ability to upregulate or acquire antibiotic resistance genes, a characteristic that enhances its spread and, makes it one of the bacteria that threatens the current antibiotic era². However, limited information remains on the genetic antimicrobial resistance determinants as well as the mobile genetic element-mediated antibiotic resistance evolution in *Acinetobacter baumannii* especially from Uganda. This work sought to fill this gap by providing the first genomic analysis of a pan-drug resistant *Acinetobacter baumannii* from Uganda, and to tell a story of mobile genetic element-mediated antibiotic resistance evolution in the isolate.

66 **Materials and methods**

67 **Isolate metadata and source of whole-genome sequences.** *Acinetobacter baumannii*
 68 SRR8291681_Uganda whole-genome sequences were obtained from the NCBI's Sequence
 69 Read Archive (SRA) repository, these had been submitted in December 2018 by J. Craig
 70 Venter Institute (JCVI) under a study entitled the whole-genome sequencing of carbapenem-
 71 resistant Gram-negative clinical isolates from Mbarara Regional Referral Hospital, Western-
 72 Uganda and BioProject: PRJNA508495 (<https://www.ncbi.nlm.nih.gov/sra/SRR8291681/>). In
 73 addition, 8 other available *Acinetobacter baumannii* whole-genome sequences for comparison
 74 were purposively obtained from the same source, and these were from; Tanzania, South Korea,
 75 United States, Canada, Germany, and Sweden. Mbarara Regional Referral Hospital is located
 76 in Mbarara Municipality, which is 286 km south west of Kampala the capital city of Uganda. It
 77 is a public hospital funded by the Government of Uganda through Ministry of Health. It is the
 78 referral hospital for south western Uganda serving 10 districts with a population of more than
 79 2.5 million people. It also receives patients from neighbouring countries of Rwanda, Tanzania
 80 and Democratic Republic of Congo.

81 **Whole-genome sequencing.** The whole genome of *Acinetobacter baumannii*
 82 SRR8291681_Uganda was sequenced using the Illumina NextSeq 500.

83 **Read quality assessment, genome assembly and annotation.** FastQC v0.11.8
 84 (<https://www.bioinformatics.babraham.ac.uk/projects/download.html>) was used to assess the
 85 quality of the reads. *De novo* genome assembly of the Illumina NextSeq 500 reads from
 86 *Acinetobacter baumannii* SRR8291681_Uganda was performed using the Unicycler v0.4.8-
 87 beta⁵² with SPAdes v3.11.1⁵³, with default parameters. Following genome assembly, gene
 88 calling and automatic functional annotation were done for all the contigs using the

Pathosystems Resource Integration Center (PATRIC 3.5.36) identifying 4,037 coding regions on the chromosome with 64 tRNA and 3 rRNA. PATRIC uses the RAST tool kit to provide an annotation of genomic features in default parameters (<https://www.patricbrc.org/>).

Identification of antibiotic resistance genes. The identification of antibiotic resistance genes from the complete Illumina NextSeq 500 assembly was performed using PATRIC 3.5.36. Additional screening for antibiotic resistance genes was performed by comparison (BLASTp; sequence identity $\geq 40\%$; E-value ≤ 0.0001) of all predicted coding regions against the PATRIC's Antibiotic Resistance Genes Database, the Comprehensive Antimicrobial Resistance Database (CARD)⁵⁴, and the National Database of Antibiotic Resistant Organisms (NDARO) (<https://www.ncbi.nlm.nih.gov/pathogens/antimicrobial-resistance/>). Furthermore, acquired antibiotic resistance genes and/or chromosomal mutations were identified using ResFinder version 3.1 (<https://cge.cbs.dtu.dk/services/ResFinder/>) at a threshold for percentage identity of 90% and a minimum length of 60%. Manual inspection of the antibiotic resistance genes was performed to improve their functional annotation based on known and correct start sites as well as to identify point mutations which may contribute to a resistant phenotype. Resistance gene loci were screened for known insertion sequences by comparison against the ISFinder database set at a minimum % coverage of 70% and a threshold for minimum % identity of 30% (<https://isfinder.biotoul.fr/>). Finally, screening for the presence of plasmids using PlasmidFinder 2.0 set at 60% minimum % coverage and the threshold for minimum % identity 95% (<https://cge.cbs.dtu.dk/services/PlasmidFinder/>) was done.

Phylogenetic analysis and multilocus sequence typing (MLST). The building of the phylogenetic tree was performed using the PATRIC server with only 1,000 genes selected randomly from the assembled contigs and default parameters including no genomic duplications and insertions. Multilocus sequence typing was performed using MLST v2.0

(<https://cge.cbs.dtu.dk/services/MLST/>). The results of phylogenetic analysis obtained from the the PATRIC server were verified using Genealogies Unbiased By recomBinations In Nucleotide Sequences (Gubbins) (<https://sanger-pathogens.github.io/gubbins/>), this iteratively identifies loci containing elevated densities of base substitutions while concurrently constructing a phylogeny based on the putative point mutations outside of these regions enabling more accurate, recombination free phylogenetic analysis of *Acinetobacter baumannii* genomes known to be full of regions acquired as a result of recombination events.

Results

Read quality assessment. FastQC v0.11.8 was used to assess the quality of the reads, *Acinetobacter baumannii* SRR8291681_Uganda was found to have a coverage of >30X, generated 6.19 million paired-end 150 bp genomic reads. The assembled genome of *Acinetobacter baumannii* SRR8291681_Uganda has N50 and L50 of 337442 and 5 respectively.

General genome features of *Acinetobacter baumannii* SRR8291681_Uganda. SRR8291681_Uganda consists of a circular chromosome spanning 4173379 base pairs in length with an average G+C content of 38.95%. The genome quality was generally good, with coarse and fine consistencies accounting for 98.7% and 97.6% respectively. The chromosome of the isolate shares close homology with *Acinetobacter baumannii* ERR1989115 and ERR1989084 strains originating from Tanzania (**Figure 1**). Multilocus sequence typing (MLST) revealed that SRR8291681_Uganda belonged to the previously reported sequence type (ST) 136³, this ST have been documented to belong to clonal complex 208 (CC208) and global complex 2 (GC2)⁴.

Figure 1. Phylogenetic tree. This shows the phylogenetic relationship between *Acinetobacter baumannii* SRR8291681_Uganda, strains from Tanzania (ERR1989115, ERR1989100, ERR1989084), those collected outside Africa (i.e. (SRR8289559_South Korea, SRR3666962_United States, SRR5739056_Canada, ERR3197698_Germany, and SRR6037664_Sweden)), as well as strains AB0057, AYE, ACICU, and AB5075.

Genetic determinants of antibiotic resistance. To determine the genetic basis of antibiotic resistance, we interrogated the genome to identify acquired and intrinsic antibiotic resistance genes (**Table 1**). The acquired resistance genes were distributed throughout the genome (**Figure 2**). Resistance to β -lactams (i.e. penams, second-, third, and fourth-generation cephalosporins, cephamycins and carbapenems) was mediated by the previously described *Acinetobacter*-derived cephalosporinases (ADCs) and carbapenem-hydrolyzing class D β -lactamase genes (CHDLs)⁵⁻⁷. Allelic copies or mutants (i.e. *lpxA* and *lpxC*) that have previously been linked with colistin resistance were also present^{8,9}. Tetracycline resistance was mediated by the previously described acquired narrow-spectrum efflux pumps, such as the major

facilitator superfamily (MFS) members¹⁰; an allelic copy or mutant (*tetR*) that has been linked with tetracycline resistance was also identified^{11,12}. Aminoglycoside resistance genes previously described were also identified^{13,14}. Macrolide resistance was mediated by chromosomally encoded macrolide-specific efflux proteins previously described¹⁵. Resistance to trimethoprim was mediated by a previously described trimethoprim resistance gene^{16,17}. Sulphonamide resistance was mediated by a previously described mobile genetic element-encoded sulphonamide resistance gene^{17,18}. Phenicol resistance was mediated by phenicol resistance genes previously described^{19,20}. Mobile genetic elements (i.e. plasmids) previously documented to be mobilized into a number of species of Gram-negative bacteria by co-resident conjugative plasmids were also present, these were the RSF1010 like IncQ broad-host-range plasmids^{21–24}; In addition, features of previously described *Acinetobacter baumannii* plasmids (i.e. pACICU1, pACICU2, and p3ABAYE)^{25,26} that mostly occur in clinical isolates were also present, these were the chromosome (plasmid) partitioning proteins ParA and ParB^{27,28}. ISFinder analysis indicated that the genome contains numerous insertion sequences (IS), the majority of the insertion sequences belonging to the IS3, IS5, IS66 and ISLre2 families²⁹.

Table 1: Acquired and selected intrinsic- antibiotic resistance genes

Antibiotic category	Genes associated with resistance*	Genetic Mechanism	Resistance Phenotype
Penams, Second- and Third- generation Cephalosporins, Cephamycins	<i>bla_{OXA-2}</i> , <i>bla_{OXA-51}</i> , <i>bla_{OXA-58}</i> , <i>bla_{OXA-88}</i> , <i>bla_{OXA-317}</i> , <i>bla_{ADC-25}</i>	Chromosomal; Mobile genetic element- mediated	β-lactamase producing

Fourth-generation Cephalosporins	<i>bla_{ADC-2}</i> , <i>bla_{ADC-25}</i> , <i>bla_{OXA-88}</i>	Chromosomal;	AmpC β-
		Mobile genetic element- mediated	lactamase producing
Carbapenems	<i>bla_{OXA-58}</i> , <i>bla_{OXA-317}</i> , and <i>bla_{OXA-51}</i>	Chromosomal;	Carbapenemase-
		Mobile genetic element-mediated	producing
Aminoglycosides	<i>aph(3'')-Ib</i> , <i>aph(6)-Id</i> , <i>aph(3')-Ia</i> , <i>aac(3)-IId</i> , <i>aac(3)-IIa</i> , <i>aph(3')-I</i> , <i>aph(3'')-I</i> , <i>aph(6)-Ic</i> , <i>aph(6)-Id</i> , <i>aac(3)-II</i> , <i>III</i> , <i>IV</i> , <i>VI</i> , <i>VIII</i> , <i>IX</i> , <i>X</i>	Mobile genetic element- mediated	Aminoglycoside resistance
Macrolides	<i>macA</i> , <i>macB</i>	Chromosomal - mediated	Macrolide resistance
Tetracyclines	<i>tetA</i> , <i>tetB</i> , <i>tetR</i>	Mobile genetic element- mediated	Tetracycline resistance

Trimethoprim	<i>dfrA1</i>	Mobile genetic element- mediated	Trimethoprim resistance
Sulphonamides	<i>sul2</i>	Mobile genetic element- mediated	Sulphonamide resistance
Phenicol	<i>floR</i> family	Mobile genetic element- mediated	Phenicol resistance
Colistin	<i>lpxA, lpxC</i>	Chromosomal – mediated	Colistin resistance

166

167 **Table 1. Antibiotic resistance in *Acinetobacter baumannii* SRR8291681_Uganda.***Several
168 additional intrinsic factors, such as porins and efflux pumps which may be involved in pan-drug
169 resistance are also encoded in the genome.

170 **Figure 2. Graphical circular map of the genome of *Acinetobacter baumannii***
171 **SRR8291681_Uganda**

1

173 **Figure 2. Circular map of the genome.** The two inner black circles indicate the G+C content
 174 plotted against the average G+C content of 38.95%; the green and purple circles show GC
 175 skew information, and the outer circles display the open reading frames (ORFs) in opposite
 176 orientations.

177 Discussion

178 This is the first genomic analysis of a pan-drug resistant *Acinetobacter baumannii* in Uganda.
 179 The definition of this isolate as pan-drug resistant was based on the definition of the Centers

for Disease Control and Prevention (CDC) and the European Centre for Disease Prevention and Control (ECDC) assessment criteria³⁰

Comparable to other Gram-negative bacteria, several intrinsic and acquired mechanisms of antibiotic resistance previously described are present in *Acinetobacter baumannii* SRR8291681_Uganda^{31–34}. These mechanisms can be classified into three broad categories, these include; (i) antibiotic-inactivating enzymes, (ii) reduced access to drug targets of the bacteria, attributable to the decrease in the outer membrane permeability caused by the loss or reduced expression of porins, over-expression of multi-drug efflux pumps, and (iii) mutations that change drug targets or cellular functions (i.e. alterations in the penicillin-binding proteins). These resistance mechanisms explain the evolution of antibiotic resistance of the pan-drug resistance genotype of *Acinetobacter baumannii* SRR8291681_Uganda.

Coupled with the intrinsic antibiotic resistance mechanisms, steps in the evolution of antibiotic resistance of *Acinetobacter baumannii* SRR8291681_Uganda likely followed in a sequential manner that started with the, (i) acquisition of multiple antibiotic resistance genes by the antibiotic-susceptible strain from antibiotic-resistant strains of bacteria through conjugation, transformation, or transduction in horizontal gene transfer^{31–35}; genes that enabled the organism to produce enzymes with the ability to degrade antibiotics, to express antibiotic efflux pumps that prevent antibiotics from reaching their intracellular targets, to modify the antibiotics' target sites, or to trigger the production of alternative metabolic pathways that enable the bypass of action of antibiotics^{31–35}, (ii) integration of the multiple antibiotic resistance genes into the host bacteria's genome or plasmids; and their distribution herein, facilitated by deoxyribonucleic acid transposons and bacterial insertion-sequences^{36,37}, and (iii) consequent vertical gene transfer of the antibiotic resistance genes in clonal spread^{38–40}.

203 In *Acinetobacter baumannii* SRR8291681_Uganda evolution of antibiotic resistance likely
 204 followed from penams to second- and third-generation cephalosporins along with the naturally
 205 occurring resistance to fourth-generation cephalosporins, to cephamycins, carbapenems and
 206 then to the polymyxins; along with resistance to other classes of antibiotics (i.e.
 207 aminoglycosides, tetracyclines, trimethoprim, sulphonamides, and phenicols; along with
 208 intrinsic macrolide resistance).

209 *Acinetobacter*-derived cephalosporinases (ADCs) precisely the extended-spectrum AmpC-
 210 type β -lactamases (ESAC) have been described to possess the ability to hydrolyze extended-
 211 spectrum cephalosporins and monobactams, this has been attributed to a Pro210Arg
 212 substitution and a duplication of an Ala residue at position 215 (inside the O-loop); however,
 213 these have also been described to have no potential of compromising the efficacy of
 214 carbapenems^{41,42}. Furthermore, *Acinetobacter baumannii* has also been described to express
 215 the *bla*_{OXA-51}, another chromosomally-encoded intrinsic β -lactamase for which many point-
 216 mutant variants have been described; these enzymes normally exhibit low-levels of
 217 carbapenemase activity and have the potential of being overproduced especially when their
 218 genes are provided with efficient promoters by insertion sequences (i.e. ISAb1 or ISAb9),
 219 making *Acinetobacter baumannii* consequently resistant to carbapenems^{34,43,44}.

220 Over several decades, carbapenems have been reserved for the treatment of known or
 221 suspected multi-drug resistant bacterial infections including those by *Acinetobacter*
 222 *baumannii*^{6,45}. However, over time, various acquired β -lactamases (i.e. carbapenem-
 223 hydrolyzing class D β -lactamase genes (CHDLs)) have been identified to mediate carbapenem
 224 resistance in *Acinetobacter baumannii*^{6,46}. Despite the ability of these to hydrolyze
 225 carbapenems, they only show but limited ability to hydrolyze extended-spectrum
 226 cephalosporins; their association with extra intrinsic resistance mechanisms, however, has

227 been described to provide a high level of carbapenem resistance⁴¹. *bla*_{OXA-58} and its variants
228 have been documented to occur in worldwide isolates of *Acinetobacter baumannii*⁴⁷. These
229 genes have been described as plasmid-encoded, as well as being associated with specific
230 insertion sequences which play a role in enhancing expression of the same; these have,
231 however, been described to not be part of transposons^{41,48}.

232 Over the years, several aminoglycoside-modifying enzymes including phosphotransferases,
233 acetyltransferases, and adenylyltransferases have been described and documented to occur in
234 *Acinetobacter baumannii*, these mediate antibiotic resistance via reducing or abolishing binding
235 of the aminoglycoside molecule to the ribosome^{41,49}. The genes encoding these enzymes have
236 also been documented to often occur in mobile genetic elements^{41,49}.

237 Tetracycline resistance has been documented to occur when reversible binding of tetracycline
238 to the 30S ribosomal subunit occurs and as a result, inhibiting protein translation^{41,49,50}. In
239 *Acinetobacter baumannii*, the *tetA* and *tetB* have been described as specific efflux pump
240 proteins that actively remove tetracycline from the cells^{41,49,50}.

241 The main mechanisms of resistance to polymyxins and colistin antibiotics involve the; (i)
242 reduction of the net negative of the outer-membrane protein by modification of lipid A, an
243 essential component of the bacterial lipopolysaccharide (LPS); and (ii) proteolytic cleavage of
244 the antibiotic compound and exclusion of peptides by a broad spectrum efflux-pump⁵¹. In
245 *Acinetobacter baumannii*, antibiotic resistance gene variants or mutants *lpxA* and *lpxC* that
246 mediate polymyxin and colistin resistance via molecular bypass have been documented^{8,9}.

247 Despite the significance of pan-drug resistance phenotypic data in defining true pan-drug
248 resistance in bacterial isolates⁵⁴, the current study was unable to obtain this data for

249 *Acinetobacter baumannii* SRR8291681_Uganda. The study therefore described the isolate as
250 pan-drug resistant as defined by CDC and ECDC basing on genomic analysis.

251 In addition, genomic analysis alone was utilized to describe mobile genetic element-mediated
252 antibiotic resistance evolution in *Acinetobacter baumannii* SRR8291681_Uganda. This was
253 informed by a similar study that determined the genetic basis and evolution of pan-drug
254 resistance in bacteria by interrogating the genome to identify acquired as well as intrinsic
255 antibiotic resistance genes⁵⁴.

256 The inability to compare the differential evolution of antimicrobial resistance in this study was
257 to the most part attributed to the fact that the only available *Acinetobacter baumannii* genomes
258 from Africa particularly Tanzania that were added as part of the analysis had earlier been
259 described as extended-spectrum β -lactamase encoding and had been confirmed as such post-
260 sequencing. These genomes therefore could have been inadequate to compare with those of
261 the carbapenem and colistin resistant *Acinetobacter baumannii* SRR8291681_Uganda.

262 **Conclusion**

263 Although limited by the absence of pan-drug resistance phenotypic data necessary in defining
264 bacteria as truly pan-drug resistant this study provided the first genomic analysis of a pan-drug
265 resistant *Acinetobacter baumannii* from Uganda and Africa and described mobile genetic
266 element-mediated antibiotic resistance in the same isolate. The study provides a basis to track
267 trends in antibiotic resistance as well as identification of emerging resistance patterns in
268 *Acinetobacter baumannii* in Uganda. Lastly, the study recommends that continued surveillance
269 using both phenotypic and genotype data be done continuously to identify country-specific data
270 on antibiotics effective for antibiotic resistant bacteria.

271 **Abbreviations**

272 WHO: World Health Organization; MLST: Multilocus sequence typing; ST: Sequence typing;
 273 ADCs: Acinetobacter-derived cephalosporinases; CHDLs: Carbapenem-hydrolyzing class D β -
 274 lactamase genes; MFS: Major facilitator superfamily; IS: Insertion sequence; CDC: Centres for
 275 Disease Control and Prevention; ECDC: European Centre for Disease Prevention and Control;
 276 ESAC: Extended-spectrum AmpC-type β -lactamases; PATRIC: Pathosystems Resource
 277 Integration Center; CARD: Comprehensive Antimicrobial Resistance Database; NDARO:
 278 National Database of Antibiotic Resistant Organisms; JCVI: J. Craig Venter Institute; CC:
 279 Clonal complex; GC: Global complex

280 **Acknowledgments**

281 Not applicable.

282 **Author contributions**

283 Conceptualization: DA, GM. Data Curation: DA, GM, IS. Formal analysis: DA, GM, IS.
 284 Methodology: DA, GM, IS. Validation: DA, GM, IS. Writing- Original draft preparation: DA, GM,
 285 IS. Writing- Review and Editing: DA, GM, IS. All authors read and approved the final
 286 manuscript.

287 **Funding**

288 Gerald Mboowa is supported through the DELTAS Africa Initiative (Grant no. DEL15011) to
 289 THRiVE-2 (Training Health Researchers into Vocational Excellence in East Africa- 2). The
 290 DELTAS Africa Initiative is an independent funding scheme of the African Academy of
 291 Sciences' (AAS) Alliance for Accelerating Excellence in Science in Africa (AESA) supported by
 292 the New Partnership for Africa's Development Planning and Coordinating Agency (NEPAD
 293 Agency) with funding from the Wellcome Trust (Grant no. 107742/Z/15/Z) and the UK
 294 government.

295 GM also is supported through the Grand Challenges Africa programme [GCA/AMR/rnd2/058].
 296 Grand Challenges Africa is a programme of the African Academy of Sciences (AAS)
 297 implemented through the Alliance for Accelerating Excellence in Science in Africa (AESA)
 298 platform, an initiative of the AAS and the African Union Development Agency (AUDA-NEPAD).
 299 GC Africa is supported by the Bill & Melinda Gates Foundation (BMGF) and The African
 300 Academy of Sciences and partners. The views expressed herein are those of the author(s) and
 301 not necessarily those of the AAS and her partners.

302 The funders had no role in study design, data collection and analysis, decision to publish, or
 303 preparation of the manuscript.

304 **Availability of data and materials**

305 The datasets used and/or analysed during the current study are available from the
 306 corresponding author on reasonable request. Genomic sequence data analyzed here is freely
 307 available on NCBI and is deposited under the BioProject ID PRJNA508495
 308 (<https://www.ncbi.nlm.nih.gov/sra/SRR8291681/>).

309 **Ethical approval and consent to participate**

310 Not applicable.

311 **Consent for publication**

312 Not applicable.

313 **Competing interests**

314 The authors declare that they have no competing interests.

315 **Author details**

316 ¹Department of Medical Microbiology, College of Health Sciences, School of Biomedical
317 Sciences, Makerere University, Kampala, 7072, Uganda

318 ²Department of Immunology and Molecular Biology, College of Health Sciences, School of
319 Biomedical Sciences, Makerere University, Kampala, 7072, Uganda

320 ³The African Center of Excellence in Bioinformatics and Data Intensive Sciences, the Infectious
321 Disease Institute, Makerere University P.O. Box 22418, Kampala, Uganda

322 **References**

323

- 324 1. World Health Organisation. list of bacteria for which new antibiotics are urgently needed.
325 2017
- 326 2. Howard, A., O'Donoghue, M., Feeney, A., & Sleator, R. D. *Acinetobacter baumannii*: an
327 emerging opportunistic pathogen. *Virulence* **3**, 243–250 (2012).
- 328 3. Huang, G. *et al.* Multilocus Sequence Typing Analysis of Carbapenem-Resistant
329 *Acinetobacter baumannii* in a Chinese Burns Institute . *Frontiers in Microbiology* **7**,
330 1717 (2016).
- 331 4. Huang, Z.-Y. *et al.* Co-existence of blaOXA-23 and blaVIM in carbapenem-resistant
332 *Acinetobacter baumannii* isolates belonging to global complex 2 in a Chinese teaching
333 hospital. *Chin. Med. J. (Engl)*. **132**, (2019).
- 334 5. Zhao, W.-H. & Hu, Z.-Q. *Acinetobacter*: a potential reservoir and dispenser for β -
335 lactamases. *Crit. Rev. Microbiol.* **38**, 30–51 (2012).
- 336 6. Kateete, D. P. *et al.* Carbapenem resistant *Pseudomonas aeruginosa* and *Acinetobacter*
18

- 337 baumannii at Mulago Hospital in Kampala, Uganda (2007–2009). *Springerplus* **5**, 1308
338 (2016).
- 339 7. Camargo, C. H. *et al.* Population structure analysis of carbapenem-resistant
340 Acinetobacter baumannii clinical isolates from Brazil reveals predominance of clonal
341 complexes 1, 15, and 79. *Antimicrob. Agents Chemother.* **60**, 2545–2547 (2016).
- 342 8. Carretero-Ledesma, M. *et al.* Phenotypic changes associated with Colistin resistance
343 due to Lipopolysaccharide loss in Acinetobacter baumannii. *Virulence* **9**, 930–942
344 (2018).
- 345 9. Zhang, W. *et al.* The role of LpxA/C/D and pmrA/B gene systems in colistin-resistant
346 clinical strains of Acinetobacter baumannii. *Front. Lab. Med.* **1**, 86–91 (2017).
- 347 10. Coyne, S., Courvalin, P. & Périchon, B. Efflux-mediated antibiotic resistance in
348 Acinetobacter spp. *Antimicrob. Agents Chemother.* **55**, 947–953 (2011).
- 349 11. Møller, T. S. B. *et al.* Relation between tetR and tetA expression in tetracycline resistant
350 Escherichia coli. *BMC Microbiol.* **16**, 39 (2016).
- 351 12. Cuthbertson, L. & Nodwell, J. R. The TetR family of regulators. *Microbiol. Mol. Biol. Rev.*
352 **77**, 440–475 (2013).
- 353 13. Shaw, K. J., Rather, P. N., Hare, R. S. & Miller, G. H. Molecular genetics of
354 aminoglycoside resistance genes and familial relationships of the aminoglycoside-
355 modifying enzymes. *Microbiol. Mol. Biol. Rev.* **57**, 138–163 (1993).
- 356 14. Ramirez, M. S. & Tolmasky, M. E. Aminoglycoside modifying enzymes. *Drug Resist.*
357 *Updat.* **13**, 151–171 (2010).
- 358 15. Okada, U. *et al.* Crystal structure of tripartite-type ABC transporter MacB from
19

Acinetobacter baumannii. *Nat. Commun.* **8**, 1336 (2017).

16. Anitha, P., Anbarasu, A. & Ramaiah, S. Computational gene network study on antibiotic resistance genes of *Acinetobacter baumannii*. *Comput. Biol. Med.* **48**, 17–27 (2014).

17. Chang-Tai, Z. *et al.* High frequency of integrons related to drug-resistance in clinical isolates of *Acinetobacter baumannii*. *Indian J. Med. Microbiol.* **29**, 118 (2011).

18. Leungtongkam, U., Thummeepak, R., Tasanapak, K. & Sitthisak, S. Acquisition and transfer of antibiotic resistance genes in association with conjugative plasmid or class 1 integrons of *Acinetobacter baumannii*. *PLoS One* **13**, e0208468 (2018).

19. Si-Tuan, N. *et al.* New eight genes identified at the clinical multidrug-resistant *Acinetobacter baumannii* DMS06669 strain in a Vietnam hospital. *Ann. Clin. Microbiol. Antimicrob.* **16**, 74 (2017).

20. Taitt, C. R. *et al.* Antimicrobial resistance determinants in *Acinetobacter baumannii* isolates taken from military treatment facilities. *Antimicrob. Agents Chemother.* **58**, 767–781 (2014).

21. Sakai, H. & Komano, T. DNA replication of IncQ broad-host-range plasmids in gram-negative bacteria. *Biosci. Biotechnol. Biochem.* **60**, 377–382 (1996).

22. Jain, A. & Srivastava, P. Broad host range plasmids. *FEMS Microbiol. Lett.* **348**, 87–96 (2013).

23. von Wintersdorff, C. J. H. *et al.* Dissemination of antimicrobial resistance in microbial ecosystems through horizontal gene transfer. *Front. Microbiol.* **7**, 173 (2016).

24. Bellanger, X., Payot, S., Leblond-Bourget, N. & Guédon, G. Conjugative and mobilizable genomic islands in bacteria: evolution and diversity. *FEMS Microbiol. Rev.* **38**, 720–760

381 (2014).

382 25. Bertini, A. *et al.* Characterization and PCR-based replicon typing of resistance plasmids
383 in *Acinetobacter baumannii*. *Antimicrob. Agents Chemother.* **54**, 4168–4177 (2010).

384 26. Liu, C.-C., Kuo, H.-Y., Tang, C. Y., Chang, K.-C. & Liou, M.-L. Prevalence and mapping
385 of a plasmid encoding a type IV secretion system in *Acinetobacter baumannii*. *Genomics*
386 **104**, 215–223 (2014).

387 27. Bignell, C. & Thomas, C. M. The bacterial ParA-ParB partitioning proteins. *J. Biotechnol.*
388 **91**, 1–34 (2001).

389 28. Hayes, F. The partition system of multidrug resistance plasmid TP228 includes a novel
390 protein that epitomizes an evolutionarily distinct subgroup of the ParA superfamily. *Mol.*
391 *Microbiol.* **37**, 528–541 (2000).

392 29. Vijayakumar, S., Kanthan, K. & Veeraraghavan, B. Whole-genome shotgun sequences
393 of seven colistin-resistant *Acinetobacter baumannii* isolates from bacteraemia. *J. Glob.*
394 *Antimicrob. Resist.* **12**, 155–156 (2018).

395 30. Magiorakos, A. *et al.* Multidrug-resistant, extensively drug-resistant and pandrug-
396 resistant bacteria: an international expert proposal for interim standard definitions for
397 acquired resistance. *Clin. Microbiol. Infect.* **18**, 268–281 (2012).

398 31. Hancock, R. E. W. Resistance mechanisms in *Pseudomonas aeruginosa* and other
399 nonfermentative gram-negative bacteria. *Clin. Infect. Dis.* **27**, S93–S99 (1998).

400 32. Tenover, F. C. Mechanisms of antimicrobial resistance in bacteria. *Am. J. Med.* **119**, S3–
401 S10 (2006).

402 33. Pfeifer, Y., Cullik, A. & Witte, W. Resistance to cephalosporins and carbapenems in
21

- 403 Gram-negative bacterial pathogens. *Int. J. Med. Microbiol.* **300**, 371–379 (2010).
- 404 34. Poirel, L. & Nordmann, P. Carbapenem resistance in *Acinetobacter baumannii*:
405 mechanisms and epidemiology. *Clin. Microbiol. Infect.* **12**, 826–836 (2006).
- 406 35. Moore, R. A. & Hancock, R. E. Involvement of outer membrane of *Pseudomonas cepacia*
407 in aminoglycoside and polymyxin resistance. *Antimicrob Agents Chemother* **30**, (1986).
- 408 36. Muñoz-López, M. & García-Pérez, J. L. DNA transposons: nature and applications in
409 genomics. *Curr. Genomics* **11**, 115–128 (2010).
- 410 37. Siguier, P., Gournay, E. & Chandler, M. Bacterial insertion sequences: their genomic
411 impact and diversity. *FEMS Microbiol. Rev.* **38**, 865–891 (2014).
- 412 38. Spratt, B. G. Exploring the concept of clonality in bacteria. in *Genomics, Proteomics, and*
413 *Clinical Bacteriology* 323–352 (Springer, 2004).
- 414 39. Poirel, L. *et al.* Nosocomial spread of OXA-58-positive carbapenem-resistant
415 *Acinetobacter baumannii* isolates in a paediatric hospital in Greece. *Clin. Microbiol.*
416 *Infect.* **12**, 1138–1141 (2006).
- 417 40. Durante-Mangoni, E. & Zarrilli, R. Global spread of drug-resistant *Acinetobacter*
418 *baumannii*: molecular epidemiology and management of antimicrobial resistance. *Future*
419 *Microbiol.* **6**, 407–422 (2011).
- 420 41. Poirel, L., Bonnin, R. A. & Nordmann, P. Genetic basis of antibiotic resistance in
421 pathogenic *Acinetobacter* species. *IUBMB Life* **63**, 1061–1067 (2011).
- 422 42. Rodríguez-Martínez, J.-M., Nordmann, P., Ronco, E. & Poirel, L. Extended-spectrum
423 cephalosporinase in *Acinetobacter baumannii*. *Antimicrob. Agents Chemother.* **54**,
424 3484–3488 (2010).

- 425 43. Turton, J. F. *et al.* The role of ISAb₁ in expression of OXA carbapenemase genes in
426 *Acinetobacter baumannii*. *FEMS Microbiol. Lett.* **258**, 72–77 (2006).
- 427 44. Figueiredo, S., Poirel, L., Papa, A., Koulourida, V., and Nordmann, P. Overexpression of
428 the naturally occurring blaOXA-51 gene in *Acinetobacter baumannii* mediated by novel
429 insertion sequence ISAb₉. *Antimicrob. Agents Chemother.* **53**, 4045–4047 (2009).
- 430 45. Fishbain J, P. A. Treatment of *Acinetobacter* infections. *Clin. Infect. Dis.* **51**, 79–84
431 (2010).
- 432 46. Poirel L, Cabanne L, Vahaboglu H, N. P. Genetic environment and expression of the
433 extended-spectrum β -lactamase blaPER-1 gene in gram-negative bacteria. *Antimicrob.*
434 *Agents Chemother.* **49**, 1708–13 (2005).
- 435 47. Weiyuan Wu , Yi He, Jian Lu, Yuemei Lu, Jinsong Wu, Y. L. Transition of blaOXA-58-
436 like to blaOXA-23-like in *Acinetobacter baumannii* Clinical Isolates in Southern China: An
437 8-Year Study. *PLoS One* **10(9)**, (2015).
- 438 48. Poirel, L., and Nordmann, P. Genetic structures at the origin of acquisition and
439 expression of the carbapenem-hydrolyzing oxacillinase gene blaOXA-58 in
440 *Acinetobacter baumannii*. *Antimicrob. Agents Chemother* **50**, 1442–1448 (2006).
- 441 49. Aliakbarzade K, Farajnia S, Nik AK, Zarei F, T. A. Prevalence of aminoglycoside
442 resistance genes in *Acinetobacter baumannii* isolates. *Jundishapur J. Microbiol.* **7**,
443 (2014).
- 444 50. Speer BS, Shoemaker NB, S. A. Bacterial resistance to tetracycline: mechanisms,
445 transfer, and clinical significance. *Clin. Microbiol. Rev.* **5**, 387–99 (1992).
- 446 51. Aruhomukama D, S. I. and M. G. Investigating colistin drug resistance: The role of high-

447 throughput sequencing and bioinformatics. *F1000Research* **8**, (2019).

448 52. Wick, R. R., Judd, L. M., Gorrie, C. L. & Holt, K. E. Unicycler: resolving bacterial genome
449 assemblies from short and long sequencing reads. *PLoS Comput. Biol.* **13**, e1005595
450 (2017).

451 53. Bankevich, A. *et al.* SPAdes: a new genome assembly algorithm and its applications to
452 single-cell sequencing. *J. Comput. Biol.* **19**, 455–477 (2012).

453 54. Zowawi, H. M. *et al.* Stepwise evolution of pandrug-resistance in *Klebsiella pneumoniae*.
454 *Sci. Rep.* **5**, 15082 (2015).

455

456