Isolation and characterization of Streptomyces bacteriophages and Streptomyces strains encoding biosynthetic arsenals

Streptomyces strains and phages for antibiotic discovery

Elizabeth T. Montaña, Jason F. Nideffer, Lauren Brumage, Marcella Erb, Alan I. Derman, John Paul Davis, Elena Estrada, Sharon Fu, Danielle Le, Aishwarya Vuppala, Cassidy Tran, Elaine Luterstein, Shivani Lakkaraju, Sriya Panchagnula, Caroline Ren, Jennifer Doan, Sharon Tran, Jamielyn Soriano, Yuya Fujita, Pranathi Gutala, Quinn Fujii, Minda Lee, Anthony Bui, Carleen Villarreal, Samuel R. Shing, Sean Kim, Danielle Freeman, Vipula Racha, Alicia Ho, Prianka Kumar, Kian Falah, Thomas Dawson, Eray Enustun, Amy Prichard, Ana Gomez, Kanika Khanna, Shelly Trigg, Kit Pogliano, Joe Pogliano*

Division of Biological Sciences, University of California, San Diego, La Jolla, California, USA.

* Corresponding author: jpogliano@ucsd.edu

Abstract

The threat to public health posed by drug-resistant bacteria is rapidly increasing, as some of healthcare's most potent antibiotics are becoming obsolete. Approximately two-thirds of the world's antibiotics are derived from natural products produced by *Streptomyces* encoded biosynthetic gene clusters. Thus, in order to identify novel gene clusters, we sequenced the genomes of four bioactive *Streptomyces* strains isolated from the soil in San Diego County and used Bacterial Cytological Profiling adapted for agar plate culturing in order to examine the mechanisms of bacterial inhibition exhibited by these strains. In the four strains, we identified 101 biosynthetic gene clusters. Some of these clusters were predicted to produce previously studied antibiotics; however, the known mechanisms of these molecules could not fully account for the antibacterial activity exhibited by the strains, suggesting that novel clusters might encode antibiotics. When assessed for their ability to inhibit the growth of clinically isolated pathogens, three *Streptomyces* strains demonstrated activity against methicillin-resistant *Staphylococcus aureus*. Additionally, due to the utility of bacteriophages for genetically manipulating bacterial strains via transduction, we also isolated four new phages (BartholomewSD, IceWarrior, Shawty, and TrvxScott) against *S. platensis*. A genomic analysis of our phages revealed nearly 200 uncharacterized proteins, including a new site-specific serine integrase that could prove to be a useful genetic tool. Sequence analysis of the *Streptomyces* strains identified CRISPR-Cas systems and specific spacer sequences that allowed us to predict phage host ranges. Ultimately, this study identified *Streptomyces* strains with the potential to produce novel chemical matter as well as integrase-encoding phages that could potentially be used to manipulate these strains.

Introduction

Antibiotic discovery is an international priority requiring immediate action.¹ The increasing prevalence of multi-drug resistant (MDR) bacterial pathogens has resulted in an increased use of last-resort antibiotics.¹⁻³ Microbes that produce natural products are the most prolific source of clinically approved antibiotics.⁴ In particular, soil dwelling Actinobacteria, notably Streptomyces, account for two-thirds of the antibiotics currently on the market.⁵⁻⁷ Despite intensive studies, however, the full potential of microbes to produce natural products has not been realized.⁸ Genome mining studies have shown that microbes encode many biosynthetic gene clusters (BGCs) that have not yet been characterized.⁸ It is widely assumed that many of these clusters produce novel natural products and that further characterization of Streptomyces bacteria increases the probability of identifying molecules with unique chemical structures and new mechanisms of action.⁹

In addition to identifying Streptomyces strains containing potentially novel BGCs, it is necessary to improve on the conventional approaches used in natural product antibiotic discovery. One of the major stumbling blocks in natural product discovery is dereplication, since the isolation of bioactive molecules often yields antibiotics that have previously been discovered.¹⁰ We recently developed Bacterial Cytological Profiling (BCP) as a new whole-cell screening technique that can be used to rapidly identify the mechanism of action (MOA) of antibiotics.¹¹⁻¹⁶ BCP can accurately identify the pathway inhibited by antibacterial compounds present in unfractionated crude organic extracts and can be used to guide the purification of molecules with specific bioactivities.^{11, 15} BCP can also be used to screen bacterial strains directly on petri plates in order to identify and prioritize those strains that produce molecules with desired MOA.¹⁵ In effect, BCP helps with the problem of dereplication by allowing for the determination of the MOA of

antibiotics synthesized by a particular *Streptomyces* strain before labor-intensive antibiotic purification efforts are performed.

Since many BGCs are not expressed under laboratory conditions, genetic methods are often used to augment their expression and facilitate the identification and purification of their products.¹⁷ Sometimes, increased expression can be achieved using techniques such as CRISPR/Cas or plasmid cloning and overexpression.¹⁷ However, there is still an occasional need to move large chromosomal regions from one strain to another via transduction in order to engineer strains optimally suited for antibiotic production. Transduction requires a phage capable of infecting the strain(s) of interest. Moreover, because phages generally display narrow host ranges¹⁸ and relatively few *Streptomyces* phages have been isolated¹⁹ compared to the large number of studied *Streptomyces* bacteria,²⁰ phages aptly suited for genetic manipulations are not available for the majority of antibiotic producing *Streptomyces* strains isolated. In addition, phage derived enzymes such as recombinases and integrases can also be used to engineer new strains.²¹⁻²⁵ Thus, studying the phages that infect antibiotic-producing *Streptomyces* strains could not only yield new transducing phages but potentially also new genetic tools for strain engineering.

Here we describe the isolation and characterization of *Streptomyces* strains and phages. We used a combination of bioinformatics and BCP to characterize the antibiotic biosynthetic potential of four *Streptomyces* strains that displayed an ability to inhibit Gram-negative and Gram-positive growth. Additionally, we isolated four new phages and assessed their abilities to infect our *Streptomyces* strains, which contained many CRISPRs. The proteins encoded by the phages were subjected to bioinformatic analyses in order to identify putative integrases that might be used for genetic manipulations. This work highlights a novel set of gene clusters and

Streptomyces sp. phages that serve as a starting point for the isolation of potentially novel natural products.

Results and Discussion

Isolation and Antibacterial Activities of Streptomyces sp.

To identify Streptomyces strains containing potentially novel BGCs, we collected 28 unique soil samples from sites across San Diego County. From these samples, we isolated a total of eight bacterial strains that were determined, based on 16s RNA sequencing, to be members of the Streptomyces genus (Fig 1, Table 1). These strains (designated JS, DF, QF2, EDE, SK, AH, ELW, and SFW) and two known species (*Streptomyces coelicolor* A3(2) and *Streptomyces platensis* AB045882) were screened using the cross-streak method for their ability to inhibit the growth of wild type *E. coli* MC4100, an efflux defective mutant *E. coli* JP313 $\Delta tolC$, and *B. subtilis* PY79. Since the production of bioactive secondary metabolites is highly dependent on growth conditions,²⁶ this screen was conducted on actinomycete isolation agar (AIA) as well as LB agar. Each of the 10 strains proved capable of inhibiting the growth of *E. coli* and/or *B. subtilis* on at least one of the tested media (Fig 2), suggesting that these strains likely produce antibiotics. As expected, however, the production of antibiotics often depended upon whether the strain was grown on AIA or LB agar. For example, strain ELW was incapable of inhibiting the growth of Gram-negative and Gram-positive bacteria when grown on AIA. However, when grown on LB agar, strain ELW inhibited the growth of both Gram-negative and Gram-positive bacteria. Conversely, strains JS and QF2 exhibited growth inhibition regardless of the media on which they were grown.

Fig 1. Phylogenetic tree of Streptomyces bacteria isolated from soil samples. This tree was constructed using PCR-amplified 16S rRNA sequences and was generated from a multiple sequence alignment using a neighbor-joining algorithm. The scale bar (0.03) indicates substitutions per nucleotide.

Table 1. Top NCBI BlastN hits of the 16S rRNA gene sequences.

Sample ID	NCBI BlastN Top 16S ribosomal RNA Hit Description	Max Score	Total Score	Query Cover	E value	Percent Identity	Accession No.
S. platensis JCM 4662	Streptomyces platensis strain JCM 4662	2748	2748	100%	0	100%	NR_024761.1
S. coelicolor A3(2)	Streptomyces coelestis strain AS 4.1594	2793	2793	98%	0	99.93%	NR_027222.1
JS	Streptomyces rochei strain NRRL B-1559	2741	2741	91%	0	99.93%	NR_116078.1
DF	Streptomyces fulvissimus strain DSM 40593	2691	2691	100%	0	99.93%	NR_103947.1
QF2	Streptomyces californicus strain NBRC 12750	2699	2699	100%	0	100%	NR_112257.1
EDE	Streptomyces pratensis strain ch24	1238	1238	99%	0	98.99%	NR_125616.1
SK	Streptomyces californicus strain NBRC 12750	1242	1242	100%	0	100%	NR_112257.1
AH	Streptomyces pratensis strain ch24	1194	1194	100%	0	100%	NR_125616.1
ELW	Streptomyces atratus strain NRRL B-16927	1138	1138	100%	0	99.84%	NR_043490.1
SFW	Streptomyces caviscabies strain ATCC 51928	2767	2767	100%	0	99.87%	NR_114493.1

Fig 2. Inhibition of bacterial growth by Streptomyces isolates. The cross-streak method was used to measure the zone of inhibition among ten Streptomyces strains against two Gram-negative *E. coli* strains (MC4100 WT, JP313 $\Delta toIC$), and Gram-positive *B. subtilis* PY79 on (a) LB and (b) actinomycete isolation agar (AIA). Error bars represent the standard deviation of three independent trials.

Mechanistic Analysis of Natural Products Produced by Four Streptomyces Isolates

Strains QF2, JS, SFW and DF all inhibited the growth of *E. coli* $\Delta toIC$ when grown on either AIA or LB agar, but in each case, the mechanism underlying inhibition was unknown. Thus, we utilized BCP to examine the mechanism of growth inhibition exhibited by the antibacterial natural products synthesized by these four Streptomyces isolates. Each of the four strains was grown on three different media (LB, AIA, or ISP2) for 5 days to allow for the synthesis and excretion of natural products into the surrounding agar. We then added exponentially growing *E. coli* cells adjacent to the Streptomyces lawn. After two hours of incubation at 30°C, the *E. coli* cells were stained with fluorescent dyes and imaged with high resolution fluorescence microscopy. *E. coli* cells incubated adjacent to each of the four Streptomyces isolates displayed characteristic cytological profiles that, in some cases, allowed for the classification of these strains according to the MOA of the natural products they produced (Fig 3).

Fig 3. BCP phenotypes of *E. coli* JP313 $\Delta tolC$ exposed to natural products produced by four *Streptomyces* soil isolates grown on different solid media. Also displayed, are *E. coli* JP313 $\Delta tolC$ untreated controls grown on the tested media (LB agar, AIA, and ISP2 agar). White arrows indicate cells with three chromosomes. BCP images were collected after staining the cells with FM4-64 (red), DAPI (blue), and SYTOX-green (green). The scale bar represents one micron.

When grown on either LB or ISP2, strain QF2 synthesized an antibiotic that caused the DNA of affected *E. coli* cells to assume a toroidal conformation (Fig 3). This phenotype is characteristic of bacteria treated with protein synthesis inhibitors such as chloramphenicol,^{11, 27} and thus, we concluded that strain QF2 is capable of synthesizing a translation-inhibiting natural product. QF2 also produced a membrane-active secondary metabolite, evidenced by visible membrane abnormalities as well as Sytox Green permeability under all tested nutrient conditions (Fig 3). Strain JS appeared to induce similar phenotypes in *E. coli*, though under different growth conditions; protein synthesis inhibition phenotypes were observed on AIA and ISP2 but not on LB. And, as was the case with strain QF2, Sytox Green permeability was observed in some cells regardless of the medium on which strain JS was grown.

Strain SFW induced distinct phenotypes in *E. coli* cells under each of the three nutrient conditions (Fig 3). On LB, a significant number of *E. coli* cells grown in the presence of strain SFW appeared to contain three chromosomes (white arrows), a phenotype that was not present in the untreated control cells. When strain SFW was grown on AIA, the *E. coli* cells became bent and lost their characteristic rod shape. Finally, strain SFW grown on ISP2 induced substantial swelling in *E. coli* cells that ultimately led to lysis. Notably, *E. coli* cells grown in the presence of strain DF exhibited nearly these same phenotypes under identical growth conditions, suggesting that these two strains produce compounds targeting similar pathways.

Genomic Analysis of Four Streptomyces Isolates

In order to better understand how strains QF2, JS, SFW and DF inhibited bacterial growth, we sequenced their genomes and aligned them to the most similar genomes in the NCBI database (Fig 4A). Sequence reads for strain DF were assembled into a single contig that was most similar to the genome of *S. fulvissimus* DSM 40593. Sequencing of strains QF2, JS, and SFW yielded multiple contigs that were aligned to the genomes of *S. globisporus* C-1027, *S. parvulus* 12434, *S. pratensis* ATCC 33331, respectively.

Fig 4. Genome characteristics of Streptomyces strains DF, SFW, QF2, and JS isolated from soil samples. (a) Circularized representations of the linear genomes of the four bacterial isolates represented by the assembled contigs obtained from genome sequencing. (b) Genomes oriented according to their threonine operons (dark blue). Predicted biosynthetic gene clusters (light blue), loci of Cas-associated protein-coding genes (green), and CRISPR arrays (purple) are shown. (c) A Venn diagram displaying the numbers of BGCs that are shared by and unique to the genomes of our isolates. Five clusters of particular importance are explicitly named.

In order to identify predicted gene clusters associated with secondary metabolism, the assembled genome sequences for strains QF2, JS, SFW and DF were annotated using RASTtk²⁸ and submitted to AntiSMASH 5.0²⁹ (Fig 4B). Each strain encoded between 18 and 34 BGCs, some of which were present in multiple strains (Fig 4C). Additionally, some of the encoded clusters closely resembled known BGCs in the MIBig repository.³⁰ For example, of the 23 putative BGCs identified in the genome of strain QF2 (Table 2), one of them (cluster 21) was similar to the viomycin BGC (Fig 5A). Viomycin inhibits protein synthesis by stabilizing tRNAs in the A site of the bacterial ribosome, inhibiting translocation.³¹ According to AntiSMASH, 66% of the genes within the viomycin BGC were similar to genes within cluster 21. However, a global pairwise alignment of cluster 21 and the viomycin BGC revealed that the nucleotide sequence of cluster 21 is actually 98.5% identical over 32.5kb of the 36kb viomycin BGC (Fig 5A). This suggests that a viomycin related molecule is synthesized by strain QF2 and may account for

strain's ability to inhibit protein synthesis in *E. coli* (Fig 3). While some of the other clusters in the QF2 genome (Table 2: clusters 2, 4, 7, 8, 9, 12, 13, 22) have significant similarity to known BGCs, no other clusters appear to produce known antibiotics.

Table 2. BGCs encoded within the draft genome sequence of strain QF2.

Strain - QF2							
Cluster	Type	Most Similar MiBIG Cluster and Predicted Percent Similarity	Antibacterial Activity	MiBIG BGC-ID	Minimum	Maximum	Length (nt.)
1	butyrolactone	Coelimycin P1 T1PKS (8%)		BGC0000038	28795	39496	10702
2	terpene	Geosmin Terpene (100%)		BGC0001181	60346	82527	22182
3	NRPS	Griseobactin NRPS (35%)		BGC0000368	101394	123548	22155
4	NRPS	Coelichelin NRPS (72%)		BGC0000325	123549	145157	21609
5	T3PKS	Herboxidiene T1PKS, T3PKS (6%)		BGC0001065	184513	202853	18341
6	terpene	Isorenieratene Terpene (57%)		BGC0000664	681001	697369	16369
7	ectoine	Ectoine Other (75%)		BGC0000853	1125565	1134445	8881
8	T2PKS	Griseorhodin T2PKS (69%)		BGC0000230	1908342	1950906	42565
9	siderophore	Desferrioxamine B Siderophore (80%)		BGC0000941	2278919	2290697	11779
10	LAP,thiopeptide	-	-	-	2690041	2722548	32508
11	ectoine, butyrolactone	Showdomycin Other (47%)	Nucleic Acid and Protein Synthesis	BGC0001778	3344033	3359401	15369
12	melanin	Melanin Other (100%)		BGC0000911	4777563	4787988	10426
13	lanthipeptide	AmfS Lanthipeptide (100%)		BGC0000496	5105643	5127766	22124
14	terpene	-	-	-	5476141	5497007	20867
15	siderophore	Ficellomycin NRPS (3%)	DNA Replication	BGC0001593	5881146	5896034	14889
16	NRPS	Vioprolide A NRPS (25%)		BGC0001822	6093895	6137611	43717
17	bacteriocin	-	-	-	6194201	6205608	11408
18	NRPS-like	-	-	-	6430967	6442494	11528
19	NRPS-like, ladderane, arylpolyene	Skyllamycin NRPS (14%)	Unknown MOA	BGC0000429	6608620	6645073	36454
20	terpene	Hopene Terpene (46%)		BGC0000663	6755130	6764031	8902
21	NRPS, T1PKS	Viomycin NRPS (66%)	Protein Synthesis	BGC0000458	6806327	6870853	64527
22	T3PKS	Alkylresorcinol T3PKS (66%)	Unknown MOA	BGC0000282	7058484	7080991	22508
23	lassopeptide	-	-	-	7260793	7282889	22097

The most similar BGCs in the MiBIG database are listed, as well as the percentage of genes in each MiBIG known cluster that have similarity to genes in the corresponding QF2 cluster. In cases where the most similar known BGC produces an antibiotic, the MOA was listed (Showdomycin,³² Ficellomycin,³³ Skyllamycin,³⁴ Viomycin,³¹ Alkylresorcinol³⁵).

Fig 5. Comparison of BGCs encoded in the genomes of bacterial soil isolates and the predicted most similar, previously characterized BGC with an antibacterial product. (a) Strain QF2, BGC 21, compared to the BGC previously described to produce the antibiotic viomycin (NCBI Acc No. AY263398.1), encoded in the WGS of *S. vinaceus* ATCC 11861. (b) Strain JS, BGC 7, compared to the BGC previously described to produce the antibacterial sesquiterpene Albaflavenone (NCBI Acc No. AL645882.2), encoded in the WGS of *S. coelicolor* A3(2). (c) Strain SFW, BGC 1, compared to the BGC previously described to produce the antibacterial beta-lactam Carbapenem MM 4550 (NCBI Acc No. KF042303.1), encoded in the WGS of *S. argenteolus* ATCC 11009. (d) Strain DF, BGC 27, compared to the BGC previously described to produce the ammonium ionophore antibiotic Nonactin (NCBI Acc No. AF074603.2), encoded in the WGS of *S. griseus subsp. griseus* ETH A7796. Cluster comparisons were constructed in Easyfig. Regions of nucleotide homology are indicated on a gray scale and genes are colored according to the putative function of the corresponding protein product.

Strain JS contained 18 putative BGCs, six of which shared significant similarity (>60% of genes in common) with a known cluster (Table 3). Of these six, however, only cluster 7 was predicted

to produce an antibiotic. All of the genes constituting a known terpene cluster that produces albaflavenone were present in cluster 7 (Fig 5B). Albaflavenone is capable of inhibiting the growth of *B. subtilis* by an unknown MOA³⁶ and has previously been isolated from *S. coelicolor* A3(2),³⁷ a close relative of strain JS. Since the MOA of albaflavenone is unknown, it's not clear whether the products of cluster 7 or of a different cluster are responsible for the inhibition of protein synthesis and/or the membrane permeability observed in *E. coli* (Fig 3).

Table 3. BGCs encoded within the draft genome sequence of strain JS.

Strain - JS									
Cluster	Type	Most Similar MiBIG Cluster and Predicted Percent Similarity	Antibacterial Activity	MiBIG BGC-ID	Minimum	Maximum	Length (nt.)		
1	T3PKS	Herboxidiene T1PKS, T3PKS (7%)	Potentially Inhibits Cell Wall Biosynthesis	BGC0001065	214109	229085	14977		
2	ectoine	Ectoine Other (100%)		BGC0000853	672467	682865	10399		
3	melanin	Melanin Other (60%)		BGC0000911	1414668	1425276	10609		
4	siderophore	Desferrioxamine B Siderophore (66%)		BGC0000941	1507497	1519344	11848		
5	furan	Methylenomycin Other (9%)	Unknown MOA	BGC0000914	2667706	2688702	20997		
6	NRPS	Ansamitocin P-3 T1PKS (7%)		BGC0001511	3021294	3076234	54941		
7	terpene	Albaflavenone Terpene (100%)		BGC0000660	3743916	3764845	20930		
8	T2PKS	Spore pigment T2PKS (66%)		BGC0000271	3800219	3857023	56805		
9	siderophore	-	-	-	4274698	4286154	11457		
10	bacteriocin	-	-	-	4471015	4482384	11370		
11	terpene	-	-	-	4488880	4508472	19593		
12	NRPS	Lipopeptide 8D1-1 & 8D1-2 NRPS (25%)	PMF Collapse	BGC0001370	4616241	4669719	53479		
13	NRPS	Lipopeptide 8D1-1 & 8D1-2 NRPS (15%)	PMF Collapse	BGC0001370	4929292	4967094	37803		
14	terpene	Hopene Terpene (76%)	Cell Wall Biosynthesis	BGC0000663	5042481	5069167	26687		
15	terpene	Lysolipin T2PKS (4%)		BGC0000242	5088579	5103665	15087		
16	T1PKS	Candicidin T1PKS (28%)		BGC0000034	5331251	5350185	18935		
17	T2PKS, butyrolactone	Kinamycin T2PKS (25%)		BGC0000236	5370217	5395391	25175		
18	T1PKS	FR-008/Levorin A3 T1PKS (28%)	DNA Synthesis	BGC0000061	5395392	5413565	18174		

The most similar BGCs in the MiBIG database are listed, as well as the percentage of genes in each MiBIG known cluster that have similarity to genes in the corresponding JS cluster. In cases where the most similar known BGC produces an antibiotic, the MOA was listed (Methylenomycin,³⁸ Albaflavenone,³⁶ Lipopeptide 8D1-1 & 8D1-2,³⁹ Lysolipin,⁴⁰ Kinamycin⁴¹).

Of the 26 putative BGCs that were identified in the genome of strain SFW, only one cluster shared a high percentage of genes in common with a known antibiotic-producing cluster (Table 4). Cluster 1 shared similarity with 62% of the genes within a known BGC that produces carbapenems (Fig 5C), a class of beta-lactam antibiotics that inhibit cell wall biogenesis.⁴² Additionally, cluster 4 contained a low percentage of genes in common with a BGC involved in the synthesis of clavulanic acid, which inhibits beta-lactamase and consequently strengthens the bactericidal activity of beta-lactams. Cluster 1 (and perhaps cluster 4) could, therefore,

contribute to the synthesis of bioactive molecules that account for the inhibition of *E. coli* cell wall biogenesis on ISP2 media (Fig 3).

Table 4. BGCs encoded within the draft genome sequence of strain SFW.

Strain - SFW							
Cluster	Type	Most Similar MiBIG Cluster and Predicted Percent Similarity	Antibacterial Activity	MiBIG BGC-ID	Minimum	Maximum	Length (nt.)
1	NRPS, blactam	Carbapenem MM 4550 Other (62%)	Cell Wall Biosynthesis	BGC0000842	280243	420495	140253
2	NRPS	Coelichelin NRPS (90%)		BGC0000325	537068	587954	50887
3	terpene	Isorenieratene Terpene (28%)		BGC0000664	601924	615201	13278
4	blactam	Clavulanic acid Other (20%)	Beta-lactamase Inhibition	BGC0000845	839861	863248	23388
5	terpene	Hopene Terpene (69%)		BGC0000663	920123	946635	26513
6	T1PKS	Sceliphrolactam T1PKS (72%)		BGC0001770	1325302	1388343	63042
7	bacteriocin	-	-	-	1598693	1609196	10504
8	lanthipeptide	Kanamycin Saccharide (1%)	Protein Synthesis	BGC0000703	1734703	1760347	25645
9	NRPS	Lipopeptide 8D1-1 & 8D1-2 NRPS (6%)	PMF Collapse	BGC0001370	1773501	1830128	56628
10	siderophore	Ficellomycin NRPS (3%)	DNA replication	BGC0001593	2107372	2120491	13120
11	terpene	-	-	-	2186197	2205874	19678
12	butyrolactone	Lactonamycin T2PKS (3%)	Protein Synthesis	BGC0000238	4018281	4029096	10816
13	NRPS	Istamycin Saccharide (11%)	Protein Synthesis	BGC0000700	4251191	4307412	56222
14	siderophore	Desferrioxamine B Siderophore (83%)		BGC0000941	4924798	4936579	11782
15	lanthipeptide	-	-	-	5321361	5346350	24990
16	terpene	-	-	-	5588840	5608518	19679
17	ectoine	Ectoine Other (100%)		BGC0000853	6072388	6080990	8603
18	T2PKS, PKS-like	Cinerubin B T2PKS (25%)	DNA Intercalation	BGC0000212	6443938	6515214	71277
19	terpene	Steffimycin T2PKS-Saccharide (16%)		BGC0000273	6560271	6580717	20447
20	terpene, ectoine	Ectoine Other (100%)		BGC0000853	6860752	6881669	20918
21	bacteriocin	-	-	-	6909859	6920014	10156
22	T3PKS	Tetronasin T1PKS (11%)	PMF Collapse	BGC0000163	7071589	7112647	41059
23	melanin	Melanin Other (100%)		BGC0000911	7208921	7219385	10465
24	T2PKS, terpene	Spore pigment T2PKS (83%)		BGC0000271	7244899	7317424	72526
25	NRPS	Rimosamide NRPS (21%)		BGC0001760	7458615	7511513	52899
26	butyrolactone	-	-	-	7592249	7602533	10285

The most similar BGCs in the MiBIG database are listed, as well as the percentage of genes in each MiBIG known cluster that have similarity to genes in the corresponding SFW cluster. In cases where the most similar known BGC produces an antibiotic, the MOA was listed (Carbapenem,⁴³ Clavulanic acid,⁴⁴ Kanamycin,⁴⁵ Lipopeptide 8D1-1 & 8D1-2,³⁹ Ficellomycin,³³ Lactonamycin,^{46, 47} Istamycin,⁴⁸⁻⁵⁰ Cinerubin,⁵¹ Tetronasin⁵²).

Strain DF encoded 34 BGCs (Table 5). Despite this rich supply of BGCs, however, we were only able to identify one cluster that likely participates in the synthesis of an antibiotic with a confirmed MOA. According to AntiSMASH v5.0, cluster 27 shared 85% gene identity with a known BGC that produces nonactin, a bioactive ionophore that disrupts membrane potential⁵³ (Fig 5D). The known clusters could not fully account for the antibacterial activity exhibited by strain DF (Fig 3), suggesting that antibiotics might be produced by novel clusters.

Table 5. BGCs encoded within the closed genome sequence of strain DF.

Strain - DF							
Cluster	Type	Most Similar MiBIG Cluster and Predicted Percent Similarity	Antibacterial Activity	MiBIG BGC-ID	Minimum	Maximum	Length (nt.)
1	ectoine	-	-	-	55194	65595	10402
2	butyrolactone	Gamma-butyrolactone (100%)	-	BGC0000038	156789	167693	10905
3	terpene	Geosmin Terpene (100%)	-	BGC0001181	196683	218891	22209
4	transAT-PKS, T1PKS, NRPS	Griseobactin NRPS (82%)	-	BGC0000368	227963	343592	115630
5	NRPS	Coelichelin NRPS (81%)	-	BGC0000325	363422	414309	50888
6	NRPS	Arsenopolyketides Other (41%)	Unknown MOA	BGC0001283	421648	466618	44971
7	T3PKS	Naringenin Terpene (100%)	Potentially Inhibits Fatty Acid Biosynthesis	BGC0001310	481558	522683	41126
8	T2PKS	Granaticin T2PKS (29%)	-	BGC0000227	874325	916838	42514
9	terpene	Steffimycin T2PKS-Saccharide (19%)	-	BGC0000273	1079402	1100700	21299
10	ectoine	Ectoine Other (100%)	-	BGC0000853	1578303	1588701	10399
11	NRPS	Laspertomycin NRPS (9%)	Cell Wall Biosynthesis	BGC0000379	2182777	2259643	76867
12	siderophore	Desferrioxamine B Siderophore (100%)	-	BGC0000941	2696558	2708336	11779
13	LAP, thiopeptide	-	-	-	3100604	3132306	31703
14	NRPS	Microsclerodermins NRPS, T1PKS (14%)	-	BGC0001019	3327941	3392550	64610
15	betalactone	Divergolide T1PKS (6%)	-	BGC0001119	3740002	3763908	23907
16	T2PKS	Rabelomycin T2PKS-Saccharide (25%)	-	BGC0000262	4318297	4360617	42321
17	lassopeptide	Keywimycin RiPP (100%)	-	BGC0001634	4419581	4442206	22626
18	T1PKS	Streptazone E T1PKS (91%)	-	BGC0001296	4973731	5038740	65010
19	lanthipeptide	AmfS Lanthipeptide (100%)	-	BGC0000496	5317536	5340199	22664
20	terpene	-	-	-	5677735	5698600	20866
21	siderophore	-	-	-	6150075	6164335	14261
22	butyrolactone	4-formylaminooxyvinylglycine Other (21%)	-	BGC0001488	6317636	6328541	10906
23	bacteriocin	-	-	-	6488518	6499849	11332
24	terpene	2-methylisoborneol Terpene (100%)	-	BGC0000658	6514280	6535368	21089
25	NRPS	Asukamycin T2PKS (12%)	Unknown MOA	BGC0000187	6618679	6679113	60435
26	arylpylene	Tetrapetalone A-D PKS (26%)	Unknown MOA	BGC0001798	6726255	6767369	41115
27	NRPS	Nonactin T2PKS (85%)	Dissipates Transmembrane Electric Potential	BGC0000252	6774441	6840334	65894
28	terpene	Hopenin Terpene (69%)	-	BGC0000663	7093211	7119781	26571
29	NRPS	SGR PTMs NRPS, T1PKS (100%)	Unknown MOA	BGC0001043	7204325	7247340	43016
30	bacteriocin	-	-	-	7260810	7271608	10799
31	melanin	Melanin Other (100%)	-	BGC0000911	7452699	7463166	10468
32	T3PKS	Herboxidiene T1PKS, T3PKS (7%)	-	BGC0001065	7495580	7536629	41050
33	terpene	Isorenieratene Terpene (100%)	-	BGC0000664	7626626	7652190	25565
34	thiopeptide, LAP, NRPS	Lactazole Thiopeptide (33%)	-	BGC0000606	7666072	7717669	51598

The most similar BGCs in the MiBIG database are listed, as well as the percentage of genes in each MiBIG known cluster that have similarity to genes in the corresponding DF cluster. In cases where the most similar known BGC produces an antibiotic, the MOA was listed (Arsenopolyketides,⁵⁴ Naringenin,⁵⁵ Laspertomycin,⁵⁶ Asukamycin,⁵⁷ Tetrapetalone,⁵⁸ Nonactin,⁵³ SGR PTMs NRPS⁵⁹).

Antimicrobial Activity of Four *Streptomyces* isolates Against Clinically Relevant Pathogens

In order to assess the relevance of antibiotics produced by strains JS, DF, SFW, and QF2, we screened their ability to inhibit the growth of three clinically isolated pathogens using the cross-streak method (Table 6). Both strain QF2 and strain JS inhibited the growth of methicillin-resistant *S. aureus* (MRSA) and efflux-deficient *P. aeruginosa* PA01. These strains did not, however, inhibit the growth of the wild-type clinical isolates *P. aeruginosa* PA01 and *P. aeruginosa* P4, which were resistant to the antibiotics produced by all four *Streptomyces* isolates. Strain DF, though incapable of inhibiting the growth of *E. coli* *tolC*⁺ (Fig 2), did inhibit the growth of MRSA and efflux-deficient *P. aeruginosa* PA01. Strain SFW was the least capable

of inhibiting the growth of clinical pathogens, producing antibiotics only effective against *E. coli* *tolC*⁺ (Fig 2).

Table 6. Inhibition of growth of clinically relevant pathogens by Streptomyces strains DF, SFW, QF2, and JS.

	Gram-Negative Bacteria			Gram-Positive Bacteria		
	<i>P. aeruginosa</i>			<i>S. aureus</i>		
	PAO1	P4	PAO1 Δefflux	MRSA USA300	TCH1516	
DF	-	-	+			+
SFW	-	-	-			-
QF2	-	-	+			+
JS	-	-	+			+

Plus signs indicate growth inhibition, while minus signs indicate pathogen growth.

Phage Isolation and Genome Sequencing

With the goal of identifying genetic tools that could be used to augment expression of the BGCs in our bacterial isolates, we decided to isolate bacteriophages using *S. platensis* as a host. This species, in particular, was chosen as a host because it is relatively well-characterized, and *S. platensis* phages capable of infecting our Streptomyces isolates could possibly be used to move (via transduction) BGCs from our isolates into a more genetically manipulatable and familiar background.^{60, 61} Thus, to increase the probability that our *S. platensis* phages could be used for this purpose, we performed the isolation using the same soil samples from which our Streptomyces strains were obtained. Four *S. platensis* bacteriophages (BartholomewSD, IceWarrior, Shawty, and TrvxScott) were successfully isolated. These phages were imaged using negative-stain transmission electron microscopy (Fig 6A) and subsequently characterized as members of the family Siphoviridae due to their long filamentous tails and icosahedral capsids.^{62, 63} Genome sequencing revealed that BartholomewSD (52.1 kb) and TrvxScott (52.6 kb) are 89% identical (Fig 6B) and belong to the BD2 subcluster of Streptomyces phages, which currently contains 20 other members.¹⁹ IceWarrior (55.5 kb) clustered in subcluster BI1 (24 members), and

Shawty (40.7 kb) clustered in BB1, a subcluster of 7 phages that includes notable members TG1 and phiC31 (Table 7).¹⁹ The BLASTp-predicted functions of the gene products encoded by these phages are shown in Table 8.

Fig 6. Characterization of four Streptomyces phages isolated from soil samples. (a) Electron micrographs of the four phages (IceWarrior, TrvxScott, BartholomewSD, and Shawty). Lysate samples were negatively stained and imaged with transmission electron microscopy (TEM). The scale bar represents 100 nm. (b) A whole-genome sequence comparison of the four phages generated by Phamerator (top to bottom: TrvxScott, BartholomewSD, Shawty, IceWarrior).

Table 7. Summary of the NCBI WGS annotations of four phage isolates.

Bacteriophage	Taxa	Genome (bp)	GC%	Genes	Cluster	Subcluster	Genbank Acc. No.
TrvxScott	unclass. Arequatrovirus	52600	67.8	81	BD	BD2	MH669016
IceWarrior	unclass. Rimavirus	55532	59.5	86	BI	BI1	MK433259
BartholomewSD	unclass. Arequatrovirus	52131	67.6	88	BD	BD2	MK460245
Shawty	unclass. Lomovskayavirus	40733	63.2	58	BB	BB1	MK433266

Table 8. Functions of the putative proteins encoded within the genomes of four phage isolates.

Streptomyces Phage TrvxScott taxon:2301575		Streptomyces phage Shawty taxon:2510521		Streptomyces phage IceWarrior taxon:2510515		Streptomyces phage BartholomewSD taxon:2510587	
CDS No.	Product	CDS No.	Product	CDS No.	Product	CDS No.	Product
1	hypothetical protein	1	terminase small subunit	1	hypothetical protein	1	hypothetical protein
2	HNH endonuclease	2	terminase large subunit	2	HNH endonuclease	2	hypothetical protein
3	thioredoxin	3	portal protein	3	hypothetical protein	3	HNH endonuclease
4	terminase small subunit	4	capsid maturation protease	4	hypothetical protein	4	tRNA-Phe
5	terminase large subunit	5	major capsid protein	5	endolysin	5	hypothetical protein
6	portal protein	6	head-to-tail adaptor	6	head-to-tail connector complex protein	6	thioredoxin
7	capsid maturation protease	7	hypothetical protein	7	hypothetical protein	7	hypothetical protein
8	scaffolding protein	8	major tail protein	8	terminase large subunit	8	terminase
9	major capsid protein	9	hypothetical protein	9	hypothetical protein	9	portal protein
10	head-to-tail connector complex protein	10	tail assembly chaperone	10	hypothetical protein	10	MuF-like minor capsid protein
11	head-to-tail connector complex protein	11	tail assembly chaperone	11	hypothetical protein	11	scaffolding protein
12	hypothetical protein	12	tape measure protein	12	portal protein	12	major capsid protein
13	head-to-tail connector complex protein	13	minor tail protein	13	hypothetical protein	13	head-to-tail adaptor
14	major tail protein	14	minor tail protein	14	capsid maturation protease	14	head-to-tail stopper
15	tail assembly chaperone	15	minor tail protein	15	hypothetical protein	15	hypothetical protein
16	tail assembly chaperone	16	minor tail protein	16	hypothetical protein	16	tail terminator
17	tape measure protein	17	hypothetical protein	17	major tail protein	17	major tail protein
18	minor tail protein	18	tail fiber	18	hypothetical protein	18	tail assembly chaperone
19	minor tail protein	19	lysine A	19	major tail protein	19	tail assembly chaperone
20	hypothetical protein	20	hypothetical protein	20	hypothetical protein	20	tape measure protein
21	hypothetical protein	21	deoxynucleoside monophosphate kinase	21	chitosanase	21	minor tail protein
22	hypothetical protein	22	immunity repressor	22	hypothetical protein	22	minor tail protein
23	minor tail protein	23	Cas4 family exonuclease	23	tape measure protein	23	hypothetical protein
24	hypothetical protein	24	hypothetical protein	24	minor tail protein	24	minor tail protein
25	lysine A	25	hypothetical protein	25	minor tail protein	25	hypothetical protein
26	hypothetical protein	26	hypothetical protein	26	hypothetical protein	26	hypothetical protein
27	hypothetical protein	27	hypothetical protein	27	hypothetical protein	27	hypothetical protein
28	hypothetical protein	28	hypothetical protein	28	hypothetical protein	28	lysine A
29	hypothetical protein	29	hypothetical protein	29	holin	29	hypothetical protein
30	exonuclease	30	HNH endonuclease	30	hypothetical protein	30	hypothetical protein
31	hypothetical protein	31	DNA primase	31	hypothetical protein	31	immunity repressor
32	hypothetical protein	32	restriction endonuclease	32	hypothetical protein	32	hypothetical protein
33	hypothetical protein	33	DNA polymerase I	33	hypothetical protein	33	Cas4 family exonuclease
34	deoxycytidylate deaminase	34	RNA polymerase sigma factor	34	hypothetical protein	34	hypothetical protein
35	DNA helicase	35	hypothetical protein	35	hypothetical protein	35	hypothetical protein
36	holliday junction resolvase	36	hypothetical protein	36	hypothetical protein	36	hypothetical protein
37	hypothetical protein	37	hypothetical protein	37	hypothetical protein	37	deoxycytidylate deaminase
38	DNA primase	38	hypothetical protein	38	hypothetical protein	38	DnaB-like helicase
39	DNA primase	39	hypothetical protein	39	hypothetical protein	39	holliday junction resolvase
40	hypothetical protein	40	ThyX-like thymidylate synthase	40	hypothetical protein	40	hypothetical protein
41	hypothetical protein	41	hypothetical protein	41	hypothetical protein	41	DNA primase
42	exonuclease	42	hypothetical protein	42	hypothetical protein	42	DNA primase
43	hypothetical protein	43	thioredoxin	43	hypothetical protein	43	hypothetical protein
44	HTH DNA binding protein	44	hypothetical protein	44	hypothetical protein	44	hypothetical protein
45	hypothetical protein	45	deoxycytidylate deaminase	45	hypothetical protein	45	hypothetical protein
46	ribonucleotide reductase	46	hypothetical protein	46	hypothetical protein	46	Mre11 family dsDNA break repair endo/exonuclease
47	DNA methylase	47	hypothetical protein	47	hypothetical protein	47	hypothetical protein
48	hypothetical protein	48	hypothetical protein	48	hypothetical protein	48	helix-turn-helix DNA binding protein
49	hypothetical protein	49	hypothetical protein	49	hypothetical protein	49	hypothetical protein
50	hypothetical protein	50	hypothetical protein	50	hypothetical protein	50	ribonucleotide reductase
51	HTH DNA binding protein	51	hypothetical protein	51	hypothetical protein	51	hypothetical protein
52	integrase	52	hypothetical protein	52	hypothetical protein	52	hypothetical protein
53	hypothetical protein	53	protein kinase	53	hypothetical protein	53	hypothetical protein
54	thymidylate synthase	54	integrase	54	hypothetical protein	54	hypothetical protein
55	hypothetical protein	55	tRNA-Asp	55	hypothetical protein	55	helix-turn-helix DNA binding protein
56	hypothetical protein	56	tRNA-Thr	56	hypothetical protein	56	integrase
57	hypothetical protein	57	hypothetical protein	57	hypothetical protein	57	hypothetical protein
58	hypothetical protein	58	HNH endonuclease	58	hypothetical protein	58	ThyX-like thymidylate synthase
59	hypothetical protein	59		59	DNA primase/polymerase	59	hypothetical protein
60	hypothetical protein	60		60	hypothetical protein	60	hypothetical protein
61	deoxynucleoside monophosphate kinase	61		61	hypothetical protein	61	hypothetical protein
62	hypothetical protein	62		62	hypothetical protein	62	hypothetical protein
63	hypothetical protein	63		63	hypothetical protein	63	hypothetical protein
64	hypothetical protein	64		64	hypothetical protein	64	hypothetical protein
65	hypothetical protein	65		65	hypothetical protein	65	deoxymononucleoside kinase
66	hypothetical protein	66		66	hypothetical protein	66	hypothetical protein
67	hypothetical protein	67		67	hypothetical protein	67	hypothetical protein
68	hypothetical protein	68		68	hypothetical protein	68	hypothetical protein
69	hypothetical protein	69		69	hypothetical protein	69	hypothetical protein
70	hypothetical protein	70		70	hypothetical protein	70	hypothetical protein
71	hypothetical protein	71		71	hypothetical protein	71	hypothetical protein
72	hypothetical protein	72		72	hypothetical protein	72	hypothetical protein
73	hypothetical protein	73		73	hypothetical protein	73	hypothetical protein
74	hypothetical protein	74		74	hypothetical protein	74	hypothetical protein
75	hypothetical protein	75		75	hypothetical protein	75	hypothetical protein
76	hypothetical protein	76		76	hypothetical protein	76	hypothetical protein
77	hypothetical protein	77		77	hypothetical protein	77	hypothetical protein
78	acetyltransferase	78		78	hypothetical protein	78	hypothetical protein
79	hypothetical protein	79		79	hypothetical protein	79	hypothetical protein
80	hypothetical protein	80		80	hypothetical protein	80	hypothetical protein
81	hypothetical protein	81		81	DNA helicase	81	hypothetical protein
		82		82	HNH endonuclease	82	hypothetical protein
		83		83	hydrolase	83	hypothetical protein
		84		84	DNA helicase	84	hypothetical protein
		85		85	helix-turn-helix DNA binding domain protein	85	hypothetical protein
		86		86	hypothetical protein	86	hypothetical protein
				87		87	hypothetical protein
				88		88	hypothetical protein

Characterization of CRISPR Elements in the Genomes of Our Streptomyces Strains

Prior to testing the ability of our phages to infect the *Streptomyces* isolates, we decided to examine the strains for complete and functional CRISPR/Cas systems. Our reasoning for this was two-fold. First, the presence of acquired spacers and their specific sequences would allow us to make predictions about whether or not our phages can infect our antibiotic-producing strains. And, second, it was conceivable that in doing so we might discover a novel CRISPR/Cas-based system. Our bioinformatic analysis identified the presence of Cas enzymes and CRISPR arrays within the genomes all four of our *Streptomyces* isolates, but the abundance of CRISPRs in each strain varied greatly (Table 9). QF2 contained the largest number of predicted CRISPRs – 39 in total, scattered around the chromosome, each containing between one and 25 spacers (Fig 4B, purple; Table 10). Some predicted spacers within these arrays matched with 94-100% identity to sequences within TrvxScott (7 spacers), BartholomewSD (4 spacers), Shawty (2 spacers), and IceWarrior (5 spacers) (Table 10, Table 11). Spacers targeted a variety of genes including those encoding minor tail proteins, tape measure proteins, deoxycytidylate deaminase, helix-turn-helix DNA binding proteins, endolysin, and capsid maturation protease (Fig 7). The large number of putative spacers in the QF2 genome targeting TrvxScott, BartholomewSD, Shawty, and IceWarrior suggests that strain QF2 has likely previously encountered and acquired resistance to each of these phages. Moreover, strain QF2 was isolated from the same soil sample as BartholomewSD, providing support for these findings. Strain QF2 also encoded seven proteins of a Type IE CRISPR-Cas system.⁶⁴⁻⁶⁸ The QF2 proteins were distantly related to the enzymes of the canonical Cas3 system in *E. coli* (Fig 8), but the operon in strain QF2 lacked two genes (Cas1 and Cas2) involved in spacer acquisition. This phenomenon, the absence of Cas1 and Cas2, has previously been reported as a common feature of Streptomycetaceae Type IE systems.⁶⁹ The presence in the QF2 genome of a Cas3 system and spacers mapping to essential

proteins in each of the genomes of our phages suggests that the strain is likely resistant to all four of our phages, and thus, transduction is unlikely with strain QF2.

Table 9. General characteristics of predicted CRISPR-Cas systems within the genomes of strains DF, SFW, QF2, and JS.

Strains	CRISPR	Spacers	Repeats	Spacers with Blastn Hits to Host Range Phage	Cas Loci	Cas- Associated Genes
DF	12	14	26	2	3	9
SFW	11	23	34	2	3	17
QF2	39	168	207	14	5	22
JS	4	8	12	0	4	20

Included in this table is the number of spacers within the genome of each bacterial strain with sequence similarity to regions within any of the four phage isolates (IceWarrior, TrvxScott, BartholomewSD, or Shawty).

Table 10. Characteristics of the 38 CRISPRs predicted in the draft genome sequence of strain QF2.

Strain QF2						
CRISPRs	Min	Max	Length (nt.)	No. Repeats	No. Spacers	Spacer Blastn Hit to Host Range Phage
1	384313	384428	116	2	1	S1 [BartholomewSD]
2	437656	437755	100	2	1	
3	835820	835922	103	2	1	
4	1307659	1307752	94	2	1	
5	1344898	1345205	308	6	5	
6	1543053	1543144	92	2	1	
7	1616159	1616542	384	5	4	S13 [BartholomewSD, TrvxScott]
8	1618208	1618291	84	2	1	
9	1833873	1833977	105	2	1	
10	1861097	1861197	101	2	1	
11	2316101	2316187	87	2	1	
12	2704894	2704990	97	2	1	
13	3015981	3016376	396	7	6	S21 [IceWarrior]
14	3106815	3107262	448	8	7	S27 [BartholomewSD, TrvxScott]
15	3112452	3113433	982	17	16	S34 [IceWarrior], S41 [TrvxScott]
16	3138610	3140161	1,552	26	25	
17	3145439	3145830	392	7	6	
18	3444685	3444779	95	2	1	
19	3507440	3507598	159	3	2	
20	3791739	3792012	274	5	4	
21	3838730	3838804	75	2	1	
22	4257871	4258080	210	4	3	S89 [IceWarrior]
23	4327550	4328916	1,367	23	22	S105 [IceWarrior], S106 [Shawty, TrvxScott]
24	4333365	4334666	1,302	22	21	S118 [TrvxScott], S131 [Shawty, TrvxScott]
25	4335904	4336481	578	10	9	S138 [BartholomewSD], S140 [TrvxScott]
26	4522773	4522879	107	2	1	
27	4528080	4528148	69	2	1	
28	4657265	4657374	110	2	1	
29	4754273	4754356	84	2	1	
30	4787509	4787642	134	3	2	
31	4987714	4987810	97	2	1	
32	5305650	5305745	96	2	1	
33	5400452	5400541	90	2	1	S151 [IceWarrior]
34	5417083	5417162	80	2	1	
35	5441923	5442032	110	2	1	
36	6552625	6552699	75	2	1	
37	6798173	6798309	137	3	2	
38	7177566	7177797	232	6	5	

Spacers with sequence similarity to any of the four phages in this study are listed next to their corresponding CRISPR and are identified according to their position relative to all other spacers within the QF2 genome.

Table 11. Spacers within the genomes of strain DF, SFW, and QF2 that have sequence similarity to at least one of the four phage isolates.

Strain DF									
CRISPR	Spacer	Blatn Hit to Host Range Phase	Score (bits) / E Val.	No. Identities (%ID)	Strand	Minimum	Maximum	Length (nt.)	Sequence
9	S11	BartholomewSD	30.2 (15) / 2.4	15/15 (100%)	Plus / Minus	4171019	4171050	32	TGCCACCCGGCCGAGCCGCTCCGCGAGGCAG
10	S12	TrvxScott	30.2 (15) / 4.9	15/15 (100%)	Plus / Plus	4684554	4684602	49	GGTGTCGCCCGCGGTGCGGTGCATGCTCTCGGCTTGAGCGGGCTGCCG
		BartholomewSD	28.2 (14) / 9.5	14/14 (100%)	Plus / Minus				
Strain SFW									
CRISPR	Spacer	Blatn Hit to Host Range Phase	Score (bits) / E Val.	No. Identities (%ID)	Strand	Minimum	Maximum	Length (nt.)	Sequence
1	S1	Shawty	28.2 (14) / 5.9	14/14 (100%)	Plus / Plus	250625	250650	26	GAGTCACGAGCCGGGCGAAGGCACGC
5	S6	IceWarrior	32.2 (16) / 0.99	19/20 (95%)	Plus / Plus	4553831	4553872	42	CGGGCGTCGACGGTGACGAGCGTCGCGCTGCTACTTCTCTTG
Strain QF2									
CRISPR	Spacer	Blatn Hit to Host Range Phase	Score (bits) / E Val.	No. Identities (%ID)	Strand	Minimum	Maximum	Length (nt.)	Sequence
1	S1	BartholomewSD	32.2 (16) / 1.2	16/16 (100%)	Plus / Plus	384347	384394	48	GCGGACGGCGCGCGCGCGGTACCCCGGTGTCCACGACGGCGCGCG
7	S13	BartholomewSD	32.2 (16) / 1.9	16/16 (100%)	Plus / Plus	1616356	1616423	68	CGACCTGCGGTACCACTCGATCCGGCGCGGTCCCATCTACAAGGGCACGGTGTCCAGCGGACCGAG
13	S21	TrvxScott	30.2 (15) / 1.9	16/16 (100%)	Plus / Plus				
		IceWarrior	30.2 (15) / 2.5	15/15 (100%)	Plus / Minus	3016077	3016109	33	CGCCGGAACCTCAAGGAGGAGACGCGCGGG
14	S27	BartholomewSD	30.2 (15) / 3.1	18/19 (94%)	Plus / Minus	3106902	3106938	37	AGGGCGTGGCGGTGCGGGGTGCGGGTGGAGCTGTGGT
		TrvxScott	30.2 (15) / 3.1	18/19 (94%)	Plus / Minus				
15	S34	IceWarrior	30.2 (15) / 2.4	15/15 (100%)	Plus / Plus	3112541	3112572	32	ACAGCGACGTCGCCCTACAACCTACGCCGCTGG
		TrvxScott	28.2 (14) / 9.5	14/14 (100%)	Plus / Plus	3112961	3112992	32	GGTGTGAACCGCTCGCGCGCGCTGCACTTGT
22	S89	IceWarrior	30.2 (15) / 2.5	15/15 (100%)	Plus / Plus	4257958	4257990	33	CGCGGGCGTCCCTTCGCCGAGGAGACCTGCCC
		IceWarrior	30.2 (15) / 2.5	15/15 (100%)	Plus / Minus	4328429	4328461	33	CATCAGCGTCTGAAGCAGCAGCGCCATCGCCTT
23	S106	Shawty	28.2 (14) / 9.5	14/14 (100%)	Plus / Plus	4328491	4328522	32	TGATCGAGCGCGGACGGGCACATCAGCGGCC
		TrvxScott	28.2 (14) / 9.5	14/14 (100%)	Plus / Minus				
		S118	28.2 (14) / 9.5	14/14 (100%)	Plus / Plus	4333692	4333723	32	GCCGCGTCCGGCTACGGCTACGGCTCCGCCCC
		S131	28.2 (14) / 9.5	14/14 (100%)	Plus / Minus	4334483	4334514	32	AACGCGCTCATGAGGCGCTGCGTTTGGCGTC
		S138	28.2 (14) / 9.5	14/14 (100%)	Plus / Plus	4336177	4336208	32	AACGCGGACGATGCGCCGTACGAGCGCGG
		S140	28.2 (14) / 9.5	14/14 (100%)	Plus / Minus	4336299	4336330	32	ATCTCGCGTCCAGACCGCTCGACGCAAT
33	S151	IceWarrior	36.2 (18) / 0.063	18/18 (100%)	Plus / Minus	5400476	5400517	42	GTGGTGGCCTCGCCGACGAGTGTCTCGGACGCTGGGCGGC

The bold portion of each spacer shares high sequence similarity with a region in the genome of the listed host range phage. In cases where a single spacer mapped to two phages, bold and underlined are used to sequences distinguish the two.

Fig 7. Genomic maps of phages showing regions containing sequence similarity to spacers found within the CRISPRs of strains QF2, DF, and SFW. (a) TrvxScott, (b) BartholomewSD, (c) Shawty, and (d) IceWarrior. Key displays putative functions of CRISPR targeted genes.

Fig 8. Class I, Type I-E CRISPR-Cas system encoded in the WGS of strain QF2. (top) The Type I-E CRISPR-Cas operon encoded by strain QG is located from 4,973,482 to 4,987,810 and includes seven genes. The Type I-E cascade is followed by CRISPR 31, consisting of two repeats and a single spacer. **(bottom)** The canonical Type I-E CRISPR-Cas system encoded in the genome of *E. coli* K-12 MG1655 is located from 2,887,219 to 2,877,618 and includes eight genes. The Type I-E cascade is followed by a CRISPR 31, consisting of five repeats and four spacers.

Specific spacers mapping to some of our phages were also discovered within the genomes of strains DF and SFW (but not JS). Strain DF contained two spacers that mapped to sequences within the genome of BartholomewSD, and one of these spacers also shared sequence similarity with a region in TrvxScott (Table 11, Table 12). Strain SFW contained two spacers – one that shared sequence similarity with Shawty and another that mapped to a sequence in IceWarrior (Table 11, Table 13). Both strains DF and SFW encoded proteins containing regions with

similarity to the RuvC and HNH endonuclease domains of known Cas enzymes. However, given the limited similarity of these putative proteins to known Cas proteins, further study is necessary to determine if they constitute novel Cas systems. If these systems are functional, we predict that strain DF is resistant to infection by TrvxScott and BartholomewSD, and strain SFW, resistant to Shawty and IceWarrior.

Table 12. Characteristics of the 12 CRISPRs predicted in the complete genome sequence of strain DF.

Strain DF						
CRISPRs	Min	Max	Length (nt.)	No. Repeats	No. Spacers	Spacer Blastn Hit to Host Range Phage
1	1491701	1491797	97	2	1	
2	1517080	1517203	124	2	1	
3	1669449	1669522	74	2	1	
4	1995371	1995463	93	2	1	
5	2013671	2013864	194	3	2	
6	2241854	2241958	105	2	1	
7	2443114	2443225	112	2	1	
8	3767692	3767796	105	2	1	
9	4170935	4171076	142	3	2	S11 [BartholomewSD]
10	4684531	4684625	95	2	1	S12 [BartholomewSD, TrvxScott]
11	4802187	4802274	88	2	1	
12	5682554	5682639	86	2	1	

Spacers with sequence similarity to any of the four phages in this study are listed next to their corresponding CRISPR and are identified according to their position relative to all other spacers within the DF genome.

Table 13. Characteristics of the 11 CRISPRs predicted in the draft genome sequence of strain SFW.

Strain SFW						
CRISPRs	Min	Max	Length (nt.)	No. Repeats	No. Spacers	Spacer Blastn Hit to Host Range Phage
1	250601	250674	74	2	1	S1 [Shawty]
2	452415	452685	271	3	2	
3	2779718	2779796	79	2	1	
4	2824197	2824275	79	2	1	
5	4553801	4554168	368	5	4	S6 [IceWarrior]
6	4731148	4731651	504	10	9	
7	5166339	5166424	86	2	1	
8	5317268	5317371	104	2	1	
9	5722922	5723016	95	2	1	
10	6532966	6533033	68	2	1	
11	7183989	7184080	92	2	1	

Spacers with sequence similarity to any of the four phages in this study are listed next to their corresponding CRISPR and are identified according to their position relative to all other spacers within the SFW genome.

398

399 A curious feature emerged from our analysis of the CRISPRs within the *Streptomyces* strains.
 400 Among the 213 predicted spacers encoded by all four bacterial strains, 18 contained sequence
 401 similarity (95-100% identity) with at least one of the four phages (Table 9). The lengths of the
 402 matching sequences (100% identity) within bacterial spacers ranged from 14 to 18 nucleotides
 403 and accounted for approximately half the length of a typical spacer. Additionally, a single spacer
 404 occasionally appeared capable of targeting two distantly related phages. These spacers
 405 contained sequences mapping to two distinct genes encoded by different viral genomes. For
 406 example, spacer 106 in CRISPR 23 of strain QF2 is 32 nucleotides in length, and it contains 14
 407 bases that share 100% identity with a region in the TrvxScott tape measure gene. These 14
 408 bases overlap (by 8 nucleotides) with another sequence that is 14 base pairs in length and shares
 409 100% identity with a region within the Shawty genome (Table 11). If these spacers functionally
 410 serve to resist infection, our analysis suggests that a single spacer may evolve to efficiently target
 411 more than one phage, thus providing broad immunity.

412

413 *Susceptibility of Streptomyces strains to infection by S. platensis phages*

414 With the hope of identifying phages that might serve as tools for transduction, we assessed the
 415 susceptibility of our *Streptomyces* isolates to infection by each of the four *S. platensis*
 416 phages. As predicted, strain QF2, with its Type IE Cas system and many CRISPRs containing
 417 spacers against our phages, could not be infected by any of our four phages. Strain DF
 418 experienced inefficient infection by TrvxScott (~2.0 x 10⁴-fold reduced plating efficiency
 419 compared to *S. platensis*) and was completely resistant to infection by BartholomewSD. In
 420 addition to these results, which were generally predicted by our CRISPR/Cas findings, we also
 421 demonstrated strain DF's resistance to infection by Shawty and susceptibility to IceWarrior (~20-
 422 fold reduced efficiency). Strain SFW was at least partially resistant to infection by all four phages:

Shawty (no infection), BartholomewSD (no infection), IceWarrior ($\sim 10^7$ -fold reduced efficiency), and TrvxScott ($\sim 1.3 \times 10^5$ -fold reduced efficiency). Finally, strain JS, despite having no spacers specifically targeting our phages, was similarly immune to infection by Shawty and BartholomewSD and partially resistant to infection by IceWarrior and TrvxScott ($\sim 10^6$ -fold and $\sim 10^5$ -fold reduced efficiency, respectively). These data are consistent with our predictions regarding the resistance of our *Streptomyces* isolates to infection by the phages against which they carry spacers, though it is the case that the presence of a spacer did not always confer complete immunity to the phage it targeted. In some cases, strains containing spacers could be infected relatively inefficiently by the targeted phage. For example, strain DF encoded a single spacer targeting TrvxScott but remained partially susceptible to infection. Strains DF, SFW, and JS were all capable of being infected by TrvxScott and IceWarrior to some degree. Thus, it remains possible that these two phages could be used for transducing BGCs into *S. platensis*.

Identification of phage integrases

In analyzing the proteins encoded within the genomes of our phages, we identified site-specific serine recombinases encoded by BartholomewSD, TrvxScott, and Shawty. The integrases of BartholomewSD and TrvxScott were nearly identical and shared similarity to integrases belonging to a number of previously studied phages. The Shawty integrase shared protein sequence similarity with the integrases of *Streptomyces* phages TG1 and phiC31 (71.3% and 51.9% respectively). The TG1 and phiC31 integrases are distinct recombinases that share 49.7% protein sequence identity and have been used extensively as tools for integrating genes of interest into specific loci within the genomes of a wide variety of organisms, ranging from soil microbes such as *Streptomyces* to multicellular animals such as *Drosophila*.²¹⁻²⁵ Thus, as a newly discovered member of this serine integrase family, the Shawty integrase could also

potentially be used to move *Streptomyces*-encoded BGCs between strains in order to facilitate augmented expression of bioactive natural products.

This work demonstrates a method for effectively studying newly isolated, antibiotic-producing bacteria and phages that may infect them. In particular, we have highlighted how BCP can be used to assess the novelty of BGCs encoded within *Streptomyces* strains, providing a solution to the problem of dereplication. Moreover, our work illustrates how the isolation and genomic analysis of phages that infect antibiotic-producing *Streptomyces* strains might yield new genetic tools such as transducing phages or integrases, which can be used to augment the expression of novel antibiotics. Ultimately, our identification of potentially novel BGCs and phage integrases serves as a foundation for further studies that could lead to the discovery of new antibiotics.

Materials and Methods

Soil sample collection and site description

Undergraduate students enrolled in the Phage Hunters Advancing Genomics and Evolutionary Science (PHAGE) class at UCSD collected soil samples for isolating bacteria and their associated phages. Soil samples (approx. 30 ml) were collected around San Diego County (32.7157° N, 117.1611° W), California, USA.

*Isolation of *Streptomyces**

Actinomycete isolation agar (AIA) plates (for one liter: sodium caseinate 2 g, L-Asparagine 0.1 g, sodium propionate 4 g, dipotassium phosphate 0.5 g, magnesium sulphate 0.1 g, glycerol 5 mL, rifampicin 50 µg mL⁻¹, nystatin 100 µg mL⁻¹, cycloheximide 100 µg mL⁻¹, agarose 15 g, pH 8.1) were used to select for Actinobacteria from soil samples. One gram of soil was added to the agar surface and streaked across the AIA plate and incubated for two days at 30°C. The plates

were investigated for individual colonies with morphologies indicative of *Streptomyces* (vegetative hyphae, aerial mycelium), and those colonies were picked and purified at least four times on AIA-CHX plates.

Phage isolation and purification

Bacteriophages were isolated from soil samples with host, *Streptomyces platensis* JCM 4664 substr. MJ1A1. An enrichment culture was prepared from 1 g soil and 2.5 ml of *S. platensis* added to 15 ml of Luria-Bertani (LB) medium (for one liter: tryptone 10 g, yeast extract 5 g, NaCl 10 g, agar 15 g, pH 7.0), followed by a 2-day incubation at 30°C with shaking. Phage were isolated from a 1.2 ml volume of enrichment culture that was centrifuged at maximum speed for 3 min, 1 ml of the resulting supernatant was filtered (0.22 µm filter), and 5 µl of the filtrate was spotted and then streaked onto an LB plate containing 100 µg/ml of cycloheximide. *S. platensis* (0.1 ml) was mixed with 4.5 ml of LB top agar 0.7%, poured over the streak plate and incubated for two days at 30°C. Resultant plaques were re-streaked onto new LB plates containing 100 µg/ml of cycloheximide about 3-4 times for phage purification.

Bacterial genomic DNA extraction and quantification for 16S rRNA PCR amplification and sequencing

An adaptation of the DNeasy® Blood & Tissue Kit (Qiagen) protocol was used for bacterial genomic DNA extraction. Strains were cultured overnight at 30°C in 5 ml of LB broth while rolling. Cells were pelleted (16,000 x g, 3 min) from 1 ml of culture, re-suspended in 180 µl of lysis buffer (prepared in house), and incubated at 37°C for 45 min after which 25 µl of proteinase K (20 mg/mL) and 200 µl Buffer AL (Qiagen) was added. The samples were vortexed at maximum speed for 20 sec, incubated at 56°C for 30 min, and 200 µl of ethanol (96-100%) was added. The samples were vortexed at maximum speed for 30 sec, added to a DNeasy Mini spin column,

centrifuged (16,000 x g, 1 min), and the supernatant was discarded. Buffer AW2 (Qigen) was added (500 µl), followed by centrifugation (20,000 x g, 3 min). The DNeasy Mini spin column was placed into a sterile 2 ml microcentrifuge tube, and the gDNA was eluted in 100 µl of AE Buffer by centrifugation (20,000 x g, 1 min) following a 1 min incubation at room temperature. The gDNA concentration was quantified (1 µl sample volume) with a Thermo Scientific™ NanoDrop™ One Microvolume UV-Vis Spectrophotometer (840274100) and stored at -20°C.

Bacterial Genomic DNA extraction for PacBio whole-genome sequencing

High molecular weight genomic DNA (20-160 kb) was extracted from four *Streptomyces* strains (DF, SFW, QF2, and JS) with the QIAGEN-Genomic-tip 500/G kit (10262) according to the manufacturer's protocol for bacteria.

Bacterial whole-genome sequencing, assembly, and annotation

The genome sequences of four *Streptomyces* strains were generated using the Pacific Biosciences RS II (PacBio RS II) single molecule real-time (SMRT) sequencing platform at the IGM Genomics Center, University of California, San Diego, La Jolla, CA. Genome sequences were assembled using the HGAP protocol integrated in the PacBio RS II sequencer (smrt analysis v2.3.0/Patch5) resulting in a variable number (n = 1-95) of contigs per genome, and ranged in size from 5.42 to 7.79 Mb. The mauve contig mover was used to order the contigs of three draft genome sequences (genomes of strains SFW, QF2, and JS) relative to a closely related reference sequence (*S. pratensis* ATCC 33331, *S. globisporus* C-1027, and *S. parvulus* 2297 respectively). DNA sequencing of strain DF resulted in a single contig and did not require reordering to restore gene synteny. Gene prediction and annotation were made with the Rapid Annotations using Subsystems Technology (RASTtk) platform.⁷⁰

Phage genomic DNA extraction

5 µl of RNase A and 5 µl of DNase I were added to 10ml of lysate, incubated at 30°C for 30 minutes, and then precipitated overnight at 4°C by the addition of 4 ml of 20% polyethylene glycol 8,000. Samples were centrifuged at 10,000 g's for 30 minutes, and pellets resuspended in Qiagen PB buffer and DNA isolated using a Qiagen plasmid DNA isolation column as recommended by the manufacturer.

Phage genome sequencing, assembly, and annotation

Genomic DNA of 4 bacteriophages (TrvxScott, BartholomewSD, Shawty, and IceWarrior) was sequenced using the Illumina MiSeq platform at the Pittsburgh Bacteriophage Institute sequencing facility. The genomes were assembled with Newbler and checked for quality with Consed. The whole genome sequences were submitted to GenBank (Acc No. MH669016, MK460245, MK433266, and MK433259). DNA Master was used for annotation, and NCBI BLASTp was used to determine the potential function of gene products. Whole genome sequence comparisons were performed in Phamerator.⁷¹

16S rRNA PCR amplification and sequencing

16S ribosomal DNA templates (~1,465 bp) were amplified using Q5 high fidelity PCR (New England Biolabs) with the universal primer set 27F (5'-AGAGTTTGATCCTGGCTCAG-3') and 1492R (5'-GGTACCTTGTTACGACTT-3').⁷² Each PCR mixture (50 µl) contained 100 ng of template gDNA, 500 pmol of each primer, and 200 µM dNTPs. PCR thermocycling conditions were as follows: 30 seconds of initial denaturation at 98°C, 30 cycles of denaturation at 98°C for 10 seconds, annealing for 15 seconds at 60°C, extension at 72°C for 1.5 minutes, and a final extension at 72°C for 5 minutes then held at 4°C. PCR products were purified with the oligonucleotide cleanup protocol as described in the Monarch PCR & DNA Cleanup Kit 5 µg user

manual (NEB #T1030). Clean PCR products were sequenced using Sanger methods by Eton Biosciences (<https://www.etonbio.com/>) and trimmed for quality before analysis.

CRISPR-Cas sequence analysis and predictions

The sequences of all four *Streptomyces* were searched for CRISPR arrays (repeats and spacers) and potentially associated Cas genes using the following software tools; CRISPR-Cas++,^{64, 65} CRISPROne,⁶⁶ CRISPRMiner2,⁶⁷ and CRISPRDetect.⁶⁸

Phylogenetic analyses of bacterial isolates

16S rRNA sequences were trimmed on both ends, (5' and 3') in Geneious Prime using the Trim Ends function with an error probability limit set at 0.05, which trims regions with more than a 5% chance of an error per base. Sequences were aligned using MUSCLE v3.8.425 with a maximum of 1,000 iterations, and profile-dependent parameters were set. Sequences were grouped based on similarity, and anchor optimized. Distance was measured using kmer4_6 for the first iteration, and subsequent distance measurements were performed with pctid_kimura. For all iterations the clustering method was neighbor joining, and the sequence weighting scheme was ClustalW with an spm objective score, 32 anchor spacing and -1 gap open score. Diagonals were set at a minimum length of 24 and a margin of 5, the minimum column anchor scores were set min best and min smoothed at 90. Finally, the hydrophobicity multiplier was set at 1.2 with a window size of 5. The tree was created with the Geneious Tree Builder using the Jukes-Cantor genetic distance model, Neighbor-Joining method to build the tree, with the outgroup set as *Chlorobium limicola* strain DSM 245. The resulting consensus tree was obtained after 1,000 iterations of bootstrap resampling, with a support threshold set at 70% and a random seed of 409,923.

Cross-streak method for assessing antibacterial production potential

From a single colony, using sterile Q-tips, *Streptomyces* isolates were streaked in a broad vertical line (2 inch) onto LB, and AIA, solid agar plates and incubated for one week at 30°C. The day before the assay, test strains (*E. coli* JP313 $\Delta tolC$, *B. subtilis* PY79, and *E. coli* MC4100) were grown in 5 ml of LB and incubated at 30°C overnight while rolling. On the day of the antibacterial screen, the overnight cultures of each test strain were diluted (1:100 in 5 ml LB) and grown to log phase OD₆₀₀ 0.15-0.2 (~1.5 hr at 30°C while rolling). A volume of 10 µl of each test strain was spotted in distinct lines almost to the edge of the *Streptomyces* line at a perpendicular angle. The plates were incubated overnight at 30°C, then investigated for the presence of zones of inhibition which were measured in millimeters.

Bacterial cytological profiling (BCP) on plates

Fluorescence microscopy and BCP on plates was performed as previously described by Nonejuie et. al.¹⁵ Briefly, *Streptomyces* strains (DF, SFW, QF2, and JS) were streaked in a vertical line down the center of LB, AIA, and ISP2 plates (for one liter: 4.0 g Difco yeast extract, 10.0 g Difco malt extract, 4.0 g dextrose, 20.0 g agar, pH 7.0), incubated for one week at 30°C. The test strain, *E. coli* JP313 $\Delta tolC$, was prepared and spotted as described above in the cross-streak method. Following a 2 hr incubation at 30°C, a 1.5 x 1.5 cm square (~2.5 cm²) piece of agar containing the *E. coli* test strain was cut and prepared for high resolution fluorescence microscopy. The cut piece of agar was placed on a microscope slide, the *E. coli* cells were stained with fluorescent dyes, a coverslip was placed on top of the stained cells then imaged.

Host range experiment

The host ranges of 4 phages were determined against the *Streptomyces* strains: QF2, DF, JS, and SFW. The experiment was blinded by assigning phages numbers i-iv and hosts letters A-D.

A lawn of *Streptomyces* in LB top agar was poured on LB CHX plates. After the top agar solidified, a grid was drawn on the bottom of the plate, and 5 μ l of pre-diluted phage samples (10^0 to 10^{-10} in phage buffer) were spotted in squares on the grid. Plaques were counted and used to calculate a titer, which was then compared to the titer obtained against *S. platensis* in order to calculate the efficiency of infection.

Transmission electron microscopy

10 μ l of lysate was applied to a copper grid, stained with 1% uranyl acetate, washed twice with phage buffer, and allowed to dry. Images were collected using a FEI Tecnai Spirit G2 BioTWIN Transmission Electron Microscope equipped with a bottom mount Eagle 4k camera.

Strains used in this study

We used the following strains: *S. platensis* JCM 4664 substr. MJ1A1, *E. coli* MC4100, *B. subtilis* PY79, *P. aeruginosa* P4, *S. aureus* MRSA USA300 TCH1516 from Texas Children's Hospital (USA300-HOU-MR), *S. coelicolor* A3(2) substr. M146, *E. coli* JP313 $\Delta tolC$, as well as two strains generously donated by Prof. Keith Poole at Queens University in Kingston, Canada – *P. aeruginosa* PA01 and *P. aeruginosa* K2733 Δ efflux ($\Delta MexAB-OprM$, $\Delta MexCD-OprJ$, $\Delta MexEF-OprN$, $\Delta MexXY-OprM$). The $\Delta tolC5$ mutation is derived from strain EW1b (CGSC #5634), and was introduced into strain JP313 by P1 transduction. JP313 was transduced to tetracycline resistance with a lysate of strain CAG18475 (*metC162::Tn10*), and the methionine requirement of the transductants was confirmed. This strain was then transduced to prototrophy with a lysate of EW1b, and these transductants were screened on MacConkey agar for the presence of the $\Delta tolC5$ mutation. EW1b and CAG18475 were obtained from the *Coli* Genetic Stock Center at Yale University.

Data Availability

All data generated or analyzed in this study are included in this article and its supplementary information files. Nucleotide sequences of the phages in this study have been deposited in the GenBank database with accession numbers: MK460245 (BartholomewSD), MH669016 (TrvxScott), MK433266 (Shawty), and MK433259 (IceWarrior).

Competing Interests

The authors declare no competing interests.

References

1. Wenciewicz, T.A., *Crossroads of Antibiotic Resistance and Biosynthesis*. J Mol Biol, 2019. **431**(18): p. 3370-3399.
2. Pogue, J., et al., *Appropriate antimicrobial therapy in the era of multidrug-resistant human pathogens*. Clinical Microbiology and Infection, 2015. **21**(4): p. 302-312.
3. Brown, E.D. and G.D. Wright, *Antibacterial drug discovery in the resistance era*. Nature, 2016. **529**(7586): p. 336-343.
4. Newman, D.J. and G.M. Cragg, *Natural products as sources of new drugs from 1981 to 2014*. Journal of natural products, 2016. **79**(3): p. 629-661.
5. Fischbach, M.A. and C.T. Walsh, *Antibiotics for emerging pathogens*. Science, 2009. **325**(5944): p. 1089-1093.
6. Hover, B.M., et al., *Culture-independent discovery of the malacidins as calcium-dependent antibiotics with activity against multidrug-resistant Gram-positive pathogens*. Nature microbiology, 2018. **3**(4): p. 415-422.
7. Ling, L.L., et al., *A new antibiotic kills pathogens without detectable resistance*. Nature, 2015. **517**(7535): p. 455-459.
8. Chen, R., H.L. Wong, and B.P. Burns, *New approaches to detect biosynthetic gene clusters in the environment*. Medicines, 2019. **6**(1): p. 32.
9. Pye, C.R., et al., *Retrospective analysis of natural products provides insights for future discovery trends*. Proceedings of the National Academy of Sciences, 2017. **114**(22): p. 5601-5606.
10. Cox, G., et al., *A Common Platform for Antibiotic Dereplication and Adjuvant Discovery*. Cell Chem Biol, 2017. **24**(1): p. 98-109.
11. Nonejuie, P., et al., *Bacterial cytological profiling rapidly identifies the cellular pathways targeted by antibacterial molecules*. Proceedings of the National Academy of Sciences, 2013. **110**(40): p. 16169-16174.

12. Lamsa, A., et al., *Rapid inhibition profiling in Bacillus subtilis to identify the mechanism of action of new antimicrobials*. ACS chemical biology, 2016. **11**(8): p. 2222-2231.
13. Lamsa, A., et al., *The Bacillus subtilis cannibalism toxin SDP collapses the proton motive force and induces autolysis*. Molecular microbiology, 2012. **84**(3): p. 486-500.
14. Peters, C.E., et al., *Rapid Inhibition Profiling Identifies a Keystone Target in the Nucleotide Biosynthesis Pathway*. ACS chemical biology, 2018. **13**(12): p. 3251-3258.
15. Nonejuie, P., et al., *Application of bacterial cytological profiling to crude natural product extracts reveals the antibacterial arsenal of Bacillus subtilis*. The Journal of antibiotics, 2016. **69**(5): p. 353.
16. Quach, D., et al., *Bacterial cytological profiling (BCP) as a rapid and accurate antimicrobial susceptibility testing method for Staphylococcus aureus*. EBioMedicine, 2016. **4**: p. 95-103.
17. Alberti, F. and C. Corre, *Editing streptomycete genomes in the CRISPR/Cas9 age*. Natural product reports, 2019. **36**(9): p. 1237-1248.
18. de Jonge, P.A., et al., *Molecular and evolutionary determinants of bacteriophage host range*. Trends in microbiology, 2019. **27**(1): p. 51-63.
19. Russell, D.A. and G.F. Hatfull, *PhagesDB: the actinobacteriophage database*. Bioinformatics, 2017. **33**(5): p. 784-786.
20. Genilloud, O., *Actinomycetes: still a source of novel antibiotics*. Natural product reports, 2017. **34**(10): p. 1203-1232.
21. Thorpe, H.M., S.E. Wilson, and M.C. Smith, *Control of directionality in the site-specific recombination system of the Streptomyces phage ϕ C31*. Molecular microbiology, 2000. **38**(2): p. 232-241.
22. Fogg, P.C.M., et al., *New Applications for Phage Integrases*. Journal of Molecular Biology, 2014. **426**(15): p. 2703-2716.
23. Morita, K., et al., *The site-specific recombination system of actinophage TG1*. FEMS Microbiol Lett, 2009. **297**(2): p. 234-40.
24. Morita, K., et al., *Site-specific genome integration in alphaproteobacteria mediated by TG1 integrase*. Applied Microbiology and Biotechnology, 2012. **93**(1): p. 295-304.
25. Venken, K.J.T., et al., *Genome engineering: Drosophila melanogaster and beyond*. Wiley interdisciplinary reviews. Developmental biology, 2016. **5**(2): p. 233-267.
26. Thakur, D., et al., *Influence of nutrition and culturing conditions for optimum growth and antimicrobial metabolite production by Streptomyces sp. 201*. Journal de Mycologie Medicale, 2009. **19**(3): p. 161-167.
27. Zimmerman, S.B., *Toroidal nucleoids in Escherichia coli exposed to chloramphenicol*. Journal of structural biology, 2002. **138**(3): p. 199-206.
28. Aziz, R.K., et al., *The RAST Server: rapid annotations using subsystems technology*. BMC genomics, 2008. **9**(1): p. 75.
29. Blin, K., et al., *antiSMASH 5.0: updates to the secondary metabolite genome mining pipeline*. Nucleic acids research, 2019. **47**(W1): p. W81-W87.
30. Kautsar, S.A., et al., *MIBiG 2.0: a repository for biosynthetic gene clusters of known function*. Nucleic acids research, 2020. **48**(D1): p. D454-D458.
31. Holm, M., et al., *Molecular mechanism of viomycin inhibition of peptide elongation in bacteria*. Proceedings of the National Academy of Sciences, 2016. **113**(4): p. 978-983.
32. Komatsu, Y. and K. Tanaka, *Mechanism of Action of Showdomycin: Part II. Effect of Showdomycin on the Synthesis of Deoxyribonucleic Acid in Escherichia coli*. Agricultural and Biological Chemistry, 1970. **34**(6): p. 891-899.
33. Reusser, F., *Ficellomycin and feldamycin; inhibitors of bacterial semiconservative DNA replication*. Biochemistry, 1977. **16**(15): p. 3406-3412.

34. Pohle, S., et al., *Biosynthetic gene cluster of the non-ribosomally synthesized cyclodepsipeptide skyllamycin: deciphering unprecedented ways of unusual hydroxylation reactions*. Journal of the American Chemical Society, 2011. **133**(16): p. 6194-6205.
35. Kanda, N., et al., *DB-2073, a new alkylresorcinol antibiotic*. The Journal of antibiotics, 1975. **28**(12): p. 935-942.
36. GÜRTLER, H., et al., *Albaflavenone, a sesquiterpene ketone with a zizaene skeleton produced by a streptomycete with a new rope morphology*. The Journal of antibiotics, 1994. **47**(4): p. 434-439.
37. Zhao, B., et al., *Biosynthesis of the sesquiterpene antibiotic albaflavenone in Streptomyces coelicolor A3 (2)*. Journal of Biological Chemistry, 2008. **283**(13): p. 8183-8189.
38. Hobbs, G., et al., *An integrated approach to studying regulation of production of the antibiotic methylenomycin by Streptomyces coelicolor A3 (2)*. Journal of bacteriology, 1992. **174**(5): p. 1487-1494.
39. Baltz, R.H., V. Miao, and S.K. Wrigley, *Natural products to drugs: daptomycin and related lipopeptide antibiotics*. Natural Product Reports, 2005. **22**(6): p. 717-741.
40. Drautz, H., W. Keller-Schierlein, and H. Zähner, *Stoffwechselprodukte von Mikroorganismen*. Archives of Microbiology, 1975. **106**(3): p. 175-190.
41. Ballard, T.E. and C. Melander, *Kinamycin-mediated DNA cleavage under biomimetic conditions*. Tetrahedron Letters, 2008. **49**(19): p. 3157-3161.
42. Nagarajan, R., et al., *β -Lactam Antibiotics from Streptomyces*. Journal of the American Chemical Society, 1971. **93**(9): p. 2308-2310.
43. Papp-Wallace, K.M., et al., *Carbapenems: past, present, and future*. Antimicrobial agents and chemotherapy, 2011. **55**(11): p. 4943-4960.
44. Neu, H.C. and K.P. Fu, *Clavulanic acid, a novel inhibitor of β -lactamases*. Antimicrobial agents and chemotherapy, 1978. **14**(5): p. 650-655.
45. Suzuki, J., T. KUNIMOTO, and M. HORI, *Effects of kanamycin on protein synthesis: inhibition of elongation of peptide chains*. The Journal of antibiotics, 1970. **23**(2): p. 99-101.
46. Zhang, X., et al., *Biosynthetic Investigations of Lactonamycin and Lactonamycin Z: Cloning of the Biosynthetic Gene Clusters and Discovery of an Unusual Starter Unit*. Antimicrobial Agents and Chemotherapy, 2008. **52**(2): p. 574.
47. Matsumoto, N., et al., *Lactonamycin, a new antimicrobial antibiotic produced by Streptomyces rishiriensis MJ773-88K4*. The Journal of antibiotics, 1999. **52**(3): p. 269-275.
48. Yamamoto, H., et al., *Ribosomal resistance of an istamycin producer, Streptomyces tenjimariensis, to aminoglycoside antibiotics*. Biochemical and biophysical research communications, 1981. **100**(3): p. 1396-1401.
49. Hoang, N.H., et al., *Istamycin aminoglycosides profiling and their characterization in Streptomyces tenjimariensis ATCC 31603 culture using high-performance liquid chromatography with tandem mass spectrometry*. Journal of separation science, 2016. **39**(24): p. 4712-4722.
50. Okami, Y., et al., *New aminoglycoside antibiotics, istamycins A and B*. The Journal of antibiotics, 1979. **32**(9): p. 964-966.
51. Hertweck, C., et al., *Type II polyketide synthases: gaining a deeper insight into enzymatic teamwork*. Natural product reports, 2007. **24**(1): p. 162-190.
52. Newbold, C.J., et al., *Effect of the novel ionophore tetrinasin (ICI 139603) on ruminal microorganisms*. Appl. Environ. Microbiol., 1988. **54**(2): p. 544-547.

53. Kusche, B.R., et al., *Alternating pattern of stereochemistry in the nonactin macrocycle is required for antibacterial activity and efficient ion binding*. Journal of the American Chemical Society, 2009. **131**(47): p. 17155-17165.
54. Cruz-Morales, P., et al., *Phylogenomic analysis of natural products biosynthetic gene clusters allows discovery of arseno-organic metabolites in model streptomycetes*. Genome biology and evolution, 2016. **8**(6): p. 1906-1916.
55. Wang, L.-H., et al., *Membrane and genomic DNA dual-targeting of citrus flavonoid naringenin against Staphylococcus aureus*. Integrative Biology, 2017. **9**(10): p. 820-829.
56. Kleijn, L.H., et al., *Total synthesis of Laspartomycin C and characterization of its antibacterial mechanism of action*. Journal of medicinal chemistry, 2016. **59**(7): p. 3569-3574.
57. Rui, Z., et al., *Biochemical and genetic insights into asukamycin biosynthesis*. Journal of Biological Chemistry, 2010. **285**(32): p. 24915-24924.
58. Li, C., X. Li, and R. Hong, *Synthetic Study on Tetrapetalones: Stereoselective Cyclization of N-Acyliminium Ion To Construct Substituted 1-Benzazepines*. Organic letters, 2009. **11**(18): p. 4036-4039.
59. Luo, Y., et al., *Activation and characterization of a cryptic polycyclic tetramate macrolactam biosynthetic gene cluster*. Nature communications, 2013. **4**(1): p. 1-8.
60. Smanski, M.J., et al., *Engineered Streptomyces platensis Strains That Overproduce Antibiotics Platensimycin and Platencin*. Antimicrobial Agents and Chemotherapy, 2009. **53**(4): p. 1299.
61. Burke, J., D. Schneider, and J. Westpheling, *Generalized transduction in Streptomyces coelicolor*. Proceedings of the National Academy of Sciences, 2001. **98**(11): p. 6289.
62. Ackermann, H.-W., *5500 Phages examined in the electron microscope*. Archives of virology, 2007. **152**(2): p. 227-243.
63. Jin, H., et al., *Capsid Structure of a Freshwater Cyanophage Siphoviridae Mic1*. Structure, 2019. **27**(10): p. 1508-1516. e3.
64. Couvin, D., et al., *CRISPRCasFinder, an update of CRISPRFinder, includes a portable version, enhanced performance and integrates search for Cas proteins*. Nucleic acids research, 2018. **46**(W1): p. W246-W251.
65. Abby, S.S., et al., *MacSyFinder: a program to mine genomes for molecular systems with an application to CRISPR-Cas systems*. PloS one, 2014. **9**(10).
66. Zhang, Q. and Y. Ye, *Not all predicted CRISPR-Cas systems are equal: isolated cas genes and classes of CRISPR like elements*. BMC bioinformatics, 2017. **18**(1): p. 92.
67. Zhang, F., et al., *CRISPRminer is a knowledge base for exploring CRISPR-Cas systems in microbe and phage interactions*. Communications Biology, 2018. **1**(1): p. 1-5.
68. Biswas, A., et al., *CRISPRDetect: a flexible algorithm to define CRISPR arrays*. BMC genomics, 2016. **17**(1): p. 356.
69. Faure, G., et al., *CRISPR-Cas in mobile genetic elements: counter-defence and beyond*. Nature Reviews Microbiology, 2019. **17**(8): p. 513-525.
70. Brettin, T., et al., *RASTtk: A modular and extensible implementation of the RAST algorithm for building custom annotation pipelines and annotating batches of genomes*. Scientific Reports, 2015. **5**(1): p. 8365.
71. Cresawn, S.G., et al., *Phamerator: a bioinformatic tool for comparative bacteriophage genomics*. BMC Bioinformatics, 2011. **12**(1): p. 395.
72. Janda, J.M. and S.L. Abbott, *16S rRNA Gene Sequencing for Bacterial Identification in the Diagnostic Laboratory: Pluses, Perils, and Pitfalls*. Journal of Clinical Microbiology, 2007. **45**(9): p. 2761.

A.

B.

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Type I-E CRISPR-Cas System

Strain QF2

E. coli K-12 MG1655

Figure 8