Novel reference transcriptomes for the sponges *Carteriospongia foliascens* and *Cliona orientalis* and associated algal symbiont *Gerakladium endoclionum*

Brian W. Strehlow^{*1,2,3,4,5,6}, Mari-Carmen Pineda^{5,6}, Carly D. Kenkel^{5,7}, Patrick Laffy⁵, Alan Duckworth^{5,6}, Michael Renton^{2,8}, Peta L. Clode^{2,3,4}, Nicole S Webster^{5,6,9}

*corresponding author

Author emails, respectively: strehow@biology.sdu.dk, mcarmen.pineda@gmail.com, ckenkel@usc.edu, plaffy@aims.gov.au, aduckworth1@gmail.com, michael.renton@uwa.edu.au, peta.clode@uwa.edu.au, N.Webster@aims.gov.au

¹ Current address: Department of Biology, Nordcee, University of Southern Denmark, Campusvej 55, 5230 Odense, Denmark

² School of Biological Sciences, University of Western Australia, 35 Stirling Hwy, Crawley WA 6009, Australia

³ Centre for Microscopy, Characterisation and Analysis, University of Western Australia, 35 Stirling Hwy, Crawley WA 6009, Australia

⁴ Oceans Institute, University of Western Australia, 35 Stirling Hwy, Crawley WA 6009, Australia

⁵ Australian Institute of Marine Science, PMB No 3, Townsville MC, Queensland 48106 Western Australian Marine Science Institution, Crawley, WA, Australia

⁶ Western Australian Marine Science Institution, 35 Stirling Hwy, Crawley WA 6009, Australia

⁷ Department of Biological Sciences, University of Southern California, 3616 Trousdale Parkway, Los Angeles, CA 90089, USA

⁸ School of Agriculture and Environment, University of Western Australia, 35 Stirling Hwy, Crawley WA 6009, Australia

⁹ Australian Centre for Ecogenomics, School of Chemistry and Molecular Biosciences, The University of Queensland, Brisbane, QLD, Australia

Abstract

Transcriptomes from sponges are important resources for studying the stress responses of these ecologically important filter feeders, the interactions between sponges and their symbionts, and the evolutionary history of metazoans. Here, we generated reference transcriptomes for two common and cosmopolitan Indo-Pacific sponge species: *Carteriospongia foliascens* and *Cliona orientalis*. We also created a reference transcriptome for the primary symbiont of *C. orientalis* – *Gerakladium endoclionum*. To ensure a full repertoire of transcripts were included, clones of each sponge species were exposed to a range of individual stressors: decreased salinity, elevated temperature, elevated suspended sediment concentrations, sediment deposition and light attenuation. RNA extracted from all treatments was pooled for each species, using equal concentrations from each clone. Sequencing of pooled RNA yielded 409 and 418 million raw reads for *C. foliascens* and *C. orientalis* holobionts (host and symbionts), respectively. Reads underwent quality trimming before assembly with Trinity. Assemblies were filtered into sponge-specific or, for *G. endoclionum*, symbiont-specific assemblies. Assemblies for *C. foliascens*, *C. orientalis*, and *G. endoclionum* contained 67,304, 82,895, and 28,670 contigs, respectively. Contigs represented 15,248-37,344 isogroups (~genes) per assembly, and N50s ranged 1,672-4,355 bp. Gene ortholog analysis verified a high level of completeness and quality for sponge-specific transcriptomes, with an average 93% of core EuKaryotic Orthologous Groups (KOGs) and 98% of single-copy metazoan core gene orthologs identified. The *G. endoclionum* assembly was partial with only 56% of core KOGs and 32% of single-copy eukaryotic core gene orthologs identified. These reference transcriptomes are a valuable resource for future molecular research aimed at assessing sponge stress responses.

Keywords

Porifera, transcriptome, sponge, *Cliona orientalis*, *Carteriospongia foliascens*, *Gerakladium endoclionum*

Data Description

Sponges, phylum Porifera, represent one of the oldest lineages of multicellular animals [1], hence investigating the transcriptomes of different sponge species can provide insight into the evolution of metazoans and their gene expression profiles. Furthermore, sponges have an uncertain future in the face of global climate change [2,3] as well as local stressors including coastal development, altered hydrological processes, and increased runoff of nutrients, pesticides and sediments [4–7]. Transcriptomic analysis of sponges that have been exposed to different environmental conditions would improve our understanding of the sponge molecular stress response pathways and enhance our ability to effectively manage these ecologically important filter feeders. Although there are approximately 9,000 described sponge species [8], to date only ~35 species have published transcriptomes [9-25] and only ~10 have published genomes [10,16,26–29].

In this study, we assembled the transcriptomes of two common and widely distributed Indo-Pacific sponge species – *Carteriospongia foliascens* and *Cliona orientalis*. Both are emerging model species that have been extensively used to study the physiological and ecological effects of environmental stressors on sponges [30–37]. *C. foliascens* and *C. orientalis* are only the second members of their respective orders (Dictyoceratida and Clionaida) to have a reference transcriptome sequenced. Whilst both *C. foliascens* and *C. orientalis* host diverse populations of bacterial symbionts, e.g. [32], *C. orientalis* additionally hosts an abundant population of eukaryotic Symbiodiniaceae, *Gerakladium endoclionum* [38,39], which comprises up to 96% of its algal symbiont community [37]. We used sequences generated from the *C. orientalis* holobiont, i.e. host and symbiont, to construct a partial reference transcriptome for *Gerakladium endoclionum*. Matching host and symbiont transcriptomes provide a valuable tool to understand the holobiont response to changing environmental conditions and determine the cause-effect pathways for declining host health with environmental change. These data contribute substantially to available poriferan genetic resources and advance the development of these two sponge species as model systems for field and laboratory studies.

Methods

Samples and sequencing

Samples of *C. foliascens* and *C. orientalis* were collected in May 2015 from Fantome Is. (S 18°41.028' E 146° 30.706) and Pelorus Is. (S 18°32.903' E 146° 29.172'), respectively, in the central Great Barrier Reef under permits G12/35236.1 and G13/35758.1. As *C. orientalis* is a bioeroding sponge that encrusts and erodes coral skeletons, five *C. orientalis* cores (~5 cm in diameter) were collected using an air-drill from a single individual, i.e. cloned, growing on a dead colony of *Porites* sp. An individual of *C. foliascens* was cut (cloned) into five pieces as in [32]. Sponges were healed and acclimated under natural light and flow-through seawater for 4 weeks before experiments were performed.

In order to capture the full complement of gene expression within the reference transcriptomes, sponges were subjected to five different treatments at the Australian Institute of Marine Science (AIMS) National Sea Simulator: i) decreased salinity, ii) elevated temperature, iii) elevated suspended sediment concentrations (SSCs) and sediment deposition, iv) light attenuation and v) no stress control. Sponge clones were used across all treatments to control for genotype, i.e. one genotype was used per species. Two clones of each species were used for each treatment. In the salinity stress treatment, salinity was decreased from 35 to 22 parts per thousand (ppt) by gradually adding flow-through reverse osmosis (RO) water to the system. Salinity was held constant at 22 ppt for 2 d with flow-through seawater maintained at 600 mL min⁻¹. In the heat stress treatment, sponges were exposed to a constant temperature of 32.5°C for 1 d using methods described in [32]. In the sediment treatment, sponges were exposed to elevated SSCs at 200 mg L⁻¹ for 1 d as in [40,41], using sediments described therein. In the deposition experiment, sedimentation was approximately 40 mg cm⁻², measured using SedPods [42] and sponges were left covered with sediment for 1 d. In the light attenuation treatment, sponges were kept in complete darkness for 2 d. Immediately after each treatment, a sample of sponge tissue (~1 cm³) was flash frozen in liquid nitrogen and stored at -80°C [43] for RNA extraction and sequencing (RNA-seq). After exposure to the decreased

salinity and darkness treatments, *Cliona orientalis* was visibly bleached after 2 d, but *C. foliascens* did not exhibit any colour change. Sponges were not visibly affected, e.g. no bleaching or necrosis, by the sediment exposure or elevated seawater temperature.

Approximately 50 mg of each sponge clone was excised and ground using a mortar and pestle. Grinding was performed under a thin layer of liquid nitrogen to limit RNA degradation. All tools were rinsed in ethanol followed by RNase Zap (Sigma-Aldrich, USA) to remove contamination and deactivate RNA degrading enzymes. Total RNA was isolated using the Zymo ZR RNA miniprep kit (Zymo Research, USA), with in-column DNase digestion, according to the manufacturer's protocol. Total RNA was subsequently cleaned using the Zymo RNA Clean and Concentrator kit (Zymo Research, USA), following the Manufacturer protocol.

Total RNA quality was checked using gel electrophoresis and spectroscopy (NanoDrop 2000c Spectrophotometer, Thermo Fisher Scientific, USA), and quantified using a Quant-iT Ribogreen Assay (Thermo Fisher Scientific, USA). For each sponge species, the RNA from individual treatments was combined in equal amounts (740 ng for *C. foliascens* and 1,440 ng for *C. orientalis*) from all sponge clones, to a final total RNA concentration of 3.7 µg in 40 µl of Dnase and Rnase free water (93 ng µl⁻¹) for *C. foliascens* and 7.2 µg in 55 µl of Dnase and Rnase free water (160 ng µl⁻¹) for *C. orientalis*. For *C. foliascens* and *C. orientalis* respectively, RNA had a ratio of absorbance at 260 nm to 280 nm (A260/A280) of 1.88 and 2.02 and an A260/A230 ratio of 1.11 and 1.67. To isolate eukaryotic messenger RNA (mRNA), a TruSeq Stranded mRNA-seq sample prep was performed prior to sequencing. The mRNA was sequenced across two lanes of Illumina HiSeq2500 at the Ramaciotti Centre for Genomics (University of New South Wales, Sydney, Australia) to generate 2 x 100 base pair (bp) paired-end (PE) rapid runs.

Transcriptome assembly and annotation

Sequencing produced 409 and 418 million raw reads for *C. foliascens* and *C. orientalis*, respectively (Table 1). Reads were trimmed and assembled using publicly available scripts [44,45] and the protocol detailed in

[46]. Briefly, reads < 50 bp long were removed along with reads containing a homopolymer run of adenine (A) longer than 9 bases using *fastx_toolkit* [47], and only reads with a PHRED quality score >20 over 80% of the read were retained. TruSeq sequencing adapters and PCR duplicates were also removed [44]. The remaining filtered, high quality reads (32.5, 50.5 million paired reads and 2.9, 8.7 million unpaired reads for *C. foliascens* and *C. orientalis*, respectively) underwent *de novo* assembly into contigs using Trinity v 2.8.5 [48]. Data processing and assembly was performed at AIMS in Townsville using a high-performance computer (HPC) and on ABACUS 2.0 at the Danish e-Infrastructure Cooperation (DeiC) National HPC Center. Following assembly, additional quality control was performed to ensure that only target transcripts, i.e. derived from *C. foliascens*, *C. orientalis* or *G. endoclonium*, were included in their respective reference transcriptomes [13,46]. First, contigs less than 400 bp were removed and ribosomal RNA (rRNA), mitochondrial RNA (mtRNA), Symbiodiniaceae, and other non-metazoan (e.g. bacteria) sequences were identified using a series of hierarchical BLAST [49] searches. Transcriptomes were further blasted (BLASTn) against the *A. queenslandica* rRNA database (SILVA: ACUQ01015651) [50], which was the most complete Poriferan rRNA database. Contigs with a bit-score >45 were removed, i.e. 9 and 10 sequences in *C. foliascens* and *C. orientalis*, respectively. This process was repeated using the *A. queenslandica* mitochondrial genome (NCBI: NC_008944.1 REF), resulting in 61 and 27 sequences being removed from the *C. foliascens* and *C. orientalis* assemblies respectively.

Remaining contigs were blasted (BLASTx) against the most complete Poriferan (*A. queenslandica*, aqu2.1_Genes_proteins.fasta) [28,51] and *Symbiodinium kawagutii* (*Symbiodinium_kawagutii*.0819.final.gene.pep) [52] predicted proteomes and the NCBI nonredundant (nr) protein database (downloaded September 2019). In order to be included in a sponge-specific assembly, contigs had to return a more significant match (E value $\leq 10^{-5}$) to the *A. queenslandica* proteome compared to blast results from the *S. kawagutii* proteome *and also* match a metazoan sequence in the nr database or have no match in the nr database, as described in [13]. Sequences with no match to either proteome were

excluded from the final sponge assemblies [13], a stricter exclusion procedure than used in prior invertebrate transcriptome assemblies [46]. For the *C. orientalis* holobiont, sequences matching the *S. kawagutii* proteome more closely than the *A. queenslandica* proteome ($E \text{ value} \leq 10^{-5}$) and matching to the phylum chromerida in the nr database (or having no match in the nr database) were included in the final *G. endoclionum* assembly. Although *C. foliascens* does not contain intracellular Symbiodiniaceae, the decontamination step was also performed in order to remove any potential algal contamination in the sample, resulting in only a few (1,520, 1% of total number of contigs) contaminating sequences being removed.

Within each of the three transcriptomes, contigs were assigned to isogroups (~genes) and given gene names and gene ontologies (GO) [53] following the protocol previously described in [44,54]. Briefly, the transcriptomes underwent BLAST pairwise sequence comparison (BLASTx) to the UniProt Knowledgebase (UniprotKB/Swiss-Prot) database [55]. Significant BLASTx results ($E \text{ value} \leq 10^{-4}$) were used by CDS_extractor_v2.pl [56] to extract and identify protein coding sequences. Functional annotations were assigned to isogroups based on orthologous comparisons to the eggNOG 4.5 database [57] using eggno-mapper [58]. Kyoto Encyclopedia of Genes and Genomes (KEGG) ids were also assigned to isogroups using the KEGG Automatic Annotation server (KAAS) [59]. The guanine-cytosine (GC) content of transcriptomes was calculated using the BBMap package (Joint Genome Institute, USA) [60]. Transcriptome completeness was assessed by Benchmarking Universal Single-Copy Orthologs (BUSCO) analysis [61] on gVolante [62].

Assembly evaluation and quality control

The holobiont assemblies of *C. foliascens* and *C. orientalis* contained 225,126 ($N50 = 1,284$) and 146,510 ($N50 = 1,949$) contigs greater than 400bp in length (Table 1). After data partitioning, 67,304 and 82,895 contigs, for *C. foliascens* and *C. orientalis* respectively, were considered the 'sponge-specific' transcriptome assemblies. The partitioned *G. endoclionum* transcriptome isolated from the *C. orientalis* holobiont comprised 28,670 contigs (Table 1). The *C. foliascens*, *C. orientalis*, and *G. endoclionum* transcriptomes

contained 15,248, 37,344, and 21,566 isogroups, respectively, with mean lengths of 3,024 (N50 = 4,355), 1,756 (N50 = 2,369), and 1,375 (N50 = 1,672) bp (Table 1). The number of isogroups identified in the *C. foliascens* and *C. orientalis* transcriptomes was comparable to previously published sponge transcriptomes which have reported ~11,000-60,000 expressed genes [15,23,63], although there is considerable variation across species. The *G. endoclionum* transcriptome, containing 28,670 isogroups was comparable in size to the previously published *S. kawagutii* genome (36,850 genes) [52] and previously published Symbiodiniaceae transcriptomes, ranging in size from 23,777-26,986 expressed genes [64]. The respective GC content of each assembly was 40.2, 45.5, and 59%, matching reported values for metazoans (35-55% [17,27,65]) and Symbiodiniaceae (45-65% [65]). For *C. foliascens* and *C. orientalis*, the percentage of genes assigned a name or GO terms was 64 and 77%, respectively (Table 1), also comparable to other sponge transcriptomes (30-70% [15]) and those of other non-model metazoans (25-62% [14,46]). In comparison to the annotated sponge transcriptomes, only 39% of *G. endoclionum* isogroups could be assigned function or GO term annotations, however this is consistent with functional annotation of other intracellular Symbiodiniaceae transcriptomes, where between 34-44% of genes were assigned GO terms [64]. The isogroups for *C. foliascens*, *C. orientalis*, and *G. endoclionum* were assigned 3,641, 5,339 and 2,191 unique KEGG annotations respectively.

The representative transcriptomes for *C. foliascens* and *C. orientalis* are considered largely complete based on BUSCO analysis (92.8% and 94.2% complete, respectively) and the representation of nearly all core eukaryotic Orthologous Groups (KOGs) (97.9% and 98.7% respectively) (Table 1). BUSCO analysis of the transcriptome of *G. endoclionum* was 32.3% complete and 56% of core KOGs were identified (Table 1). A reduced BUSCO completeness in transcriptomes isolated from intracellular Symbiodiniaceae in corals (33-42%) has been previously reported [66]. The current *G. endoclionum* transcriptome contained 86% more isogroups than the Symbiodiniaceae transcripts identified within the transcriptome assembly of the closely related sponge holobiont, *Cliona varians* [17]. *C. varians* also hosts a congeneric intracellular

photosymbiont, *Gerakladium spongiolum* [38]. Therefore, the current transcriptome for *G. endoclionum* was considered useful for future studies, at least for conditions *in hospite*.

Table 1. Assembly statistics for the *de novo* transcriptomes.

	<i>Carteriospongia foliascens</i> holobiont	<i>Cliona orientalis</i> holobiont	
N raw reads (x10 ⁶)	409	418	
N qual filtered: PE, SE (x10 ⁶)	32.5, 2.9	50.5, 8.7	
N contigs holobiont	146,510	225,126	
	<i>Carteriospongia foliascens</i>	<i>Cliona orientalis</i>	<i>Gerakladium endoclionum</i>
N contigs target species only	67,304	82,895	28,670
Mean GC content target species only	40.2	45.5	59
N genes	15,248	37,344	21,566
Mean contig length (bp)	3,024	1,756	1,375
N50 (bp)	4,355	2,369	1,672
% Annotated	64	77	39
% core KOGs	97.9	98.7	56
BUSCOs			
N complete (%)	908 (92.8)	921 (94.2)	98 (32.3)
N partial (%)	14 (1.48)	16 (1.94)	16 (5.28)
N missing (%)	56 (5.73)	38 (3.89)	189 (62.8)

Re-use potential

These reference transcriptomes were assembled to facilitate sponge holobiont research aimed at exploring how both host and symbionts respond to changing environmental conditions. The transcriptomes can be used for studies involving Tag-based RNAseq (TagSeq) [67], a highly accurate [68] and cost-effective sequencing technique for large sample sets. Output files are also formatted for Rank-based Gene Ontology analysis of gene expression data (GO_MWU, [69]), and for Functional Summary and Meta-Analysis of Gene Expression Data (KOGMWU, [70]).

Availability of supporting data

All data, including raw reads, can be accessed here:

https://www.dropbox.com/sh/82ue5l16n4xzxww/AABENUi-Cdbm_z-6x4Gj3qICa?dl=0. Raw data has also

209 been deposited on NCBI's SRA under accession numbers PRJNA639714 and PRJNA639798 for the *C.*

210 *orientalis* holobiont and *C. foliascens*, respectively.

211 **Declarations**

212 **List of abbreviations**

213 BUSCO: Benchmarking Universal Single-Copy Orthologs; KEGG: Kyoto Encyclopedia of Genes and Genomes;

214 KOG: EuKaryotic Orthologous Groups; TagSeq: Tag-based RNAseq.

215 **Ethics notes and consent for publication**

216 Not applicable.

217 **Competing interests**

218 The authors declare that they have no competing interests.

219 **Funding**

220 This research was funded by the Western Australian Marine Science Institution (WAMSI) as part of the

221 WAMSI Dredging Science Node, and made possible through investment from Chevron Australia, Woodside

222 Energy Limited, BHP Billiton as environmental offsets and by co-investment from the WAMSI Joint Venture

223 partners. The views expressed herein are those of the authors and not necessarily those of WAMSI. Brian

224 W. Strehlow was supported by a University of Western Australia (UWA) Scholarship for International

225 Research Fees, University International Stipend, and UWA Safety-Net Top-Up Scholarships. Brian W.

226 Strehlow was further supported by the Villum Investigator Grant awarded to Don Canfield (No. 16518). The

227 funders had no role in study design, data collection and analysis, decision to publish, or preparation of the

228 manuscript.

229 **Authors' contributions**

230 BWS, MCP, CDK, AD, MR, PC, and NSW conceived and designed the experiments; BWS, MCP, and CDK
 231 performed the experiments and laboratory work; BWS performed bioinformatics analyses with extensive
 232 help and training from CDK and PL. BWS wrote the first draft. BWS, MCP, CDK, PL, AD, MR, PC, and NSW
 233 contributed to revisions of various drafts and approved the final manuscript.

234 **Acknowledgements**

235 The authors are grateful to E. Botté and G. Millar for their assistance in technical aspects related to
 236 computing. They also thank the staff at the AIMS National Sea Simulator for their expertise and assistance
 237 in the tank-based experiments.

References

1. Feuda R, Dohrmann M, Pett W, Philippe H, Rota-Stabelli O, Lartillot N, et al. Improved Modeling of Compositional Heterogeneity Supports Sponges as Sister to All Other Animals. *Curr Biol.* 2017;27:3864-3870.e4.
2. IPCC. Climate Change 2014 Synthesis Report. Contrib Work Groups I, II III to Fifth Assess Rep Intergov Panel Clim Chang. 2014;1–112.
3. Bell JJ, Davy SK, Jones T, Taylor MW, Webster NS. Could some coral reefs become sponge reefs as our climate changes? *Glob Chang Biol.* 2013;19:2613–24. Available from: <http://doi.wiley.com/10.1111/gcb.12212>
4. Kroon FJ, Kuhnert PM, Henderson BL, Wilkinson SN, Kinsey-Henderson A, Abbott B, et al. River loads of suspended solids, nitrogen, phosphorus and herbicides delivered to the Great Barrier Reef lagoon. *Mar Pollut Bull. Elsevier Ltd;* 2012;65:167–81. Available from: <http://dx.doi.org/10.1016/j.marpolbul.2011.10.018>
5. Syvitski JPM, Vorosmarty CJ, Kettner AJ, Green P. Impact of Humans on the Flux of Terrestrial Sediment to the Global Coastal Ocean. *Science* (80). 2005;308:1364–8.
6. Maina J, de Moel H, Zinke J, Madin J, McClanahan T, Vermaat JE. Human deforestation outweighs future climate change impacts of sedimentation on coral reefs. *Nat Commun.* 2013;4:1–7. Available from: <http://www.nature.com/doi/10.1038/ncomms2986>
7. Stender Y, Jokiel PL, Rodgers KS. Thirty years of coral reef change in relation to coastal construction and increased sedimentation at Pele Kane Bay, Hawaii. *PeerJ.* 2014;2:e300. Available from: <https://peerj.com/articles/300>
8. Van Soest RW., Boury-Esnault N, Hooper JNA, Rützler K, de Voogd NJ, Alvarez de Glasby B, et al. World Porifera database. Accessed at <http://www.marinespecies.org/porifera>. 2020;
9. Manousaki T, Koutsouveli V, Lagnel J, Kollias S, Tsigenopoulos CS, Arvanitidis C, et al. A de novo transcriptome assembly for the bath sponge *Spongia officinalis*, adjusting for microsymbionts. *BMC Res Notes. BioMed Central;* 2019;12:10–2. Available from: <https://doi.org/10.1186/s13104-019-4843-6>
10. Belahbib H, Renard E, Santini S, Jourda C, Claverie J-M, Borchellini C, et al. New Genomic Data and Analyses Challenge the Traditional Vision of Animal Epithelium Evolution. *BMC Genomics. BMC Genomics;* 2018;1–15.
11. Leininger S, Adamski M, Bergum B, Guder C, Liu J, Laplante M, et al. Developmental gene expression provides clues to relationships between sponge and eumetazoan body plans. *Nat Commun.* 2014;5:1–15.
12. Qiu F, Ding S, Ou H, Wang D, Chen J, Miyamoto MM. Transcriptome changes during the life cycle of the red sponge, *Mycale phyllophila* (Porifera, Demospongiae, Poecilosclerida). *Genes (Basel).* 2015;6:1023–52.
13. Kitchen SA, Crowder CM, Poole AZ, Weis VM, Meyer E. De Novo Assembly and Characterization of Four Anthozoan (Phylum Cnidaria) Transcriptomes. *G3 (Bethesda).* 2015;5:2441–52. Available from: http://apps.webofknowledge.com.gate1.inist.fr/full_record.do?product=UA&search_mode=MarkedList&qid=7&SID=T1OpnZdVDNlGAFI4yu&page=1&doc=1&colName=WOS
14. Riesgo A, Andrade SCS, Sharma PP, Novo M, Pérez-Porro AR, Vahtera V, et al. Comparative description of ten transcriptomes of newly sequenced invertebrates and efficiency estimation of genomic sampling in non-model taxa. *Front Zool.* 2012;9:1–24.

- 278 15. Riesgo A, Farrar N, Windsor PJ, Giribet G, Leys SP. The analysis of eight transcriptomes from all poriferan
279 classes reveals surprising genetic complexity in sponges. *Mol Biol Evol.* 2014;31:1102–20.
- 280 16. Ryu T, Seridi L, Moitinho-Silva L, Oates M, Liew YJ, Mavromatis C, et al. Hologenome analysis of two
281 marine sponges with different microbiomes. *BMC Genomics.* 2016;17:1–11. Available
282 from: <http://dx.doi.org/10.1186/s12864-016-2501-0>
- 283 17. Riesgo A, Peterson K, Richardson C, Heist T, Strehlow B, McCauley M, et al. Transcriptomic analysis of
284 differential host gene expression upon uptake of symbionts: a case study with *Symbiodinium* and the major
285 bioeroding sponge *Cliona varians*. *BMC Genomics.* 2014;15:376. Available from:
286 <http://www.ncbi.nlm.nih.gov/pubmed/24885832>
- 287 18. Conaco C, Neveu P, Zhou H, Arcila ML, Degnan SM, Degnan BM, et al. Transcriptome profiling of the
288 demosponge *Amphimedon queenslandica* reveals genome-wide events that accompany major life cycle
289 transitions. *BMC Genomics.* 2012;13.
- 290 19. Díez-Vives C, Moitinho-Silva L, Nielsen S, Reynolds D, Thomas T. Expression of eukaryotic-like protein in
291 the microbiome of sponges. *Mol Ecol.* 2017;26:1432–51.
- 292 20. Ereskovsky A V., Richter DJ, Lavrov D V., Schippers KJ, Nichols SA. Transcriptome sequencing and
293 delimitation of sympatric *Oscarella* species (*O. carmela* and *O. pearsei* sp. nov) from California, USA. *PLoS*
294 *One.* 2017;12:1–25. Available from: <http://dx.doi.org/10.1371/journal.pone.0183002>
- 295 21. Fiore CL, Labrie M, Jarett JK, Lesser MP. Transcriptional activity of the giant barrel sponge, *Xestospongia*
296 *muta* Holobiont: Molecular evidence for metabolic interchange. *Front Microbiol.* 2015;6.
- 297 22. Germer J, Cerveau N, Jackson DJ. The Holo-Transcriptome of a Calcified Early Branching Metazoan.
298 *Front Mar Sci.* 2017;4. Available from: <http://journal.frontiersin.org/article/10.3389/fmars.2017.00081/full>
- 299 23. Guzman C, Conaco C. Comparative transcriptome analysis reveals insights into the streamlined
300 genomes of haplosclerid demosponges. *Sci Rep.* Nature Publishing Group; 2016;6:18774. Available from:
301 <http://www.nature.com/articles/srep18774>
- 302 24. Guzman C, Conaco C. Gene expression dynamics accompanying the sponge thermal stress response.
303 *PLoS One.* 2016;11:1–15. Available from: <http://dx.doi.org/10.1371/journal.pone.0165368>
- 304 25. Kenny NJ, de Goeij JM, de Bakker DM, Whalen CG, Berezikov E, Riesgo A. Towards the identification of
305 ancestrally shared regenerative mechanisms across the Metazoa: A Transcriptomic case study in the
306 Demosponge *Halisarca caerulea*. *Mar Genomics.* Elsevier; 2018;37:135–47. Available from:
307 <https://doi.org/10.1016/j.margen.2017.11.001>
- 308 26. Nichols SA, Roberts BW, Richter DJ, Fairclough SR, King N. Origin of metazoan cadherin diversity and the
309 antiquity of the classical cadherin-catenin complex. *Proc Natl Acad Sci.* 2012;109:13046–51. Available from:
310 <http://www.pnas.org/cgi/doi/10.1073/pnas.1120685109>
- 311 27. Francis WR, Eitel M, Vargas S, Adamski M, Haddock SHD, Krebs S, et al. The genome of the contractile
312 demosponge *Tethya wilhelma* and the evolution of metazoan neural signalling pathways. *bioRxiv.*
313 2017;preprint. Available from: <https://www.biorxiv.org/content/biorxiv/early/2017/04/12/120998.full.pdf>
- 314 28. Srivastava M, Simakov O, Chapman J, Fahey B, Gauthier M, Mitros T, et al. The *Amphimedon*
315 *queenslandica* genome and the evolution of animal complexity. *Nature.* 2010;466. Available from:
316 <http://dx.doi.org/10.1038/nature09201>
- 317 29. Fortunato SA V, Adamski M, Ramos OM, Leininger S, Liu J, Ferrier DEK, et al. Calcisponges have a
318 ParaHox gene and dynamic expression of dispersed NK homeobox genes. *Nature.* Nature Publishing Group;

2014;514:620–3. Available from: <http://dx.doi.org/10.1038/nature13881>

30. Abdul Wahab MA, Maldonado M, Luter HM, Jones R, Ricardo G. Effects of sediment resuspension on the larval stage of the model sponge *Carteriospongia foliascens*. *Sci Total Environ*. Elsevier B.V; 2019;695:133837. Available from: <https://doi.org/10.1016/j.scitotenv.2019.133837>

31. Bennett HM, Altenrath C, Woods L, Davy SK, Webster NS, Bell JJ. Interactive effects of temperature and pCO₂ on sponges: from the cradle to the grave. *Glob Chang Biol*. 2016;1–16. Available from: <http://doi.wiley.com/10.1111/gcb.13474>

32. Pineda M-C, Strehlow B, Duckworth A, Doyle J, Jones R, Webster NS. Effects of light attenuation on the sponge holobiont- implications for dredging management. *Sci Rep*. Nature Publishing Group; 2016;6:1–13. Available from: <http://dx.doi.org/10.1038/srep39038>

33. Pineda M-C, Strehlow B, Kamp J, Duckworth A, Jones R, Webster NS. Effects of combined dredging-related stressors on sponges: A laboratory approach using realistic scenarios. *Sci Rep*. 2017;7:1–14.

34. Pineda M-C, Strehlow B, Sternel M, Duckworth A, Haan J den, Jones R, et al. Effects of sediment smothering on the sponge holobiont with implications for dredging management. *Sci Rep*. Springer US; 2017;7:1–15. Available from: <http://dx.doi.org/10.1038/s41598-017-05243-x>

35. Pineda M-C, Strehlow B, Sternel M, Duckworth A, Jones R, Webster NS. Effects of suspended sediments on the sponge holobiont with implications for dredging management. *Sci Rep*. Springer US; 2017;7:1–15. Available from: <http://dx.doi.org/10.1038/s41598-017-05241-z>

36. Ramsby BD, Hoogenboom MO, Smith HA, Whalan S, Webster NS. The bioeroding sponge *Cliona orientalis* will not tolerate future projected ocean warming. *Sci Rep*. Springer US; 2018;1–13. Available from: www.nature.com/scientificreports

37. Ramsby BD, Hoogenboom MO, Whalan S, Webster NS. Elevated seawater temperature disrupts the microbiome of an ecologically important bioeroding sponge. *Mol Ecol*. 2018;27:2124–37.

38. Ramsby BD, Hill MS, Thornhill DJ, Steenhuizen SF, Achlatis M, Lewis AM, et al. Sibling species of mutualistic *Symbiodinium* clade G from bioeroding sponges in the western Pacific and western Atlantic oceans. *J Phycol*. 2017;53:951–60.

39. LaJeunesse TC, Parkinson JE, Gabrielson PW, Jeong HJ, Reimer JD, Voolstra CR, et al. Systematic Revision of Symbiodiniaceae Highlights the Antiquity and Diversity of Coral Endosymbionts. *Curr Biol*. Elsevier Ltd.; 2018;28:2570-2580.e6. Available from: <https://doi.org/10.1016/j.cub.2018.07.008>

40. Bessell-Browne P, Negri AP, Fisher R, Clode PL, Jones R. Impacts of turbidity on corals: the relative importance of light limitation and suspended sediments. *Mar Pollut Bull*. 2017;1–24.

41. Bessell-Browne P, Fisher R, Duckworth A, Jones R. Mucous sheet production in *Porites*: an effective bioindicator of sediment related pressures. *Ecol Indic*. Elsevier Ltd; 2017;77:276–85.

42. Field ME, Chezar H, Storlazzi CD. SedPods: A low-cost coral proxy for measuring net sedimentation. *Coral Reefs*. 2013;32:155–9.

43. Riesgo A, Pérez-Porro AR, Carmona S, Leys SP, Giribet G. Optimization of preservation and storage time of sponge tissues to obtain quality mRNA for next-generation sequencing. *Mol Ecol Resour*. 2012;12. Available from: <http://dx.doi.org/10.1111/j.1755-0998.2011.03097.x>

44. Matz M V. Assembly and annotation of transcriptomes de novo. Github. 2015;URL:<https://github.com/z0on/annotatingTranscriptomes>. Available from: <https://github.com/z0on/annotatingTranscriptomes>

360 45. Meyer E. A collection of scripts for analyzing transcriptome assemblies. Github.
361 2016;URL:https://github.com/Eli-Meyer/transcriptome_utilities. Available from: [https://github.com/Eli-](https://github.com/Eli-Meyer/transcriptome_utilities)
362 Meyer/transcriptome_utilities

363 46. Kenkel CD, Bay LK. Novel transcriptome resources for three scleractinian coral species from the Indo-
364 Pacific. *Gigascience*. 2017;6:1–4.

365 47. The Hannon Lab. FASTX-Toolkit. 2010;URL:http://hannonlab.cshl.edu/fastx_toolkit/. Available from:
366 http://hannonlab.cshl.edu/fastx_toolkit/

367 48. Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, et al. Full-length transcriptome
368 assembly from RNA-Seq data without a reference genome. *Nat Biotechnol*. 2011;29:644–52. Available
369 from: <http://www.nature.com/doi/10.1038/nbt.1883>

370 49. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *J Mol Biol*.
371 1990;215:403–10. Available from: <http://www.sciencedirect.com/science/article/pii/S0022283605803602>

372 50. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal RNA gene
373 database project: Improved data processing and web-based tools. *Nucleic Acids Res*. 2013;41.

374 51. Fernandez-Valverde SL, Calcino AD, Degnan BM. Deep developmental transcriptome sequencing
375 uncovers numerous new genes and enhances gene annotation in the sponge *Amphimedon queenslandica*.
376 *BMC Genomics*. 2015;16:387. Available from:
377 [http://apps.webofknowledge.com.gate1.inist.fr/full_record.do?product=UA&search_mode=GeneralSearch](http://apps.webofknowledge.com.gate1.inist.fr/full_record.do?product=UA&search_mode=GeneralSearch&qid=11&SID=T10pnZdVDNILGAfi4yu&page=2&doc=13&cacheurlFromRightClick=no)
378 [&qid=11&SID=T10pnZdVDNILGAfi4yu&page=2&doc=13&cacheurlFromRightClick=no](http://apps.webofknowledge.com.gate1.inist.fr/full_record.do?product=UA&search_mode=GeneralSearch&qid=11&SID=T10pnZdVDNILGAfi4yu&page=2&doc=13&cacheurlFromRightClick=no)

379 52. Lin S, Cheng S, Song B, Zhong X, Lin X, Li W, et al. The *Symbiodinium kawagutii* genome illuminates
380 dinoflagellate gene expression and coral symbiosis. *Science* (80-). 2015;350:691–4. Available from:
381 <http://www.sciencemag.org/cgi/doi/10.1126/science.aad0408>

382 53. Gene Ontology Consortium. The Gene Ontology Consortium. Gene ontology: tool for the unification of
383 biology. *Nat Genet*. 2000;25:25–9.

384 54. Dixon GB. Genomic determinants of coral heat tolerance across latitudes. *Science* (80-). 2013;353:1689–
385 99.

386 55. The UniProt Consortium. UniProt: the universal protein knowledgebase. *Nucleic Acids Res*.
387 2017;43:D2158–69. Available from: <http://nar.oxfordjournals.org/cgi/content/long/43/D1/D204>

388 56. Matz M V, Aglyamova G V. Cost-efficient genome-wide gene expression profiling. Github.
389 2017;URL:https://github.com/z0on/tag-based_RNAseq.

390 57. Huerta-Cepas J, Szklarczyk D, Forslund K, Cook H, Heller D, Walter MC, et al. EGGNOG 4.5: A hierarchical
391 orthology framework with improved functional annotations for eukaryotic, prokaryotic and viral sequences.
392 *Nucleic Acids Res*. 2016;

393 58. Huerta-Cepas J, Forslund K, Coelho LP, Szklarczyk D, Jensen LJ, Von Mering C, et al. Fast genome-wide
394 functional annotation through orthology assignment by eggNOG-mapper. *Mol Biol Evol*. 2017;

395 59. Moriya Y, Itoh M, Okuda S, Yoshizawa AC, Kanehisa M. KAAS: An automatic genome annotation and
396 pathway reconstruction server. *Nucleic Acids Res*. 2007;35.

397 60. Bushnell B. BMap: a fast, accurate, splice-aware aligner. *Jt. Genome Institute, Dep. energy*. 2014.

398 61. Simão FA, Waterhouse RM, Ioannidis P, Kriventseva E V., Zdobnov EM. BUSCO: Assessing genome
399 assembly and annotation completeness with single-copy orthologs. *Bioinformatics*. 2015;31:3210–2.

62. Nishimura O, Hara Y, Kuraku S. gVolante for standardizing completeness assessment of genome and transcriptome assemblies. *Bioinformatics*. 2017; <https://gvolante.riken.jp/analysis.html>.
63. López-Maury L, Marguerat S, Bähler J. Tuning gene expression to changing environments: from rapid responses to evolutionary adaptation. *Nat Rev Genet*. 2008;9:583–93. Available from: <http://www.nature.com/doifinder/10.1038/nrg2398>
64. Ladner JT, Barshis DJ, Palumbi SR. Protein evolution in two co-occurring types of Symbiodinium: An exploration into the genetic basis of thermal tolerance in Symbiodinium clade D. *BMC Evol Biol*. 2012;12.
65. Karimi E, Ramos M, Gonçalves JMS, Xavier JR, Reis MP, Costa R. Comparative metagenomics reveals the distinctive adaptive features of the *Spongia officinalis* endosymbiotic consortium. *Front Microbiol*. 2017;8.
66. Liu H, Stephens TG, González-Pech RA, Beltran VH, Lapeyre B, Bongaerts P, et al. *Symbiodinium* genomes reveal adaptive evolution of functions related to coral-dinoflagellate symbiosis. *Commun Biol*. Springer US; 2018;1:1–11. Available from: <http://dx.doi.org/10.1038/s42003-018-0098-3>
67. Meyer E, Aglyamova G V., Matz M V. Profiling gene expression responses of coral larvae (*Acropora millepora*) to elevated temperature and settlement inducers using a novel RNA-Seq procedure. *Mol Ecol*. 2011;20:3599–616.
68. Lohman BK, Weber JN, Bolnick DI. Evaluation of TagSeq, a reliable low-cost alternative for RNAseq. *Mol Ecol Resour*. 2016;
69. Matz M V. Rank-based Gene Ontology analysis of gene expression data. 2015; Available from: https://github.com/z0on/GO_MWU
70. Matz M V. KOGMWU: Functional Summary and Meta-Analysis of Gene Expression Data. 2019; Available from: <https://cran.r-project.org/package=KOGMWU>