Hierarchical effects of choice-related activity and neural encoding during feature selective attention

Jennifer L. Mohn^{1,2}, Joshua D. Downer^{1,3}, Kevin N. O'Connor^{1,2}, Jeffrey S. Johnson^{1,2}, and Mitchell L. Sutter^{1,2}

¹ Center for Neuroscience; University of California, Davis; Davis, CA 95618; USA

² Department of Neurobiology, Physiology and Behavior; University of California, Davis; Davis, CA 95618; USA

³ Current address: Dept. of Otolaryngology – Head and Neck Surgery; University of California, San Francisco; San Francisco, CA, 94158; USA

Corresponding Author:

Mitchell L. Sutter

Center for Neuroscience

University of California, Davis

1544 Newton Court Davis CA 95618

mlsutter@ucdavis.edu

No. of pages: 35

No. of Figures (Main): 6

No. of Tables: 1

No. of words: Abstract (222); Introduction (962); Results (1127) Discussion (1479)

Conflict of Interest: The authors declare no competing financial interest

Acknowledgements: This work was funded by NIH NIDCD grant DC002514 (MLS), NSF GRFP 1148897 (JDD) and ARCS Foundation Fellowship (JDD)

Abstract

Selective attention is necessary to sift through, form a coherent percept of, and make behavioral decisions on the vast amount of information present in most sensory environments. How and where selective attention is employed in cortex and how this perceptual information then informs the relevant behavioral decisions is still not well understood. Studies probing selective attention and decision making in visual cortex have been enlightening as to how sensory attention might work in that modality; whether or not similar mechanisms are employed in auditory attention is not yet clear. Therefore, we trained rhesus macaques on a feature selective attention task, where they switched between reporting changes in temporal (amplitude modulation, AM) and spectral (carrier bandwidth) features of a broadband noise stimulus. We investigated how the encoding of these features by single neurons in primary (A1) and secondary (lateral belt, ML) auditory cortex were affected by the different attention conditions. We found that neurons in A1 and ML showed mixed-selectivity to the sound and task features. We found no difference in AM encoding between the attention conditions. We found that choice-related activity in both A1 and ML neurons shifts between attentional conditions. This finding suggests that choice-related activity in auditory cortex does not simply reflect motor preparation or action, and supports the relationship between reported choice-related activity and the decision and perceptual process.

New & Noteworthy

We recorded from primary and secondary auditory cortex while monkeys performed a non-spatial feature attention task. Both areas exhibited rate-based choice-related activity. The manifestation of choice-related activity was attention-dependent, suggesting that choice-related activity in auditory cortex does not simply reflect arousal or motor influences, but relates to the specific perceptual choice. The lack of temporal-based choice activity is consistent with growing evidence that subcortical, but not cortical, single neurons inform decisions through temporal envelope following.

Introduction

The auditory system is often faced with the difficult challenge of encoding a specific sound in a noisy environment, such as following a conversation in a loud room. The neural mechanisms by which the auditory system attends to one sound source and ignores distracting sounds are not yet understood. Studies probing the mechanisms underlying auditory attention in cortex have been largely concerned with task engagement, wherein the effects of active performance on neural activity is compared to those of passive listening. Studies in auditory cortex (AC) utilizing this paradigm have shown that task engagement can improve behaviorally-relevant neural sound discrimination (Atiani et al. 2014; Bagur et al. 2018; Buran et al. 2014; Carcea et al. 2017; Francis et al. 2018a; Niwa et al. 2012a, 2015; von Trapp et al. 2016), modulate neuronal tuning (Fritz et al. 2003, 2007; Fritz 2005; Lee and Middlebrooks 2011; Lin et al. 2019; Yin et al. 2014), alter the structure of correlated variability within neural populations (Downer et al. 2015, 2017a), and more (Massoudi et al. 2014; Angeloni and Geffen 2018; Osmanski and Wang 2015; Sutter and Shamma 2011). Though informative, this active/passive paradigm makes it difficult to disentangle arousal and motor effects from the mechanisms more specifically employed in selectively attending to a single sound source or feature amidst auditory ‘clutter’.

Studies of the neural basis of auditory selective attention at the single neuron level are rare (Schwartz and David 2018), and non-spatial auditory feature-selective attention has been relatively unexplored (Downer et al. 2017b). Feature-selective attention, which segregates particular sound features, such as intensity or fundamental frequency, is essential for tasks such as discriminating between talkers in a noisy environment (Bregman 1994; McDermott 2009; Bizley and Cohen 2013; Shinn-Cunningham 2008; Woods and McDermott 2015). Furthermore, it can prove useful for listeners to switch between attended sound features because the most distinctive feature dimensions may vary across sources (Woods and McDermott 2015; Bregman 1994).

In visual cortex, feature-based attention has been suggested to follow a gain model similar to spatial attention, where responsivity to the attended feature increases in cells tuned to the attended feature and decreases in cells tuned to orthogonal features (Martinez-Trujillo and Treue 2004; Maunsell and Treue 2006). Studies of spatial attention in AC single neurons suggest that AC employs a mechanism similar to that reported in visual cortex, where a gain in neural activity increases when attention is directed into the receptive field of a neuron and, conversely, gain decreases when attention is directed outside the receptive field (Engle and Recanzone 2013; Lee and Middlebrooks 2011; Scott et al. 2007) . We endeavored to see if feature-selective attention in AC is also facilitated by a gain in activity in neurons tuned to an attended feature.

How and where task relevant sensory information is transformed into a decision in the brain is still largely unclear. There have been mixed reports of activity correlated to the reported decision in AC (Christison-Lagay et al. 2017; Elgueda et al. 2019; Guo et al. 2019; Niwa et al. 2012b; Runyan et al. 2017; Tsunada et al. 2016; Tsunada and Cohen 2014). This choice-related activity has been reported in some studies as early as primary auditory cortex (A1) (Atiani et al. 2014; Bathellier et al. 2012; Bizley et al. 2013; Christison-Lagay et al. 2017; Christison-Lagay and Cohen 2018; Francis et al. 2018b, 2018b; Gronskaia and von der Behrens 2019; Huang et al. 2019; Niwa et al. 2012b). As one progresses further along the auditory cortical hierarchy, there is either an increasingly larger proportion of neurons showing activity correlated to the decision, or the nature of the choice signal changes (Atiani et al. 2014; Niwa et al. 2013; Tsunada et al. 2016). Both cases suggest that the sensory evidence informing task-relevant decisions is transformed as the information moves up the processing stream (Bizley and Cohen 2013; Hackett 2011; Huang and Brosch 2020; Romanski et al. 1999).

There has also been uncertainty as to whether the reported choice activity in AC could be more reflective of motor influences than perceptual or decision-related influences. Go/No-Go tasks are typically used in auditory cortical studies, and these tasks require movement for report of one choice,

but not the other (Brosch 2005; Niwa et al. 2013); forced-choice tasks reduce this uncertainty by requiring movements for either report (Guo et al. 2019). It has been well documented that movement can modulate auditory cortical activity (Eliades and Wang 2003; Guo et al. 2019; Schneider et al. 2014). Here, we employ a Yes/No forced-choice task format in which a movement is required for both responses in order to disentangle motor-related from choice-related activity in AC.

We investigated whether a mechanism for feature-selective attention similar to feature-based attention in visual cortex is employed in primary (A1) and secondary (middle lateral belt, ML) auditory cortex using noise that was amplitude modulated (AM) or bandwidth restricted (Δ BW). Monkeys were presented sounds that varied either in spectral (Δ BW) or temporal (AM) dimensions, or both, and performed a detection task in which they reported change along one of these feature dimensions. In this study, we focus on the amplitude modulation feature, as it has been well studied and is a salient communicative sound feature for humans and other animals (Schnupp 2006; Shannon et al. 1995; Van Tasell et al. 1987; Wang et al. 2007) and can be helpful in sound source segregation (Bregman 1994; Grimault et al. 2002). Spectral content changes were used as a difficulty-matched attentional control. We hypothesized we would see a gain in AM encoding when animals were cued to attend to that feature, compared to when they were cued to attend Δ BW changes. We also examined choice-related activity in AC, hypothesizing to find a larger proportion of neurons with significant choice-related activity in higher-order AC (ML) than in A1.

Materials and Methods

Subjects.

Subjects were two adult rhesus macaques, one male (13kg, 14-16 years old), one female (7kg, 17-19 years old). All procedures were approved by the University of California–Davis Animal Care and

Use Committee and met the requirements of the United States Public Health Service policy on experimental animal care.

Stimuli.

Stimuli were constructed from broadband Gaussian (white) noise bursts (400 ms; 5 ms cosine ramped), 9 octaves in width (40 to 20480 Hz). Four different seeds were used to create the carrier noise, which was frozen across trials. To introduce variance along spectral and temporal dimensions, the spectral bandwidth of the noise was narrowed (ΔBW) and/or the noise envelope was sinusoidally amplitude modulated (AM). The extent of variation in each dimension was manipulated to measure behavioral and neural responses above and below threshold for detecting each feature.

Sound generation methods have been previously reported (O'Connor et al., 2011). Briefly, sound signals were produced using an in-house MATLAB program and a digital-to-analog converter (Cambridge Electronic Design [CED] model 1401). Signals were attenuated (TDT Systems PA5, Leader LAT-45), amplified (RadioShack MPA-200), and presented from a single speaker (RadioShack PA-110) positioned approximately 1.5 m in front of the subject centered at the interaural midpoint. Sounds were generated at a 100 kHz sampling rate. Intensity was calibrated across all sounds (Bruel & Kjaer model 2231) to 65 dB at the outer ear. It is important to note that some methods of generating ΔBW introduce variation in that sound's envelope, however we implemented a synthesis method that constructs noise using a single-frequency additive technique and thus avoids introducing envelope variations that could serve as cues for ΔBW (Strickland and Viemeister 1997).

Recording procedures.

Each animal was implanted with a head post centrally behind the brow ridge and a recording cylinder over an 18 mm craniotomy over the parietal lobe using aseptic surgical techniques (O'Connor et

al. 2005). Placement of the craniotomy was based on stereotactic coordinates of auditory cortex to allow vertical access through parietal cortex to the superior temporal plane (Saleem and Logothetis 2007).

All recordings took place in a sound attenuating, foam-lined booth (IAC: 2.9x3.2x2 meters) while subjects sat in an acoustically transparent chair (Crist Instruments). Three quartz-coated tungsten microelectrodes (Thomas Recording, 1–2 MΩ; 0.35 mm horizontal spacing; variable, independently manipulated vertical spacing) were advanced vertically to the superior surface of the temporal lobe. Extracellular signals were amplified (AM Systems model 1800), bandpass filtered between 0.3 Hz and 10 kHz (Krohn-Hite 3382), and then converted to a digital signal at a 50 kHz sampling rate (CED model 1401). During electrode advancement, auditory responsive neurons were isolated by presenting various sounds while the subject sat passively. When at least one auditory responsive single unit was well isolated, we measured neural responses to the two features while the subjects sat passively awake. At least 10 repetitions of each of the following stimuli were presented: the unmodulated noise, each level of bandwidth restriction, and each of the possible AM test modulation frequencies (described below). We also measured pure tone and bandpass noise tuning to aid in distinguishing area boundaries. After completing these tuning measures, experimental behavioral testing and recording began. When possible, tuning responses to the tested stimuli were again measured after task performance, to ensure stability of electrodes throughout the recording. Contributions of single units (SUs) to the signal were determined offline using principal components analysis-based spike sorting tools from Spike2 (CED). Spiking activity was at least 4–5 times the background noise level. Fewer than 0.1% of spike events assigned to single unit clusters fell within a 1 ms refractory period window. Only recordings in which neurons were well isolated for at least 180 trials within each condition were included in analysis here.

Cortical field assessment.

Recording locations were determined using both stereotactic coordinates (Martin and Bowden 1996) and established physiological measures (Merzenich and Brugge 1973; Rauschecker and Tian 2000; Tian and Rauschecker 2004). In each animal, we mapped characteristic frequency (CF) and sharpness of bandpass noise tuning to establish a topographic distribution of each. Tonotopic gradient reversal, BW distribution, spike latency and response robustness to pure tones was used to estimate the boundary between A1 and ML and assign single units to an area (Downer et al. 2017a; Niwa et al. 2015). Recordings were assigned to their putative cortical fields *post hoc* using recording location, tuning preferences, and latencies.

Feature attention task.

This feature attention task has been previously described in detail (Downer et al. 2017b). The subjects performed a change detection task in which only changes in the attended feature were relevant for the task. Subjects moved a joystick laterally to initiate a trial, wherein an initial sound (the S1, always the 9-octave-wide broadband, unmodulated noise) was presented, followed by a second sound (S2) after a 400ms inter-stimulus interval (ISI). The S2 could be identical to the S1, it could change by being amplitude-modulated (AM), it could change by being bandwidth restricted (Δ BW), or it could change along both feature dimensions.

Only three values of each feature (AM, Δ BW) were presented, limiting the size of the stimulus set in order to obtain reasonable power for neural data analysis. The stimulus space was further reduced by presenting only a subset of the possible co-varying stimuli. Within each recording session, we presented 13 total stimuli. To equilibrate difficulty between the two features, we presented values of each feature so that one was near threshold, one was slightly above, and one far above threshold.

Thresholds were determined for each feature independently for each subject using a range of six levels for each feature before three feature values for each animal were selected and the co-varying feature attention task began. For Monkey U, the ΔBW values were 0.375, 0.5, and 1 octave and the AM depth values were 28%, 40%, and 100%. For Monkey W, the ΔBW values were 0.5, 0.75, and 1.5 octaves and the AM depth values were 40%, 60%, and 100%.

For all analyses in which data are collapsed across subjects, ΔBW values and AM values are presented as ranks (ΔBW 0-3 and AM 0-3) (e.g., Fig. 1). Within a given session, AM sounds were presented at only a single modulation frequency. Across sessions, a small set of frequencies was used (15, 22, 30, 48, and 60 Hz). The AM frequency was selected randomly each day. Subjects were cued visually via an LED above the speaker as to which feature to attend (green or red light, counterbalanced between subjects). Additionally, each block began with 60 “instruction” trials in which the S2s presented were only altered along the attended feature dimension (i.e., sounds containing the distractor feature were not presented). Subjects were to respond with a “yes” (up or down joystick movement, counterbalanced across subjects) on any trial in which the attended feature was presented, otherwise, the correct response was “no” (opposite joystick movement). We chose upward or downward joystick movement to avoid influences on single neuron choice activity dependent on contralateral movements. Such movement related activity has been recently reported in other studies (Guo et al. 2019). Hits and correct rejections were rewarded with a drop of water or juice and misses and false alarms resulted in a penalty (3–5 s timeout).

During the test conditions, the S2 was unmodulated broadband noise (no change from S1) on 25% of the trials, co-varying on 25% of the trials, and contained only ΔBW or AM on 25% of the trials respectively. Sounds in the set were presented pseudo-randomly such that, over sets of 96 trials, the entire stimulus set was presented exhaustively (including all four random noise seeds). Block length was variable, based in part on subjects’ performance, to ensure a sufficient number of correct trials for each

stimulus. Not including instruction trials, block length was at least 180 trials and at most 360 trials, to ensure that subjects performed in each attention condition at least once during the experiment. Subjects could perform each attention condition multiple times within a session. Only sessions in which subjects completed at least 180 trials per condition (excluding instruction trials) were considered for analysis in this study.

Analysis of single neuron feature selectivity

Neurons' firing rate responses across the range of values were calculated to derive a firing rate function for each feature. Functions were categorized based on whether firing rates increased as the level of the feature increased or decreased ('increasing' vs. 'decreasing' functions). Spike counts (SC) were calculated over the entire 400ms stimulus window. SCs in response to feature-present stimuli were normalized over the entire spike count distribution across both features, including unmodulated noise, for that cell. To characterize this response function, we calculated a feature-selectivity index (FSI) for each feature as follows:

$$[1] \quad FSI_{AM} = \frac{SC_{AM>0, \Delta BW_0} - SC_{AM_0, \Delta BW_0}}{SC_{AM>0, \Delta BW_0} + SC_{AM_0, \Delta BW_0}}$$

$$[2] \quad FSI_{BW} = \frac{SC_{\Delta BW>0, AM_0} - SC_{\Delta BW_0, AM_0}}{SC_{\Delta BW>0, AM_0} + SC_{\Delta BW_0, AM_0}}$$

Where SC_x is the mean SC in response to the given set of stimuli designated by the subscript. A Kruskal–Wallis rank-sum test was performed between distributions of SCs with the feature-present (feature level greater than 0) and those with the feature-absent (feature value of 0) to determine the significance of the FSI for each neuron. Cells that had a significant FSI for a given feature were categorized as encoding that feature.

Phase projected vector strength

Vector strength (VS) is a metric that describes the degree to which the neural response is phase-locked to the stimulus (Goldberg and Brown 1969; Mardia and Jupp 2000). VS is defined as:

$$[3] \quad VS = \frac{\sqrt{(\sum_{i=1}^n \cos \theta_i)^2 + (\sum_{i=1}^n \sin \theta_i)^2}}{n}$$

Where n is the number of spikes over all trials and θ_i is the phase of each spike, in radians, calculated by:

$$[4] \quad \theta_i = 2\pi \frac{t_i \bmod p}{p}$$

Where t_i is the time of the spike (in ms) relative to the onset of the stimulus and p is the modulation period of the stimulus (in ms). When spike count is low, VS has a tendency to report as spuriously high. Phase projected Vector Strength (VS_{pp}), is a variation on VS developed to help mitigate issues with low SC trials (Yin et al. 2011). VS_{pp} is calculated by first calculating VS for each trial, then the mean phase angle of each trial is compared to the mean phase angle of all trials, and the trial VS value is penalized if out of phase with the global mean response. VS_{pp} is defined as:

$$[5] \quad VS_{pp} = VS_t \cos(\phi_t - \phi_c)$$

Where VS_{pp} is the phase-projected vector strength per trial, VS_t is the vector strength per trial, as calculated in [1], and ϕ_t and ϕ_c are the trial-by-trial and mean phase angle in radians, respectively, calculated for each stimulus by:

$$[6] \quad \phi = \arctan2\left(\frac{\sum_{i=1}^n \sin \theta_i}{\sum_{i=1}^n \cos \theta_i}\right)$$

Where n is the number of spikes per trial (for ϕ_t) or across all trials (for ϕ_c). In this report, we use VS_{pp} exclusively to measure phase-locking, as SC tended to be relatively low and VS and VS_{pp} tend to be in good agreement with the exception of low SCs where VS_{pp} tends to be more accurate than VS (Yin et al. 2011). To determine significance of VS_{pp} encoding for each neuron, a Kruskal–Wallis rank-sum test was performed between distributions of VS_{pp} values on trials with non-zero AM depths, to those from unmodulated noise trials. Of note, when we refer to the VS_{pp} in response to an unmodulated stimulus,

this is a control measurement assuming the same modulation frequency as the corresponding AM frequency from that recording session.

Analysis of neural discriminability

We applied the signal detection theory-based metric area under the receiver operating characteristic (ROCa) (Green and Swets 1974) to measure how well neurons could detect each feature. ROCa represents the probability an ideal observer can detect the presence of the target feature given only a measure of the neural responses (either firing rate or VS_{pp}). To calculate ROCa, we partitioned the trial-by-trial neural responses into two distributions: those when the target feature was present in the stimulus and trials where it was absent. Then we determined the proportion of trials in each group where the neural response exceeds a criterion value. We repeated the measure using 100 criterion values, covering the whole range of responses. The plot of the probability of exceeding the criterion for feature-present trials (neural 'hits') versus the probability of exceeding the criteria for feature-absent trials (neural false alarms) plotted for all 100 criteria as separate points creates the ROC plot. The area under this curve is the ROCa. ROCa is bounded by 0 and 1, where both extremes indicate perfect discrimination between target feature-present and -absent stimuli, and 0.5 indicates a chance level of discrimination between the two distributions.

Analysis of choice-related activity

Choice probability (CP) is an application of ROC analysis used to measure the difference between neural responses contingent on what the animal reports, for example, whether a stimulus feature is present or absent (Britten et al. 1992, 1996). Similar to ROCa described above, CP values are bounded by 0 and 1, and a CP value of 0.5 indicates no difference (or perfect overlap) in the neural responses between 'feature-present' and 'feature-absent' reports. A CP value of 1 means for every trial

that the animal reports a feature, the neuron fired more than on trials where the animal did not report the feature. A CP value of 0 means that, for every trial that the animal reports a feature, the neuron fired *less* than on trials where the animal did not report the feature. Stimuli that did not have at least 5 ‘yes’ and 5 ‘no’ responses were excluded from analyses. CP was calculated based on both firing rate and on VS_{pp} . For rate-based CP, we calculated CP both for each stimulus separately, and pooled across stimuli. We calculated this stimulus-pooled CP by first separating the ‘yes’ and ‘no’ response trials within stimulus, then converting these rates into z-scores within a stimulus, then combined these z-scored responses across stimuli. This type of z-scoring has been found to be conservative in estimating CP (Kang and Maunsell 2012). CP was calculated during both the 400ms stimulus presentation (S2) and during the response window (RW), the time after stimulus offset and prior to the response (typically ~0.2 – 3s). The significance of each neuron’s CP was determined using a permutation test (Britten et al. 1996). The neural responses were pooled between the ‘feature-present report’ and ‘feature-absent report’ distributions and random samples were taken (without replacement). CP was then calculated from this randomly sampled set. This procedure was repeated 2000 times. The p value is the proportion of CP values from these randomly sampled repeats that were greater than the CP value from the non-shuffled distributions.

Results

We recorded activity from 92 single units in A1 (57 from Monkey W, 35 from Monkey U) from 33 recording sessions and 122 single units in ML (49 from Monkey W, 73 from Monkey U) over 39 recording sessions as animals performed a feature-selective attention task.

Feature tuning

There was no significant difference in the proportions of neurons in A1 and ML that encoded AM (47.8% A1, 38.5% ML; $p = 0.08$, χ^2 test). We found a large proportion of neurons in both A1 and ML that were sensitive to the relatively small changes in ΔBW from the 9-octave wide unmodulated noise, though there was no difference in the proportion of ΔBW encoding neurons between areas (32.6% A1, 29.5% ML; $p = 0.18$, χ^2 test) (Table 1).

A large population of neurons decreased firing rate for increasing AM depth ('decreasing cells') in both A1 and ML (Table 1). We also found that nearly half of the neurons in both A1 and ML decreased firing rate for increasing ΔBW (Table 1). However, the population of neurons that *significantly* encoded AM was largely dominated by cells that increased firing rate for increasing AM depth in both A1 and ML, with only 13.6% of AM encoders classified as 'decreasing' units in A1, and 10.6% of AM encoders 'decreasing' in ML. Among significant ΔBW encoders, the population was more evenly split between 'increasing' and 'decreasing' units in both A1 and ML: 43.3% of ΔBW encoders have 'decreasing' functions in A1 vs. 30.6% in ML. In both A1 and ML, there was a significant positive correlation between AM and BW selectivity, so cells that tended to increase firing rate for increasing AM levels, also tended to increase firing rate for increasing ΔBW levels (For FSI_{AM} vs. FSI_{BW} , A1 Pearson's rho = 0.3143, $p = 0.002$; ML Pearson's rho = 0.3109, $p = 5.3 \times 10^{-4}$) (Figure 1). In this feature selective attention task, we found no significant difference between A1 and ML in the proportions of 'increasing' and 'decreasing' encoding cells for either AM ('Increasing' $p = 0.21 \chi^2$ test; 'Decreasing' $p = 0.11 \chi^2$ test) or BW ('Increasing' $p = 0.22 \chi^2$ test; 'Decreasing' $p = 0.52 \chi^2$ test).

Vector strength encoding

We found a similar proportion of cells in A1 and ML that significantly phase-locked to AM ($p = 0.77$, χ^2 test), as measured by phase-projected vector strength (VS_{pp}) (Table 1). As in previous reports

(Niwa et al. 2013), we found VS_{pp} to be weaker in ML than A1 (Figure 2, $p < 0.05$ at all AM depths, Wilcoxon rank-sum Test). In both A1 and ML, there was no significant difference in phase-locking (VS_{pp}) between the attend AM and attend ΔBW conditions ($p > 0.05$, signed-rank test, Figure 2).

Feature discriminability and context effects

We used the signal detection theory-based area under the receiver operating characteristic (ROCa) to measure how well an ideal observer could detect the presence of each sound feature based on the neural responses (either firing rate or VS_{pp}). Increases in the levels of both features tended to yield increasing ROCa (A1 AM Spearman rho = 0.15, BW Spearman rho = .06; ML AM Spearman rho = 0.13, BW Spearman rho = 0.05) (Figure 3). However, there was no significant effect of attentional condition on either feature at any level of feature modulation for either A1 (Figure 3a,c) or ML (Figure 3b,d).

VS_{pp} -based discrimination (ROCa) of AM from unmodulated sounds was better at the lowest modulation depth in A1 than in ML ($p = 0.02$, Wilcoxon Rank Sum Test, Figure 4). At the higher modulation depths, VS_{pp} -based discrimination was similar in A1 and ML ($p = 0.99$ AM depth 2; $p = 0.26$, AM depth 3; Wilcoxon Rank Sum Test; Figure 4). However, there was no significant difference in VS_{pp} discriminability between attention conditions for any modulation depth ($p > 0.05$, signed-rank test, Figure 4).

Choice-related activity

We found a similar proportion of neurons in A1 (19.5%) and ML (26.2%) with significant choice-related activity during the stimulus window ($p = 0.31$, χ^2 test). In both areas, the population of neurons with significant choice-related activity during the response window (from S2 end to joystick movement)

was larger than during the stimulus window, and the proportions of neurons were again similar between the two areas (41.3% A1, 34.4% ML, $p = 0.41$, χ^2 test).

In A1, during the attend AM condition, CP values were evenly distributed about 0.5 during both the stimulus presentation (S2 median CP = 0.50, $p = 0.87$ signed-rank test) and the response window (RW median CP = 0.49, $p = 0.43$ signed-rank test) (Figure 5a,c). In contrast, during the attend Δ BW context, the CP values tended to be lower than 0.5 during both the stimulus (S2 median CP = 0.49, $p = 0.02$ signed-rank test) and the response window (RW median CP = 0.46, $p = 4.2 \times 10^{-8}$ signed-rank test) (Figure 5b,d). That is, during the attend Δ BW condition, the population of neurons tended to decrease firing rate when reporting target feature detection, whereas during the attend AM condition, it was equally likely for a neuron to increase firing rate for a report of target detection as it was for a report of target absence. There was a significant difference in the population CP distributions between attention conditions during the RW (Attend AM median = 0.49, Attend Δ BW median = 0.46, $p = 0.004$, signed-rank test), though not during the S2 ($p = 0.06$, signed-rank test).

The choice-related activity in ML was similar to that reported above in A1 during the S2. During the attend AM condition, activity was evenly distributed about 0.5 (S2 median CP = 0.50, $p = 0.94$, signed-rank test) (Figure 6a). During the attend Δ BW condition, average CP was less than 0.5 (S2 median CP = 0.49, $p = 0.043$ signed-rank test) (Figure 6b). However, during the response window, CP values were less than 0.5 in both the attend AM condition (median CP = 0.48, $p = 0.004$ signed-rank test) and the attend Δ BW condition (median CP = 0.47, $p = 2.7 \times 10^{-5}$ signed-rank test) (Figure 6c,d). This is in contrast to A1 where CP values tended to be lower than 0.5 only in the attend BW condition. There was no significant difference in the distribution of CP values in ML neurons between the attend AM and attend Δ BW conditions during the S2 ($p = 0.15$, signed-rank test). However, there was a significant difference in the CP distribution between the attend AM and attend Δ BW conditions in ML during the response window (Attend AM median = 0.48, Attend Δ BW median = 0.47, $p = 0.033$ signed-rank test), reflecting

the population shift to CP values less than 0.5 in the attend Δ BW condition compared to the attend AM condition.

Discussion

We found a large proportion of cells in ML that decreased firing rate with increasing AM detectability, similar to previous findings in ML (Johnson et al. 2020; Niwa et al. 2013). However, unlike these previous studies where ML had a significantly larger population of cells with decreasing AM depth functions than A1, we found a similar proportion of A1 neurons with decreasing AM depth functions. Further, the majority of neurons in both A1 and ML significantly encoding AM depth had increasing AM depth functions. This suggests that the encoding of amplitude modulation can be flexible depending upon the behavioral and sensory demands of the task. In essence, with increased perceptual difficulty, stimulus/feature ambiguity, and task difficulty it may be necessary for A1 to develop a more robust and appropriate code in order to solve the task, and for ML to take on more of the sensory processing, and thus the encoding schemes look more similar between these two areas.

We also found a large population of cells in both A1 and ML that were sensitive to changes in bandwidth. This was particularly surprising as the changes in bandwidth were relatively small compared to the 9-octave wide unmodulated noise. It's possible that the Δ BW encoding we saw was due to an increasing concentration of power in the middle frequencies of the broadband noise as the level of bandwidth restriction increased. It could also be caused by decreasing power in flanking inhibitory bands. Further studies investigating if and how neurons in A1 and ML encode small changes in spectral bandwidth to broad-band sounds under power-matched conditions could be enlightening.

Using phase-projected vector strength (VS_{pp}) as a measure of temporal coding, neither ML nor A1 single neurons showed attention-related changes in VS_{pp} -based sensitivity to AM or VS-based choice-related activity. This is consistent with previous results from our lab showing smaller effects for VS_{pp} -

based attention and choice than for firing rate (Niwa et al. 2013). A recent study that could help interpretation of this result shows thalamic projections to the striatum (an area involved in decisions and possibly attention) relay information about temporally modulated sounds in the form of phase-locking, whereas cortical projections to the striatum only convey information about temporally modulated sounds with average firing rate over the stimulus (Ponvert and Jaramillo 2019).

Attending to the target-feature did not significantly improve single neuron amplitude modulation or bandwidth restriction detection in A1 or ML. This seems surprising considering the wide array of effects that have been previously reported in auditory cortex related to different tasks, and behavioral contexts (Atiani et al. 2014; Bagur et al. 2018; Buran et al. 2014; Francis et al. 2018a; Niwa et al. 2012b; Otazu et al. 2009; Lakatos et al. 2013; Angeloni and Geffen 2018; Sutter and Shamma 2011). In macaque monkeys, an improvement in both rate-based and temporal AM encoding was observed in A1 and ML neurons when animals performed a single-feature AM detection task compared to when animals passively listened to the same stimuli (Niwa et al. 2013, 2015). We did not see a similar level of encoding improvement, possibly due to the more fine-tuned form of attention needed to perform this task.

One might expect to observe smaller effects from this more selective form of attention than in a passive versus active listening task, as the difference between attending to one feature of a sound compared to another is much smaller than switching between paying attention to a sound and passive sound presentation. Furthermore, arousal, as measured with pupillometry, has recently been shown to correlate with increases in activity, gain and trial-to-trial reliability of A1 neurons (Schwartz et al. 2019), which could account for some of the effects seen in task engagement paradigms.

Feature-based attention has been shown to have gain effects on neurons tuned to the attended feature in visual cortex (Ni and Maunsell 2019; Treue and Trujillo 1999). It is possible that we did not see a similar gain effect of feature attention in AC due to the mixed-selectivity we and others (Chambers et

al. 2014) found in the encoding of these features (i.e. most neurons are sensitive to both AM and ΔBW). However, it is likely that mixed-selectivity is not the only reason we did not see a gain effect. In a study where rats performed a frequency categorization task with shifting boundaries, Jaramillo and colleagues similarly found that neurons in AC did not improve their discriminability with attentional context (Jaramillo et al. 2014). This similar lack of enhancement seen in a task where only a single feature is modulated, suggests that the mechanism for feature attention in auditory cortex could be enacted via a different mechanism.

In visual cortical studies probing *selective* feature attention – where the subject must distinguish between features within a single object, rather than object- or place-oriented, feature-based attention – results have been similarly complex. At the level of the single neuron, there have not been clear, gain-like improvements in the sensitivity to the attended feature (Chen et al. 2012; Mirabella et al. 2007; Sasaki and Uka 2009; Uka et al. 2012). Further, the effects of feature-selective attention seem to be dependent upon not just the tuning preferences of a neuron, but also the strength of its tuning (Ruff and Born 2015). These studies, along with our own, suggest that segregation of features within an object may require a different mechanism relative to object-directed, feature-based attention.

In each of the feature-selective attention studies cited above, a common observation is that single neurons in sensory cortex have mixed selectivity for the features in the task, as opposed to being uniquely responsive to one feature or another. Such mixed selectivity among single neurons may permit sophisticated, flexible computations at the population level (Fusi et al. 2016). It thus seems likely the mechanism for feature-selective attention lies not at the level of the single neuron, but rather requires the integration of activity from a larger population of neurons. A feature-selective study using ERPs found that the neural responses to identical stimuli varied when the subjects attend to different features of the stimulus (Nobre et al. 2006). The single neuron and neural circuit mechanisms underlying this effect remain unclear. One such possible mechanism might be the structure of

correlated variability within the population, which has been shown to be modulated by feature-selective attention (Downer et al. 2017b). Another study, simulating populations by pooling single-neurons across A1 recordings permitted clear segregation of these two features, as well as an enhancement in discrimination of the attended feature (Downer et al. 2020). Further studies investigating feature-selective attention at the level of populations of neurons are necessary to better understand the underlying mechanisms.

We did see an interesting difference in the distribution of choice-related activity between the attentional conditions, where the correlation between firing rate and choice shifted direction between conditions. During the attend AM context, CP was evenly distributed about 0.5 with some neurons showing significant choice activity at either extreme. In contrast, during the attend BW context, CP values were shifted towards 0, with very few neurons having significant choice-related activity greater than 0.5 (increasing firing rate for ‘feature-present’ response). Neurons in auditory cortical areas may also modulate their responses to motor events (Brosch 2005). Some previous reports on choice-related activity have been difficult to interpret, as they employed a Go/No-Go task format in which one perceptual choice required a movement and the other choice did not (Brosch 2005; Niwa et al. 2013). Therefore, the choice-related activity observed was difficult to disentangle from a general preparation to move. The task reported here was a Yes/No forced-choice task, requiring a motor response to each decision (target present versus target absent). The shift in choice-related activity between attention conditions observed in this force choice task, and another recent study (Guo et al. 2019) shows that this choice-related activity cannot simply reflect motor preparation or action. This then strengthens the possible relationship between this activity and the decision or attention process.

The lack of clear attentional improvement of single neuron feature encoding found in this study suggests one or more of the following: (1) the feature-selective attention required in this task is not implemented at the level of an individual neuron in A1 or ML; (2) the feature-selective attention

necessary for this particular task occurs at a later stage in auditory processing; (3) the mixed selectivity of single neurons in A1 and ML for these features complicates the interpretability of the effects of attention at the single neuron level, in contrast to feature-based attention neurons studied found in visual cortex (Martinez-Trujillo and Treue 2004; Maunsell 2015; Maunsell and Treue 2006). While we did not see robust differences in encoding between attentional conditions, the difference in attentional choice-related activity reveals that it is not simply reflective of motor preparation, and suggests that activity correlated to reported choice as early as A1 could be informing perceptual and decision processes.

	AM Rate Coder	Decreasing AM	Increasing AM +AM coder	Decreasing AM + AM coder	BW Coder	Decreasing BW	Increasing BW +BW Coder	Decreasing BW + BW Coder	VS Coder
A1 (n=92)	47.8%	32.6%	41.3%	6.5%	32.6%	42.5%	18.5%	14.1%	32.6%
ML (n=122)	38.5%	27.1%	34.4%	4.1%	29.5%	41.8%	20.5%	9.0%	30.33%

A.

B.

Figure 1: Single neuron feature selectivity index (FSI), a measure of how sensitive a neuron is to changes in each feature value separately. **A.** A1: a positive correlation between AM and Δ BW selectivity (Pearson $\rho = 0.3143$, $p = 0.002$) **B.** ML: positive correlation between AM and BW selectivity (Pearson's $\rho = 0.3109$, $p = 5.32 \times 10^{-4}$)

508

509

Figure 2. Average phase locking ability of single units in A1 (solid lines) and ML (dashed lines), as measured by phase projected vector strength (VS_{pp}). VS_{pp} is greater in A1 (solid) than ML (dashed) at low AM depths (AM level 1, $p = 0.01$; AM level 2, $p = 0.002$, Wilcoxon Ranked Sum), though phase locking is more similar at the highest AM depth ($p = 0.73$, Wilcoxon ranked sum). There was no significant difference in either area between attend AM (gold) attend ΔBW (blue) conditions, ($p > 0.05$ for all AM levels, Wilcoxon signed rank test).

Figure 3: Firing rate based ROCa for each feature by attention condition. Blue lines indicate attend BW condition, yellow lines indicate attend AM condition. **A.** AM encoding in A1 (38 increasing, 6 decreasing cells). **B.** AM encoding in ML (42 increasing, 5 decreasing cells) **C.** BW encoding in A1 (17 increasing, 13 decreasing cells) **D.** BW encoding in ML (25 increasing, 11 decreasing cells). There was no significant effect of attentional condition on either feature at any level of feature modulation for either A1 (**A,C**) or ML (**B,D**).

Figure 4: VS_{pp}-based discriminability (ROCa) of AM from unmodulated sounds in A1 and ML for attend AM (yellow) and attend ΔBW (blue) conditions. **A.** In A1, VS_{pp}-based discriminability of AM is not significantly different between attention conditions ($p > 0.05$, signed-rank test). **B.** In ML, VS_{pp}-based ROCa does not differ between attentional conditions ($p > 0.05$, signed-rank test). At low modulation depths (AM depth rank = 1), A1 had significantly better AM discriminability than ML ($p = 0.02$, rank sum test), however they were not significantly different at the higher modulation depths (AM ranks 2 and 3, $p > 0.05$, rank sum test).

511

Figure 6: Choice probability in ML, as in Figure 5. CP during the attend AM condition is evenly distributed about 0.5 in **A**. the stimulus window (median = 0.50, $p = 0.94$, signed-rank test). However, in the response window **C**. CP values tended to be less than 0.5 (median = 0.48, $p = 0.004$, signed-rank test). In the attend Δ BW condition, CP values tended to be less than 0.5 in both **B**. the stimulus window (median = 0.49, $p = 0.043$ signed-rank test) and **D**. the response window (median = 0.47, $p = 2.7 \times 10^{-5}$ signed-rank test). As in A1, there was a significant difference in the population CP distributions between attention conditions during the RW ($p = 0.033$, signed-rank test), though not during the S2 ($p = 0.15$, signed-rank test).

References

- Angeloni C, Geffen M.** Contextual modulation of sound processing in the auditory cortex. *Current Opinion in Neurobiology* 49: 8–15, 2018.
- Atiani S, David SV, Elgueda D, Locastro M, Radtke-Schuller S, Shamma SA, Fritz JB.** Emergent Selectivity for Task-Relevant Stimuli in Higher-Order Auditory Cortex. *Neuron* 82: 486–499, 2014.
- Bagur S, Averseng M, Elgueda D, David S, Fritz J, Yin P, Shamma S, Boubenec Y, Ostojic S.** Go/No-Go task engagement enhances population representation of target stimuli in primary auditory cortex. *Nat Commun* 9: 2529, 2018.
- Bathellier B, Ushakova L, Rumpel S.** Discrete Neocortical Dynamics Predict Behavioral Categorization of Sounds. *Neuron* 76: 435–449, 2012.
- Bizley JK, Cohen YE.** The what, where and how of auditory-object perception. *Nature Reviews Neuroscience* 14: 693–707, 2013.
- Bizley JK, Walker KMM, Nodal FR, King AJ, Schnupp JWH.** Auditory Cortex Represents Both Pitch Judgments and the Corresponding Acoustic Cues. *Current Biology* 23: 620–625, 2013.
- Bregman AS.** *Auditory Scene Analysis: The Perceptual Organization of Sound*. MIT Press, 1994.
- Britten K, Shadlen M, Newsome W, Movshon J.** The analysis of visual motion: a comparison of neuronal and psychophysical performance. *The Journal of Neuroscience* 12: 4745–4765, 1992.
- Britten KH, Newsome WT, Shadlen MN, Celebrini S, Movshon JA.** A relationship between behavioral choice and the visual responses of neurons in macaque MT. *Visual Neuroscience* 13: 87–100, 1996.
- Brosch M.** Nonauditory Events of a Behavioral Procedure Activate Auditory Cortex of Highly Trained Monkeys. *Journal of Neuroscience* 25: 6797–6806, 2005.
- Buran BN, von Trapp G, Sanes DH.** Behaviorally Gated Reduction of Spontaneous Discharge Can Improve Detection Thresholds in Auditory Cortex. *Journal of Neuroscience* 34: 4076–4081, 2014.
- Carcea I, Insanally MN, Froemke RC.** Dynamics of auditory cortical activity during behavioural engagement and auditory perception. *Nat Commun* 8: 14412, 2017.
- Chambers AR, Hancock KE, Sen K, Polley DB.** Online Stimulus Optimization Rapidly Reveals Multidimensional Selectivity in Auditory Cortical Neurons. *Journal of Neuroscience* 34: 8963–8975, 2014.
- Chen X, Hoffmann K-P, Albright TD, Thiele A.** Effect of feature-selective attention on neuronal responses in macaque area MT. *Journal of Neurophysiology* 107: 1530–1543, 2012.
- Christison-Lagay KL, Bennur S, Cohen YE.** Contribution of spiking activity in the primary auditory cortex to detection in noise. *Journal of Neurophysiology* 118: 3118–3131, 2017.
- Christison-Lagay KL, Cohen YE.** The Contribution of Primary Auditory Cortex to Auditory Categorization in Behaving Monkeys. *Front Neurosci* 12: 601, 2018.

548 **Downer JD, Niwa M, Sutter ML.** Task Engagement Selectively Modulates Neural Correlations in Primary
549 Auditory Cortex. *Journal of Neuroscience* 35: 7565–7574, 2015.

550 **Downer JD, Niwa M, Sutter ML.** Hierarchical differences in population coding within auditory cortex.
551 *Journal of Neurophysiology* 118: 717–731, 2017a.

552 **Downer JD, Rapone B, Verhein J, O'Connor KN, Sutter ML.** Feature-Selective Attention Adaptively Shifts
553 Noise Correlations in Primary Auditory Cortex. *The Journal of Neuroscience* 37: 5378–5392, 2017b.

554 **Downer JD, Verhein JR, Rapone BC, O'Connor KN, Sutter ML.** An emergent population code in primary
555 auditory cortex supports selective attention to spectral and temporal sound features. *bioRxiv*
556 2020.03.09.984773, 2020.

557 **Elgueda D, Duque D, Radtke-Schuller S, Yin P, David SV, Shamma SA, Fritz JB.** State-dependent
558 encoding of sound and behavioral meaning in a tertiary region of the ferret auditory cortex. *Nat*
559 *Neurosci* 22: 447–459, 2019.

560 **Eliades SJ, Wang X.** Sensory-Motor Interaction in the Primate Auditory Cortex During Self-Initiated
561 Vocalizations. *Journal of Neurophysiology* 89: 2194–2207, 2003.

562 **Engle JR, Recanzone GH.** Characterizing spatial tuning functions of neurons in the auditory cortex of
563 young and aged monkeys: a new perspective on old data. *Frontiers in Aging Neuroscience* 4: 36, 2013.

564 **Francis NA, Elgueda D, Englitz B, Fritz JB, Shamma SA.** Laminar profile of task-related plasticity in ferret
565 primary auditory cortex. *Sci Rep* 8: 16375, 2018a.

566 **Francis NA, Winkowski DE, Sheikhattar A, Armengol K, Babadi B, Kanold PO.** Small Networks Encode
567 Decision-Making in Primary Auditory Cortex. *Neuron* 97: 885–897.e6, 2018b.

568 **Fritz J, Shamma S, Elhilali M, Klein D.** Rapid task-related plasticity of spectrotemporal receptive fields in
569 primary auditory cortex. *Nat Neurosci* 6: 1216–1223, 2003.

570 **Fritz JB.** Differential Dynamic Plasticity of A1 Receptive Fields during Multiple Spectral Tasks. *Journal of*
571 *Neuroscience* 25: 7623–7635, 2005.

572 **Fritz JB, Elhilali M, Shamma SA.** Adaptive Changes in Cortical Receptive Fields Induced by Attention to
573 Complex Sounds. *Journal of Neurophysiology* 98: 2337–2346, 2007.

574 **Fusi S, Miller EK, Rigotti M.** Why neurons mix: high dimensionality for higher cognition. *Current Opinion*
575 *in Neurobiology* 37: 66–74, 2016.

576 **Goldberg JM, Brown PB.** Response of binaural neurons of dog superior olivary complex to dichotic tonal
577 stimuli: some physiological mechanisms of sound localization. *Journal of Neurophysiology* 32: 613–636,
578 1969.

579 **Green DM, Swets JA.** *Signal detection theory and psychophysics*. Huntington, NY: Krieger Publishing
580 Company, 1974.

581 **Grimault N, Bacon SP, Michey C.** Auditory stream segregation on the basis of amplitude-modulation
582 rate. *The Journal of the Acoustical Society of America* 111: 1340–1348, 2002.

583 **Gronskaya E, von der Behrens W.** Evoked Response Strength in Primary Auditory Cortex Predicts
584 Performance in a Spectro-Spatial Discrimination Task in Rats. *J Neurosci* 39: 6108–6121, 2019.

585 **Guo L, Weems JT, Walker WI, Levichev A, Jaramillo S.** Choice-Selective Neurons in the Auditory Cortex
586 and in Its Striatal Target Encode Reward Expectation. *J Neurosci* 39: 3687–3697, 2019.

587 **Hackett TA.** Information flow in the auditory cortical network. *Hearing Research* 271: 133–146, 2011.

588 **Huang Y, Brosch M.** Associations between sounds and actions in primate prefrontal cortex. *Brain*
589 *Research* 1738: 146775, 2020.

590 **Huang Y, Heil P, Brosch M.** Associations between sounds and actions in early auditory cortex of
591 nonhuman primates. *eLife* 8: e43281, 2019.

592 **Jaramillo S, Borges K, Zador AM.** Auditory Thalamus and Auditory Cortex Are Equally Modulated by
593 Context during Flexible Categorization of Sounds. *Journal of Neuroscience* 34: 5291–5301, 2014.

594 **Johnson JS, Niwa M, O'Connor KN, Sutter ML.** Amplitude modulation encoding in auditory cortex:
595 Comparisons between the primary and middle lateral belt regions. *bioRxiv*, 2020.
596 doi:10.1101/2020.03.05.979575.

597 **Kang I, Maunsell JHR.** Potential confounds in estimating trial-to-trial correlations between neuronal
598 response and behavior using choice probabilities. *Journal of Neurophysiology* 108: 3403–3415, 2012.

599 **Lakatos P, Musacchia G, O'Connell MN, Falchier AY, Javitt DC, Schroeder CE.** The Spectrotemporal Filter
600 Mechanism of Auditory Selective Attention. *Neuron* 77: 750–761, 2013.

601 **Lee C-C, Middlebrooks JC.** Auditory cortex spatial sensitivity sharpens during task performance. *Nature*
602 *Neuroscience* 14: 108–114, 2011.

603 **Lin P-A, Asinof SK, Edwards NJ, Isaacson JS.** Arousal regulates frequency tuning in primary auditory
604 cortex. *Proc Natl Acad Sci USA* 116: 25304–25310, 2019.

605 **Mardia KV, Jupp PE.** *Directional statistics*. Chichester; New York: J. Wiley, 2000.

606 **Martinez-Trujillo JC, Treue S.** Feature-Based Attention Increases the Selectivity of Population Responses
607 in Primate Visual Cortex. *Current Biology* 14: 744–751, 2004.

608 **Massoudi R, Van Wanrooij MM, Van Wetter SMCI, Versnel H, Van Opstal AJ.** Task-related preparatory
609 modulations multiply with acoustic processing in monkey auditory cortex. *Eur J Neurosci* 39: 1538–1550,
610 2014.

611 **Maunsell JHR.** Neuronal Mechanisms of Visual Attention. *Annual Review of Vision Science* 1: 373–391,
612 2015.

613 **Maunsell JHR, Treue S.** Feature-based attention in visual cortex. *Trends in Neurosciences* 29: 317–322,
614 2006.

615 **McDermott JH.** The cocktail party problem. *Curr Biol* 19: R1024-1027, 2009.

616 **Merzenich MM, Brugge JF.** Representation of the cochlear partition on the superior temporal plane of
617 the macaque monkey. *Brain Research* 50: 275–296, 1973.

618 **Mirabella G, Bertini G, Samengo I, Kilavik BE, Frilli D, Della Libera C, Chelazzi L.** Neurons in Area V4 of
619 the Macaque Translate Attended Visual Features into Behaviorally Relevant Categories. *Neuron* 54: 303–
620 318, 2007.

621 **Ni AM, Maunsell JHR.** Neuronal Effects of Spatial and Feature Attention Differ Due to Normalization. *J*
622 *Neurosci* 39: 5493–5505, 2019.

623 **Niwa M, Johnson JS, O'Connor KN, Sutter ML.** Active Engagement Improves Primary Auditory Cortical
624 Neurons' Ability to Discriminate Temporal Modulation. *Journal of Neuroscience* 32: 9323–9334, 2012a.

625 **Niwa M, Johnson JS, O'Connor KN, Sutter ML.** Activity Related to Perceptual Judgment and Action in
626 Primary Auditory Cortex. *Journal of Neuroscience* 32: 3193–3210, 2012b.

627 **Niwa M, Johnson JS, O'Connor KN, Sutter ML.** Differences between Primary Auditory Cortex and
628 Auditory Belt Related to Encoding and Choice for AM Sounds. *Journal of Neuroscience* 33: 8378–8395,
629 2013.

630 **Niwa M, O'Connor KN, Engall E, Johnson JS, Sutter ML.** Hierarchical effects of task engagement on
631 amplitude modulation encoding in auditory cortex. *Journal of Neurophysiology* 113: 307–327, 2015.

632 **Nobre AC, Rao A, Chelazzi L.** Selective Attention to Specific Features within Objects: Behavioral and
633 Electrophysiological Evidence. *Journal of Cognitive Neuroscience* 18: 539–561, 2006.

634 **O'Connor KN, Petkov CI, Sutter ML.** Adaptive Stimulus Optimization for Auditory Cortical Neurons.
635 *Journal of Neurophysiology* 94: 4051–4067, 2005.

636 **Osmanski MS, Wang X.** Behavioral Dependence of Auditory Cortical Responses. *Brain Topogr* 28: 365–
637 378, 2015.

638 **Otazu GH, Tai L-H, Yang Y, Zador AM.** Engaging in an auditory task suppresses responses in auditory
639 cortex. *Nature Neuroscience* 12: 646–654, 2009.

640 **Ponvert ND, Jaramillo S.** Auditory Thalamostriatal and Corticostriatal Pathways Convey Complementary
641 Information about Sound Features. *J Neurosci* 39: 271–280, 2019.

642 **Rauschecker JP, Tian B.** Mechanisms and streams for processing of “what” and “where” in auditory
643 cortex. *Proceedings of the National Academy of Sciences* 97: 11800–11806, 2000.

644 **Romanski LM, Tian B, Fritz J, Mishkin M, Goldman-Rakic PS, Rauschecker JP.** Dual streams of auditory
645 afferents target multiple domains in the primate prefrontal cortex. *Nature Neuroscience* 2: 1131–1136,
646 1999.

647 **Ruff DA, Born RT.** Feature attention for binocular disparity in primate area MT depends on tuning
648 strength. *Journal of Neurophysiology* 113: 1545–1555, 2015.

649 **Runyan CA, Piasini E, Panzeri S, Harvey CD.** Distinct timescales of population coding across cortex.
650 *Nature* 548: 92–96, 2017.

651 **Saleem K, Logothetis Nikos K.** *A combined MRI and histology atlas of the rhesus monkey brain in*
652 *stereotaxic coordinates*. Burlington, MA: Academic Press, 2007.

653 **Sasaki R, Uka T.** Dynamic Readout of Behaviorally Relevant Signals from Area MT during Task Switching.
654 *Neuron* 62: 147–157, 2009.

655 **Schneider DM, Nelson A, Mooney R.** A synaptic and circuit basis for corollary discharge in the auditory
656 cortex. *Nature* 513: 189–194, 2014.

657 **Schnupp JWH.** Plasticity of Temporal Pattern Codes for Vocalization Stimuli in Primary Auditory Cortex.
658 *Journal of Neuroscience* 26: 4785–4795, 2006.

659 **Schwartz ZP, Buran BN, David SV.** Pupil-associated states modulate excitability but not stimulus
660 selectivity in primary auditory cortex. *Journal of Neurophysiology* 191–208, 2019.

661 **Schwartz ZP, David SV.** Focal Suppression of Distractor Sounds by Selective Attention in Auditory Cortex.
662 *Cerebral Cortex* 28: 323–339, 2018.

663 **Scott BH, Malone BJ, Semple MN.** Effect of Behavioral Context on Representation of a Spatial Cue in
664 Core Auditory Cortex of Awake Macaques. *Journal of Neuroscience* 27: 6489–6499, 2007.

665 **Shannon RV, Zeng F-G, Kamath V, Wygonski J, Ekelid M.** Speech Recognition with Primarily Temporal
666 Cues. *Science* 270: 303–304, 1995.

667 **Shinn-Cunningham BG.** Object-based auditory and visual attention. *Trends in Cognitive Sciences* 12:
668 182–186, 2008.

669 **Strickland EA, Viemeister NF.** The effects of frequency region and bandwidth on the temporal
670 modulation transfer function. *The Journal of the Acoustical Society of America* 102: 1799–1810, 1997.

671 **Sutter ML, Shamma SA.** The Relationship of Auditory Cortical Activity to Perception and Behavior. In:
672 *The Auditory Cortex*, edited by Winer JA, Schreiner CE. Springer US, p. 617–641.

673 **Tian B, Rauschecker JP.** Processing of Frequency-Modulated Sounds in the Lateral Auditory Belt Cortex
674 of the Rhesus Monkey. *Journal of Neurophysiology* 92: 2993–3013, 2004.

675 **von Trapp G, Buran BN, Sen K, Semple MN, Sanes DH.** A Decline in Response Variability Improves
676 Neural Signal Detection during Auditory Task Performance. *Journal of Neuroscience* 36: 11097–11106,
677 2016.

678 **Treue S, Trujillo JCM.** Feature-based attention influences motion processing gain in macaque visual
679 cortex. *Nature* 399: 575–579, 1999.

680 **Tsunada J, Cohen YE.** Neural mechanisms of auditory categorization: from across brain areas to within
681 local microcircuits. *Frontiers in Neuroscience* 8: 161, 2014.

682 **Tsunada J, Liu ASK, Gold JI, Cohen YE.** Causal contribution of primate auditory cortex to auditory
683 perceptual decision-making. *Nature Neuroscience* 19: 135–142, 2016.

684 **Uka T, Sasaki R, Kumano H.** Change in Choice-Related Response Modulation in Area MT during Learning
685 of a Depth-Discrimination Task is Consistent with Task Learning. *Journal of Neuroscience* 32: 13689–
686 13700, 2012.

687 **Van Tasell DJ, Soli SD, Kirby VM, Widin GP.** Speech waveform envelope cues for consonant recognition.
688 *The Journal of the Acoustical Society of America* 82: 1152–1161, 1987.

689 **Wang L, Narayan R, Grana G, Shamir M, Sen K.** Cortical Discrimination of Complex Natural Stimuli: Can
690 Single Neurons Match Behavior? *Journal of Neuroscience* 27: 582–589, 2007.

691 **Woods KJP, McDermott JH.** Attentive Tracking of Sound Sources. *Current Biology* 25: 2238–2246, 2015.

692 **Yin P, Fritz JB, Shamma SA.** Rapid Spectrotemporal Plasticity in Primary Auditory Cortex during Behavior.
693 *Journal of Neuroscience* 34: 4396–4408, 2014.

694 **Yin P, Johnson JS, O'Connor KN, Sutter ML.** Coding of Amplitude Modulation in Primary Auditory Cortex.
695 *Journal of Neurophysiology* 105: 582–600, 2011.

696