Testosterone-mediated behavior shapes the emergent properties of social networks

Roslyn Dakin^{1,2}, Ignacio T. Moore³, Brent M. Horton⁴, Ben J. Vernasco³, T. Brandt Ryder^{1,5}

¹Migratory Bird Center, Smithsonian Conservation Biology Institute, National Zoological Park, Washington, DC 20013, USA

²Current address: Department of Biology, Carleton University, Ottawa, Ontario K1S 5B6, Canada

³Department of Biological Sciences, Virginia Tech, Blacksburg, Virginia 24061, USA

⁴Department of Biology, Millersville University, Millersville, Pennsylvania 17551, USA

⁵Current address: Bird Conservancy of the Rockies, Fort Collins, Colorado, 80525, USA

Correspondence: rydert@si.edu

16 Abstract

1. Social networks can vary in their organization and dynamics, with implications for
ecological and evolutionary processes. Understanding the mechanisms that drive social
network dynamics requires integrating individual-level biology with comparisons across
multiple social networks.
2. Testosterone is a key mediator of vertebrate social behavior and can influence how
individuals interact with social partners. Although the effects of testosterone on
individual behavior are well established, no study has examined whether hormone-
mediated behavior can scale up to shape the emergent properties of social networks.
3. We investigated the relationship between testosterone and social network dynamics in the
wire-tailed manakin, a lekking bird species in which male-male social interactions form
complex social networks. We used an automated proximity system to longitudinally
monitor several leks and we quantified the social network structure at each lek. Our
analysis examines three emergent properties of the networks: social specialization (the
extent to which a network is partitioned into exclusive partnerships), network stability
(the overall persistence of partnerships through time), and behavioral assortment (the
tendency for like to associate with like). All three properties are expected to promote the
evolution of cooperation. As the predictor, we analyzed the collective testosterone of
males within each social network.
4. Social networks that were composed of high-testosterone dominant males were less
specialized, less stable, and had more negative behavioral assortment, after accounting
for other factors. These results support our main hypothesis that individual-level hormone
physiology can predict group-level network dynamics. We also observed that larger leks

with more interacting individuals had more positive behavioral assortment, suggesting
 40 that small groups may constrain the processes of homophily and behavior-matching.

5. Overall, these results provide evidence that hormone-mediated behavior can shape the
 42 broader architecture of social groups. Groups with high average testosterone exhibit
 social network properties that are predicted to impede the evolution of cooperation.

44
Keywords: androgens, behavioral endocrinology, collective behavior, cooperation, dynamic
 46 networks, social networks, testosterone

Introduction

Behavioral interactions are the foundation of social network structures that can vary through time, among populations, and across species. Network structures play an important role in many ecological and evolutionary processes, including the spread of diseases (Sah *et al.* 2018; Stroeymeyt *et al.* 2018), the transmission of information and resources (Aplin *et al.* 2012; Maldonado-Chaparro *et al.* 2018), and selection on individual behavior (Ohtsuki *et al.* 2006). A major challenge is understanding how individual-level factors, such as physiological and behavioral mechanisms, scale up to drive the emergent structural properties of social groups (Krause & Ruxton 2002). Linking these two levels of analysis is difficult because it requires integrating individual-level data with repeated measures of entire social groups (Sah *et al.* 2017).

Here, we use a comparison of multiple social networks through time to investigate how hormone-mediated behavior shapes the higher-order structure of social networks. Testosterone is a steroid hormone that is well known for its influence on social behavior and its sensitivity to changes in the social environment (Wingfield *et al.* 1990; Adkins-Regan 2005; Goymann 2009). Testosterone often promotes physical aggression and other behaviors associated with social dominance (Oyegbile & Marler 2005; Fuxjager *et al.* 2010). Testosterone can also promote status-seeking behaviors in a non-aggressive context, including cooperative and gregarious behavior (Eisenegger *et al.* 2011; Boksem *et al.* 2013; Ryder *et al.* 2020). Overall, these hormone-signaling pathways are essential for the development and modulation of complex behavioral phenotypes (Cohen *et al.* 2012).

To investigate how testosterone is associated with social network dynamics, we studied the wire-tailed manakin (*Pipra filicauda*), a bird species in which the males engage in coordinated displays with each other at sites known as leks. In *P. filicauda*, males exhibit two

social status classes: the dominant males who hold territories on the leks, and subordinate, “floater” males who must acquire a territory on a lek before they can mate (Heindl 2002; Ryder *et al.* 2011a). Male social partnerships in wire-tailed manakins can be remarkably stable and typically occur between two unrelated males, most commonly, but not exclusively, between a territory-owner and a floater (Ryder *et al.* 2011a; Dakin & Ryder 2020). Two features of the manakin leks make them especially well-suited to studying the relationship between hormones and group-level social structure. First, testosterone is known to affect the social behavior of individual male wire-tailed manakins (Ryder *et al.* 2020). Second, these partnerships among males form the basis of complex and dynamic social networks that are replicated across leks, facilitating a comparative approach (Dakin & Ryder 2018, 2020). Given this background, we sought to test whether testosterone and its effects on male behavioral phenotype could drive the emergent properties of the social network.

The broader function of male-male social behavior and coordinated displays in the manakin family has been the subject of considerable study (e.g., Prum 1994; DuVal 2007; McDonald 2007; Ryder *et al.* 2008, 2009; Díaz-Muñoz *et al.* 2014). One possible function is that dyadic displays may be competitive and/or they may serve to maintain an individual’s position in a dominance hierarchy (Prum 1994; Heindl 2002). Social interactions may also represent cooperative coalitions that provide benefits to both parties (Ryder *et al.* 2008, 2009). The potential competitive and cooperative functions of male-male social behavior are not mutually exclusive, and function may be context- and/or status-specific (Ryder *et al.* 2008). Coordinated displays may also be a vestige of ancestral cooperative behavior (i.e., the behavior may have been directly beneficial to both parties in the past, and it persists today, even if it no longer has adaptive benefits; Prum 1994). Given this background and recent evidence that testosterone

modulates male social behavior in wire-tailed manakins (Ryder *et al.* 2020), we focused this
 94 study on three emergent properties of the social networks formed by male-male interactions that
 can influence the evolution and maintenance of cooperation (Fig. 1).

96 The first property we examined, social specialization, seeks to capture the exclusivity of
 the relationships between partners in a social network. In the context of manakin behavior,
 98 highly specialized networks are well-partitioned among specific territory-owner and floater
 relationships, as illustrated in Fig. 1A. In humans, social specialization has been found to
 100 maximize the ability of a team to successfully perform a challenging task (Jehn & Shah 1997). In
 manakins, we expect specialization to improve the familiarity of social partners and the
 102 behavioral coordination of their displays. Greater specialization is also expected to minimize
 conflict over mating and territorial ascension opportunities (Schjelderup-Ebbe 1922; McDonald
 104 1993).

The second property, network stability (Fig. 1B), quantifies the average persistence of
 106 social partnerships through time (Poisot *et al.* 2012; Dakin & Ryder 2020). Coordinated displays
 in manakins require the synchronization of complex behaviors, and previous empirical work
 108 indicates that longer partnership tenure has a positive effect on display coordination (Trainer &
 McDonald 1995; Trainer *et al.* 2002). Greater temporal stability of partnerships also increases
 110 the opportunity for familiarity and reciprocity within a social network (Trivers 1971; Roberts &
 Sherratt 1998; Croft *et al.* 2006).

112 The third property, behavioral assortment (Fig. 1C), captures the extent to which males
 interact with partners who express similar behaviors (i.e., is like associated with like? Croft *et al.*
 114 2006; Farine 2014). At the proximate level, positive assortment may represent the outcome of
 generalized reciprocity and/or partner choice (Fowler & Christakis 2010; Dakin & Ryder 2018).

At the ultimate level, positive assortment has also been shown to promote the evolution of cooperation (Ohtsuki *et al.* 2006). To quantify the overall behavioral assortment of the manakin networks, we focused on two correlated metrics of social behavior within the network: “strength” (a male’s frequency of daily social interactions) and “degree” (his daily number of social partnerships; Dakin & Ryder 2018). We used a composite measure of assortment that averaged the assortativity indices of these two phenotypes.

As a potential predictor of specialization, stability, and assortment, we quantified the collective testosterone of the manakin leks (Fig. 2; Akinola *et al.* 2016). Because hormone-behavior relationships are status-specific in manakins and many other species (Eisenegger *et al.* 2011; Boksem *et al.* 2013; Ryder *et al.* 2020), we analyzed collective testosterone for each of the two status classes separately. High individual testosterone levels are associated with reduced sociality in dominant males, but increased sociality in subordinate males (Ryder *et al.* 2020). We therefore predicted that the average (or collective) testosterone of territorial males in a group would be negatively associated with its specialization, stability, and assortment. In contrast, we predicted that the collective testosterone of floater males in a group would be positively associated with these three emergent properties of the social network.

Materials and Methods

Study Population

We studied wire-tailed manakins (*Pipra filicauda*) at the Tiputini Biodiversity Station in Orellana Province, Ecuador (0° 38’ S, 76° 08’ W). This population of *P. filicauda* has been studied and individuals color-banded annually since 2002 (e.g., Ryder *et al.* 2008, 2009). The present study was conducted on 11 leks during peak breeding activity (December-March) across

three field seasons: 2015-16, 2016-17 and 2017-18. All research was approved by the
 140 Smithsonian ACUC (protocols #12-23, 14-25, and 17-11) and the Ecuadorean Ministry of the
 Environment (MAE-DNB-CM-2015-0008).

142 *Testosterone Assessment*

144 Male manakins were captured using mist-nets on the leks as described in (Ryder *et al.* 2020). We
 deployed up to 16 mist-nets simultaneously at a given lek, with the intention of capturing every
 146 male on the lek. We rotated the nets between leks with the goal of capturing each male up to
 three times per field season. Each mist-net was checked on a 30-minute schedule with variation
 148 resulting from capture rate (i.e., multiple birds being caught on the same net run; Vernasco *et al.*
 2019; Ryder *et al.* 2020). The amount of time a male spends in the net has a subtle, but
 150 significant, negative effect on his circulating testosterone (Vernasco *et al.* 2019). Therefore, we
 used video monitoring to determine the duration of time that each bird was in the mist-net prior
 152 to blood sampling, so that it could be accounted for in further analyses (mean = 17.5 minutes, SD
 = 10.3 minutes, range = 1–72 minutes). Following removal of a male from the mist-net, a small
 154 blood sample (< 125uL) was taken from the brachial vein and stored on ice prior to being
 centrifuged at 10,000 rpm for 5 min, as described in previous studies (Ryder *et al.* 2011b;
 156 Vernasco *et al.* 2019; Ryder *et al.* 2020). Plasma volume was measured to the nearest 0.25 ul and
 stored in 0.75 ml of 100% ethanol (Goymann *et al.* 2007). Testosterone was double extracted
 158 from the plasma using dichloromethane. Following extraction, a direct radioimmunoassay was
 used to measure the total plasma androgen concentration (ng/ml) adjusted by the extraction
 160 efficiency and plasma volume of each sample (Eikenaar *et al.* 2011; Ryder *et al.* 2011b).
 Hormone assays were conducted annually, and the detection limits were 0.12, 0.08, and 0.09

ng/ml for 2015-16, 2016-17 and 2017-18, respectively; any sample that fell below the assay-specific limit of detection was assigned that limit as its testosterone concentration as a most conservative estimate. The extraction efficiency for all samples was between 62-73%, and the intra-assay coefficients of variation were 6.6%, 11.6%, and 9.2% for 2015-16, 2016-17 and 2017-18, respectively; the inter-assay coefficient of variation was 19.5%.

Behavioral Recording

We used an automated data-logging system to monitor male-male interactions on the display territories of the leks (Ryder *et al.* 2012; Dakin & Ryder 2018; Ryder *et al.* 2020). The territories on these leks are specifically used for male-male coordinated social displays (as described in Schwartz & Snow 1978). At the beginning of each field season, male manakins were outfitted with coded nano-tags (NTQB-2, Lotek Wireless; 0.35 g). The tags transmitted a unique VHF signal ping once per 20 s for three months. In total, 296 tag deployments were performed on 180 individuals (mean 1.7 field seasons per male $\pm$ SD 0.7), 178 of whom also had hormone data (mean number of hormone samples per male = 3 $\pm$ SD 1.5). Approximately 10 days ($\pm$ SD 7) after tagging and sampling was completed at a given lek, a proximity data-logger (SRX-DL800, Lotek Wireless) was deployed within each display territory at the lek to record tagged males within a detection radius of 30 m (a distance that corresponds to the typical diameter of a manakin display territory; Heindl 2002; Dakin & Ryder 2018).

Proximity recording sessions ran from 06:00 to 16:00 for ~6 consecutive days ($\pm$ SD 1 day) and were performed ~3 times per field season at a given lek. Occasionally, the length of a recording session was extended due to extenuating circumstances such as inclement weather. The recording sessions were scheduled to be distributed evenly throughout each field season at

each lek, to minimize any confounding seasonal effects. Each recording session represents an
 186 observation of the social network at a given lek. In total, we conducted 86 recording sessions
 (29,760 data-logger hours) representing repeated measures of the social activity at 11 leks during
 188 three field seasons (see Fig. S1 in the supplement for additional details on the sampling regime).

Prior to data-logger deployment, each territory was also observed on non-recording days
 190 to identify the territory-holder based on his color-bands, following previous studies (Ryder *et al.*
 2008, 2009). These status assignments were subsequently verified in the proximity data.

Data Processing and Statistical Analysis

194 All computational and statistical analyses were performed in R (R Core Team 2018). Network
 illustrations were made using the igraph package (Csardi *et al.* 2018).

Social Interactions

198 Social networks were constructed by first defining interactions between two males that occurred
 on the display territories. To do this, a computational algorithm was used to identify joint
 200 detections, wherein two males were located at the same display territory within a pre-defined
 spatial and temporal threshold (Ryder *et al.* 2008, 2012; Dakin & Ryder 2018). For the temporal
 202 threshold, the two males had to occur < 45 s apart. This temporal threshold was chosen to allow
 for the fact that each tag pinged with a 20 second pulse rate, such that overlapping individuals
 204 could have up to a 40 s gap between their respective pings. For the spatial threshold, the two
 males had to have a difference in received signal strength values (ΔRSSI) < 10. This threshold
 206 corresponded to a typical distance between 0 to 5 m apart in a ground-truthing experiment
 (Dakin & Ryder 2018). Hence, according to our definition, a social interaction is initiated only

after males come within this approximate spatial threshold. We chose this spatial threshold because it is close enough to permit visual and acoustic contact during typical social behaviors (such as those described in Schwartz & Snow 1978). After completing our study, we also performed a sensitivity analysis to verify that our main results were robust to alternative spatial threshold definitions (see supplement for details).

Any repeated co-occurrence of the two males within 5 minutes was considered to be part of the same social interaction, but after a gap of ≥ 5 minutes, it was considered to be a new interaction between those two males. The average duration of social interactions defined by this method was $5.1 \text{ min} \pm \text{SD } 13.3$, $8.8 \text{ min} \pm \text{SD } 20.1$, and $7.0 \text{ min} \pm \text{SD } 20.5$ in the three respective field seasons, further indicating that these were sustained social interactions, rather than random encounters.

In total, we identified 36,885 social interactions over the three field seasons of study. These interactions were used to define a weighted social network for each lek recording session. The nodes in the network were the individual males, and the edges were weighted by the number of social interactions between each pair of males. An earlier validation study compared social interactions that were detected by the proximity system with those that were directly observed for 11 males (Ryder *et al.* 2012), and confirmed that all of the interactions detected by the proximity system were also directly observed.

Null Model Validation of the Social Networks

Two broad classes of methods have been described for building social networks: (i) networks that are built based purely on the proximity of animals unrelated to their behavioral context (“gambit of the group”), and (ii) networks that are built based on specific behavioral criteria that

are directly observed (Franks *et al.* 2010; Croft *et al.* 2011; Farine 2015). Although the
 232 interactions in our study were not directly observed, they were recorded at specialized display
 perches that have a known function in male-male social interactions (Schwartz & Snow 1978;
 234 Ryder *et al.* 2011a). Hence, they do not qualify as the gambit of the group. To further
 demonstrate this point, we performed a pre-network permutation of the raw data to determine
 236 whether the observed network edges occurred more often than expected by chance (Farine 2017).
 This analysis was highly conservative in that it preserved key features of the data including male
 238 visit rates to specific territories within each recording session and lek; it is described in detail in
 the supplement and Fig. S2. The results demonstrated that 95% of the observed network edges
 240 had a greater edge weight than expected under stringent permutation conditions. Moreover, these
 preferred edges had edge weights that were 10-fold to 50-fold greater than expected by chance
 242 under these stringent conditions (Fig. S2). This provides an additional validation to our approach,
 because it indicates that the observed network edges were nearly always preferred, even relative
 244 to other possible interactions within the same lek. These results are also consistent with a
 previous validation study indicating that these methods capture male-male coalition partnerships
 246 that are directly observed (Ryder *et al.* 2012).

Although pre-network data permutations are sometimes used to derive adjusted
 248 association indices, or to prune networks prior to further analysis, we did not take this approach
 for several reasons. First, the statistical rarity of a relationship in our system does not *a priori*
 250 define the importance of any one social interaction, especially given that all of the interactions
 occurred on display territories with specialized function. A single interaction between two rarely
 252 interacting males may have been highly consequential (e.g., if one of those males had a highly
 influential hormone-behavioral phenotype). Conversely, a single interaction between two

frequently interacting males may have been relatively unimportant. It would be unwarranted to assume that partnership rarity indicates the importance of any single interaction in this context. Second, and perhaps more importantly, the goal of this study was a comparison across multiple networks. Some networks are genuinely less preferential, and more random, than others. If we modified the network edges based on permutation-based indices, it would disproportionately prune the truly random networks, introducing a source of bias that would be contrary to our main goal. Hence, all further analyses are based on networks where the edge weights are given by the observed number of interactions, as this method is most appropriate for our study system and aims. Below, we also describe a separate node-label permutation that provided an additional check on our final statistical analyses.

Social Specialization

To quantify social specialization, we sought a metric that would capture the extent to which a network was partitioned into exclusive social relationships (as opposed to a network made up of non-specific or non-exclusive partnerships). To do this, we used a network metric of specialization that is commonly used in community ecology called H_2' (Blüthgen *et al.* 2006). An advantage of H_2' is that it is standardized against a theoretical maximum, based on the overall activity levels of different nodes and Shannon entropy (Blüthgen *et al.* 2006); this makes it possible to compare the extent of specialization across different bipartite networks in a standardized way. To apply this metric to our manakin data, we converted each lek's social network into its bipartite adjacency matrix (Fig. 1A), with floaters along one axis, and territory-holders on the other, and then calculated social specialization as H_2' using the bipartite package (Dormann *et al.* 2019). Higher values of specialization indicate that the network is well-

partitioned (i.e., made up of exclusive relationships), as illustrated in Fig. 1A. We chose to focus
 278 on floater-territorial specialization because these two social classes are well-defined and floater-
 territorial partnerships tend to be the most common in this species (Fig. S3; see also Ryder *et al.*
 280 2011a).

Our measure social specialization at the network level, H_2' , can also be related to the
 282 exclusivity of social partnerships at the individual level, as used in (Sih *et al.* 2009; Edenbrow *et al.*
al. 2011; Dakin & Ryder 2018). All else being equal, a highly specialized network is expected to
 284 be made up of individuals who are relatively more exclusive and/or more important towards their
 partners, *sensu* (Sih *et al.* 2009; Dakin & Ryder 2018). Because H_2' has the properties described
 286 in the previous paragraph, it is more appropriate as the network-level metric of social
 specialization. Note that it is possible, and perhaps even common, for the edge weights in a
 288 highly specialized network to be relatively invariant if the network is not fully connected, as
 shown in the example manakin networks in Fig. 1A. Hence, specialization based on H_2' is not in
 290 principle related to the network-average coefficient of variation of each individual's edge
 weights (Maldonado-Chaparro *et al.* 2018).

Network Stability

294 We define network stability as the average persistence of network edges through time (Dakin &
 Ryder 2020). To quantify the stability of manakin networks, we compared each lek's social
 296 network from a given recording session to its subsequent recording session within the same field
 season (Fig. 1B). Network stability was then calculated as the number of male-male partnerships
 298 (binary network edges) shared by both time points divided by the number of partnerships at
 either time point (Dakin & Ryder 2020). Higher values of stability indicate greater persistence of

social relationships within the network, independent of any changes in the representation of particular males (nodes) (Poisot *et al.* 2012). To focus this measure on the persistence of strong relationships, we computed network stability of partnerships that occurred at least six times following (Dakin & Ryder 2020). The threshold of six was chosen because it corresponds to an average rate of one social interaction per day in our data. We conducted an additional sensitivity analysis to verify that alternative thresholds for the stability calculation (greater or less than six) did not change our main results (see supplement for details). Previous work using this metric of stability has shown that the wire-tailed manakin social networks are far more stable than expected by chance (Dakin & Ryder 2020).

Behavioral Assortment

Assortment refers to the extent to which individuals associate with similar partners (Fig. 1C); it can be due to partner choice (homophily), shared environments, or the social transmission of behavior (Dakin & Ryder 2018). Assortment was quantified using Newman's assortativity index, which is a correlation coefficient for the statistical association among linked nodes within a network. It ranges from -1 (a negative association), through 0 (no association), and up to $+1$ (a positive association). To quantify the assortment of social behaviors, we first determined the daily frequency of two behaviors for each male: his number of social interactions per day (strength), and his number of unique social partnerships per day (degree). Strength and degree are both repeatable measures of a male's social behavior in our study population (Dakin & Ryder 2018). We used the average log-transformed values of each male's strength and degree within the recording session, and then calculated the assortativity coefficient for the entire social network using the algorithm for weighted networks in the assortnet package (Farine 2016).

Because assortativity values for strength and degree were highly correlated (Pearson's $r = 0.78$, $p < 0.0001$, $n = 86$ networks), we took the average of these two values as the measure of overall behavioral assortment within the social network. Note that we used log-transformed values of strength and degree because these two variables are strongly positively skewed (Dakin & Ryder 2018; Ryder *et al.* 2020), and assortativity is based on a Pearson's correlation. Finally, we also computed the assortativity of the two discrete status classes (floater and territory-holder), to ensure that our analysis of behavioral assortment was not solely driven by status-assortment.

Collective Testosterone

To understand how hormones might predict network properties, we derived a measure of collective testosterone of each social network. This was based on the hormonal trait that was the best predictor of social behavior in our previous study, referred to as "mean testosterone" (Ryder *et al.* 2020). A male's mean testosterone is his average residual circulating testosterone. It is calculated using a linear regression of log-transformed testosterone to statistically account for the effects of field season, Julian date, time of day when captured, and duration of restraint, all of which may influence point estimates of a male's baseline hormone level (Vernasco *et al.* 2019). Hence, mean testosterone represents a standardized measure of a male's circulating testosterone, independent of his capture conditions (Fig. 2A). Next, to determine collective testosterone, we took the average mean testosterone for each social network, weighted by the interaction frequency (strength) of the males within the network (Fig. 2B). Collective testosterone is thus a group-level characteristic that is weighted towards the males that made the greatest contribution to group social structure. In other words, networks with low collective testosterone are made up of mostly low-testosterone individuals, whereas networks with high collective testosterone are

made up of mostly high-testosterone individuals. We calculated collective testosterone separately for each status class, because the effects of hormones on social behavior are status-dependent (Eisenegger *et al.* 2011; Boksem *et al.* 2013; Ryder *et al.* 2020).

Statistical Analysis

To evaluate the hypothesis that collective testosterone predicts network properties, we analyzed mixed-effects models of the social network properties in the lme4 package (Bates *et al.* 2018). The three response variables were social specialization, network stability, and behavioral assortment (Fig. 1). We used Akaike's Information Criterion (AIC) to compare four candidate models for each response variable, as follows: (1) collective testosterone of territory-holders + collective testosterone of floaters; (2) collective testosterone of territory-holders; (3) collective testosterone of floaters, and (4) no testosterone predictors. All of the models included additional fixed effects to account for field season (a categorical variable with three levels), the average Julian date of the recording session, the average number of recorded hours per territory, and the size of the social network (number of individuals), as well as a random effect of lek to account for repeated measures.

Model comparison was performed on models fit with maximum likelihood, and the best-fit models were re-fit using restricted estimation of maximum likelihood (REML) to derive parameter estimates (Zuur *et al.* 2009). We used the lmerTest package to compute p-values for parameter estimates in the mixed-effects models based on Satterthwaite's method (Kuznetsova *et al.* 2018). We also report Nakagawa and Schielzeth's R^2_{LMM} values as an estimate of effect size (Nakagawa & Schielzeth 2013). We verified that all models met the assumptions of linear regression analyses. First, we checked that the Pearson residuals met the assumption of

normality. We visually inspected the partial residual plots for each fixed and random effect, to confirm that there were no outliers or departures from the assumption of homoscedasticity. To check for multicollinearity, we used the performance package (Lüdtke *et al.* 2019) to calculate variance inflation factors (VIFs), and we verified that all VIFs were between 1 – 1.8.

Sample Sizes and Exclusions

In two field seasons (2016-17 and 2017-18), we performed an experiment as part of a separate study to test the influence of transiently-elevated testosterone on individuals ($n = 5$ individuals in 2016-17 and $n = 4$ in 2017-18; Ryder *et al.* 2020). The results of that experiment demonstrated that elevated testosterone caused a temporary decrease in the frequency and the number of social partnerships in the altered males (Ryder *et al.* 2020). It is important to note that this experiment was not designed to test emergent properties at the network level, because it was conducted on a limited scale with only one or two individuals temporarily altered within each lek. We therefore excluded the 6 post-manipulation networks from the main analysis in this study. We verified that when we included these manipulated leks, all of our main conclusions were unchanged.

After excluding the 6 post-manipulation observations, 80 of the original 86 recording sessions remained. Table 1 summarizes the sample sizes for the network-level analyses. In 10 cases, specialization could not be calculated because the bipartite network did not have sufficient data to determine H_2' . Stability could not be calculated in 26 cases, when either the recording session occurred at the end of a field season, or when there were insufficient partnerships that met the criteria for the stability calculation.

Node-label Permutation Analysis

To evaluate the possibility that our results could be influenced by other properties of the leks that were independent of hormonal traits, we also performed a statistical permutation of the post-network data (Farine 2017). The purpose of this analysis was to verify that the results were driven by (and sensitive to) the relative contribution and social position of different males. To do this, we performed a statistical permutation that randomized the node labels (male IDs) within each of the social networks, retaining network topology, and leaving each male's testosterone traits unchanged. Hence, this analysis preserved which males were present in which recording session, but it randomized the relative position and contribution of each male. After generating 1,000 of these node-label permutation datasets, we recalculated specialization, stability, and assortment, and then refit the top models from our mixed-model analysis. We then compared the slope estimates from the observed data to those derived from 1,000 node-label permutations. As a one-sided p-value, we calculated the proportion of slope estimates from the node-label permutations that were more negative than the corresponding estimate from the observed data.

Results

The lek social networks had 14 males on average, but there was considerable variation among leks ranging from 3 to 43 individuals (Fig. S1). Additional descriptive statistics are provided in Table 1. Although network size has a lognormal distribution (Fig. S1), all other emergent properties of the social structure were approximately normally distributed (Fig. S4).

Table 1. Descriptive statistics for manakin social networks. Means and standard deviations (SD) are provided for each field season. The bottom row provides the sample sizes for the number of social networks analyzed (n_{obs}) at each lek (N_{lek}). Additional data is provided in the supplement in Fig. S1 and Fig. S4.

Field season	Network size (# birds)	Social specialization	Network stability	Behavioral assortment	Collective T (terr.)	Collective T (floa.)
15-16	14.5 (8.9)	0.50 (0.22)	0.43 (0.25)	0.01 (0.36)	0.44 (0.42)	-0.31 (0.56)
16-17	17.8 (10.5)	0.30 (0.24)	0.47 (0.20)	0.06 (0.30)	0.49 (0.24)	-0.22 (0.37)
17-18	10.3 (4.5)	0.42 (0.30)	0.42 (0.27)	0.02 (0.37)	0.43 (0.40)	-0.70 (0.46)
Sample size	$n_{\text{obs}} = 80$ $N_{\text{lek}} = 11$	$n_{\text{obs}} = 70$ $N_{\text{lek}} = 11$	$n_{\text{obs}} = 54$ $N_{\text{lek}} = 11$	$n_{\text{obs}} = 80$ $N_{\text{lek}} = 11$	$n_{\text{obs}} = 80$ $N_{\text{lek}} = 11$	$n_{\text{obs}} = 80$ $N_{\text{lek}} = 11$

We found that the collective testosterone of territorial males could predict all three emergent properties of the social networks (specialization, stability, and assortment). The leks with greater representation of high-testosterone territorial males were less specialized, less stable, and more negatively assorted (Fig. 3; all $p \leq 0.03$ in mixed-effects models). The slope coefficients for these three relationships were also all greater than expected under a node-label permutation (inset panels, Fig. 3; all $p < 0.02$). Model selection results for the observed data indicated considerable uncertainty in the best-fit model for each of the three network properties analyzed (Table S1). However, the collective testosterone of territory-holders was a significant predictor in all of the best-supported models (Table S2). In contrast, the collective testosterone of floater males was not a significant predictor of network properties in any of the best-fit models (Table S2).

We did not detect any significant effects of Julian date on network properties within our study period, but we did observe some significant year-to-year differences (e.g., social specialization and behavioral assortment in Table S2). Assortment was the only property that was significantly related to network size and recording effort (Table S2). All else being equal, behavioral assortment was more positive in larger networks, and it was more negative in networks that had longer recording sessions.

$R^2_{\text{LMM}(m)}$ provides an estimate of the proportion of variance explained by the fixed effects in a model (Table S1). The $R^2_{\text{LMM}(m)}$ for assortment in the best-fit model was 0.52. This indicates that about 52% the variation in behavioral assortment could be explained by the combined associations with collective testosterone, network size, sampling effort, and annual variation (Table S2). The $R^2_{\text{LMM}(m)}$ for stability was 0.12, indicating that collective testosterone and the other predictors (network size, sampling effort, year) could explain about 12% the variation in that metric (Table S2). Finally, for specialization, the $R^2_{\text{LMM}(m)}$ indicated that the combined effects of testosterone and these other sources of variation together explained about 20% of the variation (Table S2).

Discussion

The collective testosterone of the dominant, territory-holding males within a lek was associated with multiple emergent properties of the social network (Fig. 3). Variation in collective testosterone is a function of both the number of high-testosterone males and their frequency of social interactions. Our results indicate that the hormone-mediated behavior of these individuals may affect all three social network properties of specialization, stability and assortment. This indicates that the effects of testosterone on dominant males may mediate an extended phenotype with the power to shape social structure (Dawkins 1982). In contrast, the collective testosterone of floater males was not significantly associated with any of the emergent network properties (Table S2).

How can the relationship between hormone levels and network properties be explained in terms of individual mechanisms? Given that testosterone has antagonistic effects on the sociality of territorial males (Ryder *et al.* 2020; Vernasco *et al.* 2020), we hypothesize that the behavior of

high-testosterone individuals can cause several features of the social organization to break down.

Our previous results showed that high-testosterone dominant individuals have a reduced ability to attract and maintain social partners (Ryder *et al.* 2020). We propose that this weakening of social relationships may cause floater males to prospect elsewhere for new partners, negatively affecting both the stability and specialization of the lek network as a whole. Likewise, high-testosterone dominant males may inhibit the processes of social contagion, reciprocity, and/or behavioral matching that can cause positive behavioral assortment (Dakin & Ryder 2018). Recent studies have found that in other social animals, sparse and specialized social networks can be associated with fitness benefits (Stroeymeyt *et al.* 2018); hence, a breakdown to this organization may incur costs (Maldonado-Chaparro *et al.* 2018). Testing our proposed mechanism for the link between hormone-mediated behavior and network dynamics will require direct observation of the individual social behaviors that occur within dyads, and how these behaviors change through time. Because our current data cannot assess the fine-scale valence of social interactions, further studies are needed that combine direct observation with high-throughput data on social network dynamics.

We included several additional parameters in our analyses to account for social network size and sampling effort. Although it was not one of our main hypotheses, we noted that network size was positively associated with behavioral assortment (Table S2). In other words, males were more likely to associate with behaviorally similar partners in larger leks, whereas they were more likely to associate with dissimilar partners in smaller leks. Effects of group size on assortment have been noted in a few other studies, although the form of this relationship varies (Griffiths & Magurran 1997; Ilmarinen *et al.* 2017; McDonald *et al.* 2017). In manakins and other lekking systems, larger leks are known to have heightened display activity and higher female visitation

rates (Lank & Smith 1992; Durães *et al.* 2009). This raises the possibility that social facilitation and heightened activity may be associated with increased homophily and/or behavioral matching (either through partner choice, contagion, or reciprocity). Another plausible explanation is that smaller social groups may constrain these behavioral processes, by making it more difficult to find or match a suitable partner. This hypothesis could be explored in experiments on captive systems and simulation models.

Behavioral assortment was also more positive in networks that were recorded for less time in our study (in other words, leks where males associated with behaviorally similar partners tended to be recorded for fewer days). Recording time was designed to be approximately even among leks (Fig. S1), with the most common reason for an extended recording duration being inclement weather that extended the recording session. Therefore, we speculate that inclement weather may have reduced the amount of behavioral matching (assortment) while also affecting recording time. Although we did not collect data on weather at each lek as this was not our main goal, the question of how inclement weather affects group-level social dynamics is an interesting one that merits further study.

Conducting experimental tests of causation at the level of whole social networks remains a major challenge in ecology (Pinter-Wollman *et al.* 2014; James *et al.* 2009). Although our node-label permutation analysis provides evidence that our results are not merely due to structural differences among networks, we cannot rule out the possibility that high-testosterone individuals chose to participate in certain networks due to other factors that may also influence the emergent properties of the network (e.g., environmental quality and/or female activity). It is important to note that in this system, as in many other wild animals, controlled experimental manipulations of the broader social network structure are not yet possible (Zyphur *et al.* 2009;

Akinola *et al.* 2016). Nevertheless, our data here indicate that the increased prevalence of dominant, high-testosterone individuals can predict changes social dynamics and subsequent higher-order network structure. These findings establish that hormone-behavior relationships are not limited to one individual, but instead hormones have population-level consequences (McClintock 1981; Robinson 1992).

Increasing evidence demonstrates that the structure and stability of social networks is often associated with benefits—and costs—during foraging, breeding, and disease outbreaks (Silk *et al.* 2010; Maldonado-Chaparro *et al.* 2018; Riehl & Strong 2018; Stroeymeyt *et al.* 2018). Given the widespread influence of steroid hormones on social interactions across vertebrates (Adkins-Regan 2005), we expect that collective testosterone will be broadly associated with social network properties in other systems. The direction of these effects may depend on the context of the behavioral interactions that form the social network, and whether these interactions are primarily cooperative or competitive in nature. Taken together with previous studies, we propose that testosterone-mediated behavior can alter social network dynamics in ways that often impede the evolution of stable social relationships and cooperation.

520 Figure Legends

522 **Figure 1. Example social networks illustrating social specialization, network stability, and**
behavioral assortment. (A) Social specialization is measured using the bipartite form of the
524 social network. In the example on the left, the associations between floater males and territory-
holders are poorly partitioned, creating a network with a relatively low specialization. On the
526 right, there is greater partitioning, such that each floater male maintains a smaller number of
associations with specific territory-holding males. (B) Network stability is measured by
528 evaluating the persistence of partnerships from one recording session (t_1) to the next (t_2). The
example at the top of (B) has a lower stability than the example on the bottom. (C) Behavioral
530 assortment measures the tendency of like to associate with like. In (C), nodes are shaded to
indicate a continuous gradient of daily behavior (in this case, each male's average daily strength
532 is shown). In the example network on the left, the males tend to associate with behaviorally
dissimilar partners, yielding a negative behavioral assortment. In the example on the right, the
534 males tend to associate with behaviorally similar partners, yielding a positive assortment. See
Table 1 for additional data.

536

Figure 2. Collective testosterone of manakin social networks. (A) Circulating testosterone
538 varies among individual males. This graph shows repeated measures from 210 individual male
manakins, sorted along the x-axis by mean testosterone, a standardized measure of a male's
540 average circulating testosterone level (Ryder *et al.* 2020). Opacity is used to denote the two male
status classes (with subordinate floaters colored semi-transparent, and dominant territory-holders
542 colored opaque). Color ramping is used to denote each individual's hormone phenotype. (B-D)

These data were used to define collective testosterone in the present study. Node colors indicate an individual's mean testosterone following the scale in (A). The collective testosterone of each social network is calculated as the average of the individual hormone phenotypes, weighted by strength, and is shown below each example. Collective testosterone was determined for each status class separately. See Table 1 for additional data.

Figure 3. Collective testosterone predicts the structure of the social network. The networks

with higher-testosterone dominant males were less specialized, less stable over time, and had more negative behavioral assortment. Each plot shows the partial residuals from a statistical analysis that also accounts for field season, the average Julian date of the recording session, the number of recording hours, network size, and lek identity. Because floater and territorial males can differ in behavior, the analysis of behavioral assortment also accounted for status assortment within each network. Different symbols are used to indicate repeated measures of 11 different leks, colored according to the collective testosterone scale in Fig. 2. Inset panels show the results of node-label permutation tests. In each case, the slope in the best-fit model (solid vertical line) is significantly more negative than expected based on the distribution of permuted estimates (grey distribution, dotted line is at 0). See Tables S1-S2 for additional data.

560 **Acknowledgments**

We thank Camilo Alfonso, Brian Evans, David and Consuelo Romo, Kelly Swing, Diego
562 Mosquera, Gabriela Vinueza, and the Tiputini Biodiversity Station of the Universidad San
Francisco de Quito. Funding was provided by the National Science Foundation (NSF) IOS
564 1353085 and the Smithsonian Migratory Bird Center.

566 **Author Contributions**

RD, ITM, BMH, and TBR designed the study. ITM, BMH, BJV, and TBR collected the data.
568 RD, ITM, and TBR analyzed the data. RD and TBR wrote the manuscript. All authors edited the
manuscript.

570

Declaration of Interests

572 The authors declare no competing interests.

574 **Data Availability**

All data and R scripts necessary to reproduce this study are available for download at:
576 <https://figshare.com/s/13a311662fee686fa4f3>

578 **Supporting Information**

Supplemental text, Figures S1-S5, and Table S1-S3 in the attached PDF

580

References

- Adkins-Regan, E. (2005). *Hormones and Animal Social Behavior*. Princeton University Press.
- Akinola, M., Page-Gould, E., Mehta, P.H. & Lu, J.G. (2016). Collective hormonal profiles predict group performance. *Proceedings of the National Academy of Sciences*, 113, 9774–9779.
- Aplin, L.M., Farine, D.R., Morand-Ferron, J. & Sheldon, B.C. (2012). Social networks predict patch discovery in a wild population of songbirds. *Proceedings of the Royal Society B: Biological Sciences*, 279, 4199–4205.
- Bates, D., Maechler, M., Bolker, B., Walker, S., Christensen, R.H.B., Singmann, H., et al. (2018). *lme4 1.1-18-1: linear mixed-effects models using “Eigen” and S4*.
- Blüthgen, N., Menzel, F. & Blüthgen, N. (2006). Measuring specialization in species interaction networks. *BMC Ecology*, 6, 9.
- Boksem, M.A.S., Mehta, P.H., Van den Bergh, B., van Son, V., Trautmann, S.T., Roelofs, K., et al. (2013). Testosterone inhibits trust but promotes reciprocity. *Psychological Science*, 24, 2306–2314.
- Cohen, A.A., Martin, L.B., Wingfield, J.C., McWilliams, S.R. & Dunne, J.A. (2012). Physiological regulatory networks: ecological roles and evolutionary constraints. *Trends in Ecology & Evolution*, 27, 428–435.
- Croft, D.P., James, R., Thomas, P.O.R., Hathaway, C., Mawdsley, D., Laland, K.N., et al. (2006). Social structure and co-operative interactions in a wild population of guppies (*Poecilia reticulata*). *Behavioral Ecology and Sociobiology*, 59, 644–650.
- Croft, D.P., Madden, J.R., Franks, D.W. & James, R. (2011). Hypothesis testing in animal social networks. *Trends in Ecology & Evolution*, 26, 502–507.
- Csardi, G., Nepusz, T. & and 84 other coauthors. (2018). *igraph 1.2.2: network analysis and visualization*.
- Dakin, R. & Ryder, T.B. (2018). Dynamic network partnerships and social contagion drive cooperation. *Proceedings of the Royal Society B: Biological Sciences*, 285, 20181973.
- Dakin, R. & Ryder, T.B. (2020). Reciprocity and behavioral heterogeneity govern the stability of social networks. *Proceedings of the National Academy of Sciences*, 117, 2993–2999.
- Dawkins, R. (1982). *The Extended Phenotype: The Gene as the Unit of Selection*. W. H. Freeman and Company, San Francisco.
- Díaz-Muñoz, S.L., DuVal, E.H., Krakauer, A.H. & Lacey, E.A. (2014). Cooperating to compete: altruism, sexual selection and causes of male reproductive cooperation. *Animal Behaviour*, 88, 67–78.
- Dormann, C.F., Fruend, J. & Gruber, B. (2019). *bipartite 2.13: visualising bipartite networks and calculating some (ecological) indices*.
- Durães, R., Loiselle, B.A., Parker, P.G. & Blake, J.G. (2009). Female mate choice across spatial scales: influence of lek and male attributes on mating success of blue-crowned manakins. *Proceedings of the Royal Society B: Biological Sciences*, 276, 1875–1881.
- DuVal, E.H. (2007). Adaptive advantages of cooperative courtship for subordinate male lance-tailed manakins. *The American Naturalist*, 169, 423–432.
- Edenbrow, M., Darden, S.K., Ramnarine, I.W., Evans, J.P., James, R. & Croft, D.P. (2011). Environmental effects on social interaction networks and male reproductive behaviour in guppies, *Poecilia reticulata*. *Animal Behaviour*, 81, 551–558.

624 Eikenaar, C., Whitham, M., Komdeur, J., van der Velde, M. & Moore, I.T. (2011). Endogenous
testosterone is not associated with the trade-off between paternal and mating effort.
626 *Behavioral Ecology*, 22, 601–608.

Eisenegger, C., Haushofer, J. & Fehr, E. (2011). The role of testosterone in social interaction.
628 *Trends in Cognitive Sciences*, 15, 263–271.

Farine, D.R. (2014). Measuring phenotypic assortment in animal social networks: weighted
630 associations are more robust than binary edges. *Animal Behaviour*, 89, 141–153.

Farine, D.R. (2015). Proximity as a proxy for interactions: issues of scale in social network
632 analysis. *Animal Behaviour*, 104, e1–e5.

Farine, D.R. (2016). *assortnet 0.12: calculate the assortativity coefficient of weighted and binary
634 networks*.

Farine, D.R. (2017). A guide to null models for animal social network analysis. *Methods in
636 Ecology and Evolution*, 8, 1309–1320.

Fowler, J.H. & Christakis, N.A. (2010). Cooperative behavior cascades in human social networks.
638 *Proceedings of the National Academy of Sciences*, 107, 5334–5338.

Franks, D.W., Ruxton, G.D. & James, R. (2010). Sampling animal association networks with the
640 gambit of the group. *Behavioral Ecology and Sociobiology*, 64, 493–503.

Fuxjager, M.J., Forbes-Lorman, R.M., Coss, D.J., Auger, C.J., Auger, A.P. & Marler, C.A. (2010).
642 Winning territorial disputes selectively enhances androgen sensitivity in neural
pathways related to motivation and social aggression. *Proceedings of the National
644 Academy of Sciences*, 107, 12393–12398.

Goymann, W. (2009). Social modulation of androgens in male birds. *General and Comparative
646 Endocrinology*, 9th International Symposium on Avian Endocrinology, 163, 149–157.

Goymann, W., Schwabl, I., Trappschuh, M. & Hau, M. (2007). Use of ethanol for preserving
648 steroid and indoleamine hormones in bird plasma. *General and Comparative
Endocrinology*, 150, 191–195.

Griffiths, S.W. & Magurran, A.E. (1997). Schooling preferences for familiar fish vary with group
650 size in a wild guppy population. *Proceedings of the Royal Society of London. Series B:
652 Biological Sciences*, 264, 547–551.

Heindl, M. (2002). Social organization on leks of the wire-tailed manakin in Southern Venezuela.
654 *The Condor*, 104, 772–779.

Ilmarinen, V.-J., Vainikainen, M.-P., Verkasalo, M.J. & Lönnqvist, J.-E. (2017). Homophilous
656 friendship assortment based on personality traits and cognitive ability in middle
childhood: the moderating effect of peer network size. *European Journal of Personality*,
658 31, 208–219.

James, R., Croft, D.P. & Krause, J. (2009). Potential banana skins in animal social network
660 analysis. *Behavioral Ecology and Sociobiology*, 63, 989–997.

Jehn, K.A. & Shah, P.P. (1997). Interpersonal relationships and task performance: an
662 examination of mediation processes in friendship and acquaintance groups. *Journal of
Personality and Social Psychology*, 72, 775–790.

664 Krause, J. & Ruxton, G.D. (2002). *Living in Groups*. OUP Oxford.

Kuznetsova, A., Brockhoff, P.B. & Christensen, R.H.B. (2018). *lmerTest 3.0-1: tests in linear
666 mixed effects models*.

- Lank, D.B. & Smith, C.M. (1992). Females prefer larger leks: field experiments with ruffs
668 (*Philomachus pugnax*). *Behavioral Ecology and Sociobiology*, 30, 323–329.
- Lüdecke, D., Makowski, D. & Waggoner, P. (2019). *performance 0.4.2: assessment of regression*
670 *models performance*.
- Maldonado-Chaparro, A.A., Alarcón-Nieto, G., Klarevas-Irby, J.A. & Farine, D.R. (2018).
672 Experimental disturbances reveal group-level costs of social instability. *Proceedings of*
the Royal Society B: Biological Sciences, 285, 20181577.
- McClintock, M.K. (1981). Social control of the ovarian cycle and the function of estrous
674 synchrony. *Integrative and Comparative Biology*, 21, 243–256.
- McDonald, D.B. (1993). Delayed plumage maturation and orderly queues for status: a manakin
676 mannequin experiment. *Ethology*, 94, 31–45.
- McDonald, D.B. (2007). Predicting fate from early connectivity in a social network. *Proceedings*
678 *of the National Academy of Sciences*, 104, 10910–10914.
- McDonald, G.C., Farine, D.R., Foster, K.R. & Biernaskie, J.M. (2017). Assortment and the analysis
680 of natural selection on social traits. *Evolution*, 71, 2693–2702.
- Nakagawa, S. & Schielzeth, H. (2013). A general and simple method for obtaining R² from
682 generalized linear mixed-effects models. *Methods in Ecology and Evolution*, 4, 133–142.
- Ohtsuki, H., Hauert, C., Lieberman, E. & Nowak, M.A. (2006). A simple rule for the evolution of
684 cooperation on graphs and social networks. *Nature*, 441, 502–505.
- Oyegbile, T.O. & Marler, C.A. (2005). Winning fights elevates testosterone levels in California
686 mice and enhances future ability to win fights. *Hormones and Behavior*, 48, 259–267.
- Pinter-Wollman, N., Hobson, E.A., Smith, J.E., Edelman, A.J., Shizuka, D., de Silva, S., *et al.*
688 (2014). The dynamics of animal social networks: analytical, conceptual, and theoretical
690 advances. *Behavioral Ecology*, 25, 242–255.
- Poisot, T., Canard, E., Mouillot, D., Mouquet, N. & Gravel, D. (2012). The dissimilarity of species
692 interaction networks. *Ecology Letters*, 15, 1353–1361.
- Prum, R.O. (1994). Phylogenetic analysis of the evolution of alternative social behavior in the
694 manakins (Aves: Pipridae). *Evolution*, 48, 1657–1675.
- R Core Team. (2018). *R 3.5.1: A Language and Environment for Statistical Computing*. R
696 Foundation for Statistical Computing, Vienna, Austria.
- Riehl, C. & Strong, M.J. (2018). Stable social relationships between unrelated females increase
698 individual fitness in a cooperative bird. *Proceedings of the Royal Society B: Biological*
Sciences, 285, 20180130.
- Roberts, G. & Sherratt, T.N. (1998). Development of cooperative relationships through
700 increasing investment. *Nature*, 394, 175.
- Robinson, G.E. (1992). Regulation of division of labor in insect societies. *Annual Review of*
702 *Entomology*, 37, 637–665.
- Ryder, T.B., Blake, J.G., Parker, P.G. & Loiselle, B.A. (2011a). The composition, stability, and
704 kinship of reproductive coalitions in a lekking bird. *Behavioral Ecology*, 22, 282–290.
- Ryder, T.B., Dakin, R., Vernasco, B.J., Evans, B.S., Horton, B.M. & Moore, I.T. (2020).
706 Testosterone modulates status-specific patterns of cooperation in a social network. *The*
708 *American Naturalist*.
- Ryder, T.B., Horton, B.M. & Moore, I.T. (2011b). Understanding testosterone variation in a
710 tropical lek-breeding bird. *Biology Letters*, 7, 506–509.

- Ryder, T.B., Horton, B.M., Tillaart, M. van den, Morales, J.D.D. & Moore, I.T. (2012). Proximity data-loggers increase the quantity and quality of social network data. *Biology Letters*, 8, 917–920.
- Ryder, T.B., McDonald, D.B., Blake, J.G., Parker, P.G. & Loiselle, B.A. (2008). Social networks in the lek-mating wire-tailed manakin (*Pipra filicauda*). *Proceedings of the Royal Society B: Biological Sciences*, 275, 1367–1374.
- Ryder, T.B., Parker, P.G., Blake, J.G. & Loiselle, B.A. (2009). It takes two to tango: reproductive skew and social correlates of male mating success in a lek-breeding bird. *Proceedings of the Royal Society B: Biological Sciences*, 276, 2377–2384.
- Sah, P., Leu, S.T., Cross, P.C., Hudson, P.J. & Bansal, S. (2017). Unraveling the disease consequences and mechanisms of modular structure in animal social networks. *Proceedings of the National Academy of Sciences*, 114, 4165–4170.
- Sah, P., Mann, J. & Bansal, S. (2018). Disease implications of animal social network structure: A synthesis across social systems. *Journal of Animal Ecology*, 87, 546–558.
- Schjelderup-Ebbe, T. (1922). Beiträge zur sozialpsychologie des haushuhns. [Observation on the social psychology of domestic fowls.]. *Zeitschrift für Psychologie und Physiologie der Sinnesorgane. Abt. 1. Zeitschrift für Psychologie*, 88, 225–252.
- Schwartz, P. & Snow, D.W. (1978). Display and related behavior of the wire-tailed manakin. *Living Bird*, 17, 51–78.
- Sih, A., Hanser, S.F. & McHugh, K.A. (2009). Social network theory: new insights and issues for behavioral ecologists. *Behav Ecol Sociobiol*, 63, 975–988.
- Silk, J.B., Beehner, J.C., Bergman, T.J., Crockford, C., Engh, A.L., Moscovice, L.R., et al. (2010). Strong and consistent social bonds enhance the longevity of female baboons. *Current Biology*, 20, 1359–1361.
- Stroeymeyt, N., Grasse, A.V., Crespi, A., Mersch, D.P., Cremer, S. & Keller, L. (2018). Social network plasticity decreases disease transmission in a eusocial insect. *Science*, 362, 941–945.
- Trainer, J.M. & McDonald, D.B. (1995). Singing performance, frequency matching and courtship success of long-tailed manakins (*Chiroxiphia linearis*). *Behavioral Ecology and Sociobiology*, 37, 249–254.
- Trainer, J.M., McDonald, D.B. & Learn, W.A. (2002). The development of coordinated singing in cooperatively displaying long-tailed manakins. *Behavioral Ecology*, 13, 65–69.
- Trivers, R.L. (1971). The evolution of reciprocal altruism. *The Quarterly Review of Biology*, 46, 35–57.
- Vernasco, B.J., Horton, B.M., Moore, I.T. & Ryder, T.B. (2020). Reduced cooperative behavior as a cost of high testosterone in a lekking passerine bird. *Behavioral Ecology*, 31, 401–410.
- Vernasco, B.J., Horton, B.M., Ryder, T.B. & Moore, I.T. (2019). Sampling baseline androgens in free-living passerines: Methodological considerations and solutions. *General and Comparative Endocrinology*, Endocrinology of Neotropical Vertebrates, 273, 202–208.
- Wingfield, J.C., Hegner, R.E., Dufty, Alfred M. & Ball, G.F. (1990). The “challenge hypothesis”: theoretical implications for patterns of testosterone secretion, mating systems, and breeding strategies. *The American Naturalist*, 136, 829–846.
- Zuur, A., Ieno, E.N., Walker, N., Saveliev, A.A. & Smith, G.M. (2009). *Mixed Effects Models and Extensions in Ecology with R*. Springer Science & Business Media.

756 Zyphur, M.J., Narayanan, J., Koh, G. & Koh, D. (2009). Testosterone–status mismatch lowers
collective efficacy in groups: evidence from a slope-as-predictor multilevel structural
equation model. *Organizational Behavior and Human Decision Processes*, Biological
758 Basis of Business, 110, 70–79.

Figure 1

Figure 2

Figure 3