CD13 is a Critical Regulator of Cell-cell Fusion in Osteoclastogenesis

Mallika Ghosh^{1*}, Ivo Kalajzic², Hector Leonardo Aguila^{1,3} and Linda H Shapiro,^{1*}

Centers for ¹Vascular Biology, ²Regenerative Medicine and Skeletal Development

³Departments of Immunology, University of Connecticut Medical School, Farmington, Connecticut 06032, USA

***Corresponding authors:**

Linda H Shapiro, PhD
Center for Vascular Biology
University of Connecticut Medical School
Farmington, CT 06030
Email: lshapiro@uchc.edu
Telephone: 860-679-4373
Fax:860-679-1201

Mallika Ghosh, PhD
Center for Vascular Biology
University of Connecticut Medical School
Farmington, CT 06030
Email: mghosh@uchc.edu
Telephone: 860-679-1189
Fax:860-679-1201

Conflict of interest: The authors have declared that no conflict of interest exists.

Abstract

In vertebrates, bone formation is dynamically controlled by the activity of two specialized cell types: the bone-generating osteoblasts and bone-degrading osteoclasts. Osteoblasts produce the soluble receptor activator of NF κ B ligand (RANKL) that binds to its receptor RANK on the surface of osteoclast precursor cells to promote osteoclastogenesis, a process that involves cell-cell fusion and assembly of molecular machinery to ultimately degrade the bone. CD13 is a transmembrane aminopeptidase that is highly expressed in cells of myeloid lineage has been shown to regulate dynamin-dependent receptor endocytosis and recycling and is a necessary component of actin cytoskeletal organization. In the present study, we show that CD13-deficient mice display a normal distribution of osteoclast progenitor populations in the bone marrow, but present a low bone density phenotype. Further, the endosteal bone formation rate is similar between genotypes, indicating a defect in osteoclast-specific function *in vivo*. Loss of CD13 led to exaggerated *in vitro* osteoclastogenesis as indicated by significantly enhanced fusion of bone marrow-derived multinucleated osteoclasts in the presence of M-CSF and RANKL, resulting in abnormally large cells with remarkably high numbers of nuclei with a concomitant increase in bone resorption activity. Similarly, we also observed increased formation of multinucleated giant cells (MGC) in CD13^{KO} bone marrow progenitor cells stimulated with IL-4 and IL-13, suggesting that CD13 may regulate cell-cell fusion events via a common pathway, independent of RANKL signaling. Mechanistically, while expression levels of the fusion-regulatory proteins dynamin and DC-STAMP are normally downregulated as fusion progresses in fusion-competent mononucleated progenitor cells, in the absence of CD13 they are uniformly sustained at high levels, even in mature multi-nucleated osteoclasts. Taken together, we conclude that CD13 may regulate cell-cell fusion by controlling expression and localization of key fusion proteins that are critical for both osteoclast and MGC fusion.

Introduction

Osteoclastogenesis is a critical process for skeletal growth and development that is tightly regulated by differentiation of myeloid progenitor cells into osteoclasts, which are specialized cells in the bone marrow whose major function is bone resorption¹. This highly-regulated process is responsible for bone modeling and remodeling that ultimately translates into maintenance of bone integrity, skeletal growth and repair^{2,3}. Deregulation of the process leads to dramatic outcomes^{4,5}. Loss of osteoclast (OC) generation/function gives rise to elevated bone formation without remodeling, resulting in osteopetrosis characterized by high bone mass and growth impairment,

while gain of OC generation/function results in exacerbated bone degradation that, without equivalent coupling to bone formation, leads to osteoporosis, characterized by low bone mass, bone weakness and high predisposition to fractures with poor healing progression. OCs form from committed monocyte progenitors via initial signals provided by two central bone marrow-derived cytokines: macrophage colony stimulating factor (M-CSF) and receptor activator of NF κ B ligand (RANKL), initiating a highly-organized program of commitment towards terminal differentiation and function⁶⁻⁸. This process involves proteins that participate in cell-cell fusion which are critical to the generation of functionally active OCs. Despite our current knowledge of events leading to the formation of OCs, understanding the regulation of these processes in homeostatic as well as their de-regulation in pathological conditions is unclear⁹⁻¹¹.

CD13 is a transmembrane metalloprotease widely expressed in all cells of the myeloid lineage, as well as activated endothelial cells, hematopoietic progenitor and stem cells¹²⁻¹⁶. Previous studies have implicated CD13 in vasculogenesis, tumor cell invasion, inflammatory trafficking and as a receptor for human corona virus¹⁷⁻¹⁹. Our novel observations have clearly shown that independent of its peptidase activity, CD13 is a critical molecule that assembles the molecular machinery enabling diverse cellular processes such as cell-cell adhesion, migration, membrane organization and dynamin-mediated receptor endocytosis and recycling of cell surface proteins^{14,15,20-22}. Taken together, CD13 regulates many of the activities that have been described to be critical for osteoclastogenesis and cell-cell fusion. In the current study, we demonstrate that despite a relatively normal distribution of hematopoietic components in bone marrow and periphery, CD13^{KO} mice have reduced bone mass with increased OC numbers per bone surface area but normal bone formation parameters, indicating that osteoclastogenesis is compromised in the absence of CD13. In addition, the in vitro induction of CD13-deficient myeloid progenitors generated from bone marrow and spleen resulted in increased numbers of OCs that were considerably larger in size, contained many more nuclei and resorbed bone more efficiently than those from wild type progenitors. We confirmed that CD13-deficient macrophages can also hyper-fuse to generate elevated multinucleated giant cells (MGCs) which again, were larger and contained more nuclei than those generated from wild type macrophages, suggesting that CD13 is a component of common fusion pathways shared by OC and MGC. Furthermore, we demonstrated that expression of fusion proteins which is typically downregulated in mature osteoclasts post-fusion²³ is abnormally preserved in osteoclasts lacking CD13. These findings are in agreement with our data showing that CD13 is a mediator of homotypic cell interaction and a regulator of molecular events defining cell membrane organization, fluidity and movement, all processes critical to cell-cell fusion^{14,15,20}. We hypothesize that CD13 is a negative regulator of

cell-cell fusion in osteoclastogenesis and giant cell formation and potentially, a universal modulator of membrane fusion and is a novel target for therapeutic intervention in pathological conditions mediated by defects in cell-cell fusion.

Results

Bone density is reduced CD13^{KO} mice *in vivo*

Based on the notion that the high, sustained expression of CD13 in all cells of the myeloid lineage reflects its important role in myeloid cell biology, we and others have demonstrated that it contributes to many fundamental cellular processes that impact myeloid cell function in various tissues. These studies prompted our current focus on the myeloid cells of the bone, the osteoclasts. We initially examined the effect of a global loss of CD13 on the phenotype of developing bone. Analysis of bone micro-architecture and function in the cortical and trabecular bone isolated from 8-10 wks. old WT and CD13^{KO} male mice by μ -CT and histomorphometric analysis revealed that the femur cortical and trabecular bone density and thickness in CD13^{KO} mice is significantly reduced (1.5 fold) compared to WT animals (**FIG. 1A, B**). **Immunohistochemical analysis of TRAP stained frozen sections of the femur clearly indicated elevated TRAP⁺ osteoclasts (in purple) in CD13^{KO} femur compared to the wildtype counterpart (FIG. 1C).** Reduced bone volume/total volume (BV/TV, **1D**), number of osteoclasts per bone surface (OC/BS, **1E**) and per bone perimeter (OC/B.Pm, **1F**) with similar endosteal bone formation rates per bone surface (BFR/BS, **1G**), together, strongly suggested a defect in bone remodeling in the absence of CD13. Osteoblast cultures from BM stained for alkaline phosphatase activity indicated that alkaline phosphatase (ALP)⁺ osteoblast surface per well was equivalent between genotypes (data not shown) supporting a contribution of CD13 to osteoclast activity *in vivo* and *in vitro*.

OCP populations with osteoclastogenic potential are similar in WT and CD13^{KO} bone marrow.

Previously we have shown that the distribution of the hematopoietic population comprised of early hematopoietic progenitors, myelo-erythroid progenitors, common myeloid progenitors, and granulocyte macrophage progenitors in CD13^{KO} mice were similar to wildtype animals¹². Cells that can generate bone-resorbing osteoclast reside in both bone marrow and peripheral hematopoietic organs²⁴. To determine if differences in osteoclast progenitor (OCP) frequency are responsible for the loss of bone mass in the absence of CD13, we analyzed the distribution of

primary OCP in both the bone marrow microenvironment and spleen in WT and CD13^{KO} mice. Flow cytometric analysis (**FIG. 2A,B**) revealed that the OCP profile indicated by CD3-, B220-, NK1.1-, CD11b-/lo, CD115+, CD117+ in the BM (**A**, WT vs. CD13^{KO}; 1.7 vs. 1.95) and CD3-, B220-, NK1.1-, CD11b+, Ly6G-, Ly6C+, CD115+ in spleen (**B**, WT vs. CD13^{KO}; 0.24 vs. 0.238)²⁴ is similar between genotypes, indicating that the absence of CD13 does not change the differentiation potential of myeloid cells to osteoclast progenitors.

In vitro osteoclastogenesis is exaggerated in CD13^{KO} cells.

Stimulation of monocyte lineage-committed hematopoietic progenitors with two principal bone marrow cytokines, M-CSF and RANKL, triggers the expression of molecules involved in cell-cell fusion and functional bone resorption⁶⁻⁸ and the generation of multinucleated osteoclasts. The normal hematopoietic profiles in CD13^{KO} mice, taken together with the defective skeletal phenotype, suggested that CD13 may participate in osteoclastogenesis in vitro. Analysis of flow sorted, bone marrow-derived OCP (CD3- B220- NK1.1- CD11b -/lo CD115+ Ly6C+, **FIG. S1A**) differentiated to mature OC in the presence of recombinant M-CSF (30ng/ml) and RANKL (30ng/ml) stained with TRAP, revealed significant increases in the number of osteoclasts containing >3 nuclei per field (2.8 fold), the average OC size (4-fold) and the number of nuclei/OC (2.5-fold) in CD13^{KO} cells grown on plastic (not shown) or bovine cortical bone slices (**FIG. 3A-D**) compared to those generated from normal WT progenitors. This difference was evident by d3, suggesting that the lack of CD13 accelerates OC fusion and multinucleation but not OCP proliferation rates, as the cell density (total number of nuclei/dish) was not significantly different between genotypes (**FIG. 3G**). Furthermore, we examined the flow sorted spleen-derived OCP (CD3- B220- NK1.1- CD11b+ Ly6G- Ly6C+ CD115+), which were allowed to proliferate in presence of M-CSF for d3, followed by differentiation into mature OCs by RANKL treatment for an additional 3 days. Similar to BM-derived cells, CD13^{KO}-derived splenic cells produced OCs with larger area (3-fold), more cells with >3 nuclei (2.5-fold) and an increased number of nuclei per cell (3-fold) compared to WT mice (**FIG. 4A-D**). To assess OC bone resorptive capacity, we plated WT and CD13^{KO} flow-sorted bone marrow-derived OCP on Osteo assay plates (Corning) in the presence of recombinant M-CSF and RANKL and allowed them to mature to OC over time. At d10, OCs were removed, individual or multiple resorption pit areas were imaged and the area of resorption quantified by ImageJ. As expected, the increase in OC nuclei/cell positively correlates with resorption and CD13^{KO} OCs showed increased resorption areas (3-fold) compared to WT, confirming that the elevated osteoclastogenesis in CD13^{KO} mice translates into

exaggerated functional activity (**FIG. 3E, F**). Our in vitro data is consistent with our in vivo bone phenotype in the absence of CD13.

Macrophage giant cell formation (MGC) is enhanced in CD13^{KO} cells.

In addition to osteoclasts and independent of RANKL, monocyte progenitors can differentiate into macrophages which also undergo cell-cell fusion to form large, multi-nucleated macrophage giant cells (MGC)^{25,26}. To evaluate if CD13 also regulates MGC fusion, we expanded sorted BM myeloid progenitor cells (CD3- B220- NK1.1- CD11b -/lo CD115+ Ly6C+) in M-CSF for 5 days, followed by the addition of IL-4 (30ng/ml) and IL-13 (30ng/ml) for 3 days to promote macrophage fusion. Images of Giemsa stained cells indicated a significant increase in the number of cells per field with >3 nuclei (2.5-fold), increased area/cell and a greater number of nuclei/cell (5-fold) of MGCs in CD13^{KO} cultures compared to WT (**FIG. 5A-D**). Immunofluorescence and confocal microscopy further confirmed CD13 expression in CD68+ MGC (**FIG. 5E**). In agreement with CD13 as a cell surface mediator of fusion in macrophages, CD13 and the fusion protein DC-STAMP (DCST1)²⁷ are co-localized in MGC stimulated with IL-4/IL-13 (**FIG. 5F**). Our data strongly suggest that CD13 regulates a common cell fusion event independent of RANKL signaling²⁸.

Expression of fusion proteins is dysregulated in CD13^{KO} mature osteoclasts

Pertinent to our observations, it is believed that since OCs and macrophages are derived from common progenitors, some of the molecular mechanisms mediating their fusion and multinucleation may be shared²⁸. In particular, the small GTPase dynamin 2 (the major isoform in OC) and the fusion protein DC-STAMP are common and critical regulators of both osteoclast and MGC fusion^{23,27}. Our recent studies have shown that CD13 is a potent negative regulator of dynamin-dependent endocytosis of a variety of receptors²⁰⁻²², suggesting that CD13 may participate in cell fusion by regulating endocytic processes. Indeed, immunoblot (**FIG. 6A**) and immunofluorescence (**FIG. 7**) analyses of flow-sorted WT BM-OCPs differentiated to OC with M-CSF and RANKL demonstrated high levels of dynamin and DC-STAMP expression by d2-post differentiation, which was subsequently reduced by 3d, when cell fusion and maturation into multinucleated WT osteoclasts is complete, as previously reported²³. However, while dynamin and DC-STAMP are highly expressed in CD13^{KO} OCP, rather than being downregulated, this strong expression is maintained in mature multinucleated OCs (**FIG. 6A,7**), suggesting that CD13 may impact fusion by regulating the levels of key fusion and/or endocytic molecules critical for OC and MGC fusion. In addition, immunoblot analysis of cell lysates obtained from WT bone marrow-derived progenitor cells stimulated with M-CSF and RANKL over 3d indicated that CD13

is highly expressed in myeloid progenitor cells but its expression level is unaltered upon stimulation with M-CSF and RANKL over time (d0-3) (**FIG. 6B**), consistent with CD13 regulating fusion mechanisms independent of RANKL signaling. Further investigation into the relationship between CD13 and these and other common regulators of myeloid cell fusion such as OC-STAMP, the macrophage fusion receptor, osteoclast receptor $\alpha v \beta 3$ integrin²⁸ will clarify common mechanisms regulating CD13-dependent myeloid cell fusion as well as the fusion of other lineages including satellite stem cells.

Discussion

The fusion of plasma membranes is essential to and indispensable for many physiologic processes such as fertilization through sperm/egg fusion²⁹, muscular development through myoblast fusion³⁰, skeletal development and maintenance of skeletal integrity through formation of multinucleated osteoclasts (OC); and control of certain viral infections and spreading through the formation of macrophage multinucleated giant cells (MGC)³¹. Thus, this important biological process directly defines the course of many pathological processes including infertility, skeletal defects (osteoporosis and osteopetrosis), failure of skeletal repair, failure of maintenance of prosthetic implants as well as fusion of host plasma membrane and virus fusion protein in viral diseases. Clearly, potential common regulators and mechanisms would be attractive therapeutic targets in these disorders.

Two of the cell types that undergo fusion, osteoclasts and multinucleated giant cells, are derived from a common progenitor from bone marrow and periphery. These cells are rendered fusion competent by common molecular mediators and ultimately mediate specialized functions in specific microenvironments. While OC can undergo fusion in both normal or pathological states such as Paget disease³², macrophages undergo fusion to form MGC primarily under inflammatory conditions such as chronic granulomatous disease³³. However, the fact that the fusion of OC and MGCs are governed by common signaling mechanisms again makes these and their component molecules attractive targets for therapeutic intervention. Interestingly, we have previously shown that in response to ischemic injury, CD13^{KO} skeletal muscle satellite cells fused more readily than WT cells to form multinucleated myoblasts, suggesting that CD13 may also participate in fusion of other cell types, thus affecting processes such as skeletal muscle repair¹⁶.

Osteoclastogenesis comprises many steps from the commitment and survival of osteoclast progenitor cells, their differentiation into mononuclear pre-osteoclasts that fuse to generate multinucleated mature osteoclasts and finally activation of osteoclasts for bone resorption. Among the different steps, osteoclast fusion is thought to be the critical step in this phenomenon. Our

data clearly indicate that osteoclast progenitor survival, differentiation and proliferation is not dependent on CD13 expression, suggesting that CD13 may be involved in the fusion mechanism to generate multinucleated osteoclasts.

The OC fusion process itself involves pre-fusion events; apposition of membranes via cell-adhesion molecules and the formation of an unstable intermediate stalk structure³⁴. Fusion entails hemifusion of adjacent monolayers, remodeling of the phospholipid bilayer, mixing of membrane lipids, stalk elongation and formation of a bilayered diaphragm connecting the two cells. The post-fusion disruption of this diaphragm forms the fusion pore which expands in a dynamin-dependent manner producing a fused cell with continuous cytoplasm containing multiple nuclei. Studies have implicated various membrane-associated processes as critical to osteoclastogenesis such as clustering of membrane tetraspanins³⁵ and endocytosis as well as specific molecules, such as DC-STAMP²⁷, OC-STAMP³⁶, dynamin and ATP6V0d2 (ATPase, H⁺ transporting V0 subunit d2)³⁷. Recently, it has been shown that expression and localization membrane lipid species such as phosphatidylethanolamine³⁸ and phosphatidylserine³⁹ are also essential for proper cell-cell fusion. We have shown that CD13 regulates dynamin-and clathrin-mediated endocytosis^{21,22}, recycling of cell surface proteins²⁰ and Src activation⁴⁰, each of which have been shown to participate in OC fusion. Our data showing an aberrant persistence of fusion protein expression post-fusion in CD13^{KO} OCs suggests that CD13 may regulate the expression of these endocytic and fusion proteins during fusion, perhaps by mediating their internalization, endocytic trafficking, recycling and/or localization.

In addition, organizers of actin-based protrusions are pivotal in both osteoclast and macrophage fusion. Recently, we have reported that CD13 is a critical signaling platform that links the plasma membrane to dynamic mediators of actin cytoskeletal assembly and rearrangement²⁰. It remains to be established if CD13 localization is required at the site of cell-cell fusion to enable cells to be fusion competent and that its expression on both cells is imperative for fusion.

In conclusion, in the present study we demonstrate that CD13 expression controls osteoclastogenesis specifically at the level of cell-cell fusion. Considering the diversity and importance of pathologies that are influenced by cell-cell fusion, identification of CD13-dependent molecular mechanisms and signaling that regulate myeloid fusion will provide novel therapeutic approaches in fusion pathologies.

Legends

FIG 1. Loss of CD13 leads to reduced bone volume and increased number of osteoclasts in vivo. μ CT and histomorphometric analysis of trabecular (A) and cortical (B) bone with (C), immunohistochemistry of TRAP⁺ osteoclasts in the femur indicated in purple (arrow), (D), BV/TV; Bone volume/Tissue volume (E), Oc/BS; % OCs per bone surface (F), OcS/BPm; % Osteoclasts per bone perimeter and (G), endosteal bone formation rate relative to bone surface area (BFR/BS) of WT and CD13^{KO} are shown. Data represents +/- SEM of two independent experiments. N=6/genotype. **,p<0.01.

FIG 2. Differentiation of hematopoietic cells to osteoclast progenitors is independent of CD13. Flow cytometric analysis of Osteoclast progenitor population from BM (A) and Spleen (B) indicate similar distribution of progenitor cells between WT and CD13^{KO} mice. Data represents +/- SEM of three independent experiments. N=6/genotype.

FIG 3. Increased TRAP⁺ multinucleated OCs in absence of CD13 in vitro. A). Primary murine bone marrow derived OCs size and number of nuclei/OC grown on bovine cortical bone slices in CD13^{KO} cells compared to WT at d3. **B).** Average # of cell with > 3 nuclei/field, **C).** Average cell area and **D).** average size of nuclei per OCs in CD13^{KO} are significantly larger than the WT cells. **E).** Area of resorption is significantly higher in CD13^{KO} than WT OCs grown on Osteo assay plates (Corning) for d10 by phase contrast imaging & quantified by Image J **(F).** Data represents +/- SEM of three independent experiments. N=6/genotype, **,p<0.01, *,p<0.05.

FIG 4. Increased TRAP⁺ spleen derived multinucleated OCs in absence of CD13 in vitro. A). Primary murine spleen derived OCs size and number of nuclei/OC grown on dentine slices in CD13^{KO} cells in presence of M-CSF and RANKL compared to WT at d3. **B).** Average # of cell with >3 nuclei/field, **C).** cell area and **D)** average size of nuclei per OCs in CD13^{KO} are significantly larger than the WT cells. Data represents +/- SEM of two independent experiments. N=6/genotype, **,p<0.01, *,p<0.05.

FIG 5. Increased Multinucleated Giant Cell formation in CD13^{KO}. A). Giemsa staining indicated increased size and number of MGCs in CD13^{KO} compared to wildtype counterpart in presence of M-CSF and IL-4/IL-13 after d5. **B).** Quantification of MGC with > 3 nuclei/field **C).** Average area and **D).** Number of nuclei/MGC. Data represents +/- SEM of two independent experiments. N=6/genotype, **,p<0.01. **E).** CD13 is expressed in CD68⁺ WT MGC and **F).** co-localize with fusion protein DCST1 after IL-4/IL-13 addition. CD68; green. CD13; red. Magnification 63X oil.

FIG 6. Evaluation of CD13 and fusion-promoting protein expression in M-CSF/RANKL stimulated BM-derived osteoclast by Immunoblot analysis. (A). Expression of the fusion-promoting proteins dynamin and DC-STAMP (DCST1) is aberrantly sustained in CD13^{KO} but not in WT multinucleated osteoclasts in presence of M-CSF and RANKL. **(B).** CD13 expression is

unaltered in BM-derived cells in response to M-CSF and RANKL stimulation over time. Data represents average of two isolates. N=2/genotype.

FIG 7. Immunofluorescence analysis of Dynamin and DC-STAMP (DCST1) in OC. Fusion proteins dynamin and DCST1 expression is maintained in CD13^{KO} multinucleated mature osteoclasts but not in WT cells (**). High level of dynamin colocalizes with actin and DCST1 in OCPs (*). Data represents average of three independent experiments. N=3/genotype. Dynamin, DCST1;green. Phalloidin; red. Magnification 63X oil.

Supplemental Data

FIG S1. Flow sorting of primary murine bone marrow and spleen from WT and CD13^{KO} mice. Live CD11b^{lo} CD115^{hi} Ly6G⁺ (CD3/B220/Nk1.1)⁻ cells were sorted to homogeneity using BD FACS ARIA and analyzed by FACS Diva. Data represents average of three independent experiments.

Materials and Methods

Animals

Global Wildtype and CD13^{KO} (C57BL/6J) male mice were generated and housed at the Gene Targeting and Transgenic Facility at University of Connecticut School of Medicine ¹². All procedures were performed in accordance with the guidelines and regulations approved by the Institutional Animal Care and Use Committee.

Reagents

Recombinant mouse M-CSF, RANKL, IL-4 and IL-13 were purchased from R&D Systems. Acid Phosphatase, Leukocyte (TRAP) Kit was purchased from Sigma-Aldrich. Giemsa/ MAY-GRÜNWARD Stain solution was purchased from Sigma-Aldrich. Toluidine Blue was obtained from EMD Millipore. Bovine cortical bone slices were a kind gift from Dr. Joseph Lorenzo, University of Connecticut School of Medicine⁴¹. Osteoplates were purchased from Corning.

Antibodies

Antibodies to DC-STAMP1 (DCST1; Biorbyt, orb2242, rabbit polyclonal Ab), Dynamin (Abcam, ab3457, rabbit polyclonal Ab), Phalloidin-TRITC (Sigma-Aldrich, P1951), CD13 (SL13, Millipore, MABC950, rat monoclonal Ab)

Flow cytometry

Antibodies for phenotypic analyses and sorting by flow cytometry used are as follows²⁴; anti-CD3 (145-2C11), anti-B220 (RA3-6B2), anti-NK1.1 (PK136), anti-CD11b (M1/70), anti-CD115 (AFS98), anti-Ly6C (Al-21). All antibodies were purchased from Biolegend, BD Biosciences, e-Biosciences. UV Blue Live dead dye was purchased from Life Technologies. Labeling of cells for flow cytometry and sorting was performed as described. Briefly, flow cytometry on live cells were performed using BD-FACS Aria (BD Biosciences) and data analyzed with FlowJo software.

In vivo analysis of WT and CD13^{KO} mice

Micro-Computed Tomography and Histomorphometry^{41,42}

Cortical and trabecular bone from 8-10 wks. old WT and CD13^{KO} mice male were isolated and scanned using micro-computed tomography (μ CT) system and 3-D analysis and reconstruction were performed as described to measure the trabecular bone volume (BV/TV, %). For static histomorphometry, trabecular volume was measured as described. Osteoclasts were identified by multi-nucleated TRAP+ cells adjacent to the bone surface.

Alkaline Phosphatase assay⁴²

Osteoblast culture generated from bone marrow were stained for the alkaline phosphatase activity as described.

Isolation of hematopoietic progenitor population from bone marrow and spleen

Bone marrow cells were obtained by flushing femur and tibia from WT or CD13^{KO} mice with 10ml 1xPBS and 2% heat inactivated FBS, followed by RBC lysis and filtering through 40um cell strainer (BD Biosciences). Total live cells counted with Countess Automated cell counter (Thermo Fisher Scientific) were stained with antibody cocktail at 4 degree C. Cells from mouse spleen was obtained by gentle crushing the organ between frosted microscopic slides in cold 10ml 1x PBS and 2% heat inactivated FBS.

Generation and Culture of Osteoclast progenitors from bone marrow or spleen

Cells from mouse BM or spleen were stained with Ab cocktail containing anti-(CD3, B220, NK1.1, CD115, Ly6C) Ab and subjected to single-cell sorting by BD FACS Aria²⁴. Sorted progenitors at a density of 10,000-50,000 cells/well were seeded in 96-well dish in α -MEM containing 10%FBS, 1% Penicillin-Streptomycin, 30ng/ml M-CSF and 30ng/ml RANKL at 37 degree C with 5% CO₂ for 0-10d. Multi-nucleated osteoclasts were stained with Tartrate-Resistant Acid Phosphatase staining and assessed by counting cells with more than three nuclei. Average area of osteoclast was measured by ImageJ software.

TRAP staining⁴²

Osteoclast progenitor cells in α -MEM (GIBCO BRL) containing 10%FBS, 1% Penicillin-Streptomycin, 30ng/ml M-CSF and 30ng/ml RANKL grown on plastic or UV-sterilized, devitalized bovine cortical bone slices (placed in 96-well dishes), at a density of 50,000 cells/well for indicated time were fixed in 2.5% Glutaraldehyde and Tartrate-Resistant Acid Phosphatase in osteoclasts were stained using TRAP staining kit according to manufacturer's instruction (Sigma-Aldrich).

Bone resorption assay⁴²

Osteoclast progenitors were seeded on Osteo Assay plate (Corning) at a density of 50,000 cells/well in α -MEM containing 10%FBS, 1% Penicillin-Streptomycin, 30ng/ml M-CSF and 30ng/ml RANKL for d10. Surface pit formation was measured by removing cells with 100ul of 10% bleach solution at RT for 5 min. Wells were washed with deionized water and allowed to dry. Cluster of pits formed was imaged using a light microscope (Olympus Scientific) and the area of resorption was measured by ImageJ software.

Generation of Multinucleated Giant Cells (MGC) and Giemsa/ MAY-GRÜNWARD Staining

BM sorted Osteoclast progenitors were seeded on 96-well dishes at a density of 50,000 cells/well in α -MEM containing 10%FBS, 30ng/ml M-CSF for d5 followed by addition of and 30ng/ml IL-4 and 30ng/ml IL-13 for an additional d3. Cells were washed and fixed with 100% methanol followed by staining with Giemsa stain according to manufacturer's instruction (Sigma-Aldrich).

Immunofluorescence and microscopy

Osteoclast progenitors or MGC grown on glass coverslips that were previously coated with 5 μ g/ml fibronectin for indicated time period. Cells were fixed in 4% paraformaldehyde (Electron Microscopy Sciences) at RT for 30 min, permeabilized with 0.1% Triton-X-100 in PBS at RT for 5 min. Cells were blocked with blocking buffer containing 5% goat or donkey serum/5% BSA/1xPBS at RT for 1h followed by incubation with primary Ab in blocking buffer at 4 degree C for overnight. Cells were washed and treated with secondary Ab (1:1200) and DAPI (nuclear stain) in blocking buffer at RT for 1h. Coverslips were mounted with ProLong Gold antifade mounting medium (Life Technologies), visualized at excitation wavelength of 488nm (Alexa 488), 543nm (Alexa 594 or TRITC) and 405nm (DAPI) and imaged by Zeiss LSM 880 confocal fluorescence microscope.

Immunoblot analysis

Cell lysates from flow-sorted bone marrow osteoclast progenitors grown in presence of M-CSF and RANKL over time were harvested in 1% NP40 lysis buffer containing 1X cOmplete Protease Inhibitor cocktail (Roche). Samples were separated by SDS-PAGE and transferred to nitrocellulose membrane, blocked in 1XTBST containing 5% bovine serum albumin, treated with primary Ab followed by appropriate secondary Ab and imaged by ChemiDoc Imaging system (Biorad). β actin was used as loading control.

Statistical analysis

Statistical analysis was performed using unpaired, two-tailed Student's t test using GraphPad Prism software and results are representative of mean $\pm$ SD. Differences at $p \leq 0.05$ were considered significant.

Disclosures: All animal experiments in this study were reviewed and approved by Animal Care Committee at University of Connecticut Medical School. All the authors do not have any conflict of interest.

Author contributions:

MG, IK, HLA and LHS designed the research plan. MG, IK and HLA conducted experiments and analyzed the data. MG, HLA and LHS wrote the manuscript.

Acknowledgments

We thank Dr. Joseph Lorenzo for the cortical bone slices and for the use of his light microscope. We also thank Susan Staurovsky and Evan Jellison from CCAM and Flow cytometry core facilities respectively at UConn Health, for providing technical assistance. This work was supported by National Institutes of Health grants R01HL127449 and R01HL125186 (to LHS and MG)

- 1 Teitelbaum, S. L. Bone resorption by osteoclasts. *Science* **289**, 1504-1508, doi:10.1126/science.289.5484.1504 (2000).
- 2 Tsukasaki, M. & Takayanagi, H. Osteoimmunology: evolving concepts in bone-immune interactions in health and disease. *Nat Rev Immunol* **19**, 626-642, doi:10.1038/s41577-019-0178-8 (2019).
- 3 Madel, M. B. *et al.* Immune Function and Diversity of Osteoclasts in Normal and Pathological Conditions. *Front Immunol* **10**, 1408, doi:10.3389/fimmu.2019.01408 (2019).
- 4 Boudin, E., Fijalkowski, I., Hendrickx, G. & Van Hul, W. Genetic control of bone mass. *Mol Cell Endocrinol* **432**, 3-13, doi:10.1016/j.mce.2015.12.021 (2016).
- 5 Krakow, D. Skeletal dysplasias. *Clin Perinatol* **42**, 301-319, viii, doi:10.1016/j.clp.2015.03.003 (2015).
- 6 Arai, F. *et al.* Commitment and differentiation of osteoclast precursor cells by the sequential expression of c-Fms and receptor activator of nuclear factor kappaB (RANK) receptors. *J Exp Med* **190**, 1741-1754, doi:10.1084/jem.190.12.1741 (1999).
- 7 Fogg, D. K. *et al.* A clonogenic bone marrow progenitor specific for macrophages and dendritic cells. *Science* **311**, 83-87, doi:10.1126/science.1117729 (2006).
- 8 Jacquin, C., Gran, D. E., Lee, S. K., Lorenzo, J. A. & Aguila, H. L. Identification of multiple osteoclast precursor populations in murine bone marrow. *J Bone Miner Res* **21**, 67-77, doi:10.1359/JBMR.051007 (2006).
- 9 Mensah, K. A., Ritchlin, C. T. & Schwarz, E. M. RANKL induces heterogeneous DC-STAMP(lo) and DC-STAMP(hi) osteoclast precursors of which the DC-STAMP(lo) precursors are the master fusogens. *J Cell Physiol* **223**, 76-83, doi:10.1002/jcp.22012 (2010).
- 10 Miyamoto, H. *et al.* Osteoclast stimulatory transmembrane protein and dendritic cell-specific transmembrane protein cooperatively modulate cell-cell fusion to form osteoclasts and foreign body giant cells. *J Bone Miner Res* **27**, 1289-1297, doi:10.1002/jbmr.1575 (2012).
- 11 Yagi, M. *et al.* DC-STAMP is essential for cell-cell fusion in osteoclasts and foreign body giant cells. *J Exp Med* **202**, 345-351, doi:10.1084/jem.20050645 (2005).
- 12 Winnicka, B. *et al.* CD13 is dispensable for normal hematopoiesis and myeloid cell functions in the mouse. *J Leukoc Biol* **88**, 347-359, doi:10.1189/jlb.0210065 (2010).

- 13 Look, A. T., Ashmun, R. A., Shapiro, L. H. & Peiper, S. C. Human myeloid plasma membrane glycoprotein CD13 (gp150) is identical to aminopeptidase N. *J Clin Invest* **83**, 1299-1307, doi:10.1172/JCI114015 (1989).
- 14 Petrovic, N. *et al.* CD13/APN regulates endothelial invasion and filopodia formation. *Blood* **110**, 142-150, doi:10.1182/blood-2006-02-002931 (2007).
- 15 Mina-Osorio, P. *et al.* CD13 is a novel mediator of monocytic/endothelial cell adhesion. *J Leukoc Biol* **84**, 448-459, doi:10.1189/jlb.1107802 (2008).
- 16 Rahman, M. M. *et al.* CD13 promotes mesenchymal stem cell-mediated regeneration of ischemic muscle. *Frontiers in physiology* **4**, 402, doi:10.3389/fphys.2013.00402 (2014).
- 17 Nomura, R. *et al.* Human coronavirus 229E binds to CD13 in rafts and enters the cell through caveolae. *J Virol* **78**, 8701-8708, doi:10.1128/JVI.78.16.8701-8708.2004 (2004).
- 18 Saiki, I. *et al.* Role of aminopeptidase N (CD13) in tumor-cell invasion and extracellular matrix degradation. *Int J Cancer* **54**, 137-143, doi:10.1002/ijc.2910540122 (1993).
- 19 Bhagwat, S. V. *et al.* CD13/APN is activated by angiogenic signals and is essential for capillary tube formation. *Blood* **97**, 652-659, doi:10.1182/blood.v97.3.652 (2001).
- 20 Ghosh, M. *et al.* CD13 tethers the IQGAP1-ARF6-EFA6 complex to the plasma membrane to promote ARF6 activation, beta1 integrin recycling, and cell migration. *Sci Signal* **12**, doi:10.1126/scisignal.aav5938 (2019).
- 21 Ghosh, M., Subramani, J., Rahman, M. M. & Shapiro, L. H. CD13 restricts TLR4 endocytic signal transduction in inflammation. *J Immunol* **194**, 4466-4476, doi:10.4049/jimmunol.1403133 (2015).
- 22 Ghosh, M., McAuliffe, B., Subramani, J., Basu, S. & Shapiro, L. H. CD13 regulates dendritic cell cross-presentation and T cell responses by inhibiting receptor-mediated antigen uptake. *J Immunol* **188**, 5489-5499, doi:10.4049/jimmunol.1103490 (2012).
- 23 Shin, N. Y. *et al.* Dynamin and endocytosis are required for the fusion of osteoclasts and myoblasts. *J Cell Biol* **207**, 73-89, doi:10.1083/jcb.201401137 (2014).
- 24 Jacome-Galarza, C. E., Lee, S. K., Lorenzo, J. A. & Aguila, H. L. Identification, characterization, and isolation of a common progenitor for osteoclasts, macrophages, and dendritic cells from murine bone marrow and periphery. *J Bone Miner Res* **28**, 1203-1213, doi:10.1002/jbmr.1822 (2013).
- 25 Helming, L. & Gordon, S. The molecular basis of macrophage fusion. *Immunobiology* **212**, 785-793, doi:10.1016/j.imbio.2007.09.012 (2007).
- 26 Helming, L. & Gordon, S. Macrophage fusion induced by IL-4 alternative activation is a multistage process involving multiple target molecules. *Eur J Immunol* **37**, 33-42, doi:10.1002/eji.200636788 (2007).
- 27 Yagi, M., Miyamoto, T., Toyama, Y. & Suda, T. Role of DC-STAMP in cellular fusion of osteoclasts and macrophage giant cells. *J Bone Miner Metab* **24**, 355-358, doi:10.1007/s00774-006-0697-9 (2006).
- 28 Pereira, M. *et al.* Common signalling pathways in macrophage and osteoclast multinucleation. *J Cell Sci* **131**, doi:10.1242/jcs.216267 (2018).
- 29 Ogle, B. M., Cascalho, M. & Platt, J. L. Biological implications of cell fusion. *Nat Rev Mol Cell Biol* **6**, 567-575, doi:10.1038/nrm1678 (2005).

- 30 Pajcini, K. V., Pomerantz, J. H., Alkan, O., Doyonnas, R. & Blau, H. M. Myoblasts and macrophages share molecular components that contribute to cell-cell fusion. *J Cell Biol* **180**, 1005-1019, doi:10.1083/jcb.200707191 (2008).
- 31 Bracq, L. *et al.* T Cell-Macrophage Fusion Triggers Multinucleated Giant Cell Formation for HIV-1 Spreading. *J Virol* **91**, doi:10.1128/JVI.01237-17 (2017).
- 32 Roodman, G. D. & Windle, J. J. Paget disease of bone. *J Clin Invest* **115**, 200-208, doi:10.1172/JCI24281 (2005).
- 33 Seitzer, U., Haas, H. & Gerdes, J. A human in vitro granuloma model for the investigation of multinucleated giant cell and granuloma formation. *Histol Histopathol* **16**, 645-653, doi:10.14670/HH-16.645 (2001).
- 34 Brukman, N. G., Uygur, B., Podbilewicz, B. & Chernomordik, L. V. How cells fuse. *J Cell Biol* **218**, 1436-1451, doi:10.1083/jcb.201901017 (2019).
- 35 Takeda, Y. *et al.* Tetraspanins CD9 and CD81 function to prevent the fusion of mononuclear phagocytes. *J Cell Biol* **161**, 945-956, doi:10.1083/jcb.200212031 (2003).
- 36 Khan, U. A., Hashimi, S. M., Bakr, M. M., Forwood, M. R. & Morrison, N. A. Foreign body giant cells and osteoclasts are TRAP positive, have podosome-belts and both require OC-STAMP for cell fusion. *J Cell Biochem* **114**, 1772-1778, doi:10.1002/jcb.24518 (2013).
- 37 Kim, K., Lee, S. H., Ha Kim, J., Choi, Y. & Kim, N. NFATc1 induces osteoclast fusion via up-regulation of Atp6v0d2 and the dendritic cell-specific transmembrane protein (DC-STAMP). *Mol Endocrinol* **22**, 176-185, doi:10.1210/me.2007-0237 (2008).
- 38 Irie, A., Yamamoto, K., Miki, Y. & Murakami, M. Phosphatidylethanolamine dynamics are required for osteoclast fusion. *Sci Rep* **7**, 46715, doi:10.1038/srep46715 (2017).
- 39 Verma, S. K. *et al.* Cell-surface phosphatidylserine regulates osteoclast precursor fusion. *J Biol Chem* **293**, 254-270, doi:10.1074/jbc.M117.809681 (2018).
- 40 Subramani, J. *et al.* Tyrosine Phosphorylation of CD13 Regulates Inflammatory Cell-Cell Adhesion and Monocyte Trafficking. *J Immunol* **191**, 3905-3912, doi:10.4049/jimmunol.1301348 (2013).
- 41 Grcevic, D. *et al.* The Long Pentraxin 3 Plays a Role in Bone Turnover and Repair. *Front Immunol* **9**, 417, doi:10.3389/fimmu.2018.00417 (2018).
- 42 Novak, S. *et al.* Osteoclasts Derive Predominantly from Bone Marrow-Resident CX3CR1(+) Precursor Cells in Homeostasis, whereas Circulating CX3CR1(+) Cells Contribute to Osteoclast Development during Fracture Repair. *J Immunol* **204**, 868-878, doi:10.4049/jimmunol.1900665 (2020).

FIG 1. Loss of CD13 leads to reduced bone volume and increased number of osteoclasts in vivo. μ CT and histomorphometric analysis of trabecular (A) and cortical (B) bone with (C), immunohistochemistry of TRAP⁺ osteoclasts (arrow) in the femur indicated in purple, (D), BV/TV; Bone volume/Tissue volume (E), Oc/BS; % OCs per bone surface (F), OcS/BPm; % Osteoclast per bone perimeter and (G), endosteal bone formation rate relative to bone surface area (BFR/BS) of WT and CD13^{KO} are shown. Data represents $\pm$ SEM of two independent experiments. N=6/genotype. **, $p < 0.01$.

FIG 2. Differentiation of hematopoietic cells to osteoclast progenitors is independent of CD13. Flow cytometric analysis of Osteoclast progenitor populations from BM (A) and Spleen (B) indicate a similar distribution of progenitor cells between WT and CD13^{KO} mice. Data represents +/- SEM of three independent experiments. N=6/genotype.

FIG 3. TRAP⁺ multinucleated OCs are increased in the absence of CD13 in vitro. **A).** Primary murine bone marrow derived OCs size and number of nuclei/OC grown on bovine cortical bone slices in CD13^{KO} cells compared to WT at d3. **B).** Average # of cells with >3 nuclei/field, **C).** Average cell area and **D).** average size of nuclei per OCs in CD13^{KO} are significantly larger than the WT cells. **E).** Area of resorption is significantly higher in CD13^{KO} than WT OCs grown on osteoplates for d10 by phase contrast imaging & quantified by Image J (**F**). Data represents +/- SEM of three independent experiments. N=6/genotype, **,p<0.01, *,p<0.05.

FIG 4. Increased TRAP⁺ splenic multinucleated OCs in the absence of CD13 in vitro. A). Primary murine spleen-derived OCs size and number of nuclei/OC grown on cortical bone slices in CD13^{KO} cells in the presence of M-CSF and RANKL compared to WT at d3. **B).** Average # of cells with >3 nuclei/field, **C).** cell area and **D)** average size of nuclei per OC in CD13^{KO} are significantly larger than the WT cells. Data represents +/- SEM of two independent experiments. N=6/genotype, **, p<0.01.

FIG 5. Increased Multinucleated Giant Cell formation in CD13^{KO} cells. **A).** Giemsa staining indicated increased size and number of MGCs in CD13^{KO} compared to WT in the presence of M-CSF and IL-4/IL-13 after 5 days. **B).** Quantification of MGC with >3 nuclei/field **C).** Average area and **D).** Number of nuclei/MGC. Data represents +/- SEM of two independent experiments. N=6/genotype, **,p<0.01. **E).** CD13 is expressed in CD68⁺ WT MGC and **F).** co-localize with fusion protein DCST1 after IL-4/IL-13 addition. CD68; green. CD13; red. Magnification 63X oil.

FIG 6. Evaluation of CD13 and fusion-promoting protein expression in M-CSF/RANKL stimulated BM-derived osteoclast by immunoblot analysis. (A). Expression of the fusion-promoting proteins dynamin and DC-STAMP (DCST1) is aberrantly sustained in CD13^{KO} but not in WT multinucleated osteoclasts in presence of M-CSF and RANKL. **(B).** CD13 expression is unaltered in BM-derived cells in response to M-CSF and RANKL stimulation over time. Data represents average of two isolates. N=2/genotype.

FIG 7. Immunofluorescence analysis of DC-STAMP (DCST1) in OC. Expression of the fusion-promoting proteins dynamin and DCST1 is maintained in CD13^{KO} multinucleated mature osteoclasts but not in WT cells (**). High levels of dynamin co-localize with actin and DCST1 in OCPs (*). Data represents average of three independent experiments. N=3/genotype. Dynamin, DCST1; green. Phalloidin; red. Magnification 63X oil.