bioRxiv

Trial-by-trial variability in cortical responses exhibits scaling in spatial correlations predicted from critical dynamics

Tiago L. Ribeiro¹, Shan Yu^{1,2,3}, Daniel A. Martin^{4,5}, Daniel Winkowski⁶, Patrick Kanold⁶, Dante R. Chialvo^{4,5}, &
Dietmar Plenz¹

¹Section on Critical Brain Dynamics, National Institute of Mental Health, National Institutes of Health, Bethesda,
MD, 20892, USA

²Brainnetome Center, Institute of Automation, and ³CAS Center for Excellence in Brain Science and Intelligence
Technology, Chinese Academy of Sciences, Beijing 100190, China

⁴Center for Complex Systems & Brain Sciences (CEMSC3), Instituto de Ciencias Físicas, (ICIFI) Escuela de Ciencia y
Tecnología, Universidad Nacional de San Martín (UNSAM), San Martín, (1650) Buenos Aires, Argentina

⁵Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Godoy Cruz 2290, Buenos Aires, Argentina

⁶Department of Biology, University of Maryland, College Park, MD 20742, USA

Pages/Figures: 33 pages, 5 figures

Suppl. Material: 5 pages, 3 figures

Running Title: Scale-free correlations in neuronal activity

Correspondence: Dietmar Plenz, Ph.D., Section on Critical Brain Dynamics, National Institute of
Mental Health, Porter Neuroscience Research Center, Rm 3A-1000, 35 Convent
Drive, Bethesda, MD 20892. office: (301) 402-2249, fax: (301) 480-7480,
plenzd@mail.nih.gov

Keywords: mouse, primary cortex, cortex, 2-photon imaging, nonhuman primate, prefrontal and
premotor cortex, high-density microelectrode array, local field potential, correlation
functions, neuronal avalanches, network simulations, criticality

Abstract

Simple sensory stimuli or motor outputs engage large populations of neurons in the mammalian cortex. When stimuli or outputs repeat, the robust population response contrasts with fluctuating responses of individual neurons, known as trial-by-trial variability. To understand this apparent discrepancy, a detailed identification of the underlying spatiotemporal correlations is required. Here, we analyze spatial correlations in the instantaneous fluctuations between neurons relative to the neuronal population. Using 2-photon imaging of visual and auditory responses in primary cortices of awake mice, we show that these correlations grow linearly with the size of the observed cortical area. We extend these observations to the cortical mesoscale by analyzing local field potentials in behaving nonhuman primates. In network simulations, we demonstrate this linear growth in spatial correlation to emerge at criticality. Our findings suggest that trial-by-trial variability is a signature of critical dynamics in cortex maintaining robust, long-range spatial correlations among neurons.

13

Introduction

Even simple stimuli or movements engage large numbers of neurons in the mammalian cortex. These robust population responses are contrasted by the heterogeneity and fluctuations of single neuron responses observed during repeated stimuli or motor outputs. This response variability or trial-by-trial variability has been consistently found *in vivo* for neurons that are non-selective to a particular stimulus feature (Dewese & Zador, 2004) as well as highly selective neurons (Heggelund & Albus, 1978; Tolhurst *et al.*, 1983; Vogels *et al.*, 1989; Shadlen & Newsome, 1998) and *in vitro* under network and stimulus conditions of reduced complexity (e.g. (Haroush & Marom, 2019)). While part of the variability has been attributed to single neuron properties, e.g. through a non-linear transfer between the membrane potential

and firing rate (Carandini, 2004; Charles *et al.*, 2018), numerous findings point to a significant role of the network dynamics to the observed variability. In particular, response variability is shared among neurons (Cohen & Kohn, 2011) through shared gain (Goris *et al.*, 2014), includes the interactions between selective as well as non-selective neurons (Kotekal & MacLean, 2019), is supported by local cortical motifs such as fan-in (Dechery & MacLean, 2018) and depends on the balance of excitation and inhibition (Haroush & Marom, 2019). This potential role in network dynamics is in line with early findings that response variability strongly correlates with ongoing synaptic activity preceding a stimulus (Arieli *et al.*, 1996).

The diversity and variability encountered in single neuron responses to even simple, repeated stimuli raises the question whether there exist certain principles that underlie this variability in the face of coherent network responses. These principles in order to be general should be independent from the specific stimulus given or motor output achieved, which translates into the general question of how individual elements, by interacting locally, can achieve a coherent, global population response. Correlation length measures can be used to identify the relationship between global, coherent system responses and fluctuating system components (Wilson, 1979). By studying the correlation in the fluctuations of system components, one can determine the distance at which they behave independently. It is well-established that the correlation length diverges at criticality (Wilson, 1976) in the thermodynamic limit. For finite systems, however, this asymptote behavior can be demonstrated by showing that the correlation length grows with system size. This relationship between correlation length and system size has been identified in the context of bird flocks, where a flock still maintains a coherent trajectory in space despite fluctuations in single bird trajectories (Cavagna *et al.*, 2010; Bialek *et al.*, 2012; Bialek *et al.*, 2014). It was also demonstrated for resting activity of the human brain that the correlation length of the blood oxygenated level dependent (BOLD) signal scales linearly with the size of the brain region measured (Fraiman & Chialvo, 2012), a finding in line with expectations that the brain at rest maintains a critical state.

It has been argued (Chialvo, 2010) and shown experimentally (Shew & Plenz, 2013) that neuronal networks confer multiple benefits for being in the critical state, such as high dynamic range and maximal information capacity. For the bird flock, the critical state has been suggested to confer fast and coherent changes in flight direction in response to external perturbations, such as a predator attack (Bialek *et al.*, 2014). For the brain, one might expect that external perturbations from sensory input or internal perturbations such as self-initiated motor commands lead to population responses that maintain the critical state. If this was correct, the correlation length for sensory and motor responses in the brain should scale with the size of the observed population of neurons and should be independent of feature selectivity. Here, we show experimentally a linear scaling in correlation length in the fluctuations of neuronal evoked responses in cortex. Using numerical simulations of a network model, we demonstrate this scaling to be found when the model is tuned to be critical. Our results identify a specific form of scaling in the trial-by-trial variability for sensory and motor responses in cortex *in vivo* and suggest this variability is a signature of critical dynamics that maintains robust, long-range correlations between neurons. Finally, by comparing experimental and numerical correlation functions, we are able to estimate the interaction length (i.e. the distance at which neurons interact physically) in visual cortex.

Results

We investigated the extent in the spatial correlation of neuronal activity at the cellular and mesoscale level in cortex using two different experimental approaches. At cellular resolution, we studied pair-wise correlations between individual neurons using 2-photon imaging (2PI), whereas at the mesoscale in nonhuman primates we measured the local field potential (LFP) using high-density microelectrode arrays. We analyzed ongoing as well as activity obtained during sensory/motor processing for each scale. We will first present results obtained at the cellular level in awake mice followed by a comparison at the mesoscale

in behaving nonhuman primates. Simulations of neuronal activity position our results in the context of subcritical, critical and supercritical dynamical regimes.

72 **Linear scaling of correlation lengths in mouse primary cortex**

73 For cellular analysis, spiking activity in pyramidal cells of mouse primary visual cortex (V1; $n = 7$ mice)
 74 was recorded using 2PI over an area of $L \times L = \sim 400 \mu\text{m} \times 400 \mu\text{m}$ in superficial cortical layers at a depth
 75 of $\sim 100 - 150 \mu\text{m}$ from the cortical surface. Mice were quietly resting during recording while 2 s-lasting
 76 drifting gratings were presented every 4 s (Fig. 1a). As can be seen in the variance over average response
 77 plot (Fig. 1b), the Fano factor was above one for most cells, indicating high response variability in single
 78 neuron responses. About 20% of neurons were tuned to stimulus direction with a selectivity index (DSI;
 79 Fig. 1c; see Materials and Methods) larger than 0.4 and an average response profile (Fig. 1d) depicting
 80 minimal response to the orthogonal orientation. Importantly, all single neuron responses were embedded
 81 in spatio-temporal clusters of activity that distributed in cluster size according to power laws, the hallmark
 82 of neuronal avalanches (Beggs & Plenz, 2003; Bellay *et al.*, 2015). This scale-invariant organization in
 83 activity clusters was found during stimulation regardless of the directional drift as well as during
 84 luminance-matched gray screen presentation (Fig. 1e; $p < 0.05$ for power law test; see Materials and
 85 Methods).

86 The response variability encountered in single neurons, their tuning diversity and scale-invariant
 87 grouping in the form of avalanches points to a fluctuation-dominated dynamic regime of cortex, in which
 88 individual elements seem to be difficult to relate to stable population responses. One potential answer to
 89 this problem could lie in critical dynamics, which maintain some order in the presence of strong
 90 fluctuations (Chialvo 2010). Therefore, to gain deeper insight into the dynamical regime of the cortex
 91 during the relatively short-lasting (compared to the scale of large avalanches) evoked responses and, at

Fig. 2: Correlations in activity fluctuations between V1 neurons grow with the size of the cortical area observed during visual presentations. **a** Schematic to measure pair-wise correlation between two V1 neurons (*open circles*) separated by distance r for different fields of view (*colored squares*), superimposed on a typical $400 \times 400 \mu\text{m}$ 2-photon image showing YC2.6 expressing pyramidal neurons in V1 ($\sim 120 \mu\text{m}$ cortical depth). **b** Average spatial correlation in $\Delta R/R$ for V1 neurons decays according to a power law with exponential decay during visual presentations in line with prediction for critical dynamics. *Squares*: data (mean $\pm$ standard deviation). *Line*: power law with exponential decay fit. *Inset*: Trial-shuffling (*gray*) results in near-zero correlations independent of distance. For comparison, original correlation (*black*) added in log-linear coordinates. *Bars* indicate standard deviation. **c** Example comparison of fluctuations in neuronal responses relative to the population response as a function of window extent. *Left*: Individual, single $\Delta R/R \text{ Ca}^{2+}$ traces for two neurons (see panel **a**). Corresponding population activity fluctuates more in small windows (*blue*) compared to larger windows (*orange*; see panel **a**). *Right*: Subtracting the instantaneous population activity from individual neuronal responses reveals the corresponding fluctuations around the mean for each neuron as a function of window extent. Correlation value for each window extent is shown. **d** Correlation in the fluctuations around the mean decays with distance and crosses zero. The zero crossing defines the correlation length ξ , which is seen to increase with window length (*arrows*). Averages for $n = 7$ animals and all gratings directions (*left*) and gray screen (*right*). *Colors*: window length. *Broken line*: zero correlation. **e** Correlations are destroyed by trial-shuffling in time (*short dash*) or shuffling of ROI position (*long dash*), as revealed from the near-zero values obtained, independent of distance ($L = 343 \mu\text{m}$ case shown).

92

93

the same time, avoid the statistical influence of the strong drive created by the stimulus, we needed an

alternative approach to the statistical distribution of observed activity patterns. Here we assessed the dynamics of the cortical network by analyzing how correlations in the evoked responses scale with the size of the observed system. We first directly calculated how the average pairwise correlation between neurons changes as a function of cortical distance r for the full recording window. This correlation was well described by a power law with exponent close to -1 and exponential cut-off, in line with expectations for critical dynamics (Cavagna *et al.*, 2010) measured within a finite size window (Fig. 2b; $p < 0.05$ for power law test). Importantly, when activity from each of the trials for a given stimulus angle was randomly shuffled for each cell independently (trial shuffling; see Materials and Methods), the correlations dropped to zero regardless of the distance between the cells. This indicates that the total activity induced by the stimulus does not contribute to the distance related decay in correlations. We then assessed how the fluctuations of neuronal responses around the instantaneous population mean change with distance r between cells, by subtracting the population average as a function of time from each neuron's time series (Fig. 2c). The correlation in fluctuations as a function of distance r , $C(r)$, was assessed for varying length L of the observation window (Fig. 2d). This variation in L systematically changes the population mean as exemplified in Fig. 2c for two window lengths. Indeed, nearby neurons deviate in similar fashion from population activity and $C(r)$ decayed similarly, both for drifting gratings as well as during gray screen. At a certain distance, defined as the correlation length, ξ (Fig. 2d, arrows) (Cavagna *et al.*, 2010; Fraiman & Chialvo, 2012), $C(r)$ turns negative. We found that $C(r)$ decayed similarly with distance for different $L < 400 \mu\text{m}$ (Fig. 2d, different colors) and ξ was shorter for smaller windows.

Deviations from the population average of cells further apart than ξ were found, on average, to be anti-correlated for any window size. This anti-correlation is to be expected to a certain degree as by definition, at any given time some cells will exhibit activity above the population mean, while others will be below it. This fact and the general drop in correlations with distance will lead to a crossing through zero

correlation, allowing to estimate ξ . We used trial shuffling as a control to demonstrate that ξ indeed captures emergent correlations in trial-by-trial fluctuations among neurons and does not reflect an artificial position due to subtracting the instantaneous mean. Trial shuffling maintains the common input introduced by the stimulus and distances among neurons, while removing coordinated changes of individual neuronal response around the population response. Trial shuffling results in $C(r)$ being close to zero regardless of distance between neuron pairs, and therefore ξ cannot be estimated. This is true for both drifting gratings stimulus and gray-screen periods (Fig. 2e). Similarly, ξ is not defined after randomly permuting cell positions while keeping instant trial-by-trial correlations intact. These controls carried out for the full window size confirm ξ as a correlation length measure for neuronal activity that emerges from interactions between the observed neurons.

We note that a common way to obtain correlation length estimates from a system is precisely with the fit performed in Fig. 2b, by measuring how quickly correlations (without the population subtraction) drop with distance between system components after the power law regime. More specifically, the characteristic decay of the exponential portion of such a correlation spatial function can be used as a measure for the correlation length. However, as can be seen in Fig. 2b, the correlation function becomes very noisy for larger distances, reflecting the small number of pairs of neurons that far apart. Consequently, our estimates for the correlation length using that method become very unreliable ($\xi \sim 110.97 \pm 58.55$; 95% confidence interval). With the alternative method (Cavagna *et al.*, 2010), which has previously been employed in neuronal activity data (Fraiman & Chialvo, 2012), we achieved a full collapse for all L by rescaling $C(r)$ with its corresponding value ξ (Fig. 3a; $p < 0.05$; see Materials and Methods) and obtained a linear relationship for ξ over L (Fig. 3b), i.e. spatial correlations in V1 grow linearly with the observed cortical area. This increase holds regardless of sensory stimulus presented (Fig. 3b, different colors; $p < 0.05$ chi-square test for both cases; see Materials and Methods) and for subsets of cells responsive to the stimulus

as well as those that are not (Fig. 3b, inset; $p < 0.05$ chi-square test for both cases), indicating that all cells, independent of their tuning-selectivity, follow the same correlation principles. Linear scaling with window size is also obtained when spike density estimates are being used (Suppl. Fig. S1). Our findings suggest that fluctuations around the mean between V1 neurons are scale-invariant, a hallmark of criticality.

Fig. 3: Linear growth in correlation length between neurons in primary visual and auditory cortex during sensory stimulation. **a** Scale-invariant spatial correlation function in V1 during sensory stimulation obtained by collapsing correlation functions from different window extents (cf. Fig. 2d; average from $n = 7$ mice). Rescaled correlations $C \times \xi^\gamma$ as function of distance normalized by correlation length ξ . *Black solid line*: average. *Left/right*: Drifting gratings/gray screen. **b** A linear growth in correlation length ξ with window extent L is found for drifting gratings (red) as well as gray screen (gray) presentations ($n = 7$ mice). *Inset*: tuned (orange) as well as untuned (green) neurons exhibit similar linear growth in spatial correlations. *Lines*: linear regression. **c** Neurons in primary auditory cortex of passively listening mice exhibit scale-invariant spatial correlation functions for both tone ON and OFF conditions (left and right, respectively). **d** Corresponding linear growth in correlation length ξ as function of L , for tone ON (red) and OFF (blue) ($n = 11$ mice). *Lines*: linear regression.

We extended these findings from V1 to the primary auditory cortex in awake mice. Mice passively listened to short (1 s) auditory tones semi-randomly presented from 8 different frequencies and 3 different volume levels every 3 – 5 s (Bowen et al., 2019). Evoked responses were recorded using 2PI in pyramidal neurons

from superficial layers ($n = 11$ mice). Evoked neural responses were analyzed as described for visual cortex. Correlation functions $C(r)$ were collapsed based on correlation length ξ and recording window size (Fig. 3c). In line with what was found for V1, ξ grew linearly with L for both tone ON and tone OFF conditions in auditory cortex (Fig. 3d; $p < 0.05$ chi-square test for both cases).

Linear scaling of correlation lengths in nonhuman primates at cortical mesoscale

For the mesoscale level, i.e. analysis beyond several hundred μm , we employed microelectrode arrays (MEAs; 10×10 electrodes without corners; $400 \mu\text{m}$ interelectrode distance) and recorded the local field potential (LFP) in prefrontal (PF; monkey A) and premotor (PM; monkey B) cortex over an area of $\sim 4 \text{ mm} \times 4 \text{ mm}$ (Fig. 4a; for details see (Yu *et al.*, 2017)). Monkey A was trained in a visual-motor mapping task (Fig. 4b, bottom; see Materials and Methods), while monkey B performed a self-initiated movement task (Fig. 4b; top). We applied the methods described in the previous section to investigate whether the same results could be observed in this larger scale. Amplitude fluctuations in the LFP were obtained by subtracting the instantaneous average on the squared subarrays of width L from each electrode LFP. We then obtained $C(r)$ between pairs of electrodes for the subarray of width L in multiples of the inter electrode distance ($400 \mu\text{m}$). $C(r)$ decayed with distance (Fig. 4c) in remarkably similar shapes for different L , allowing for successful collapse of the curves (Fig. 4d). The collapsed functions were similar among both monkeys (Fig. 4d, left vs. right), during baseline or task-evoked activity (Fig. 4d, top vs. bottom). As found for primary visual and auditory cortex, ξ grew linearly with L for both cortical regions and was similar between baseline and motor/sensory processing epochs (Fig. 4e; $p < 0.05$ chi-square test for all cases). Importantly, this is not the case when trial shuffling is employed, as that procedure leads to a non-decaying near-zero correlation function (Fig. 4c, bottom), also in line with results obtained from mice.

Neuronal network model demonstrates linear growth of correlation lengths unique to critical dynamics

We simulated a neural network (see Materials and Methods and refs. (Kinouchi & Copelli, 2006; Ribeiro et al., 2014) for more details) to establish to which extent the results obtained in the experimental data can be used as an identifier for criticality. Specifically, we explore the impact of windowed access to a larger system, as this is currently the only way to examine finite-size effects on brain activity measures.

Furthermore, given the less than an order of magnitude spatial range we were able to access with 2PI and high-density microelectrode arrays, we estimated the expected deviation in correlation length estimates

when networks move away from criticality. In the simulations we performed the correlation analysis exactly as described for the experimental data using a network of 1000×1000 units, with neurons interacting up to a distance $I_c = 20$. As can be seen in Fig. 5a, the correlations in the fluctuations were highest and extended for the longest distance in the critical regime (Fig. 5a, middle). Furthermore, the growth of ξ with the observed grid extent L was bounded for sub- and supercritical regimes (Fig. 5a, left and right), while in the critical regime, it increased with L . In fact, Fig. 5b shows that ξ peaked at criticality ($\sigma \simeq 1$; see Materials and Methods), regardless of the extent of the observed grid (Fig. 5b, different colors). We note that the absolute difference in the value of correlations from our simulations compared to experimental data simply results from the smoothing procedure employed in the latter (see Materials and Methods). In summary, the linear scaling of ξ as a function of L was unique to criticality, with sub- and supercritical systems presenting asymptotic behavior for ξ over L (Fig. 5c, different colors) and our simulations support our use of observation windows of different size as a proxy of system size in the analysis of correlations of critical systems. This approach was important so that the simulations properly mimic what is observed from the brain: only a small subset of the full network is then further subsampled in order to obtain the scaling of the correlation length. In Supplemental Figure S2, we provide further demonstration that this windowing approach (as opposed to measuring the correlation length as function of system size) is a valid method using the paradigmatic Ising model. Finally, in Fig. 5d we demonstrate that the susceptibility χ also peaks at criticality.

Altogether, these results demonstrate that at the critical regime, the range at which the fluctuations from two units in the network are correlated extends much further than the range at which units are connected, the interaction length I_c (see Materials and Methods). In contrast, for a system far away from the critical point, correlations cannot extend significantly beyond I_c (Fig. 5c, dashed line at 20 coincides

with asymptote value for ξ at $\sigma = 0.16$). However, as can be seen in Fig. 5c, even near the critical regime the linear growth of the correlation length is not present at smaller scales.

Finally, we studied how $\xi(L)$ changed as a function of L_c for critical systems. Linear scaling was not obeyed for grid extents shorter than L_c , for which the curve abruptly changed from the asymptotic slope to a sharper decay (Suppl. Fig. S3a, *left*; see also Supplementary Materials). Indeed, scaling ξ and L by L_c , we obtained a collapse of all curves (Suppl. Fig. S3a, *right*). This scaling function clearly identified two regimes: scale free behavior took place for $L > L_c$, while ξ increased much more abruptly for increasing $L < L_c$, due to direct connections between units. We observed similar trends for individual mice in V1 when replotting the correlation lengths as a function of window extent with the linear regime being preceded by a steeper increase (Suppl. Fig. S3b, *left*). We estimated the point where the scaling abruptly changed for each mouse. They ranged from ~ 100 to $180 \mu\text{m}$ for V1 networks. These estimates were used to successfully collapse all data (Suppl. Fig. S3, *right*).

Discussion

Here, we used a correlation approach to assess the dynamical regime of the cortex during information processing epochs. The correlation length, a measure of how far apart neurons in cortex are positively correlated was shown to scale approximately linearly with the size of the observed window. If the distance over which neuronal activities correlate were to be finite, e.g. $200 \mu\text{m}$, our approach would have revealed an upper bound in window size, or cortical area, for which the correlation length saturated. Instead, both of our experimental approaches, first, for up to $400 \mu\text{m}$ at the level of individual neuronal firing in primary visual and auditory cortex using 2-photon imaging and second, for up to 4 mm in the local field potential in awake nonhuman primates using high-density arrays, demonstrated that correlation length simply grows linearly with the size of the cortical area observed. Our simulations confirmed that our approach using

windowing identifies linear growths in networks with critical dynamics, for which correlation length growths unbounded up to the final observed size.

We evaluated the correlation length as function of subsamples of the recorded region, i.e. compact windows, and not as a function of system size, which is the more common approach in physics. While it is known that windowing differs to a certain degree from finite size effects (Chen *et al.*, 2011), we demonstrated for two different models that our method leads to the similar results. Given that the finite-size changes cannot be executed for real brains, windowing is therefore a realistic, alternative approach. Further support of our windowing method has been provided in BOLD fMRI recordings for the whole brain. In that study (Fraiman & Chialvo, 2012) it was shown that the linear growth in correlation length with the size of separate functional areas of the brain also describes the behavior of the correlation length when different subsets of multiple areas are added together in the analysis.

We used the fluctuations of local neuronal activity around the instantaneous mean of the observed network size to evaluate individual trials. The subtraction of the (observed) population average before calculating correlations has been successfully applied in the context of bird flocks (Cavagna *et al.*, 2010), where one needs to evaluate how birds move in relation to one-another, disregarding the overall movement of the flock. This approach is much more common in physics (Wilson, 1976; Wilson, 1979) and, to our knowledge, has been applied here for the first time to single neurons and local neuronal populations in the wake animal. One might reconcile this common method by realizing that during sensory input or motor output, many neurons integrate similar inputs that consequently drive the activity of the network as a whole. Accordingly, subtracting the instantaneous population average reduces this ‘drive’ component, allowing the correlation analysis to focus on how activity in neurons changes in relation to one-another, independent from the general sensory/motor responses observed. This interpretation is thoroughly supported by our trial-shuffling results, in which the specific inter-neuron relationships that differ from

trial-by-trial are removed and our correlation analysis correctly predicted zero correlations. We note that this method is not advisable for small number of neurons in which subtracting the average introduces dominant anti-correlations artificially decreasing correlation lengths for small windows. We mitigated this effect by imposing a minimum number of units in a given window before considering it for analysis. Small windows also suffer from the influence of the interaction length. Specifically, when the interaction length is on the order of the observed system size, the growth of the correlation length naturally is much faster with system size. We used this to our advantage, which allowed us to estimate the typical interaction range for superficial layers in the primary visual cortex.

That the cortical state is known to affect trial-by-trial variability (Kisley & Gerstein, 1999) and correlation among neurons (Cohen & Kohn, 2011; Rosenbaum *et al.*, 2016) has been a long-standing observation. It is also well established that part of this variability originates from the cortical network itself (Kara *et al.*, 2000; Sadagopan & Ferster, 2012; Goris *et al.*, 2014; Schölvink *et al.*, 2015). Our demonstration of linear scaling in correlation length suggests critical dynamics as the framework that captures the intracortical correlation structure underlying this variability. Critical dynamics has been a fundamental driver in understanding optimization of information processing in complex systems in light of the evidence that fluctuations or variability are high at criticality (e.g. refs. (Shew *et al.*, 2011; Fraiman & Chialvo, 2012; Tkačik *et al.*, 2013; Karimipour *et al.*, 2017)). Decades ago, it was suggested that critical dynamics optimize information transfer in gene-regulation networks (Kauffman, 1969; Sole *et al.*, 1999; Rämö *et al.*, 2007; Nykter *et al.*, 2008). Since then, the criticality hypothesis (Beggs & Plenz, 2003; Chialvo, 2010; Mora & Bialek, 2011; Plenz, 2012; Hesse & Gross, 2014; Marković & Gros, 2014; Plenz & Niebur, 2014; Bettger, 2017; Cocchi *et al.*, 2017; Muñoz, 2018) has gained much ground in the field of neuroscience. Highly desirable aspects of information processing have been shown to improve at criticality such as the maximization of mutual information between stimulus input and output (Kinouchi

& Copelli, 2006; Shew *et al.*, 2009; Shew & Plenz, 2013; Gautam *et al.*, 2015; Bortolotto *et al.*, 2016), increased information capacity (i.e. the number of possible internal states a network can establish) (Haldeman & Beggs, 2005; Shew *et al.*, 2011; Tkačik *et al.*, 2015), improved stimulus discrimination (Shriki & Yellin, 2016; Clawson *et al.*, 2017), and the ability of neurons to flexibly change synchronization while maintaining an overall robust degree of phase-locking (Jantzen *et al.*, 2009; Yang *et al.*, 2012; Kelso *et al.*, 2013; Kirst *et al.*, 2017). Accordingly, our findings support the notion that trial-by-trial variability rather than reflecting pure noise, represents an intrinsic property of cortical dynamics during information processing.

Previous reports relied solely on the calculation of avalanche statistics in examining how close cortical networks are to critical dynamics. Such statistics is sensitive to non-stationarities in neuronal activity as found in evoked responses (Yu *et al.*, 2017), which might explain the lack of avalanche statistics during sensory processing in *ex vivo* turtle (Shew *et al.*, 2015) and the human MEG (Arviv *et al.*, 2015). Power laws statistics as found for avalanches can in principle arise in critically balanced systems in the absence of correlations (see e.g. neutral avalanches (Martinello *et al.*, 2017) and alternative models (Williams-García *et al.*, 2014; Aitchison *et al.*, 2016; Ioffe & Berry, 2017; Touboul & Destexhe, 2017). Our present results, therefore, which are based on the scaling of spatial correlations, provide a new experimental underpinning that cortical processing is in line with critical dynamics (Chialvo, 2010; Shew & Plenz, 2013).

Materials and Methods

All procedures followed the Institute of Laboratory Animal Research (part of the National Research Council of the National Academy of Sciences) guidelines and were approved by the NIMH Animal Care and Use Committee or by the University of Maryland Institutional Animal Care and Use Committee.

294 **Mouse surgery and preparation**

295 Wild type (C57/Bl6, Jackson Laboratory) mice were housed under a reversed 12 h-light/12 h-dark cycle
 296 with ad libitum access to food and water. Imaging experiments were generally performed near the end of
 297 the light and beginning of the dark cycle. A custom-made titanium head bar was surgically implanted onto
 298 the skull of the mice under isoflurane anesthesia (4% induction, 1-1.5% maintenance). A circular
 299 craniotomy (~3 mm) was made above the area of interest (visual or auditory cortex), followed by injection
 300 of a virus containing the genetically encoded calcium indicator (YC2.6 for V1; GCaMP6s for A1) at a
 301 depth of ~250-300 μm . After that, a cranial window composed of two 3 mm diameter coverslips glued to
 302 a 5 mm coverslip was implanted and the entire area (except for the window) was sealed with dental cement
 303 (Goldey *et al.*, 2014; Bowen *et al.*, 2019).

304

305 **Visual stimulation and response measures**

306 Visual stimuli were prepared in Matlab using the Psychophysics Toolbox (Kleiner *et al.*, 2007) and
 307 delivered via a monitor (Dell, 60 Hz refresh rate) placed ~25 cm in front of the contra-lateral eye of the
 308 mouse. The stimulus was composed of moving gratings at 8 different directions presented for 2 s at
 309 maximum contrast, 0.04 cycles per degree and 2 cycles per sec. Stimuli were interspaced by gray screen
 310 (matched for average luminance) for 2 s. Each direction was presented 20 times in randomized order, for
 311 a total of 160 iterations. We calculated the direction selectivity index using the common definition: $DSI =$
 312 $(R_P - R_O)/(R_P + R_O)$, where R_P and R_O are the responses to the preferred and opposite direction,
 313 respectively. Significance of DSI for each cell was assessed by comparing the values obtained from the
 314 original data with those obtained from shuffling the inter-spike intervals.

315 **Acoustic stimulation**

316 Sound stimuli were synthesized in Matlab using custom software, passed through a multifunction
317 processor (RX6, TDT), attenuated (PA5, Programmable Attenuator), and delivered via ES1 speaker placed
318 ~5 cm directly in front of the mouse. The sound system was calibrated between 2.5 and 80 kHz and showed
319 a flat (± 3 dB) spectrum over this range. Overall SPL at 0 dB attenuation was ~90 dB SPL on average (for
320 tones). Sounds were played at a range of sound levels (40-80 dB SPL). Auditory stimuli consisted of
321 sinusoidal amplitude-modulated (SAM) tones (20 Hz modulation, cosine phase), ranging from 3 to 48
322 kHz. The frequency resolution was 2 tones/octave (0.5 octave spacing). Each of these tonal stimuli was
323 repeated 5 times with a 6 second inter-stimulus interval, for a total of 135 iterations (Bowen *et al.*, 2019).

325 **Two-photon imaging and analysis**

326 Images were acquired by a scanning microscope (Bergamo II series, B248, Thorlabs) coupled to a pulsed
327 femtosecond Ti:Sapphire 2-photon laser with dispersion compensation (Vision S, Coherent). The
328 microscope was controlled by ThorImageLS software. The wavelength was tuned to either 830 nm or 940
329 nm in order to excite YC2.6 or GCaMP6s, respectively. Signals were collected through a 16 $\times$ 0.8 NA
330 microscope objective (Nikon). Emitted photons were directed through 525/50 nm (green) and 607/70 nm
331 (red) band filters (for GCaMP6s) or 535/22 nm (yellow) and 479/40 nm (cyan) band filters (for YC2.6)
332 onto GaAsP photomultiplier tubes. The field of view was ~400 x 400 μ m. Imaging frames of 512 $\times$ 512
333 pixels were acquired at 30 Hz by bidirectional scanning of an 8 kHz resonant scanner. Beam turnarounds
334 at the edges of the image were blanked with a Pockels cell. The average power for imaging was <70 mW,
335 measured at the sample. The obtained images were corrected for motion using dft registration software
336 with Matlab (Guizar-Sicairos *et al.*, 2008). Regions of interest (ROI) were identified from the average
337 image of the motion corrected sequence using custom code. For each labeled neuron, raw fluorescence

signals (cyan and yellow for YC2.6; green for GCaMP6s) over time were extracted from the ROI overlying the soma. The mean ratiometric signal (R ; YC2.6) or single fluorescence (F ; GCaMP6s) in each ROI was calculated across frames and converted to a relative fluorescence measure ($\Delta R/R_0$ or $\Delta F/F_0$). The baseline signal R_0 (or F_0) was estimated by using a sliding window that calculated the average fluorescence of points less than the 10th-percentile during the previous 1.3-second window (40 frames).

Monkey behavioral training and electrophysiological setup

Experiments were described previously (Yu *et al.*, 2017). In short, two adult rhesus monkeys (*Macaca mulatta*) were surgically implanted with a titanium head post. After recovery, they were trained to sit head-fixed in a primate chair for behavioral performance. In the cue-initiated task, monkey A (male, 9 years old, 8 kg) had to press a bar in front of the chair upon presentation of the ‘trial-initiation’ cue on a computer screen. After ~2 s, the initiation cue was followed by an ‘instruction’ cue, for the duration of 1 s. Upon cue disappearance, monkey A had to release the bar and reach with his right arm to one of two specialized feeders, depending on which of two possible cues were presented (Mitz *et al.*, 2001). Approaching the incorrect feeder rapidly triggered a proximity sensor to sequester the food rewards in both feeders, which prevented the monkey from obtaining a reward on that trial. The inter trial interval was 3 – 5 s. In the self-initiated motor task, monkey B (female, 8 years old, 7 kg) had to move her right arm to touch a pad placed ~30 cm in front of the monkey chair after which a food reward was given. After the monkeys learned their respective tasks, a multi-electrode array (MEA; 96 channels - 10×10 without corners, inter-electrode distance: 400 μ m; electrode length: 1 mm for monkey A and 0.55 mm for monkey B; BlackRock Microsystems) was chronically implanted in the arm representative region of the left prefrontal area (area 46, monkey A) or the left premotor cortex (monkey B). The LFP (1 – 100 Hz band pass filtered; 2 kHz sampling frequency) was obtained from the implanted MEA. Electrophysiological signals as well as the

timing of behaviorally relevant events, e.g. touching the pad, presentation of visual cues, etc., were stored for off-line analysis.

Numerical simulations

We simulated a neural network, as described previously (Kinouchi & Copelli, 2006; Ribeiro *et al.*, 2014). In short, each neuron can be in one of three states at each time step: 0 for resting, 1 for active, and 2 for refractory. The model considers S^2 neurons on a square lattice. Each neuron outputs to K other neurons, selected with an exponentially decaying probability function of the Euclidian distance r between them ($P_{Conn} \sim e^{-r/R_0}$, with $R_0 = 5$). A spatial cutoff is set in the interaction distance: neurons cannot directly connect at distances greater than $I_c = 4R_0$ spatial units (thus, defining an interaction length I_c). Furthermore, to reduce small S effects, we employed periodic boundary conditions. Results were computed on a square grid of length L . Simulations parameters were always such that S was much larger than L to reduce finite-size artifacts and to better mimic experimental data. A small Poisson drive ($h = 10^{-7}$ per time step) to each neuron determined the overall rate of firing activity. The present results were robust over a wide range of h values (e.g., $h = 10^{-9}$ to 10^{-4} per time step). The control parameter of the model determines the branching of the neural activity and was defined as $\sigma = K \times T$, where T is the probability that an active neuron (i.e., in state 1) can excite each one of the K neighbors that it connects to. Therefore, as shown previously, the model can be made critical by selecting a transmission probability T such that $\sigma \sim 1$, for any given K . Susceptibility was defined as the area under the correlation vs. distance curve, for a given window size observed, up to the correlation length.

Considering the focus of the present study, we note that there are four length scales in the model. The *interaction length* (here called I_c) is the scale at which neurons can interact via direct connections. *System length* (called S) determines system size (the network is composed of S^2 neurons). The third one is the

window length L ($L \leq S$), called *grid length*, which determines how many neurons we will measure from (i.e., L^2 neurons). The last scale is the *correlation length* ξ , the longest distance at which on average the activity of any two given neurons may remain positively correlated. It is calculated from fluctuations on activity, following Cavagna's work (Cavagna *et al.*, 2010) (see section below). To avoid confusions, we remark that the term *interaction* is reserved here to denote *direct connections* and *correlations* to the mathematical result from computing *correlations of neural activity*. Simulations for varying interaction length (I_c) were carried out using the same model described above. When changing I_c , the decaying parameter used for making connections R_0 was changed accordingly (such that the relation $I_c = 4R_0$ still holds).

Avalanche analysis

Avalanche analysis for the 2PI data was performed as described previously (Beggs & Plenz, 2003; Bellay *et al.*, 2015). In short, estimated spikes (Deneux *et al.*, 2016) from all neurons were pooled together to create a population activity series. Avalanches are defined by contiguous non-zero population activity preceded and followed by blank frames (frames with zero population activity). Avalanche sizes were defined by the total number of spikes throughout their lifetime. Power-law goodness of fit was evaluated through a p-value calculated from the log-likelihood ratio when comparing power law, exponential and lognormal fits, as described previously (Clauset *et al.*, 2009; Klaus *et al.*, 2011; Bellay *et al.*, 2015).

Correlation analysis

The correlation of the fluctuations as function of distance (Cavagna *et al.*, 2010) was calculated as

$$C(r) = \frac{1}{C_0} \frac{\sum_{i,j} u_i u_j \delta(r - r_{ij})}{\sum_{i,j} \delta(r - r_{ij})}, \quad (1)$$

where $\delta(r - r_{ij})$ is a smoothed Dirac δ function defining all pairs of neurons located at mutual distance r , r_{ij} is the Euclidean distance from the i -th neuron's spatial location to the spatial location of neuron j , and u_i is the value of the signal v of neuron i at time t , after subtracting the overall mean of signals v from neurons inside the observation window of size L at that time t : $u_i(t) = v_i(t) - \bar{v}(t)$. To ensure that $C(r = 0) = 1$, the normalization factor $1/C_0$ was used. We note that since the instantaneous average is subtracted, $C(r)$ is not equivalent to the most commonly used pairwise Pearson correlation function.

The objective of computing the $C(r)$ is to determine the correlation length ξ , which is defined as the point where the correlations of the fluctuations reaches zero, i.e. $C(\xi) = 0$. Since system size, i.e. cortex size in our experimental data, was fixed, we investigated how ξ changes with system size subsampled by our recordings and considered neurons/electrodes within a window of length L (this proxy is validated by computing correlation lengths both as function of increasing system and window sizes in a model (Suppl. Fig. S2). More specifically, for the 2PI data in mice, fields of view ranging from $\sim 40 \times 40 \mu\text{m}$ (windows with fewer than 5 units were ignored to avoid bias introduced by the average subtraction procedure when the number of units is too small) to the maximum possible size were considered, while for the monkey LFP data the smallest subarray considered was 3×3 . To reduce noise effects, results were averaged across all possible subregions for any given size. The time series were smoothed in the time domain (using Matlab routine *medfilt1.m* with 20 samples for the mice 2PI data, 8 samples for the monkey LFP data). This smoothing procedure improved statistics without changing the results qualitatively. In order to more precisely estimate the zero-crossing point for the experimental data, we fit 3rd order polynomial functions to the $C(r)$ curves around the zero-crossing.

To quantify linear growth in correlation length ξ as function of window length L , we first obtained a linear regression of the $\xi(L)$ data followed by chi-square statistics $\chi_c^2 = \sum_i [\xi(L_i) - R(L_i)]^2 / R(L_i)$, where L_i is the i^{th} measured value of L and $R(L_i)$ is the linear regression value at L_i . χ_c^2 can be used to obtain a p-

value that estimates how likely the data fit the linear regression that well by chance from the chi-square distribution. We rescaled the correlation *vs.* distance curves by normalizing the distances by the correlation length and by rescaling the correlations by the correlation length to the power of γ , defined as the slope of the curves at zero-crossing (Cavagna *et al.*, 2010).

Trial shuffling and spatial shuffling

Trial shuffling for the V1 data was obtained by randomly permuting the responses from each of the 8 presented directions separately. This was done for each neuron independently. Therefore, in each trial of the trial shuffled dataset activity from each cell corresponds to a response to the same stimulus presented in the original data but taken from different presentations of that stimulus. For example, suppose we presented stimulus 5 in trial 1. In the shuffled data, the response of neuron 1 in that trial may be taken from the 10th presentation of stimulus 5, while response of neuron 2 may be taken from the 3rd presentation. Spatial shuffling was performed by randomly permuting cell positions, leaving everything else unchanged.

Acknowledgements

We thank members of the Plenz lab for lively discussions. This research was supported by the Division of the Intramural Research Program (DIRP) of the National Institute of Mental Health (NIMH), USA, ZIA MH00297 and the BRAIN initiative Grant U19 NS107464-01. D.A.M. acknowledges financial support from ANPCyT Grant No. PICT-2016-3874 (Argentina). This research utilized the computational resources of Biowulf (<http://hpc.nih.gov>) at the National Institutes of Health (NIH), USA, and UnCaFiQT-INIFTA (SNCAD), Argentina.

Contributions

T.L.R., D.R.C. and D.P. conceived and planned the study; T.L.R., S.Y., D.W., P.K. and D.P. oversaw and carried out experiments; T.L.R., D.A.M., D.R.C and D.P. designed the models and carried out the simulations; T.L.R. took the lead in the analysis; T.L.R., D.R.C. and D.P. wrote the paper.

Conflicts of Interest

The authors declare no competing financial interests.

References

- Aitchison, L., Corradi, N. & Latham, P.E. (2016) Zipf's law arises naturally when there are underlying, unobserved variables. *PLOS Computational Biology*, **12**, e1005110.
- Arieli, A., Sterkin, A., Grinvald, A. & Aertsen, A. (1996) Dynamics of ongoing activity: Explanation of the large variability in evoked cortical responses. *Science*, **273**, 1868-1871.
- Arviv, O., Goldstein, A. & Shriki, O. (2015) Near-critical dynamics in stimulus-evoked activity of the human brain and its relation to spontaneous resting-state activity. *The Journal of Neuroscience*, **35**, 13927-13942.
- Beggs, J.M. & Plenz, D. (2003) Neuronal avalanches in neocortical circuits. *J Neurosci*, **23**, 11167-11177.
- Bellay, T., Klaus, A., Seshadri, S. & Plenz, D. (2015) Irregular spiking of pyramidal neurons organizes as scale-invariant neuronal avalanches in the awake state. *eLife*, **4**, e07224.
- Bettinger, J.S. (2017) Comparative approximations of criticality in a neural and quantum regime. *Progress in Biophysics and Molecular Biology*, **131**, 445-462.

- 479 Bialek, W., Cavagna, A., Giardina, I., Mora, T., Pohl, O., Silvestri, E., Viale, M. & Walczak, A.M.
480 (2014) Social interactions dominate speed control in poising natural flocks near criticality. **111**,
481 7212-7217.
482
- 483 Bialek, W., Cavagna, A., Giardina, I., Mora, T., Silvestri, E., Viale, M. & Walczak, A.M. (2012)
484 Statistical mechanics for natural flocks of birds. **109**, 4786-4791.
485
- 486 Bortolotto, G.S., Girardi-Schappo, M., Gonsalves, J.J., Pinto, L.T. & Tragtenberg, M.H.R. (2016)
487 Information processing occurs via critical avalanches in a model of the primary visual cortex.
488 *Journal of Physics: Conference Series*, **686**, 012008.
489
- 490 Bowen, Z., Winkowski, D.E., Seshadri, S., Plenz, D. & Kanold, P.O. (2019) Neuronal avalanches in
491 input and associative layers of auditory cortex. *Frontiers in Systems Neuroscience*, **13**.
492
- 493 Carandini, M. (2004) Amplification of trial-to-trial response variability by neurons in visual cortex.
494 *PLoS Biol*, **2**, E264.
495
- 496 Cavagna, A., Cimorelli, A., Giardina, I., Parisi, G., Santagati, R., Stefanini, F. & Viale, M. (2010) Scale-
497 free correlations in starling flocks. *Proc Natl Acad Sci U S A*, **107**, 11865-11870.
498
- 499 Charles, A.S., Park, M., Weller, J.P., Horwitz, G.D. & Pillow, J.W. (2018) Dethroning the Fano factor: a
500 flexible, model-based approach to partitioning neural variability. *Neural Computation*, **30**, 1012-
501 1045.
502
- 503 Chen, Y.-J., Papanikolaou, S., Sethna, J.P., Zapperi, S. & Durin, G. (2011) Avalanche spatial structure
504 and multivariable scaling functions: Sizes, heights, widths, and views through windows. **84**.
505
- 506 Chialvo, D.R. (2010) Emergent complex neural dynamis. *Nat. Phys.*, **6**, 744-750.
507
- 508 Clauset, A., Shalizi, C.R. & Newman, M.E.J. (2009) Power-law distributions in empirical data. *Siam*
509 *Review*, **51**, 661-703.

510

511 Clawson, W.P., Wright, N.C., Wessel, R. & Shew, W.L. (2017) Adaptation towards scale-free dynamics
512 improves cortical stimulus discrimination at the cost of reduced detection. *PLOS Computational*
513 *Biology*, **13**, e1005574.

514

515 Cocchi, L., Gollo, L.L., Zalesky, A. & Breakspear, M. (2017) Criticality in the brain: A synthesis of
516 neurobiology, models and cognition. *Progress in Neurobiology*, **158**, 132-152.

517

518 Cohen, M.R. & Kohn, A. (2011) Measuring and interpreting neuronal correlations. *Nat. Neurosci.*, **14**,
519 811-819.

520

521 Dechery, J.B. & MacLean, J.N. (2018) Functional triplet motifs underlie accurate predictions of single-
522 trial responses in populations of tuned and untuned V1 neurons. *PLOS Computational Biology*,
523 **14**, e1006153.

524

525 Deneux, T., Kaszas, A., Szalay, G., Katona, G., Lakner, T., Grinvald, A., Rozsa, B. & Vanzetta, I. (2016)
526 Accurate spike estimation from noisy calcium signals for ultrafast three-dimensional imaging of
527 large neuronal populations in vivo. *Nat Commun*, **7**.

528

529 Deweese, M.R. & Zador, A.M. (2004) Shared and private variability in the auditory cortex. *J*
530 *Neurophysiol*, **92**, 1840-1855.

531

532 Fraiman, D. & Chialvo, D.R. (2012) What kind of noise is brain noise: anomalous scaling behavior of
533 the resting brain activity fluctuations. *Front Physiol*, **3**, 307.

534

535 Gautam, S.H., Hoang, T.T., McClanahan, K., Grady, S.K. & Shew, W.L. (2015) Maximizing sensory
536 dynamic range by tuning the cortical state to criticality. *PLoS Comput Biol*, **11**, e1004576.

537

538 Goldey, G.J., Roumis, D.K., Glickfeld, L.L., Kerlin, A.M., Reid, R.C., Bonin, V., Schafer, D.P. &
539 Andermann, M.L. (2014) Removable cranial windows for long-term imaging in awake mice. *Nat*
540 *Protoc*, **9**, 2515-2538.

541

542 Goris, R.L.T., Movshon, J.A. & Simoncelli, E.P. (2014) Partitioning neuronal variability. *Nature*
543 *Neuroscience*, **17**, 858.

544

545 Guizar-Sicairos, M., Thurman, S.T. & Fienup, J.R. (2008) Efficient subpixel image registration
546 algorithms. *Opt Lett*, **33**, 156-158.

547

548 Haldeman, C. & Beggs, J.M. (2005) Critical branching captures activity in living neural networks and
549 maximizes the number of metastable States. *Phys Rev Lett*, **94**, 058101.

550

551 Haroush, N. & Marom, S. (2019) Inhibition increases response variability and reduces stimulus
552 discrimination in random networks of cortical neurons. *Scientific Reports*, **9**, 4969.

553

554 Heggelund, P. & Albus, K. (1978) Response variability and orientation discrimination of single cells in
555 striate cortex of cat. *Experimental brain research. Experimentelle Hirnforschung.*
556 *Expérimentation cérébrale*, **32**, 197-211.

557

558 Hesse, J. & Gross, T. (2014) Self-organized criticality as a fundamental property of neural systems. **8**.

559

560 Ioffe, M.L. & Berry, M.J., II (2017) The structured ‘low temperature’ phase of the retinal population
561 code. *PLOS Computational Biology*, **13**, e1005792.

562

563 Jantzen, K.J., Steinberg, F.L. & Kelso, J.A.S. (2009) Coordination dynamics of large-scale neural
564 circuitry underlying rhythmic sensorimotor behavior. **21**, 2420-2433.

565

566 Kara, P., Reinagel, P. & Reid, R.C. (2000) Low response variability in simultaneously recorded retinal,
567 thalamic, and cortical neurons. *Neuron*, **27**, 635-646.

568

569 Karimipani, Y., Ma, Z., Miller, J.-E.K., Yuste, R. & Wessel, R. (2017) Neocortical activity is stimulus-
570 and scale-invariant. *PloS one*, **12**, e0177396-e0177396.

571

- 572 Kauffman, S. (1969) Homeostasis and differentiation in random genetic control networks. *Nature*, **224**,
573 177-178.
- 574
- 575 Kelso, J.A.S., Dumas, G. & Tognoli, E. (2013) Outline of a general theory of behavior and brain
576 coordination. *Neural Networks*, **37**, 120-131.
- 577
- 578 Kinouchi, O. & Copelli, M. (2006) Optimal dynamical range of excitable networks at criticality. *Nature*
579 *Physics*, **2**, 348-352.
- 580
- 581 Kirst, C., Modes, C.D. & Magnasco, M.O. (2017) Shifting attention to dynamics: Self-reconfiguration of
582 neural networks. *Current Opinion in Systems Biology*, **3**, 132-140.
- 583
- 584 Kisley, M.A. & Gerstein, G.L. (1999) Trial-to-trial variability and state-dependent modulation of
585 auditory-evoked responses in cortex. *Journal of Neuroscience*, **19**, 10451-10460.
- 586
- 587 Klaus, A., Yu, S. & Plenz, D. (2011) Statistical analyses support power law distributions found in
588 neuronal avalanches. *PLoS One*, **6**, e19779.
- 589
- 590 Kleiner, M., Brainard, D. & Pelli, D. (2007) What's new in Psychtoolbox-3? *Perception*, **36**, 14-14.
- 591
- 592 Kotekal, S. & MacLean, J.N. (2019) Recurrent interactions can explain the variance in single trial
593 responses. *bioRxiv*, 635359.
- 594
- 595 Marković, D. & Gros, C. (2014) Power laws and self-organized criticality in theory and nature. *Physics*
596 *Reports*, **536**, 41-74.
- 597
- 598 Martinello, M., Hidalgo, J., Maritan, A., di Santo, S., Plenz, D. & Muñoz, M.A. (2017) Neutral theory
599 and scale-free neural dynamics. *Physical Review X*, **7**, 041071.
- 600

- Mitz, A.R., Boring, S.A., Wise, S.P. & Lebedev, M.A. (2001) A novel food-delivery device for neurophysiological and neuropsychological studies in monkeys. *Journal of Neuroscience Methods*, **109**, 129-135.
- Mora, T. & Bialek, W.J.J.o.S.P. (2011) Are biological systems poised at criticality? , **144**, 268-302.
- Muñoz, M.A. (2018) Colloquium: Criticality and dynamical scaling in living systems. *Reviews of Modern Physics*, **90**, 031001.
- Nykter, M., Price, N.D., Larjo, A., Aho, T., Kauffman, S.A., Yli-Harja, O. & Shmulevich, I. (2008) Critical networks exhibit maximal information diversity in structure-dynamics relationships. *Phys Rev Lett*, **100**, 058702.
- Plenz, D. (2012) Neuronal avalanches and coherence potentials. *The European Physical Journal Special Topics*, **205**, 259-301.
- Plenz, D. & Niebur, E. (2014) *Criticality in Neural Systems*. Wiley Blackwell.
- Rämö, P., Kauffman, S., Kesseli, J. & Yli-Harja, O. (2007) Measures for information propagation in Boolean networks. *Physica D*, **227**, 100-104.
- Ribeiro, T.L., Ribeiro, S., Belchior, H., Caixeta, F. & Copelli, M. (2014) Undersampled critical branching processes on small-world and random networks fail to reproduce the statistics of spike avalanches. *Plos One*, **9**.
- Rosenbaum, R., Smith, M.A., Kohn, A., Rubin, J.E. & Doiron, B. (2016) The spatial structure of correlated neuronal variability. *Nature Neuroscience*, **20**, 107.
- Sadagopan, S. & Ferster, D. (2012) Feedforward origins of response variability underlying contrast invariant orientation tuning in cat visual cortex. *Neuron*, **74**, 911-923.

- Schölvinck, M.L., Saleem, A.B., Benucci, A., Harris, K.D. & Carandini, M. (2015) Cortical state determines global variability and correlations in visual cortex. *The Journal of Neuroscience*, **35**, 170.
- Shadlen, M.N. & Newsome, W.T. (1998) The variable discharge of cortical neurons: implications for connectivity, computation, and information coding. *J. Neurosci.*, **18**, 3870-3896.
- Shew, W.L., Clawson, W.P., Pobst, J., Karimipani, Y., Wright, N.C. & Wessel, R. (2015) Adaptation to sensory input tunes visual cortex to criticality. *Nat Phys*, **11**, 659-663.
- Shew, W.L. & Plenz, D. (2013) The functional benefits of criticality in the cortex. *Neuroscientist*, **19**, 88-100.
- Shew, W.L., Yang, H., Petermann, T., Roy, R. & Plenz, D. (2009) Neuronal avalanches imply maximum dynamic range in cortical networks at criticality. *J Neurosci*, **29**, 15595-15600.
- Shew, W.L., Yang, H., Yu, S., Roy, R. & Plenz, D. (2011) Information capacity and transmission are maximized in balanced cortical networks with neuronal avalanches. *J Neurosci*, **31**, 55-63.
- Shriki, O. & Yellin, D. (2016) Optimal information representation and criticality in an adaptive sensory recurrent neuronal network. *PLOS Computational Biology*, **12**, e1004698.
- Sole, R.V., Luque, B. & Kauffman, S. (1999) Phase transition in random networks with multiple states *arXiv e-prints*.
- Tkačik, G., Marre, O., Mora, T., Amodei, D., Berry II, M.J. & Bialek, W. (2013) The simplest maximum entropy model for collective behavior in a neural network. *Journal of Statistical Mechanics: Theory and Experiment*, **2013**, P03011.
- Tkačik, G., Mora, T., Marre, O., Amodei, D., Palmer, S.E., Berry, M.J. & Bialek, W. (2015) Thermodynamics and signatures of criticality in a network of neurons. **112**, 11508-11513.

663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688

- Tolhurst, D.J., Movshon, J.A. & Dean, A.F. (1983) The statistical reliability of signals in single neurons in cat and monkey visual cortex. *Vision Res*, **23**, 775-785.
- Touboul, J. & Destexhe, A. (2017) Power-law statistics and universal scaling in the absence of criticality. *Physical Review E*, **95**, 012413.
- Vogels, R., Spileers, W. & Orban, G. (1989) The response variability of striate cortical neurons in the behaving monkey. *Experimental brain research. Experimentelle Hirnforschung. Expérimentation cérébrale*, **77**, 432-436.
- Williams-García, R.V., Moore, M., Beggs, J.M. & Ortiz, G. (2014) Quasicritical brain dynamics on a nonequilibrium Widom line. *Physical Review E*, **90**, 062714.
- Wilson, K.G. (1976) *Phase Transitions and Critical Phenomena*. Academic Press, New York.
- Wilson, K.G. (1979) Problems in Physics with Many Scales of Length. *Sci Am*, **241**, 158-&.
- Yang, H., Shew, W.L., Roy, R. & Plenz, D. (2012) Maximal variability of phase synchrony in cortical networks with neuronal avalanches. *J Neurosci*, **32**, 1061-1072.
- Yu, S., Ribeiro, T.L., Meisel, C., Chou, S., Mitz, A., Saunders, R. & Plenz, D. (2017) Maintained avalanche dynamics during task-induced changes of neuronal activity in nonhuman primates. *Elife*, **6**.

Trial-by-trial variability in cortical responses exhibits scaling in spatial correlations **predicted from critical dynamics**

Tiago L. Ribeiro et al.

Supplementary Material

Scaling of correlation lengths in mouse V1 based on spike density estimates

Supplemental Figure S1: Correlation length scales linearly with window extent for spiking data. Correlation as a function of window extent obtained from estimated spike times from V1 in mice. *Line* represents a linear regression.

Correlation length scaling for increasing window size in the Ising model

In critical systems, the correlation length is known to scale with system size. Because in biological systems changing system size is often impractical, in this work we introduce a proxy which uses windows of different sizes. In order to validate that approach, we test the concept in the paradigmatic 2D Ferromagnetic Ising model, which undergoes a critical transition at temperature $T \approx 2.3$. The simulations

17 (lasting at least 10^5 Montecarlo steps) used two setups: in the first, the correlation function was computed
 18 in the standard way, from a model running on square lattices of increasing $S = 16, 32, 64$ and 128 . In the
 19 second setup, a relatively large $S = 300$ square lattice (i.e. 300×300 spins) was simulated, and the
 20 correlation function was computed from square windows of smaller sizes $L = 4, 8, 16, 32, 64$ and 128 .
 21 Representative results for three different temperatures are shown: subcritical ($T = 1.6$, Fig. S2a), critical
 22 ($T = 2.3$, Fig. S2b) and supercritical ($T = 3.0$, Fig. S2c). Fig. S2d shows the correlation lengths computed
 23 from both setups are very similar. Therefore, the curves obtained changing system size S or changing
 24 window size L produce similar correlation lengths. These results validate the use of window size as a
 25 proxy of system size in the analysis of correlations of critical systems.

Supplemental Figure S2: The ferromagnetic 2D Ising model.

a Correlation as a function of distance for the subcritical regime ($T = 1.6$). Color code: different window sizes (L from 4 to 128) in a system of size $S = 300$. Dotted lines: fixed $S = L = 64$ and 128 , as indicated. **b** Results for the critical regime ($T = 2.3$). **c** Results for supercritical regime ($T = 3.0$). **d** Correlation length as a function of distance for $T = 1.6$, $T = 2.3$ and $T = 3$ and different conditions. Empty circles and dashed lines are used for systems with $S = 300$. Filled symbols with continuous lines are used for systems and windows of sizes $S = L$.

28 **Estimating the interaction length from experimental data**

29 We studied how the scaling law changed as a function of the interaction length for critical systems. We fit
30 a power law of exponent 1 with an extended exponential part to the data as follows $\xi(L) = CL\{1 -$
31 $\exp[-(L/L_0)^\varepsilon]\}$, where $C = 0.27$ is a constant that controls the slope of the curves (in linear coordinates),
32 $\varepsilon = 3$ is a constant that controls how sharp the decay for small L is and L_0 is the cross-over point where
33 the behavior changes from the exponential to the linear growth, our estimate for the interaction length.
34 The values for C and ε were found empirically and represent a good match for the curves obtained in the
35 simulations.

36 As shown in Fig. S3a (left), the linear scaling was not obeyed for grid extents shorter than I_c , for which
37 the curve abruptly changed from the asymptotic slope to a sharper decay. If this change in behavior was
38 solely due to the influence of direct connections between units (as opposed to influence of critical
39 dynamics) then we should be able to obtain a universal curve that is invariant to the range at which units
40 can be connected. Indeed, scaling ξ and L by I_c we obtained a collapse of all curves (see Fig. S3a, right).
41 This scaling function clearly identified two regimes: scale free behavior took place for $L > I_c$, while ξ
42 increased much more abruptly for increasing $L < I_c$, due to direct connections between units. We observed
43 similar trends for individual mice in V1 when replotting the correlation lengths as a function of window
44 extent with the linear regime being preceded by a steeper increase (Fig. S3b, left). The point where the
45 scaling abruptly changed differed between mice (Fig. S3b, different colors) and we used a function that
46 mimics the behavior observed for the collapsed curve in Fig. S3a (right) to estimate the interaction length
47 for each mouse (see Materials and Methods). These estimates were used to collapse all functions (Fig.
48 S3b, right). They ranged from ~ 100 to $180 \mu\text{m}$ among mice and are a good estimate for the local distance
49 at which pyramidal cells typically connect in superficial layers in V1 (Levy & Reyes, 2012; Seeman *et*
50 *al.*, 2018).

Supplemental Figure S3: Interaction length in experimental data can be estimated from correlation length growth with observed window size. **a Left:** Correlation length ξ as a function of L for simulations at criticality for 3 different values of interaction length I_c (color code) in log-log coordinates. *Dashed line:* linear growth as a visual guide. **Right:** After rescaling grid extents and correlation lengths by the interaction length, all curves collapse into a single scaling function that highlights the change in the linear behavior at $L = I_c$. In all cases, system size is 1000 x 1000 with connectivity $K = 16$. **b Same as panel a,** but for correlation lengths obtained from individual mice in V1. We estimated the point where the linear growth of the correlation length breaks for each mouse (see Materials and Methods) and used those values to rescale the curves as in panel **a (right)**. The collapse obtained suggests that the estimated points ($\sim 119 \mu\text{m}$ for mouse 1 – orange; $\sim 143 \mu\text{m}$ for mouse 2 – purple; $\sim 108 \mu\text{m}$ for mouse 3 – pink; $\sim 179 \mu\text{m}$ for mouse 4 – dark green; $\sim 180 \mu\text{m}$ for mouse 5 – light green) represent well the interaction length for these mice.

53 **References**

- 54 Levy, R.B. & Reyes, A.D. (2012) Spatial Profile of Excitatory and Inhibitory Synaptic Connectivity in
55 Mouse Primary Auditory Cortex. *Journal of Neuroscience*, **32**, 5609-5619.
56
- 57 Seeman, S.C., Campagnola, L., Davoudian, P.A., Hoggarth, A., Hage, T.A., Bosma-Moody, A., Baker,
58 C.A., Lee, J.H., Mihalas, S., Teeter, C., Ko, A.L., Ojemann, J.G., Gwinn, R.P., Silbergeld, D.L.,
59 Cobbs, C., Phillips, J., Lein, E., Murphy, G., Koch, C., Zeng, H.K. & Jarsky, T. (2018) Sparse
60 recurrent excitatory connectivity in the microcircuit of the adult mouse and human cortex. *Elife*,
61 **7**, e37349.
62