1 **Enlarged cortical cells and reduced cortical cell file number improve growth under**
2 **suboptimal nitrogen, phosphorus and potassium availability**
3 Xiyu Yang, Miranda Niemiec, Jonathan P. Lynch*
4 Department of Plant Science, The Pennsylvania State University, University Park, PA, U.S.A.,
5 16802
6 Xiyu Yang: xby5025@psu.edu
7 Miranda Niemiec: mdn22@psu.edu
8 Jonathan P. Lynch: jpl4@psu.edu
9 *For Correspondence. Email: jpl4@psu.edu, Tel.: +1 8148632256
10 Date of Submission:
11 Number of Figures: 9
12 Word Count:
13
14
15
16
17

Title: Enlarged cortical cells and reduced cortical cell file number improve growth under suboptimal nitrogen, phosphorus and potassium availability

Short title: Root cortical anatomy and nutrient capture

Xiyu Yang, Miranda Niemiec, Jonathan P. Lynch*

Department of Plant Science, The Pennsylvania State University, University Park, PA, U.S.A., 16802

Author contributions: XY conducted the experiments, analyzed data, and wrote the article. MN contributed to data analysis, figure creation, and writing. JPL conceived and supervised the project and contributed to data analysis and writing. JPL agrees to serve as corresponding author.

One sentence summary: Functional-structural modeling indicates that enlarged root cortical cells and reduced cortical cell file number decrease root maintenance cost, permitting greater soil exploration, resource capture, and plant growth under suboptimal nitrogen, phosphorus and potassium availability.

Abstract

Reduced cortical cell files (CCFN) and enlarged cortical cells (CCS) reduce root maintenance costs. We used *OpenSimRoot*, a functional-structural model, to test the hypothesis that larger CCS, reduced CCFN, and their interactions with root cortical aerenchyma (RCA), are useful adaptations to suboptimal soil N, P, and K availability. Interactions of CCS and CCFN with lateral root branching density (LRBD) and increased carbon availability were evaluated under limited N, P and K availability. The combination of larger CCS and reduced CCFN increases the growth of maize up to 105%, 106%, and 144%, respectively, under limited N, P, or K availability. Interactions among larger CCS, reduced CCFN, and greater RCA results in combined growth benefits of up to 135%, 132%, and 161% under limited N, P, and K levels, respectively. Under low phosphorus and potassium availability, increased LRBD approximately doubles the utility of larger CCS and reduced CCFN. The utility of larger CCS and reduced CCFN is reduced by greater C availability as may occur in future climate scenarios. Our results support the hypothesis that larger CCS, reduced CCFN, and their interactions with RCA could increase nutrient acquisition by reducing root respiration and root nutrient demand. Phenetic synergisms may exist between CCS, CCFN, and LRBD. Natural genetic variation in CCS and CCFN merit consideration for breeding cereal crops with improved nutrient acquisition, which is critical for global food security.

Keywords: *Zea mays*, root cortical aerenchyma, cortical cell size, cortical cell file number, nutrient acquisition efficiency, *OpenSimRoot*, functional-structural plant model

Abbreviations:

Cortical cell size (CCS)

Cortical cell file number (CCFN)

Root cortical aerenchyma (RCA)

Steep, Cheap and Deep (SCD)

56 Lateral root branching density (LRBD)

57 Root hair length (RHL)

58 Basal root growth angle (BRGA)

59 **Introduction**

60 The development of crops with reduced fertilizer requirements is needed in global
61 agriculture to reduce the environmental, economic, and energy costs of crop production in high-
62 input agroecosystems and increase crop production in low-input agroecosystems (Koevoets et al.,
63 2016; Lynch, 2019). One avenue towards this goal is via selection for root phenotypes that reduce
64 the metabolic cost of soil exploration (Lynch, 2015). The metabolic costs of root tissues can be
65 estimated as the investment of limiting resources, mainly carbohydrates and limiting mineral
66 nutrients, in root growth and maintenance, and are important drivers of tolerance to edaphic stress
67 (Chimungu et al., 2015a; Lynch, 2015; Postma and Lynch, 2011a; Saengwilai et al., 2014; Zhu et
68 al., 2010a). The carbon cost of soil exploration includes carbon expenditure in root tissue
69 construction and maintenance and ion uptake and assimilation (Nielsen et al., 1994, 2001). Of these,
70 maintenance respiration is the largest carbon cost over time. Root metabolic costs at low nutrient
71 availability are significantly greater than rates at high nutrient availability (Lambers et al., 2008;
72 Nielsen et al., 2001). Root respiration is also a major cause of growth reduction under nutrient stress
73 (Postma and Lynch, 2011), the consumption of carbon by root respiration can exceed 50% of daily
74 photosynthesis under suboptimal nutrient levels (Ho et al., 2005; Lambers and Oliveira, 2020;
75 Nielsen et al., 2001). Therefore, phenes, i.e., the basic unit of the phenotype, and phene states, i.e.,
76 the status of specific phenes (Lynch, 2011; Pieruschka and Poorter, 2012; York et al., 2013), that
77 reduce maintenance respiration allow more internal resources to be allocated to better root
78 establishment, thus improving crop growth under limited nutrient availability, and therefore present
79 opportunities for the development of crops with reduced nutrient requirements (Lynch, 2015).

80 The “Steep, Cheap and Deep” (SCD) ideotype proposes maize root phenotypes to optimize
81 water and N capture under limited availability of those resources (Lynch, 2013). This ideotype
82 consists of root anatomical, architectural and physiological phenes that increase root depth, and
83 improve the acquisition of resources from deep soil domains. By determining the proportion of
84 respiring to non-respiring root tissue affecting the carbon and nutrient cost of tissue construction
85 and maintenance, root anatomy regulates the metabolic cost of soil exploration and therefore is an
86 important factor in the effects of edaphic stress on root and whole plant development (Fan et al.,
87 2003; Jaramillo et al., 2013; Mano et al., 2006). The “topsoil foraging” ideotype for P capture (Ho
88 et al., 2004; Lynch, 2011; Lynch and Brown, 2001; Richardson et al., 2011; Wang et al., 2010,
89 Lynch, 2019) has been useful as a breeding goal in developing soybean and common bean cultivars
90 that can enhance P acquisition in low phosphorus and drought environments (Burridge et al., 2019),
91 with similar application for enhanced P acquisition in maize (Zhu et al., 2005), given that P is
92 immobile in the soil strata, and is concentrated in the topsoil. Phene states that create a greater root
93 surface area in the topsoil, such as shallow root angle (Lynch and Brown, 2001; Rubio et al., 2003;
94 Zhu et al., 2005; Rangarajan et al., 2018) many hypocotyl-borne roots (Miller et al., 2003;
95 Walk et al., 2006; Rangarajan et al., 2018), dense lateral branching (Zhu and Lynch, 2004;
96 Jia et al., 2018), greater production of axial roots (Miguel et al., 2015, Walk et al., 2006;
97 Rangarajan et al., 2018), RCA formation (Postma and Lynch, 2011a,b) and root hair formation
98 (Zhu et al., 2010b; Miguel et al., 2015), have greater capacity of intercepting P, thus enhancing P
99 uptake in the topsoil. This strategy may also be relevant to improving K acquisition in the topsoil
100 under low K availability, as K is also relatively immobile (Lynch, 2019). Anatomical phene states

that contribute to reduced metabolic cost were also found to exhibit synergism with architectural phenes (Postma and Lynch, 2011a).

The formation of root cortical aerenchyma (RCA), the enlarged intercellular spaces that form through either programmed cell death or cell separation (Evans, 2003), is generally increased in response to hypoxia (Jackson and Armstrong, 1999) and various edaphic stresses, including suboptimal availability of phosphorus, nitrogen, sulfur, and water (Bouranis et al., 2003; Drew et al., 1989; Fan et al., 2003; Konings and Verschuren, 1980; Zhu et al., 2010a; Saengwilai et al., 2014; Chimungu et al., 2015; Galindo-Castañeda et al., 2019). RCA formation alleviates the limitation of hypoxia for root respiration with improved oxygen transport (Jackson and Armstrong, 1999). The utility of RCA formation to maintain greater growth rates under various soil nutrient and drought stresses by remobilizing nutrients from the root cortex and reducing maintenance respiration has been demonstrated in several previous studies (Chimungu et al., 2015a; Fan et al., 2003; Galindo-Castañeda et al., 2019; Jaramillo et al., 2013; Postma and Lynch, 2011a; Saengwilai et al., 2014; Zhu et al., 2010a). However, the dynamic interaction between RCA and other anatomical phenes and their effects on growth requires further examination, as RCA formation reduces the proportion of root volume occupied by living cortical tissue, which is more metabolically demanding than stelar tissue (Lynch, 2013). Chimungu et al. (2014a, b) reported that reduction in the number of concentric layers of parenchyma cells in the cortex of the maize root, or cortical cell file number (CCFN), and increased volume of individual cortical parenchyma cells, or cortical cell size (CCS), could decrease the metabolic costs of root growth and maintenance, in terms of both the carbon cost of root respiration and the nutrient content of cortical tissue. In contrasting maize lines exposed to water deficit stress in controlled environments and the field, larger CCS and reduced CCFN were associated with reduced root respiration, deeper rooting, greater water capture, improved plant water status, and hence greater growth and yield (Chimungu et al., 2014a, b). However, the physiological utilities of larger cortical cells, reduced CCFN, and their interaction with RCA and root architectural phenes under nutrient deficiencies, are not known.

The utility of a root phene state under stress may be dependent on its interactions with other architectural and anatomical phenes. Phene synergism refers to the phenomenon where the combined effect of two or more phenes is greater than the additive sum of their individual effects. For example, in low P soils, common bean genotypes with long root hairs (RHL) and shallow basal root growth angle (BRGA) had three-fold greater biomass accumulation than genotypes with short root hairs and steep root angle, while only 89% greater biomass was contributed by RHL alone, and 58% by shallow BRGA alone (Miguel et al., 2015). In another study, RCA formation in lateral roots in genotypes with increased lateral root branching density had greater benefits for phosphorus acquisition (Postma and Lynch, 2011b) than the effect of RCA alone. Integration of anatomical phenes and architectural phenes of maize root systems are important for plant growth and nitrogen acquisition (York et al., 2013; York and Lynch, 2015). These potential synergisms may be useful for breeding crops with greater edaphic stress tolerance. However, interactions among phenes may also be antagonistic, i.e., the functional response of phene states in combination is worse than that expected from the sum of their responses in isolation. For example, at low soil N levels, a phenotype with increased LRBD combined with RCA formation caused 42% reduction in shoot dry weight, compared to the expected additive effects of this phenotype, which indicates a functional antagonism (Postma and Lynch, 2011b; York et al., 2013).

A quantitative understanding of the functional dependence of one phene on the expression of other phenes and interactions among phenes and environmental factors is important for probing phenotypic diversity and breeding utility. We hypothesize that larger CCS and reduced CCFN, in combination with RCA formation, would decrease root respiration and tissue nutrient content, which would result in greater root growth, more efficient acquisition of soil N, P and K, and better

149 root and whole plant establishment under suboptimal N, P, and K availability. We also hypothesize
150 that the combined benefit of RCA, CCS, and CCFN is additive and is greater than the benefit of
151 RCA alone. *OpenSimRoot*, a functional-structural plant model, was used to evaluate: (1) the utility
152 of CCS, CCFN, and RCA under suboptimal N, P, and K availability, (2) potential synergism
153 between CCS, CCFN, and LRBD, and (3) the benefit of CCS and CCFN under conditions of greater
154 carbon availability as may occur with elevated atmospheric CO₂ concentration.

155 Results

156 When maize was grown under N stress in solution culture, IBM201 (genotype with reduced
157 CCFN) showed lower N concentration in the roots, IBM30 (genotype with larger CCS) showed
158 lower N concentration only in stems. Under sufficient P availability, both IBM201 and IBM30
159 showed reduced P concentration in the roots. In addition, IBM201 showed lower P concentration
160 in leaves. Under K stress, IBM201 showed reduced K concentration in root tissues but also
161 increased K concentration in the stems (Fig. 1).

162 In the simulations, increased RCA, reduced CCFN, and larger CCS had positive effects on
163 plant growth under limiting soil nitrogen, potassium and phosphorus as simulated independently
164 (Fig. 2, Fig. 3, Fig. 4). Plants with larger CCS and reduced CCFN had greater rooting depth and
165 greater uptake rate of nitrate at deeper soil strata as well (Fig. 2). Improved plant growth by larger
166 and reduced CCFN was highly dependent on the intensity of nutrient stress and the specific nutrient
167 simulated. Generally, at intermediate deficiency (i.e., plant dry weight at 30% - 50% of an
168 unstressed reference), RCA formation, larger CCS and reduced CCFN exhibited the greatest
169 beneficial effect when potassium was the limiting resource, while at severe deficiency (i.e., plant
170 dry weight at 1% - 25% of an unstressed reference), the greatest beneficial effect was found when
171 nitrogen and phosphorus were the limiting resources. Under N and P stress the utility of all three
172 phenotypes generally decreased with increasing nutrient availability, while under potassium stress
173 they benefited the plant the most at intermediate deficiency. Reduced nutrient content in root tissue
174 contributed more towards the total improvement in growth of phenotypes with larger CCS and
175 reduced CCFN than did reduced respiration. The total benefits of larger CCS or reduced CCFN
176 were greater than summing respective benefits introduced by reduced nutrient content and reduced
177 respiration (Fig. 3), and under extreme P and K stresses, were greater than the total benefit of RCA.
178 For example, under extremely limiting nutrient levels (21 kg/ha N, 0.05 kg/ha P), large CCS
179 increased biomass 47% (N stress) and 56% (P stress), while the reduction in respiration contributed
180 only 16% (N stress) and 14% (P stress), and reduced nutrient content only 18% (N stress) and 23%
181 (P stress). Under moderate K stress (1.9 kg/ha), reduction in respiration caused by larger CCS
182 contributed 30% and reduced nutrient content 18% towards growth benefits, while enabling both
183 functions resulted in a 69% growth enhancements, 21% higher than the additive terms of the two
184 functions, indicating synergism.

185 Predicted benefits of reducing respiration and nutrient concentration would increase in
186 general as cell size increases and file number decreases within the range of observed variation (Fig.
187 5). At extremely low soil nutrient availabilities (10% of sufficient soil nitrate and potassium
188 availabilities, 1% of sufficient soil phosphorus availability), larger CCS and reduced CCFN did not
189 achieve the most substantial enhancement of plant growth, which were found at moderately low
190 soil nutrient availability (20% of sufficient soil nitrate availability, 8% of sufficient soil phosphorus
191 availability and 25% of sufficient soil potassium availability).

192 We simulated the timing and development of nitrogen and phosphorus stress in plants with
193 only RCA formation, with both RCA formation and larger CCS, or with both RCA formation and
194 reduced CCFN independently. Plants with either larger CCS or reduced CCFN present along with

195 RCA formation were slightly less nitrogen and phosphorus stressed in that nutrient stress was
196 delayed approximately 1-3 additional days by both phenotypes (Fig. 6). With decreases in nutrient
197 availability, stresses developed earlier and were more severe in the phenotypes without RCA
198 formation, or larger CCS, or reduced CCFN than the ones with these phenotypes. RCA formation,
199 CCS and CCFN could alleviate nitrogen or phosphorus stress in terms of both duration and severity
200 of stress.

201 We used a high RCA phenotype with large CCS and reduced CCFN to simulate the
202 beneficial effects of large CCS and reduced CCFN before and after RCA formation, and their
203 interactions after RCA formation under nitrogen, phosphorus and potassium deficiency (Fig. 7).
204 Both CCS and CCFN had initial benefits at the very beginning of growth, and both continuously
205 increased dry weight until RCA formation, which replaced root cortical cells and cell files with
206 large intercellular spaces. Over time, by reducing nutrient content in the root, larger CCS and
207 reduced CCFN contributed more to growth enhancement than that of respiration reduction under
208 N or P stress, while respiration reduction contributed more under K stress. RCA formation was at
209 a minimal level initially, but increased substantially under suboptimal levels of all three nutrients
210 at 15 DAG, which corresponded to the time nutrient stress was perceived due to exhausted seed
211 reserves. After the substantial increase in RCA formation, RCA was responsible for the majority
212 of benefits under nutrient stress. We also tested potential additive effects of RCA formation, large
213 CCS and reduced CCFN, since the majority of benefits of RCA do not overlap with those of large
214 CCS and reduced CCFN over time during growth. Benefits of both large CCS and reduced CCFN
215 after RCA formation at 15 DAG were reduced. After 15 DAG, the majority of benefits were
216 contributed by RCA. In this case, the combination of all three phenotypes at their most carbon-
217 efficient level, i.e., the observed level that showed greatest reduction in the carbon cost of root
218 maintenance, achieved growth benefits up to 135%, 132% or 161% under low nitrate, phosphorus
219 or potassium availabilities.

220 We simulated the utility of CCS and CCFN under nitrogen and phosphorus deficiency with
221 varied lateral root branching density (LRBD) in phenotypes with either the largest root cortical
222 cells, or fewest root cortical cell files. Both large CCS and reduced CCFN achieved greater growth
223 enhancements in phenotypes with half the reference LRBD under low soil nitrate availability. The
224 contrary was evident for phenotypes grown under both low and medium soil phosphorus
225 availability, where greater benefits were observed when plants with doubled LRBD compared to
226 the reference phenotype. In soils with intermediate nitrate availability, phenotypes with normal
227 LRBD had the greatest benefit compared to half or doubled LRBD (Fig. 8).

228 With greater carbon availability, simulated by increasing light utilization efficiency in the
229 canopy module, the benefits of large CCS under low soil nitrogen or phosphorus
230 availability declined (Fig. 9).

231 Discussion

232 Our results align with previous findings that RCA formation, which reduces the metabolic
233 cost of soil exploration in terms of nutrient and C investment, improves plant growth under
234 conditions of suboptimal availability of N, P and K (Postma and Lynch, 2011; Saengwilai et al.,
235 2014; Galindo-Castañeda et al., 2019), and support the hypothesis that larger CCS and reduced
236 CCFN increase soil nutrient acquisition by reducing root metabolic costs. The combined benefits
237 of RCA formation, larger CCS and reduced CCFN for growth are greater than the benefit of RCA
238 formation alone. Larger CCS and reduced CCFN reduce the metabolic cost of soil exploration
239 under drought stress (Chimungu et al., 2014a, b), and are predicted by our results to alleviate
240 nitrogen, phosphorus and potassium stress as well. No literature has reported CCS or CCFN

alleviating potassium stress. However, given that both larger CCS and reduced CCFN reduce root respiration (Chimungu et al., 2014a, b), we believe that they may have utility under potassium stress. These results indicate that all three phenes may have substantial utility on infertile soils, suggesting that cultivars with high RCA formation, large cortical cells and reduced cortical cell files would have reduced fertilizer requirements in intensive agriculture and would yield better in low-input systems. Our results focus on maize but we propose that they should be generally applicable to other grass species, which like most monocots lack secondary growth and so have a persistent cortex.

Typically, multiple edaphic stresses occur simultaneously (St. Clair and Lynch, 2010), although it is difficult to reflect concurrent stresses in the model because of potential interactions among plant stress responses (Dathe et al., 2013). In such environments, tradeoffs between nutrient acquisition strategies for specific nutrients, and between other plant physiological functions, are challenging (Hu et al., 2014; Lynch and St. Clair, 2004; Rubio et al., 2003). For example, the strategy of enhancing topsoil foraging proved to be critical for adaptations to low phosphorus soil in common bean (Lynch and Brown, 2001) and maize (Zhu et al., 2005); the “Steep, Cheap and Deep” ideotype proposes that root phenotypes capable of rapid exploration of deep soil strata would optimize soil nitrate and water capture in maize (Lynch, 2013). However, given the limited amount of carbon and nutrients available for root maintenance, extreme inclinations towards one strategy may be detrimental for the other, and optimization of resource allocation between deep soil exploration and topsoil exploration is critical. For example, the optimal lateral root branching density in maize is dependent upon nitrate and phosphorus availability (Postma et al., 2014). Sparse but long LRBD is optimal for nitrate uptake, while dense but short LRBD is optimal for phosphorus uptake. In another study (Dathe et al., 2016), axial root growth angle exhibit significant effects on nitrogen acquisition in maize, where extreme phenotypes have narrow intervals of optimal performances – extremely shallow root systems only increase N acquisition under reduced precipitation, while dimorphic phenotypes that combined shallow seminal roots with deep crown roots performed well in all environments. High LRBD introduces strong competition among roots for nitrate capture, therefore decreasing nitrate uptake since the carbon budget of the whole plant does not grant greater root length. Carbon budget and root competition does not impact phosphorus uptake as significantly as nitrate uptake, therefore increasing root length by increasing LRBD is optimal for phosphorus uptake. In reality, most genotypes have a balanced LRBD to meet the demand of both nitrate and phosphorus acquisition. Maize genotypes with high RCA formation could also inhibit radial phosphorus transport due to the reduction of living tissue (Hu et al., 2014).

The utilities of RCA, larger CCS and reduced CCFN were greater in plants that were experiencing moderate potassium stress (25% of potential growth) than in plants under severe potassium stress (less than 10% of potential growth). Similarly, RCA, larger CCS and reduced CCFN did not achieve optimal growth enhancement under extremely low soil nitrate and phosphorus availability (1% to 8% of potential growth under N stress, 1% to 5% of potential growth under P stress), but showed greater benefits under less severe stress. This decline is caused by reduction in the utility of the respiration reduction function. With extremely limited nutrient availability, root respiration per ion absorbed increases, inhibiting root and shoot growth to compensate for greater respiration. The reduction in respiration is more important in potassium deficient plants than in nitrogen deficient or phosphorus deficient plants, as carbon is relatively more limiting in potassium stressed plants, which differs from the cases of nitrogen and phosphorus (Postma and Lynch, 2011a). Substantial reductions in photosynthetic assimilation caused by nutrient deficiency impose carbon limitations under N, P or K stress. However, in potassium stressed plants, an adaptive response in carbon partitioning between roots and shoots is absent,

288 which is present in nitrogen or phosphorus stressed plants, that can allocate more carbon to root
289 growth.

290 The model predicts a decline in the benefit of larger CCS and reduced CCFN at extremely
291 low nutrient availability. If we consider the amount of nutrient required for the construction of a
292 root segment as the cost, then the cost is greater in low nutrient soils than in fertile soils (Postma
293 and Lynch, 2011b). Nutrient uptake under severe deficiency could be limited by both carbon and
294 the deficient nutrient, while the cost of root tissue construction and respiration remain high, causing
295 an increase in the cost-benefit ratio for nutrient uptake (Nielsen et al., 2001). In *OpenSimRoot*, the
296 benefit of larger CCS and reduced CCFN is dependent on the cost – benefit ratio of root segments.
297 Under extremely scarce nutrient availability, the decline in benefits is caused by the high cost –
298 benefit ratio of root growth. Therefore, if nutrient availabilities fall below a threshold where the
299 stressed plant does not have sufficient nutrient stores for tissue construction, causing extra
300 allocation of carbon to the root system, thus deteriorating the photosynthetic and nutritional status
301 of the plant, then the utility of RCA formation, CCS, CCFN, or other phenes that reduce metabolic
302 costs, would decrease. In an extreme theoretical environment where all nutrients in soil are depleted,
303 the utility of these phenes becomes nil.

304 While the potential for RCA formation is genetically controlled, RCA formation is highly
305 responsive to edaphic stress (Fan et al., 2003). Substantial variation in CCS and CCFN, however,
306 were observed among RILs in empirical studies, but were not as plastic to edaphic stress as RCA
307 formation (Chimungu et al., 2014a, b), and could be beneficial starting at the very beginning of
308 growth under edaphic stress. In our simulation, both large CCS and reduced CCFN exhibited
309 benefits before the formation of RCA in response to nutrient limitation. As cortical cells and cell
310 files were replaced by RCA formation, we expect the majority of benefits from these three phenes
311 do not overlap over time. Therefore, we predict a simplification of the additive effects among the
312 three phenes to be present under N, P or K stress, and observed increased benefits due to the
313 combination of all three phene states compared to that of RCA formation alone. In reality, however,
314 we expect a more complicated interaction between RCA formation, CCS and CCFN. RCA
315 formation, as a response to nutrient stress, manifested 12-13 days after germination (Postma and
316 Lynch, 2011). RCA formation in mid-root and apical regions were less than that of basal regions
317 of a root (Fan et al., 2003). The variation in the spatial and temporal distribution of RCA at both
318 the single root scale and the root system scale is dynamic (Burton et al., 2013), and such variation
319 can cause changes in nitrogen uptake kinetics (York et al., 2016). Therefore, in reality, the
320 interaction between RCA formation, CCS and CCFN could not be represented by a simplified
321 expectation of additive effects.

322 Plants have developed multiple alternate strategies to increase nutrient uptake under severe
323 nutrient stress, such as root hair formation, root exudation, and mycorrhizal colonization. Root hair
324 formation has a relatively low cost – benefit ratio, but can increase phosphorus uptake significantly
325 (Bates and Lynch, 2001; Nielsen et al., 1994, 2001; Zhu and Lynch, 2004; Miguel et al., 2015).
326 Mycorrhizal colonization increased phosphorus efficiency significantly at low phosphorus
327 availability compared to non-colonized plants (Ning and Cumming, 2001), but did not increase
328 plant dry weight significantly due to the increased maintenance and growth respiration of the fungal
329 tissue (Nielsen et al., 2001), indicating that root carbon costs are a major limitation to plant growth
330 under phosphorus stress. Additionally, when the carbon cost of root growth is removed, simulated
331 plant growth increased under P stress (Postma and Lynch, 2011). Our results support the general
332 hypothesis that the metabolic costs of soil exploration in terms of the carbon and nutrient
333 investment in root tissue over time becomes increasingly important as the availability of critical
334 soil resources declines.

Chimungu (2014) reported, in both drought-stressed and non-stressed plants, significantly thicker roots when larger cortical cells and more cortical cell files were present. In addition, Burton (2010) observed greater RCA formation in thicker root classes in non-stressed plants. The ability of roots to penetrate compacted soil and root depth are correlated to both root anatomical phenes such as RCA, CCS and CCFN, as well as root diameter, where deeper-rooting plants in compacted soil showed reduced CCFN and increased RCA formation. Additionally, root thickening in the form of root cortical area expansion were closely related to soil mechanical impedance in some genotypes (Chimungu et al., 2015b; Vanhees et al., 2020). Smaller outer band cortical cells could reduce the risk of root collapse when encountering increased mechanical impedance. RCA formation is also negatively correlated with root bending strength, while smaller distal root cortical cells, more cortical cells, and more CCFN increase the strength of root and reduce root collapsing during penetration of soil (Whiteley et al., 1982; Clark et al., 2003; Jin et al., 2013;) since they contribute to the construction of a thicker root. Root cortical cell diameter is a pivotal and heritable trait in determining the carbon cost of penetrating compacted soil, where large cell diameter correlated with reduced carbon cost of root growth, especially under high soil mechanical impedance; the plasticity of this trait allowed the enlargement of root cortical cells was a common response when roots encounter compacted soil (Colombi et al., 2019). These observations from empirical studies suggest that, although RCA formation, larger CCS and reduced CCFN reduce the metabolic cost of soil exploration, they may affect root penetration of hard soil domains, which could affect their ability to acquire resources located in deeper soil profiles.

An important merit of simulation modeling is the ability to test hypotheses and probe scenarios that are inaccessible to empirical studies. Simulation modeling makes it possible to isolate and test the objects of study, in this case, larger CCS and reduced CCFN, from interactions with many other biotic and abiotic factors, which is difficult to avoid in empirical studies (Dunbabin et al., 2013; Postma et al., 2014). It would be infeasible to test separately how larger CCS and reduced CCFN reduce respiration and nutrient content in an empirical study, while in modeling, different functions of specific phenes could be isolated and examined without being confounded by other functions, which was critical to several previous studies (Postma and Lynch, 2011a; 2011b). In other cases, modeling also allows us to study phenes that are otherwise difficult to manipulate in real plants, such as changing nutrient uptake kinetics (York et al., 2016), or examining root competition in time and space in the ‘three sisters’ polyculture (Postma et al., 2014) where empirical measurement is impractical. *In silico* approach allows the flexibility of conducting thousands of simulations in factorial designs (in this study, over 3,300 runs) which would be difficult to conduct empirically. Although such models are designed with assumptions and simplifications of the actual scenarios or mechanisms they simulate, and often (as in the present case) rely on empirical data as input parameters, it does not nullify the value of models as a useful research tool to provide a preliminary insight into root anatomy, architecture, physiological processes and interactions with other factors of interest, and a compliment to field studies even when empirical data are present. In our case, *OpenSimRoot* is capable of simulating a comprehensive range of phenes and phenotypes in specific environments that can be customized. Because of its heuristic nature, *OpenSimRoot* focuses on the validity of simulating physiological processes, rather than the alignment with empirical studies that predictive models emphasize.

Our results suggest potential areas where structural-functional plant models could be improved. The duration of simulation could be parameterized for longer periods, which could potentially enable models to simulate the full life cycle of plants, and demonstrate the dynamics of physiological processes. Interactions and dynamics among root phene states deserve more attention. Some other parameters, such as soil hardness, microbial associations, interplant competition (Postma and Lynch, 2011), and interspecific interactions in cropping systems (Postma and Lynch,

2012) are important for understanding ecosystem functioning on a greater scale, and may have consequences for the utility of root anatomical phenes such as RCA, CCS and CCFN. The phene aggregate of reduced CCFN and larger CCS, along with RCA formation, is defined as reduced living cortical area (LCA). Plants with reduced LCA had decreased root segment respiration, reduced P concentration in root tissues, and greater rooting depth, indicating lower carbon cost of root growth, which resulted in increased biomass and resource capture under P stress (Chimungu et al., 2014b; Galindo-Castañeda et al., 2019).

390 **Conclusions**

Quantitative evidence that larger CCS and reduced CCFN are adaptive phene states for multiple nutrient stresses are presented. The utilities of larger CCS and reduced CCFN in soils with suboptimal nitrogen, phosphorus and potassium availability are dependent on nutrient availability, phene functions, and interactions among phenes. We propose that larger CCS and reduced CCFN are complementary to RCA formation in terms of growth enhancements as the majority of benefits of these three phene states do not overlap in time. We expect tradeoffs for RCA formation, larger CCS and reduced CCFN to be present, as all three phene states are related to soil penetration and root proliferation. This aspect merits further investigation. Functional-structural plant models like *OpenSimRoot* can be used to simulate variations in these anatomical phenes and thereby evaluate their utilities under multiple edaphic stresses, and have the potential to provide a holistic understanding of the roles of root phenotypes for plant fitness. These results indicate that large CCS and reduced CCFN merit investigation as breeding targets for maize and possibly other cereal crops, since the development of crop cultivars with improved soil resource acquisition remains a critical strategy for improving the sustainability of intensive agriculture and for improving the productivity of low-input agroecosystems.

406 **Materials and Methods**

We used *OpenSimRoot* (Lynch et al., 1997), a functional-structural plant model with focus on root architecture and soil resource acquisition, to simulate the formation of RCA, variation in CCS and CCFN, and their physiological utility in maize growing with varied nitrate, phosphorus, or potassium availability in the soil. We also evaluated potential additive effects among RCA, CCS, and CCFN. In addition, we conducted a pair of two factor solution culture experiments to examine the variation in tissue N, P and K concentration in genotypes with contrast in CCS or CCFN.

413 *Solution culture study*

Four maize genotypes (IBM population IBM178, IBM201, IBM365, IBM30) were used in the solution culture study. Genotypes were selected for contrasting CCS (IBM365 and IBM30), and CCFN (IBM178 and IBM201) based on preliminary screening.

Genotypes were planted in four replications in total under both high and low N or K availabilities in solution culture, with sufficient P availability across all treatments in a greenhouse at the Penn State University campus located at University Park, PA, USA (40.8148° N, 77.8653° W), with two replications planted on June 4th, and two more replications planted on June 13th, 2017. Eight 100-liter non-transparent plexi glass solution culture tanks were used, within each tank, two replications of the four genotypes were planted. Nutrient solution was based on and modified upon the Hoagland solution (Johnson et al., 1957). N concentration in N stress treatments was 160 μmol/L, and K concentration in K stress treatments was 60 μmol/L, P concentration was 2 mmol/L.

Plants were grown for 24 days after transplanting, or 32 days after germination in growth chamber to avoid RCA formation confounding the effect of larger CCS and reduced CCFN. Upon harvest, 10 cm long root segments from base and tip of the second and the third whorl nodal roots were collected to conduct anatomical analysis and respiration measurements. The rest of the root system, along with leaves and stems, were separated and dried in oven at 60°C to measure root and shoot dry weight. Tissues were then ground and sent to the Agricultural Analytical Services Laboratory at the Penn State University for P and K content analysis. N content analysis was conducted with a 2400 CHNS/O Series II element analyzer (PerkinElmer). Nutrient content data was used to parameterize *OpenSimRoot* to include how tissue nutrient content was reduced by larger CCS and reduced CCFN in the simulations.

Model description

OpenSimRoot simulates the three-dimensional root architecture and soil resource acquisition of a root system over time. The root system is described as distinct root classes represented by a growing number of root nodes and segments as the root system develops (Lynch et al. 1997). Root growth is based on a carbon source-sink model, where the carbon partition protocol has been described by Postma and Lynch (2011a). Shoot growth and photosynthesis is simulated using LINTUL (Spitters and Schapendonk, 1990). Nutrient uptake is simulated for each root segment in comparison with the optimal (o) and minimal (m) nutrient requirements of the plant. Nutrient deficiency, or stress factor, is defined as when nutrient uptake falls below the optimal nutrient requirement. The stress factor influences shoot development and photosynthetic efficiency depending on the nutrient simulated. We used the Barber-Cushman model (Itoh and Barber, 1983; Postma and Lynch, 2011a) to simulate phosphorus uptake, and linked *OpenSimRoot* to the three-dimensional hydrological model SWMS3D (Simunek et al., 1995) to simulate nitrate and potassium uptake. The Barber-Cushman model is considered to be inadequate for nitrate and potassium uptake as these nutrients are relatively mobile (Postma and Lynch, 2011b), and the Barber-Cushman model does not simulate leaching and ignores root competition in three dimensions. The SWMS3D model is not ideal for simulating the phosphorus depletion zones at root surface (Postma and Lynch, 2011b), as computational demands required by the resulting substantial number of finite element (FEM) nodes (Hardelauf et al., 2007) are considerable, and the phosphorus depletion zone would be artificially enlarged in the SWMS3D model.

Variation in RCA formation, CCS and CCFN in maize are simulated for each root segment with empirical parameters retrieved from Burton (2010), and Chimungu (2014a, b). The percentage RCA for different root classes is well described by Fan et al. (2003). We simplified CCS and CCFN simulation by assuming they are uniformly distributed across root classes. The addition of CCS and CCFN were implemented as new model input files, no specific modification were made to the computational codes of OSR to accommodate this addition. RCA formation is allowed to combine nutrient remobilization and respiration reduction and is based on regression between the amount of RCA and nutrient content and root respiration of empirical measurements by Fan et al. (2003). *OpenSimRoot* does not explicitly represent root anatomy, so CCS and CCFN variation is represented by reducing modeled root respiration and tissue nutrient content.

Effects of nutrient stress on growth

In *OpenSimRoot*, the nutrient stress factor module is implemented to affect the potential leaf area expansion rate and light use efficiency (LUE) independently as in the LINTUL model. The nutrient stress factor functions as a growth regulator between root and shoot growth. The nutrient stress factor negatively impacts light use efficiency and resulted in reduced carbon available for plant growth. Reduction in the potential leaf area expansion rate caused by the stress factor resulted in reduced sink strength of the shoot, and consequently greater allocation of carbon

to root growth. Nutrient-specific stress response was used to determine the effect of internal nutrient concentrations (nitrogen, potassium and phosphate) on the two parameters. In this study, potassium stress strongly reduces LUE (Zhao et al., 2001) but does not affect the potential leaf area expansion rate (Cakmak et al., 1994). Suboptimal phosphate strongly reduces the potential leaf area expansion rate but is trivial in affecting LUE (Lynch et al., 1991). Inorganic nitrogen strongly affects both parameters (Uhart and Andrade, 1995).

Distribution of RCA formation, CCS and CCFN within the root system

We assumed that RCA formation starts behind the elongation zone of a root and develops over time until reaching a maximum. Therefore, the greatest amount of RCA formation can be found close to the base of a root, which aligns with Fan et al. (2003) but disagrees with Bouranis et al. (2006), Lenochova et al. (2009), and Burton (2010). RCA formation is reduced in the first 5 cm of the root (Bouranis et al., 2006), which is a small part of the total root length and we expect the effect on total RCA formation to be small. We used the maximum amount of RCA formation in the literature, which is 39% of root cross-section area at 20 days after germination (Fan et al., 2003) in the model.

Variation in CCS and CCFN was observed in the mid cortical band of roots by Chimungu et al. (2014a, b). In reality, the spatial distribution of CCS and CCFN are not uniform in either the area cross-sectioned, or across different root classes. To demonstrate the effect of observed respiration reduction of these phenes, we assumed that CCS and CCFN variation are uniform regardless of root class and location in the area cross-sectioned. Parameterization of these phenes are based on the genotypic variation described by Chimungu et al. (2014a, b). CCS varies between 101 μm^2 and 533 μm^2 , and CCFN varies between 8 and 17 in maize.

Interactions between RCA formation, LRBD, CCS and CCFN

Living cortical area (LCA; Jaramillo et al., 2013) is proposed as a good predictor of root respiration, and a critical determinant of root metabolic cost, which involves the phenes in this study. Interactions between LCA components requires further demonstration. We simulated the extremes of variation in RCA formation, where RCA takes up between 0% to 39% of the root cross sectional area, CCS and CCFN to probe additive effects under low nitrogen and phosphorus availability. Significant genetic variation exists in lateral root branching density (LRBD) (Trachsel et al., 2011). We varied the LRBD parameter to the extremes reported in this study, between 4 to 16 lateral roots/cm on axial roots, to examine if potential synergism between LRBD, CCS and CCFN under low soil nitrogen and phosphorus availability to further test the utility of CCS and CCFN in an integrated genotype.

System description, parameterization, and runs

We simulated growth of 40 days after germination of a single maize plant, which represents a uniform monoculture plant community with a between-row spacing of 60 cm and a within row spacing of 26 cm. Aboveground competition was simulated by a shading function (Postma and Lynch, 2011). Parameterization was based on input parameters used in previous simulation studies with *OpenSimRoot* (Postma and Lynch, 2011a; 2011b). From empirical measurements from Chimungu et al. (2014a, b), we parameterized how larger CCS and reduced CCFN reduced root respiration. We parameterized how tissue nutrient content varied between contrasting phenotypes by conducting a solution culture study (see above). All simulations were performed on the Penn State supercomputing clusters aci-b, with the following variables: (1) CCS and CCFN; (2) the functions of RCA formation; (3) lateral root branching density with CCS and CCFN held constant; (4) atmospheric CO₂ pressure between ambient values of 400ppm, and up to four-fold (1600 ppm); (5) the availability of nitrate, phosphorus and potassium in the soil, from low to sufficient; and (6)

518 “max and min RCA”, “max and min CCS” and “max and min CCFN” reference genotypes. To
 519 account for stochasticity in growth rates and root branching frequencies, each scenario was
 520 simulated with four replications each with *OpenSimRoot*’s random number generator initialized to
 521 different values, with the graph showing the mean value. The variation of phenes in this study were
 522 based on empirical studies to avoid extrapolation towards unrealistic conditions. Appendix A
 523 contains a summary of the model parameterizations.

524 *Statistical analysis*

525 Empirical data from the solution culture study were analyzed by paired Student’s *t* tests in
 526 R 3.4.1 (R Core Team, 2017). We did not conduct significance test on the simulation results, as
 527 such tests were not reliable in simulation studies, as the ease of replication in computer simulations
 528 allows for any effect size to be found significant if there are enough replicates. Biological
 529 significance, rather than the statistical significance, should be the main focus of simulation
 530 experiments (White et al., 2014).

531

532

533 **Literature cited**

- 534
- 535 Bates, T. R., and Lynch, J. P. (2001). Root hairs confer a competitive advantage under low
- 536 phosphorus availability. *Plant and Soil*, 236(2), 243–250.
- 537 <https://doi.org/10.1023/A:1012791706800>
- 538 Bouranis, D. L., Chorianopoulou, S. N., Siyiannis, V. F., Protonotarios, V. E., and Hawkesford,
- 539 M. J. (2003). Aerenchyma formation in roots of maize during sulphate starvation. *Planta*,
- 540 217(3), 382–391. <https://doi.org/10.1007/s00425-003-1007-6>
- 541 Bouranis, Dimitris L., Chorianopoulou, S. N., Kollias, C., Maniou, P., Protonotarios, V. E.,
- 542 Siyiannis, V. F., and Hawkesford, M. J. (2006). Dynamics of Aerenchyma Distribution in
- 543 the Cortex of Sulfate-deprived Adventitious Roots of Maize. *Annals of Botany*, 97(5),
- 544 695–704. <https://doi.org/10.1093/aob/mcl024>
- 545 Burrridge, J. D., Findeis, J. L., Jochua, C. N., Miguel, M. A., Mubichi-Kut, F. M., Quinhentos, M.
- 546 L., Xerinda, S. A., and Lynch, J. P. (2019). A case study on the efficacy of root
- 547 phenotypic selection for edaphic stress tolerance in low-input agriculture: Common
- 548 bean breeding in Mozambique. *Field Crops Research*, 244, 107612.
- 549 <https://doi.org/10.1016/j.fcr.2019.107612>
- 550 Burton, A. (2010) Phenotypic evaluation and genetic basis of anatomical and architectural root
- 551 traits in the genus *Zea*. PhD Thesis, Pennsylvania State University, USA.
- 552 Burton, A. L., Brown, K. M., and Lynch, J. P. (2013). Phenotypic Diversity of Root Anatomical
- 553 and Architectural Traits in *Zea* Species. *Crop Science*, 53(3), 1042–1055.
- 554 <https://doi.org/10.2135/cropsci2012.07.0440>
- 555 Cakmak, I., Hengeler, C., and Marschner, H. (1994). Partitioning of shoot and root dry matter and
- 556 carbohydrates in bean plants suffering from phosphorus, potassium and magnesium

557 deficiency. *Journal of Experimental Botany*, 45(9), 1245–1250.
558 <https://doi.org/10.1093/jxb/45.9.1245>

559 Chimungu, J. G., Brown, K. M., and Lynch, J. P. (2014a). Large Root Cortical Cell Size
560 Improves Drought Tolerance in Maize. *Plant Physiology*, 166(4), 2166–2178.
561 <https://doi.org/10.1104/pp.114.250449>

562 Chimungu, J. G., Brown, K. M., and Lynch, J. P. (2014b). Reduced Root Cortical Cell File
563 Number Improves Drought Tolerance in Maize. *Plant Physiology*, 166(4), 1943–1955.
564 <https://doi.org/10.1104/pp.114.249037>

565 Chimungu, J. G., Loades, K. W., and Lynch, J. P. (2015). Root anatomical phenes predict root
566 penetration ability and biomechanical properties in maize (*Zea Mays*). *Journal of*
567 *Experimental Botany*, 66(11), 3151–3162. <https://doi.org/10.1093/jxb/erv121>

568 Chimungu, J. G., Maliro, M. F. A., Nalivata, P. C., Kanyama-Phiri, G., Brown, K. M., and Lynch,
569 J. P. (2015). Utility of root cortical aerenchyma under water limited conditions in tropical
570 maize (*Zea mays* L.). *Field Crops Research*, 171, 86–98.
571 <https://doi.org/10.1016/j.fcr.2014.10.009>

572 Colombi, T., Herrmann, A. M., Vallenback, P., and Keller, T. (2019). Cortical Cell Diameter Is
573 Key To Energy Costs of Root Growth in Wheat. *Plant Physiology*, 180(4), 2049–2060.
574 <https://doi.org/10.1104/pp.19.00262>

575 Clark, L. J., Whalley, W. R., and Barraclough, P. B. (2003). *How do roots penetrate strong soil?*
576 12.

577 Dathe, A., Postma, J. A., and Lynch, J. P. (2013). Modeling Resource Interactions Under
578 Multiple Edaphic Stresses. In D. Timlin and L. R. Ahuja (Eds.), *Advances in Agricultural*
579 *Systems Modeling* (pp. 273–294). American Society of Agronomy and Soil Science
580 Society of America. <https://doi.org/10.2134/advagricsystmodel4.c12>

581 Dathe, A., Postma, J. A., Postma-Blaauw, M. B., & Lynch, J. P. (2016). Impact of axial root
582 growth angles on nitrogen acquisition in maize depends on environmental conditions.
583 *Annals of Botany*, 118(3), 401–414. <https://doi.org/10.1093/aob/mcw112>

584 Drew, M. C., He, C.-J., and Morgan, P. W. (1989). Decreased Ethylene Biosynthesis, and
585 Induction of Aerenchyma, by Nitrogen- or Phosphate-Starvation in Adventitious Roots of
586 *Zea mays* L. *Plant Physiology*, 91(1), 266–271. <https://doi.org/10.1104/pp.91.1.266>

587 Dunbabin, V. M., Postma, J. A., Schnepf, A., Pagès, L., Javaux, M., Wu, L., Leitner, D., Chen, Y.
588 L., Rengel, Z., and Diggle, A. J. (2013). Modelling root–soil interactions using three–
589 dimensional models of root growth, architecture and function. *Plant and Soil*, 372(1–2),
590 93–124. <https://doi.org/10.1007/s11104-013-1769-y>

591 Evans, D. E. (2003). Aerenchyma formation: Tansley review. *New Phytologist*, 161(1), 35–49.
592 <https://doi.org/10.1046/j.1469-8137.2003.00907.x>

593 Fan, M., Zhu, J., Richards, C., Brown, K. M., and Lynch, J. P. (2003). Physiological roles for
594 aerenchyma in phosphorus-stressed roots. *Functional Plant Biology*, 30(5), 493.
595 <https://doi.org/10.1071/FP03046>

596 Galindo-Castañeda, T., Brown, K. M., Kulda, G. A., Roth, G. W., Wenner, N. G., Ray, S.,
597 Schneider, H., and Lynch, J. P. (2019). Root cortical anatomy is associated with
598 differential pathogenic and symbiotic fungal colonization in maize. *Plant, Cell and*
599 *Environment*, 42(11), 2999–3014. <https://doi.org/10.1111/pce.13615>

600 Hardelauf, H., Javaux, M., Herbst, M., Gottschalk, S., Kasteel, R., Vanderborght, J., and
601 Vereecken, H. (2007). PARSWMS: A Parallelized Model for Simulating Three-
602 Dimensional Water Flow and Solute Transport in Variably Saturated Soils. *Vadose Zone*
603 *Journal*, 6(2), 255–259. <https://doi.org/10.2136/vzj2006.0156>

604 Ho, M. D., McCannon, B. C., and Lynch, J. P. (2004). Optimization modeling of plant root
605 architecture for water and phosphorus acquisition. *Journal of Theoretical Biology*,
606 226(3), 331–340. <https://doi.org/10.1016/j.jtbi.2003.09.011>

607 Ho, M. D., Rosas, J. C., Brown, K. M., and Lynch, J. P. (2005). Root architectural tradeoffs for
608 water and phosphorus acquisition. *Functional Plant Biology*, 32(8), 737.
609 <https://doi.org/10.1071/FP05043>

610 Hu, B., Henry, A., Brown, K. M., and Lynch, J. P. (2014). Root cortical aerenchyma inhibits
611 radial nutrient transport in maize (*Zea mays*). *Annals of Botany*, 113(1), 181–189.
612 <https://doi.org/10.1093/aob/mct259>

613 Itoh, S., and Barber, S. A. (1983). A numerical solution of whole plant nutrient uptake for soil-
614 root systems with root hairs. *Plant and Soil*, 70(3), 403–413.
615 <https://doi.org/10.1007/BF02374895>

616 Jackson, M. B., and Armstrong, W. (1999). Formation of Aerenchyma and the Processes of Plant
617 Ventilation in Relation to Soil Flooding and Submergence. *Plant Biology*, 1(3), 274–287.
618 <https://doi.org/10.1111/j.1438-8677.1999.tb00253.x>

619 Jaramillo, R. E., Nord, E. A., Chimungu, J. G., Brown, K. M., and Lynch, J. P. (2013). Root
620 cortical burden influences drought tolerance in maize. *Annals of Botany*, 112(2), 429–
621 437. <https://doi.org/10.1093/aob/mct069>

622 Jia X, Liu P, Lynch JP. 2018. Greater lateral root branching density in maize (*Zea mays* L.)
623 improves phosphorus acquisition from low phosphorus soil. *Journal of Experimental*
624 *Botany* 69: 4961–4970.

625 Jin, K., Shen, J., Ashton, R. W., Dodd, I. C., Parry, M. A. J., and Whalley, W. R. (2013). How do
626 roots elongate in a structured soil? *Journal of Experimental Botany*, 64(15), 4761–4777.
627 <https://doi.org/10.1093/jxb/ert286>

628 Johnson, C. M., Stout, P. R., Broyer, T. C., and Carlton, A. B. (1957). Comparative chlorine
629 requirements of different plant species. *Plant and Soil*, 8(4), 337–353.
630 <https://doi.org/10.1007/BF01666323>

631 Koevoets, I. T., Venema, J. H., Elzenga, J. Theo. M., and Testerink, C. (2016). Roots
632 Withstanding their Environment: Exploiting Root System Architecture Responses to
633 Abiotic Stress to Improve Crop Tolerance. *Frontiers in Plant Science*, 07.
634 <https://doi.org/10.3389/fpls.2016.01335>

635 Konings, H., and Verschuren, G. (1980). Formation of aerenchyma in roots of *Zea mays* in
636 aerated solutions, and its relation to nutrient supply. *Physiologia Plantarum*, 49(3), 265–
637 270. <https://doi.org/10.1111/j.1399-3054.1980.tb02661.x>

638 Lambers, H., and Oliveira, R. S. (2020). *Plant Physiological Ecology*. Springer.
639 <https://public.ebookcentral.proquest.com/choice/publicfullrecord.aspx?p=5996299>

640 Lambers, H., Raven, J., Shaver, G., and Smith, S. (2008). Plant nutrient-acquisition strategies
641 change with soil age. *Trends in Ecology and Evolution*, 23(2), 95–103.
642 <https://doi.org/10.1016/j.tree.2007.10.008>

643 Lenochová, Z., Soukup, A., and Votrubová, O. (2009). Aerenchyma formation in maize roots.
644 *Biologia Plantarum*, 53(2), 263–270. <https://doi.org/10.1007/s10535-009-0049-4>

645 Liao, H., Rubio, G., Yan, X., Cao, A., Brown, K. M., and Lynch, J. P. (2002). Effect of
646 phosphorus availability on basal root shallowness in common bean. In D. S. Powlson, G.
647 L. Bateman, K. G. Davies, J. L. Gaunt, and P. R. Hirsch (Eds.), *Interactions in the Root*
648 *Environment: An Integrated Approach* (pp. 69–79). Springer Netherlands.
649 https://doi.org/10.1007/978-94-010-0566-1_7

650 Lynch, J., Läuchli, A., and Epstein, E. (1991). Vegetative Growth of the Common Bean in
651 Response to Phosphorus Nutrition. *Crop Science*, 31(2), 380–387.
652 <https://doi.org/10.2135/cropsci1991.0011183X003100020031x>

653 Lynch, J. P. (2011). Root Phenotypes for Enhanced Soil Exploration and Phosphorus Acquisition:
654 Tools for Future Crops. *Plant Physiology*, 156(3), 1041–1049.
655 <https://doi.org/10.1104/pp.111.175414>

656 Lynch, J. P. (2013). Steep, cheap and deep: An ideotype to optimize water and N acquisition by
657 maize root systems. *Annals of Botany*, 112(2), 347–357.
658 <https://doi.org/10.1093/aob/mcs293>

659 Lynch, J. P. (2015). Root phenotypes that reduce the metabolic costs of soil exploration:
660 Opportunities for 21st century agriculture: New roots for agriculture. *Plant, Cell and*
661 *Environment*, 38(9), 1775–1784. <https://doi.org/10.1111/pce.12451>

662 Lynch, J. P. (2019). Root phenotypes for improved nutrient capture: An underexploited
663 opportunity for global agriculture. *New Phytologist*, 223(2), 548–564.
664 <https://doi.org/10.1111/nph.15738>

665 Lynch, J. P., and Brown, K. M. (2001). *Topsoil foraging – an architectural adaptation of plants*
666 *to low phosphorus availability*. 13.

667 Lynch, J. P., Nielsen, K. L., Davis, R. D., and Jablonsky, A. G. (1997). *SimRoot: Modelling and*
668 *visualization of root systems*. 14.

669 Lynch, J. P., and St.Clair, S. B. (2004). Mineral stress: The missing link in understanding how
670 global climate change will affect plants in real world soils. *Field Crops Research*, 90(1),
671 101–115. <https://doi.org/10.1016/j.fcr.2004.07.008>

672 Mano, Y., Omori, F., Takamizo, T., Kindiger, B., Bird, R. McK., and Loaisiga, C. H. (2006).
673 Variation for Root Aerenchyma Formation in Flooded and Non-Flooded Maize and
674 Teosinte Seedlings. *Plant and Soil*, 281(1–2), 269–279. [https://doi.org/10.1007/s11104-](https://doi.org/10.1007/s11104-005-4268-y)
675 [005-4268-y](https://doi.org/10.1007/s11104-005-4268-y)

676 Miguel, M. A., Postma, J. A., and Lynch, J. P. (2015). Phenological Synergism between Root Hair
677 Length and Basal Root Growth Angle for Phosphorus Acquisition. *Plant Physiology*,
678 167(4), 1430–1439. <https://doi.org/10.1104/pp.15.00145>

679 Miller, C. R., Ochoa, I., Nielsen, K. L., Beck, D., and Lynch, J. P. (2003). Genetic variation for
680 adventitious rooting in response to low phosphorus availability: Potential utility for
681 phosphorus acquisition from stratified soils. *Functional Plant Biology*, 30(9), 973.
682 <https://doi.org/10.1071/FP03078>

683 Nielsen, K. L., Eshel, A., and Lynch, J. P. (2001). The effect of phosphorus availability on the
684 carbon economy of contrasting common bean (*Phaseolus vulgaris* L.) genotypes. 11.

685 Nielsen, K. L., Lynch, J. P., Jablonsky, A. G., and Curtis, P. S. (1994). Carbon cost of root
686 systems: An architectural approach. *Plant and Soil*, 165(1), 161–169.
687 <https://doi.org/10.1007/BF00009972>

688 Ning, J., and Cumming, J. R. (2001). Arbuscular mycorrhizal fungi alter phosphorus relations of
689 broomsedge (*Andropogon virginicus* L.) plants. *Journal of Experimental Botany*,
690 52(362), 1883–1891. <https://doi.org/10.1093/jexbot/52.362.1883>

691 Pieruschka, R., and Poorter, H. (2012). Phenotyping plants: Genes, phenes and machines.
692 *Functional Plant Biology*, 39(11), 813. https://doi.org/10.1071/FPv39n11_IN

693 Postma, Johannes A., and Lynch, J. P. (2011a). Theoretical evidence for the functional benefit of
694 root cortical aerenchyma in soils with low phosphorus availability. *Annals of Botany*,
695 107(5), 829–841. <https://doi.org/10.1093/aob/mcq199>

696 Postma, Johannes Auke, and Lynch, J. P. (2011b). Root Cortical Aerenchyma Enhances the
697 Growth of Maize on Soils with Suboptimal Availability of Nitrogen, Phosphorus, and
698 Potassium. *Plant Physiology*, 156(3), 1190–1201. <https://doi.org/10.1104/pp.111.175489>

699 Postma, Johannes A., and Lynch, J. P. (2012). Complementarity in root architecture for nutrient
700 uptake in ancient maize/bean and maize/bean/squash polycultures. *Annals of Botany*,
701 110(2), 521–534. <https://doi.org/10.1093/aob/mcs082>

702 Postma, Johannes A., Schurr, U., and Fiorani, F. (2014). Dynamic root growth and architecture
703 responses to limiting nutrient availability: Linking physiological models and
704 experimentation. *Biotechnology Advances*, 32(1), 53–65.
705 <https://doi.org/10.1016/j.biotechadv.2013.08.019>

706 Rangarajan H, Postma J, Lynch JP. (2018). Co-optimization of axial root phenotypes for nitrogen
707 and phosphorus acquisition in common bean. *Annals of Botany* 122: 485–499.

708 Richardson, A. E., Lynch, J. P., Ryan, P. R., Delhaize, E., Smith, F. A., Smith, S. E., Harvey, P.
709 R., Ryan, M. H., Veneklaas, E. J., Lambers, H., Oberson, A., Culvenor, R. A., and
710 Simpson, R. J. (2011). Plant and microbial strategies to improve the phosphorus
711 efficiency of agriculture. *Plant and Soil*, 349(1–2), 121–156.
712 <https://doi.org/10.1007/s11104-011-0950-4>

713 Rubio, G., Liao, H., Yan, X., and Lynch, J. P. (2003). Topsoil Foraging and Its Role in Plant
714 Competitiveness for Phosphorus in Common Bean. *Crop Science*, 43(2), 598.
715 <https://doi.org/10.2135/cropsci2003.0598>

716 Sablowski, R., and Carnier Dornelas, M. (2014). Interplay between cell growth and cell cycle in
717 plants. *Journal of Experimental Botany*, 65(10), 2703–2714.
718 <https://doi.org/10.1093/jxb/ert354>

719 Saengwilai, P., Nord, E. A., Chimungu, J. G., Brown, K. M., and Lynch, J. P. (2014). Root
720 Cortical Aerenchyma Enhances Nitrogen Acquisition from Low-Nitrogen Soils in Maize.
721 *PLANT PHYSIOLOGY*, 166(2), 726–735. <https://doi.org/10.1104/pp.114.241711>

722 Simunek, J., Huang, K., and van Genuchten, M. (1995). *The SWMS3D Code for Simulating Water*
723 *Flow and Solute Transport in Three-Dimensional Variably-Saturated Media*.

724 Spitters, C. J. T., and Schapendonk, A. H. C. M. (1990). Evaluation of breeding strategies for
725 drought tolerance in potato by means of crop growth simulation. *Plant and Soil*, 123(2),
726 193–203. <https://doi.org/10.1007/BF00011268>

727 St.Clair, S. B., and Lynch, J. P. (2010). The opening of Pandora’s Box: Climate change impacts
728 on soil fertility and crop nutrition in developing countries. *Plant and Soil*, 335(1–2), 101–
729 115. <https://doi.org/10.1007/s11104-010-0328-z>

730 Trachsel, S., Kaeppler, S. M., Brown, K. M., and Lynch, J. P. (2011). Shovelomics: High
731 throughput phenotyping of maize (*Zea mays* L.) root architecture in the field. *Plant and*
732 *Soil*, 341(1–2), 75–87. <https://doi.org/10.1007/s11104-010-0623-8>

733 Uhart, S. A., and Andrade, F. H. (1995). Nitrogen Deficiency in Maize: I. Effects on Crop
734 Growth, Development, Dry Matter Partitioning, and Kernel Set. *Crop Science*, 35(5),
735 1376–1383. <https://doi.org/10.2135/cropsci1995.0011183X003500050020x>

736 Vanhees, D.J., Loades, K.W., Bengough, A.G., Mooney, S. J., and Lynch, J. P. (2020). Root
737 anatomical traits contribute to deeper rooting of maize under compacted field conditions.
738 *Journal of Experimental Botany*. <https://doi.org/10.1093/jxb/eraa165>

739 Walk, T.C., Jaramillo, R., Lynch, J. P. (2006). Architectural tradeoffs between adventitious and
740 basal roots for phosphorus acquisition. *Plant and Soil* 279: 347–366.

741 Wang, X., Yan, X., and Liao, H. (2010). Genetic improvement for phosphorus efficiency in
742 soybean: A radical approach. *Annals of Botany*, 106(1), 215–222.
743 <https://doi.org/10.1093/aob/mcq029>

744 White, J. W., Rassweiler, A., Samhour, J. F., Stier, A. C., and White, C. (2014). Ecologists
745 should not use statistical significance tests to interpret simulation model results. *Oikos*,
746 123(4), 385–388. <https://doi.org/10.1111/j.1600-0706.2013.01073.x>

747 Whiteley, G. M., Hewitt, J. S., and Dexter, A. R. (1982). The buckling of plant roots. *Physiologia*
748 *Plantarum*, 54(3), 333–342. <https://doi.org/10.1111/j.1399-3054.1982.tb00268.x>

749 York, L. M., and Lynch, J. P. (2015). Intensive field phenotyping of maize (*Zea mays* L.) root
750 crowns identifies phenes and phene integration associated with plant growth and nitrogen
751 acquisition. *Journal of Experimental Botany*, 66(18), 5493–5505.
752 <https://doi.org/10.1093/jxb/erv241>

753 York, L. M., Nord, E. A., and Lynch, J. P. (2013). Integration of root phenes for soil resource
754 acquisition. *Frontiers in Plant Science*, 4. <https://doi.org/10.3389/fpls.2013.00355>

755 York, L. M., Silberbush, M., and Lynch, J. P. (2016). Spatiotemporal variation of nitrate uptake
756 kinetics within the maize (*Zea mays* L.) root system is associated with greater nitrate
757 uptake and interactions with architectural phenes. *Journal of Experimental Botany*,
758 67(12), 3763–3775. <https://doi.org/10.1093/jxb/erw133>

759 Zhao, D., Oosterhuis, D. M., and Bednarz, C. W. (2001). Influence of Potassium Deficiency on
760 Photosynthesis, Chlorophyll Content, and Chloroplast Ultrastructure of Cotton Plants.
761 *Photosynthetica*, 39(1), 103–109. <https://doi.org/10.1023/A:1012404204910>

762 Zhu, J., Brown, K. M., and Lynch, J. P. (2010a). Root cortical aerenchyma improves the drought
763 tolerance of maize (*Zea mays* L.). *Plant, Cell and Environment*.
764 <https://doi.org/10.1111/j.1365-3040.2009.02099.x>

765 Zhu, J., Zhang, C., Lynch, J. P. (2010b). The utility of phenotypic plasticity of root hair length for
766 phosphorus acquisition. *Functional Plant Biology* 37: 313–322.

767 Zhu, J., Kaeppler, S. M., and Lynch, J. P. (2005). Topsoil foraging and phosphorus acquisition
768 efficiency in maize (*Zea mays*). *Functional Plant Biology*, 32(8), 749.
769 <https://doi.org/10.1071/FP05005>

770 Zhu, J., and Lynch, J. P. (2004). The contribution of lateral rooting to phosphorus acquisition
771 efficiency in maize (*Zea mays*) seedlings. *Functional Plant Biology*, 31(10), 949.
772 <https://doi.org/10.1071/FP04046>

773

774 Figures

792 Figure 1. Variation in tissue nutrient concentration among maize genotypes contrasting in CCFN
793 and CCS. Unit of tissue nutrient concentration is $\mu\text{mol/g}$ dry weight. IBM178 is a many CCFN
794 genotype, IBM201 is a reduced CCFN genotype, IBM365 is a small CCS genotype, IBM30 is a
795 larger CCS genotype. In the nutrient solution, N concentration is 160 $\mu\text{mol/L}$, and K concentration
796 is 60 $\mu\text{mol/L}$ under N or K stress respectively. P concentration is 360 $\mu\text{mol/L}$. Error bars represent
797 standard deviation of measurements from four replications.

798
799
800
801
802

Figure 2. Visualized output of the simulated growth of maize root systems at 42 D.A.G. Plant growth was simulated under moderate N stress (42 kg/ha). Plant A represents a few CCFN genotype (8 cortical cell files), plant B represents a large CCS genotype (533 microns), and plant C represents a reference genotype with increased CCFN (17 cortical cell files) and reduced CCS (101 microns). The axis represents the rooting depth of the three plants in centimeters.

831 Figure 3. The benefits of RCA formation, larger CCS and small CCFN under suboptimal
832 availability of N, P and K. Stresses are expressed as the relative plant dry weight at 40 days after
833 germination compared to a non-stressed reference plant on the x axis. Benefits are expressed as
834 increase in plant dry weight due to the presence of the phenes compared to a reference
835 phenotype. The phenes were at the maximum beneficial level, i.e., maximum RCA formation,
836 largest cortical cells and least cortical cell files simulated independently.
837

838

839 Figure 4. Total plant dry weight showing the utility of larger CCS (533 μm^2) and reduced CCFN
840 (8 cell files) vs the reference phenotype (101 μm^2 CCS and 17 CCFN) under N, P and K stress
841 (plant dry weight 10% of unstressed) at 40 days after germination. Error bars represent standard
842 deviation in four repeated runs. Variation was caused by stochasticity in modeled root growth rate,
843 root branching frequency, and root growth angle.

844

845

846

Figure 5. Sensitivity analysis of CCS and CCFN variation on benefits introduced by reducing respiration. Different lines correspond with percent sufficient soil nutrient availabilities as indicated. Cortical cell size and cell file number values used for simulations were based on empirical data from the literature and were within the range observed empirically. Benefits are expressed as increase in plant dry weight due to the presence of the phene states compared to the model default phenotype. N level (10% = 21 kg/ha, 20% = 42 kg/ha, 30% = 63 kg/ha, 40% = 84 kg/ha, 80% = 168 kg/ha), P level (01% = 0.05 kg/ha, 08% = 0.4 kg/ha, 30% = 1.5 kg/ha, 50% = 2.5 kg/ha, 80% = 4 kg/ha), K level (10% = 0.5 kg/ha, 25% = 1.2 kg/ha, 40% = 1.9 kg/ha, 80% = 3.8 kg/ha).

858
859 Figure 6. Nitrogen and phosphorus stress as affected by contrasting CCS and CCFN on a scale from
860 0-1. Stress is calculated as $1 - (\mu - m)/(o - m)$, where μ is the amount of nitrate or phosphorus
861 being uptaken, o is the optimal nitrate or phosphorus content in the plant, and m is the minimal
862 nitrate or phosphorus content in the plant. 0 indicates no stress, 1 indicates the most severe stress.
863 HN = 210 kg/ha, MN = 84 kg/ha, LN = 21kg/ha, HP = 5kg/ha, MP = 2kg/ha, LP = 0.5kg/ha.
864

865
866
867

868
869

870 Figure 7. Relative benefit of having RCA formation, larger CCS and small CCFN simultaneously
871 in a simulated plant over time at 42 kg/ha of soil nitrate, 0.5 kg/ha of soil phosphorus and 1.5kg/ha
872 of soil potassium availability. Different lines correspond to relative benefits for respiration and
873 nutrient content, similarly described in figure 2. The gray line indicates the hypothetical additive
874 benefit when RCA formation achieves the optimal growth enhancement

876 Figure 8. Interactions between larger CCS and Lateral Root Branching Density (LRBD) (A, C and
877 E), and reduced CCFN and LRBD (B, D, and F) under low or medium soil nitrogen, phosphorus
878 or potassium availability. CCS and CCFN used in this scenario are both at the least level of
879 metabolic carbon demand (largest cell size, reduced cell files). Three levels of Lateral Root
880 Branching Density (4, 8, and 16 roots/cm) are shown, the range of which was based on Trachsel et
881 al. (2010), as used by Postma and Lynch (2011). Low and medium nitrate levels are 21 kg/ha and
882 84 kg/ha respectively. Low and medium phosphorus levels are 0.5 kg/ha and 2 kg/ha respectively.
883 Low and medium potassium levels are 0.5 kg/ha and 1.9 kg/ha respectively.

884

885

886

887

Figure 9. Benefits of CCS and CCFN on plants grown under nitrogen or phosphorus deficiencies as affected by elevated atmospheric CO₂ concentration, as simulated by increased canopy light use efficiency. Soil phosphorus level was 0.4kg/ha, nitrate level 21kg/ha, and potassium level 1kg/ha.

920 Appendix

921

922 OpenSimRoot Parameterization

923 OpenSimRoot uses a hierarchical xml formatted input file which is graphically presented
924 below. The hierarchy places the parameters in a context. For example, the parameter 'specific leaf
925 area' belongs to the shoot of a specific genotype. In OpenSimRoot parameters can be a single
926 value, a value drawn from a distribution, or the result of an interpolation table. We have tried to
927 base all our parameters on our own measurements or data from the literature. We have indicated
928 the sources behind the parameters. Note that in many cases, we used more than one source and in
929 some cases we had to convert the measurements using assumptions. A common assumption we
930 made is that the value was equal for all root classes and or for all genotypes.

```
931     1 'environment'
932     1.1 'atmosphere'
933         1.1.1 'evaporation' [cm]=f{'time'} [day] x,y pairs : {0 0 1 0.05 2 0.1 3 0.1 4 0.05 5 0.05 6
934 0.1 7 0.05 8 0.05 9 0.1 10 0.1 11 0.05 12 0.1 13 0.1 14 0.05 15 0.04 16 0.03 17 0.02 18 0.09 19
935 0.09 20 0.04 21 0.09 22 0.09 23 0.04 24 0.03 25 0.02 26 0.02 27 0.08 28 0.03 29 0.08 30 0.03 31
936 0.08 32 0.07 33 0.07 34 0.07 35 0.03 36 0.02 37 0.01 38 0 39 0 40 0 41 0 42 0.06}
937         1.1.2 'irradiation' = 4000 [umol.cm-2.day-1]
938         1.1.3 'precipitation' [cm]=f{'time'} [day] x,y pairs : {0 0 1 0 2 1 3 0.29 4 0 5 0 6 0.61 7 0 8
939 0 9 0.25 10 0.03 11 0 12 0.64 13 0.33 14 0 15 0 16 0 17 0 18 1.8 19 0.2 20 0 21 2.84 22 0.38 23 0
940 24 0 25 0 26 0 27 0.18 28 0 29 0.46 30 0 31 1.35 32 0.13 33 0.23 34 0.25 35 0 36 0 37 0 38 0 39 0
941 40 0 41 0 42 1.42} (Rocksprings, PA, weather station data June 2009)
942     1.2 'dimensions'
943         1.2.1 'max corner' = 13 0 30 [cm]
944         1.2.2 'min corner' = -13 -150 -30 [cm]
945     1.3 'silt-loam soil'
946         1.3.1 'bulk density' [g.cm-3]=f{'depth'} [cm] x,y pairs : {-200 1.51 -65 1.51 -47 1.4 -30
947 1.42 -16 1.29 -5 1.24 0 1.24} (M.B. Postma, University Park, unpublished)
948         1.3.2 'nitrate'
949             1.3.2.1 'adsorption coefficient' = 0 [umol.cm-1]
950             1.3.2.2 'buffer power' [-]=f{'depth'} [cm] x,y pairs : {-1000 0.4 1000 0.4}
951             1.3.2.3 'concentration' [umol.ml-1]=f{'depth'} [cm] x,y pairs : {-1000 1.59 -55 1.59 -45
952 1.67 -35 2.17 -25 3.15 -15 4.02 -5 2.36 0 2.8 0.01 0 100 0} (M.B. Postma, University Park,
953 unpublished)
954             1.3.2.4 'diffusion coefficient' [cm2.day-1]=f{'depth'} [cm] x,y pairs : {-1000 0.07 -0 0.07
955 1e-05 1e-08 1000 1e-08}
956             1.3.2.5 'longitudinal dispersivity' = 1 [cm]
957             1.3.2.6 'r1-r0' = 4 [cm]
958             1.3.2.7 'saturated diffusion coefficient' = 1.6416 [cm2.day-1]
959             1.3.2.8 'transverse dispersivity' = 0.5 [cm]
960         1.3.3 'organic' (Yang and Janssen 2000)
961             1.3.3.1 'C/N ratio microbes' = 10 [g.g-1] 1.3.3.2 'C/N ratio' [g.g-1]=f{'depth'} [cm] x,y
962 pairs : {-10000 13 0 13}
963             1.3.3.3 'assimilation efficiency microbes' = 1 [-]
964             1.3.3.4 'carbon content' [g.g-1]=f{'depth'} [cm] x,y pairs : {-200 0.005 -40 0.005 -30 0.01
965 -10 0.02
966             0 0.02}
```

967 1.3.3.5 'initial relative mineralisation rate' [g.g-1.year-1]=f{'depth'} [cm] x,y pairs :{-
 968 1000 0 -25 0
 969 -10 0.037 0 0.037} (Postma, University Park, Unpublished)
 970 1.3.3.5.1 'multiplier' = 0.1 [-]
 971 1.3.3.6 'speed of aging' = 0.46 [-]
 972 1.3.3.7 'time offset' = 30 [day]
 973 1.3.4 'phosphorus' (S. A. Barber 1995; Bhadoria et al. 1991)
 974 1.3.4.1 'adsorption coefficient' = 1333.3 [umol.cm-1]
 975 1.3.4.2 'buffer power' [-]=f{'depth'} [cm] x,y pairs :{-1000 400 1000 400}
 976 1.3.4.3 'concentration' [umol.ml-1]=f{'depth'} [cm] x,y pairs :{-1000 0.00024 -30
 977 0.00025 -29 0.00175 0 0.00175 0.0001 0 1000 0}
 978 1.3.4.4 'diffusion coefficient' [cm2.day-1]=f{'depth'} [cm] x,y pairs :{-1000 0.00019872
 979 1000 0.00019872}
 980 1.3.4.5 'longitudinal dispersivity' = 0 [cm]
 981 1.3.4.6 'r1-r0' = 0.3 [cm]
 982 1.3.4.7 'saturated diffusion coefficient' = 0.00495 [cm2.day-1]
 983 1.3.4.8 'transverse dispersivity' = 0 [cm]
 984 1.3.5 'potassium' (Claassen et al. 1986; S. A. Barber 1995; Dunham and Nye 1976)
 985 1.3.5.1 'adsorption coefficient' = 33.3 [umol.cm-1]
 986 1.3.5.2 'buffer power' [-]=f{'depth'} [cm] x,y pairs :{-1000 10 1000 10}
 987 1.3.5.3 'concentration' [umol.ml-1]=f{'depth'} [cm] x,y pairs :{-1000 0.05 -30 0.05 -29
 988 0.15 0 0.15 1e-05 0 1000 0}
 989 1.3.5.4 'diffusion coefficient' [cm2.day-1]=f{'depth'} [cm] x,y pairs :{-1000 0.067 1000
 990 0.067}
 991 1.3.5.5 'longitudinal dispersivity' = 1 [cm]
 992 1.3.5.6 'r1-r0' = 1.5 [cm]
 993 1.3.5.7 'saturated diffusion coefficient' = 1.56 [cm2.day-1]
 994 1.3.5.8 'transverse dispersivity' = 0.5 [cm]
 995 1.3.6 'water'
 996 1.3.6.1 'initial hydraulic head' [cm]=f{'depth'} [cm] x,y pairs :{-200 0 -151 -50 -50 -150 -
 997 45 -155 -40 -160 -35 -165 -30 -170 -25 -175 -20 -180 -15 -190 -10 -200 -5 -220 -2 -240 -1 -300 -
 998 0 -400} (M.B. Postma, University Park, Unpublished)
 999 1.3.6.2 'residual water content' [100%]=f{'depth'} [cm] x,y pairs :{-300 0.067 0 0.067}
 1000 (Hodnett and Tomasella 2002)
 1001 1.3.6.3 'saturated conductivity' [cm.day-1]=f{'depth'} [cm] x,y pairs :{-300 10.8 0 10.8}
 1002 (Luo et al. 2008)
 1003 1.3.6.4 'saturated water content' [100%]=f{'depth'} [cm] x,y pairs :{-300 0.39 -65 0.39 -
 1004 35 0.39 -25 0.43 -15 0.45 0 0.46} (Luo et al. 2008)
 1005 1.3.6.5 'van genuchten:alpha' [-.cm-1]=f{'depth'} [cm] x,y pairs :{-300 0.02 0 0.02}
 1006 (Hodnett and Tomasella 2002)
 1007 1.3.6.6 'van genuchten:n' [-]=f{'depth'} [cm] x,y pairs :{-300 1.41 0 1.41} (Hodnett and
 1008 Tomasella 2002)
 1009 1.3.6.7 'volumetric water content in Barber Cushman' = 0.3 [cm3.cm-3]
 1010 1.4 'loamy-sand soil'
 1011 1.4.1 'water'
 1012 1.4.1.1 'initial hydraulic head' [cm]=f{'depth'} [cm] x,y pairs :{-200 -0 -90 -110 -32 -168
 1013 -28 -172 -0 -200}
 1014 1.4.1.2 'residual water content' [100%]=f{'depth'} [cm] x,y pairs :{-300 0.057 0 0.057}
 1015 1.4.1.3 'saturated conductivity' [cm.day-1]=f{'depth'} [cm] x,y pairs :{-200 400 0 400}

1016 1.4.1.4 'saturated water content' [100%]=f{'depth'} [cm] x,y pairs :{-200 0.339 -32 0.339
1017 -28 0.399 0 0.399}
1018 1.4.1.5 'van genuchten:alpha' [-.cm-1]=f{'depth'} [cm] x,y pairs :{-300 0.033 -30 0.033 -
1019 28 0.038 0 0.038}
1020 1.4.1.6 'van genuchten:n' [-]=f{'depth'} [cm] x,y pairs :{-200 1.6024 -32 1.6024 -28
1021 1.3757 0 1.3757}
1022 1.4.1.7 'volumetric water content in Barber Cushman' = 0.18 [cm³.cm-3]
1023 1.4.2 see silt-loam soil for other parameters
1024 2 'plant parameters'
1025 2.1.1 'braceroots'
1026 2.1.1.1 'aerenchyma formation' [100%]=f{'age'} [day] x,y pairs :{0 0 3 0 5 0.1 10 0.25 20
1027 0.393 1000 0.393} (Fan, Zhu, et al. 2003)
1028 'cell size' [um²]={101 150 200 250 300 350 400 450 500 533} (Chimungu et al., 2014a)
1029 'cell file number' [#]={8 9 10 11 12 13 14 15 16 17}
1030 2.1.1.2 'branch list'
1031 2.1.1.2.1 'lateral of crown roots'
1032 2.1.1.2.1.1 'allow branches to form above ground' = 0 [-]
1033 2.1.1.2.1.2 'branching frequency' [cm]=f{'uniform distribution'} minimum=0.1
1034 maximum=0.3
1035 2.1.1.2.1.3 'branching spatial offset' = 12 [cm]
1036 2.1.1.2.1.4 'length root tip' = 10.93 [cm]
1037 2.1.1.2.1.5 'number of branches/whorl' = 1 [#]
1038 2.1.1.3 'branching angle' = 140 [degrees]
1039 2.1.1.4 'density' = 0.094 [g.cm-3] (Pahlavanian and Silk 1988)
1040 2.1.1.5 'diameter' [cm]=f{'age'} [day] x,y pairs :{0 0.4 8 0.4 15 0.15 24 0.1 100 0.1}
1041 2.1.1.6 'gravitropism.v2' [cm]=f{'uniform distribution'} minimum=-0.01 maximum=-
1042 0.005
1043 2.1.1.7 'growth rate' [cm.day-1]=f{'age'} [day] x,y pairs :{0 0.01 5 1 10 4.5 17 4.5 22 0
1044 1000 0}
1045 2.1.1.8 'length root tip without xylem vessels' = 2 [cm]
1046 2.1.1.9 'longitudinal growth rate multiplier' [cm]=f{'uniform distribution'} minimum=0.7
1047 maximum=1
1048 2.1.1.10 'nitrate'
1049 2.1.1.10.1 'Cmin' = 0.001 [umol.ml-1]
1050 2.1.1.10.2 'Imax' [umol.cm-2.day-1]=f{'age'} [day] x,y pairs :{0 1.21 2 2.1 40 2.1}
1051 2.1.1.10.3 'Km' [umol.ml-1]=f{'age'} [day] x,y pairs :{0 0.0157 2 0.0522 40 0.0522}
1052 2.1.1.10.4 'minimal nutrient concentration' = 600 [umol.g-1]
1053 2.1.1.10.5 'optimal nutrient concentration' = 1200 [umol.g-1]
1054 2.1.1.11 'number of xylem poles' = 40 [-]
1055 2.1.1.11 'phosphorus' (S. A. Barber 1995)
1056 2.1.1.11.1 'Cmin' = 0.0002 [umol.ml-1]
1057 2.1.1.11.2 'Imax' = 0.0555 [umol.cm-2.day-1]
1058 2.1.1.11.3 'Km' = 0.00545 [umol.ml-1]
1059 2.1.1.11.4 'minimal nutrient concentration' = 30 [umol.g-1]
1060 2.1.1.11.5 'optimal nutrient concentration' = 60 [umol.g-1]
1061 2.1.1.12 'potassium' (Barber 1995)
1062 2.1.1.12.1 'Cmin' = 0.002 [umol.ml-1]
1063 2.1.1.12.2 'Imax' = 0.467 [umol.cm-2.day-1]
1064 2.1.1.12.3 'Km' = 0.014 [umol.ml-1]

1065 2.1.1.12.4 'minimal nutrient concentration' = 117 [umol.g-1] 2.1.1.11.5 'optimal nutrient
 1066 concentration' = 234 [umol.g-1] (Silk et al. 1986)
 1067 2.1.1.13 'reduction in respiration due to aerenchyma' [100%]=f{'aerenchymaFormation'}
 1068 [100%] x,y pairs : {0 0 0.3 0.7 0.6 1} (Fan et al. 2003)
 1069 'reduction in respiration due to cell size' [100%]=f{cellSize} [100%] x,y pairs : {101 0
 1070 150 0.07 200 0.14 250 0.17 300 0.25 350 0.32 400 0.37 450 0.43 500 0.51 533 0.57}
 1071 'reduction in respiration due to file number' [100%]=f{fileNumber} [100%] x,y pairs : {17
 1072 0 16 0.05 15 0.13 14 0.17 13 0.23 12 0.3 11 0.35 10 0.42 9 0.49 8 0.52}
 1073 2.1.1.14 'regular topology' = 4 [-]
 1074 2.1.1.15 'relative carbon cost of exudation' [g.cm-1.day-1]=f{'age'} [day] x,y pairs : {0 5e-
 1075 06 100 5e-06} (Groleau-Renaud et al. 1998)
 1076 2.1.1.16 'relative respiration' [g.g-1.day-1]=f{'age'} [day] x,y pairs : {0 0.09 2 0.04 6 0.04
 1077 1000 0.04} (Fan et al., 2003)
 1078 2.1.1.17 'root class id' = 102 [-]
 1079 2.1.1.18 'root hair density' [#.cm-2]=f{'age'} [day] x,y pairs : {0 2000 1 2000 2 2000 10
 1080 2000 30 0 2000 0} (Zhu, Kaeppler, et al. 2005; Mackay and S. Barber 1985)
 1081 2.1.1.19 'root hair diameter' = 0.0005 [cm]
 1082 2.1.1.20 'root hair length' [cm]=f{'age'} [day] x,y pairs : {0 0 1 0 2 0.028 2000 0.028}
 1083 2.4.15.20 (Zhu et al. 2005; Mackay and S. Barber 1985)
 1084 2.1.1.21 'soil impedance.v2' [cm]=f{'uniform distribution'} minimum=-0.03
 1085 maximum=0.03
 1086 2.1.2 'braceroots2'
 1087 2.1.2.1 'aerenchyma formation' [100%]=f{'age'} [day] x,y pairs : {0 0 3 0 5 0.1 10 0.25 20
 1088 0.393 1000 0.393} (Fan et al. 2003)
 1089 'cell size' [um^2]={101 150 200 250 300 350 400 450 500 533} (Chimungu et al., 2014a)
 1090 'cell file number' [#]={8 9 10 11 12 13 14 15 16 17}
 1091 2.1.2.2 'branch list'
 1092 2.1.2.2.1 'lateral of crown roots'
 1093 2.1.2.2.1.1 'allow branches to form above ground' = 0 [-]
 1094 2.1.2.2.1.2 'branching frequency' [cm]=f{'uniform distribution'} minimum=0.1
 1095 maximum=0.4
 1096 2.1.2.2.1.3 'branching spatial offset' = 15 [cm]
 1097 2.1.2.2.1.4 'length root tip' = 10.93 [cm]
 1098 2.1.2.2.1.5 'number of branches/whorl' = 1 [#]
 1099 2.1.2.3 'branching angle' = 130 [degrees]
 1100 2.1.2.4 'density' = 0.094 [g.cm-3]
 1101 2.1.2.5 'diameter' [cm]=f{'age'} [day] x,y pairs : {0 0.5 9 0.5 16 0.2 24 0.1 100 0.1}
 1102 2.1.2.6 'gravitropism.v2' [cm]=f{'uniform distribution'} minimum=-0.01 maximum=-
 1103 0.005
 1104 2.1.2.7 'growth rate' [cm.day-1]=f{'age'} [day] x,y pairs : {0 0.01 5 1 10 4.5 17 4.5 22 0
 1105 1000 0}
 1106 2.1.2.8 'length root tip without xylem vessels' = 2 [cm]
 1107 2.1.2.9 'longitudinal growth rate multiplier' [cm]=f{'uniform distribution'} minimum=0.7
 1108 maximum=1
 1109 2.1.2.10 'nitrate'
 1110 2.1.2.10.1 'Cmin' = 0.001 [umol.ml-1]
 1111 2.1.2.10.2 'Imax' [umol.cm-2.day-1]=f{'age'} [day] x,y pairs : {0 1.21 2 2.1 40 2.1}
 1112 2.1.2.10.3 'Km' [umol.ml-1]=f{'age'} [day] x,y pairs : {0 0.0157 2 0.0522 40 0.0522}
 1113 2.1.2.10.4 'minimal nutrient concentration' = 600 [umol.g-1]

1114 2.1.2.10.5 'optimal nutrient concentration' = 1200 [umol.g-1]
 1115 2.1.2.11 'number of xylem poles' = 48 [-]
 1116 2.1.2.12 'phosphorus' (Barber 1995)
 1117 2.1.2.12.1 'Cmin' = 0.0002 [umol.ml-1]
 1118 2.1.2.12.2 'Imax' = 0.0555 [umol.cm-2.day-1]
 1119 2.1.2.12.3 'Km' = 0.00545 [umol.ml-1]
 1120 2.1.2.12.4 'minimal nutrient concentration' = 30 [umol.g-1]
 1121 2.1.2.12.5 'optimal nutrient concentration' = 60 [umol.g-1]
 1122 2.1.2.13 'potassium' (Barber 1995)
 1123 2.1.2.13.1 'Cmin' = 0.002 [umol.ml-1]
 1124 2.1.2.13.2 'Imax' = 0.467 [umol.cm-2.day-1]v2.1.2.13.3 'Km' = 0.014 [umol.ml-1]
 1125 2.1.2.13.4 'minimal nutrient concentration' = 117 [umol.g-1]
 1126 2.1.2.13.5 'optimal nutrient concentration' = 234 [umol.g-1] (Silk et al. 1986)
 1127 2.1.2.14 'reduction in respiration due to aerenchyma' [100%]=f{'aerenchymaFormation'}
 1128 [100%]
 1129 x,y pairs :{0 0 0.3 0.7 0.6 1} (Fan et al. 2003)
 1130 'reduction in respiration due to cell size' [100%]=f{'cellSize'} [100%] x,y pairs :{101 0
 1131 150 0.07 200 0.14 250 0.17 300 0.25 350 0.32 400 0.37 450 0.43 500 0.51 533 0.57}
 1132 'reduction in respiration due to file number' [100%]=f{'filename'} [100%] x,y pairs :{17
 1133 0 16 0.05 15 0.13 14 0.17 13 0.23 12 0.3 11 0.35 10 0.42 9 0.49 8 0.52}
 1134 2.1.2.15 'regular topology' = 3 [-]
 1135 2.1.2.16 'relative carbon cost of exudation' [g.cm-1.day-1]=f{'age'} [day] x,y pairs :{0 5e-
 1136 06 100
 1137 5e-06} (Groleau-Renaud et al. 1998)
 1138 2.1.2.17 'relative respiration' [g.g-1.day-1]=f{'age'} [day] x,y pairs :{0 0.09 2 0.04 6 0.04
 1139 1000
 1140 0.04} (Fan et al., 2003)
 1141 2.1.2.18 'root class id' = 102 [-]
 1142 2.1.2.19 'root hair density' [#.cm-2]=f{'age'} [day] x,y pairs :{0 2000 1 2000 2 2000 10
 1143 2000 30 0
 1144 2000 0} (Zhu et al. 2005; Mackay and S. Barber 1985)
 1145 2.1.2.20 'root hair diameter' = 0.0005 [cm]
 1146 2.1.2.21 'root hair length' [cm]=f{'age'} [day] x,y pairs :{0 0 1 0 2 0.028 2000 0.028}
 1147 2.4.15.20
 1148 (Zhu et al. 2005; Mackay and S. Barber 1985)
 1149 2.1.2.22 'soil impedance.v2' [cm]=f{'uniform distribution'} minimum=-0.03
 1150 maximum=0.03
 1151 2.1.3 'finelateral'
 1152 2.1.3.1 'aerenchyma formation' [100%]=f{'age'} [day] x,y pairs :{0 0 3 0 5 0.1 10 0.25 20
 1153 0.393 1000 0.393} (Fan et al. 2003)
 1154 'cell size' [um^2]={101 150 200 250 300 350 400 450 500 533} (Chimungu et al., 2014a)
 1155 'cell file number' [#]={8 9 10 11 12 13 14 15 16 17}
 1156 2.1.3.2 'branch list'
 1157 2.1.3.2.1 'finelateral2'
 1158 2.1.3.2.1.1 'allow branches to form above ground' = 0 [-]
 1159 2.1.3.2.1.2 'branching frequency' [cm]=f{'uniform distribution'} minimum=0.4
 1160 maximum=0.6
 1161 2.1.3.2.1.3 'length root tip' = 1.5 [cm]
 1162 2.1.3.3 'branching angle' = 62.83 [degrees]

1163 2.1.3.4 'density' = 0.094 [g.cm-3]
 1164 2.1.3.5 'diameter' = 0.025 [cm]
 1165 2.1.3.6 'gravitropism.v2' = 0 0 0 [cm]
 1166 2.1.3.7 'growth rate' [cm.day-1]=f{'age'} [day] x,y pairs :{0 0.01 1 0.35 6 0 1000 0}
 1167 2.1.3.8 'length root tip without xylem vessels' = 2 [cm]
 1168 2.1.3.9 'longitudinal growth rate multiplier' [cm]=f{'normal distribution'} minimum=0.5
 1169 maximum=1.5 mean=1 stdev=0.1
 1170 2.1.3.10 'nitrate'
 1171 2.1.3.10.1 ' Cmin' = 0.0017 [umol.ml-1]
 1172 2.1.3.10.2 ' Imax' = 1.27 [umol.cm-2.day-1]
 1173 2.1.3.10.3 ' Km' = 0.0027 [umol.ml-1]
 1174 2.1.3.10.4 'minimal nutrient concentration' = 600 [umol.g-1]
 1175 2.1.3.10.5 'optimal nutrient concentration' = 1200 [umol.g-1]
 1176 2.1.3.11 'number of xylem poles' = 4 [-]
 1177 2.1.3.12 'phosphorus' (Barber 1995)
 1178 2.1.3.12.1 ' Cmin' = 0.0002 [umol.ml-1]
 1179 2.1.3.12.2 ' Imax' = 0.0555 [umol.cm-2.day-1]
 1180 2.1.3.12.3 ' Km' = 0.00545 [umol.ml-1]
 1181 2.1.3.12.4 'minimal nutrient concentration' = 30 [umol.g-1]
 1182 2.1.3.12.5 'optimal nutrient concentration' = 60 [umol.g-1]
 1183 2.1.3.13 'potassium' (Barber 1995)
 1184 2.1.3.13.1 ' Cmin' = 0.002 [umol.ml-1]
 1185 2.1.3.13.2 ' Imax' = 0.467 [umol.cm-2.day-1]
 1186 2.1.3.13.3 ' Km' = 0.014 [umol.ml-1]
 1187 2.1.3.13.4 'minimal nutrient concentration' = 117 [umol.g-1]
 1188 2.1.3.13.5 'optimal nutrient concentration' = 234 [umol.g-1] (Silk et al. 1986) (Silk et al.
 1189 1986)
 1190 2.1.3.14 'reduction in respiration due to aerenchyma' [100%]=f{'aerenchymaFormation'}
 1191 [100%] x,y pairs :{0 0 0.3 0.7 0.6 1} (Fan et al. 2003)
 1192 'reduction in respiration due to cell size' [100%]=f{'cellSize'} [100%] x,y pairs :{101 0
 1193 150 0.07 200 0.14 250 0.17 300 0.25 350 0.32 400 0.37 450 0.43 500 0.51 533 0.57}
 1194 'reduction in respiration due to file number' [100%]=f{'filenumber'} [100%] x,y pairs :{17
 1195 0 16 0.05 15 0.13 14 0.17 13 0.23 12 0.3 11 0.35 10 0.42 9 0.49 8 0.52}
 1196 2.1.3.15 'relative carbon cost of exudation' [g.cm-1.day-1]=f{'age'} [day] x,y pairs :{0 5e-
 1197 06 100
 1198 1e-06}
 1199 2.1.3.16 'relative respiration' [g.g-1.day-1]=f{'age'} [day] x,y pairs :{0 0.09 2 0.04 6 0.04
 1200 1000
 1201 0.04} (Fan et al., 2003)
 1202 2.1.3.17 'root class id' = 98 [-]
 1203 2.1.3.18 'root hair density' [#.cm-2]=f{'age'} [day] x,y pairs :{0 2000 1 2000 2 2000 10
 1204 2000 30 0
 1205 2000 0} (Zhu et al. 2005; Mackay and S. Barber 1985)
 1206 2.1.3.19 'root hair diameter' = 0.0005 [cm]
 1207 2.1.3.20 'root hair length' [cm]=f{'age'} [day] x,y pairs :{0 0 1 0 2 0.028 2000 0.028}
 1208 2.4.15.20
 1209 (Zhu et al. 2005; Mackay and S. Barber 1985)
 1210 2.1.3.21 'soil impedance.v2' [cm]=f{'uniform distribution'} minimum=-0.05
 1211 maximum=0.05

1212 2.1.4 'finelateral2'

1213 2.1.4.1 'aerenchyma formation' [100%]=f{'age'} [day] x,y pairs :{0 0 3 0 5 0.1 10 0.25 20

1214 0.393 1000 0.393} (Fan et al. 2003)

1215 'cell size' [um²]={101 150 200 250 300 350 400 450 500 533} (Chimungu et al., 2014a)

1216 'cell file number' [#]={8 9 10 11 12 13 14 15 16 17}

1217 2.1.4.2 'branch list'

1218 2.1.4.3 'branching angle' = 62.83 [degrees]

1219 2.1.4.4 'density' = 0.094 [g.cm-3]

1220 2.1.4.5 'diameter' = 0.015 [cm]

1221 2.1.4.6 'gravitropism.v2' = 0 0 0 [cm]

1222 2.1.4.7 'growth rate' [cm.day-1]=f{'age'} [day] x,y pairs :{0 0.001 1 0.28 4 0 1000 0}

1223 2.1.4.8 'length root tip without xylem vessels' = 2 [cm]

1224 2.1.4.9 'longitudinal growth rate multiplier' [cm]=f{'normal distribution'} minimum=0.5

1225 maximum=1.5 mean=1 stdev=0.1

1226 2.1.4.10 'nitrate'

1227 2.1.4.10.1 'Cmin' = 0.0017 [umol.ml-1]

1228 2.1.4.10.2 'Imax' = 1.27 [umol.cm-2.day-1]

1229 2.1.4.10.3 'Km' = 0.0027 [umol.ml-1]

1230 2.1.4.10.4 'minimal nutrient concentration' = 600 [umol.g-1]

1231 2.1.4.10.5 'optimal nutrient concentration' = 1200 [umol.g-1]

1232 2.1.4.11 'number of xylem poles' = 4 [-]

1233 2.1.4.12 'phosphorus' (Barber 1995)

1234 2.1.4.12.1 'Cmin' = 0.0002 [umol.ml-1]

1235 2.1.4.12.2 'Imax' = 0.0555 [umol.cm-2.day-1]

1236 2.1.4.12.3 'Km' = 0.00545 [umol.ml-1]

1237 2.1.4.12.4 'minimal nutrient concentration' = 30 [umol.g-1]

1238 2.1.4.12.5 'optimal nutrient concentration' = 60 [umol.g-1]

1239 2.1.4.13 'potassium' (Barber 1995)

1240 2.1.4.13.1 'Cmin' = 0.002 [umol.ml-1]

1241 2.1.4.13.2 'Imax' = 0.467 [umol.cm-2.day-1]

1242 2.1.4.13.3 'Km' = 0.014 [umol.ml-1]

1243 2.1.4.13.4 'minimal nutrient concentration' = 117 [umol.g-1]

1244 2.1.4.13.5 'optimal nutrient concentration' = 234 [umol.g-1] (Silk et al. 1986)

1245 2.1.4.14 'reduction in respiration due to aerenchyma' [100%]=f{'aerenchymaFormation'}

1246 [100%] x,y pairs :{0 0 0.3 0.7 0.6 1} (Fan et al. 2003)

1247 'reduction in respiration due to cell size' [100%]=f{cellSize} [100%] x,y pairs :{101 0

1248 150 0.07 200 0.14 250 0.17 300 0.25 350 0.32 400 0.37 450 0.43 500 0.51 533 0.57}

1249 'reduction in respiration due to file number' [100%]=f{filenumber} [100%] x,y pairs :{17

1250 0 16 0.05 15 0.13 14 0.17 13 0.23 12 0.3 11 0.35 10 0.42 9 0.49 8 0.52}

1251 2.1.4.15 'relative carbon cost of exudation' [g.cm-1.day-1]=f{'age'} [day] x,y pairs :{0 5e-

1252 06 100

1253 1e-06}

1254 2.1.4.16 'relative respiration' [g.g-1.day-1]=f{'age'} [day] x,y pairs :{0 0.09 2 0.04 6 0.04

1255 1000

1256 0.04} (Fan et al., 2003)

1257 2.1.4.17 'root class id' = 98 [-]

1258 2.1.4.18 'root hair density' [#.cm-2]=f{'age'} [day] x,y pairs :{0 2000 1 2000 2 2000 10

1259 2000 30 0

1260 2000 0} (Zhu et al. 2005; Mackay and S. Barber 1985)

1261 2.1.4.19 'root hair diameter' = 0.0005 [cm]
1262 2.1.4.20 'root hair length' [cm]=f{'age'} [day] x,y pairs : {0 0 1 0 2 0.028 2000 0.028}
1263 2.4.15.20
1264 (Zhu et al. 2005; Mackay and S. Barber 1985)
1265 2.1.4.21 'soil impedance.v2' [cm]=f{'uniform distribution'} minimum=-0.05
1266 maximum=0.05
1267 2.1.5 'mesocotyl'
1268 2.1.5.1 'aerenchyma formation' [100%]=f{'age'} [day] x,y pairs : {0 0 100 0}
1269 'cell size' [um^2]={101 150 200 250 300 350 400 450 500 533} (Chimungu et al., 2014a)
1270 'cell file number' [#]={8 9 10 11 12 13 14 15 16 17}
1271 2.1.5.2 'branch list'
1272 2.1.5.2.1 'braceroots'
1273 2.1.5.2.1.1 'allometric scaling' = 1 [-]
1274 2.1.5.2.1.2 'branching spatial offset' = 4 [cm]
1275 2.1.5.2.1.3 'branching time offset' = 25 [day]
1276 2.1.5.2.1.4 'number of branches/whorl' = 14 [#]
1277 2.1.5.2.2 'braceroots2'
1278 2.1.5.2.2.1 'allometric scaling' = 1 [-]
1279 2.1.5.2.2.2 'branching delay' = 14 [day]
1280 2.1.5.2.2.3 'branching frequency' = 5 [cm]
1281 2.1.5.2.2.4 'branching spatial offset' = 7 [cm]
1282 2.1.5.2.2.5 'branching time offset' = 36 [day]
1283 2.1.5.2.2.6 'number of branches/whorl' = 20 [#]
1284 2.1.5.2.3 'nodalroots'
1285 2.1.5.2.3.1 'branching spatial offset' = 1.5 [cm]
1286 2.1.5.2.3.2 'branching time offset' = 9 [day]
1287 2.1.5.2.3.3 'number of branches/whorl' = 3 [#]
1288 2.1.5.2.4 'nodalroots2'
1289 2.1.5.2.4.1 'allometric scaling' = 1 [-]
1290 2.1.5.2.4.2 'branching spatial offset' = 1.9 [cm]
1291 2.1.5.2.4.3 'branching time offset' = 16 [day]
1292 2.1.5.2.4.4 'number of branches/whorl' = 4 [#]
1293 2.1.5.2.5 'nodalroots3'
1294 2.1.5.2.5.1 'allometric scaling' = 1 [-]
1295 2.1.5.2.5.2 'branching spatial offset' = 2.1 [cm]
1296 2.1.5.2.5.3 'branching time offset' = 20 [day]
1297 2.1.5.2.5.4 'number of branches/whorl' = 5 [#]
1298 2.1.5.2.6 'nodalroots4'
1299 2.1.5.2.6.1 'allometric scaling' = 1 [-]
1300 2.1.5.2.6.2 'branching spatial offset' = 2.3 [cm]
1301 2.1.5.2.6.3 'branching time offset' = 23 [day]
1302 2.1.5.2.6.4 'number of branches/whorl' = 6 [#]
1303 2.1.5.3 'density' = 0.094 [g.cm-3]
1304 2.1.5.4 'diameter' = 0.15 [cm]
1305 2.1.5.5 'gravitropism' = -1 [-]
1306 2.1.5.6 'gravitropism.v2' [cm]=f{'uniform distribution'} minimum=0.5 maximum=1
1307 2.1.5.7 'growth rate' [cm.day-1]=f{'age'} [day] x,y pairs : {0 1 1 2 2 2 3 2 5 0 1000 0}
1308 2.1.5.8 'length root tip without xylem vessels' = 2 [cm]
1309 2.1.5.9 'nitrate'

1310 2.1.5.9.1 'Cmin' = 0 [umol.ml-1]
 1311 2.1.5.9.2 'Imax' = 0 [umol.cm-2.day-1]
 1312 2.1.5.9.3 'Km' = 1 [umol.ml-1]
 1313 2.1.5.9.4 'minimal nutrient concentration' = 600 [umol.g-1]
 1314 2.1.5.9.5 'optimal nutrient concentration' = 1200 [umol.g-1]
 1315 2.1.5.10 'number of xylem poles' = 61 [-]
 1316 2.1.5.11 'phosphorus' (Barber 1995)
 1317 2.1.5.11.1 'Cmin' = 0.0002 [umol.ml-1]
 1318 2.1.5.11.2 'Imax' = 0.0555 [umol.cm-2.day-1]
 1319 2.1.5.11.3 'Km' = 0.00545 [umol.ml-1]
 1320 2.1.5.11.4 'minimal nutrient concentration' = 30 [umol.g-1]
 1321 2.1.5.11.5 'optimal nutrient concentration' = 60 [umol.g-1]
 1322 2.1.5.12 'potassium' (Barber 1995)
 1323 2.1.5.12.1 'Cmin' = 0.002 [umol.ml-1]
 1324 2.1.5.12.2 'Imax' = 0.467 [umol.cm-2.day-1]
 1325 2.1.5.12.3 'Km' = 0.014 [umol.ml-1]
 1326 2.1.5.12.4 'minimal nutrient concentration' = 117 [umol.g-1]
 1327 2.1.5.12.5 'optimal nutrient concentration' = 234 [umol.g-1] (Silk et al. 1986)
 1328 2.1.5.13 'reduction in respiration due to aerenchyma' [100%]=f{'aerenchymaFormation'}
 1329 [100%]
 1330 x,y pairs :{0 0 0.3 0.7 0.6 1} (Fan et al. 2003)
 1331 'reduction in respiration due to cell size' [100%]=f{cellSize} [100%] x,y pairs :{101 0
 1332 150 0.07 200 0.14 250 0.17 300 0.25 350 0.32 400 0.37 450 0.43 500 0.51 533 0.57}
 1333 'reduction in respiration due to file number' [100%]=f{filenumber} [100%] x,y pairs :{17
 1334 0 16 0.05 15 0.13 14 0.17 13 0.23 12 0.3 11 0.35 10 0.42 9 0.49 8 0.52}
 1335 2.1.5.14 'relative carbon cost of exudation' [g.cm-1.day-1]=f{'age'} [day] x,y pairs :{0 0
 1336 100 0}
 1337 2.1.5.15 'relative respiration' [g.g-1.day-1]=f{'age'} [day] x,y pairs :{0 0.09 2 0.04 6 0.04
 1338 1000
 1339 0.04} (Fan et al., 2003)
 1340 2.1.5.16 'root class id' = 97 [-]
 1341 2.1.5.17 'root hair density' [#.cm-2]=f{'age'} [day] x,y pairs :{0 0 2000 0}
 1342 2.1.5.18 'root hair diameter' = 0.0005 [cm]
 1343 2.1.5.19 'root hair length' [cm]=f{'age'} [day] x,y pairs :{0 0 1 0 2 0.028 2000 0.028}
 1344 2.4.15.20
 1345 (Zhu et al. 2005; Mackay and S. Barber 1985)
 1346 2.1.5.20 'soil impedance' = 0.3 [-]
 1347 2.1.5.21 'soil impedance.v2' [cm]=f{'uniform distribution'} minimum=-0.3 maximum=0.3
 1348 2.1.5.22 'top boundary' = 0 [-]
 1349 2.1.6 'lateral'
 1350 2.1.6.1 'aerenchyma formation' [100%]=f{'age'} [day] x,y pairs :{0 0 3 0 5 0.1 10 0.25 20
 1351 0.3931000 0.393} (Fan et al. 2003)
 1352 'cell size' [um^2]={101 150 200 250} (Chimungu et al., 2014a)
 1353 'cell file number' [#]={4 5 6}
 1354 2.1.6.2 'bottom boundary' = 1 [-]
 1355 2.1.6.3 'branch list'
 1356 2.1.6.3.1 'finelateral'
 1357 2.1.6.3.1.1 'allow branches to form above ground' = 0 [-]
 1358 2.1.6.3.1.2 'branching frequency' [cm]=f{'uniform distribution'} minimum=0.15

1359 maximum=0.35
 1360 2.1.6.3.1.3 'length root tip' = 4 [cm]
 1361 2.1.6.4 'branching angle' = 90 [degrees]
 1362 2.1.6.5 'density' = 0.094 [g.cm-3]
 1363 2.1.6.6 'diameter' = 0.04 [cm]
 1364 2.1.6.7 'gravitropism.v2' = 0 0 0 [cm]
 1365 2.1.6.8 'growth rate' [cm.day-1]=f{'age'} [day] x,y pairs :{0 0.01 1 0.2 3 0.4 7 1 11 0
 1366 1000 0}
 1367 2.1.6.9 'length root tip without xylem vessels' = 2 [cm]
 1368 2.1.6.10 'longitudinal growth rate multiplier' [cm]=f{'normal distribution'} minimum=0.1
 1369 maximum=2 mean=0.7 stdev=0.3
 1370 2.1.6.11 'nitrate'
 1371 2.1.6.11.1 'Cmin' = 0.0017 [umol.ml-1]
 1372 2.1.6.11.2 'Imax' = 1.27 [umol.cm-2.day-1]
 1373 2.1.6.11.3 'Km' = 0.0027 [umol.ml-1]
 1374 2.1.6.11.4 'minimal nutrient concentration' = 600 [umol.g-1]
 1375 2.1.6.11.5 'optimal nutrient concentration' = 1200 [umol.g-1]
 1376 2.1.6.12 'number of xylem poles' = 4 [-]
 1377 2.1.6.13 'phosphorus' (Barber 1995)
 1378 2.1.6.13.1 'Cmin' = 0.0002 [umol.ml-1]
 1379 2.1.6.13.2 'Imax' = 0.0555 [umol.cm-2.day-1]
 1380 2.1.6.13.3 'Km' = 0.00545 [umol.ml-1]
 1381 2.1.6.13.4 'minimal nutrient concentration' = 30 [umol.g-1]
 1382 2.1.6.13.5 'optimal nutrient concentration' = 60 [umol.g-1]
 1383 2.1.6.14 'potassium' (Barber 1995)
 1384 2.1.6.14.1 'Cmin' = 0.002 [umol.ml-1]
 1385 2.1.6.14.2 'Imax' = 0.467 [umol.cm-2.day-1]
 1386 2.1.6.14.3 'Km' = 0.014 [umol.ml-1]
 1387 2.1.6.14.4 'minimal nutrient concentration' = 117 [umol.g-1]
 1388 2.1.6.14.5 'optimal nutrient concentration' = 234 [umol.g-1] (Silk et al. 1986)
 1389 2.1.6.15 'reduction in respiration due to aerenchyma' [100%]=f{'aerenchymaFormation'}
 1390 [100%] x,y pairs :{0 0 0.3 0.7 0.6 1} (Fan et al. 2003)
 1391 'reduction in respiration due to cell size' [100%]=f{'cellSize'} [100%] x,y pairs :{101 0
 1392 150 0.07 200 0.14 250 0.17 300 0.25 350 0.32 400 0.37 450 0.43 500 0.51 533 0.57}
 1393 'reduction in respiration due to file number' [100%]=f{'filenumber'} [100%] x,y pairs :{17
 1394 0 16 0.05 15 0.13 14 0.17 13 0.23 12 0.3 11 0.35 10 0.42 9 0.49 8 0.52}
 1395 2.1.6.16 'relative carbon cost of exudation' [g.cm-1.day-1]=f{'age'} [day] x,y pairs :{0 5e-
 1396 06 100
 1397 3e-06}
 1398 2.1.6.17 'relative respiration' [g.g-1.day-1]=f{'age'} [day] x,y pairs :{0 0.09 2 0.04 6 0.04
 1399 1000
 1400 0.04} (Fan et al., 2003)
 1401 2.1.6.18 'root class id' = 98 [-]
 1402 2.1.6.19 'root hair density' [#.cm-2]=f{'age'} [day] x,y pairs :{0 2000 1 2000 2 2000 10
 1403 2000 30 0
 1404 2000 0} (Zhu et al. 2005; Mackay and S. Barber 1985)
 1405 2.1.6.20 'root hair diameter' = 0.0005 [cm]
 1406 2.1.6.21 'root hair length' [cm]=f{'age'} [day] x,y pairs :{0 0 1 0 2 0.028 2000 0.028}
 1407 2.4.15.20
 (Zhu et al. 2005; Mackay and S. Barber 1985)

1408 2.1.6.22 'soil impedance.v2' [cm]=f{'uniform distribution'} minimum=-0.1 maximum=0.1
1409 2.1.6.23 'top boundary' = 1 [-]
1410 2.1.7 'lateral of crown roots'
1411 2.1.7.1 'aerenchyma formation' [100%]=f{'age'} [day] x,y pairs :{0 0 3 0 5 0.1 10 0.25 20
1412 0.393 1000 0.393} (Fan et al. 2003)
1413 'cell size' [um^2]={101 150 200 250 300 350 400 450 500 533} (Chimungu et al., 2014a)
1414 'cell file number' [#]={8 9 10 11 12 13 14 15 16 17}
1415 2.1.7.2 'branch list'
1416 2.1.7.2.1 'lateral'
1417 2.1.7.2.1.1 'allow branches to form above ground' = 0 [-]
1418 2.1.7.2.1.2 'branching frequency' [cm]=f{'uniform distribution'} minimum=0.25
1419 maximum=0.35
1420 2.1.7.2.1.3 'length root tip' = 5 [cm]
1421 2.1.7.3 'branching angle' = 90 [degrees]
1422 2.1.7.4 'density' = 0.094 [g.cm-3]
1423 2.1.7.5 'diameter' = 0.07 [cm]
1424 2.1.7.6 'gravitropism' = 0 [-]
1425 2.1.7.7 'gravitropism.v2' = 0 0 0 [cm]
1426 2.1.7.8 'growth rate' [cm.day-1]=f{'age'} [day] x,y pairs :{0 0.1 1 0.5 3 1.2 12 1.2 18 0
1427 1000 0}
1428 2.1.7.9 'length root tip without xylem vessels' = 2 [cm]
1429 2.1.7.10 'longitudinal growth rate multiplier' [cm]=f{' normal distribution'} minimum=0.1
1430 maximum=1 mean=0.4 stdev=0.3
1431 2.1.7.11 'nitrate'
1432 2.1.7.11.1 ' Cmin' = 0.0017 [umol.ml-1]
1433 2.1.7.11.2 ' Imax' = 1.27 [umol.cm-2.day-1]
1434 2.1.7.11.3 ' Km' = 0.0027 [umol.ml-1]
1435 2.1.7.11.4 'minimal nutrient concentration' = 600 [umol.g-1]
1436 2.1.7.11.5 'optimal nutrient concentration' = 1200 [umol.g-1]
1437 2.1.7.12 'number of xylem poles' = 4 [-]
1438 2.1.7.13 'phosphorus' (Barber 1995)
1439 2.1.7.13.1 ' Cmin' = 0.0002 [umol.ml-1]
1440 2.1.7.13.2 ' Imax' = 0.0555 [umol.cm-2.day-1]
1441 2.1.7.13.3 ' Km' = 0.00545 [umol.ml-1]
1442 2.1.7.13.4 'minimal nutrient concentration' = 30 [umol.g-1]
1443 2.1.7.13.5 'optimal nutrient concentration' = 60 [umol.g-1]
1444 2.1.7.14 'potassium' (Barber 1995)
1445 2.1.7.14.1 ' Cmin' = 0.002 [umol.ml-1]
1446 2.1.7.14.2 ' Imax' = 0.467 [umol.cm-2.day-1]
1447 2.1.7.14.3 ' Km' = 0.014 [umol.ml-1]
1448 2.1.7.14.4 'minimal nutrient concentration' = 117 [umol.g-1]
1449 2.1.7.14.5 'optimal nutrient concentration' = 234 [umol.g-1] (Silk et al. 1986)
1450 2.1.7.15 'reduction in respiration due to aerenchyma' [100%]=f{'aerenchymaFormation'}
1451 [100%] x,y pairs :{0 0 0.3 0.7 0.6 1} (Fan et al. 2003)
1452 'reduction in respiration due to cell size' [100%]=f{'cellSize'} [100%] x,y pairs :{101 0
1453 150 0.07 200 0.14 250 0.17 300 0.25 350 0.32 400 0.37 450 0.43 500 0.51 533 0.57}
1454 'reduction in respiration due to file number' [100%]=f{'filenumber'} [100%] x,y pairs :{17
1455 0 16 0.05 15 0.13 14 0.17 13 0.23 12 0.3 11 0.35 10 0.42 9 0.49 8 0.52}

1456 2.1.7.16 'relative carbon cost of exudation' [g.cm-1.day-1]=f{'age'} [day] x,y pairs :{0 5e-
1457 06 100
1458 4e-06}
1459 2.1.7.17 'relative respiration' [g.g-1.day-1]=f{'age'} [day] x,y pairs :{0 0.09 2 0.04 6 0.04
1460 1000
1461 0.04} (Fan et al., 2003)
1462 2.1.7.18 'root class id' = 98 [-]
1463 2.1.7.19 'root hair density' [#.cm-2]=f{'age'} [day] x,y pairs :{0 2000 1 2000 2 2000 10
1464 2000 30 0
1465 2000 0} (Zhu et al. 2005; Mackay and S. Barber 1985)
1466 2.1.7.20 'root hair diameter' = 0.0005 [cm]
1467 2.1.7.21 'root hair length' [cm]=f{'age'} [day] x,y pairs :{0 0 1 0 2 0.028 2000 0.028}
1468 2.4.15.20
1469 (Zhu et al. 2005; Mackay and S. Barber 1985)
1470 2.1.7.22 'soil impedance' = 0.05 [-]
1471 2.1.7.23 'soil impedance.v2' [cm]=f{'uniform distribution'} minimum=-0.05
1472 maximum=0.05
1473 2.1.8 'nodalroots'
1474 2.1.8.1 'aerenchyma formation' [100%]=f{'age'} [day] x,y pairs :{0 0 3 0 5 0.1 10 0.25 20
1475 0.393 1000 0.393} (Fan et al. 2003)
1476 'cell size' [um^2]={101 150 200 250 300 350 400 450 500 533} (Chimungu et al., 2014a)
1477 'cell file number' [#]={8 9 10 11 12 13 14 15 16 17}
1478 2.1.8.2 'branch list'
1479 2.1.8.2.1 'lateral'
1480 2.1.8.2.1.1 'allow branches to form above ground' = 0 [-]
1481 2.1.8.2.1.2 'branching frequency' [cm]=f{'uniform distribution'} minimum=0.1
1482 maximum=0.3
1483 2.1.8.2.1.3 'length root tip' = 10.93 [cm]
1484 2.1.8.3 'branching angle' = 160 [degrees]
1485 2.1.8.4 'density' = 0.094 [g.cm-3]
1486 2.1.8.5 'diameter' [cm]=f{'age'} [day] x,y pairs :{0 0.12 10 0.09 100 0.09}
1487 2.1.8.6 'gravitropism.v2' [cm]=f{'uniform distribution'} minimum=-0.01 maximum=-
1488 0.005
1489 2.1.8.7 'growth rate' [cm.day-1]=f{'age'} [day] x,y pairs :{0 0.01 1 1 3 4.5 28 4.5 38 0
1490 1000 0}
1491 2.1.8.8 'length root tip without xylem vessels' = 2 [cm]
1492 2.1.8.9 'longitudinal growth rate multiplier' [cm]=f{'normal distribution'} minimum=0.6
1493 maximum=1.2 mean=1 stdev=0.1
1494 2.1.8.10 'nitrate'
1495 2.1.8.10.1 'Cmin' = 0.001 [umol.ml-1]
1496 2.1.8.10.2 'Imax' [umol.cm-2.day-1]=f{'age'} [day] x,y pairs :{0 1.21 2 2.1 40 2.1}
1497 2.1.8.10.3 'Km' [umol.ml-1]=f{'age'} [day] x,y pairs :{0 0.0157 2 0.0522 40 0.0522}
1498 2.1.8.10.4 'minimal nutrient concentration' = 600 [umol.g-1]
1499 2.1.8.10.5 'optimal nutrient concentration' = 1200 [umol.g-1]
1500 2.1.8.11 'number of xylem poles' = 10 [-]
1501 2.1.8.12 'phosphorus' (Barber 1995)
1502 2.1.8.12.1 'Cmin' = 0.0002 [umol.ml-1]
1503 2.1.8.12.2 'Imax' = 0.0555 [umol.cm-2.day-1]
1504 2.1.8.12.3 'Km' = 0.00545 [umol.ml-1]

1505 2.1.8.12.4 'minimal nutrient concentration' = 30 [umol.g-1]
1506 2.1.8.12.5 'optimal nutrient concentration' = 60 [umol.g-1]
1507 2.1.8.13 'potassium' (Barber 1995)
1508 2.1.8.13.1 'Cmin' = 0.002 [umol.ml-1]
1509 2.1.8.13.2 'Imax' = 0.467 [umol.cm-2.day-1]
1510 2.1.8.13.3 'Km' = 0.014 [umol.ml-1]
1511 2.1.8.13.4 'minimal nutrient concentration' = 117 [umol.g-1]
1512 2.1.8.13.5 'optimal nutrient concentration' = 234 [umol.g-1] (Silk et al. 1986)
1513 2.1.8.14 'reduction in respiration due to aerenchyma' [100%]=f{'aerenchymaFormation'}
1514 [100%] x,y pairs : {0 0 0.3 0.7 0.6 1} (Fan et al. 2003)
1515 'reduction in respiration due to cell size' [100%]=f{cellSize} [100%] x,y pairs : {101 0
1516 150 0.07 200 0.14 250 0.17 300 0.25 350 0.32 400 0.37 450 0.43 500 0.51 533 0.57}
1517 'reduction in respiration due to file number' [100%]=f{fileNumber} [100%] x,y pairs : {17
1518 0 16 0.05 15 0.13 14 0.17 13 0.23 12 0.3 11 0.35 10 0.42 9 0.49 8 0.52}
1519 2.1.8.15 'regular topology' = 3 [-]
1520 2.1.8.16 'relative carbon cost of exudation' [g.cm-1.day-1]=f{'age'} [day] x,y pairs : {0 5e-
1521 06 100
1522 5e-06} (Groleau-Renaud et al. 1998)
1523 2.1.8.17 'relative respiration' [g.g-1.day-1]=f{'age'} [day] x,y pairs : {0 0.09 2 0.04 6 0.04
1524 1000
1525 0.04} (Fan et al., 2003)
1526 2.1.8.18 'root class id' = 101 [-]
1527 2.1.8.19 'root hair density' [#.cm-2]=f{'age'} [day] x,y pairs : {0 2000 1 2000 2 2000 10
1528 2000 30 0
1529 2000 0} (Zhu et al. 2005; Mackay and S. Barber 1985)
1530 2.1.8.20 'root hair diameter' = 0.0005 [cm]
1531 2.1.8.21 'root hair length' [cm]=f{'age'} [day] x,y pairs : {0 0 1 0 2 0.028 2000 0.028}
1532 2.4.15.20
1533 (Zhu et al. 2005; Mackay and S. Barber 1985)
1534 2.1.8.22 'soil impedance.v2' [cm]=f{'uniform distribution'} minimum=-0.02
1535 maximum=0.02
1536 2.1.8.23 'topology offset' = 0 [-]
1537 2.1.9 'nodalroots2'
1538 2.1.9.1 'aerenchyma formation' [100%]=f{'age'} [day] x,y pairs : {0 0 3 0 5 0.1 10 0.25 20
1539 0.393 1000 0.393} (Fan et al. 2003)
1540 'cell size' [um^2]={101 150 200 250 300 350 400 450 500 533} (Chimungu et al., 2014a)
1541 'cell file number' [#]={8 9 10 11 12 13 14 15 16 17}
1542 2.1.9.2 'branch list'
1543 2.1.9.2.1 'lateral'
1544 2.1.9.2.1.1 'allow branches to form above ground' = 0 [-]
1545 2.1.9.2.1.2 'branching frequency' [cm]=f{'uniform distribution'} minimum=0.1
1546 maximum=0.3
1547 2.1.9.2.1.3 'length root tip' = 10.93 [cm]
1548 2.1.9.3 'branching angle' = 150 [degrees]
1549 2.1.9.4 'density' = 0.094 [g.cm-3]
1550 2.1.9.5 'diameter' [cm]=f{'age'} [day] x,y pairs : {0 0.14 10 0.09 100 0.09}
1551 2.1.9.6 'gravitropism.v2' [cm]=f{'uniform distribution'} minimum=-0.01 maximum=-
1552 0.005

1553 2.1.9.7 'growth rate' [cm.day-1]=f{'age'} [day] x,y pairs :{0 0.01 1 1 3 4.5 28 4.5 38 0
1554 1000 0}
1555 2.1.9.8 'length root tip without xylem vessels' = 2 [cm]
1556 2.1.9.9 'longitudinal growth rate multiplier' [cm]=f{'normal distribution'} minimum=0.6
1557 maximum=1.2 mean=1 stdev=0.1
1558 2.1.9.10 'nitrate'
1559 2.1.9.10.1 ' Cmin' = 0.001 [umol.ml-1]
1560 2.1.9.10.2 ' Imax' [umol.cm-2.day-1]=f{'age'} [day] x,y pairs :{0 1.21 2 2.1 40 2.1}
1561 2.1.9.10.3 ' Km' [umol.ml-1]=f{'age'} [day] x,y pairs :{0 0.0157 2 0.0522 40 0.0522}
1562 2.1.9.10.4 'minimal nutrient concentration' = 600 [umol.g-1]
1563 2.1.9.10.5 'optimal nutrient concentration' = 1200 [umol.g-1]
1564 2.1.9.11 'number of xylem poles' = 18 [-]
1565 2.1.9.12 'phosphorus' (Barber 1995)
1566 2.1.9.12.1 ' Cmin' = 0.0002 [umol.ml-1]
1567 2.1.9.12.2 ' Imax' = 0.0555 [umol.cm-2.day-1]
1568 2.1.9.12.3 ' Km' = 0.00545 [umol.ml-1]
1569 2.1.9.12.4 'minimal nutrient concentration' = 30 [umol.g-1]
1570 2.1.9.12.5 'optimal nutrient concentration' = 60 [umol.g-1]
1571 2.1.9.13 'potassium' (Barber 1995)
1572 2.1.9.13.1 ' Cmin' = 0.002 [umol.ml-1]
1573 2.1.9.13.2 ' Imax' = 0.467 [umol.cm-2.day-1]
1574 2.1.9.13.3 ' Km' = 0.014 [umol.ml-1]
1575 2.1.9.13.4 'minimal nutrient concentration' = 117 [umol.g-1]
1576 2.1.9.13.5 'optimal nutrient concentration' = 234 [umol.g-1] (Silk et al. 1986)
1577 2.1.9.14 'reduction in respiration due to aerenchyma' [100%]=f{'aerenchymaFormation'}
1578 [100%] x,y pairs :{0 0 0.3 0.7 0.6 1} (Fan et al. 2003)
1579 'reduction in respiration due to cell size' [100%]=f{'cellSize'} [100%] x,y pairs :{101 0
1580 150 0.07 200 0.14 250 0.17 300 0.25 350 0.32 400 0.37 450 0.43 500 0.51 533 0.57}
1581 'reduction in respiration due to file number' [100%]=f{'filenumber'} [100%] x,y pairs :{17
1582 0 16 0.05 15 0.13 14 0.17 13 0.23 12 0.3 11 0.35 10 0.42 9 0.49 8 0.52}
1583 2.1.9.15 'regular topology' = 0 [-]
1584 2.1.9.16 'relative carbon cost of exudation' [g.cm-1.day-1]=f{'age'} [day] x,y pairs :{0 5e-
1585 06 100
1586 5e-06} (Groleau-Renaud et al. 1998)
1587 2.1.9.17 'relative respiration' [g.g-1.day-1]=f{'age'} [day] x,y pairs :{0 0.09 2 0.04 6 0.04
1588 1000 0.04} (Fan et al., 2003)
1589 2.1.9.18 'root class id' = 101 [-]
1590 2.1.9.19 'root hair density' [#.cm-2]=f{'age'} [day] x,y pairs :{0 2000 1 2000 2 2000 10
1591 2000 30 0
1592 2000 0} (Zhu et al. 2005; Mackay and S. Barber 1985)
1593 2.1.9.20 'root hair diameter' = 0.0005 [cm]
1594 2.1.9.21 'root hair length' [cm]=f{'age'} [day] x,y pairs :{0 0 1 0 2 0.028 2000 0.028}
1595 2.4.15.20
1596 (Zhu et al. 2005; Mackay and S. Barber 1985)
1597 2.1.9.22 'soil impedance.v2' [cm]=f{'uniform distribution'} minimum=-0.02
1598 maximum=0.02
1599 2.1.9.23 'topology offset' = 0 [-]
1600 2.1.10 'nodalroots3'

1601 2.1.10.1 'aerenchyma formation' [100%]=f{'age'} [day] x,y pairs :{0 0 3 0 5 0.1 10 0.25
1602 20 0.393 1000 0.393} (Fan et al. 2003)
1603 'cell size' [um²]={101 150 200 250 300 350 400 450 500 533} (Chimungu et al., 2014a)
1604 'cell file number' [#]={8 9 10 11 12 13 14 15 16 17}
1605 2.1.10.2 'branch list'
1606 2.1.10.2.1 'lateral'
1607 2.1.10.2.1.1 'allow branches to form above ground' = 0 [-]
1608 2.1.10.2.1.2 'branching frequency' [cm]=f{'uniform distribution'} minimum=0.1
1609 maximum=0.3
1610 2.1.10.2.1.3 'length root tip' = 10.93 [cm] 2.1.10.3 'branching angle' = 140 [degrees]
1611 2.1.10.4 'density' = 0.094 [g.cm-3]
1612 2.1.10.5 'diameter' [cm]=f{'age'} [day] x,y pairs :{0 0.16 10 0.1 100 0.1}
1613 2.1.10.6 'gravitropism.v2' [cm]=f{'uniform distribution'} minimum=-0.01 maximum=-
1614 0.005
1615 2.1.10.7 'growth rate' [cm.day-1]=f{'age'} [day] x,y pairs :{0 0.01 1 1 3 4.5 28 4.5 38 0
1616 1000 0}
1617 2.1.10.8 'length root tip without xylem vessels' = 2 [cm]
1618 2.1.10.9 'longitudinal growth rate multiplier' [cm]=f{'normal distribution'} minimum=0.6
1619 maximum=1.2 mean=1 stdev=0.1
1620 2.1.10.10 'nitrate'
1621 2.1.10.10.1 'Cmin' = 0.001 [umol.ml-1]
1622 2.1.10.10.2 'Imax' [umol.cm-2.day-1]=f{'age'} [day] x,y pairs :{0 1.21 2 2.1 40 2.1}
1623 2.1.10.10.3 'Km' [umol.ml-1]=f{'age'} [day] x,y pairs :{0 0.0157 2 0.0522 40 0.0522}
1624 2.1.10.10.4 'minimal nutrient concentration' = 600 [umol.g-1]
1625 2.1.10.10.5 'optimal nutrient concentration' = 1200 [umol.g-1]
1626 2.1.10.11 'number of xylem poles' = 24 [-]
1627 2.1.10.12 'phosphorus' (Barber 1995)
1628 2.1.10.12.1 'Cmin' = 0.0002 [umol.ml-1]
1629 2.1.10.12.2 'Imax' = 0.0555 [umol.cm-2.day-1]
1630 2.1.10.12.3 'Km' = 0.00545 [umol.ml-1]
1631 2.1.10.12.4 'minimal nutrient concentration' = 30 [umol.g-1]
1632 2.1.10.12.5 'optimal nutrient concentration' = 60 [umol.g-1]
1633 2.1.10.13 'potassium' (Barber 1995)
1634 2.1.10.13.1 'Cmin' = 0.002 [umol.ml-1]
1635 2.1.10.13.2 'Imax' = 0.467 [umol.cm-2.day-1]
1636 2.1.10.13.3 'Km' = 0.014 [umol.ml-1]
1637 2.1.10.13.4 'minimal nutrient concentration' = 117 [umol.g-1]
1638 2.1.10.13.5 'optimal nutrient concentration' = 234 [umol.g-1] (Silk et al. 1986)
1639 2.1.10.14 'reduction in respiration due to aerenchyma' [100%]=f{'aerenchymaFormation'}
1640 [100%] x,y pairs :{0 0 0.3 0.7 0.6 1} (Fan et al. 2003)
1641 'reduction in respiration due to cell size' [100%]=f{'cellSize'} [100%] x,y pairs :{101 0
1642 150 0.07 200 0.14 250 0.17 300 0.25 350 0.32 400 0.37 450 0.43 500 0.51 533 0.57}
1643 'reduction in respiration due to file number' [100%]=f{'filenumber'} [100%] x,y pairs :{17
1644 0 16 0.05 15 0.13 14 0.17 13 0.23 12 0.3 11 0.35 10 0.42 9 0.49 8 0.52}
1645 2.1.10.15 'regular topology' = 0 [-]
1646 2.1.10.16 'relative carbon cost of exudation' [g.cm-1.day-1]=f{'age'} [day] x,y pairs :{0
1647 5e-06 100
1648 5e-06} (Groleau-Renaud et al. 1998)

1649 2.1.10.17 'relative respiration' [g.g-1.day-1]=f{'age'} [day] x,y pairs :{0 0.09 2 0.04 6
1650 0.04 1000
1651 0.04} (Fan et al., 2003)
1652 2.1.10.18 'root class id' = 101 [-]
1653 2.1.10.19 'root hair density' [#.cm-2]=f{'age'} [day] x,y pairs :{0 2000 1 2000 2 2000 10
1654 2000 30 0
1655 2000 0} (Zhu et al. 2005; Mackay and S. Barber 1985)
1656 2.1.10.20 'root hair diameter' = 0.0005 [cm]
1657 2.1.10.21 'root hair length' [cm]=f{'age'} [day] x,y pairs :{0 0 1 0 2 0.028 2000 0.028}
1658 2.4.15.20
1659 (Zhu et al. 2005; Mackay and S. Barber 1985)
1660 2.1.10.22 'soil impedance.v2' [cm]=f{'uniform distribution'} minimum=-0.02
1661 maximum=0.02
1662 2.1.10.23 'topology offset' = 0 [-]
1663 2.1.11 'nodalroots4'
1664 2.1.11.1 'aerenchyma formation' [100%]=f{'age'} [day] x,y pairs :{0 0 3 0 5 0.1 10 0.25
1665 20 0.393 1000 0.393} (Fan et al. 2003)
1666 'cell size' [um^2]={101 150 200 250 300 350 400 450 500 533} (Chimungu et al., 2014a)
1667 'cell file number' [#]={8 9 10 11 12 13 14 15 16 17}
1668 2.1.11.2 'branch list'
1669 2.1.11.2.1 'lateral'
1670 2.1.11.2.1.1 'allow branches to form above ground' = 0 [-]
1671 2.1.11.2.1.2 'branching frequency' [cm]=f{'uniform distribution'} minimum=0.1
1672 maximum=0.3
1673 2.1.11.2.1.3 'length root tip' = 10.93 [cm]
1674 2.1.11.3 'branching angle' = 130 [degrees]
1675 2.1.11.4 'density' = 0.094 [g.cm-3]
1676 2.1.11.5 'diameter' [cm]=f{'age'} [day] x,y pairs :{0 0.2 10 0.11 100 0.11}
1677 2.1.11.6 'gravitropism.v2' [cm]=f{'uniform distribution'} minimum=-0.01 maximum=-
1678 0.005
1679 2.1.11.7 'growth rate' [cm.day-1]=f{'age'} [day] x,y pairs :{0 0.01 1 1 3 4.5 28 4.5 38 0
1680 1000 0}
1681 2.1.11.8 'length root tip without xylem vessels' = 2 [cm]
1682 2.1.11.9 'longitudinal growth rate multiplier' [cm]= f{'normal distribution'} minimum=0.6
1683 maximum=1.2 mean=1 stdev=0.1
1684 2.1.11.10 'nitrate'
1685 2.1.11.10.1 'Cmin' = 0.001 [umol.ml-1]
1686 2.1.11.10.2 'Imax' [umol.cm-2.day-1]=f{'age'} [day] x,y pairs :{0 1.21 2 2.1 40 2.1}
1687 2.1.11.10.3 'Km' [umol.ml-1]=f{'age'} [day] x,y pairs :{0 0.0157 2 0.0522 40 0.0522}
1688 2.1.11.10.4 'minimal nutrient concentration' = 600 [umol.g-1]
1689 2.1.11.10.5 'optimal nutrient concentration' = 1200 [umol.g-1]
1690 2.1.11.11 'number of xylem poles' = 32 [-]
1691 2.1.11.12 'phosphorus' (Barber 1995)
1692 2.1.11.12.1 'Cmin' = 0.0002 [umol.ml-1]
1693 2.1.11.12.2 'Imax' = 0.0555 [umol.cm-2.day-1]
1694 2.1.11.12.3 'Km' = 0.00545 [umol.ml-1]
1695 2.1.11.12.4 'minimal nutrient concentration' = 30 [umol.g-1]
1696 2.1.11.12.5 'optimal nutrient concentration' = 60 [umol.g-1]
1697 2.1.11.13 'potassium' (Barber 1995)

1698 2.1.11.13.1 'Cmin' = 0.002 [umol.ml-1]
1699 2.1.11.13.2 'Imax' = 0.467 [umol.cm-2.day-1]
1700 2.1.11.13.3 'Km' = 0.014 [umol.ml-1]
1701 2.1.11.13.4 'minimal nutrient concentration' = 117 [umol.g-1]
1702 2.1.11.13.5 'optimal nutrient concentration' = 234 [umol.g-1] (Silk et al. 1986)
1703 2.1.11.14 'reduction in respiration due to aerenchyma' [100%]=f{'aerenchymaFormation'}
1704 [100%] x,y pairs :{0 0 0.3 0.7 0.6 1} (Fan et al. 2003)
1705 'reduction in respiration due to cell size' [100%]=f{'cellSize'} [100%] x,y pairs :{101 0
1706 150 0.07 200 0.14 250 0.17 300 0.25 350 0.32 400 0.37 450 0.43 500 0.51 533 0.57}
1707 'reduction in respiration due to file number' [100%]=f{'fileNumber'} [100%] x,y pairs :{17
1708 0 16 0.05 15 0.13 14 0.17 13 0.23 12 0.3 11 0.35 10 0.42 9 0.49 8 0.52}
1709 2.1.11.15 'relative carbon cost of exudation' [g.cm-1.day-1]=f{'age'} [day] x,y pairs :{0
1710 5e-06 100
1711 5e-06} (Groleau-Renaud et al. 1998)
1712 2.1.11.16 'relative respiration' [g.g-1.day-1]=f{'age'} [day] x,y pairs :{0 0.09 2 0.04 6
1713 0.04 1000
1714 0.04} (Fan et al., 2003)
1715 2.1.11.17 'root class id' = 101 [-]
1716 2.1.11.18 'root hair density' [#.cm-2]=f{'age'} [day] x,y pairs :{0 2000 1 2000 2 2000 10
1717 2000 30 0
1718 2000 0} (Zhu et al. 2005; Mackay and S. Barber 1985)
1719 2.1.11.19 'root hair diameter' = 0.0005 [cm]
1720 2.1.11.20 'root hair length' [cm]=f{'age'} [day] x,y pairs :{0 0 1 0 2 0.028 2000 0.028}
1721 2.4.15.20
1722 (Zhu et al. 2005; Mackay and S. Barber 1985)
1723 2.1.11.21 'soil impedance.v2' [cm]=f{'uniform distribution'} minimum=-0.02
1724 maximum=0.02
1725 2.1.12 'primary root'
1726 2.1.12.1 'aerenchyma formation' [100%]=f{'age'} [day] x,y pairs :{0 0 3 0 5 0.1 10 0.25
1727 20 0.393 1000 0.393} (Fan et al. 2003)
1728 'cell size' [um^2]={101 150 200 250 300 350 400 450 500 533} (Chimungu et al., 2014a)
1729 'cell file number' [#]={8 9 10 11 12 13 14 15 16 17}
1730 2.1.12.2 'branch list'
1731 2.1.12.2.1 'lateral'
1732 2.1.12.2.1.1 'allow branches to form above ground' = 0 [-]
1733 2.1.12.2.1.2 'branching frequency' [cm]=f{'uniform distribution'} minimum=0.25
1734 maximum=0.45
1735 2.1.12.2.1.3 'length root tip' = 10.93 [cm]
1736 2.1.12.2.2 'seminal'
1737 2.1.12.2.2.1 'allow branches to form above ground' = 0 [-]
1738 2.1.12.2.2.2 'branching frequency' = 1 [cm]
1739 2.1.12.2.2.3 'branching time offset' = 1 [day]
1740 2.1.12.2.2.4 'max number of branches' = 5 [#]
1741 2.1.12.2.2.5 'number of branches/whorl' = 5 [#]
1742 2.1.12.3 'branching angle' = 0 [degrees]
1743 2.1.12.4 'density' = 0.094 [g.cm-3]
1744 2.1.12.5 'diameter' = 0.065 [cm]
1745 2.1.12.6 'gravitropism' = 0.01 [-]

1746 2.1.12.7 'gravitropism.v2' [cm]=f{'uniform distribution'} minimum=-0.015 maximum=-
1747 0.005
1748 2.1.12.8 'growth rate' [cm.day-1]=f{'age'} [day] x,y pairs : {0 4.5 28 4.5 38 0 1000 0}
1749 2.1.12.9 'length root tip without xylem vessels' = 2 [cm]
1750 2.1.12.10 'nitrate'
1751 2.1.12.10.1 ' Cmin' = 0.001 [umol.ml-1]
1752 2.1.12.10.2 ' Imax' [umol.cm-2.day-1]=f{'age'} [day] x,y pairs : {0 2.3 2 1.92 40 1.92}
1753 2.1.12.10.3 ' Km' [umol.ml-1]=f{'age'} [day] x,y pairs : {0 0.0105 2 0.0161 40 0.0161}
1754 2.1.12.10.4 'minimal nutrient concentration' = 600 [umol.g-1]
1755 2.1.12.10.5 'optimal nutrient concentration' = 1200 [umol.g-1]
1756 2.1.12.11 'number of xylem poles' = 8 [-]
1757 2.1.12.12 'phosphorus' (Barber 1995)
1758 2.1.12.12.1 ' Cmin' = 0.0002 [umol.ml-1]
1759 2.1.12.12.2 ' Imax' = 0.0555 [umol.cm-2.day-1]
1760 2.1.12.12.3 ' Km' = 0.00545 [umol.ml-1]
1761 2.1.12.12.4 'minimal nutrient concentration' = 30 [umol.g-1]
1762 2.1.12.12.5 'optimal nutrient concentration' = 60 [umol.g-1]
1763 2.1.12.13 'potassium' (Barber 1995)
1764 2.1.12.13.1 ' Cmin' = 0.002 [umol.ml-1]
1765 2.1.12.13.2 ' Imax' = 0.467 [umol.cm-2.day-1]
1766 2.1.12.13.3 ' Km' = 0.014 [umol.ml-1]
1767 2.1.12.13.4 'minimal nutrient concentration' = 117 [umol.g-1]
1768 2.1.12.13.5 'optimal nutrient concentration' = 234 [umol.g-1] (Silk et al. 1986)
1769 2.1.12.14 'radial hydraulic conductivity' [cm.day-1.hPa-1]=f{'time since planting'} [day]
1770 x,y pairs : {0 0 1 0.000216 10 0.000216 20 0.000216 30 0.000116 40 5e-05 60 0}
1771 2.1.12.15 'reduction in respiration due to aerenchyma' [100%]=f{'aerenchymaFormation'}
1772 [100%] x,y pairs : {0 0 0.3 0.7 0.6 1} (Fan et al. 2003)
1773 'reduction in respiration due to cell size' [100%]=f{'cellSize'} [100%] x,y pairs : {101 0
1774 150 0.07 200 0.14 250 0.17 300 0.25 350 0.32 400 0.37 450 0.43 500 0.51 533 0.57}
1775 'reduction in respiration due to file number' [100%]=f{'filenumber'} [100%] x,y pairs : {17
1776 0 16 0.05 15 0.13 14 0.17 13 0.23 12 0.3 11 0.35 10 0.42 9 0.49 8 0.52}
1777 2.1.12.16 'relative carbon cost of exudation' [g.cm-1.day-1]=f{'age'} [day] x,y pairs : {0
1778 5e-06 100
1779 5e-06} (Groleau-Renaud et al. 1998)
1780 2.1.12.17 'relative respiration' [g.g-1.day-1]=f{'age'} [day] x,y pairs : {0 0.09 2 0.04 6
1781 0.04 1000
1782 0.04} (Fan et al., 2003)
1783 2.1.12.18 'root class id' = 100 [-]
1784 2.1.12.19 'root hair density' [#.cm-2]=f{'age'} [day] x,y pairs : {0 2000 1 2000 2 2000 10
1785 2000 30 0
1786 2000 0} (Zhu et al. 2005; Mackay and S. Barber 1985)
1787 2.1.12.20 'root hair diameter' = 0.0005 [cm]
1788 2.1.12.21 'root hair length' [cm]=f{'age'} [day] x,y pairs : {0 0 1 0 2 0.028 2000 0.028}
1789 2.4.15.20
1790 (Zhu et al. 2005; Mackay and S. Barber 1985)
1791 2.1.12.22 'soil impedance' = 0.05 [-]
1792 2.1.12.23 'soil impedance.v2' [cm]=f{'uniform distribution'} minimum=-0.05
1793 maximum=0.05
1794 2.1.13 'resources'

1795 2.1.13.1 'carbon to dry weight ratio' = 0.45 [100%]
1796 2.1.13.2 'carbon allocation2 leafs factor' [100%]=f{'time'} [day] x,y pairs :{0 1 10 0.7 20
1797 0.45 33
1798 0.42 40 0.4 60 0.4}
1799 2.1.13.3 'carbon allocation2 roots factor' [100%]=f{'time'} [day] x,y pairs :{0 1 1 1 6 0.4
1800 20 0.2
1801 40 0.17 80 0.17}
1802 2.1.13.4 'carbon cost of nitrate uptake' = 1.392e-05 [g.umol-1]
1803 2.1.13.5 'max carbon allocation2 shoot' = 0.82 [100%]
1804 2.1.13.6 'nitrate'
1805 2.1.13.6.1 'initial nutrient uptake' = 285 [umol]
1806 2.1.13.7 'phosphorus' (Barber 1995)
1807 2.1.13.7.1 'initial nutrient uptake' = 20 [umol]
1808 2.1.13.8 'potassium' (Barber 1995)
1809 2.1.13.8.1 'initial nutrient uptake' = 27 [umol]
1810 2.1.13.9 'reserve allocation rate' [%day-1]=f{'time'} [day] x,y pairs :{0 0.01 1 0.02 2
1811 0.04 3 0.04
1812 10 0.2 11 0.2 1000 0.2}
1813 2.1.13.10 'seed size' = 0.15 [g]
1814 2.1.14 'seminal'
1815 2.1.14.1 'aerenchyma formation' [100%]=f{'age'} [day] x,y pairs :{0 0 3 0 5 0.1 10 0.25
1816 20 0.393 1000 0.393} (Fan et al. 2003)
1817 'cell size' [um²]={101 150 200 250 300 350 400 450 500 533} (Chimungu et al., 2014a)
1818 'cell file number' [#]={8 9 10 11 12 13 14 15 16 17}
1819 2.1.14.2 'branch list'
1820 2.1.14.2.1 'lateral'
1821 2.1.14.2.1.1 'allow branches to form above ground' = 0 [-]
1822 2.1.14.2.1.2 'branching frequency' [cm]=f{'uniform distribution'} minimum=0.05
1823 maximum=0.25
1824 2.1.14.2.1.3 'length root tip' = 10.93 [cm]
1825 2.1.14.3 'branching angle' = 90 [degrees]
1826 2.1.14.4 'density' = 0.094 [g.cm-3]
1827 2.1.14.5 'diameter' = 0.085 [cm]
1828 2.1.14.6 'gravitropism' = 0.004 [-]
1829 2.1.14.7 'gravitropism.v2' [cm]=f{'uniform distribution'} minimum=-0.035 maximum=-
1830 0.025
1831 2.1.14.8 'growth rate' [cm.day-1]=f{'age'} [day] x,y pairs :{0 0.01 1 0.5 2 4.5 28 4.5 38 0
1832 100 0}
1833 2.1.14.9 'length root tip without xylem vessels' = 2 [cm]
1834 2.1.14.10 'longitudinal growth rate multiplier' [cm]= f{'normal distribution'}
1835 minimum=0.6
1836 maximum=1.2 mean=1 stdev=0.1
1837 2.1.14.11 'nitrate'
1838 2.1.14.11.1 'Cmin' = 0.001 [umol.ml-1]
1839 2.1.14.11.2 'Imax' [umol.cm-2.day-1]=f{'age'} [day] x,y pairs :{0 2.3 2 1.92 40 1.92}
1840 2.1.14.11.3 'Km' [umol.ml-1]=f{'age'} [day] x,y pairs :{0 0.0105 2 0.0161 40 0.0161}
1841 2.1.14.11.4 'minimal nutrient concentration' = 600 [umol.g-1]
1842 2.1.14.11.5 'optimal nutrient concentration' = 1200 [umol.g-1]
1843 2.1.14.12 'number of xylem poles' = 6 [-]

1844 2.1.14.13 'phosphorus' (Barber 1995)
1845 2.1.14.13.1 'Cmin' = 0.0002 [umol.ml-1]
1846 2.1.14.13.2 'Imax' = 0.0555 [umol.cm-2.day-1]
1847 2.1.14.13.3 'Km' = 0.00545 [umol.ml-1]
1848 2.1.14.13.4 'minimal nutrient concentration' = 30 [umol.g-1]
1849 2.1.14.13.5 'optimal nutrient concentration' = 60 [umol.g-1] 2.1.14.14 'potassium' (Barber
1850 1995)
1851 2.1.14.14.1 'Cmin' = 0.002 [umol.ml-1]
1852 2.1.14.14.2 'Imax' = 0.467 [umol.cm-2.day-1]
1853 2.1.14.14.3 'Km' = 0.014 [umol.ml-1]
1854 2.1.14.14.4 'minimal nutrient concentration' = 117 [umol.g-1]
1855 2.1.14.14.5 'optimal nutrient concentration' = 234 [umol.g-1] (Silk et al. 1986)
1856 2.1.14.15 'reduction in respiration due to aerenchyma' [100%]=f{'aerenchymaFormation'}
1857 [100%] x,y pairs :{0 0 0.3 0.7 0.6 1} (Fan et al. 2003)
1858 'reduction in respiration due to cell size' [100%]=f{'cellSize'} [100%] x,y pairs :{101 0
1859 150 0.07 200 0.14 250 0.17 300 0.25 350 0.32 400 0.37 450 0.43 500 0.51 533 0.57}
1860 'reduction in respiration due to file number' [100%]=f{'filenumber'} [100%] x,y pairs :{17
1861 0 16 0.05 15 0.13 14 0.17 13 0.23 12 0.3 11 0.35 10 0.42 9 0.49 8 0.52}
1862 2.1.14.16 'regular topology' = 1 [-]
1863 2.1.14.17 'relative carbon cost of exudation' [g.cm-1.day-1]=f{'age'} [day] x,y pairs :{0
1864 5e-06 100
1865 5e-06} (Groleau-Renaud et al. 1998)
1866 2.1.14.18 'relative respiration' [g.g-1.day-1]=f{'age'} [day] x,y pairs :{0 0.09 2 0.04 6
1867 0.04 1000
1868 0.04} (Fan et al., 2003)
1869 2.1.14.19 'root class id' = 99 [-]
1870 2.1.14.20 'root hair density' [#.cm-2]=f{'age'} [day] x,y pairs :{0 2000 1 2000 2 2000 10
1871 2000 30 0
1872 2000 0} (Zhu et al. 2005; Mackay and S. Barber 1985)
1873 2.1.14.21 'root hair diameter' = 0.0005 [cm]
1874 2.1.14.22 'root hair length' [cm]=f{'age'} [day] x,y pairs :{0 0 1 0 2 0.028 2000 0.028}
1875 2.4.15.20
1876 (Zhu et al. 2005; Mackay and S. Barber 1985)
1877 2.1.14.23 'soil impedance' = 0.02 [-]
1878 2.1.14.24 'soil impedance.v2' [cm]=f{'uniform distribution'} minimum=-0.04
1879 maximum=0.04
1880 2.1.15 'shoot'
1881 2.1.15.1 'area per plant' = 1600 [cm2]
1882 2.1.15.2 'extinction coefficient' = 0.85 [-]
1883 2.1.15.3 'leaf area expansion rate' [cm2.day-1]=f{'time'} [day] x,y pairs :{0 0 2 0 2.38
1884 2.32 2.77 3.24 3.15 3.93 3.54 4.41 3.92 4.72 4.3 4.87 4.69 4.89 5.07 4.81 5.45 4.64 5.84 4.41 6.22
1885 4.14 6.61 3.84 6.99 3.55 7.37 3.27 7.76 3.02 8.14 2.83 8.53 2.71 8.91 2.66 9.29 2.71 9.68 2.88
1886 10.06 3.16 10.44 3.58 10.83 4.15 11.21 4.87 11.6 5.76 11.98 6.82 12.36 8.07 12.75 9.5 13.13
1887 11.13 13.52 12.96 13.9 14.99 14.28 17.23 14.67 19.68 15.05 22.35 15.43 25.22 15.82 28.32 16.2
1888 31.62 16.59 35.14 16.97 38.87 17.35 42.81 17.74 46.95 18.12 51.29 18.51 55.83 18.89 60.55
1889 19.27 65.45 19.66 70.53 20.04 75.76 20.42 81.16 20.81 86.69 21.19 92.36 21.58 98.15 21.96
1890 104.05 22.34 110.04 22.73 116.11 23.11 122.24 23.49 128.42 23.88 134.63 24.26 140.86 24.65
1891 147.08 25.03 153.28 25.41 159.42 25.8 165.51 26.18 171.5 26.57 177.39 26.95 183.14 27.33
1892 188.73 27.72 194.13 28.1 199.33 28.48 204.29 28.87 208.98 29.25 213.38 29.64 217.45 30.02

1893 221.18 30.4 224.52 30.79 227.44 31.17 229.92 31.56 231.91 31.94 233.39 33.09 234.36 50
1894 234.36 80 0} (Zhang and Postma, University Park, unpublished)
1895 2.1.15.4 'light use efficiency' = 3.8×10^{-7} [g.umol⁻¹] (Stirling et al. 1994; Postma,
1896 University Park,unpublished)
1897 2.1.15.5 'nitrate'
1898 2.1.15.5.1 'leaf minimal nutrient concentration' [umol.g⁻¹]=f{'time'} [day] x,y pairs :{0
1899 1200 80
1900 800}
1901 2.1.15.5.2 'leaf optimal nutrient concentration' [umol.g⁻¹]=f{'time'} [day] x,y pairs :{0
1902 2500 80
1903 1500} (Zhang and Postma, University Park, Unpublished; Alexandrova and Donovan 2003;
1904 Chevalier and Schrader 1977)
1905 2.1.15.5.3 'stem minimal nutrient concentration' = 400 [umol.g⁻¹]
1906 2.1.15.5.4 'stem optimal nutrient concentration' = 800 [umol.g⁻¹]
1907 2.1.15.6 'phosphorus'
1908 2.1.15.6.1 'leaf minimal nutrient concentration' = 35 [umol.g⁻¹]
1909 2.1.15.6.2 'leaf optimal nutrient concentration' = 70 [umol.g⁻¹] (Zhang and Postma,
1910 University
1911 Park, unpublished)
1912 2.1.15.6.3 'stem minimal nutrient concentration' = 15 [umol.g⁻¹]
1913 2.1.15.6.4 'stem optimal nutrient concentration' = 30 [umol.g⁻¹]
1914 2.1.15.7 'potassium'
1915 2.1.15.7.1 'leaf minimal nutrient concentration' = 273 [umol.g⁻¹]
1916 2.1.15.7.2 'leaf optimal nutrient concentration' = 508 [umol.g⁻¹] (Leigh and Jones 1984)
1917 2.1.15.7.3 'stem minimal nutrient concentration' = 117 [umol.g⁻¹]
1918 2.1.15.7.4 'stem optimal nutrient concentration' = 250 [umol.g⁻¹]
1919 2.1.15.8 'relative potential transpiration' = 100 [cm³.g⁻¹] (Baldocchi 1994)
1920 2.1.15.9 'relative respiration rate leafs' = 0.04 [g.g⁻¹.day⁻¹] (Postma, University Park,
1921 Unpublished)
1922 2.1.15.10 'relative respiration rate stems' = 0.02 [g.g⁻¹.day⁻¹]
1923 2.1.15.11 'specific leaf area' [g.cm⁻²]=f{'time'} [day] x,y pairs :{0 0.0015 24 0.0026 50
1924 0.0032
1925 100 0.0032} (van Heemst 1988; Jacob and Lawlor 1991; Jaramillo, University Park,
1926 unpublished)
1927 2.1.16 'stress impact factors'
1928 2.1.16.1 'impact on:leaf area expansion rate'
1929 2.1.16.1.1 'impact by:nitrate' [-]=f{'nitrate stress factor'} [-] x,y pairs :{0 0 0.3 0.1 1 1}
1930 (Sinclair and Horie 1989)
1931 2.1.16.1.2 'impact by:phosphorus' [-]=f{'phosphorus stress factor'} [-] x,y pairs :{0 0 1 1}
1932 (Lynch et al. 1991; Usuda and Shimogawara 1991) (Lynch et al. 1991; Usuda and Shimogawara
1933 1991)
1934 2.1.16.1.3 'impact by:potassium' [-]=f{'potassium stress factor'} [-] x,y pairs :{0 0 0.2 0.5
1935 1 1}
1936 2.1.16.2 'impact on:photosynthesis'
1937 2.1.16.2.1 'impact by:nitrate' [-]=f{'nitrate stress factor'} [-] x,y pairs :{0 0 0.4 0.5 1 1}
1938 (Sinclair
1939 and Horie 1989)
1940 2.1.16.2.2 'impact by:phosphorus' [-]=f{'phosphorus stress factor'} [-] x,y pairs :{0 0.5
1941 0.5 1 1

- 1942 1} (Lynch et al. 1991; Usuda and Shimogawara 1991)
- 1943 2.1.16.2.3 'impact by:potassium' [-]=f{'potassium stress factor'} [-] x,y pairs :{0 0 1 1}
- 1944
- 1945 Cited literature in appendix
- 1946 Alexandrova, P. and Donovan, D., 2003. Nitrogen content and its uptake by maize as influenced by
- 1947 some meteorological elements and fertilization. *International Agrophysics* , 17(2), p.41–46.
- 1948 Baldocchi, D., 1994. A comparative study of mass and energy exchange rates over a closed C3
- 1949 (wheat) and an open C4 (corn) crop: II. CO2 exchange and water use efficiency. *Agricultural and*
- 1950 *Forest Meteorology*, 67(3-4), pp.291-321.
- 1951 Barber, S.A., 1995. *Soil nutrient bioavailability: a mechanistic approach* , John Wiley and Sons,
- 1952 New York, USA.
- 1953 Bhadoria, P.B.S. et al., 1991. Soil Phosphate Diffusion Coefficients: Their Dependence on
- 1954 Phosphorus Concentration and Buffer Power. *Soil Sci Soc Am J* , 55(1), pp.56-60.
- 1955 Burton, A.L., 2010. Phenotypic Evaluation and Genetic Basis of Anatomical and Architectural
- 1956 Root Traits in the Genus Zea. PhD dissertation. University Park: Penn State. Available at:
- 1957 <http://etda.libraries.psu.edu/theses/approved/PSUonlyIndex/ETD-5741/index.html> [Accessed
- 1958 February 9, 2011].
- 1959 Chevalier, P. and Schrader, L.E., 1977. Genotypic Differences in Nitrate Absorption and
- 1960 Partitioning of N Among Plant Parts in Maize. *Crop Sci* , 17(6), pp.897-901.
- 1961 Claassen, N., Syring, K. and Jungk, A., 1986. Verification of a mathematical model by simulating
- 1962 potassium uptake from soil. *Plant and Soil*, 95(2), pp.209-220.
- 1963 Dunham, R.J. and Nye, P.H., 1976. The Influence of Soil Water Content on the Uptake of Ions by
- 1964 Roots. III. Phosphate, Potassium, Calcium and Magnesium Uptake and Concentration Gradients
- 1965 in Soil. *Journal of Applied Ecology*, 13(3), pp.967-984.
- 1966 Fan, M., Zhu, J., et al., 2003. Physiological roles for aerenchyma in phosphorus-stressed roots.
- 1967 *Functional Plant Biology*, 30(5), pp.493-506.
- 1968 Groleau-Renaud, V., Plantureux, S. and Guckert, A., 1998. Influence of plant morphology on root
- 1969 exudation of maize subjected to mechanical impedance in hydroponic conditions. *Plant and Soil*,
- 1970 201(2), pp.231-239.
- 1971 van Heemst, H.D.J., 1988. Plant data values required for simple crop growth simulation models:
- 1972 review and bibliography. *Simulation Report CABO-TT*, Wageningen , 17, pp.0-100.
- 1973 Hodnett, M.G. and Tomasella, J., 2002. Marked differences between van Genuchten soil water-
- 1974 retention parameters for temperate and tropical soils: a new water-retention pedo-transfer
- 1975 functions developed for tropical soils. *Geoderma* , 108(3-4), pp.155-180.
- 1976 Jacob, J. and Lawlor, D.W., 1991. Stomatal and mesophyll limitations of photosynthesis in
- 1977 phosphate deficient sunflower, maize and wheat plants. *Journal of Experimental Botany*, 42(8),
- 1978 pp.1003-1011.
- 1979 Leigh, R.A. and Jones, R.G.W., 1984. A Hypothesis Relating Critical Potassium Concentrations
- 1980 for
- 1981 Growth to the Distribution and Functions of this Ion in the Plant Cell. *New Phytologist* , 97(1),
- 1982 pp.1-13.
- 1983 Luo, L., Lin, H. and Halleck, P., 2008. Quantifying Soil Structure and Preferential Flow in Intact
- 1984 Soil Using X-ray Computed Tomography. *Soil Sci Soc Am J* , 72(4), pp.1058-1069.
- 1985 Lynch, J.P., Lauchli, A. and Epstein, E., 1991. Vegetative growth of the common bean in
- 1986 response to phosphorus nutrition. *Crop Science*, 31(2), pp.380-387.
- 1987 Mackay, A. and Barber, S., 1985. Effect of soil moisture and phosphate level on root hair growth
- 1988 of corn roots. *Plant and Soil*, 86(3), pp.321-331.
- 1989 Pahlavanian, A.M. and Silk, W.K., 1988. Effect of Temperature on Spatial and Temporal Aspects
- 1990 of Growth in the Primary Maize Root. *Plant Physiology*, 87(2), pp.529-532.

- 1991 Silk, W.K. et al., 1986. Spatial Distributions of Potassium, Solutes, and Their Deposition Rates in
- 1992 the Growth Zone of the Primary Corn Root. *Plant Physiol.*, 82(3), pp.853-858.
- 1993 Sinclair, T.R. and Horie, T., 1989. Leaf Nitrogen, Photosynthesis, and Crop Radiation Use
- 1994 Efficiency: A Review. *Crop Science*, 29(1), pp.90-98.
- 1995 Stirling, C.M. et al., 1994. Changes in the photosynthetic light response curve during leaf
- 1996 development of field grown maize with implications for modelling canopy photosynthesis.
- 1997 *Photosynthesis Research*, 42(3), pp.217-225.
- 1998 Usuda, H. and Shimogawara, K., 1991. Phosphate Deficiency in Maize. I. Leaf Phosphate Status,
- 1999 Growth, Photosynthesis and Carbon Partitioning. *Plant and Cell Physiology*, 32(4), pp.497-504.
- 2000 Yang, H.S. and Janssen, B.H., 2000. A mono-component model of carbon mineralization with a
- 2001 dynamic rate constant. *European Journal of Soil Science*, 51(3), pp.517-529.
- 2002 Zhu, J., Kaeppler, S.M. and Lynch, J.P., 2005. Mapping of QTL controlling root hair length in
- 2003 maize (*Zea mays* L.) under phosphorus deficiency. *Plant and Soil* , 270(1), pp.299-310.