Title

Supramolecular assembly of the *E. coli* LdcI upon acid stress

Authors

Matthew Jessop*, Clarissa Liesche*, Jan Felix*, Ambroise Desfosses, Megghane Baulard, Virgile Adam, Angélique Fraudeau, Karine Huard, Grégory Effantin, Jean-Philippe Kleman, Maria Bacia-Verloop, Dominique Bourgeois, Irina Gutsche[#]

¹Institut de Biologie Structurale, Univ Grenoble Alpes, CEA, CNRS, IBS, 71 Avenue des martyrs, F-38044 Grenoble, France

*: equal contribution, #: corresponding author.

Significance statement

Bacteria possess a sophisticated arsenal of defence mechanisms that allow them to survive in adverse conditions. Adaptation to acid stress and hypoxia is crucial for the enterobacterial transmission in the gastrointestinal tract of their human host. When subjected to low pH, *E. coli* and many other enterobacteria activate a proton-consuming resistance system based on the acid-stress inducible lysine decarboxylase LdcI. Here we develop generally-applicable tools to uncover the spatial localisation of LdcI inside the cell by super-resolution fluorescence microscopy, and investigate the *in vitro* supramolecular organisation of this enzyme by cryo-EM. We build on these results to propose a mechanistic model for LdcI function and offer tools for further *in vivo* investigations.

Abstract

Pathogenic and commensal bacteria often have to resist the harsh acidity of the host stomach. The inducible lysine decarboxylase LdcI buffers the cytosol and the local extracellular environment to ensure enterobacterial survival at low pH. Here, we investigate the acid-stress response regulation of *E. coli* LdcI by combining biochemical and biophysical characterisation with negative stain and cryo-electron microscopy, and wide-field and super-resolution fluorescence imaging. Due to deleterious effects of fluorescent protein fusions on native LdcI decamers, we opt for three-dimensional localisation of nanobody-labelled endogenous wild-type LdcI in acid-stressed *E. coli* cells, and show that it organises into distinct patches at the cell periphery. Consistent with recent hypotheses that *in vivo* clustering of metabolic enzymes often reflects their polymerisation as a means of stimulus-induced regulation, we show that LdcI assembles into filaments *in vitro* at physiologically relevant low pH. We solve the structures of these filaments and of the LdcI decamer formed at neutral pH by cryo-electron microscopy, and reveal the molecular determinants of LdcI polymerisation, confirmed by mutational analysis. Finally, we propose a model for LdcI function inside the enterobacterial cell, providing a structural and mechanistic basis for further investigation of the role of its supramolecular organisation in the acid stress response.

Introduction

Cell survival requires the adaptation of metabolism to changing environmental demands. Biochemical regulation of metabolic enzymes by cellular metabolites has been intensely studied for many decades. In addition, a growing body of recent light microscopy observations highlights the spatial regulation of enzymes by stimulus-induced phase separation into distinct loci – liquid droplets, amyloid-like aggregates or ordered polymers (1). In eukaryotes, the observed condensation of fluorescently-labelled metabolic enzymes is triggered by stress, including medium acidification, hypoxia and nutrient limitation. Enterobacteria such as *Escherichia coli*, *Salmonella* and *Vibrio* encounter these types of conditions in the host gastrointestinal tract (2). One of the key enterobacterial proteins expressed during the acid stress response, upon oxygen limitation and regulated by the nutrient stress alarmone guanosine tetraphosphate (ppGpp), is the acid stress-inducible lysine decarboxylase LdcI (3–6). LdcI has been scrutinised since the early 1940s because of direct links between the efficiency of the acid stress response and pathogenicity (7–9). This enzyme transforms lysine into cadaverine while consuming protons and producing CO₂, thereby contributing to buffering of the bacterial cytosol and the extracellular medium under acid stress conditions to promote bacterial survival. While both the structure and the function of LdcI have been thoroughly studied (6, 10), nothing is known about its localisation inside the bacterial cell, because its fluorescent protein fusions were previously described to form inclusion bodies (11).

Whereas the overwhelming majority of super-resolution fluorescence imaging is focused on eukaryotes, optical studies of bacterial systems are nearly exclusively centred on large macromolecular complexes with obvious superstructure such as cytoskeletal, cell division, chromosome partitioning, RNA degradation and secretion machineries (12–14). However, the relevance of the documented patchy or long-range helical localisation of some of these assemblies is now questioned and requires re-evaluation. Indeed, the vast majority of these studies were based on labelling with either fluorescent proteins or epitope tags shown to be able to induce artefactual associations and localisations (13–16).

A handful of examples of regulation of bacterial metabolic enzymes by phase separation through stimulus-triggered polymerisation concern well-conserved oligomeric proteins involved in nucleotide and amino acid metabolism, such as CTP synthase (17, 18) and glutamine synthetase (19). Interestingly, these enzymes are also able to assemble into filaments *in vitro*, and their *in vivo* condensates, detected both in bacteria and in eukaryotes, have been suggested to correspond to the polymerised state of the enzymes. Other examples of bacterial metabolic enzymes purified as polymers from bacterial extracts or forming polymers *in vitro* are the aldehyde-alcohol dehydrogenase AdhE (20) and the hydrogen-dependent CO₂ reductase HDCR (21).

Specific to bacteria, LdcI is a decamer composed of five dimers tightly arranged into pentameric double-rings (PDB ID: 3N75) (6). Interestingly, *in vitro* these double-rings were observed by negative stain electron microscopy (ns-EM) to stack on top of one another into filament-like structures (6, 8) but only at non-physiological pH below 5, under which LdcI should not be significantly expressed. Here, to address the spatial localisation of the *E. coli* LdcI, we start by critically evaluating labelling artefacts with an ambition to define the optimal constructs for subsequent chromosomal manipulations. To this end, we overexpress and purify different

fluorescent protein (FP) fusions of Ldcl, perform their structural and biochemical characterisation *in vitro*, and observe the distribution of the overexpressed constructs *in vivo*. This methodological section enables us to propose a workflow that brings together examinations of *in vivo* protein localisations with *in vitro* biochemical and ns-EM characterisations of the purified FP fusions in order to ensure artefact-free optical imaging investigations. This analysis is followed by unveiling a patchy distribution of endogenous Ldcl inside the *E. coli* cell upon acid stress. Because this condensation of Ldcl into patches is reminiscent of stress-induced polymerisation, in particular in yeast, we investigate the *in vitro* architecture of Ldcl at pH 5.7. We show that at this physiological pH, corresponding to the optimum of the Ldcl enzymatic activity, Ldcl is a polymer assembled of stacked decamers, and solve the polymer structure by cryo-electron microscopy (cryo-EM). For comparison, we also determined the cryo-EM structure of the Ldcl decamer at neutral pH. In addition, mutational analysis of the Ldcl stack-forming interfaces allowed identification of critical residues involved in stack formation. Finally, we discuss the observed Ldcl localisation pattern in the light of the wealth of available functional and imaging data, and offer a structural and mechanistic basis of supramolecular Ldcl assembly, which will aid in the design of future experiments linking Ldcl stack formation to *E. coli* acid stress fitness.

Results

Fluorescent protein fusions affect Ldcl structure without modifying localisation of the overexpressed fusion constructs

Because of the small size of the *E. coli* cell, we opted for super-resolution microscopy imaging (22) and set out to localise Ldcl inside the cell upon acid stress by either Photoactivation Localisation Microscopy (PALM) or Stochastic Optical Reconstruction Microscopy (STORM) (23–25). *A priori*, PALM seemed more relevant because this technique relies on genetically-encoded FPs fused to the protein of interest, and is therefore unmatched in terms of labelling specificity and efficiency. In addition, PALM does not require delivery of fluorescent molecules across the cell wall, and enables live cell imaging and single molecule tracking.

Considering that Ldcl is an acid stress response enzyme, we opted for either mGeosM (26) or Dendra2_{T69A} (27) as FP markers because of their relatively low pKa values, high monomericity and high fluorescent quantum yields. In the Ldcl decamer structure, the N-termini are oriented inwards and towards the central pore of the double ring, while the C-termini point to the ring periphery and are readily accessible from the outside. Therefore, on the one side, one could assume that an N-terminal labelling with an FP would be likely to interrupt the Ldcl tertiary structure. On the other side, the only well characterised binding partner of Ldcl, the AAA+ ATPase RavA, is known to interact precisely with the C-terminal β -sheet of Ldcl (28). Assembly of two copies of double-pentameric rings of Ldcl and five copies of hexameric RavA spirals results in a huge 3.3 MDa macromolecular cage of intricate architecture and largely unknown function (28–30). The Ldcl-RavA cage is proposed to assist assembly of specific respiratory complexes in *E. coli* and to counteract acid stress under starvation conditions by alleviating ppGpp-dependent inhibition of Ldcl (31–33). Thus, although these functions still require further investigation, preservation of the RavA-binding propensity should be one of the criteria for assessing the suitability of an Ldcl-FP fusion. Therefore, both N- and C-terminal fusion constructs with either mGeosM or Dendra2_{T69A}

attached to Ldcl via an appropriate linker were cloned into dedicated plasmids and overexpressed at conditions optimised for Ldcl overproduction (see Methods for details). Expression of fusion proteins was immediately detected by wide-field fluorescence imaging that showed a similar distribution for the four fusions (Figure 1A). Each construct was then purified in order to assess its structural integrity and RavA binding capacity *in vitro*, with a goal of defining the most suitable construct for the subsequent creation of a corresponding chromosomal fusion.

The N-terminal Dendra2_{T69A}-Ldcl fusion formed exclusively dimers, confirming the structure-based prediction that the fluorescent tag at this position would perturb the dimer-dimer interaction (Figure 1B, Supplementary Figure 1). Admittedly, native Ldcl was shown to dissociate into dimers *in vitro* at pH above approximately 7.5, but in this pH range Ldcl is not supposed to be expressed in the cell and therefore this dissociation may be irrelevant. Surprisingly, in contrast to Dendra2_{T69A}-Ldcl, mGeosM-Ldcl assembled into regular symmetric non-native higher-order oligomers, with a dramatically altered quaternary structure (Figure 1B, Supplementary Figure 1). These oligomers were built of three Ldcl tetramers, bridged together by additional densities. Noteworthy, mEos2, the fluorescent protein from which mGeosM was derived, crystallises as a tetramer (PDB ID: 3S05) that can be straightforwardly fitted into the Ldcl-bridging densities (Figure 1B). This illustrates that despite the fact that mGeosM was designed as a monomeric FP (with the first “m” explicitly standing for monomeric), some residual oligomeric tendency is still maintained. This propensity of mGeosM to oligomerise when bound to Ldcl may be driven by avidity effects – as Ldcl dimers begin to assemble into a decamer, the local concentration of mGeosM increases to such a point that oligomerisation becomes energetically favourable, despite the apparent monomer behaviour of mGeosM in gel filtration (26). This is also in line with the known propensity of mEos2, from which mGeosM is derived, to form tetramers at high concentration (34). To conclude, both N-terminal fusions induced non-native assembly of Ldcl, and therefore neither were appropriate for determining the native localisation of this enzyme in the cell.

As expected, C-terminal Ldcl fluorescent fusions formed decamers with protruding densities that can be attributed to flexibly attached FPs (Figure 1B, Supplementary Figure 1). Nevertheless, despite the long flexible linker between Ldcl and the FP, these constructs were unable to interact with RavA as shown by Bio-Layer Interferometry (BLI) binding studies (see Methods and Supplementary Figure 1 for details). This means that the functionality of these fusions cannot be considered as entirely retained. Thus, none of the four fusions were suitable for an in-depth imaging analysis under conditions of native Ldcl expression upon acid stress that we planned to undertake. Altogether, we demonstrated that structural integrity and unaltered interaction with known partners are useful read-outs for functional preservation. Based on this result we propose that, when feasible, purification and structure-function analysis of FP fusions should be performed prior to interpretation of the protein localisation inferred from observation of FP fusions by optical methods. This workflow can be used for example in cases where chromosomal manipulation for assessment of intact function is difficult and/or the phenotype is condition-dependent or unclear, whereas basic *in vitro* biochemical and ns-EM characterisation are efficiently set up.

Immunofluorescence of LdcI in *E. coli* under acid stress reveals its supramolecular organisation

Because none of the FP fusions possessed the properties of native LdcI, we decided to turn to STORM imaging of the endogenous enzyme. One of the drawbacks of STORM is the requirement for exogenous labelling and therefore the fixation and permeabilisation of cells. These procedures are notoriously known to potentially affect cell morphology, and therefore, when possible, the usage of live cell imaging with FP fusions (as for example in PALM), or observation of unfixed samples under cryogenic conditions would be ideal. Moreover, fixation, permeabilisation and specific exogenous labelling in bacteria present unique challenges because of the complex, often multi-layered cell wall (35, 36). A significant advantage of STORM however is the possibility of direct imaging of the wild-type (WT) protein, which should circumvent the dangers associated with FP fusions described above. The three prerequisites for imaging of WT systems by immuno-based labelling in general and of LdcI in particular are (i) availability of an antibody or nanobody (antibody fragment derived from heavy-chain-only camelid antibody) coupled to an organic fluorescent dye and directed towards the native protein, (ii) precise knowledge of endogenous expression conditions, (iii) validation of an efficient permeabilisation and immunolabelling technique that enables the antibody/nanobody to enter the cell and specifically target the protein of interest without altering the native organization of the cell ultrastructures.

As a first step towards STORM imaging of endogenous LdcI, we probed an anti-LdcI nanobody, hereafter called anti-LdcI-Nb. The complex between purified *E. coli* LdcI and the nanobody was purified by size exclusion chromatography (see Methods), and imaged by ns-EM. The resulting 3D map of the LdcI/anti-LdcI-Nb complex demonstrates binding of anti-LdcI-Nb to each LdcI monomer in the decamer and reveals the location of the interaction site which is clearly distinct from the RavA binding site (Figure 2A, Supplementary Figure 2). This nanobody was thus identified as a suitable labelling agent for LdcI imaging in *E. coli* cells, and labelled with the fluorescent dye AlexaFluor647 (AF647) or AlexaFluor488 (AF488) (see Methods).

Consistently with published data, LdcI expression could be induced by a pH shift experiment (i.e. transfer of bacteria from pH 7.0 into a pH 4.6 growth medium) in the presence of lysine under oxygen-limiting conditions (Supplementary Figure 3A). While the WT strain grew well under these conditions and efficiently buffered the extracellular medium up to pH 6.2 in approximately 1.5 to two hours (Supplementary Figure 3A) concomitantly with the increase in the level of LdcI expression (Supplementary Figure 3B, C), the growth and the acid stress response of the $\Delta ldcI$ mutant strain were severely impaired (Supplementary Figure 3A). Since the peak of LdcI expression by the wild type cells was achieved between one and two hours after exposure to acid stress (Supplementary Figure 3B, C), a time point of 90 minutes was chosen for the subsequent labelling and imaging experiments. The specificity and performance of anti-LdcI-Nb in immunofluorescence labelling of permeabilised *E. coli* cells were characterised by flow cytometry and wide-field fluorescence imaging (Supplementary Figure 3D, E). Both techniques demonstrated that in the absence of LdcI expression no specific fluorescence is seen, confirming the suitability of anti-LdcI-Nb for immunofluorescence studies. Thus, the above-mentioned prerequisites for immuno-based imaging of cellular LdcI have been fulfilled. Noteworthy, the LdcI expression profile

highlighted a considerable asset of the usage of STORM instead of PALM for visualisation of the endogenously expressed Ldcl: indeed, the transient nature of the expression and the necessity of work under oxygen-limiting conditions may have created difficulties due to the longer maturation time of the FPs under these conditions.

Initial characterisation of the cellular distribution of Ldcl 90 minutes after exposure of *E. coli* cells to acid stress was carried out by wide-field fluorescence imaging. Based on these images, it appeared that natively expressed Ldcl did not display a homogeneous cytoplasmic distribution but rather showed a patchy localisation pattern (Figure 2B). 3D STORM imaging subsequently provided a more detailed view of this patchy distribution. As shown in Figure 3, Supplementary Figure 4, and Supplementary Movies 1-5, the labelling density was lower in the centre of the bacterium. This indicates a propensity for Ldcl to cluster near the cell periphery, towards the inner membrane and the cell poles, rather than being distributed homogeneously through the volume. Recent investigations argue that, in most cases, *in vivo* clustering of metabolic enzymes corresponds to their polymerised states, and represents an efficient means of regulation of enzymatic activity and metabolic homeostasis in response to a stimulus (1). Thus, structure determination of these polymers is a crucial step towards the understanding of regulation mechanisms. We previously documented that *in vitro* at pH below 5 and high concentration, Ldcl decamers indeed tend to stack (6). However, at pH 5.7, optimal for Ldcl expression and enzymatic activity and likely corresponding to the internal pH upon acid stress, the oligomeric state of Ldcl has not yet been addressed by EM. Supposing that the high local concentration of Ldcl, clustered in patches in the *E. coli* cell under acid stress conditions, is likely to induce enzyme polymerisation via stack formation, we next examined the Ldcl assembly state at pH 5.7 by cryo-EM and discovered that it does indeed form straight polymers (Figure 4).

Structural determinants of Ldcl stack formation revealed by cryo-EM

In order to understand the molecular determinants of Ldcl polymerisation at low pH and to provide a framework for the analysis of Ldcl function under acid stress, we solved the 3D structures of the Ldcl decamer at pH 7.0 and of Ldcl stacks at pH 5.7 by cryo-EM (See Methods, Figure 4, Supplementary Figures 5-6, Supplementary Table 1). The 2.8 Å resolution structure of the Ldcl decamer at pH 7.0 (Figure 4A) is extremely similar (Supplementary Table 2 & 3) to the Ldcl crystal structure solved at pH 8.5 in an inhibited ppGpp-bound state (6). However, contrary to pH 7.0 and even pH 6.2 where Ldcl is still predominantly decameric (10), Ldcl forms straight rigid filaments on the cryo-EM grid at pH 5.7, which corresponds to the pH of maximum Ldcl enzymatic activity (Figure 4B). The structure of a three-decamer stack was solved to a resolution of 3.3 Å, revealing the structural details of acid stress-induced Ldcl polymerisation (Figure 4C). Ldcl decamers stack tightly on top of one other, with negligible rotation between decamers along the stack. Each dimer fits snugly in the inter-dimer groove of decamers above and below.

A comparison of Ldcl decamer structures taken from the Ldcl stack cryo-EM map (at pH 5.7), the Ldcl decamer cryo-EM map (at pH 7.0) and the crystal structure of decameric Ldcl crystallised with bound ppGpp at pH 8.5 (PDB ID: 3N75) reveals some remarkable differences between the stack structure and the two decamer structures (Figure 4D, E, Supplementary Table 2 & 3). While the three structures do not show any major differences at the monomer level, a

structural alignment of an Ldcl dimer extracted from the Ldcl decamer structures with a dimer extracted from the Ldcl stack structure uncovers a rigid body-like rotation between monomers around a hinge region located at the monomer-to-monomer interface (Supplementary Table 2 & 3, Figure 4DE). This rotation results in a 5° tilt when comparing the N-terminal wing-domains in Ldcl dimers, and an overall slightly decreased diameter of the central cavity inside the stacked Ldcl rings (Figure 4D, E), which may contribute to the tight packing of each dimer into the grooves of an opposing decamer in the stack.

A careful examination of the Ldcl stack structure shows that two major inter-decamer interfaces situated at a two-fold symmetry axis perpendicular to the stack direction contribute to stack formation (Figure 5A). In particular, the first interface (Figure 5B) is formed between residues K422, D460, R468, D470, and E482 situated in the ppGpp-binding domain (amino acids 418-564), and residues N314, D316, and G352 from the PLP-binding domain (amino acids 184-417). D460 from one decamer makes an electrostatic interaction with K422' of a neighbouring decamer in the stack. R468 is sandwiched between D316' and E344' from a neighbouring decamer, and makes electrostatic interactions through its η_1 and η_2 nitrogen atoms with D316'. In addition, D470 interacts with the backbone of G352', and E482 forms hydrogen bonds with N314'. The second interface (Figure 5B) is formed between residue N94 of the wing domain (amino acids 1-130) of one set of opposing dimers, and a stretch of four residues in the ppGpp-binding domain – T444, E445, S446 and D447 - at the end of helix α_{16} from a second set of opposing dimers. The wing domain residue N94 makes hydrogen bonds with E445' of an opposing dimer. A second charged residue, D447, interacts with the backbone of T444', and is held in place by R97 from the wing domain of a neighbouring dimer in the same decamer.

Considering that the Ldcl polymerisation is induced by acid stress, we wondered which residues in the interface would be sensitive to pH changes. Surprisingly, most of the side chains involved in the inter-decamer interface are charged arginine (pKa 13), aspartate and glutamate residues (pKa of 4 and 3 respectively), which do not change protonation state in the pH window relevant for Ldcl activity (pH 5-7) (37). Nonetheless, other residues, situated outside the interaction interfaces, may drive stack formation through pH-dependent interactions that would in turn lead to the observed inter-monomer rotation and the associated constriction of the Ldcl central cavity, coupled to the alignment of complementary contacts at interfaces one and two (Figure 4D, E). We note for example that H694 should be protonated in the Ldcl stack structure at pH 5.7 but deprotonated in the two ring structures at pH 7.0 and pH 8.5, and that an electrostatic interaction between H694 and D192 situated in the linker region is present in the stack structure only (Figure 5E).

To validate the observed interactions at the inter-decamer interface, and to assess the individual importance of key residues involved in Ldcl stack formation, we constructed four Ldcl point mutants (R468E, R97E, H694A and H694N), two double mutants (R97E/R468E and E445A/D447A) and one triple mutant (E445A/D447A/R468E). The mutants were purified following the protocol for wild-type Ldcl (see Methods), diluted into a buffer at pH 5.7 and observed by ns-EM (Figure 6, Supplementary Figure 7). Although the grid preparation procedure for ns-EM yields stacks that are shorter and more curved and distorted when compared to the cryo-EM data (Figure 4B, Figure 6), our previous observations of the five paralogous *E. coli* amino acid decarboxylases

justify the validity of this approach for a qualitative comparative analysis (6, 37). ns-EM images make immediately apparent that the E445A/D447A double mutant does not have a significantly altered capability of stack formation at pH 5.7 when compared to WT Ldcl, whereas the single R468E mutation is sufficient to completely abolish stack formation. The R97E mutant has a moderate destabilising effect and displays fewer and smaller stacks than the WT Ldcl. Consistently with the major role of R468 in the Ldcl stack formation, the R97E/R468E double mutant and E445A/D447A/R468E triple mutant exclusively occur as decamers at low pH. Altogether, these results reveal that R468 is one of the key determinants of Ldcl stack formation. Finally, similarly to the R97E mutant, a modest destabilisation of stack formation is observed for the two histidine mutants, H694A and H694N, favouring our hypothesis that H694 may have an influence on the propensity of Ldcl polymerisation at low pH (Figure 6, Supplementary Figure 7).

Discussion

Our synergistic approach, combining several *in vitro* techniques including biochemical characterisation of purified fluorescent protein fusions, ns-EM observation of mutants, low resolution ns-EM reconstruction and high resolution cryo-EM analysis, with *in vivo* flow cytometry, wide-field and 3D-STORM imaging, provides insights into the supramolecular Ldcl assembly upon acid stress. This work adds to the very few known examples of regular polymerisation as means of regulation of enzymes involved in amino acid metabolism in bacteria. The cryo-EM structure of the Ldcl stacks presented here offers a molecular framework for future investigation of the role of the Ldcl polymerisation in the acid stress response.

The high resolution cryo-EM structures of the stack at pH 5.7 and of the decamer at pH 7.0 complement the previously solved crystal structure of the ppGpp-bound Ldcl decamer obtained at pH 8.5. Previous serendipitous co-crystallisation of Ldcl with the nutrient stress response alarmone ppGpp led to assessment of the effect of ppGpp on Ldcl activity, and to a proposal that ppGpp would act as an Ldcl inhibitor that prevents excessive lysine consumption upon nutrient limitation during acid stress (6). In addition, a ppGpp-dependent disassembly of the Ldcl stacks had been previously observed but could not be structurally explained since the ppGpp binding site is situated between two neighbouring dimers inside the Ldcl ring (6). Our cryo-EM structures show that one of the residues involved both in ppGpp binding and in the stack formation is arginine 97 (R97). In the crystal structure of ppGpp-bound Ldcl, R97 makes a stacking interaction with the guanosine imidazole ring of ppGpp, while in Ldcl stacks R97 is involved in a key interaction at the second interface (Figure 5D), where it locks D447 in a conformation allowing interactions between helices $\alpha 16$ from opposing Ldcl decamers. Binding of ppGpp to Ldcl interferes with the R97-D447 interaction, thereby most likely prohibiting correct positioning of D447 at the tip of helix $\alpha 16$, and resulting in a disruption of the second stack interface, leading to a moderate yet notable stack destabilisation (Figure 6, Supplementary Figure 7). Furthermore, our cryo-EM structure of ppGpp-free Ldcl decamers at neutral pH enables to rule out the effect of ppGpp on the differences observed between the Ldcl stack structure at pH 5.7 and the ppGpp-Ldcl crystal structure at pH 8.5. Indeed, despite the absence of ppGpp, R97 is still oriented towards the ppGpp binding site and away from the inter-decamer interface in the pH 7.0 decamer cryo-EM map (Figure 5D). This suggests that the conformational changes in Ldcl driving stack formation are mostly driven by low

pH and not by the absence of ppGpp, although the D192 and H694 hinge residues are similarly far apart in the pH 8.5 and pH 7.0 structures (Figure 5E). Our current work provides a solid experimental and structural basis for a future closer evaluation of the hypothesised role of ppGpp in LdcI regulation *in vitro* and *in vivo* (6)

Although the optical imaging experiments performed in this study do not allow quantitative evaluation of the nature of the clusters of endogenous LdcI at the bacterial periphery, they would presumably correspond to LdcI assembled into stacks. Possible reasons for such assembly may be to provide an efficient way to locally increase the LdcI concentration and to enhance its activity. But why would LdcI, an apparently highly soluble protein, be driven towards the inner membrane? What would be the advantage for acid-stressed *E. coli* cells to increase the concentration of LdcI via stack formation in these particular peripheral locations? Localisation of proteins to specific sites in the bacterial membrane was shown to be generally driven by chemical factors such as the phospholipid composition of the lipid microdomains, and by physical factors such as the degree of local curvature or the electric potential of the membrane (38). An attractive hypothesis would be that as an acid stress response protein performing a proton-consuming enzymatic reaction, LdcI may be attracted to proton sinks formed by anionic phospholipids which compartmentalise oxidative phosphorylation (OXPHOS) complexes for efficient functioning in bacterial respiration and adaptation to environmental changes. Indeed, OXPHOS complexes were often described to be unevenly distributed in the membrane in the form of mobile patches (39–43), providing evidence for highly dynamic and spatially organised bioenergetic membranes in *E. coli* cells (43). In addition, certain bacterial flotillins, which are essential scaffold proteins of the functional membrane microdomains, equivalent to the lipid rafts of eukaryotic cells, also show a patchy distribution and were shown to interact with specific OXPHOS complexes (43, 44). In this regard, two different lines of evidence would be interesting to note. First, LdcI was described to co-purify with a partially assembled Complex I (45), whereas the LdcI-binding partner RavA, as well as ViaA, the second protein expressed from the *ravAviaA* operon and which also interacts with RavA, were shown to interact with both Complex I and fumarate reductase (32, 33). Second, the other *E. coli* PLP-dependent lysine decarboxylase LdcC, exercising the role of cadaverine biosynthesis irrespectively of acid stress (46), neither binds RavA nor forms stacks (29, 37), in spite of its 69% of identity with LdcI. It is interesting to note that in LdcC the key stack-forming residue R468 has been substituted for an alanine while the interacting D316 has been preserved.

Our STORM images suggest that individual patches have a peripheral distribution with a long-range stripy or pseudo-helical organisation (Figure 3, Supplementary Figure 4, Supplementary Movies 1-5). While similar distributions have been documented for bacterial cytoskeletal, cell division, chromosome partitioning, RNA degradation and secretion machineries, an eventual impact of labelling on these distributions, demonstrated specifically for the YFP-MreB (47), warrants caution in their interpretation (13, 15). Here however, we observed endogenous wild-type LdcI in cells fixed prior to their permeabilisation and labelled with anti-LdcI-Nb, which means that the resulting pattern is likely real. In addition, examination of some published images of OXPHOS patches (for example Llorente-Garcia et al., 2014) also hints to a similar organisation. Excitingly, anionic phospholipid-specific dyes and fluorescently-labelled antibiotics specific for nascent peptidoglycan synthesis upon cell elongation were also proposed to be distributed on

helical or stripe patterns (38, 48–51). It is therefore tempting to imagine that inside the bacterial cell, Ldcl has a tendency to follow a general path upon polymerisation governed by the underlying chiral patterns in the cell envelope (52).

Finally, from the methodological view, our work convincingly illustrates that different FP fusion constructs can share the same cellular distribution in spite of a completely different structure, necessitating caution when inferring intact function from the preservation of the protein localisation inside the cell. Our findings emphasize the importance of characterising FP fusions using both biochemical and structural techniques, such as ns-EM, to ensure that the FP tag disrupts neither structure nor function of the target protein.

Methods

Expression constructs

For fluorescence studies, several FP fusion constructs were generated starting from an available plasmid containing the coding sequence of Ldcl (Uniprot entry P0A9H3), cloned in the pET22b(+) vector with a C-terminal 6xHis-tag (6). All constructs were generated using the Gibson cloning strategy and verified by sequencing analysis. The Gibson assembly was performed using 0,4U T5 exonuclease, 2,5U Phusion polymerase and 400U Taq ligase (New England Biolabs) in 1X ISO buffer (100mM Tris-HCl pH 7,5, 10 mM MgCl₂, 0.8 mM dNTP mix, 10mM DTT, 50 mg PEG-8000, 1 mM NAD). 7.5 µL of the GIBSON Master Mix was mixed with 2.5 µL DNA, containing circa 100 ng of vector. The mix was incubated for 60 min at 50°C. Transformations were performed in Top10 competent bacteria (One Shot™ TOP10 Chemically Competent *E. coli*, Invitrogen) and selected using 100 µg/mL ampicillin or 50 µg/mL kanamycin sulphate (Euromedex). Agarose Gel purification and DNA plasmid extraction kits were purchased from Macherey-Nagel.

Dendra2_{T69A}-Ldcl and mGeosM-Ldcl were both cloned in the pET-TEV vector containing an N-terminal 6xHis-tag, and a TEV cleavage site between Dendra2_{T69A} or mGeosM and the Ldcl gene. Ldcl-Dendra2_{T69A} and Ldcl-mGeosM were both cloned in the pET22b(+) vector with Ldcl followed by either Dendra2_{T69A} or mGeosM containing an uncleavable C-terminal 6XHis-tag.

The anti-Ldcl-Nb was obtained from the nanobody generation platform of the AFMB laboratory (Marseille, France). A llama (*Llama glama*) was immunized with the purified wild type Ldcl. Lymphocytes were isolated from blood samples, a nanobody phage display library was generated, and enrichment of antigen-specific clones was performed using standard procedures (53, 54). Sequences of three positive clones were subcloned in the pHEN6 vector containing the pelB leader sequence from *Erwinia carotovora* for secretion into the periplasm, and a C-terminal 6xHis-tag (kindly provided by Dr. Aline Desmyter, AFMB Marseille). One of the clones was used for the present study to yield anti-Ldcl-Nb.

Plasmids, primers and cloning strategy are summarised in Supplementary Table 4.

Protein purification

Ldcl-FP fusions and Ldcl mutants were expressed in BL21(DE3) cells grown in LB medium supplemented with 100 µg/mL ampicillin or 50 µg/mL kanamycin sulphate. Protein expression was induced using 40 µM IPTG (Euromedex) and carried out overnight at 18°C. The Ldcl-FP fusions

and Ldcl mutants were purified as previously described for wild-type Ldcl (28, 29, 37), in a final buffer containing 25 mM Tris (pH 7.5), 0.3 M NaCl, 5% glycerol, 1 mM DTT and 0.1 mM PLP.

The anti-Ldcl-Nb was expressed in *E. coli* WK6 cells following the protocol described by (53), and purified with immobilised metal affinity chromatography (IMAC, using a Ni-NTA column) followed by Size exclusion chromatography (SEC) using a a superdex 75 Increase 10/300GL column (GE-Healthcare) equilibrated with a buffer containing 25 mM Tris pH 7.4 and 0.3 M NaCl.

In order to characterise the Ldcl complex with anti-Ldcl-Nb, the two proteins were mixed at a 1:5 molar ratio and submitted to size exclusion chromatography as carried out for Ldcl alone but without DTT in the buffer. The top of the peak was taken for subsequent ns-EM analysis.

ns-EM on Ldcl-FP fusions, Ldcl/anti-Ldcl-Nb and Ldcl mutants

FP fusion samples after gel filtration were diluted to a concentration of approximately 0.025 mg/mL. 3 μ L was applied to the clean side of carbon on a carbon–mica interface and stained with 2% uranyl acetate (mGeosM-Ldcl, Ldcl-mGeosM, Ldcl-Dendra2_{T69A}, Ldcl/anti-Ldcl-Nb, Ldcl mutants) or 2% sodium silicotungstate (Dendra2_{T69A}-Ldcl). Images were collected on a 120 kV Tecnai T12 microscope with an Orius 1000 camera (Gatan) or on a 200 kV Tecnai F20 electron microscope with either a OneView camera (Gatan) or a Ceta camera (Thermo Scientific). All images were collected with a defocus range of approximately -1.0 μ m to -2.0 μ m and with pixel sizes between 2.29 Å/pixel and 3.42 Å/pixel.

Image processing – Ldcl-FP and Ldcl/anti-Ldcl-Nb

36 micrographs of Dendra2_{T69A}-Ldcl with a pixel size of 2.73 Å/pixel, 368 micrographs of mGeosM-Ldcl with a pixel size of 2.29 Å/pixel, 23 micrographs of Ldcl-Dendra2_{T69A}, 92 micrographs of Ldcl-mGeosM with a pixel size of 3.42 Å/pixel, and 124 micrographs of Ldcl/anti-Ldcl-Nb with a pixel size of 2.82 Å/pixel were used for image analysis.

CTF estimation was performed with CTFFIND3 (55). Semi-automatic particle selection was carried out with BOXER (56), with box sizes of 98 x 98 pixels for Dendra2_{T69A}-Ldcl, 180 x 180 pixels for mGeosM-Ldcl, 128 x 128 pixels for Ldcl-Dendra2_{T69A} and Ldcl-mGeosM, and 112 x 112 pixels for Ldcl/anti-Ldcl-Nb respectively. Particle extraction followed by several rounds of cleaning by 2D classification in RELION-1.4 (57), resulted in the following number of particles for each dataset: Dendra2_{T69A}-Ldcl = 7140, mGeosM-Ldcl = 5514, Ldcl-Dendra2_{T69A} = 832, Ldcl-mGeosM = 12,211 and Ldcl/anti-Ldcl-Nb = 14,075.

For Dendra2_{T69A}-Ldcl, initial model generation was carried out in RELION-2.1 (58) without any symmetry applied. For mGeosM-Ldcl, initial model generation was carried out in RELION-2.1 with either C1, C3 or D3 symmetry applied. The results of all three calculations being very similar, the model with applied D3 symmetry was selected. For Ldcl-Dendra2_{T69A}, Ldcl-mGeosM and Ldcl/anti-Ldcl-Nb, the previously-determined Ldcl decamer structure (PDB ID: 3N75) was filtered to 60 Å and used as an initial model for 3D refinement.

3D refinement was carried out for each dataset with applied C2 symmetry for Dendra2_{T69A}-Ldcl, D3 symmetry for mGeosM-Ldcl, and D5 symmetry for Ldcl-Dendra2_{T69A}, Ldcl-mGeosM and Ldcl/anti-Ldcl-Nb. Rigid body fitting of Ldcl (PDB ID: 3N75), mEos2 (PDB ID: 3S05) and Dendra2 (PDB ID: 2VZX) crystal structures was then carried out in Chimera (59) for the five datasets.

Cryo-EM on Ldcl stacks (pH 5.7)

Wild-type Ldcl was purified as previously described (10) from an *E. coli* strain impaired in the production of ppGpp (MG1655 $\Delta relA\Delta spoT$) in order to avoid any serendipitous ppGpp binding. Purified Ldcl was diluted to a final concentration of approximately 0.25 mg/mL in a buffer containing 25 mM MES (pH 5.7), 0.3 M NaCl, 5% glycerol, 1 mM DTT and 0.1 mM PLP. 3 μ L of the sample was applied to a glow-discharged R2/1 300 mesh holey carbon copper grid (Quantifoil Micro Tools GmbH) and plunge-frozen in liquid ethane using a Vitrobot Mark IV (FEI) operated at 100% humidity. Datasets were recorded at the European Synchrotron Radiation Facility (ESRF) in Grenoble, France (60), on a Titan Krios microscope (Thermo Scientific) equipped with a BioQuantum LS/967 energy filter (Gatan) and a K2 summit direct electron detector (Gatan) operated in counting mode. A total of 2564 movies of 30 frames were collected with a total exposure of 6 s, total dose of 29.3 e⁻/Å² and a slit width of 20 eV for the energy filter. All movies were collected at a magnification of 130,000x, corresponding to a pixel size of 1.052 Å/pixel at the specimen level. A summary of cryo-EM data collection parameters can be found in Supplementary Table 1.

Image Processing – Ldcl stacks (pH 5.7)

Motion correction and dose-weighting of the recorded movies were performed using MotionCor2 (61). CTF parameters were determined on the aligned and dose-weighted sums using CTFFIND4 (62). After manual inspection of the dose-weighted sums, the best 558 (21.7%) micrographs were selected for further processing. Ldcl stacks were manually picked using e2helixboxer in EMAN2 (63). A total of 15,165 Ldcl-stack particles were extracted in RELION-3.0 (64) with an extract size of 320 pixels, resulting in boxes containing three Ldcl decamers, and with the --helix option with an outer diameter of 160 pixels and a helical rise of 77 pixels. After particle extraction, per-particle CTF correction was performed using Gctf (65). Extracted particles were subjected to 2D classification in RELION-3.0, resulting in a cleaned dataset containing 15,157 particles. Initial 3D refinement with imposed D5 symmetry was carried out in RELION-3.0, using an initial model generated by manually stacking three Ldcl decamers (PDB ID: 3N75) in Chimera (59) and low-pass filtering the resulting Ldcl-stack to 40 Å. The resulting 4.3 Å resolution 3D reconstruction, along with the cleaned particle stack, was subsequently imported into CryoSPARC (66). A final homogeneous 3D refinement in CryoSPARC, using a dynamic mask and imposing D5 symmetry, resulted in a map with a resolution of 3.28 Å based on the 0.143 gold-standard Fourier shell correlation (FSC) criterion (67). The final map was sharpened using a B-factor of -96 Å². A local resolution estimation of the final 3D reconstruction was calculated in RELION-3.0. A summary of cryo-EM data collection parameters and image processing steps for Ldcl stacks can be found in Supplementary Table 1 and Supplementary Figure 4, with local resolution and FSC curves shown in Supplementary Figure 5.

Cryo-EM on Ldcl decamers (pH 7.0)

Purified Ldcl was dialysed into a buffer at pH 7.0 and diluted to a concentration of 0.25 mg/mL. 3 μ L of diluted sample was applied to a glow-discharged (R2/1 300 mesh holey carbon

copper grid (Quantifoil Micro Tools GmbH) and plunge-frozen in liquid ethane using a Vitrobot Mark IV (FEI) operated at 100% humidity. Images were recorded on a Glacios microscope (Thermo Scientific) equipped with a Falcon II direct electron detector (Thermo Scientific). A total of 2772 movies of 29 frames were collected with a total exposure of 1.5 s and a total dose of 45 e⁻/Å². All movies were collected at a magnification of 116,086x, corresponding to a pixel size of 1.206 Å/pixel at the specimen level.

Image Processing – Ldcl decamers (pH 7.0)

Motion correction was carried out using patch motion correction in CryoSPARC, discarding the first two frames. Initial CTF estimation was then carried out on summed frames using CTFFIND4. A subset of 600 micrographs were subjected to automatic picking using the blob picker in CryoSPARC, resulting in ~238,000 picked particles. Particles were then extracted with a box size of 256 x 256 and subjected to 2D classification. Particles from the best classes showing clear secondary structural features for Ldcl were selected for homogeneous refinement (with applied D5 symmetry) against EMD-3204 low-pass filtered to 30 Å, resulting in a reconstruction with a resolution of 4.2 Å (FSC = 0.143). This reconstruction was then used to create templates for picking the entire dataset using the template picker in CryoSPARC, after filtering to 12 Å. ~796,000 particles were extracted and subjected to 2D classification, and the best ~428,000 particles were subjected to heterogeneous refinement with applied D5 symmetry against the 4.2 Å map, resulting in one higher-resolution class corresponding to ~229,000 particles. These particles were subjected to homogeneous refinement with applied D5 symmetry, resulting in a map with a resolution of 2.76 Å (FSC = 0.143) which was then sharpened with a B-factor of -173 Å². A summary of cryo-EM data collection parameters and image processing steps for Ldcl decamers can be found in Supplementary Table 1 and Supplementary Figure 4, with local resolution and FSC curves shown in Supplementary Figure 5.

Fitting of structures and refinement

For fitting of atomic models in the 3D reconstructions of Ldcl at pH 5.7 (stack) or pH 7.0 (decamer), two (for the stack) or one copy (for the decamer) of the Ldcl X-ray crystal structure (PDB ID: 3N75) were first rigid-body fitted in the corresponding 3D reconstructions using Chimera. Refinement was performed using the Phenix software package (68) and was identical for both 3D reconstructions. A first round of real space refinement was carried out with enabled rigid-body, global minimization, local grid search and ADP refinement parameters, and imposing rotamer, Ramachandran, NCS and reference model (PDB ID: 3N75) restraints. A final round of real space refinement was then performed using the same settings, but without rigid body refinement and without applying reference restraints, setting the 'nonbonded_weight' parameter to 4000 and disabling 'local_grid_search'. A summary of refinement and model validation statistics can be found in Supplementary Table 1.

pH shift experiment

Stationary phase cultures, which were grown overnight from single colonies in LB medium, were diluted to OD600 ~ 0.01 and re-grown at 37°C within approximately 1h 45 min in fresh LB

medium to an OD600 of 0.1. From this culture, 14 mL were transferred to 15 mL falcon tubes and pelleted by centrifugation at RT for 5 min. The supernatant was decanted, whereby systematically around 200 μ L LB remained in the falcon tube. The pellet of bacteria was resuspended and afterwards, LB-4.6 containing 30 mM L-lysine was added to the cells up to 14 mL. To prepare LB-4.6 medium, LB powder (Sigma-Aldrich) was completely dissolved in distilled water by stirring for 30 min. The pH of 4.6 was then adjusted using HCl. After autoclaving, sterile filtered L-lysine was added to LB-4.6. For this, L-lysine was dissolved in an aliquot of LB-4.6, sterile filtered and mixed with the remaining LB-4.6. 30 mM L-lysine were used. In order to grow the culture under oxygen-limiting conditions, the lid of the falcon tube was closed and the tubes were placed at 37°C on a shaker (150 rpm). After defined time-points, aliquots were taken for OD600 measurement, pH measurement, SDS-PAGE or immunofluorescence. For each time-point, a new tube was opened and not reused further.

Western Blotting

SDS-PAGE was performed with a Biorad electrophoresis chamber using standard 12% reducing SDS-PAGE gels. Proteins were transferred to a nitrocellulose membrane (Biorad) using a Trans-Blot Turbo Transfer System (Biorad). The membrane was blocked for 1h using 5% BSA in TBS supplemented with 0.1% Tween (TBS-T). Afterwards, the membrane was incubated for 1h with an anti-Ldcl antibody (Qalam-Antibodies) in BSA/TBS-T (1:5000). The membrane was subsequently washed 3 x 10 min in TBS-T and incubated for 1 h with HRP-coupled anti-rabbit antibody (1:10000 in BSA/TBS-T). Finally the membrane was washed 3 x 10 min in TBS-T. The membrane was rinsed once with TBS, prior to detection of antibody-labelled proteins using ECL reagent (GE Healthcare). Incubation and detection of antibody was performed at 25 °C.

Nanobody labelling

For the labelling reaction, 50 μ L nanobody (i.e. 200 μ g) was pipetted into a 1.5 mL tube, placed on ice and supplemented with 5.5 μ L of 1 M bicarbonate buffer at pH 8.3. 100 μ g of Alexa-647 or Alexa-488 NHS ester dye (Life Technologies, A37573 and A20000) were dissolved in 10 μ L DMSO to final concentration of 10 mg/mL. 5 μ L (i.e. 40 nmol) dye in DMSO was added to the protein and incubated for 1 h at room temperature in a shaking block, covered with an aluminium foil to protect the dye from the light. Excess dye was removed by iterative buffer exchange using a 3K spin column (Amicon-Ultra-4 Centrifugal Filters Ultracell 3K, Millipore UFC800396) to PBS. The degree of labelling was inferred from measuring the OD.

Cell preparation for immunofluorescence staining of *E. coli* cells with nanobodies

For immunofluorescence staining, cells were subjected to a pH shift in order to induce Ldcl expression, as described in the “pH shift experiment” section above. 90 minutes after the pH shift, the OD600 of the cell culture was measured, and the volume of cells corresponding to OD = 4, with OD = 1 corresponding to about 8×10^8 cells per mL, were collected by centrifugation. After removal of LB by pipetting, cells were resuspended in 2 mL of 4 % FA in PBS (made from 16% Formaldehyde solution, Methanol-free from Thermo Scientific). Falcons were placed on a rotor for constant agitation for 45 min at room temperature. After fixation, cells were collected by

centrifugation and the solution was removed by pipetting. Cells were then resuspended in 14 mL PBS (Gibco, Thermo Scientific) to remove and dilute the fixative. Cells were permeabilised for 10 min using 2 mL 0.1 % Triton X-100 in PBS and subsequently washed three times with 10 mL PBS. Finally, cells were transferred to 1.5 mL tubes, centrifuged and resuspended in 200 μ L 1% BSA/PBS (BSA/PBS solution was dissolved for 30 min and sterile-filtered to avoid clumps). After 30 min of incubation, 0.5 μ g anti-Ldcl-Nb, labelled with the dedicated dye, was added to the 200 μ L bacteria-BSA/PBS suspension. Cells were incubated with the labelled anti-Ldcl-Nb for 16 h at 4°C. The next day, cells were washed three times with 1 mL PBS, centrifuged to remove antibody solution and resuspended in 250 μ L PBS. When needed, Hoechst 33342 (Sigma-Aldrich) was added to a final concentration of about 100 ng/mL.

Wide-field imaging and flow cytometry

For epifluorescence imaging, 2 μ L of cells were placed between a glass slide and a coverslip, which have been carefully pressed together, and observed using an inverted IX81 microscope, with a UPLFLN 100X oil immersion objective (N.A. 1.3) (Olympus), using the appropriate specific excitation and emission filters for AF488 (GFP-3035B set, Semrock) and DAPI (DAPI-5060B set, Semrock). Acquisitions were performed with Velocity software (QuorumTechnologies™) with a sCMOS 2048x2048 camera (Hamamatsu ORCA Flash 4, 16 bits/pixel) achieving a final magnification of 64 nm per pixel. For flow cytometry, 50 μ L of cells suspensions were injected in a MACSQuant VYB flow cytometer (Miltenyi Biotech, Bergish Gladbach, Germany) using the 488 nm excitation and 525 (50) nm emission channel (B1). AF488 positive populations were estimated after forward scatter (FSC) and side scatter (SSC) gating on the cells. Data were further processed with MACSQuantify software (Miltenyi Biotech).

STORM imaging

For Single Molecule Localization Microscopy, cells were transferred to a glucose buffer containing 50 mM NaCl, 150 mM Tris (pH 8.0), 10% Glucose, 100 mM MEA (Mercaptoethylamine) and 1x Glox. Glox was prepared as a 10x stock and contained 1 μ M catalase and 2.3 μ M glucoseoxidase. Menzel glass slides (Thermo Scientific) and precision coverslips (1.5H from ThorLabs) were cleaned for 30 min using an UV-ozone cleaning device (HELIOS-500, UVOTECH Systems). 2 μ L of immuno-labelled cells were placed onto a glass slide and covered with the coverslip, then cells were carefully spread by pressing glass slides firmly together and the sides were sealed with transparent nail polish to avoid evaporation. Mounted samples were imaged on a homemade SMLM set up based on an IX81 microscope (Olympus). STORM was performed by focusing a 643 nm excitation laser beam (Toptica Diode laser) to the back focal plane of an oil immersion UAPON100X (N.A. 1.49) objective. The intensity of the laser was tuned using an Acousto-Optical Tunable Filter (OATF, Quanta Tech). Acquisition was obtained with a 16 bits/pixel Evolve 512 EMCCD (Photometrics) using Metamorph (Molecular Devices), for a final pixel size of 123 nm. 3D-STORM based on point-spread-function astigmatism (69) was performed using a cylindrical lens (LJ1516L1-A, Thorlabs) placed in the detection light path. STORM datasets consisting of about 30,000 frames, using a 643 nm laser power density of 3 kW/cm² and a 405 nm laser power density of up to 1 W/cm² and a framerate of 50 ms, were acquired with an EMCCD gain set to 200. 3D

point spread function calibration was achieved using tetraspec beads. Finally, data were processed with the Thunderstorm plugin (70) in ImageJ (71). 3D-images were rendered with Visp (72) using a minimum neighbour density threshold of 20 to 28. A total of about 25,000 localisations with a median localisation precision of ~17 nm were recorded per cell. A quantitative evaluation of molecular copy number from STORM data is notoriously difficult due to pronounced and potentially environmentally-dependent blinking of AF647. However, assuming an average of 3 blinking events per molecule and 100% efficient labelling, one can estimate the number of Ldcl decamers in the range 500-1000 per cell, which would agree very well with earlier independent estimations (6, 31). Therefore, although the Ldcl labelling efficiency *in cellulo* was not precisely determined in this study, it is likely to be quite high, which is also consistent with our *in vitro* structural data on the Ldcl-nanobody complex.

Biolayer interferometry measurements

For BLI binding studies, RavA with a biotinylated C-terminal AviTag was expressed and purified as previously described (30). BLI experiments were performed in 1x HBS pH at 7.0 (25 mM HEPES, 300 mM NaCl, 10 mM MgCl₂, 10% glycerol) supplemented with 1x kinetics buffer (0.1% w/v BSA, 0.02% v/v Tween-20), 1mM ADP, 1mM DTT and 0.1 mM PLP. Experiments were performed using the BLItz System instrument (FortéBio), operated at room temperature. Before the start of each BLI experiment, RavA-AviTag was incubated with 1 mM ADP for 10 min. Streptavidin-coated biosensors (FortéBio) were functionalised with biotinylated RavA-AviTag, then quenched with 10 µg/mL biocytin. Experiments with the wild-type Ldcl are from (30). For C-terminal Ldcl-FP fusions, pins were dipped in wells containing a range of Ldcl-FP concentrations from 0 to 1000 nM, with no binding signal recorded at any concentration of Ldcl-FP.

Data and code availability

Cryo-EM maps, along with the corresponding fitted atomic structures, have been submitted to the EMDB and PDB with accession codes EMD-10850 and PDB-6YN6 for Ldcl stacks (pH 5.7), and EMD-10849 and PDB-6YN5 for Ldcl decamers (pH 7.0).

Acknowledgements

We thank Guy Schoehn for establishing and managing the cryo-electron microscopy platform and for providing training and support, and Rose-Laure Revel-Goyet and Françoise Lacroix for the support and access to the M4D Cell imaging Platform. We are grateful to Aymeric Peuch for help with the usage of the EM computing cluster and Daniel Thédie, Kévin Floc'h and Joel Beaudoin for discussions. We thank Alain Roussel and Aline Desmyter for the nanobody production, and Julien Perard for help with the cloning of the Dendra2_{T69A}-Ldcl construct. We acknowledge the European Synchrotron Radiation Facility for provision of beam time on CM01. This work was funded by the European Union's Horizon 2020 research and innovation programme under grant agreement No 647784 to IG. The nanobody generation platform of the AFMB laboratory (Marseille, France) was supported by the French Infrastructure for Integrated Structural Biology (FRISBI) ANR-10-INSB-05-01. For electron and fluorescence microscopy studies, this work used the platforms of the Grenoble Instruct centre (ISBG; UMS 3518 CNRS-CEA-UJF-EMBL)

with support from FRISBI (ANR-10-INSB-05-02) and GRAL (ANR-10-LABX-49-01) within the Grenoble Partnership for Structural Biology (PSB). The electron microscope facility (Glacios electron microscope) is supported by the Rhône-Alpes Region (CIBLE and FEDER), the FRM, the CNRS, the University of Grenoble and the GIS-IBISA. JF was supported by a long-term EMBO fellowship (ALTF441-2017) and a Marie Skłodowska-Curie actions Individual Fellowship (789385, RespViRALI).

Author Contributions

C.L., M.B., V.A., A.F. and K.H. cloned constructs, M.J., A.F. and K.H. purified proteins. M.J. and I.G. performed ns-EM imaging and analysis. A.D., G.E. and M.B.-V. performed cryo-EM imaging. J.F. and A.D. performed cryo-EM analysis. M.J. and J.F. built models resulting from EM maps. M.J., J.F. and I.G. analysed structures and interpreted data. C.L. and M.B. performed optical imaging of overexpression constructs with input of V.A., J.-P.K. and D.B. C.L. performed nanobody characterisation for optical imaging with input from J.-P.K. C.L. analysed endogenous expression and performed optical imaging of endogenous LdcI with input of V.A., J.-P.K. and D.B. C.L. and D.B. analysed STORM Images. M.J., C.L., J.F. and I.G. analysed the data and prepared the figures and tables. C.L. and D.B. contributed to the design of the optical imaging part of the project together with I.G. I.G. designed, supervised and funded the overall study. M.J., J.F. and I.G. wrote the manuscript with significant input from C.L. and contributions from all of the authors.

References

1. M. Prouteau, R. Loewith, Regulation of Cellular Metabolism through Phase Separation of Enzymes. *Biomolecules* **8**, 160 (2018).
2. F. C. Fang, E. R. Frawley, T. Tapscott, A. Vázquez-Torres, Bacterial Stress Responses during Host Infection. *Cell Host Microbe* **20**, 133–143 (2016).
3. N. Watson, D. S. Dunyak, E. L. Rosey, J. L. Slonczewski, E. R. Olson, Identification of elements involved in transcriptional regulation of the Escherichia coli cad operon by external pH. *J. Bacteriol.* **174**, 530–540 (1992).
4. J. W. Foster, Escherichia coli acid resistance: Tales of an amateur acidophile. *Nat. Rev. Microbiol.* **2**, 898–907 (2004).
5. P. L. Moreau, The lysine decarboxylase CadA protects Escherichia coli starved of phosphate against fermentation acids. *J. Bacteriol.* **189**, 2249–2261 (2007).
6. U. Kanjee, *et al.*, Linkage between the bacterial acid stress and stringent responses: The structure of the inducible lysine decarboxylase. *EMBO J.* **30**, 931–944 (2011).
7. E. F. Gale, The Bacterial Amino Acid Decarboxylases. *Adv. Enzymol. - Relat. Areas Mol. Biol.* **6**, 1–32 (1946).
8. D. L. Sabo, E. A. Boeker, B. Byers, H. Waron, E. H. Fischer, Purification and Physical Properties of Inducible Escherichia coli Lysine Decarboxylase. *Biochemistry* **13**, 662–670 (1974).
9. E. F. Gale, H. M. R. Epps, The effect of the pH of the medium during growth on the enzymic activities of bacteria (Escherichia coli and Micrococcus lysodeikticus) and the biological significance of the changes produced. *Biochem. J.* **36**, 600–618 (1942).
10. E. Kandiah, *et al.*, Structural insights into the Escherichia coli lysine decarboxylases and molecular

determinants of interaction with the AAA+ ATPase RavA. *Sci. Rep.* **6** (2016).

11. B. Zhao, Investigation of the Regulation of the Lysine Decarboxylase LdcI Activity by the Alarmones ppGpp and MoxR Family AAA + ATPase RavA (2012).
12. I. V. Surovtsev, C. Jacobs-Wagner, Subcellular Organization: A Critical Feature of Bacterial Cell Replication. *Cell* **172**, 1271–1293 (2018).
13. W. Margolin, The price of tags in protein localization studies. *J. Bacteriol.* **194**, 6369–6371 (2012).
14. A. Gahlmann, W. E. Moerner, Exploring bacterial cell biology with single-molecule tracking and super-resolution imaging. *Nat. Rev. Microbiol.* **12**, 9–22 (2014).
15. D. Landgraf, B. Okumus, P. Chien, T. A. Baker, J. Paulsson, Segregation of molecules at cell division reveals native protein localization. *Nat. Methods* **9**, 480–482 (2012).
16. L. M. Costantini, M. Fossati, M. Francolini, E. L. Snapp, Assessing the Tendency of Fluorescent Proteins to Oligomerize Under Physiologic Conditions. *Traffic* **13**, 643–649 (2012).
17. M. Ingerson-Mahar, A. Briegel, J. N. Werner, G. J. Jensen, Z. Gitai, The metabolic enzyme CTP synthase forms cytoskeletal filaments. *Nat. Cell Biol.* **12**, 739–746 (2010).
18. E. M. Lynch, *et al.*, Human CTP synthase filament structure reveals the active enzyme conformation. *Nat. Struct. Mol. Biol.* **24**, 507–514 (2017).
19. T. G. Frey, D. Eisenberg, F. A. Eiserling, Glutamine synthetase forms three and seven stranded helical cables. *Proc. Natl. Acad. Sci. U. S. A.* **72**, 3402–3406 (1975).
20. G. Kim, *et al.*, Aldehyde-alcohol dehydrogenase forms a high-order spiroosome architecture critical for its activity. *Nat. Commun.* **10** (2019).
21. K. Schuchmann, J. Vonck, V. Müller, A bacterial hydrogen-dependent CO₂ reductase forms filamentous structures. *FEBS J.* **283**, 1311–1322 (2016).
22. Z. Yao, R. Carballido-López, Fluorescence Imaging for Bacterial Cell Biology: From Localization to Dynamics, From Ensembles to Single Molecules. *Annu. Rev. Microbiol.* **68** (2014).
23. M. J. Rust, M. Bates, X. Zhuang, Sub-diffraction-limit imaging by stochastic optical reconstruction microscopy (STORM). *Nat. Methods* **3**, 793–795 (2006).
24. E. Betzig, *et al.*, Imaging intracellular fluorescent proteins at nanometer resolution. *Science* **313**, 1642–5 (2006).
25. S. T. Hess, T. P. K. Girirajan, M. D. Mason, Ultra-high resolution imaging by fluorescence photoactivation localization microscopy. *Biophys. J.* **91**, 4258–4272 (2006).
26. H. Chang, *et al.*, A unique series of reversibly switchable fluorescent proteins with beneficial properties for various applications. *Proc. Natl. Acad. Sci.* **109**, 4455–4460 (2012).
27. R. Berardozi, V. Adam, A. Martins, D. Bourgeois, Arginine 66 Controls Dark-State Formation in Green-to-Red Photoconvertible Fluorescent Proteins. *J. Am. Chem. Soc.* **138**, 558–565 (2016).
28. H. Malet, *et al.*, Assembly principles of a unique cage formed by hexameric and decameric E. coli proteins. *Elife* **3**, e03653 (2014).
29. J. Snider, *et al.*, Formation of a distinctive complex between the inducible bacterial lysine decarboxylase and a novel AAA+ ATPase. *J. Biol. Chem.* **281**, 1532–1546 (2006).
30. M. Jessop, *et al.*, Structural insights into ATP hydrolysis by the MoxR ATPase RavA and the LdcI-RavA cage-like complex. *Commun. Biol.* **3**, 46 (2020).
31. M. El Bakkouri, *et al.*, Structure of RavA MoxR AAA+ protein reveals the design principles of a molecular cage modulating the inducible lysine decarboxylase activity. *Proc. Natl. Acad. Sci.* **107**,

22499–22504 (2010).

32. K. S. Wong, *et al.*, The MoxR ATPase RavA and its cofactor ViaA interact with the NADH: Ubiquinone oxidoreductase I in Escherichia coli. *PLoS One* **9**, e85529 (2014).

33. K. S. Wong, V. Bhandari, S. C. Janga, W. A. Houry, The RavA-ViaA Chaperone-Like System Interacts with and Modulates the Activity of the Fumarate Reductase Respiratory Complex. *J. Mol. Biol.* **429**, 324–344 (2017).

34. M. Zhang, *et al.*, Rational design of true monomeric and bright photoactivatable fluorescent proteins. *Nat. Methods* **9** (2012).

35. C. Coltharp, J. Xiao, Superresolution microscopy for microbiology. *Cell. Microbiol.* **14**, 1808–1818 (2012).

36. K. Cramer, *et al.*, Visualization of Bacterial Protein Complexes Labeled with Fluorescent Proteins and Nanobody Binders for STED Microscopy. *Int. J. Mol. Sci.* **20**, 3376 (2019).

37. U. Kanjee, I. Gutsche, S. Ramachandran, W. A. Houry, The enzymatic activities of the escherichia coli basic aliphatic amino acid decarboxylases exhibit a pH zone of inhibition. *Biochemistry* **50**, 9388–9398 (2011).

38. I. Barák, K. Muchová, The role of lipid domains in bacterial cell processes. *Int. J. Mol. Sci.* **14**, 4050–4065 (2013).

39. T. Lenn, M. C. Leake, C. W. Mullineaux, Clustering and dynamics of cytochrome bd-I complexes in the Escherichia coli plasma membrane in vivo. *Mol. Microbiol.* **70**, 1397–1407 (2008).

40. H. Erhardt, *et al.*, Organization of the Escherichia coli aerobic enzyme complexes of oxidative phosphorylation in dynamic domains within the cytoplasmic membrane. *Microbiologyopen* **3**, 316–326 (2014).

41. I. Llorente-Garcia, *et al.*, Single-molecule in vivo imaging of bacterial respiratory complexes indicates delocalized oxidative phosphorylation. *Biochim. Biophys. Acta - Bioenerg.* **1837**, 811–824 (2014).

42. J. N. Werner, *et al.*, Quantitative genome-scale analysis of protein localization in an asymmetric bacterium. *Proc. Natl. Acad. Sci.* **106**, 7858–7863 (2009).

43. A. Magalon, F. Alberge, Distribution and dynamics of OXPHOS complexes in the bacterial cytoplasmic membrane. *Biochim. Biophys. Acta - Bioenerg.* **1857**, 198–213 (2016).

44. M. Bramkamp, D. Lopez, Exploring the Existence of Lipid Rafts in Bacteria. *Microbiol. Mol. Biol. Rev.* **79**, 81–100 (2015).

45. H. Erhardt, *et al.*, Disruption of individual nuo-genes leads to the formation of partially assembled NADH:ubiquinone oxidoreductase (complex I) in Escherichia coli. *Biochim. Biophys. Acta* **1817**, 863–71 (2012).

46. Y. Yamamoto, Y. Miwa, K. Miyoshi, J. I. Furuyama, H. Ohmori, The Escherichia coli ldcC gene encodes another lysine decarboxylase, probably a constitutive enzyme. *Genes Genet. Syst.* **72**, 167–172 (1997).

47. M. T. Swulius, G. J. Jensen, The helical mreB cytoskeleton in Escherichia coli MC1000/pLE7 is an artifact of the N-terminal yellow fluorescent protein tag. *J. Bacteriol.* **194**, 6382–6386 (2012).

48. I. Barák, K. Muchová, A. J. Wilkinson, P. J. O'Toole, N. Pavlendová, Lipid spirals in Bacillus subtilis and their role in cell division. *Mol. Microbiol.* **68**, 1315–1327 (2008).

49. E. Mileyskaya, W. Dowhan, Cardiolipin membrane domains in prokaryotes and eukaryotes. *Biochim. Biophys. Acta - Biomembr.* **1788**, 2084–2091 (2009).

50. I. Fishov, C. L. Woldringh, Visualization of membrane domains in Escherichia coli. *Mol. Microbiol.* **32**, 1166–1172 (1999).
51. R. A. Daniel, J. Errington, Control of cell morphogenesis in bacteria: Two distinct ways to make a rod-shaped cell. *Cell* **113**, 767–776 (2003).
52. S. Wang, L. Furchtgott, K. C. Huang, J. W. Shaevitz, Helical insertion of peptidoglycan produces chiral ordering of the bacterial cell wall. *Proc. Natl. Acad. Sci. U. S. A.* **109** (2012).
53. E. Pardon, *et al.*, A general protocol for the generation of Nanobodies for structural biology. *Nat. Protoc.* **9**, 674–693 (2014).
54. A. Desmyter, *et al.*, Viral infection modulation and neutralization by camelid nanobodies. *Proc. Natl. Acad. Sci. U. S. A.* (2013) <https://doi.org/10.1073/pnas.1301336110>.
55. J. A. Mindell, N. Grigorieff, Accurate determination of local defocus and specimen tilt in electron microscopy. *J. Struct. Biol.* **142**, 334–347 (2003).
56. S. J. Ludtke, P. R. Baldwin, W. Chiu, EMAN: Semiautomated Software for High-Resolution Single-Particle Reconstructions. *J. Struct. Biol.* **128**, 82–97 (1999).
57. S. H. W. Scheres, RELION: Implementation of a Bayesian approach to cryo-EM structure determination. *J. Struct. Biol.* **180**, 519–530 (2012).
58. S. H. W. Scheres, “Processing of Structurally Heterogeneous Cryo-EM Data in RELION” in *Methods in Enzymology*, (2016), pp. 125–157.
59. E. F. Pettersen, *et al.*, UCSF Chimera--a visualization system for exploratory research and analysis. *J. Comput. Chem.* **25**, 1605–12 (2004).
60. E. Kandiah, *et al.*, CM01: A facility for cryo-electron microscopy at the European synchrotron. *Acta Crystallogr. Sect. D Struct. Biol.* **75** (2019).
61. S. Q. Zheng, *et al.*, MotionCor2: Anisotropic correction of beam-induced motion for improved cryo-electron microscopy. *Nat. Methods* **14**, 331–332 (2017).
62. A. Rohou, N. Grigorieff, CTFFIND4: Fast and accurate defocus estimation from electron micrographs. *J. Struct. Biol.* **192**, 216–221 (2015).
63. G. Tang, *et al.*, EMAN2: An extensible image processing suite for electron microscopy. *J. Struct. Biol.* **157**, 38–46 (2007).
64. J. Zivanov, *et al.*, New tools for automated high-resolution cryo-EM structure determination in RELION-3. *Elife* **7**, 1–22 (2018).
65. K. Zhang, Gctf: Real-time CTF determination and correction. *J. Struct. Biol.* **193**, 1–12 (2016).
66. A. Punjani, J. L. Rubinstein, D. J. Fleet, M. A. Brubaker, CryoSPARC: Algorithms for rapid unsupervised cryo-EM structure determination. *Nat. Methods* **14**, 290–296 (2017).
67. P. B. Rosenthal, R. Henderson, Optimal determination of particle orientation, absolute hand, and contrast loss in single-particle electron cryomicroscopy. *J. Mol. Biol.* **333**, 721–745 (2003).
68. P. D. Adams, *et al.*, PHENIX: A comprehensive Python-based system for macromolecular structure solution. *Acta Crystallogr. Sect. D Biol. Crystallogr.* **66**, 213–221 (2010).
69. B. Huang, W. Wang, M. Bates, X. Zhuang, Three-dimensional super-resolution imaging by stochastic optical reconstruction microscopy. *Science (80-.)*. **319** (2008).
70. M. Ovesný, P. Křížek, J. Borkovec, Z. Švindrych, G. M. Hagen, ThunderSTORM: A comprehensive ImageJ plug-in for PALM and STORM data analysis and super-resolution imaging. *Bioinformatics* **30** (2014).

- 840 71. C. A. Schneider, W. S. Rasband, K. W. Eliceiri, NIH Image to ImageJ: 25 years of image analysis.
841 *Nat. Methods* **9** (2012).
- 842 72. M. El Beheiry, M. Dahan, ViSP: Representing single-particle localizations in three dimensions. *Nat.*
843 *Methods* **10** (2013).
- 844
- 845
- 846

Figure 1.

Figure 1. Fluorescent protein fusions affect LdcI structure without significantly altering cellular localisation of overexpressed constructs. A) Wide-field fluorescence microscopy images of *E. coli* cells overexpressing the fluorescent fusion proteins in A). Green fluorescence = FP-LdcI, blue fluorescence = DAPI-stained DNA. **B)** ns-EM 3D maps of fluorescently-tagged LdcI, with fitted models. Left – Dendra2_{T69A}-LdcI forms dimers, with fluorescent barrels located next to the N-terminus of LdcI as expected. Second from left – mGeosM-LdcI forms large non-native oligomers, composed of three LdcI tetramers bridged by tetramers of mGeosM. Second from right and right – both C-terminal fluorescent fusions LdcI-mGeosM and LdcI-Dendra2_{T69A} form decamers, with protrusions at the C-terminus (coloured in red) attributed to flexibly-linked fluorescent proteins. Fitted PDB models are as follows – LdcI: 3N75; Dendra2: 2VZX; mGeosM: 3S05 (mEos2 crystal structure).

Figure 2.

Figure 2. Anti-Ldcl-Nb is a useful tool to probe cellular localisation of endogenous Ldcl under acid stress conditions. **A)** Ns-EM 3D reconstruction of the Ldcl decorated by anti-Ldcl-Nb. An Ldcl decamer (gold, PDB ID: 3N75) and 10 nanobodies (blue, PDB ID: 1MEL, anti-lysozyme nanobody) are fitted in the density, with the nanobody binding at the top and bottom of the decameric ring. The C-terminal RavA binding site is indicated in red, and is in a spatially distinct location from the bound nanobodies. Scale bar = 100Å. **B)** Wide-field fluorescence microscopy image of wild-type *E. coli* MG1655 cells grown at pH 4.6 for 90 minutes stained with anti-Ldcl-Nb labelled with AF488. Inset – zoom of the image showing punctuate fluorescence patterns. Scale bar = 5 µm.

Figure 3.

Figure 3. 3D STORM imaging of *E. coli* cells stained with anti-Ldcl-Nb reveal a patchy pseudo-helically arranged distribution of endogenous Ldcl upon acid stress. 3D-STORM imaging of native Ldcl in wild-type *E. coli* cells 90 minutes after exposure to acid stress (see Methods) with AF647-conjugated anti-Ldcl-Nb. Points are coloured according to localisation density, with brighter points corresponding to higher localisation density. The centre panel shows four cells in the field of view, looking down the z-axis. Left and right panels show side and tilted views respectively. Scale bar = 1 μ m.

Figure 4.

Figure 4. Cryo-EM analysis of the Ldcl decamer at neutral pH and Ldcl stacks formed in acid stress conditions. A) Cryo-EM reconstruction of the Ldcl decamer at pH 7.0 from side (left) and top (right) views. Scale bar = 50 Å. **B)** Cryo-EM micrograph of Ldcl stacks at pH 5.7, scale bar = 100 nm. Inset – 2D class averages displaying clear secondary structural features, scale bar = 100 Å. **C)** Top (above) and side (below) views of the cryo-EM reconstruction of a three-decamer Ldcl

stack. Four dimers are coloured either blue/gold or green/coral, corresponding to the colouring of the atomic model presented in Figure 5. Scale bar = 50 Å. **D)** Overlay of Ldcl dimers at pH 8.5 (PDB ID: 3N75, shown in lilac), 7.0 (shown in cyan) and 5.7 (shown in gold). Alignment was carried out on a single monomer in the dimer pair. There is a 5° shift in the angle between the wing domains of Ldcl at pH 5.7 and pH 7.0/8.5. **E)** Comparison of the central decamer pore diameter between Ldcl at pH 8.5, 7.0 and 5.7, showing a 5 Å decrease in the pore size upon stack formation at low pH.

Figure 5.

Figure 5. Structural insights into molecular determinants of the Ldcl polymerisation under acid stress conditions. **A)** Atomic model of a two-decamer Ldcl stack at pH 5.7. Dimers are coloured as shown in the cryo-EM map in Figure 4. Dotted boxes on the two-decamer stack indicate the locations of the zooms shown in panels B-E. **B)** Close-up of the first decamer-decamer interface, which includes the key stack-forming residue R468. **C)** Close-up of the second decamer-decamer interface. **D)** Overlay of the Ldcl decamer structures at pH 8.5 and 7.0 with the Ldcl stack structure at pH 5.7, focussed on R97. R97 in the Ldcl stack (green) adopts a different conformation compared to the one in the pH 8.5 crystal structure (purple, with ppGpp bound) and the pH 7.0 cryo-EM map (cyan, without ppGpp bound). Despite the absence of ppGpp in the pH 7.0 sample, R97 is still oriented towards the ppGpp binding site. **E)** Comparison between the H694-D192' distance in the Ldcl stack at pH 5.7 (coloured gold and blue), the Ldcl decamer at pH 8.5 (left, coloured purple) and pH 7.0 (right, coloured cyan). Key residues are labelled for all panels, and interactions are shown with dotted lines.

Figure 6.

Figure 6. Mutational analysis of the predicted molecular determinants of the Ldcl polymerisation under acid stress conditions Cropped negative stain EM micrographs of wild-type and mutant Ldcl at pH 5.7, scale bar = 100 nm. Ldcl_{WT} polymerises at pH 5.7, as does the double mutant Ldcl_{E445A/D447A}. In contrast, the single mutation R468E abolishes stack formation completely. Both Ldcl_{R97E} and Ldcl_{H694A} are able to polymerise, however the stacks tend to be shorter than for Ldcl_{WT}. Micrographs for Ldcl_{R97E/R468E}, Ldcl_{E445A/D447A/R468E}, and Ldcl_{H694N} were also collected but are not shown here. Ldcl_{E445A/D447A/R468E} and Ldcl_{R97E/R468E} behaved like Ldcl_{R468E} and remained entirely decameric at low pH, while Ldcl_{H694N} displayed similar behaviour to Ldcl_{H694A}.

Supplementary Figure 1.

Supplementary Figure 1. Characterisation of **Dendra2_{T69A}-LdcI** and **mGeosM-LdcI** fusion constructs by negative stain EM and BLI. **A**) Ns-EM micrographs of N-terminal FP-LdcI fusions Dendra2_{T69A}-LdcI (left) and mGeosM-LdcI (right), with 2D class averages inset (scale bar = 100 Å). Dendra2_{T69A}-LdcI only forms dimers, compared to mGeosM-LdcI which forms large non-native dodecamers. **B**) Ns-EM micrographs of C-terminal LdcI-FP fusions LdcI-Dendra2_{T69A} (left) and LdcI-mGeosM (right), with 2D class averages inset (scale bar = 100 Å). Both fusions form decamers, with flexible-linked fluorophores visible as weak densities around the outside of the decameric ring in the 2D class averages. **C**) BLI binding curves of wild-type LdcI (left, [LdcI_{WT}] = 500, 250, 125, 62.5, and 31.25 nM) and LdcI_{WT} (right, 500 nM), LdcI-Dendra2_{T69A}-LdcI (right, 500 nM) and LdcI-mGeosM-LdcI (right, 500 nM) against RavA-AVI. In contrast to wild-type LdcI, which binds RavA-AVI with a 46.3 nM affinity, there is no measurable interaction between either of the C-terminal LdcI-FP fusions and RavA-AVI.

Supplementary Figure 2.

Supplementary Figure 2. Ldcl interacts with anti-Ldcl-Nb and RavA at distinct sites. A) Ns-EM micrograph of Ldcl + anti-Ldcl-Nb (left), with 2D class averages (right) of Ldcl alone and Ldcl + anti-Ldcl-Nb (scale bar = 100 Å). **B)** *In silico* model of anti-Ldcl-Nb (represented by the anti-lysozyme nanobody crystal structure, PDB ID: 1MEL) superimposed over the cryo-EM map of the Ldcl-RavA complex (Jessop et al., 2020) (EMDB ID: 4469, PDB ID: 6Q7L). The nanobody (coloured magenta) would occupy a distinct location to the RavA binding site in the Ldcl-RavA cage-like complex (coloured gold and blue).

Supplementary Figure 3.

Supplementary Figure 3. Immunolabelling of endogenous LdcI in *E. coli* upon acid stress.

Characterisation of LdcI expression using anti-LdcI-Nb. **A**) Left – growth curve of wild-type (red) and LdcI knock-out (grey) cells after exposure to acid stress (pH 4.6), showing compromised growth in LdcI KO cells. Right – measured pH of the growth medium for wild-type and LdcI KO cells after exposure to acid stress (pH 4.6). Wild-type cells are able to buffer the medium under acid stress, whereas LdcI knock-out cells are not. Each data point represents the average of 3 independent measurements taken from 2 biological replicates, error bars correspond to the calculated standard deviation. **B**) Left – western blot showing LdcI expression after exposure to acid stress after 0, 1, 3 and 6 hours, and overnight (ON). LdcI expression is maximal soon after exposure to acid stress, tailing off after three hours. Right – western blot against LdcI in $\Delta ldcI$ and WT MG1655 cells, 0 minutes and 90 minutes after exposure to acid stress, showing a lack of LdcI expression in $\Delta ldcI$ cells and a lack of baseline expression before exposure to acid stress in WT cells. M = molecular weight marker. **C**) Flow cytometry measurements carried out after exposure of wild-type cells to acid stress show that LdcI is maximally expressed one to two hours after exposure to acid stress, before tailing off. **D**) Flow cytometry measurements of wild-type and LdcI knock-out cells labelled with anti-LdcI-Nb-AF488 after 90 minutes of growth under three conditions – pH 4.6 in the absence of oxygen and lysine, neutral pH in the absence of oxygen and lysine, and

neutral pH in the presence of oxygen but absence of lysine. No fluorescence is seen above baseline for the Ldcl knock-out cells under any condition. For wild-type cells, fluorescence is only seen under acid stress conditions. **E)** Wide-field fluorescence microscopy of Ldcl knock-out MG1655 cells (left) and wild-type MG1655 cells (right), 90 minutes after exposure to acid stress. The staining of anti-Ldcl-Nb-AF488 is specific, shown by the lack of background in the knock-out strain. Scale bar = 2µm.

Supplementary Figure 4.

Supplementary Figure 4. Galleries of 3D-STORM images of the four individual *E. coli* cells shown in Figure 3. Different orientations of each cell related by 90° rotations around the long axis of the bacterium are shown in order to facilitate visualisation of the patchy distribution of Ldcl.

Supplementary Figure 5.

Supplementary Figure 5. Cryo-EM processing pipelines. A) Cryo-EM processing pipeline for Ldcl decamers at pH 7.0. Unless otherwise indicated, processing steps were carried out in CryoSPARC. **B)** Cryo-EM processing pipeline for Ldcl stacks at pH 5.7. Software packages used at each step are indicated. For a summary of data collection and processing parameters, and refinement and validation statistics, see Supplementary Table 1.

Supplementary Figure 6.

Supplementary Figure 6. Assessment of the resolution of the cryo-EM maps. 3D reconstructions of **A**) the Ldcl decamer at pH 7.0 **B**) the Ldcl stack at pH 5.7, coloured according to the local resolution. Gold-standard FSC curves with the estimated resolution at FSC = 0.143 are shown on the right.

Supplementary Figure 7.

Supplementary Figure 8. Mutational analysis of the predicted molecular determinants of LdcI polymerisation under acid stress conditions. Full uncropped ns-EM micrographs of wild type and mutant LdcI at pH 5.7, scale bars = 100 nm. Three representative micrographs are shown for each construct. Micrographs for LdcI_{R97E/R468E}, LdcI_{E445A/D447A/R468E}, and LdcI_{H694N} were also collected but are not shown here. LdcI_{E445A/D447A/R468E} and LdcI_{R97E/R468E} behaved like LdcI_{R468E} and remained entirely decameric at low pH, while LdcI_{H694N} displayed similar behaviour to LdcI_{H694A}.

Supplementary Table 1. Cryo-EM data collection, refinement and validation statistics

	Ldcl stacks (pH 5.7)	Ldcl decamers (pH 7.0)
	EMDB 10850 PDB 6YN6	EMDB 10849 PDB 6YN5
Data collection and processing		
Magnification	130,000	116,086
Voltage (kV)	300	200
Electron exposure (e-/Å ²)	29.3	45.0
Defocus range (µm)	1.0 – 2.5	1.0 – 3.0
Pixel size (Å)	1.052	1.206
Symmetry imposed	D5	D5
Initial particle images (no.)	15,165	~796,000
Final particle images (no.)	15,157	~229,000
Map resolution (Å)	3.3	2.8
FSC threshold	0.143	0.143
Map resolution range (Å)	69.0-3.3	51.2-2.8
Refinement		
Initial model used (PDB code)	3N75	3N75
Model resolution (Å) (masked)	3.7	2.9
FSC threshold	0.5	0.5
Model resolution range (Å)	∞-3.7	∞-2.9
Map sharpening B factor (Å ²)	-96 Å ⁻²	-173 Å ⁻²
Model composition		
Non-hydrogen atoms	113,760	57,074
Protein residues	14,220	7,110
Ligands	0	0
B factors (Å ⁻²)		
Protein	91.22	36.51
Ligand	-	-
R.m.s. deviations		
Bond lengths (Å)	0.009	0.012
Bond angles (°)	1.131	1.268
Validation		
MolProbity score	1.59	1.28
Clashscore	4.78	2.83
Poor rotamers (%)	0.49	0.19
Ramachandran plot		
Favored (%)	95.04	96.74
Allowed (%)	4.96	3.26
Outliers (%)	0.00	0.00

Supplementary Table 2. Root-mean-square-deviation (RMSD) between extracted Ldcl monomer structures at pH 8.5 (PDB ID: 3N75), 7.0 and 5.7.

	Ldcl decamer, pH 8.5	Ldcl decamer, pH 7.0	Ldcl decamer, pH 5.7
Ldcl decamer, pH 8.5	0	0.24 Å (670 aligned atoms)	0.61 Å (610 aligned atoms)
Ldcl decamer, pH 7.0	0.24 Å (670 aligned atoms)	0	0.56 Å (604 aligned atoms)
Ldcl decamer, pH 5.7	0.61 Å (610 aligned atoms)	0.56 Å (604 aligned atoms)	0

Supplementary Table 3. Root-mean-square-deviation (RMSD) between extracted Ldcl dimer structures at pH 8.5 (PDB ID: 3N75), 7.0 and 5.7.

	Ldcl decamer, pH 8.5	Ldcl decamer, pH 7.0	Ldcl decamer, pH 5.7
Ldcl decamer, pH 8.5	0	0.30 Å (1328 aligned atoms)	1.51 Å (1281 aligned atoms)
Ldcl decamer, pH 7.0	0.30 Å (1328 aligned atoms)	0	1.41 Å (1287 aligned atoms)
Ldcl decamer, pH 5.7	1.51 Å (1281 aligned atoms)	1.41 Å (1287 aligned atoms)	0

Supplementary Table 4, A

Strains or plasmids	Relevant characteristics	Source/reference
<u>E. coli Strains</u>		
<u>Genotype and Use</u>		
BL21(DE3)	F ⁻ <i>ompT hsdSB</i> (r _B ⁻ , m _B ⁻) <i>gal dcm</i> (DE3): transformation and high level protein expression using a T7 RNA polymerase-IPTG induction system	Novagen product reference 69450
WK6	Δ(lac-proAB) <i>galE strA nal</i> F' [<i>lacI</i> ^q ΔM15, <i>proAB</i>]: expression of an anti-Ldcl Nanobody	Kindly provided by Dr. Aline Desmyter
TOP10	F ⁻ <i>mcrA</i> Δ(<i>mrr-hsdRMS-mcrBC</i>) φ80/ <i>lacZ</i> ΔM15 Δ <i>lacX74 recA1 araD139 Δ(ara-leu)</i> 7697 <i>galU galK</i> λ– <i>rpsL</i> (StrR) <i>endA1 nupG</i> : high-efficiency cloning and plasmid propagation	Invitrogen product reference C404003
DH5α	F ⁻ φ80/ <i>lacZ</i> ΔM15 Δ(<i>lacZYA-argF</i>)U169 <i>recA1 endA1 hsdR17</i> (r _K ⁻ , m _K ⁺) <i>phoA supE44</i> λ– <i>thi-1 gyrA96 relA1</i> : strain used for general cloning and sub-cloning applications	Invitrogen product reference 18265017
<u>Plasmids</u>		
<u>Plasmids for the construction of Dendra2_{T69A}-Ldcl_pET/TEV</u>		
Ldcl_pET/TEV	kanamycin resistant, bacterial expression vector, contains Ldcl with a TEV cleavable N-terminal His-tag	This work
Dendra2 _{T69A} _pRSETB	ampicillin resistant, high-level prokaryotic expression controlled by the strong bacteriophage T7 promoter, produces T7 lysozyme, contains Dendra2 _{T69A} with an N-terminal His-tag	J Am Chem Soc. 2016 Jan 20;138(2):558-65. doi: 10.1021/jacs.5b09923. Epub 2016 Jan 6.
Result: Dendra2 _{T69A} -Ldcl_pET/TEV	kanamycin resistant, bacterial expression vector, contains a Dendra2 _{T69A} -Ldcl fusion with an N-terminal His-tag and a TEV cleavage site between Dendra2 _{T69A} and Ldcl	This work
<u>Plasmids for the construction of mGeosM</u>		
mEos2-pRSETA	ampicillin resistant, high-level prokaryotic expression controlled by the strong bacteriophage T7 promoter, produces T7 lysozyme, contains mEos2 with an N-terminal His-tag	Nat Methods. 2009 Feb . 6(2):131-3. doi: 10.1038/nmeth.1296 A bright and photostable photoconvertible fluorescent protein. McKinney SA, Murphy CS, Hazelwood KL, Davidson MW, Looger LL. Nat Methods. 2009 Feb . 6(2):131-3. 10.1038/nmeth.1296
<u>Plasmids for the construction of mGeosM-Ldcl_pET/TEV</u>		
mGeosM-pRSETA	ampicillin resistant, high-level prokaryotic expression controlled by the strong bacteriophage T7 promoter, produces T7 lysozyme, contains mGeosM with an N-terminal His-tag	Proc Natl Acad Sci U S A. 2012 Mar 20;109(12):4455-60. doi: 10.1073/pnas.1113770109. Epub 2012 Feb 28
Ldcl_pET/TEV	kanamycin resistant, bacterial expression vector, contains Ldcl with a TEV cleavable N-terminal His-tag	This work
Result: mGeosM-Ldcl_pET/TEV	kanamycin resistant, bacterial expression vector, contains an mGeosM-Ldcl fusion with an N-terminal His-tag and a TEV cleavage site between mGeosM and Ldcl	This work

**Plasmids for the construction
of Ldcl-**

Dendra2_{T69A}_pET22b(+)

Ldcl-Dendra2_{T69A}_pET/TEV kanamycin resistant, bacterial expression vector, contains an Ldcl-Dendra2_{T69A} fusion
with a TEV cleavable N-terminal His-tag

This work

Dendra2_{T69A}_pRSETB see above

[J Am Chem Soc. 2016 Jan 20;138\(2\):558-65. doi: 10.1021/jacs.5b09923. Epub 2016 Jan 6.](#)

Ldcl_pET22b(+)

see above

[EMBO J. 2011 Mar 2;30\(5\):931-44. doi: 10.1038/emboj.2011.5. Epub 2011 Jan 28](#)

Result: Ldcl-
Dendra2_{T69A}_pET22b(+)

kanamycin resistant, bacterial expression vector, contains an Ldcl-Dendra2_{T69A} fusion
with a non-cleavable C-terminal His-tag

This work

**Plasmids for the construction
of Ldcl-mGeosM_pET22b(+)**

Ldcl-Dendra2_{T69A}_pET/TEV see above

This work

mGeosM-Ldcl_pET/TEV see above

This work

Ldcl_pET22b(+)

see above

[EMBO J. 2011 Mar 2;30\(5\):931-44. doi: 10.1038/emboj.2011.5. Epub 2011 Jan 28](#)

Result: Ldcl-
mGeosM_pET22b(+)

kanamycin resistant, bacterial expression vector, contains an Ldcl-mGeosM fusion
with a non-cleavable C-terminal His-tag

This work

Supplementary Table 4, B

Primers	Sequences (5'-3')
<i>Dendra2_{T69A}-Ldcl</i>	
V_Ldcl_Dendra2_Fw	TTAATTCCTGGGTGTTTCATGTGATGATGATGATGATGGCTGCT
V_Ldcl_Dendra2_Rv	CCCTGCCAGCCAGGTGTGGAGCAGCGGCGAAAAACC
I_Dendra2_Ldcl_Fw	GCCATCATCATCATCATCACATGAACACCCCGGAATTAACC
I_Dendra2_Ldcl_Rv	TACAGGTTTTCGCCGCTGCTCCACACCTGGCTGGGC
<i>mEos2 -> mGeosM</i>	
mEos2-H62M_Fw	ATGTACGGCAACAGGGTATTCGCC
mEos2-H62M_Rv	GAATGCAGTGGTCAGGATATCAAAGG
mEos2-H62M_2_Fw	CCTGACCACTGCATTATGTACGGCAACAGGGTATTCG
mEos2-H62M_2_Rv	CGAATACCTGTTGCCGTACATGAATGCAGTGGTCAGG
<i>mGeosM-Ldcl</i>	
V_Ldcl_mGeos_Fw	TCTGGCTTAATCGCACTCATGTGATGATGATGATGATGGCTGCT
V_Ldcl_mGeos_Rv	TGCCTGACAATGCCAGACGAAGCAGCGGCGAAAAACC
I_mGeos_Ldcl_Fw	GCCATCATCATCATCATCACATGAGTGCATTAAAGCCAGACA
I_mGeos_Ldcl_Rv	TACAGGTTTTCGCCGCTGCTTCGTCTGGCATTGTCAGGCAAT
<i>Ldcl-Dendra2_{T69A} & Ldcl-mGeosM</i>	
CL_Primer_1	AGCCAGGTGTGGTAAGATCCGGCTGCTAACAAAGC
CL_Primer_2	TGAACCAACCGCTGAGCCTTTTTTGCTTTCTTTCAATACCTTAACGG
CL_Primer_3	GGCTCAGGCGGTGGTTCAGATCTGTACGACGATGACGATAAG
CL_Primer_4	TTACCACACCTGGCTGGG
CL_Primer_7	GGTGGTTCGGGTTTCAGGTAACACCCCGGAATTAACCTG
CL_Primer_8	TGAACCCGATGATCCCGACACACCTGGCTGGGCAG
CL_Primer_25	GCAAAATGAACGAGAACCTGC
CL_Primer_26	acctgaaccgaaccaccATCTGAACCAACCGCTGAG
CL_Primer_27	tcgggatcatcggttcaAATTCGAGCTCCGTCGACAAG
CL_Primer_28	GCAGGTTCTCGTTCAATTTGC
CL_Primer_19	gtgagcaaaggcgaagaagatATGAAGATCAAACCTCCGTATGGAAGGC
CL_Primer_20	tttatacagttcatcatgccTCCAGAATGAGCAACAGCATGC
CL_Primer_21	atcttcttcgctttgctcacgtgatgatgatgatggctgctg
CL_Primer_22	ggcatggatgaactgtataaaagcagcggcGAAAACCTGTATTTTC
CL_Primer_29	GGTGGTTCGGGTTTCAGGTgtgagcaaaggcgaagaagatATG
CL_Primer_30	TGAACCCGATGATCCCGAttatacagttcatcatgccTCC

Supplementary Table 4, C

Construction of Dendra2_{T69A}Ldcl_pET/TEV

1. PCR using template Ldcl_pET/TEV with primers V_Ldcl_Dendra2_Fw and V_Ldcl_Dendra2_Rv to obtain amplification of the Ldcl containing plasmide
2. PCR using template Dendra2_{T69A}-pRSETB with primers I_Dendra2_Ldcl_Fw and I_Dendra2_Ldcl_Rv to obtain amplification of the Dendra2_{T69A} gene
3. Gibson assembly using the two obtained fragments

Construction of mGeosM-Ldcl_pET/TEV

1. Mutagenesis of mEos2 to obtain the mGeosM variant
 1. A. PCR on template mEos2-pRSETA using primers mEos2-H62M_Fw and mEos2-H62M_Rv
 1. B. Second PCR on the clone obtained in 1.A. with primers mEos2-H62M_2_Fw et mEos2-H62M_2_Rv to obtain mGeosM-pRSETA
2. Construction of an mGeosM-Ldcl fusion
 - 2.A. PCR using template Ldcl_pET/TEV with primers V_Ldcl_mGeos_Fw and V_Ldcl_mGeos_Rv to obtain amplification of the Ldcl containing plasmid
 - 2.B. PCR using template mGeosM-pRSETA with primers I_mGeos_Ldcl_Fw and I_mGeos_Ldcl_Rv to obtain amplification of the mGeosM gene
 - 2.C. Gibson assembly using the two obtained fragments

Construction of Ldcl-Dendra2_{T69A}_pET22b(+)

1. Construction of Ldcl-Dendra2_{T69A}_pET/TEV
 1. A. PCR using template Ldcl_pET/TEV with primers CL_Primer_1 and CL_Primer_2 to obtain amplification of the Ldcl containing plasmid
 1. B. PCR using template Dendra2_{T69A}-pRSETB with primers CL_Primer_3 and CL_Primer_4 to obtain amplification of the Dendra2_{T69A} gene
 1. C. Gibson assembly using the two obtained fragments
2. Construction of Ldcl-Dendra2_{T69A}_pET22b(+)
 2. A. PCR using template Ldcl-Dendra2_{T69A}_pET/TEV with primers CL_Primer_25 and CL_Primer_26 to obtain amplification of the Ldcl gene
 2. B. PCR using template Dendra2_{T69A}-pRSETB with primers CL_Primer_7 and CL_Primer_8 to obtain amplification of the Dendra2_{T69A} gene
 2. C. PCR using template Ldcl_pET22b(+) with primers CL_Primer_27 and CL_Primer_28 to obtain amplification of the pET22b(+) plasmid
 2. D. Gibson assembly using the three obtained fragments

Construction of Ldcl-mGeosM_pET22b(+)

1. Construction of mGeosM-Ldcl_pET/TEV
 1. A. PCR using template mGeosM-Ldcl_pET/TEV with primers CL_Primer_21 and CL_Primer_22 to obtain amplification of the Ldcl containing plasmid
 1. B. PCR using template mGeosM-pRSETA with primers CL_Primer_19 and CL_Primer_20 to obtain amplification of the mGeosM gene
 1. C. Gibson assembly using the two obtained fragments
2. Construction of Ldcl-mGeosM_pET22b(+)
 2. A. PCR using template Ldcl-Dendra2_{T69A}_pET/TEV with primers CL_Primer_25 and CL_Primer_26 to obtain amplification of the Ldcl gene
 2. B. PCR using template mGeosM-Ldcl_pET/TEV with primers CL_Primer_29 and CL_Primer_30 to obtain amplification of the mGeosM gene
 2. C. PCR using template Ldcl_pET22b(+) with primers CL_Primer_27 and CL_Primer_28 to obtain amplification of the pET22b(+) plasmid
 2. D. Gibson assembly using the three obtained fragments