A stomatal model of anatomical tradeoffs between gas exchange and pathogen colonization

Christopher D. Muir^{1*}

¹ School of Life Sciences, University of Hawaii, Honolulu, Hawaii, USA

Correspondence*:
Christopher D. Muir
cdmuir@hawaii.edu

2 ABSTRACT

3 Stomatal pores control both leaf gas exchange and are one route for infection of internal plant
4 tissues by many foliar pathogens, setting up the potential for tradeoffs between photosynthesis
5 and defense. Anatomical shifts to lower stomatal density and/or size may also limit pathogen
6 colonization, but such developmental changes could permanently reduce the gas exchange
7 capacity for the life of the leaf. I developed and analyzed a spatially explicit model of pathogen
8 colonization on the leaf as a function of stomatal size and density, anatomical traits which partially
9 determine maximum rates of gas exchange. The model predicts greater stomatal size or density
10 increases the probability of colonization, but the effect is most pronounced when the fraction of
11 leaf surface covered by stomata is low. I also derived scaling relationships between stomatal size
12 and density that preserves a given probability of colonization. These scaling relationships set up
13 a potential anatomical conflict between limiting pathogen colonization and minimizing the fraction
14 of leaf surface covered by stomata. Although a connection between gas exchange and pathogen
15 defense has been suggested empirically, this is the first mathematical model connecting gas
16 exchange and pathogen defense via stomatal anatomy. A limitation of the model is that it does not
17 include variation in innate immunity and stomatal closure in response to pathogens. Nevertheless,
18 the model makes predictions that can be tested with experiments and may explain variation in
19 stomatal anatomy among plants. The model is generalizable to many types of pathogens, but
20 lacks significant biological realism that may be needed for precise predictions.

21 **Keywords:** anatomy, leaf gas exchange, model, pathogen, photosynthesis, scaling, stomata, tradeoff

INTRODUCTION

22 Stomata evolved to regulate gas exchange in and out of the leaf (Hetherington and Woodward, 2003; Berry
23 et al., 2010; Chater et al., 2017), but many foliar pathogens take advantage of these chinks in the leaf
24 cuticular armor to infect prospective hosts (Zeng et al., 2010; McLachlan et al., 2014; Melotto et al., 2017).
25 The stomatal and mesophyll conductance to CO₂ are two major limits to photosynthesis (Lawson et al.,
26 2018; Flexas et al., 2018) that are partially determined by stomatal anatomy. Since CO₂ conductance
27 limits photosynthesis (Farquhar and Sharkey, 1982; Jones, 1985) and pathogen infection can reduce fitness
28 (Gilbert, 2002), this sets up a potential tradeoff between increased photosynthesis and defense against
29 pathogens mediated by stomatal anatomy (McKown et al., 2014; Tateda et al., 2019; Dutton et al., 2019;
30 Fetter et al., 2019). For example, plants could increase photosynthetic rate by developing more stomata,
31 but more stomata could result in more pathogen colonization. The optimal stomatal density, size, and
32 arrangement on the leaf will depend on the fitness gains from increased gas exchange and fitness losses

imposed by foliar pathogens, both of which depend on the environment. In the next two paragraphs I will review the relationship between stomatal anatomy, gas exchange, and foliar pathogen colonization. Then I will discuss why two anatomical traits, stomatal size and density, might be crucial components of a broader tradeoff between photosynthesis and pathogen defense.

The stomatal density and maximum pore area set an anatomical upper limit on stomatal conductance (Brown and Escombe, 1900; Parlange and Waggoner, 1970; Franks and Farquhar, 2001; Franks and Beerling, 2009b; Lehmann and Or, 2015; Sack and Buckley, 2016; Harrison et al., 2019), but stomatal shape, distribution, and patterning also affect gas exchange. Smaller guard cells and dumbbell-shaped stomata of grasses can respond faster to environmental changes (Drake et al., 2013), but responsiveness is further modulated by subsidiary cell anatomy and physiology (Franks and Farquhar, 2007; Raissig et al., 2017; Gray et al., 2020). Stomatal clustering reduces gas exchange and photosynthesis because adjacent stomata interfere with one another (Dow et al., 2014b), diffusion shells overlap (Lehmann and Or, 2015), and limitations on lateral diffusion of CO₂ in the mesophyll (Lawson and Blatt, 2014 and references therein). However, sparse clusters of small stomata could allow a leaf with low rates of gas exchange to have faster stomatal response compared to a leaf with large, low-density stomata (Papanatsiou et al., 2017). Leaves with stomata on both lower and upper surfaces (amphistomatic or amphistomatous) supply more CO₂ to the mesophyll than hypostomatic (aka hypostomatous) leaves that only have stomata on the lower surface (Parkhurst, 1978; Gutschick, 1984; Parkhurst and Mott, 1990; Oguchi et al., 2018). In addition to anatomy, the pore area shrinks and expands in response to internal and external factors to regulate gas exchange dynamically (Buckley, 2019). For example, stomata typically open during the day and close at night in C₃/C₄ plants, but the opposite is true for CAM plants. Shade, high vapor pressure deficits, dry soil and other factors can cause stomata to (partially) close even in the middle of the day. Variation in how stomata respond to internal and external signals may explain as much of the variation in gas exchange across leaves as anatomy (Lawson and Blatt, 2014).

Many types of foliar pathogens, including viruses (Murray et al., 2016), bacteria (Melotto et al., 2006; Underwood et al., 2007), protists (Fawke et al., 2015), and fungi (Hoch et al., 1987; Zeng et al., 2010) use stomatal pores to gain entry into the leaf. For example, rust fungi hyphae recognize the angle at which guard cells project from the leaf surface and use it as a cue for appressorium formation (Allen et al., 1991). Oomycete pathogens can target open stomata on a leaf (Kiefer et al., 2002). Plants can limit colonization through innate immunity, called stomatal defense (recently reviewed in Melotto et al., 2017), by closing stomata after they recognize microbe-associated molecular patterns (MAMPs) on pathogen cells. Some bacterial pathogens have responded by evolving the ability to prevent stomatal closure, increasing their colonization of the leaf interior (Melotto et al., 2006). In addition to stomatal closure, anatomical changes in stomatal density and/or size might provide another layer of defense against pathogen colonization. For example, infection increases in leaves with higher stomatal density (McKown et al., 2014; Tateda et al., 2019; Dutton et al., 2019; Fetter et al., 2019). The positive effect of stomatal density on infection suggests that infection is limited by the number or size of locations for colonization, meaning that many individual pathogens must usually be unable to find stomata or other suitable locations for colonization. This is actually somewhat surprising given the ability of some pathogens to search for and sense stomata (see above).

Stomatal anatomy could be a key link between gas exchange and pathogen colonization. Although many anatomical factors and stomatal movement affect gas exchange (see above), here I focus on the density and size of stomata in a hypostomatous leaf. Stomatal size refers to both the area of guard cells when fully open, from which one can calculate the pore area for gas exchange (see Model). For simplicity, I

77 model a hypostomatous leaf, but consider the implications for amphistomatous leaves in the Discussion.
78 Stomatal size and density not only determine the theoretical maximum stomatal conductance ($g_{s,max}$), but
79 is also proportional to the operational stomatal conductance ($g_{s,op}$) in many circumstances (Franks et al.,
80 2009, 2014; Dow et al., 2014a; McElwain et al., 2016; Murray et al., 2019). $g_{s,op}$ is the actual stomatal
81 conductance of plants in the field and is almost always below $g_{s,max}$ because stomata are usually not fully
82 open. Although they are not the same, the strong empirical relationship between $g_{s,max}$ and $g_{s,op}$ means that
83 anatomical $g_{s,max}$ can be used as a proxy for $g_{s,op}$ without explicitly modeling dynamic changes in stomatal
84 aperture (see Discussion). Stomatal size and density have also been measured on many more species than
85 stomatal responsiveness, which may make it easier to test predictions.

86 After a pathogen reaches a host, it must survive on the leaf surface and colonize the interior (Beattie and
87 Lindow, 1995; Tucker and Talbot, 2001). For analytical tractability, I restrict the focus here to colonization
88 by a pathogen using a random search on a leaf without stomatal defense (i.e. a leaf that cannot recognize
89 pathogens and close stomata). Obviously, these simplifications ignore a lot of important plant-pathogen
90 interaction biology. In the Discussion, I delve further into these limitations and suggest future work to
91 overcome these limitations. In order for pathogen-mediated selection on stomatal anatomy, I assume that
92 the pathogen reduces host fitness once it colonizes (Gilbert, 2002). Susceptible hosts can lose much of
93 their biomass or die, but even resistant hosts must allocate resources to defense or reduce photosynthesis
94 because of defoliation, biotrophy, or necrosis around sites of infection (Bastiaans, 1991; Mitchell, 2003).

95 The purpose of this study is to develop a theoretical framework to test whether variation in stomatal size
96 and density arises from a tradeoff between gas exchange and pathogen colonization. Since stomatal size and
97 density affect both gas exchange and pathogen colonization, selection to balance these competing demands
98 could shape stomatal size-density scaling relationships. Botanists have long recognized that stomatal size
99 and density are inversely correlated (Weiss, 1865; Tichá, 1982; Hetherington and Woodward, 2003; Sack et
100 al., 2003; Franks and Beerling, 2009a; Brodribb et al., 2013; Boer et al., 2016), but the evolutionary origin
101 of this relationship is not yet known. Here I argue that deleterious effects of pathogen infection could shape
102 selection on this relationship. Explanations for inverse size-density scaling are usually cast in terms of
103 preserving $g_{s,max}$ and/or stomatal cover (f_S), defined as the fraction of epidermal area allocated to stomata
104 (Boer et al., 2016), because there are many combinations of stomatal size and density that have same $g_{s,max}$
105 or same f_S :

$$g_{s,max} = bmDS^{0.5} \quad (1)$$

$$f_S = DS. \quad (2)$$

106 D and S are stomatal density and size, respectively (see Table 1 for a glossary of mathematical
107 symbols and units). b and m are assumed to be biophysical and morphological constants, *sensu* Sack and
108 Buckley (2016) (see Supplementary Material). f_S is proportional to the more widely used stomatal pore area
109 index [Sack et al. (2003); see Supplementary Material]. If size and density also affect pathogen colonization,
110 then selection from foliar pathogens could significantly alter the size-density scaling relationship. The
111 empirical size-density scaling relationship is linear on a log-log scale, determined by an intercept α and
112 slope β :

$$D = e^{\alpha} S^{-\beta}; \quad (3)$$

$$d = \alpha - \beta S. \quad (4)$$

113 For brevity, $d = \log(D)$ and $s = \log(S)$. Rearranging Equations 1 and 2, a scaling relationship where
114 $\beta = 0.5$ preserves $g_{s,\max}$ while $\beta = 1$ preserves f_S .

115 How would adding pathogens alter these predicted scaling relationships? For simplicity, consider two
116 environments, one without foliar pathogens and one with lots. In the absence of foliar pathogens, we
117 expect size-density scaling to preserve $g_{s,\max}$, f_S , or some least-cost combination of them. What happens
118 when we introduce pathogens? If stomatal size and density increase pathogen colonization, then selection
119 will favor reduced size and/or density. This would change the intercept α but not the slope. The effect
120 of foliar pathogens on the slope depends on the relationship between size, density, and probability of
121 colonization. If the probability of colonization is proportional to the product of *linear* stomatal size ($S^{0.5}$)
122 and density ($\propto DS^{0.5}$ as for $g_{s,\max}$) then it has the same effect on the slope as $g_{s,\max}$ because there are many
123 combinations of D and $S^{0.5}$ that have same probability of colonization. If the probability of colonization is
124 proportional to the product of *areal* stomatal size (S) and density ($\propto DS$ as for f_S) then it has the same
125 effect on the slope as f_S because there are many combinations of D and S that have same probability
126 of colonization. Alternatively, the probability of colonization may have a different scaling relationship
127 (neither 0.5 nor 1) or may be nonlinear on a log-log scale. Unlike $g_{s,\max}$ and f_S , we do not have theory to
128 predict a stomatal size-density relationship that preserves the probability of colonization.

129 In summary, the physical relationship between stomatal size, density, and conductance is well established
130 (Harrison et al., 2019). Size and density also likely affect the probability of pathogen colonization, but
131 we do not have a theoretical model that makes quantitative predictions. The inverse stomatal size-density
132 relationship has usually been explained in terms of preserving stomatal conductance and/or stomatal cover,
133 but selection by pathogens might alter scaling. To address these gaps, the goals of this study are to 1)
134 introduce a spatially explicit model pathogen colonization on the leaf surface; 2) use the model to predict
135 the relationship between $g_{s,\max}$, f_S , and the probability of colonization; 3) work out what these relationships
136 predict about stomatal size-density scaling. I analyzed an idealized, spatially explicit Model of how a
137 pathogen lands on a leaf and finds a stomate to colonize the leaf using a random search. To my knowledge,
138 this is the first model that makes quantitative predictions about the relationship between stomatal anatomy,
139 the probability of colonization, and their impact on stomatal size-density scaling.

MODEL

140 In this section, I introduce a spatially explicit model of successful pathogen colonization on a leaf surface.
141 I explain the model structure and assumptions here; the Materials and Methods section below describes
142 how I analyzed the model to address the goals of the study. For generality, I refer to a generic “pathogen”
143 that lands on leaf and moves to a stomate. The model is agnostic to the type of pathogen (virus, bacterium,
144 fungus, etc.) and the specific biological details of how it moves. For example, motile bacterial cells can
145 land and move around (Beattie and Lindow, 1995) whereas fungi may germinate from a cyst and grow until
146 they form an appressorium for infection (Tucker and Talbot, 2001). These very different tropic movements
147 on the leaf are treated identically here. I do not model photosynthesis explicitly, but assume that stomatal
148 conductance limits carbon fixation, even though the relationship is nonlinear. I used Sympy version 1.6.1
149 (Meurer et al., 2017) for symbolic derivations.

Table 1 | Glossary of mathematical symbols. The columns indicate the mathematical Symbol used in the paper, the associated symbol used in R scripts, scientific Units, and a verbal Description.

Symbol	R	Units	Description
D	D	mm^{-2}	stomatal density
d	d	mm^{-2}	stomatal density (log-scale, $d = \log D$)
f_s	f_s	none	stomatal cover ($f_s = DS$)
$g_{s,\max}$	g_smax	$\text{mol m}^{-2} \text{s}^{-1}$	theoretical maximum stomatal conductance
$g_{s,\text{op}}$	g_sop	$\text{mol m}^{-2} \text{s}^{-1}$	operational stomatal conductance
H	H	μm^{-1}	death rate of pathogen on leaf surface
R	R	μm	stomatal radius ($S = \pi R^2$)
S	S	μm^2	stomatal size
s	s	μm^2	stomatal size (log-scale, $s = \log S$)
θ_i	theta_i	radians	angles between pathogen (x_p, y_p) and lines tangent to the circumference of stomate i
U	U	μm	interstomatal distance
v_i	v_i	μm	distance between pathogen (x_p, y_p) and stomate i
x_i, y_i	x_i, y_i	μm	position of stomate i
x_p, y_p	x_p, y_p	μm	starting position of pathogen

Spatial representation of stomata

Stomata develop relatively equal spacing to minimize resistance to lateral diffusion (Morison et al., 2005), allow space between stomata (Dow et al., 2014b), and prevent stomatal interference (Lehmann and Or, 2015). Here I assume that stomata are arrayed in an equilateral triangular grid with a density D and size (area) S on the abaxial surface only, since most leaves are hypostomatous [Muir (2015); but see Discussion]. This assumption ignores veins, trichomes, and within-leaf variation in stomatal density. Stomata are therefore arrayed in an evenly spaced grid (Figure 1a). The interstomatal distance U , measured as the distance from the center of one stomata to the next, is the maximal diagonal of the hexagon in μm that forms an equal area boundary between neighboring stomata. The area of a hexagon is $A_{\text{hexagon}} = \frac{\sqrt{3}}{2} U^2$. By definition the stomatal density is the inverse of this area, such that $D = A_{\text{hexagon}}^{-1} = \frac{2}{\sqrt{3}} U^{-2}$. Therefore, interstomatal distance can be derived from the stomatal density as:

$$D = \frac{2}{\sqrt{3}} U^{-2}$$

$$U = \left(\frac{2}{\sqrt{3}} D^{-1} \right)^{0.5}$$

For example, if the density is $D = 10^2 \text{ mm}^{-2} = 10^{-4} \mu\text{m}^{-2}$, then U is $107.5 \mu\text{m}$. Parkhurst (1994) described this result previously. I also make the simplifying assumption that stomata are perfectly circular with radius R when fully open. This may be approximately true for fully open stomata with kidney-shaped guard cells (Sack and Buckley, 2016 and references therein). Although I assume stomata are circular here, in calculating $g_{s,\max}$, I assume typical allometric relationships between length, width, and pore area [Sack and Buckley (2016); see Supplementary Material].

Figure 1. A spatially explicit model of stomatal anatomy and pathogen colonization. **a.** Stomata are assumed to be in a homogenous equilateral triangular grid, which means that we can extrapolate from **b.** a focal triangle to the entire leaf. The circles represent idealized stomata; the grey lines between them are for visualization. **c.** By symmetry, a single focal region within the focal triangle can be modeled and extrapolated to the rest of the triangle. **d.** The model assumes that a pathogen, depicted as a grey rod, lands somewhere on the leaf surface and will successfully locate a stomate if it moves at the correct angle, depicted by the grey polygons.

170 Spatial representation of pathogen search

171 Since stomata are arrayed in a homogeneous grid, we can focus on single focal triangle (Figure 1b-c).
 172 Suppose that an individual pathogen (e.g. bacterial cell or fungal spore) lands at a uniform random position
 173 within the focal triangle and must arrive at a stomate to colonize. If it lands on a stomate, then it infects the
 174 leaf with probability 1; if it lands between stomata, then it infects the leaf with probability p_{locate} . This

175 is the probability that it locates a stomate, which I will derive below. The probabilities of landing on or
176 between a stomate are f_S and $1 - f_S$, respectively. Hence, the total probability of colonization is:

$$p_{\text{colonize}} = f_S + (1 - f_S)p_{\text{locate}}. \quad (5)$$

177 I assume that the pathogen cannot sense where stomata are and orients at random, thereafter traveling in
178 that direction. If it successfully locates a stomate, it colonizes the leaf, but otherwise does not infect. If
179 there is a high density of stomata and/or large stomata, the probability of locating a stomate increases. By
180 assuming that stomata form an equilateral triangular grid (see above), we can extrapolate what happens
181 in the focal triangle (Figure 1b) by symmetry. Further, since an equilateral triangle can be broken up into
182 six identical units (Figure 1c), we can simply calculate p_{locate} in this focal area. This implicitly assumes
183 that the probability of colonizing stomata outside the focal area is 0 because they are too far away. This
184 assumption may be unrealistic for larger pathogens, such as fungi, whose hyphae can travel longer distances
185 on the leaf surface (Brand and Gow, 2012). In Appendix 1: Spatially implicit model I derive a simpler, but
186 spatially *implicit* model that relaxes the assumption the pathogens must colonize a stomate within their
187 focal triangle.

188 Consider a pathogen that lands in position (x_p, y_p) within the triangle. The centroid of the triangle is
189 at position (x_c, y_c) and a reference stomate is at position $(0, 0)$ (Figure 2a). Therefore $x_c = U/2$ and
190 $y_c = \sqrt{3}U/6$. The other stomata are at positions $(U/2, \sqrt{3}U/2)$ and $(U, 0)$ (Figure 2). x_p and y_p are
191 defined as the horizontal and vertical distances, respectively, from the pathogen to the reference stomate at
192 position $(0, 0)$.

193 Given that the pathogen starts at position (x_p, y_p) , what's the probability of contacting one of the stomata
194 at the vertices of the focal triangle? I assume the probability of contacting a stomate is equal to the
195 proportion of angular directions that lead to a stomate (Figure 1d). I solved this by finding the angles
196 $(\theta_1, \theta_2, \theta_3)$ between lines that are tangent to the outside of the three stomata and pass through (x_p, y_p)
197 (Figure 2a). If stomate i is centered at (x_i, y_i) , the two slopes of tangency as function of pathogen position
198 are:

$$t_{i,1}(x_p, y_p) = \frac{-Re_{i,2}(x_p, y_p) + e_{i,3}(x_p, y_p)}{e_{i,1}(x_p, y_p)} \quad (6)$$

$$t_{i,2}(x_p, y_p) = \frac{Re_{i,2}(x_p, y_p) + e_{i,3}(x_p, y_p)}{e_{i,1}(x_p, y_p)} \quad (7)$$

199 where

$$e_{i,1}(x_p, y_p) = (R^2 - x_i^2 + 2x_i x_p - x_p^2), \quad (8)$$

$$e_{i,2}(x_p, y_p) = \sqrt{-e_{i,1} + (y_i - y_p)^2}, \quad (9)$$

$$e_{i,3}(x_p, y_p) = -x_i y_i + x_i y_p + x_p y_i - x_p y_p. \quad (10)$$

Figure 2. Spatial representation of stomata and pathogen. **a.** The pathogen starts at a uniform random position within the focal region denoted (x_p, y_p) . Within the focal triangle, the reference stomate is at position $(0,0)$ by definition, and other stomatal positions are determined by the interstomatal distance U . **b.** Within the focal region, a pathogen can land within the stomate (white circle with grey outline and radius R) or in the grey area. The outer borders of this area are shown and depend on R and U . For a given position x , there is a minimum y -value ($y_{\min}$, dashed line) and maximum y -value ($y_{\max}$, solid line).

200 Note that $i \in \{1, 2, 3\}$, indexing the three stomata in the focal triangle. The angle in radians between
 201 $t_{i,1}(x_p, y_p)$ and $t_{i,2}(x_p, y_p)$ is:

$$\theta_i(x_p, y_p) = \arctan\left(\frac{t_{i,1}(x_p, y_p) - t_{i,2}(x_p, y_p)}{1 + (t_{i,1}(x_p, y_p)t_{i,2}(x_p, y_p))}\right) \quad (11)$$

I further assumed that the longer distance a pathogen must travel, the less likely it would be to locate a stomate. For example, if stomata are at very low density, then a pathogen may die before it reaches a stomate because of UV, desiccation, or another factor. I included this effect by assuming the probability of reaching a stomate declines exponentially at rate H with the Euclidean distance $v_i(x_p, y_p)$ between the pathogen location and the edge of stomata i , which is distance R from its center at x_i, y_i :

$$v_i(x_p, y_p) = \sqrt{(x_i - x_p)^2 + (y_i - y_p)^2} - R. \quad (12)$$

The probability of locating a stomate as a function of pathogen position (x_p and y_p) is the sum of the angles divided by 2π , discounted by their distance from the stomate:

$$f_{\text{locate}}(x_p, y_p) = \frac{1}{2\pi} \sum_{i=1}^3 e^{-Hv_i(x_p, y_p)} \theta_i(x_p, y_p) \quad (13)$$

When there is no pathogen death ($H = 0$), p_{locate} is the fraction of angles that lead from (x_p, y_p) to a stomate. When $H > 0$, p_{locate} is proportional to this fraction, but less than it depending on stomatal density, size, and starting location of the pathogen.

To obtain the average p_{locate} , we must integrate $f_{\text{locate}}(x_p, y_p)$ over all possible starting positions (x_p, y_p) within the focal area. The focal area is a 30-60-90 triangle with vertices at the center of the reference stomate $(0, 0)$, the midpoint of baseline $(U/2, 0)$, and the centroid of the focal triangle $(U/2, \sqrt{3}/6U)$ (Figure 1c). Colonization occurs with probability 1 if the pathogen lands in the reference stomate, so we need to integrate the probability of colonization if it lands elsewhere. This region extends from the edge of the stomate, at $\sqrt{3}/2R$ to $U/2$ (Figure 2b). At any x , we integrate from the bottom of the focal area ($y_{\min}$) to the top ($y_{\max}$):

$$y_{\min} = f(x) = \begin{cases} \sqrt{R^2 - x^2}, & \text{if } \frac{\sqrt{3}}{2}R < x < R \\ 0, & \text{if } R \leq x \leq \frac{U}{2} \end{cases} \quad (14)$$

$$y_{\max} = f(x) = \frac{\sqrt{3}}{3}x \quad (15)$$

The integral is:

$$p_{\text{locate}} = \frac{1}{a_{\text{focal}}} \int_{\frac{\sqrt{3}}{2}R}^{U/2} \int_{y_{\min}}^{y_{\max}} f_{\text{locate}}(x, y) dx dy \quad (16)$$

a_{focal} is the area of the focal region depicted in grey in Figure 2b:

$$a_{\text{focal}} = \frac{U^2}{8\sqrt{3}} - \frac{\pi R^2}{12}$$

MATERIALS AND METHODS

The Model calculates a probability of host colonization (Equation 5) as a function of stomatal density, size, and position of a pathogen on the leaf. I solved p_{colonize} by importing symbolic derivations from Sympy into R with **reticulate** version 1.16 (Ushey et al., 2020) and used the `integral2()` function in the **pracma** package version 2.2.9 (Borchers, 2019) for numerical integration. I used R version 4.0.2 (R Core Team, 2020) for all analyses and wrote the paper in **rmarkdown** version 2.3 (Xie et al., 2018; Allaire et al., 2020). Citations for additional R software packages are in Appendix 2. Source code is deposited on GitHub (<https://github.com/cdmuir/stomata-tradeoff>) and will be archived on Zenodo upon publication.

What is the relationship between stomatal size, density, and colonization?

I calculated p_{colonize} over a biologically plausible grid of stomatal size and density for hypostomatous species based on Boer et al. (2016). Stomatal density (D) ranges from $10^1 - 10^{3.5} \text{ mm}^{-2}$; stomatal size (S) ranges from $10^1 - 10^{3.5} \mu\text{m}^2$. I only considered combinations of size and density where stomatal cover (f_S) was less than $1/3$, which is close to the upper limit in terrestrial plants (Boer et al., 2016). I crossed stomatal traits with three levels of $H \in \{0, 0.01, 0.1\}$. When $H = 0$, a pathogen persists indefinitely on the leaf surface. $H = 0.01$ and $H = 0.1$ correspond to low and high death rates, respectively. These values are not necessarily realistic, but illustrate qualitatively how a hostile environment on the leaf surface alters model predictions.

How do pathogens alter optimal stomatal size-density scaling?

The stomatal size-density scaling relationship can be explained in terms of preserving a constant stomatal conductance ($g_{s,\text{max}}$) that is proportional to $DS^{0.5}$ when bm is constant (Equation 1). In other words, there are infinitely many combinations of D and $S^{0.5}$ with the same $g_{s,\text{max}}$. If $g_{s,\text{max}}$ is held constant at C_g , then the resulting size-density scaling relationship on a log-log scale is:

$$d = c_g - 0.5s$$

where lowercase variables are log-transformed equivalents of their uppercase counterparts (Table 1). The scaling exponent $\beta_g = 0.5$ preserves C_g .

Next, suppose there is a scaling exponent β_p that preserves p_{colonize} for the product DS^{β_p} . If $\beta_p = 0.5$, then p_{colonize} is always proportional to $g_{s,\text{max}}$. If $\beta_p > 0.5$, small, densely packed stomata would be more resistant to colonization (lower p_{colonize}) compared to larger, sparsely spaced stomata with the same $g_{s,\text{max}}$. If $\beta_p < 0.5$, small, densely packed stomata would be less defended (higher p_{colonize}) compared to larger, sparsely spaced stomata with the same $g_{s,\text{max}}$. I refer to the three outcomes ($\beta_p = 0.5$, $\beta_p < 0.5$, and $\beta_p > 0.5$) as iso-, hypo-, and hyper-conductance, respectively. I was unable to solve analytically for β_p , so I numerically calculated isoclines of p_{colonize} over the grid of D and S values described in the preceding subsection. I numerically calculated the scaling relationships at a constant $p_{\text{colonize}} \in \{0.025, 0.1, 0.4\}$ for $H \in \{0, 0.01, 0.1\}$.

RESULTS

Nonlinear relationships between colonization, stomatal cover, and conductance

The probability of colonization (p_{colonize}) is not simply proportional to stomatal cover (f_S). At low f_S , p_{colonize} increases rapidly relative to f_S at first (Figure 3a). At higher f_S , p_{colonize} increases linearly with f_S . When pathogens persist indefinitely ($H = 0$), any combination of stomatal size (S) and density (D) with the same f_S have the same effect on p_{colonize} . When $H > 0$, pathogens are less likely to land close enough

to a stomate to infect before dying, so p_{colonize} is closer to f_S (Figure 3a). The maximum p_{colonize} under the range of parameters considered was ~ 0.6 when $H = 0$ and f_S is at its maximum value of $1/3$. When f_S is low, p_{colonize} is also low. The relationship between p_{colonize} , f_S , and $g_{s,\text{max}}$ is qualitatively similar in the spatially implicit model, but the values for p_{colonize} are substantially higher because pathogens can potentially colonize any stomate on the leaf rather than only those in the focal triangle (see Appendix 1: Spatially implicit model for more detail). Bear in mind that this is the probability for a single individual searching randomly; if enough individuals reach the leaf and/or they can actively find stomata, it's almost certain that at least some will colonize the leaf. However, reducing p_{colonize} may help plants limit the damage since fewer total individual pathogens will colonize the leaf interior.

p_{colonize} is not directly proportional to f_S because it depends on D and S in quantitatively different ways (Figure S1). For the same f_S , leaves with greater D have higher p_{colonize} (Figure 3a). Holding f_S constant, leaves with lower D and higher S will have a greater distance (v_i) between a pathogen and its stomata. When $H > 0$, this extra distance leads more pathogens to die before they can find a stomate. However, this result is inconsistent with the spatially implicit model (Appendix 1) because S and D have identical effects on f_S .

In contrast to f_S , p_{colonize} increases at a greater than linear rate with stomatal conductance ($g_{s,\text{max}}$). Greater D (smaller S) is associated with lower p_{colonize} for a given value of $g_{s,\text{max}}$ (Figure 3b). This happens because p_{colonize} increases approximately linearly with S whereas $g_{s,\text{max}}$ is proportional to $S^{0.5}$. Therefore, p_{colonize} increases exponentially with $g_{s,\text{max}}$ at all stomatal densities, but the rate of growth is lower at greater D for a given value of $g_{s,\text{max}}$.

Hyper-conductance size-density scaling

The scaling relationship between S and D that preserves p_{colonize} is always greater than 0.5 (hyper-conductance), but usually less than 1. When $H = 0$, the scaling relationship is essentially 1 (Figure 4), which means that an increase f_S leads to a proportional increase in p_{colonize} . Because the scaling relationship is greater than 0.5, leaves with greater stomatal density will have lower p_{colonize} than leaves lower stomatal density but the same $g_{s,\text{max}}$. In other words, increasing D and lowering S allows plants to reduce p_{colonize} while maintaining $g_{s,\text{max}}$. The scaling relationship is slightly less than 1, but still greater than 0.5, when $H > 0$ (Figure 4). In this area of parameter space, lower stomatal density can reduce f_S while p_{colonize} is constant, but this will still result in lower $g_{s,\text{max}}$. In the spatially implicit model, the size-density scaling exponent was always exactly 1 except when $H = 0$ (Appendix 1).

DISCUSSION

Stomatal density and size set the upper limit on gas exchange in leaves (Harrison et al., 2019) and is often closely related to operational stomatal conductance in nature (Murray et al., 2019). Despite the fact that many foliar pathogens infect through stomata, the relationship between stomatal anatomy and resistance to foliar pathogens is less clear than it is for gas exchange. I used a spatially explicit model of a pathogen searching for a stomate to colonize a host. From this Model, I derived predictions about the relationship between stomatal anatomy and the probability of colonization, a component of disease resistance. The model predicts that the probability of colonization is not always proportional to the surface area of leaf covered by stomata (f_S), as one might intuitively predict. If the leaf surface is a hostile environment and pathogens have a limited time to search, lower stomatal density decreases the probability of colonization even if f_S is constant. However, $g_{s,\text{max}}$ decreases proportionally more than the probability of colonization. The model highlights the potential for conflicting demands of minimizing pathogen colonization, minimizing stomatal cover, and maintaining stomatal conductance. Including the effect of

Figure 3. The probability of colonization increases with both stomatal cover and conductance I simulated the probability of colonization (p_{colonize} , y -axis) over a range of stomatal densities and sizes (see [Materials and Methods]), but a subset of results are shown here. Stomatal size and density determine stomatal cover (f_s ; Equation 2) and theoretical maximum stomatal conductance ($g_{s, \text{max}}$; Equation 1). **a.** p_{colonize} initially increases rapidly with f_s (x -axis), then slows down to a linear relationship. Overall, p_{colonize} is lower when pathogens can die on the leaf surface ($H > 0$). The relationship between f_s and p_{colonize} is the same regardless of stomatal density when $H = 0$ (upper facet), which is why the lines overlap. When $H > 0$, higher density (solid lines) increase p_{colonize} (lower facets). **b.** p_{colonize} increases exponentially with $g_{s, \text{max}}$ at all stomatal densities, but p_{colonize} is much lower at higher densities for a given $g_{s, \text{max}}$. The relationship between $g_{s, \text{max}}$ and p_{colonize} is similar for all values of H .

301 anatomy on pathogen colonization therefore has the potential to change our understanding of how stomatal
302 size-density scaling evolves in land plants.

Figure 4. Log-log scaling relationships between stomatal density (D , x-axis) and size (S , y-axis) that preserve the probability of colonization (p_{colonize}). In each panel, solid lines indicate values of D and S where p_{colonize} is 0.025 (lowest line), 0.1, or 0.4 (highest line). For reference, dashed grey lines show scaling relationships that preserve f_S ($\beta = 1$, slope $= -1/\beta = -1$) and $g_{S,max}$ ($\beta = 0.5$, slope $= -1/\beta = -2$) drawn through the centroid of the plotting region. When the death rate on the leaf surface is low ($H = 0$), the scaling exponent is very close to $\beta = 1$. When $H > 0$, $0.5 < \beta < 1$ and is slightly nonlinear on a log-log scale.

303 The model predicts that in most cases, increasing stomatal cover should lead to a proportional increase
304 in colonization, which agrees with empirical studies (e.g. McKown et al. (2014); Tateda et al. (2019);
305 Dutton et al. (2019); Fetter et al. (2019)). It also makes new, testable predictions that are less intuitive
306 (Table 2). At very low f_S , there is a rapid increase in colonization (Figure 3a). If there are no stomata, the
307 probability of colonization is 0, so the first few stomata dramatically increase the probability. This is less
308 likely to be significant for abaxial (lower) leaf surfaces, which usually have most of the stomata (Salisbury,
309 1928; Metcalfe and Chalk, 1950; Mott et al., 1984; Peat and Fitter, 1994; Jordan et al., 2014; Muir, 2015;
310 Bucher et al., 2017; Drake et al., 2019). However, many adaxial (upper) leaf surfaces have zero or very few
311 stomata. Using adaxial leaf surfaces, it should be possible to test if small changes in stomatal size or density
312 have a larger effect on pathogen colonization when f_S is low. Such experiments could use natural genetic
313 variation (McKown et al., 2014) or mutant lines (Dow et al., 2014b). The nonlinear increase in p_{colonize} is
314 less apparent when $H > 0$ (Figure 3a). A more hostile microenvironment (e.g. drier, higher UV) should
315 therefore reduce the effect of increased size or density at low f_S . If true, the diminishing marginal effect
316 of f_S on colonization could explain why stomatal ratio on the upper and lower surface is bimodal (Muir,
317 2015). The initial cost of adaxial (upper) stomata is relatively high, but if the benefits outweigh the costs,
318 then equal stomatal densities on each surface maximize CO_2 supply for photosynthesis (Parkhurst, 1978;
319 Gutschick, 1984; Parkhurst and Mott, 1990). The costs and benefits will certainly vary with environmental
320 conditions as well. Future work should extend this model, which considered hypostomatous leaves, to

address stomatal size and density in amphistomatous leaves, since leaf surfaces may differ in the type of pathogens present and microenvironment (McKown et al., 2014; Fetter et al., 2019).

Table 2 | New testable model predictions and suggested experiments to test them.

Model Prediction	How to test it
Increasing stomatal size and/or density will have a larger effect on pathogen colonization in leaves with low stomatal cover.	Compare the effect of changing stomatal size and/or density on pathogen colonization in leaves with low and high stomatal cover.
Increasing stomatal size and/or density will have a smaller effect on pathogen colonization in epiphytic environments more hostile to pathogens.	Compare the effect of changing stomatal size and/or density on pathogen colonization in more hostile environments (e.g. drier, higher light)
When selection against pathogen colonization stronger, the stomatal size-density scaling exponent should be lower	Measure stomatal anatomy in environments that differ in pathogen colonization using comparative or experimental approaches

An effect of stomatal size and density on foliar pathogen colonization could change our understanding of stomatal size-density scaling. Since allocating leaf epidermis to stomata may be costly (Franks and Farquhar, 2007; Assmann and Zeiger, 1987; Dow et al., 2014b; Lehmann and Or, 2015; Baresch et al., 2019), selection should favor leaves that achieve a desired $g_{s,max}$ while minimizing f_s (Boer et al., 2016). Because of their different scaling exponents (Equation 1, 2), smaller, densely packed stomata can achieve the same $g_{s,max}$ at minimum f_s . However, many leaves have larger, sparsely packed stomata. Incorporating pathogen colonization may explain why. If pathogens have a limited time to find stomata before dying ($H > 0$), then the scaling exponent between size and density that keeps $p_{colonize}$ constant is between 0.5 and 1, the scaling exponents for $g_{s,max}$ and f_s , respectively (Figure 4). Greater density of smaller stomata can increase $g_{s,max}$ while keeping $p_{colonize}$ constant, but this will increase f_s . Conversely, f_s could decrease while keeping $p_{colonize}$ constant, but this will decrease $g_{s,max}$. This sets up the potential for conflict between competing goals. The optimal stomatal size and density will therefore depend on the precise costs and benefits of infection, stomatal conductance, and stomatal cover. This may explain why many leaves have large, sparsely packed stomata despite the fact that they could achieve the same $g_{s,max}$ and lower f_s with smaller, more densely packed stomata.

The model examines the probability of colonization for a single pathogen. The calculated probabilities of colonization should not be interpreted as exact predictions, but rather as depicting qualitative relationships between stomatal anatomy and infection severity. The energetic cost and lost photosynthetic capacity (closed stomata, necrosis, etc.) of dealing with a pathogen is assumed to be proportional to the amount of infection. The actual fitness cost will be modulated by the number of pathogens landing on the leaf and the cost of infection, all else being equal. In environments with fewer or less virulent pathogens, the fitness cost of infection will be less than in environments with more abundant, virulent pathogens. The model is less relevant to very susceptible host plants that can be severely damaged or killed by a small number of colonizations that spread unchecked throughout the host tissue.

CONCLUSION

The model makes two non-intuitive predictions. First, the effect of increased stomatal density or size on susceptibility to foliar pathogens is greatest when stomatal cover is very low. Second, maximizing

350 disease resistance sets up a potential conflict between minimizing stomatal cover and maximizing stomatal
351 conductance. The first prediction is consistent with results in *Populus trichocarpa* (McKown et al., 2014)
352 and may be relatively straightforward to test experimentally with adaxial (upper) stomata that occur at low
353 and moderate densities within the same or closely related species (Muir et al., 2014; Fetter et al., 2019).
354 The second prediction about size-density scaling is more complex because we would need to know the
355 relationships between colonization, stomatal cover, stomatal conductance, and fitness in natural conditions.
356 There is growing evidence that stomata mediate tradeoffs between photosynthesis and defense in *Populus*
357 *trichocarpa* (McKown et al., 2019), but testing these predictions in a variety of species will help determine
358 whether pathogens have played an important role shaping stomatal anatomy in land plants.

ACKNOWLEDGMENTS

359 I would like to thank Athena McKown and an anonymous referee for valuable feedback that improved this
360 manuscript.

FUNDING

361 I am grateful startup funds from the University of Hawaii for supporting this work.

CONFLICT OF INTEREST STATEMENT

362 The author declares that the research was conducted in the absence of any commercial or financial
363 relationships that could be construed as a potential conflict of interest.

364

SUPPLEMENTARY MATERIAL

365 $g_{s,max}$ calculation

366 I calculated $g_{s,max}$ (Equation 1) to water vapor at a reference leaf temperature ($T_{leaf} = 25^\circ \text{C}$) following
367 Sack and Buckley (2016). They defined a biophysical and morphological constant as:

$$b = D_{wv}/v$$

$$m = \frac{\pi c^2}{j^{0.5}(4hj + \pi c)}$$

368 b is the diffusion coefficient of water vapor in air (D_{wv}) divided by the kinematic viscosity of dry air (v).
369 $D_{wv} = 2.49 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$ and $v = 2.24 \times 10^{-2} \text{ m}^3 \text{ mol}^{-1}$ at 25° (Monteith and Unsworth, 2013). For
370 kidney-shaped guard cells, $c = h = j = 0.5$.

371 f_s is proportional to the stomatal pore area index

372 The stomatal pore area index (SPI; Sack et al., 2003) is calculated as the product of the stomatal density
373 and guard cell length (GCL) squared:

$$\text{SPI} = D \times \text{GCL}^2$$

374 Assuming that the stomatal radius R is half the GCL, then stomatal size S is equivalent to:

$$S = \pi R^2 = \frac{\pi \times \text{GCL}^2}{4}$$

375 Based on equation 2, it follows that f_s and SPI are proportional:

$$f_s = \frac{\pi \times \text{SPI}}{4}$$

Figure S1. The probability of colonization increases with both stomatal size (S) and density (D) I simulated the probability of colonization (p_{colonize} , y -axis) over a range of S , D , and H (see [Materials and Methods]) **a.** Each line shows how p_{colonize} increases with S (x -axis, log-scale) for selected values of $D \in \{10, 100, 1000\} \text{ mm}^{-2}$. **b.** Each line shows how p_{colonize} increases with D (x -axis, log-scale) for selected values of $S \in \{10, 100, 1000\} \mu\text{m}^2$. The facets show results for different values of H

Figure S2. The probability of colonization increases with both stomatal cover and conductance in the spatially implicit model. As in Figure 3, I simulated the probability of colonization (p_{colonize} , y-axis) over a range of stomatal densities and sizes (see [Materials and Methods]), but a subset of results are shown here. Stomatal size and density determine stomatal cover (f_s ; Equation 2) and theoretical maximum stomatal conductance ($g_{s, \text{max}}$; Equation 1). **a.** p_{colonize} initially increases rapidly with f_s (x-axis), then slows down to a linear relationship. Overall, p_{colonize} is lower when pathogens can die on the leaf surface ($H > 0$). The relationship between f_s and p_{colonize} is the same regardless of stomatal density for all values of H , which is why the lines overlap. **b.** p_{colonize} increases sigmoidally with $g_{s, \text{max}}$ at all stomatal densities, but p_{colonize} is lower at higher densities for a given $g_{s, \text{max}}$. The relationship between $g_{s, \text{max}}$ and p_{colonize} is similar for all values of $H > 0$.

Figure S3. Log-log scaling relationships between stomatal density (D , x-axis) and size (S , y-axis) that preserve the probability of colonization (p_{colonize}) in the spatially implicit model. As in Figure 4, in each panel, solid lines indicate values of D and S where p_{colonize} is 0.25 (lowest line), 0.5, or 0.75 (highest line). For reference, dashed grey lines show scaling relationships that preserve f_s ($\beta = 1$, slope = $-1/\beta = -1$) and $g_{s,\text{max}}$ ($\beta = 0.5$, slope = $-1/\beta = -2$) drawn through the centroid of the plotting region. The scaling exponent is unity $\beta = 1$ when $H > 0$.

APPENDIX 1: SPATIALLY IMPLICIT MODEL

A limitation of the spatially explicit model is that a pathogen could only infect stomata in the focal triangle where it landed. Here I analyze an alternative, spatially implicit, model that relaxes this assumptions. Instead, I assume that a pathogen can potentially infect any stomate on the leaf. It searches through a random walk and has a continuous, constant probability of encountering a stomate that is determined by stomatal cover (f_s). If $f_s \ll 1$, this can be modeled as a homogeneous Poisson process and the distance x a pathogen must travel before reaching a stomate follows an exponential distribution:

$$f(x) = f_s e^{-f_s x}$$

Given the constant death rate per unit distance H , the probability of surviving to distance x is e^{-Hx} . The probability of locating a stomate the probability of surviving to distance x multiplied by $f(x)$ and integrated over all x from 0 to ∞ :

$$\begin{aligned} p_{\text{locate}} &= \int_0^{\infty} f(x) e^{-Hx} dx \\ &= \int_0^{\infty} f_s e^{-(f_s + H)x} dx \\ &= \frac{f_s}{H + f_s} \end{aligned}$$

Substituting p_{locate} above into Equation 5:

$$\begin{aligned} p_{\text{colonize}} &= f_s + (1 - f_s) p_{\text{locate}} \\ &= f_s + (1 - f_s) \frac{f_s}{H + f_s} \\ &= f_s \left(1 + \frac{1 - f_s}{H + f_s} \right) \end{aligned}$$

With the spatially implicit model, because pathogens can potentially reach any stomate on the leaf, p_{colonize} is greater than that in the spatially explicit model for the same value of H . For example, if the pathogen can search forever ($H = 0$), then it will always colonize ($p_{\text{colonize}} = 1$; Figure S2a). But even when $H > 0$, p_{colonize} is significantly higher than in the spatially implicit than spatially explicit model for the same f_s because pathogens can potentially colonize any stomate on the leaf.

Whereas the spatially explicit model probably underestimates p_{locate} for pathogens that can search over long distances, the implicit model overestimates because it assumes that the probability of encountering a stomate is constant (i.e. homogeneous Poisson process). This is not true because stomata are discrete areas on the leaf. If a pathogen is searching far away from a stomate, its probability of encountering a stomate in the near future is lower than that for a pathogen searching near a stomate. This should be modeled as a *nonhomogenous* Poisson process. Future work should derive p_{locate} for the stomatal anatomies presented here under a nonhomogeneous process.

399 Despite the quantitative differences in the the spatially explicit and implicit models, they have similar
 400 qualitative properties when $H > 0$, which is reasonable since the leaf surface is a relatively hostile
 401 environment for most pathogens (see Introduction). In both models, p_{colonize} increases with, but is higher
 402 than f_S . In the spatially explicit model, size-density scaling that preserves p_{colonize} is 1 when $H = 0$ and
 403 slightly less than 1 otherwise (Figure 4). In the spatially implicit model, the scaling coefficient is always 1.
 404 Rearranging the equation for p_{colonize} above and substituting $f_S = DS$, the following relationship holds:

$$SD = \frac{p_{\text{colonize}}H}{1 - p_{\text{colonize}} + H}$$

405 Since H is a constant the right-hand side of the equation above is constant for a given value of p_{colonize} .
 406 Hence the β_p that would preserve the relationship above is simply 1 (Figure S3).

REFERENCES

- 407 Allaire, J., Xie, Y., McPherson, J., Luraschi, J., Ushey, K., Wickham, H., Cheng, J., Chang, W., and
408 Iannone, R. (2020). Rmarkdown: Dynamic Documents for R. Available at: <https://github.com/rstudio/rmarkdown>.
409 *rstudio/rmarkdown*.
- 410 Allen, E. A., Hazen, B. E., Hoch, H. C., Kwon, Y., Leinhos, G. M. E., Staples, R. C., Stumpf, M. A., and
411 Terhune, B. T. (1991). Appressorium Formation in Response to Topographical Signals by 27 Rust Species.
412 *Phytopathology* 81, 323. doi:10.1094/Phyto-81-323.
- 413 Assmann, S. M., and Zeiger, E. (1987). “Guard Cell Bioenergetics,” in *Stomatal Function*, eds. E. Zeiger,
414 G. D. Farquhar, and I. R. Cowan (Stanford University Press), 163–193.
- 415 Baresch, A., Crifò, C., and Boyce, C. K. (2019). Competition for epidermal space in the evolution of
416 leaves with high physiological rates. *New Phytologist* 221, 628–639. doi:10.1111/nph.15476.
- 417 Bastiaans, L. (1991). Ratio Between Virtual and Visual Lesion Size as a Measure to Describe Reduction
418 in Leaf Photosynthesis of Rice Due to Leaf Blast. *Phytopathology* 81, 611. doi:10.1094/Phyto-81-611.
- 419 Beattie, G. A., and Lindow, S. E. (1995). The secret life of foliar bacterial pathogens on leaves. *Annual*
420 *Review of Phytopathology* 33, 145–172.
- 421 Berry, J. A., Beerling, D. J., and Franks, P. J. (2010). Stomata: Key players in the earth system, past and
422 present. *Current Opinion in Plant Biology* 13, 232–239. doi:10.1016/j.pbi.2010.04.013.
- 423 Boer, H. J. de, Price, C. A., Wagner-Cremer, F., Dekker, S. C., Franks, P. J., and Veneklaas, E. J.
424 (2016). Optimal allocation of leaf epidermal area for gas exchange. *New Phytologist* 210, 1219–1228.
425 doi:10.1111/nph.13929.
- 426 Borchers, H. W. (2019). Pracma: Practical Numerical Math Functions. R package version 2.2.9. Available
427 at: <https://CRAN.R-project.org/package=pracma>.
- 428 Brand, A., and Gow, N. A. R. (2012). “Tropic Orientation Responses of Pathogenic Fungi,” in
429 *Morphogenesis and Pathogenicity in Fungi*, eds. J. Pérez-Martín and A. Di Pietro (Berlin, Heidelberg:
430 Springer Berlin Heidelberg), 21–41. doi:10.1007/978-3-642-22916-9_2.
- 431 Brodribb, T. J., Jordan, G. J., and Carpenter, R. J. (2013). Unified changes in cell size permit coordinated
432 leaf evolution. *New Phytologist* 199, 559–570. doi:10.1111/nph.12300.
- 433 Brown, H. T., and Escombe, F. (1900). Static diffusion of gases and liquids in relation to the assimilation
434 of carbon and translocation in plants. *Proceedings of the Royal Society of London* 67, 124–128.
- 435 Bucher, S. F., Auerswald, K., Grün-Wenzel, C., Higgins, S. I., Garcia Jorge, J., and Römermann, C.
436 (2017). Stomatal traits relate to habitat preferences of herbaceous species in a temperate climate. *Flora*
437 229, 107–115. doi:10.1016/j.flora.2017.02.011.
- 438 Buckley, T. N. (2019). How do stomata respond to water status? *New Phytologist* 224, 21–36.
439 doi:10.1111/nph.15899.
- 440 Chater, C. C. C., Caine, R. S., Fleming, A. J., and Gray, J. E. (2017). Origins and Evolution of Stomatal
441 Development. *Plant Physiology* 174, 624–638. doi:10.1104/pp.17.00183.
- 442 Dow, G. J., Bergmann, D. C., and Berry, J. A. (2014a). An integrated model of stomatal development and
443 leaf physiology. *New Phytologist* 201, 1218–1226.

- 444 Dow, G. J., Berry, J. A., and Bergmann, D. C. (2014b). The physiological importance of developmental
445 mechanisms that enforce proper stomatal spacing in *Arabidopsis thaliana*. *New Phytologist* 201, 1205–1217.
446 doi:10.1111/nph.12586.
- 447 Drake, P. L., Boer, H. J., Schymanski, S. J., and Veneklaas, E. J. (2019). Two sides to every leaf: Water and
448 CO_2 transport in hypostomatous and amphistomatous
449 leaves. *New Phytologist* 222, 1179–1187. doi:10.1111/nph.15652.
- 450 Drake, P. L., Froend, R. H., and Franks, P. J. (2013). Smaller, faster stomata: Scaling of stomatal
451 size, rate of response, and stomatal conductance. *Journal of Experimental Botany* 64, 495–505.
452 doi:10.1093/jxb/ers347.
- 453 Dutton, C., Hōrak, H., Hepworth, C., Mitchell, A., Ton, J., Hunt, L., and Gray, J. E. (2019).
454 Bacterial infection systemically suppresses stomatal density. *Plant, Cell & Environment* 42, 2411–2421.
455 doi:10.1111/pce.13570.
- 456 Farquhar, G. D., and Sharkey, T. D. (1982). Stomatal Conductance and Photosynthesis. *Annual Review of*
457 *Plant Physiology* 33, 317–345. doi:10.1146/annurev.pp.33.060182.001533.
- 458 Fawke, S., Doumane, M., and Schornack, S. (2015). Oomycete Interactions with Plants: Infection
459 Strategies and Resistance Principles. *Microbiology and Molecular Biology Reviews* 79, 263–280.
460 doi:10.1128/MMBR.00010-15.
- 461 Fetter, K. C., Nelson, D. M., and Keller, S. R. (2019). Trade-offs and selection conflicts in hybrid
462 poplars indicate the stomatal ratio as an important trait regulating disease resistance. University of Vermont
463 doi:10.1101/814046.
- 464 Flexas, J., Cano, F. J., Carriquí, M., Coopman, R. E., Mizokami, Y., Tholen, D., and Xiong, D. (2018).
465 “CO₂ Diffusion Inside Photosynthetic Organs,” in *The Leaf: A Platform for Performing Photosynthesis*
466 *Advances in Photosynthesis and Respiration*, eds. W. W. Adams III and I. Terashima (Cham: Springer
467 International Publishing), 163–208. doi:10.1007/978-3-319-93594-2_7.
- 468 Franks, P. J., and Beerling, D. J. (2009a). CO₂ -forced evolution of plant gas exchange capacity and
469 water-use efficiency over the Phanerozoic. *Geobiology* 7, 227–236. doi:10.1111/j.1472-4669.2009.00193.x.
- 470 Franks, P. J., and Beerling, D. J. (2009b). Maximum leaf conductance driven by CO₂ effects on stomatal
471 size and density over geologic time. *Proceedings of the National Academy of Sciences* 106, 10343–10347.
- 472 Franks, P. J., Drake, P. L., and Beerling, D. J. (2009). Plasticity in maximum stomatal conductance
473 constrained by negative correlation between stomatal size and density: An analysis using *Eucalyptus*
474 *globulus*. *Plant, Cell & Environment* 32, 1737–1748. doi:10.1111/j.1365-3040.2009.002031.x.
- 475 Franks, P. J., and Farquhar, G. D. (2001). The effect of exogenous abscisic acid on stomatal development,
476 stomatal mechanics, and leaf gas exchange in *Tradescantia virginiana*. *Plant Physiology* 125, 935–942.
- 477 Franks, P. J., and Farquhar, G. D. (2007). The Mechanical Diversity of Stomata and Its Significance in
478 Gas-Exchange Control. *Plant Physiology* 143, 78–87. doi:10.1104/pp.106.089367.
- 479 Franks, P. J., Royer, D. L., Beerling, D. J., Van de Water, P. K., Cantrill, D. J., Barbour, M. M.,
480 and Berry, J. A. (2014). New constraints on atmospheric CO₂ concentration for the Phanerozoic:
481 Franks et al.: New constraints on Phanerozoic CO₂. *Geophysical Research Letters* 41, 4685–4694.
482 doi:10.1002/2014GL060457.

- 483 Gilbert, G. S. (2002). Evolutionary Ecology of Plant Diseases in Natural Ecosystems. *Annual Review of*
484 *Phytopathology* 40, 13–43. doi:10.1146/annurev.phyto.40.021202.110417.
- 485 Gray, A., Liu, L., and Facette, M. (2020). Flanking Support: How Subsidiary Cells Contribute to Stomatal
486 Form and Function. *Frontiers in Plant Science* 11. doi:10.3389/fpls.2020.00881.
- 487 Gutschick, V. P. (1984). Photosynthesis model for C₃ leaves incorporating CO₂ transport, propagation of
488 radiation, and biochemistry 1. Kinetics and their parameterization. *Photosynthetica* 18, 549–568.
- 489 Harrison, E. L., Arce Cubas, L., Gray, J. E., and Hepworth, C. (2019). The influence of
490 stomatal morphology and distribution on photosynthetic gas exchange. *The Plant Journal*, tpj.14560.
491 doi:10.1111/tpj.14560.
- 492 Hetherington, A. M., and Woodward, F. I. (2003). The role of stomata in sensing and driving
493 environmental change. *Nature* 424, 901–908. doi:10.1038/nature01843.
- 494 Hoch, H. C., Staples, R. C., Whitehead, B., Comeau, J., and Wolf, E. D. (1987). Signaling for Growth
495 Orientation and Cell Differentiation by Surface Topography in Uromyces. *Science, New Series* 235,
496 1659–1662. Available at: <http://www.jstor.org/stable/1698314>.
- 497 Jones, H. G. (1985). Partitioning stomatal and non-stomatal limitations to photosynthesis. *Plant, Cell &*
498 *Environment* 8, 95–104. doi:10.1111/j.1365-3040.1985.tb01227.x.
- 499 Jordan, G. J., Carpenter, R. J., and Brodribb, T. J. (2014). Using fossil leaves as
500 evidence for open vegetation. *Palaeogeography, Palaeoclimatology, Palaeoecology* 395, 168–175.
501 doi:10.1016/j.palaeo.2013.12.035.
- 502 Kiefer, B., Riemann, M., Büche, C., Kassemeyer, H.-H., and Nick, P. (2002). The host guides
503 morphogenesis and stomatal targeting in the grapevine pathogen *Plasmopara viticola*. *Planta* 215, 387–393.
504 doi:10.1007/s00425-002-0760-2.
- 505 Lawson, T., and Blatt, M. R. (2014). Stomatal Size, Speed, and Responsiveness Impact on Photosynthesis
506 and Water Use Efficiency. *Plant Physiology* 164, 1556–1570. doi:10.1104/pp.114.237107.
- 507 Lawson, T., Terashima, I., Fujita, T., and Wang, Y. (2018). “Coordination Between Photosynthesis and
508 Stomatal Behavior,” in *The Leaf: A Platform for Performing Photosynthesis* Advances in Photosynthesis
509 and Respiration., eds. W. W. Adams III and I. Terashima (Cham: Springer International Publishing),
510 141–161. doi:10.1007/978-3-319-93594-2_6.
- 511 Lehmann, P., and Or, D. (2015). Effects of stomata clustering on leaf gas exchange. *New Phytologist* 207,
512 1015–1025. doi:10.1111/nph.13442.
- 513 McElwain, J. C., Yiotis, C., and Lawson, T. (2016). Using modern plant trait relationships between
514 observed and theoretical maximum stomatal conductance and vein density to examine patterns of plant
515 macroevolution. *New Phytologist* 209, 94–103. doi:10.1111/nph.13579.
- 516 McKown, A. D., Guy, R. D., Quamme, L., Klápště, J., La Mantia, J., Constabel, C. P., El-Kassaby,
517 Y. A., Hamelin, R. C., Zifkin, M., and Azam, M. S. (2014). Association genetics, geography and
518 ecophysiology link stomatal patterning in *Populus trichocarpa* with carbon gain and disease resistance
519 trade-offs. *Molecular Ecology* 23, 5771–5790. doi:10.1111/mec.12969.
- 520 McKown, A. D., Klápště, J., Guy, R. D., Corea, O. R. A., Fritsche, S., Ehrling, J., El-Kassaby, Y. A., and
521 Mansfield, S. D. (2019). A role for *speechless* in the integration of leaf stomatal patterning with the growth
522 vs disease trade-off in poplar. *New Phytologist*, nph.15911. doi:10.1111/nph.15911.

- 523 McLachlan, D. H., Kopischke, M., and Robatzek, S. (2014). Gate control: Guard cell regulation by
524 microbial stress. *New Phytologist* 203, 1049–1063. doi:10.1111/nph.12916.
- 525 Melotto, M., Underwood, W., Koczan, J., Nomura, K., and He, S. Y. (2006). Plant Stomata Function in
526 Innate Immunity against Bacterial Invasion. *Cell* 126, 969–980. doi:10.1016/j.cell.2006.06.054.
- 527 Melotto, M., Zhang, L., Oblessuc, P. R., and He, S. Y. (2017). Stomatal Defense a Decade Later. *Plant*
528 *Physiology* 174, 561–571. doi:10.1104/pp.16.01853.
- 529 Metcalfe, C. R., and Chalk, L. (1950). *Anatomy of the dicotyledons, Vols. 1 & 2*. First. Oxford: Oxford
530 University Press.
- 531 Meurer, A., Smith, C. P., Paprocki, M., Čertík, O., Kirpichev, S. B., Rocklin, M., Kumar, A., Ivanov, S.,
532 Moore, J. K., Singh, S., et al. (2017). SymPy: Symbolic computing in Python. *PeerJ Computer Science* 3,
533 e103. doi:10.7717/peerj-cs.103.
- 534 Mitchell, C. E. (2003). Trophic control of grassland production and biomass by pathogens. *Ecology*
535 *Letters* 6, 147–155. doi:10.1046/j.1461-0248.2003.00408.x.
- 536 Monteith, J. L., and Unsworth, M. H. (2013). *Principles of environmental physics: Plants, animals, and*
537 *the atmosphere*. 4th ed. Amsterdam ; Boston: Elsevier/Academic Press.
- 538 Morison, J. I. L., Emily Gallouët, Lawson, T., Cornic, G., Herbin, R., and work(s): N. R. B. R. (2005).
539 Lateral Diffusion of CO₂ in Leaves Is Not Sufficient to Support Photosynthesis. *Plant Physiology* 139,
540 254–266. Available at: <http://www.jstor.org/stable/4281859>.
- 541 Mott, K. A., Gibson, A. C., and O’Leary, J. W. (1984). The adaptive significance of amphistomatic leaves.
542 *Plant, Cell & Environment* 5, 455–460.
- 543 Muir, C. D. (2015). Making pore choices: Repeated regime shifts in stomatal ratio. *Proceedings of the*
544 *Royal Society B: Biological Sciences* 282, 20151498. doi:10.1098/rspb.2015.1498.
- 545 Muir, C. D., Hangarter, R. P., Moyle, L. C., and Davis, P. A. (2014). Morphological and anatomical
546 determinants of mesophyll conductance in wild relatives of tomato (*solanum* sect. *Lycopersicon* , sect.
547 *Lycopersicoides* ; Solanaceae). *Plant, Cell & Environment* 37, 1415–1426. doi:10.1111/pce.12245.
- 548 Murray, M., Soh, W. K., Yiotis, C., Spicer, R. A., Lawson, T., and McElwain, J. C. (2019). Consistent
549 relationship between field-measured stomatal conductance and theoretical maximum stomatal conductance
550 in C₃ woody angiosperms in four major biomes. *International Journal of Plant Sciences*, 706260.
551 doi:10.1086/706260.
- 552 Murray, R. R., Emblow, M. S. M., Hetherington, A. M., and Foster, G. D. (2016). Plant virus infections
553 control stomatal development. *Scientific Reports* 6, 34507. doi:10.1038/srep34507.
- 554 Oguchi, R., Onoda, Y., Terashima, I., and Tholen, D. (2018). “Leaf Anatomy and Function,” in *The Leaf:*
555 *A Platform for Performing Photosynthesis* Advances in Photosynthesis and Respiration., eds. W. W. Adams
556 III and I. Terashima (Cham: Springer International Publishing), 97–139. doi:10.1007/978-3-319-93594-2_5.
- 557 Papanatsiou, M., Amtmann, A., and Blatt, M. R. (2017). Stomatal clustering in Begonia associates with
558 the kinetics of leaf gaseous exchange and influences water use efficiency. *Journal of Experimental Botany*
559 68, 2309–2315. doi:10.1093/jxb/erx072.
- 560 Parkhurst, D. F. (1994). Diffusion of CO₂ and Other Gases Inside Leaves. *New Phytologist*
561 126, 449–479. Available at: <http://www.jstor.org/stable/2557929>.

- 562 Parkhurst, D. F. (1978). The Adaptive Significance of Stomatal Occurrence on One or Both Surfaces of
563 Leaves. *The Journal of Ecology* 66, 367. doi:10.2307/2259142.
- 564 Parkhurst, D. F., and Mott, K. A. (1990). Intercellular Diffusion Limits to CO₂ Uptake in Leaves: Studies
565 in Air and Helox. *Plant Physiology* 94, 1024–1032. doi:10.1104/pp.94.3.1024.
- 566 Parlange, J.-Y., and Waggoner, P. E. (1970). Stomatal Dimensions and Resistance to Diffusion. *Plant*
567 *Physiology* 46, 337–342. doi:10.1104/pp.46.2.337.
- 568 Peat, H. J., and Fitter, A. H. (1994). A comparative study of the distribution and density of stomata in the
569 British flora. *Biological Journal of the Linnean Society* 52, 377–393.
- 570 Raissig, M. T., Matos, J. L., Anleu Gil, M. X., Kornfeld, A., Bettadapur, A., Abrash, E., Allison, H.
571 R., Badgley, G., Vogel, J. P., Berry, J. A., et al. (2017). Mobile MUTE specifies subsidiary cells to build
572 physiologically improved grass stomata. *Science* 355, 1215–1218. doi:10.1126/science.aal3254.
- 573 R Core Team (2020). *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R
574 Foundation for Statistical Computing Available at: <http://www.R-project.org/>.
- 575 Sack, L., and Buckley, T. N. (2016). The developmental basis of stomatal density and flux. *Plant*
576 *Physiology* 171, 2358–2363. doi:10.1104/pp.16.00476.
- 577 Sack, L., Cowan, P. D., Jaikumar, N., and Holbrook, N. M. (2003). The 'hydrology' of leaves: Co-
578 ordination of structure and function in temperate woody species. *Plant, Cell and Environment* 26, 1343–
579 1356. doi:10.1046/j.0016-8025.2003.01058.x.
- 580 Salisbury, E. J. (1928). On the Causes and Ecological Significance of Stomatal Frequency, with Special
581 Reference to the Woodland Flora. *Philosophical Transactions of the Royal Society B: Biological Sciences*
582 216, 1–65. doi:10.1098/rstb.1928.0001.
- 583 Tateda, C., Obara, K., Abe, Y., Sekine, R., Nekoduka, S., Hikage, T., Nishihara, M., Sekine, K.-T., and
584 Fujisaki, K. (2019). The Host Stomatal Density Determines Resistance to *Septoria gentianae* in Japanese
585 Gentian. *Molecular Plant-Microbe Interactions* 32, 428–436. doi:10.1094/MPMI-05-18-0114-R.
- 586 Tichá, I. (1982). Photosynthetic characteristics during ontogenesis of leaves 7. Stomata density and sizes.
587 *Photosynthetica* 16, 375–471.
- 588 Tucker, S. L., and Talbot, N. J. (2001). Surface attachment and pre-penetration stage development by plant
589 pathogenic fungi. *Annual Review of Phytopathology* 39, 385–417. doi:10.1146/annurev.phyto.39.1.385.
- 590 Underwood, W., Melotto, M., and He, S. Y. (2007). Role of plant stomata in bacterial invasion. *Cellular*
591 *Microbiology* 9, 1621–1629. doi:10.1111/j.1462-5822.2007.00938.x.
- 592 Ushey, K., Allaire, J. J., and Tang, Y. (2020). Reticulate: Interface to 'Python'. Available at: <https://CRAN.R-project.org/package=reticulate>.
- 594 Weiss, A. (1865). Untersuchungen über die Zahlen- und Grössenverhältnisse der Spaltöffnungen.
595 *Jahrbücher für Wissenschaftliche Botanik* 4, 125–196.
- 596 Xie, Y., Allaire, J. J., and Grolemond, G. (2018). *R Markdown: The definitive guide*. Boca Raton: Taylor
597 & Francis, CRC Press.
- 598 Zeng, W., Melotto, M., and He, S. Y. (2010). Plant stomata: A checkpoint of host immunity and pathogen
599 virulence. *Current Opinion in Biotechnology* 21, 599–603. doi:10.1016/j.copbio.2010.05.006.