

APPENDIX

Laurent Jutras-Dubé

Department of Physics
McGill University
3600 rue University
Montreal, QC H3A 2T8
Canada

Ezzat El-Sherif

Department of Biology
Friedrich-Alexander-Universität Erlangen-Nürnberg
Staudtstraße 5
Erlangen 91058
Germany

Paul François

Department of Physics
McGill University
3600 rue University
Montreal, QC H3A 2T8
Canada

1 A two-enhancer model reproduces dynamical features of Tribolium segmentation

In [1], two of us proposed a model of Tribolium segmentation relying on the interplay of two sets of enhancers. In short, two sets of enhancers (static $S(P)$, dynamic $D(P)$) were used, where the role of parameter g is played by morphogen *Caudal* (*cad*) [2] (Figure 1–figure supplement 2). $S(P)$ encodes a multistable system and $D(P)$ a sequential cascade of genetic expression of gap genes (*hb*, *Kr*, *mlpt*, *gt*). This system was found to implement a “speed gradient” model, where the speed of traveling waves of gap genes from posterior to anterior depended on the level of *cad* concentration (Figure 1–figure supplement 2B–D). This led to robust patterning of the embryo (Figure 1–figure supplement 2E) but the mathematical origin of the speed gradient was not explained.

To better understand the underlying dynamics of the system, we consider the time courses of multiple cells at different positions and thus with different final fates. Figure 1–figure supplement 2F shows the projection of the cells’ dynamics on a 2D plane corresponding to the first two genes expressed in the cascade (*Kr* and *hb*), as well as a typical flow for different values of *cad* while keeping the other genes (*mlpt*, *gt* and X) at zero. Importantly, for $cad = 0.13$, we see the appearance of a new fixed point (green disk on Figure 1–figure supplement 2F).

We make four observations:

- The flow of the system is canalized. The trajectories of the cells stay very close to one another in phase space.
- As *cad* is lowered, the new fixed point appears very close to the common trajectory of all cells (Figure 1–figure supplement 2F, top row), and clearly separates the trajectories of cells ending up at different fates (Figure 1–figure supplement 2F, bottom row).
- When *cad* further decreases, the new fixed point moves in the high *hb*, low *Kr* region, corresponding to the eventual fate of Cell 1.
- When the new fixed point appears, the flow of cells past this fixed point is slowed down (Figure 1–figure supplement 2G).

23 These four observations offer a concise explanation to the “speed gradient” model: as the system gets closer to the
 24 bifurcation happening at $cad = 0.13$, the system is slowing down because of the future fixed points appearing on the
 25 trajectory. Intuitively, this is due to the fact that a fixed point corresponds to a frozen state, and thus to an infinite
 26 time-scale (static). When cad varies, the system has to interplay between a non-zero time-scale (dynamics) and such
 27 infinite time-scale, and it thus makes sense *a priori* that in between, the time-scale of the system diverges. This
 28 mechanism is close in principle to the critical timing proposed in [3].

29 **2 List of the functions used for the dynamics of each model**

30 **2.1 Gene network models**

31 In the gene network models, biochemical interactions between genes are modeled explicitly. Ordinary differential
 32 equations (ODEs) represent the dynamics of the concentration of the proteins that are encoded by the genes in the
 33 network. The deterministic part of the dynamics is composed of a protein production term and a protein degradation
 34 term. The production rate of a given protein can be altered by the interactions between the genes. Hill functions are use
 35 to model repression and activation of the production of a given protein by the genes. When multiple genes affect the
 36 concentration of a protein, the Hill functions corresponding to each interaction are multiplied. In the simulations, we set
 37 to 1 the maximal production rate of all proteins. Similarly, we set the degradation rate of all proteins to 1. In Eq. 1 of
 38 the main text, $C(P)$ encodes the degradation term, and $\Theta_S(g) S(P) + \Theta_D(g) D(P)$ represents the production term.

39 **2.1.1 3-gene models**

40 The proteins associated to the 3 genes are named arbitrarily A , B and C :

$$P = \begin{bmatrix} A \\ B \\ C \end{bmatrix} \quad C(P) = \begin{bmatrix} -A \\ -B \\ -C \end{bmatrix} \quad D(P) = \begin{bmatrix} \frac{1}{1 + (B/K_D^{B-A})^5} \\ \frac{1}{1 + (C/K_D^{C-B})^5} \\ \frac{1}{1 + (A/K_D^{A-C})^5} \end{bmatrix} \quad S(P) = \begin{bmatrix} \frac{1}{1 + (B/K_S^{B-A})^5} \frac{1}{1 + (C/K_S^{C-A})^5} \\ \frac{1}{1 + (C/K_S^{C-B})^5} \frac{1}{1 + (A/K_S^{A-B})^5} \\ \frac{1}{1 + (A/K_S^{A-C})^5} \frac{1}{1 + (B/K_S^{B-C})^5} \end{bmatrix} \quad (1)$$

41 Table 1 lists the values of the parameters used in the repression interactions of all versions of the 3-gene models: the
 42 symmetric version used to generate the results of Figure 2, Figure 2–figure supplements 1 and 2, Figure 3, Figure
 43 3–figure supplement 1 and Figure 7–figure supplement 2, the version with a weak asymmetry used in Figure 5, the
 44 version with a strong asymmetry used in Figure 5–figure supplement 1 and the version with a randomized asymmetry
 45 used in Figure 5–figure supplement 2. In the latter version, we randomly picked the values of the repression interactions
 46 of the static term $S(P)$ from a Gaussian distribution with mean 0.4 and standard deviation 0.04. Table 2 lists the weights
 47 $\Theta_D(g)$ and $\Theta_S(g)$ used for all 3-gene models: Models 1 and 2 used to generate the results of Figure 2, Figure 2–figure

48 supplement 1, Figure 3, Figure 5, Figure 5–figure supplements 1 and 2 and Figure 7–figure supplement 2, as well as
 49 Models 1 and 2 with Hill functions for the weights $\Theta_D(g)$ and $\Theta_S(g)$, used in Figure 2–figure supplement 1, Figure
 50 3–figure supplement 1 and Figure 7–figure supplement 2.

Table 1: Parameter values for the repression interactions of the 3-gene models

Model version	$K_D^{B \rightarrow A}$	$K_D^{C \rightarrow B}$	$K_D^{A \rightarrow C}$	$K_S^{B \rightarrow A}$	$K_S^{C \rightarrow A}$	$K_S^{C \rightarrow B}$	$K_S^{A \rightarrow B}$	$K_S^{A \rightarrow C}$	$K_S^{B \rightarrow C}$
Symmetric	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4
Weak asymmetry	0.4	0.4	0.4	0.36	0.36	0.4	0.4	0.4	0.4
Strong asymmetry	0.4	0.4	0.4	0.32	0.32	0.36	0.36	0.4	0.4
Randomized asymmetry	0.4	0.4	0.4	0.3825	0.3560	0.4334	0.4102	0.3802	0.4038

Table 2: Weights of the dynamic and static terms of the 3-gene models

Weights	Model 1	Model 2	Model 1 with Hill functions	Model 2 with Hill functions
$\Theta_D(g)$	g^2	g	$\frac{(g/0.4)^5}{1+(g/0.4)^5}$	$\frac{(g/0.4)^5}{1+(g/0.4)^5}$
$\Theta_S(g)$	$(1-g)^2$	$1-g$	$\frac{1}{1+(g/0.6)^5}$	$\frac{1}{1+(g/0.4)^5}$

51 2.1.2 Model of Tribolium segmentation

52 In the model of Figure 1–figure supplement 2, the interactions between *hunchback* (*hb*), *Krüppel* (*Kr*), *mille-pattes*
 53 (*mlpt*), *giant* (*gt*) and an unidentified gene *X* are modeled (see the supplement of [1]). Note that the role of parameter *g*
 54 is played by *caudal* (*cad*) in the model for Tribolium segmentation.

$$P = \begin{bmatrix} hb \\ Kr \\ mlpt \\ gt \\ X \end{bmatrix} \quad C(P) = \begin{bmatrix} -hb \\ -Kr \\ -mlpt \\ -gt \\ -X \end{bmatrix} \quad D(P) = \begin{bmatrix} \frac{(hb/0.2)^5}{1+(hb/0.2)^5} \frac{1}{1+(Kr/0.12)^5} \\ \frac{(hb/0.4)^5}{1+(hb/0.4)^5} \frac{1}{1+(mlpt/0.25)^5} \frac{1}{1+(gt/0.01)^5} \\ \frac{(Kr/0.4)^5}{1+(Kr/0.4)^5} \frac{1}{1+(gt/0.3)^5} \\ \frac{(mlpt/0.4)^5}{1+(mlpt/0.4)^5} \frac{1}{1+(X/0.08)^5} \\ \frac{(gt/0.4)^5}{1+(gt/0.4)^5} \end{bmatrix} \quad (2)$$

$$S(P) = \begin{bmatrix} \frac{(hb/0.4)^5}{1+(hb/0.4)^5} \frac{1}{1+(Kr/0.4)^5} \\ \frac{(Kr/0.4)^5}{1+(Kr/0.4)^5} \frac{1}{1+(hb/0.01)^5} \\ \frac{(mlpt/0.4)^5}{1+(mlpt/0.4)^5} \\ \frac{(gt/0.4)^5}{1+(gt/0.4)^5} \\ \frac{(X/0.4)^5}{1+(X/0.4)^5} \end{bmatrix} \quad \Theta_D(cad) = 3 \frac{cad}{1+cad} \quad \Theta_S(cad) = \frac{1}{1+cad} \quad (3)$$

55 2.2 Gene-free models

56 In the gene-free model, ODEs encode flows in an abstract 2D phase space. The two geometric variables are named
57 arbitrarily y and z :

$$P = \begin{bmatrix} y \\ z \end{bmatrix} \quad C(P) = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \quad D(P) = \begin{bmatrix} y(1 - \sqrt{y^2 + z^2}) - z \\ z(1 - \sqrt{y^2 + z^2}) + y \end{bmatrix} \quad S(P) = \begin{bmatrix} (y_0 - y)(y_1 - y)(y_2 - y) \\ -z \end{bmatrix} \quad (4)$$

58 where parameter y_0 (resp. y_1 and y_2) controls the position of the unstable fixed point (resp. the stable fixed points)
59 of the static term along the y axis. Parameter y_0 is set to 0 in the symmetric version of the model used in Figure 4,
60 Figure 4–figure supplements 1, 2, 3, 4 and 5, Figure 4–movie supplements 1, 2, 3 and 4, Figure 6, Figure 7, Figure
61 7–figure supplement 1 and Figure 7–movie supplement 1, as well as in the asymmetric versions of Figure 6–figure
62 supplement 1 and Figure 7–movie supplement 1. In Figure 6, parameter y_0 is set to 0.05 and 0.1 to model different
63 levels of asymmetry in the basins of attraction. In Figure 7–figure supplement 1, parameter y_0 is set to 0.02 for Model 1
64 and 0.05 for Model 2 to obtain a similar level of asymmetry in the final pattern generated by the two models. We set
65 $y_1 = -1$ and $y_2 = 1$ for all versions of the gene-free models, except for the asymmetric versions used to generate the
66 results of Figure 6–figure supplement 1 and Figure 7–movie supplement 1. In the former, the stable fixed points of the
67 static module are placed outside the region delimited by the limit cycle of the dynamic module by setting $y_1 = -2$ and
68 $y_2 = 2.5$, while in the latter, we set $y_1 = -1.75$ and $y_2 = 1$.

69 To obtain a supercritical Hopf bifurcation with the gene-free model, we followed a similar approach than for the
70 3-gene model. We reasoned that the sum of the weights of the dynamic and static modules should become smaller
71 than a degradation-like term for values of g around 0.5. For this reason, an "intermediate term" $I(P) = [-z \quad -y]^T$ is
72 introduced in the ODE. The intermediate term is weighted by the function $\Theta_I(g)$. Eq. 1 of the main text thus becomes:

$$\dot{P} = \Theta_D(g) D(P) + \Theta_I(g) I(P) + \Theta_S(g) S(P) + \eta(g, P) \quad (5)$$

73 Recall that in a given cell, only the dynamic module should be present at the beginning of the simulation, when $g = 1$.
74 Similarly, only the static module should be present at the end of the simulation, when $g = 0$. Therefore, we set the
75 weight of the intermediate module equal to $g(1 - g)$, which is zero at both $g = 1$ and $g = 0$. Since this weight is of the
76 order 2 in g , we make the weights of the dynamic and static modules of the order 3 in g to ensure that they become
77 smaller than the weight of the intermediate term for g around 0.5. To obtain subcritical Hopf bifurcations with the
78 gene-free model, we used a slightly different dynamic module:

$$D(P) = \begin{bmatrix} y \sqrt{y^2 + z^2} (1 - \sqrt{y^2 + z^2}) - z \\ z \sqrt{y^2 + z^2} (1 - \sqrt{y^2 + z^2}) + y \end{bmatrix} \quad (6)$$

79 Table 3 lists the weights used for all gene-free models: Model 1 used to generate the results of Figure 4–figure
80 supplements 1, 2 and 5, Figure 4–movie supplement 1, Figure 6, Figure 6–figure supplement 1, Figure 7–figure
81 supplements 1 and 2, and Figure 7–movie supplement 1, Model 2 used to generate the results of Figure 4, Figure
82 4–figure supplements 1 and 5, Figure 4–movie supplement 2, Figure 6, Figure 6–figure supplement 1, Figure 7, Figure
83 7–figure supplements 1 and 2, and Figure 7–movie supplement 1, Model 3 used to generate the results of Figure 4–figure
84 supplements 3 and 5, Figure 4–movie supplement 3 and Figure 7–figure supplement 2, and Model 4 used to generate
85 the results of Figure 4–figure supplements 4 and 5, Figure 4–movie supplement 4 and Figure 7–figure supplement 2.

Table 3: Weights of the dynamic, static and intermediate terms of the gene-free models

Weights	Models 1 and 3	Models 2 and 4
$\Theta_D(g)$	g^3	g
$\Theta_S(g)$	$(1 - g)^3$	$1 - g$
$\Theta_I(g)$	$g(1 - g)$	0

86 2.3 Infinite-period scenarios of Figure 1 and Figure 7

87 The infinite-period scenario of Figure 1B-F is a simplified version of the model of the appendix of [4]. The dynamics of
88 the phase of the oscillators are modeled directly using the following ODE:

$$\dot{\phi} = \omega(g) = \frac{\pi}{2} g^2 \quad (7)$$

89 The infinite-period scenario of Figure 7A-E is the 1D model of coupled oscillators from [5]. In brief, the dynamics of
 90 the phase of the oscillators are described by the following ODE:

$$\dot{\phi}(x, t) = \omega(x, t) + \frac{\epsilon}{2a^2} \left(\sin[\phi(x - a, t - \tau) - \phi(x, t)] + \sin[\phi(x + a, t - \tau) - \phi(x, t)] \right) \quad (8)$$

91 where ϵ represents the coupling strength between a cell and its 2 nearest neighbors, a is the average cell diameter (cd),
 92 and τ is the time delay in the coupling. The spatio-temporal profile of the frequency of the oscillators $\omega(x, t)$ is given
 93 by the following formula:

$$\omega(x, t) = \omega_\infty \left(1 - e^{-(x-vt)/\sigma} \right) \quad (9)$$

94 where ω_∞ represents the characteristic intrinsic frequency of the oscillators, v is the speed at which the spatial frequency
 95 profile moves along the posterior direction, and σ controls the spatial steepness of the frequency profile. Table 4 lists
 96 the parameter values used to generate the results of Figure 7A-E. See [5] for more details.

Table 4: Parameter values for the ODE of the phase oscillators in the infinite-period scenario of Figure 7

ϵ [cd ² /min]	a [cd]	τ [min]	ω_∞ [min ⁻¹]	v [cd/min]	σ [cd]
0.07	1	0	0.3886	0.255	36

97 2.4 Hopf scenario of Figure 1

98 The Hopf scenario of Figure 1G-K is the cell-autonomous model evolved *in silico* in [6]. The model describes the
 99 dynamics of two proteins, the effector protein E and the repressor protein R , under the control of morphogen g via
 100 ODEs with time delays:

$$\dot{E} = \left(\max \left[\frac{E^{n_1}}{E^{n_1} + E_E^{n_1}}, \frac{g^{n_2}}{g^{n_2} + g_E^{n_2}} \right] \frac{S_E}{1 + (R/R_E)^{n_3}} \right)_{t-\tau_E} - \delta_E E \quad (10)$$

$$\dot{R} = \left(\frac{g^{n_4}}{g^{n_4} + g_R^{n_4}} \frac{S_R}{1 + (R/R_R)^{n_5}} \right)_{t-\tau_R} - \delta_R R \quad (11)$$

101 The subscript of a closed parenthesis indicates the time at which the expression inside the parenthesis is evaluated. If no
 102 such parenthesis with a subscript is present in a given expression, this expression is evaluated at time t . The values of
 103 all parameters are given in Tables 5 and 6.

Table 5: Parameter values for the ODE of the effector protein E in the Hopf scenario of Figure 1

S_E	R_E	g_E	E_E	τ_E	δ_E	n_1	n_2	n_3
0.7176	0.4942	0.0678	0.3213	0.48	0.8538	3	4.3549	4.5321

Table 6: Parameter values for the ODE of the repressor protein R in the Hopf scenario of Figure 1

S_R	R_R	g_R	τ_R	δ_R	n_4	n_5
0.9422	0.1156	0.5047	3.92	0.9759	3.2136	4.522

104 2.5 Van der Pol oscillator of Figure 7

105 The schematic of an "asymmetric wave" profile shown on Figure 7D is generated with the following ODEs describing a
 106 Van der Pol oscillator:

$$\dot{y} = \mu \left(y - \frac{y^3}{3} - z \right) \quad (12)$$

$$\dot{z} = \frac{y}{\mu} \quad (13)$$

107 where parameter μ is set to 2.5. The schematic of a "symmetric wave" profile shown on Figure 7D is a sine function.

108 3 Spatio-temporal profile of the control parameter for each model

109 For all models except the model for Tribolium segmentation and the infinite-period scenario of Figure 7A-E, the follow-
 110 ing function is used to describe the spatio-temporal profile of the input g , which is treated either as the concentration of
 111 a morphogen in the gene network models, or as an abstract control parameter in the gene-free models:

$$g(x, t) = H(x - vt) = \min \left[e^{s(x-vt+x_{\text{osc}})}, 1 \right] \quad (14)$$

112 where parameter s controls the steepness of the gradient and v represents the speed at which the gradient moves along
 113 the antero-posterior axis. Parameter x_{osc} allows to generate a few oscillations inside the first simulated cell before g
 114 starts decreasing. Note that the position vector x is normalized in all our simulations, such that positions are constrained
 115 from 0 to 1. Table 7 lists the values of the parameters used for the gradients of all models (except the model for
 116 Tribolium segmentation): the gradients of the infinite-period scenario and of the Hopf scenario used to generate the
 117 results of Figure 1 B-F and Figure 1 G-K, respectively, the shallow gradient used in the 3-gene models of Figure 2,
 118 Figure 2–figure supplements 1 and 2, Figure 3, Figure 3–figure supplement 1, Figure 5, Figure 5–figure supplements 1
 119 and 2, and Figure 7–figure supplement 2, the steep gradient used in the 3-gene models of Figure 5 and Figure 5–figure
 120 supplements 1 and 2, and the gradients used in the gene-free models of Figure 4, Figure 4–figure supplements 2, 3,

121 4 and 5, Figure 6, Figure 6–figure supplement 1, Figure 7, Figure 7–figure supplements 1 and 2 and Figure 7–movie
 122 supplement 1.

Table 7: Parameter values for the spatio-temporal profile of input g

Model	s	v	x_{osc}
Infinite-period scenario of Figure 1	0.5	0.08	0.2
Hopf scenario of Figure 1	0.5	3	0
3-gene models, shallow gradient	1	0.05	0.2
3-gene models, steep gradient	2.5	0.05	0.2
Gene-free models (Figure 4, its supplements and Figure 7–figure supplement 2)	0.5	0.035	0.2
Gene-free models (Figure 6 and its supplement)	1	0.036	0
Gene-free models (Figure 7 and Figure 7–figure supplement 1)	6	0.0042	0
Gene-free models (Figure 7–movie supplement 1)	0.9	0.04	0

123 In the model for Tribolium segmentation, the role of input g is played by the maternal gene cad . The dynamics of cad
 124 is modelled with a Hill function:

$$cad(x, t) = \frac{(x/x^*(t))^{n(t)}}{1 + (x/x^*(t))^{n(t)}} \quad (15)$$

125 where the time dependencies of parameters $x^*(t)$ and $n(t)$ encode respectively the regression of the morphogen gradient
 126 along the antero-posterior axis, and the gradual increase in the steepness of the morphogen gradient:

$$x^*(t) = \max[0.4, 0.4 + 0.2(t - 2)] \quad ; \quad n(t) = \max[4, 4e^{(t-2)}] \quad (16)$$

127 4 Integration schemes

128 4.1 Euler algorithm for deterministic simulations

129 Eq. 1 of the main text can be integrated via the Euler algorithm to obtain a time series representing the deterministic
 130 dynamics of vector P :

$$P(t + dt) = P(t) + \left(\Theta_D(g(t)) D(P(t)) + \Theta_S(g(t)) S(P(t)) + C(P(t)) \right) dt \quad (17)$$

131 The Euler algorithm, which is equivalent to approximating the temporal derivative of P by a first-order finite difference,
 132 was used to perform deterministic simulations of all versions of the 3-gene models (Figure 2, Figure 2–figure supple-
 133 ments 1 and 2, Figure 5, Figure 5–figure supplements 1 and 2 and Figure 7–figure supplement 2). A similar version

134 of this algorithm that includes the intermediate term was used for deterministic simulations of the gene-free models
 135 (Figure 4, Figure 4–figure supplements 2, 3 and 4, Figure 6, Figure 6–figure supplement 1, Figure 7, Figure 7–figure
 136 supplements 1 and 2 and Figure 7–movie supplement 1). The Euler algorithm was also used to perform simulations
 137 of the infinite-period and Hopf scenarios (Figure 1 and Figure 7). On the other hand, deterministic simulations of the
 138 model for *Tribolium* segmentation were carried out via the `lsoda` integrator from the `scipy` library in Python (Figure
 139 1–figure supplement 2).

140 4.2 Langevin equation for stochastic simulations of the 3-gene models

141 The stochastic nature of chemical reactions, due at least partly to the finite number of molecules involved in these
 142 reactions, introduces fluctuations in protein concentrations in single cells. To generate the results of Figure 3 and Figure
 143 3–figure supplement 1, noise was introduced in the 3-gene models in a chemically realistic and mathematically rigorous
 144 way by following the method of [7]. In the generic formulation of the present problem, there are N molecular species
 145 S_i , $i = 1, \dots, N$, that can interact through M different reactions R_j , $j = 1, \dots, M$. Let $X_i(t)$ represent the number of
 146 S_i molecules at time t . Then, the vector $X(t) \equiv [X_1(t) \ \dots \ X_N(t)]$ represents the state of the whole system of N
 147 molecules at time t . For each reaction R_j , a propensity function a_j is defined such that if the system is in state X at time
 148 t , then $a_j(X) dt$ is the probability that one R_j reaction will occur in the next infinitesimal time interval dt , i.e. between
 149 t and $t + dt$. For each reaction R_j , a state-change vector ν_j is defined such that its i th component ν_{ji} represents the
 150 change in the number of S_i molecules produced by one R_j reaction. Once the M propensity functions and state-change
 151 vectors are defined, the time evolution of the state vector $X(t)$ is found via the N deterministic reaction rate equations:

$$\dot{X}_i(t) = \sum_{j=1}^M \nu_{ji} a_j(X(t)) \quad \text{for } i = 1, \dots, N \quad (18)$$

152 The numerical integration of these rate equations can be performed via the Euler algorithm:

$$X_i(t + dt) = X_i(t) + \sum_{j=1}^M \nu_{ji} a_j(X(t)) dt \quad \text{for } i = 1, \dots, N \quad (19)$$

153 The stochastic form of this simulation algorithm is given by the chemical Langevin equation:

$$X_i(t + dt) = X_i(t) + \sum_{j=1}^M \nu_{ji} a_j(X(t)) dt + \sum_{j=1}^M N_j(t) \nu_{ji} \sqrt{a_j(X(t))} dt \quad \text{for } i = 1, \dots, N \quad (20)$$

154 where $N_1(t), \dots, N_M(t)$ are M independent Gaussian random variables with mean and variance equal to 0 and 1,
 155 respectively, and that are not correlated in time. In the 3-gene models, the role of vector X is played by P . Note that
 156 re-scaling the numbers of proteins X_i by constant factors corresponds to multiplying both sides of Eq. 18 to 20 by
 157 that constant factor (as long as the state-change vectors ν_j are also re-scaled). Therefore, Eq. 18 to 20 are still valid
 158 when simulating protein concentrations scaled from 0 to 1 instead of absolute numbers of proteins. Furthermore, the

159 reactions of the 3-gene models are encoded in the protein production and degradation terms. The propensities of the
 160 protein production and degradation terms are respectively $\Theta_D(g) D(P) + \Theta_S(g) S(P)$ and P . Eq. 19 thus becomes eq.
 161 17, and eq. 20 can be re-written as the following expression:

$$P_i(t + dt) = P_i(t) + \left(\Theta_D(g(t)) D_i(P(t)) + \Theta_S(g(t)) S_i(P(t)) - P_i(t) \right) dt \quad i = 1, 2, 3$$

$$+ \left(N_i^{\text{prod}}(t) \sqrt{\Theta_D(g(t)) D_i(P(t)) + \Theta_S(g(t)) S_i(P(t))} - N_i^{\text{deg}}(t) \sqrt{P_i(t)} \right) \sqrt{dt} \quad (21)$$

162 where $N^{\text{prod}}(t) = [N_1^{\text{prod}}(t), N_2^{\text{prod}}(t), N_3^{\text{prod}}(t)]$ and $N^{\text{deg}}(t) = [N_1^{\text{deg}}(t), N_2^{\text{deg}}(t), N_3^{\text{deg}}(t)]$ are 2 vectors, each containing
 163 3 independent Gaussian random variables with mean 0 and variance 1. This equation can be simplified by leveraging
 164 the fact that the sum of Gaussian random variables with mean 0 and different variances is equal to a single Gaussian
 165 random variable with mean 0 and a variance equal to the sum of the variances:

$$P_i(t + dt) = P_i(t) + \left(\Theta_D(g(t)) D_i(P(t)) + \Theta_S(g(t)) S_i(P(t)) - P_i(t) \right) dt \quad i = 1, 2, 3$$

$$+ \left(N_i(t) \sqrt{\Theta_D(g(t)) D_i(P(t)) + \Theta_S(g(t)) S_i(P(t)) + P_i(t)} \right) \sqrt{dt} \quad (22)$$

166 where $N(t) = [N_1(t), N_2(t), N_3(t)]$ is a vector containing 3 independent Gaussian random variables with mean 0 and
 167 variance 1. Note that a different independent random variable is used for each protein, since the production term of
 168 each protein is due to a different combination of repression interactions. To control the level of noise, a parameter Ω is
 169 introduced in the previous equation such that increasing Ω decreases the level of noise:

$$P_i(t + dt) = P_i(t) + \left(\Theta_D(g(t)) D_i(P(t)) + \Theta_S(g(t)) S_i(P(t)) - P_i(t) \right) dt \quad i = 1, 2, 3$$

$$+ \left(\frac{N_i(t)}{\sqrt{\Omega}} \sqrt{\Theta_D(g(t)) D_i(P(t)) + \Theta_S(g(t)) S_i(P(t)) + P_i(t)} \right) \sqrt{dt} \quad (23)$$

170 Since noise arises at least partly from the stochastic nature of single reactions between a finite number of proteins,
 171 increasing the concentration of proteins is expected to buffer the intrinsic chemical noise. Therefore, the noise level
 172 is expected to decrease as the protein concentration is increased. The following mathematical derivation shows that
 173 parameter Ω can be interpreted as the typical concentration of proteins in the system, such that increasing the protein
 174 concentration corresponds to increasing the value of parameter Ω . First, let's take a look at the stochastic integration
 175 algorithm for protein A and write explicitly the maximal production rate ρ_A and the degradation rate δ_A :

$$\begin{aligned}
A^+ = & A + \left(\rho_A \left(\Theta_D(g) \frac{1}{1 + (B/K_D^{B \rightarrow A})^5} + \Theta_S(g) \frac{1}{1 + (B/K_S^{B \rightarrow A})^5} \frac{1}{1 + (C/K_S^{C \rightarrow A})^5} \right) - \delta_A A \right) dt \\
& + \frac{N_1}{\sqrt{\Omega}} \sqrt{ \rho_A \left(\Theta_D(g) \frac{1}{1 + (B/K_D^{B \rightarrow A})^5} + \Theta_S(g) \frac{1}{1 + (B/K_S^{B \rightarrow A})^5} \frac{1}{1 + (C/K_S^{C \rightarrow A})^5} \right) + \delta_A A } \sqrt{dt} \quad (24)
\end{aligned}$$

176 where a + superscript on a protein concentration indicates that this variable is evaluated at time $t + dt$ and the absence
177 of a superscript on a variable indicates that it is evaluated at time t . Multiplying both sides of the equation by Ω leads to
178 the following expression:

$$\begin{aligned}
\Omega A^+ = & \Omega A + \left(\Omega \rho_A \left(\Theta_D(g) \frac{1}{1 + (B/K_D^{B \rightarrow A})^5} + \Theta_S(g) \frac{1}{1 + (B/K_S^{B \rightarrow A})^5} \frac{1}{1 + (C/K_S^{C \rightarrow A})^5} \right) - \Omega \delta_A A \right) dt \quad (25) \\
& + N_1 \sqrt{ \Omega \rho_A \left(\Theta_D(g) \frac{1}{1 + (B/K_D^{B \rightarrow A})^5} + \Theta_S(g) \frac{1}{1 + (B/K_S^{B \rightarrow A})^5} \frac{1}{1 + (C/K_S^{C \rightarrow A})^5} \right) + \Omega \delta_A A } \sqrt{dt}
\end{aligned}$$

179 Now, let's re-scale all quantities that have the units of protein concentration by a factor of Ω . To achieve this, we
180 define the re-scaled variables $A^* = \Omega A$, $B^* = \Omega B$ and $C^* = \Omega C$, as well as re-scaled parameters $\rho_{A^*} = \Omega \rho_A$,
181 $K_D^{B^* \rightarrow A^*} = \Omega K_D^{B \rightarrow A}$, $K_S^{B^* \rightarrow A^*} = \Omega K_S^{B \rightarrow A}$ and $K_S^{C^* \rightarrow A^*} = \Omega K_S^{C \rightarrow A}$:

$$\begin{aligned}
A^{**} = & A^* + \left(\rho_{A^*} \left(\Theta_D(g) \frac{1}{1 + (B^*/K_D^{B^* \rightarrow A^*})^5} + \Theta_S(g) \frac{1}{1 + (B^*/K_S^{B^* \rightarrow A^*})^5} \frac{1}{1 + (C^*/K_S^{C^* \rightarrow A^*})^5} \right) - \delta_A A^* \right) dt \quad (26) \\
& + N_1 \sqrt{ \rho_{A^*} \left(\Theta_D(g) \frac{1}{1 + (B^*/K_D^{B^* \rightarrow A^*})^5} + \Theta_S(g) \frac{1}{1 + (B^*/K_S^{B^* \rightarrow A^*})^5} \frac{1}{1 + (C^*/K_S^{C^* \rightarrow A^*})^5} \right) + \delta_A A^* } \sqrt{dt}
\end{aligned}$$

182 A similar procedure can be followed for proteins B and C . Therefore, multiplying the stochastic term of the Langevin
183 equation for all proteins by $1/\sqrt{\Omega}$ is equivalent to re-scaling all variables and parameters that have the units of a protein
184 concentration by a factor of Ω . Since we set the maximal production rates and the degradation rates of all proteins to
185 1, the typical concentration of proteins A , B and C is normalized to 1. Re-scaling all protein concentrations and all
186 parameters with units of protein concentration by a factor of Ω thus corresponds to setting the typical concentration of
187 proteins to Ω . In conclusion, parameter Ω of equation 23 indeed corresponds to the typical concentration of proteins.

188 4.3 Cell-to-cell coupling in the 3-gene models

189 A strategy that a cell can use to fight the intrinsic noise in protein concentrations is to evaluate the protein expression
190 state of its neighbors and change its own protein expression state accordingly. In the stochastic simulations of the 3-gene
191 models, cell-to-cell communication is modelled via a diffusion term included in the differential equations describing the

192 dynamics of the set of protein concentrations. The higher the concentration of a given protein is in a given simulated
 193 cell, the more this protein will diffuse to neighboring simulated cells. Diffusion thus models the process of adjusting
 194 the protein concentration of a given cell according to the protein concentration of surrounding cells. The dynamics of
 195 vector P in the 3-gene models is therefore given by the following differential equation:

$$\frac{\partial P}{\partial t} = \Theta_D(g) D(P) + \Theta_S(g) S(P) - P + \eta(g, P) + D \frac{\partial^2 P}{\partial x^2} \quad (27)$$

196 where the diffusion constant D controls the strength of cell-to-cell coupling. The complete stochastic simulation
 197 algorithm for the 3-gene model thus becomes:

$$P_i(x, t + dt) = P_i(x, t) + \left(\Theta_D(g(x, t)) D_i(P(x, t)) + \Theta_S(g(x, t)) S_i(P(x, t)) - P_i(x, t) + D \frac{\partial^2 P_i}{\partial x^2} \right) dt \\ + \left(\frac{N_i(x, t)}{\sqrt{\Omega}} \sqrt{\Theta_D(g(x, t)) D_i(P(x, t)) + \Theta_S(g(x, t)) S_i(P(x, t)) + P_i(x, t)} \right) \sqrt{dt} \quad (28)$$

198 for $i = 1, 2, 3$. Note that diffusion is not included in the stochastic term, since diffusion of proteins is not a reaction in
 199 itself. In the simulations, the second spatial derivative is approximated by a second-order central finite difference with
 200 reflective boundaries.

201 4.4 Stochastic simulations of the gene-free models

202 Since the gene-free models simulate the dynamics of abstract variables that do not represent explicitly protein con-
 203 centrations, the variance of the noise is held independent of the state of the system. The stochastic algorithm used to
 204 generate the results of Figure 4–figure supplement 5 is therefore the following:

$$P_i(t + dt) = P_i(t) + \left(\Theta_D(g(t)) D_i(P(t)) + \Theta_I(g(t)) I_i(P(t)) + \Theta_S(g(t)) S_i(P(t)) - P_i(t) \right) dt + \frac{1}{\sqrt{\Omega}} N_i(t) \sqrt{dt} \quad (29)$$

205 where $i = 1, 2$, and $N(t) = [N_1(t), N_2(t)]$ is a vector containing 2 independent Gaussian random variables with mean 0
 206 and variance 1. Parameter Ω is still included to control the level of noise, but it cannot be interpreted as the typical
 207 concentration of proteins in the system since the gene-free models do not simulate explicitly protein interactions.

208 5 Mathematical formula for the mutual information

209 In deterministic simulations, the initial phase of the genetic oscillation inside a given cell determines in which part of the
 210 pattern this cell will end up. This is not necessarily the case in stochastic simulations. To quantify the robustness to noise
 211 of a given model for specific values of parameter Ω (and of the diffusion constant D in the case of the 3-gene models) it

212 is required to define a metric that measures the accuracy with which the initial phase of the genetic oscillations inside a
 213 cell predicts the region of the pattern in which this cell will end up. The specific metric used in Figure 3, Figure 3–figure
 214 supplement 1, Figure 4–figure supplement 5 and Figure 5–figure supplement 2 is the mutual information between the
 215 initial phase of the oscillator and the final protein expression state of the simulated cells. The mutual information $I(x, y)$
 216 between two discrete variables x and y is given by the following expression:

$$I(x, y) = \sum_{y \in Y} \sum_{x \in X} p(x, y) \log \left(\frac{p(x, y)}{p(x) p(y)} \right) \quad (30)$$

217 where X and Y are the sets of possible values for x and y , respectively. Intuitively, the mutual information between
 218 two variables quantifies the amount of information obtained on the value of the first variable by knowing the value
 219 of the second variable (and vice versa). If the logarithm is in base 2, the units of the mutual information are bits. To
 220 measure how precisely the phase of the oscillator is read to form the final pattern, variable x is set to the phase of the
 221 oscillation in protein expression at the beginning of the simulation ϕ_i , and variable y is set to the protein expression
 222 state at the end of the simulation P_f :

$$I(\phi_i, P_f) = \sum_{P_f} \sum_{\phi_i} p(\phi_i, P_f) \log \left(\frac{p(\phi_i, P_f)}{p(\phi_i) p(P_f)} \right) \quad (31)$$

$$= \sum_{P_f} \sum_{\phi_i} p(P_f | \phi_i) p(\phi_i) \log \left(\frac{p(P_f | \phi_i) p(\phi_i)}{p(\phi_i) p(P_f)} \right) \quad (32)$$

$$= \sum_{P_f} \sum_{\phi_i} p(P_f | \phi_i) p(\phi_i) \log \left(\frac{p(P_f | \phi_i)}{\sum_{\phi_i} p(P_f | \phi_i) p(\phi_i)} \right) \quad (33)$$

223 To get the second equality, the fact that $p(x, y) = p(x|y)p(y)$ for any two variables x and y was used to get rid of the joint
 224 probability $p(\phi, R_i)$, which is not straightforward to evaluate directly. Similarly, the fact that $p(y) = \sum_{x \in X} p(x, y) =$
 225 $\sum_{x \in X} p(x|y) p(y)$ for any two variables x and y was used to get rid of $p(P_f)$, which is less easy to compute than $p(\phi_i)$.
 226 Indeed, ϕ_i is sampled uniformly in the simulations of the 3-gene and gene-free models, since the speed of regression
 227 of the input g is constant throughout the simulations. In the 3-gene models, the different phases ϕ_i are defined as the
 228 different states of protein expression along the oscillation cycle generated by the dynamic module ($g = 1$). A uniform
 229 sample of ϕ_i is obtained by sampling this oscillation cycle at constant time intervals for a total time length of one period.
 230 In the gene-free models, the different phases ϕ_i are defined as the different sets of (y, z) values along the oscillation
 231 cycle generated by the dynamic module ($g = 1$). Since the oscillations are on the unit circle (centered at the origin) and
 232 have a constant speed along the cycle, sampling uniformly the angles from the positive y axis (starting at 0 and stopping
 233 at 2π) generates a uniform sample of ϕ_i .

234 6 Description of the source codes

235 All codes are written in the python3 programming language (except for two Mathematica notebooks). Commented
236 jupyter notebooks can be found on Github at the following address: [https://github.com/laurentjutradsdube/
237 Dual-Regime-Geometry-for-Embryonic-Patterning](https://github.com/laurentjutradsdube/Dual-Regime-Geometry-for-Embryonic-Patterning). This repository also contains folders with the source data
238 files, as well as the source codes used to generate the data files.

239 • **3-gene_det.ipynb**

240 This notebook performs deterministic simulations of the symmetric 3-gene Models 1 and 2, and deterministic
241 simulations of the 3-gene Models 1 and 2 with Hill functions for the weights of the dynamic and static modules.
242 It also performs a bifurcation analysis of these models using the data found in the XPPAUTO_data folder,
243 which also contains the .ode files used to generate the data with the XPP AUTO software [8]. Figure 2, Figure
244 2–figure supplements 1 and 2, and Figure 7–figure supplement 2 show the results obtained with this notebook.

245 • **3-gene_stoch.ipynb**

246 This notebook performs stochastic simulations of the symmetric 3-gene Models 1 and 2, and stochastic
247 simulations of the 3-gene Models 1 and 2 with Hill functions for the weights of the dynamic and static modules.
248 It also generates plots of the mutual information using the data found in the Mutual_info_data folder, which
249 also contains the python codes used to generate the data. Figure 3 and Figure 3–figure supplement 1 show the
250 results obtained with this notebook.

251 • **3-gene_asym.ipynb**

252 This notebook performs deterministic simulations of the asymmetric 3-gene Models 1 and 2. It also performs
253 a bifurcation analysis of these models and generates plot of the mutual information using the data found in the
254 XPPAUTO_data and Mutual_info_data folders, respectively. Figure 5 and Figure 5–figure supplements 1
255 and 2 show the results obtained with this notebook.

256 • **Gene-free_det.ipynb**

257 This notebook performs deterministic simulations of the symmetric gene-free Models 1, 2, 3 and 4. It
258 also performs a bifurcation analysis of these models and generates flow plots using the data found in the
259 XPPAUTO_data and Mathematica_data folders, respectively. Figure 4, Figure 4–figure supplements 1, 2, 3
260 and 4, Figure 4–movie supplements 1, 2, 3 and 4, and Figure 7–figure supplement 2 show the results obtained
261 with this notebook.

262 • **Gene-free_stoch.ipynb**

263 This notebook performs stochastic simulations of the symmetric gene-free Models 1, 2, 3 and 4. It also
264 generates the mutual information plots using the data found in the Mutual_info_data folder. Figure 4–figure
265 supplement 5 shows the results obtained with this notebook.

266 • **Gene-free_asym.ipynb**

267 This notebook performs deterministic simulations of the asymmetric gene-free Models 1 and 2. It also performs

268 a bifurcation analysis of these models using the data found in the XPPAUTO_data folder. Moreover, it generates
269 plots of the flow and of the spatial wave profiles. Figure 6, Figure 6–figure supplement 1, Figure 7 and Figure
270 7–figure supplement 1 show the results obtained with this notebook.

271 • **Hopf_scenario_Fig1.ipynb**

272 This notebook performs deterministic simulations of the gene network model evolved *in silico* in [6]. Results
273 are shown on Figure 1. It also performs a bifurcation analysis of this model, shown on Figure 1–figure
274 supplement 1.

275 • **Infinite-period_scenario_Fig1.ipynb**

276 This notebook performs deterministic simulations of the infinite-period model of Figure 1, which is a simplified
277 version of the model in the appendix of [4].

278 • **Infinite-period_scenario_Fig7.ipynb**

279 This notebook performs deterministic simulations of the infinite-period model of Figure 7, which is adapted
280 from [5].

281 • **Tribolium_model.ipynb**

282 This notebook performs deterministic simulations of the model for Tribolium segmentation from [1]. It also
283 generates flow plots and computes the speed of the cells in phase space. Figure 1–figure supplement 2 shows
284 the results obtained with this notebook.

285 **References**

- 286 [1] Xin Zhu, Heike Rudolf, Lucas Healey, Paul François, Susan J Brown, Martin Klingler, and Ezzat El-Sherif. Speed
287 regulation of genetic cascades allows for evolvability in the body plan specification of insects. *Proceedings of the*
288 *National Academy of Sciences*, 114(41):E8646–E8655, 2017.
- 289 [2] Ezzat El-Sherif, Michalis Averof, and Susan J Brown. A segmentation clock operating in blastoderm and germband
290 stages of tribolium development. *Development*, 139(23):4341–4346, 2012.
- 291 [3] Daniel E Tufcea and Paul François. Critical timing without a timer for embryonic development. *Biophysical*
292 *journal*, 109(8):1724–1734, 2015.
- 293 [4] Isabel Palmeirim, Domingos Henrique, David Ish-Horowicz, and Olivier Pourquié. Avian hairy gene expression
294 identifies a molecular clock linked to vertebrate segmentation and somitogenesis. *Cell*, 91(5):639–648, 1997.
- 295 [5] Luis G Morelli, Saúl Ares, Leah Herrgen, Christian Schröter, Frank Jülicher, and Andrew C Oates. Delayed
296 coupling theory of vertebrate segmentation. *HFSP journal*, 3(1):55–66, 2009.
- 297 [6] Paul François, Vincent Hakim, and Eric D Siggia. Deriving structure from evolution: metazoan segmentation.
298 *Molecular systems biology*, 3(1), 2007.

- 299 [7] Daniel T Gillespie. Approximate accelerated stochastic simulation of chemically reacting systems. *The Journal of*
300 *Chemical Physics*, 115(4):1716–1733, 2001.
- 301 [8] Bard Ermentrout. Xppaut. In *Computational Systems Neurobiology*, pages 519–531. Springer, 2012.