Elucidating essential genes in plant-associated *Pseudomonas protegens* Pf-5 using transposon insertion sequencing

Belinda K Fabian^a, Christie Foster^a, Amy J Asher^a, Liam DH Elbourne^a, Amy K Cain^{a,c}, Karl A Hassan^b, Sasha G Tetu^{a,c#}, Ian T Paulsen^{a,c#}

^a Department of Molecular Sciences, Macquarie University, Sydney Australia

^b School of Environmental and Life Sciences, University of Newcastle, Newcastle, Australia

^c ARC Centre of Excellence in Synthetic Biology, Macquarie University, Sydney Australia

Running title: Essential genes in plant-associated *P. protegens* Pf-5

Address correspondence to Ian Paulsen, ian.paulsen@mq.edu.au or Sasha Tetu, sasha.tetu@mq.edu.au.

Keywords: gene essentiality, genome-wide analysis, *in silico* essential genes, plant growth promoting rhizobacteria, rhizosphere, TraDIS, transposon mutagenesis

Abstract

Gene essentiality studies have been performed on numerous bacterial pathogens, but essential gene sets have been determined for only a few plant-associated bacteria. *Pseudomonas protegens* Pf-5 is a plant-commensal, biocontrol bacteria that can control disease-causing pathogens on a wide range of crops. Work on Pf-5 has mostly focused on secondary metabolism and biocontrol genes, but genome-wide approaches such as high-throughput transposon mutagenesis have not yet been used in this species. Here we generated a dense *P. protegens* Pf-5 transposon mutant library and used transposon-directed insertion site sequencing (TraDIS) to identify 446 genes essential for growth on rich media. Genes required for fundamental cellular machinery were enriched in the essential gene set, while genes related to nutrient biosynthesis, stress responses and transport were under-represented. Comparison of the essential gene sets of Pf-5 and *P. aeruginosa* PA14, an opportunistic human pathogen, provides insight into the biological processes important for their different lifestyles. Key differences include cytochrome c biogenesis, formation of periplasmic disulfide bonds, lipid biosynthesis, ribonuclease activity, lipopolysaccharides and cell surface structures. Comparison of the Pf-5 *in silico* predicted and *in vitro* determined essential gene sets highlighted the essential cellular functions that are over- and underestimated by each method. Expanding essentiality studies into bacteria with a range of lifestyles can improve our understanding of the biological processes important for survival and growth in different environmental niches.

Importance

Essential genes are those crucial for survival or normal growth rates in an organism. Essential gene sets have been identified in numerous bacterial pathogens, but only a few plant-associated bacteria. Employing genome-wide approaches, such as transposon insertion sequencing, allows for the concurrent analysis of all genes of a bacterial species and rapid determination of essential gene sets. We have used transposon insertion sequencing to systematically analyze thousands of *Pseudomonas protegens* Pf-5 genes and gain insights into gene functions and interactions that are not readily available using traditional methods. Comparing Pf-5 essential genes with those of *P. aeruginosa* PA14, an opportunistic human pathogen, provides insight into differences in gene essentiality which may be linked to their different lifestyles.

Introduction

Pseudomonas is a ubiquitous and extremely diverse genus with species occupying a range of niches and lifestyles, spanning from opportunistic human pathogens, such as *P. aeruginosa*, through to plant growth promoting strains, such as *P. protegens* Pf-5 (1). Pseudomonads have a core set of common genes (2), but the mechanisms that underly their niche specializations are not well understood (3). Determining the essential gene complement of related microbes residing in different niches has the potential to shed light on differences in the biochemical functions required for survival and growth in specific environments. As observed in other bacterial taxa, each pseudomonad is expected to have a number of common essential genes together with an additional set of strain-specific essential genes, which may reflect differences in lifestyle (4, 5).

Essential genes are those crucial for survival or normal growth rates in an organism (6, 7). When bacterial genes are manipulated, essential genes are considered to be those for which mutations cannot be made as deletion of these genes will be lethal (8). Experimentally determining which bacterial genes are essential is important for understanding the mechanisms that control bacterial growth, identifying the mechanisms by which microbes specialize for their environmental niches, and can assist with validating computational models of gene essentiality (7, 9, 10). While gene essentiality studies have been performed on numerous bacterial pathogens, essential gene sets have been determined for only a few plant-associated bacteria, including *Herbaspirillum seropedicae* SmR1, a plant growth promoting endophyte, and three nitrogen-fixing root endosymbionts (11, 12).

P. protegens Pf-5 (hereafter referred to as Pf-5) is a plant-commensal, biocontrol bacteria originally isolated from the roots of cotton plants (13). Pf-5 is known to produce a range of secondary metabolites with antibacterial and antifungal activities (14) and can control disease-causing pathogens on a wide range of crops, including cotton, wheat, cucumber and tomatoes (15-18). Analysis of the Pf-5 genome has shed light on many potential functions and molecular systems utilized in its rhizospheric lifestyle (14, 19). Work on specific genes and gene networks has indicated functions for a number of Pf-5 genes, particularly regulatory and metabolic genes involved in the production of secondary metabolites. However, genome-wide approaches such as high-throughput transposon mutagenesis have not yet been used to elucidate gene function or essentiality in this important organism.

Here, we define the essential genome of *P. protegens* Pf-5 using transposon-directed insertion site sequencing (TraDIS). This method combines high-density random transposon

insertion mutagenesis with high-throughput sequencing to concurrently link genotype and phenotype for thousands of genes (20, 21). TraDIS and other transposon mutagenesis techniques have successfully been used to determine the essential gene sets of a wide range of bacteria (22). We also compare the Pf-5 essential gene set with that of *P. aeruginosa* PA14 (hereafter referred to as PA14), an opportunistic human pathogen, providing insight into biological processes critical for their different lifestyles.

Materials and Methods

Transposon mutant library generation and sequencing

Pseudomonas protegens Pf-5 was isolated from soil of a cotton field in Texas, USA (13) and a complete genome sequence has been generated for this organism (19; ENA accession number CP000076). Construction of a *P. protegens* Pf-5 dense Tn5 mutant library was carried out using the technique previously described (20, 21). Briefly, a custom transposome was constructed using EZ-Tn5 transposase (EpiCentre) and a transposon carrying a kanamycin (Km) resistance cassette isolated from the plasmid pUT_Km (23). The custom transposome was electroporated into freshly prepared electrocompetent Pf-5 cells, and the cells were plated on LB-Km agar (16 ug mL⁻¹). Approximately 125,000 colonies were collected from each of four independent batches, combined and stored as glycerol stocks at -80°C. Genomic DNA was isolated from two aliquots of stock containing approximately 2.8 x 10⁹ cells and the transposon insertion sites were sequenced using the methods described previously (21) on an Illumina MiSeq platform to obtain 52 bp single-end genomic DNA reads.

Bioinformatic analysis

The transposon insertion sites were mapped to the Pf-5 genome and statistically analyzed using the Bio-Tradis pipeline (21). A 1 bp mismatch in the transposon tag was allowed and insertions in the extreme 3' end (final 10%) of each gene were discounted as they may not inactivate the gene. Reads with more than one mapping location were mapped to a random matching location to avoid repetitive elements artificially appearing to be essential (0.4% of reads; 21). The pipeline calculates an insertion index value for each gene. This is the number of transposon insertion sites per gene normalized by the length of that gene. A linear regression of the gene insertion indexes of the replicates was completed in R (24). There was a correlation co-efficient of $R^2 = 0.88$ ($p < 2.2 \times 10^{-16}$) between the insertion indexes of the replicates which validates the reproducibility of our replicates and is consistent with the high reproducibility between independent replicates in transposon insertion sequencing studies (25). Essential genes are those with an insertion index lower than the essentiality cut-off value determined by the Bio-Tradis pipeline (26). In this study we required essential genes to have an insertion index lower than the cut-off value in both replicates.

Essential gene analysis

A Cluster of Orthologous Groups (COG) code (27) and Kyoto Encyclopedia of Genes and Genomes (KEGG) Orthology (KO) term (28, 29) for each Pf-5 gene was gathered using eggNOG mapper v1 (30, 31; Dataset S1). We used eggNOG mapper v2 (31, 32) to find this information for 18 essential genes where there were no orthologs identified by eggNOG mapper v1. COG functional category enrichment analysis of the essential gene set compared to the whole gene set was conducted using Fisher's exact test ($p < 0.05$) and corrected for multiple testing using a 5% false-discovery rate (FDR; 33). The sum of all categories does not

equal the total number of genes in the genome as some genes are assigned multiple COG codes. The KEGG mapper tool (34) was used to map the essential genes to KEGG pathways.

The presence and type of signal peptides in essential genes were identified using SignalP v5.0 (Dataset S1; 35) and the presence of transmembrane domains in proteins was determined using TMHMM v2.0 (Dataset S1; 36). As a transmembrane helix close to the N terminus is likely to be a signal peptide, proteins were only classed as membrane proteins if transmembrane helices were detected outside the first 50 residues. Enrichment for genes with signal peptides or transmembrane helices in the essential gene set compared to the whole genome was tested using Fisher's exact test ($p < 0.05$) with Bonferroni correction for testing multiple values.

Pf-5 essential genes were also compared with PA14 essential genes determined by Poulsen and colleagues, which were based on growth in LB media (5). The Poulsen study determined the essential protein-coding genes using two statistical methods: the family-wise error rate (FWER) method, which identified 437 genes as essential, and the false discovery rate (FDR) method, which identified an additional 159 genes as essential (total of 596 essential genes). Orthologs of Pf-5 essential genes in PA14 were determined using Proteinortho v5 (37) and these were used to compare Pf-5 essential genes to the sets from PA14 derived from each statistical method.

In silico prediction of Pf-5 essential genes was conducted using Geptop 2.0 (<http://cefg.uestc.cn/geptop/>) with amino acid sequences and an essentiality score cutoff of 0.24 (38). The predicted Pf-5 essential genes were compared with the *in vitro* determined

Pf-5 essential genes. The gene PFL_0842 was originally annotated as a pseudogene, so it was not assessed by Geptop (marked 'N/A' in Dataset S1).

Data availability

All sequence data generated in this study have been submitted to the EBI European Nucleotide Archive (<https://www.ebi.ac.uk/ena/>) under the project accession number PRJEB39292, within which are the two samples analyzed here ERR4327923 and ERR4327924.

Results and Discussion

Identification of *P. protegens* Pf-5 essential genes

Our dense mutant library of *P. protegens* Pf-5 was generated using random saturation mutagenesis with a Tn5 transposon containing a kanamycin resistance cassette. Over 500,000 transposon mutants were pooled in the construction of the library and the analysis of sequencing data showed there are ~256,000 unique transposon insertion sites in the mutant library (Table 1). Transposon insertions occurred evenly throughout the genome (Figure 1a), with an average of one transposon insertion every ~27 bp and an average of 45 transposon insertion sites in each non-essential protein-coding gene. Multiple insertion sites in a gene are independent evidence that a gene is not essential under the specific conditions used (39, 40). Rarefaction analysis showed that the sequencing reached saturation in terms of the number of unique transposon sites identified in the library (Figure 1c).

Figure 1. *P. protegens* Pf-5 transposon mutant library overview. (A) Distribution of transposon insertion sites across the genome. The outermost circle in grey shows the Pf-5 genome in Mbp. The length of the blue bars in the middle circle represents the number of sequencing reads at each unique transposon insertion location. The inner circle in orange shows the locations of genes characterized as essential with TraDIS. Circular figure created using the R package circlize (41). (B) Correlation of gene insertion indexes for the two replicates of the library. Insertion index is calculated as the number of transposon insertion sites in a gene divided by the gene length. A line through the origin with a slope of 1 is also shown. (C) Rarefaction analysis showing the relationship between sequencing depth and the

number of insertion sites in the transposon mutant library. Analysis conducted using the seq_saturation_test.py script available at https://github.com/francesca-short/tradis_scripts (42) and visualized using the R package ggplot2 (43). (D) A section of the Pf-5 genome containing both essential (orange) and non-essential (grey) genes. The frequency of sequence reads at each transposon insertion site is capped at 1 and genes are annotated with Pf-5 gene names or locus tag numbers. The genes *groL* and *groS* have very few or no insertions and are essential genes. The genes PFL_4840, PFL_4841 and PFL_4842 have a high transposon insertion density and are therefore classed as non-essential genes. Transposon insertion sites are visualized using Artemis (44).

Table 1. *Pseudomonas protegens* Pf-5 transposon mutant library metrics based on Bio-Tradis analysis.

Replicate	No. of reads with matching transposon tag	No. (%) of aligned reads	No. of unique insertion sites	Insertion index for essentiality cut-off
1	1,320,631	1,278,951 (96.8%)	242,827	0.0076
2	1,555,998	1,504,986 (96.7%)	256,020	0.0089

We found 446 out of 6,109 coding sequences (7.3%) were critical for Pf-5 survival and growth on LB agar (Dataset S1). Other pseudomonads that have their essential gene sets characterized also have similar proportions of their genomes determined to be essential genes. For example, *P. aeruginosa* PA14 has 596 out of 5894 protein-coding genes (10%) identified as essential when grown on LB (5) and 6% of the *P. aeruginosa* PAO1 genome was found to be essential for growth on a complex medium (45). The few plant-associated bacteria that have their essential gene sets identified have similar proportions of essential

genes, for example *H. seropedicae* SmR1, a plant endophyte, required 372 out of 4804 protein-coding genes (7.7%) for survival and growth on rich media (12) and 5.6% of the genome of the root endophyte *Rhizobium leguminosarum* bv. *viciae* 3841 was determined to be essential (11). Pf-5 and other plant-associated bacteria often have large genomes with numerous stress response and biosynthetic pathways that are important for survival in their highly variable environmental niches (19, 46, 47). Many of these pathways were likely not critical for Pf-5 survival in this study due to the nutrient-rich and low stress conditions.

Transposon insertion density was slightly higher for genes located near the origin of replication compared to genes located closer to the terminus of replication (Figure 1a). This is regularly observed in genomic and transcriptomic studies due to ongoing DNA replication in the bacterial population. In keeping with this, the density of essential genes was also somewhat lower towards the replication terminus (Figure 1a). Lower numbers of essential genes in terminus regions may also reflect common patterns of genome arrangement; for example, terminus regions of bacterial genomes are often poorly conserved and experience higher rates of lateral gene transfer and rearrangement (48, 49).

The set of genes determined to be essential under the test conditions included PFL_0842, previously annotated as a pseudogene. Analysis showed that this sequence likely represents a functional gene, as it has positional orthologs with the same genomic context (part of an operon with *nusA* and *infB*) in numerous *Pseudomonas* species annotated as *rimP* ribosome maturation factor (50). Based on this, PFL_0842 has been annotated as *rimP* in this study.

Functions of essential genes

A functional overview of the Pf-5 genome was obtained by classifying the genes using Clusters of Orthologous Groups (COG) categories (27). 87.3% of Pf-5 coding genes were able to be assigned a COG code. Several functional categories were significantly over-represented in the set of essential genes relative to the overall genome, most notably translation (J); cell cycle control, cell division and chromosome partitioning (D); coenzyme transport and metabolism (H); nucleotide transport and metabolism (F) as well as intracellular trafficking, secretion, and vesicular transport (U); cell wall/membrane/envelope biogenesis (M); and replication, recombination and repair (L; Figure 2). Other functional categories were significantly under-represented in the set of essential genes including secondary metabolite biosynthesis, transport and catabolism (Q); amino acid transport and metabolism (E); inorganic ion transport and metabolism (P); transcription (K); and signal transduction mechanisms (T; Figure 2). Genes with unknown function (S) make up a significantly lower proportion of the essential gene set (10.6%) compared to the whole Pf-5 genome (22%; Figure 2).

Figure 2. Cluster of Orthologous Groups (COG) functional enrichment analysis of

P. protegens Pf-5 essential genes compared to the whole genome. COG category

enrichment was tested using Fisher's exact test and corrected for multiple testing using a

5% false-discovery rate; * indicates significant enrichment or depletion with $p < 0.05$; six

COG categories are not included in the figure: categories A and B contain 2 and 5 non-

essential genes, respectively, and categories R, W, Y and Z contain no genes.

The pattern of enrichment in essential genes belonging to COG categories for fundamental

cellular machinery, such as translation and ribosome structure, coenzyme transport and

metabolism and cell envelope biogenesis, reflects the fact that these functions are critical

for survival and growth. In contrast, genes belonging to the COG category of transcription

were notably under-represented in the essential gene set, presumably as transcriptional

regulatory genes are typically required under changing environmental conditions (51), and

therefore were not essential during growth in stable laboratory conditions with rich media.

248

249 At the metabolic pathway level, most of the key pathways were essential in Pf-5, including
250 the TCA cycle, fatty acid biosynthesis, gluconeogenesis, peptidoglycan biosynthesis and
251 heme biosynthesis, with all or most of the genes within each of these pathways observed
252 within the essential gene set. Genes required for glycolysis, however, were not essential in
253 this study, presumably as the mutant library was grown on LB media which contains high
254 concentrations of peptides and amino acids which feed into other metabolic pathways.
255 Similarly, under-representation of genes required for amino acid and inorganic ion transport
256 and metabolism was observed, likely due to these biosynthetic pathways not being required
257 for growth on rich media; this has also been observed in other gene essentiality studies
258 across a wide range of bacterial taxa (52). Secondary metabolite biosynthesis, transport and
259 catabolism genes were also under-represented in the essential gene set, potentially due to
260 the limitation of TraDIS methodology in the detection of genes related to public goods (53).

261

262 Although Pf-5 encodes 780 transporter proteins (54), only four transport systems were
263 essential: the LptBFG, MsbA, LolCDE, and CcmABC complexes. The first three of these
264 transport systems are involved in the biogenesis of the outer membrane, while CcmABC is a
265 heme chaperone-release system required for the biogenesis of cytochrome c (55, 56). The
266 non-essentiality of other transporters was likely due to functional redundancy, the presence
267 of gene duplicates, or that they were superfluous for growth in rich media.

268

269 Among the set of essential genes, a number remain annotated as hypothetical genes (35
270 genes or 7% of the essential gene set). The essentiality of these hypothetical genes indicates
271 they are associated with as yet unknown functions that are critical for Pf-5 survival and

growth on rich media. Given this, characterization of these genes would be of particular interest and may provide further useful insights into fundamental biological processes in this plant-associated bacteria.

Essential genes associated with mobile genetic elements

The Pf-5 genome contains six prophage regions (Prophage 01-06) and two large genomic islands (PFGI-1 and -2), together comprising a total of 226 genes (57). Ten of the essential Pf-5 genes reside within these genome regions (Dataset S1). This includes PFL_4679, encoding type IV pilus biogenesis protein PilR which is located within ICE-type genomic island PFGI-1. This gene is part of a *pil* cluster, orthologs of which in PA14 are also located within a genomic island and have been found to be involved with conjugative transfer of the excised island to recipients (58, 59), but were not found to be essential in PA14 Tn-Seq experiments using rich medium (5). As *pilR* was the sole essential gene from this cluster in Pf-5, it may be that mutants of this gene were not viable due to accumulation of a toxic intermediate product such as non-polymerized pilin monomers, rather than due to impaired pilus formation capacity, which would presumably select against mutation of genes for the other pilus components.

The F-pyocin-like Prophage 01 contains the essential gene PFL_1210 which encodes the transcriptional regulator PrtR (57), a Cro/C1-like repressor of pyocin production, which has been previously found to be essential for *P. aeruginosa* PAO1 in rich media (45, 60). Essentiality of the *prtR* gene was presumably important for control of production of these polypeptide toxins to prevent self-lysis. Other prophage regions contain additional regulators observed to be essential, including two more Cro/C1-type transcriptional

repressors (PFL_2126 in Prophage 04 and PFL_3780 in Prophage 06) and a peptidase S24-like protein/LexA-like repressor (PFL_1986 in Prophage 03; 57), which may similarly be required to prevent production of toxins or phage lytic conversion.

Of the remaining five MGE-associated essential genes, two have only predicted functions: the putative ATP-dependent nuclease PFL_1842 in Prophage 02 and the putative nuclease PFL_4984 within PFGI-2 (57); the remainder are conserved hypothetical proteins and potentially important targets for future characterization efforts. The presence of multiple essential genes within these MGE regions shows the capacity for such horizontally inherited material to acquire critical functions within bacterial cells, either in aiding stability of the MGE within the genome (for example, toxin/antitoxin systems) or contributing other conditionally essential functions.

Protein subcellular localization of essential genes

Enrichment analysis for genes containing signal peptide sequences showed that genes that encode proteins containing lipoprotein and general secretion (sec) signal peptides were significantly under-represented in the essential gene set when compared to the Pf-5 genome as a whole (Table 2). Similarly, genes encoding proteins with transmembrane helices were significantly under-represented in the Pf-5 essential gene set (Table 2). This under-representation in the essential gene set is consistent with the subcellular localization of essential genes of 20 other gram-negative bacteria (61). As transmembrane proteins and those containing signal peptides are secreted outside cells or anchored in cell membranes they are often involved in interactions with other microorganisms and hosts (62) and therefore may not have been required under axenic conditions.

Table 2. Bioinformatic prediction of genes with signal peptide and transmembrane helices in the Pf-5 essential gene set and the whole Pf-5 genome.

Protein subcellular localization	Number in essential gene set	Proportion of essential genes	Number in whole genome	Proportion of all genes
SP	19	4.3% [†]	673	11.0%
TAT	3	0.7%	55	0.9%
LIPO	8	1.8% [†]	260	4.3%
Total signal peptides	30	6.7%	988	16.2%
Transmembrane proteins [^]	40	9.0% [†]	1089	17.8%

SP = Sec signal peptide; TAT = Tat signal peptide; LIPO = Lipoprotein signal peptide

Enrichment of these genes in essential gene set was tested using Fisher's exact test with Bonferroni correction for testing multiple values.

[†] indicates significant underrepresentation in the essential gene set with $p < 0.05$

[^] one or more transmembrane helices detected outside the first 50 residues

Comparison of essential genes in *P. protegens* Pf-5 and *P. aeruginosa* PA14

Based on our analysis, there were 446 essential protein-coding genes in Pf-5 when grown on LB media. A similar Tn-Seq study in PA14 has recently investigated essential protein-coding genes for this opportunistic human pathogen on LB media (5). This paper utilized two statistical methods to identify essential genes, the first based on the family-wise error rate (FWER) method which identified 437 genes as essential and the second using a false discovery rate (FDR) method which identified 596 genes as essential. We compared our set of Pf-5 essential genes with both sets from Poulsen et al. (5) and found the FDR set more closely aligned with our essential gene calls (Dataset S1). The FDR set also included genes

expected to be essential, such as F_0F_1 ATPase, cell division and ribosomal protein genes, which were not identified by the family-wise error rate analysis (5).

Using the FDR set from Poulsen et al. (5) we investigated the similarities and differences between the essential gene sets of Pf-5 and PA14, finding that most (80%) of Pf-5 essential genes overlap with those of PA14, while each species also had a number of unique essential genes (Figure 3). The majority of the 357 overlapping essential genes relate to basic cellular functions, such as translation, cell envelope biogenesis, co-enzyme transport and metabolism, energy production, replication and recombination, and lipid transport and metabolism.

Figure 3. Comparison of the essential protein-coding genes of *P. protegens* Pf-5 with the essential genes of *P. aeruginosa* PA14 determined using a false discovery rate adjustment method (5). The PA14 orthologs of seven Pf-5 essential genes did not have their essentiality assessed by Poulsen et al. (5). Venn diagram created using Multiple List Comparator (www.molbiotools.com/listcompare.html).

Genes essential only in Pf-5

There were 38 genes within the Pf-5 essential set which were not flagged as essential in PA14 (Figure 3). This set included genes related to energy generation (cytochrome *c* biogenesis), formation of periplasmic disulfide bonds, three ribosomal proteins, amino acid biosynthesis, cell surface structures and hypothetical proteins.

Two-thirds of the genes in the cytochrome *c* maturation pathway are essential in Pf-5 (*ccmACEF*), but this pattern was not observed in PA14, where only one of these was in the essential set (*ccmB*). The *ccmABCDEFG* genes encode proteins that carry out post-translational modifications on *c*-type cytochromes to facilitate their binding to heme (63). The essentiality of this set of genes in Pf-5 is consistent with the importance of generating cytochrome *c* to facilitate heme production.

Two of the genes essential only in Pf-5 encode proteins involved in the formation of disulfide bonds (*dsbD* and *trxB_1*). DsbD is a thiol:disulfide interchange protein which is the sole provider of periplasmic reducing power and has a role in cytochrome *c* maturation (64). In *Escherichia coli* DsbD is important, but not essential for cytochrome *c* maturation as developing cytochromes can use either the DsbD-independent or dependent pathways (64). The PA14 *dsbD* ortholog *dipZ2* and its paralog *dipZ* were both determined to be non-essential when grown on rich media (5). The essentiality of *dsbD* in Pf-5, but not in PA14, may reflect a different balance in the proportion of maturing *c*-type cytochromes that pass through each pathway. The Pf-5 gene *trxB_1* that encodes thioredoxin reductase was essential, but its paralog *trxB_2* was non-essential. In PA14 there are also two paralogs, *trxB_1* and *trxB_2*, but both of these genes were non-essential (5). *Trx* genes in bacteria

supply reducing power to DsbD and play a role in stress responses, such as oxidative stress in *Pseudomonas syringae* pv. *tomato* (65). The difference in essentiality of the two Pf-5 *trxB* paralogs suggests that TrxB_1 may be the main provider of reducing power to DsbD and that TrxB_2 may be required under stress or other conditions.

Three ribosomal protein genes *rpmE_2*, *rpmF* and *rpmI* were essential in Pf-5, but not in PA14. In Pf-5 *rpmE_2* codes for a C+ form of the 50S ribosomal protein L31 which is stable when bound to a zinc ion; in contrast, *rpmE_1* encodes a C- form of the protein which lacks metal chelating capacity (66). The essentiality of *rpmE_2* in Pf-5 reflects the zinc replete conditions in this study, while *rpmE_1* has been found to be conditionally essential for Pf-5 in zinc limited conditions (66). In PA14, both of the genes annotated as *rpmE* were non-essential when grown on LB (PA14_66710 and PA14_17700; 5). The genes that encode 50S ribosomal proteins L32 (*rpmF*) and L35 (*rpmI*), whilst not essential in PA14 (5), have been found to be essential in a number of other bacteria. For example, *rpmF* was essential in *Burkholderia thailandensis* E264 (67) and both genes were essential in *Acinetobacter baylyi* ADP1 (68).

Genes encoding three proteins responsible for the formation of homoserine and its conversion to threonine (*thrBC* and *hom*) were essential in Pf-5, but not in PA14. These amino acids are important intermediates in amino acid biosynthesis and are precursors for the formation of methionine, serine, glycine and cysteine (69). These results suggest that PA14 was better able to take up threonine from the media in these conditions than Pf-5.

The gene *eda* which encodes 2-dehydro-3-deoxyphosphogluconate aldolase/4-hydroxy-2-oxoglutarate aldolase was essential in Pf-5 but its PA14 ortholog PA14_23090 was non-essential. *Eda* performs a key role in the catabolism of glucose and conversion into pyruvate through the Entner-Doudoroff pathway (70). This difference in essentiality is likely due to redundancy for this function in PA14; an *eda* paralog *kdgA* (PA14_23620) was also non-essential.

Three genes related to lipopolysaccharide formation (*kdsC*, *PFL_0526*, *PFL_0563*) were essential only in Pf-5. Similarly, in PA14 there are a number of essential genes that encode proteins important for lipopolysaccharide formation and other cell surface features that were not essential in Pf-5 (*htrB*, *omlA*, *wbpLV*, *rfbACD*, *pstB*, *wecB*). This difference in essential genes related to the cell membrane and cell surface likely reflects species-specific differences in many of these components, such as O-antigen, the polysaccharide component of lipopolysaccharide that extends from the surface of Gram-negative cells (71).

There were also differences in the essentiality of some conserved hypothetical proteins that are homologous in Pf-5 and PA14. The proteins encoded by PFL_1792 and PFL_2999 were essential in Pf-5, whereas their orthologs PA14_25620 and PA14_01220, respectively, were non-essential in PA14. Likewise, there were 16 essential conserved hypothetical proteins in PA14 but their orthologs were non-essential in Pf-5 (Dataset S1).

Pf-5 essential genes with no PA14 orthologs

There were 43 genes essential in Pf-5 that do not have homologs in PA14 (Figure 3), including genes related to cell membrane and surface structures, pyoluteorin biosynthesis, a

TonB complex, and 25 conserved hypothetical proteins. The unique nature of the seven genes related to the cell surface and cell membrane (PFL_2356, PFL_5099-5103 and PFL_5030) is consistent with the species specificity of these components. The essential gene *p/tL* from the pyoluteorin biosynthesis gene cluster has no PA14 homolog as this cluster is not present in PA14.

Genes that encode TonB complexes in both species were essential, but the genes that form some of the TonB systems are unique to Pf-5 and PA14. In Pf-5 the TonB1 complex (*tonB1*, *exbB1* and *exbD1*) was essential and there is no homologous TonB complex in PA14. In PA14 *tonB* (no Pf-5 homolog) and *exbD2* (homolog to PFL_2822) were essential. Many bacterial species only have one TonB system, but some species have multiple TonB systems with different functional specificities, for example *Vibrio cholerae* CA401S (72) and *P. aeruginosa* PAO1 (73). Both Pf-5 and PA14 possess multiple TonB complexes; Pf-5 has six annotated TonB complexes, while PA14 has two. This difference in essentiality of TonB systems of Pf-5 and PA14 is consistent with the differing functionalities of TonB systems observed in other species.

Genes essential only in PA14

There are 240 PA14 essential genes unique to PA14; only 141 of these have Pf-5 orthologs (Figure 3). This includes genes that encode proteins involved in four biosynthetic pathways (aromatic amino acids, biotin, lysine and lipids), cell division, homologous recombination, ribonuclease activity and 16 conserved hypothetical proteins.

Six genes in the aromatic amino acid biosynthetic pathway were essential in PA14, whereas all genes in this pathway were non-essential in Pf-5. This pathway, known as the shikimate pathway, produces chorismate which is the last common precursor of the aromatic amino acids tryptophan, tyrosine and phenylalanine as well as vitamins E and K and some siderophores (74, 75). The essentiality of *aroABCEK* and *pheA* show that chorismate is an important branch-point metabolite for PA14 survival and growth. In contrast, the non-essentiality of these genes in Pf-5 suggests that this pathway was not required under these experimental conditions and Pf-5 obtained aromatic amino acids from the media.

Genes encoding five proteins in the biotin biosynthesis pathway (*bioABCDF*) and its transcriptional repressor (*birA*) were essential in PA14, but non-essential in Pf-5. Biotin, also known as vitamin H, is a cofactor for enzymes involved in central metabolism carboxylation reactions (76). These results suggest that Pf-5 was better able to take up biotin from the media than PA14.

Four genes that encode proteins involved in cell division were essential in PA14, but non-essential in Pf-5 (*ftsK*, *minCD*, *mreD*, *PFL_0438*). Cell division proteins are essential for the normal replication and viability of bacterial cells; for example, *minCD* encodes proteins that ensure that cell division occurs in the middle of the cell not at a polar site (77) and *mreD* encodes a shape protein which is involved in the rod shape of cells. When *mreD* is knocked out cells take on a spherical form (78). Other rod-shaped bacteria, such as *A. baylyi* ADP1, have been able to dispense with these genes under lab conditions in essentiality studies (68) but *ftsK* and *mreD* have been reported to be essential in other studies in *P. aeruginosa* (45). The non-essentiality of these genes in Pf-5 suggests that cells without these proteins

survived in the short timeframe of this experiment, albeit presumably with abnormal morphologies.

Five genes that code for lipid metabolism proteins were essential in PA14, but not Pf-5. The genes *fadA* and *fadB* encode the two peptides that form the fatty acid oxidation complex. This complex is part of the β -oxidation cycle which is responsible for the degradation of long chain fatty acids into acetyl-coenzyme A (79). In Pf-5 there is a single copy of *fadA*, but multiple copies of *fadB* (*fadB*, *fadB1x* and *fadB2x*). In *P. aeruginosa* PAO1 expression of each of the *fadAB* operon homologues is induced by the presence of different fatty acids (80, 81). There are also three acyl-CoA dehydrogenase family proteins (PFL_0245, PFL_2615 and PFL_5687) that were essential in PA14, but non-essential in Pf-5. The difference in the essentiality of these five genes in PA14 and Pf-5 suggests that the two species have different availabilities of fatty acids and maintain different balances in their lipid metabolism to achieve membrane homeostasis.

Genes *recBCD* and *ruvA* that encode homologous recombination enzymes were essential in PA14 but non-essential in Pf-5. A relatively small number of bacterial essential gene studies have identified these genes as essential (22), despite the importance of Rec-mediated repair of double stranded DNA breaks. The non-essentiality of *recBCD* and *ruvA* suggests this function is not essential for Pf-5 under laboratory conditions and timescales.

Genes that encode four ribonucleases were essential in PA14, but non-essential in Pf-5 (*rnc*, *rne*, *rnt* and PFL_3322). Ribonucleases (or RNases) have dual functions: they are involved in both the maturation and degradation of rRNAs, tRNAs, sRNAs and mRNAs (82). In *E. coli*

there is evidence to suggest these ribonucleases may have overlapping functionalities (83), so these may have been non-essential in Pf-5 due to redundancy.

The genes *dapA_1* and *dapF_2* encoding proteins in the lysine biosynthetic pathway (84) were in the essential set in PA14 but not Pf-5. This difference in essentiality may occur as PA14 has a single copy of each gene, whereas Pf-5 has redundancy for these functions (two copies of each gene). A similar pattern is observed with the gene *fabF* which codes for beta-ketoacyl-acyl-carrier-protein synthase II, a part of the fatty acid biosynthesis pathway and a vital enzyme in the biogenesis of phospholipid membranes (85). The two copies of this gene in Pf-5, *fabF_1* and *fabF_2*, were non-essential. Of the two PA14 copies of this gene *fabF1* was essential and *fabF2* was non-essential (5).

Comparison of *in vitro* determined and *in silico* predicted essential genes

In parallel with the development of transposon library-based approaches for determining gene essentiality, computational tools have been developed to predict essential gene sets based on information such as gene orthology and phylogeny (38). Here we compared the set of essential genes identified *in vitro* by TraDIS with computational predictions by Geptop 2.0 (38). Geptop 2.0 identified 406 protein-coding genes as essential for growth and survival of Pf-5 (Dataset S1). When compared with the 446 essential protein-coding genes identified by TraDIS on rich media there were 308 genes in common, 138 genes only identified by TraDIS, and 98 genes that were only in the computationally predicted set (Figure 4). The 308 genes identified as essential by both methodologies include core cellular functions such as the processing of information (replication, translation and transcription), energy production, cell division and maintenance of the cell envelope.

Figure 4. Comparison of *P. protegens* Pf-5 essential protein-coding genes identified *in vitro* by TraDIS and the essential genes predicted *in silico* by Geptop 2.0 (38). Venn diagram created using Multiple List Comparator (www.molbiotools.com/listcompare.html).

Out of the 138 essential genes identified only by TraDIS, 14 are related to lipopolysaccharide biosynthesis and cell surface structures, including *rfaCF*, *waaGP* and a range of lipoprotein, O-antigen and glycosyltransferase genes. The absence of these genes from the predicted essential gene set is consistent with the species-specific nature of these pathways and the use of evolutionary conservation in the Geptop computational approach (38). Thirty-two essential genes identified only by TraDIS are annotated as hypothetical proteins. As Geptop uses orthology and phylogeny to predict essential genes it is unsurprising that hypothetical genes were not included in the predicted essential gene set.

There are a number of genes our work indicates are essential that have not been identified by Geptop. For example, *fbp* encoding fructose-1,6-bisphosphatase, cell division genes (*zipA*, *ftsB* and *minE*), *fis* encoding the DNA binding protein Fis, cytochrome c biogenesis (*ccmABCEF*), genes encoding proteins involved in the maintenance of membrane stability (*pal* and *tolABQR*), sulfur relay genes (*tusABCDE*) and iron-sulfur cluster genes (*hscAB*, *iscA* and ferredoxin genes *fdx* and PFL_5869). These genes have been found to be essential in

other organisms such as *E. coli*, *Haemophilus influenzae* and *Shewanella oneidensis* (86-88). This suggests there may be functional categories of genes that are systematically missed by Geptop.

Ninety-eight Pf-5 protein-coding genes were identified as essential by Geptop but not found to be essential *in vitro* by TraDIS. There are three major reasons these genes were not identified as essential in the *in vitro* data. Firstly, there is some redundancy in the genome for essential functions. For example, the genes *ddlAB* encode D-alanine--D-alanine ligases which condense two molecules of D-alanine, an essential step in peptidoglycan biosynthesis (89, 90), yet neither gene was essential *in vitro*, presumably as loss of function of each single gene can be tolerated. Similarly, *map_1* and *map_2* both encode methionine aminopeptidases which perform the essential function of cleaving methionine residues from the N-terminal of nascent proteins (91), but loss of either of these genes did not preclude growth *in vitro*. Many of the genes encoding subunits of NADH-quinone oxidoreductase were non-essential in Pf-5 *in vitro* (*nuoBDGHIJKLMN*) but were predicted to be essential by Geptop. In addition to the *nuo* complex, which encodes a type I NADH dehydrogenase, there are two other NADH dehydrogenases encoded in the Pf-5 genome (92). These three NADH dehydrogenases provide Pf-5 with respiratory flexibility, such as in *P. aeruginosa* where two NADH dehydrogenases were redundant in aerobic conditions (93). As TraDIS library construction generates single knockout mutants there is limited capacity to identify essential genes where there is functional redundancy in the genome, which is a recognized limitation of this technique (94).

Secondly, some genes encode products with essential functions in bacterial cells, but these functions are only performed under certain conditions. For example, the highly conserved Clp proteolytic system degrades misfolded and abnormal proteins which accumulate in response to environmental stresses (95). This essential function is reflected in the Geptop prediction of *clpP* and *clpX* as essential, but these genes were not identified as essential by TraDIS, presumably as environmental stresses were low under our library growth conditions. This pattern is also observed with *dnaJ* which encodes a molecular chaperone which helps to ensure the correct folding of proteins, particularly under heat shock (96, 97), and the highly conserved gene *polA* encoding DNA Polymerase I which repairs damaged DNA during replication (98). These proteins perform important functions but were not essential under the stable conditions and short duration of this study. The media and growth conditions under which transposon libraries are created also influences the essentiality of genes (7). The rich media used in our *in vitro* experiments may have resulted in some genes that were predicted to be essential *in silico* not being identified as essential by TraDIS. For example, three amino acid biosynthetic genes (*aroBCK*) were predicted to be essential by Geptop, but were non-essential *in vitro*, likely due to the ability of Pf-5 to acquire amino acids from the media.

Lastly, the experimental timeframe is likely also a factor in some genes being identified by TraDIS as non-essential. These genes may have important roles at certain stages of cell or population growth and therefore are identified as essential based on the phylogenetic approach used by Geptop. For example, the genes *ftsHXX* and *parAB* perform important roles in cell division (99) and were predicted to be essential by Geptop, but they were determined to be non-essential by TraDIS. This indicates cells with these individual gene

disruptions may undergo abnormal cell division, potentially resulting in aberrant morphologies, but the cells were not lost completely from the population. It is likely that such altered morphologies may become problematic in a longer-term study.

Conclusion

In this study we created a saturated transposon mutant library and used TraDIS to successfully identify 446 genes that were essential for *P. protegens* Pf-5, a plant-associated bacteria, to survive and grow on rich media. The essential gene set showed enrichment of genes required for fundamental cellular machinery, which is consistent with the composition of essential gene sets in other bacteria. Genes related to nutrient biosynthesis, stress responses and transport were under-represented, potentially due to the specific growth conditions used in this study as well as functional redundancy within the genome.

We identified key differences between the essential gene sets of the plant-associated pseudomonad Pf-5 and the well-studied opportunistic pathogen PA14. These include genes related to energy generation (cytochrome *c* biogenesis), formation of periplasmic disulfide bonds, lipid biosynthesis, ribonuclease activity, lipopolysaccharides and cell surface structures. This information highlights differences in the processes required for survival and growth of pseudomonads that occupy different environmental niches.

Our comparison of the essential gene sets determined *in silico* and via the *in vitro* TraDIS approach shows that the prediction of essential genes by Geptop on the basis of conservation through evolutionary time overestimates the essentiality of some cellular functions and underestimates others. Despite this, there is still substantial overlap in the

genes identified as essential by these two methods. While both techniques have recognized limitations, the information from TraDIS studies could be used to evaluate and improve *in silico* predictive models for essential genes.

Using TraDIS to systematically analyze thousands of genes provides insights into gene functions and interactions that are not readily available using traditional methods. The Pf-5 transposon mutant library will enable high-throughput studies in a range of growth conditions, such as competition with soil microbes or stress tolerance. Expanding essentiality studies beyond bacterial pathogens improves our understanding of the biological processes important for survival and growth in different environmental niches.

Acknowledgements

The TraDIS sequencing for this study was conducted at The Ramaciotti Centre for Genomics at the University of New South Wales. *Pseudomonas protegens* Pf-5 was obtained from Joyce Loper (Oregon State University). The authors thank Dr Francesca Short (Macquarie University) for advice on TraDIS data analysis and Professor Joyce Loper (Oregon State University) and Dr Virginia Stockwell (USDA) for helpful discussions. This work is supported by an Australian Research Council Discovery Grant (#DP160103746) to IP, ST and KH. IP is supported by an Australian Research Council Laureate Fellowship (#FL140100021), ST is supported by an Australian Research Council Discovery Early Career Researcher Award (#DE150100009) and BF is the recipient of an Australian Government Research Training Pathway Scholarship.

References

1. Peix A, Ramírez-Bahena MH, Velázquez E. 2018. The current status on the taxonomy of *Pseudomonas* revisited: An update. *Infect, Genet Evol* 57:106-116.
2. Hesse C, Schulz F, Bull CT, Shaffer BT, Yan Q, Shapiro N, Hassan KA, Varghese N, Elbourne LDH, Paulsen IT, Kyrpides N, Woyke T, Loper JE. 2018. Genome-based evolutionary history of *Pseudomonas* spp. *Environ Microbiol* 20:2142-2159.
3. Silby MW, Winstanley C, Godfrey SAC, Levy SB, Jackson RW. 2011. *Pseudomonas* genomes: Diverse and adaptable. *FEMS Microbiol Rev* 35:652-680.
4. Danchin A, Fang G. 2016. Unknown unknowns: essential genes in quest for function. *Microb Biotechnol* 9:530-540.
5. Poulsen BE, Yang R, Clatworthy AE, White T, Osmulski SJ, Li L, Penaranda C, Lander ES, Shoshitaishvili N, Hung DT. 2019. Defining the core essential genome of *Pseudomonas aeruginosa*. *Proc Natl Acad Sci USA* 116:10072-10080.
6. Goodall E, Robinson A, Johnston I, Jabbari S, Turner K, Cunningham A, Lund P, Cole J, Henderson I. 2018. The essential genome of *Escherichia coli* K-12. *mBio* 9:e02096-17.
7. Rancati G, Moffat J, Typas A, Pavelka N. 2018. Emerging and evolving concepts in gene essentiality. *Nat Rev Genet* 19:34-49.
8. Bailey J, Cass J, Gasper J, Ngo ND, Wiggins P, Manoil C. 2019. Essential gene deletions producing gigantic bacteria. *PLoS Genet* 15:e1008195.
9. Joyce AR, Reed JL, White A, Edwards R, Osterman A, Baba T, Mori H, Lesely SA, Palsson B, Agarwalla S. 2006. Experimental and computational assessment of conditionally essential genes in *Escherichia coli*. *J Bacteriol* 188:8259-8271.
10. Peters JM, Colavin A, Shi H, Czarny TL, Larson MH, Wong S, Hawkins JS, Lu CHS, Koo BM, Marta E, Shiver AL, Whitehead EH, Weissman JS, Brown ED, Qi LS, Huang KC, Gross CA. 2016. A comprehensive, CRISPR-based functional analysis of essential genes in bacteria. *Cell* 165:1493-1506.
11. Perry BJ, Yost CK. 2014. Construction of a *mariner*-based transposon vector for use in insertion sequence mutagenesis in selected members of the *Rhizobiaceae*. *BMC Microbiol* 14:298.
12. Rosconi F, de Vries SPW, Baig A, Fabiano E, Grant AJ. 2016. Essential Genes for *In Vitro* Growth of the Endophyte *Herbaspirillum seropedicae* SmR1 as Revealed by Transposon Insertion Site Sequencing. *Appl Environ Microbiol* 82:6664-6671.

13. Howell CR, Stipanovic RD. 1979. Control of *Rhizoctonia solani* on cotton seedlings with *Pseudomonas fluorescens* and with an antibiotic produced by the bacterium. *Phytopathology* 69:480-482.
14. Loper JE, Kobayashi DY, Paulsen IT. 2007. The Genomic Sequence of *Pseudomonas fluorescens* Pf-5: Insights Into Biological Control. *Phytopathology* 97:233-238.
15. Howell CR, Stipanovic RD. 1980. Suppression of *Pythium ultimum*-Induced Damping-Off of Cotton Seedlings by *Pseudomonas fluorescens* and its Antibiotic, Pyoluteorin. *Phytopathology* 70:712-715.
16. Kraus J, Loper JE. 1992. Lack of evidence for a role of antifungal metabolite production by *Pseudomonas fluorescens* Pf-5 in biological-control of *Pythium* damping-off of cucumber. *Phytopathology* 82:264-271.
17. Pfender WF, Kraus J, Loper JE. 1993. A genomic region from *Pseudomonas fluorescens* Pf-5 required for pyrrolnitrin production and inhibition of *Pyrenophora tritici-repentis* in wheat straw. *Phytopathology* 83:1223-1228.
18. Saravanan S, Muthumanickam P, Saravanan TS, Santhaguru K. 2013. Antagonistic potential of fluorescent *Pseudomonas* and its impact on growth of tomato challenged with phtopathogens. *Afr Crop Sci J* 21:29-36.
19. Paulsen IT, Press CM, Ravel J, Kobayashi DY, Myers GSA, Mavrodi DV, DeBoy RT, Seshadri R, Ren Q, Madupu R, Dodson RJ, Durkin AS, Brinkac LM, Daugherty SC, Sullivan SA, Rosovitz MJ, Gwinn ML, Zhou L, Schneider DJ, Cartinhour SW, Nelson WC, Weidman J, Watkins K, Tran K, Khouri H, Pierson EA, Pierson LS, Thomashow LS, Loper JE. 2005. Complete genome sequence of the plant commensal *Pseudomonas fluorescens* Pf-5. *Nat Biotechnol* 23:873-878.
20. Langridge GC, Phan MD, Turner DJ, Perkins TT, Parts L, Haase J, Charles I, Maskell DJ, Peters SE, Dougan G, Wain J, Parkhill J, Turner AK. 2009. Simultaneous assay of every *Salmonella* Typhi gene using one million transposon mutants. *Genome Res* 19:2308-2316.
21. Barquist L, Mayho M, Cummins C, Cain AK, Boinett CJ, Page AJ, Langridge GC, Quail MA, Keane JA, Parkhill J. 2016. The TraDIS toolkit: Sequencing and analysis for dense transposon mutant libraries. *Bioinformatics* 32:1109-1111.

22. Luo H, Lin Y, Gao F, Zhang C-T, Zhang R. 2014. DEG 10, an update of the database of essential genes that includes both protein-coding genes and noncoding genomic elements. *Nucleic Acids Res* 42:D574-D580.
23. Herrero M, De Lorenzo V, Timmis KN. 1990. Transposon Vectors Containing Non-Antibiotic Resistance Selection Markers for Cloning and Stable Chromosomal Insertion of Foreign Genes in Gram-Negative Bacteria. *J Bacteriol* 172:6557-6567.
24. R Core Team. 2018. R: A language and environment for statistical computing, R Foundation for Statistical Computing, Vienna, Austria, <https://www.R-project.org>.
25. Barquist L, Boinett CJ, Cain AK. 2013. Approaches to querying bacterial genomes with transposon-insertion sequencing. *RNA Biol* 10:1161-1169.
26. Barquist L, Langridge GC, Turner DJ, Phan MD, Turner AK, Bateman A, Parkhill J, Wain J, Gardner PP. 2013. A comparison of dense transposon insertion libraries in the *Salmonella* serovars Typhi and Typhimurium. *Nucleic Acids Res* 41:4549-4564.
27. Tatusov RL, Galperin MY, Natale DA, Koonin EV. 2000. The COG database: a tool for genome-scale analysis of protein functions and evolution. *Nucleic Acids Res* 28:33-36.
28. Kanehisa M, Goto S. 2000. KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids Res* 28:27-30.
29. Kanehisa M, Sato Y, Furumichi M, Morishima K, Tanabe M. 2019. New approach for understanding genome variations in KEGG. *Nucleic Acids Res* 47:D590-D595.
30. Huerta-Cepas J, Szklarczyk D, Forslund K, Cook H, Heller D, Walter MC, Rattei T, Mende DR, Sunagawa S, Kuhn M, Jensen LJ, Von Mering C, Bork P. 2016. eggNOG 4.5: A hierarchical orthology framework with improved functional annotations for eukaryotic, prokaryotic and viral sequences. *Nucleic Acids Res* 44:D286-D293.
31. Huerta-Cepas J, Forslund K, Coelho LP, Szklarczyk D, Jensen LJ, Von Mering C, Bork P. 2017. Fast genome-wide functional annotation through orthology assignment by eggNOG-mapper. *Mol Biol Evol* 34:2115-2122.
32. Huerta-Cepas J, Szklarczyk D, Heller D, Hernández-Plaza A, Forslund SK, Cook H, Mende DR, Letunic I, Rattei T, Jensen LJ, Von Mering C, Bork P. 2019. eggNOG 5.0: A hierarchical, functionally and phylogenetically annotated orthology resource based on 5090 organisms and 2502 viruses. *Nucleic Acids Res* 47:D309-D314.

33. Benjamini Y, Hochberg Y. 1995. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *J Roy Stat Soc Ser B (Stat Method)* 57:289-300.
34. Kanehisa M, Sato Y. 2020. KEGG Mapper for inferring cellular functions from protein sequences. *Protein Sci* 29:28-35.
35. Almagro Armenteros JJ, Tsirigos KD, Sønderby CK, Petersen TN, Winther O, Brunak S, von Heijne G, Nielsen H. 2019. SignalP 5.0 improves signal peptide predictions using deep neural networks. *Nat Biotechnol* 37:420-423.
36. Krogh A, Larsson B, von Heijne G, Sonnhammer ELL. 2001. Predicting Transmembrane Protein Topology with a Hidden Markov Model: Application to Complete Genomes. *J Mol Biol* 305:567-580.
37. Lechner M, Findeiß S, Steiner L, Marz M, Stadler PF, Prohaska SJ. 2011. Proteinortho: Detection of (Co-)orthologs in large-scale analysis. *BMC Bioinformatics* 12:124.
38. Wen Q-F, Liu S, Dong C, Guo H-X, Gao Y-Z, Guo F-B. 2019. Geptop 2.0: An Updated, More Precise, and Faster Geptop Server for Identification of Prokaryotic Essential Genes. *Front Microbiol* 10:1-6.
39. Chao MC, Abel S, Davis BM, Waldor MK. 2016. The design and analysis of transposon insertion sequencing experiments. *Nat Rev Microbiol* 14:119-128.
40. Zhao L, Anderson MT, Wu W, Harry HL, Bachman MA. 2017. TnseqDiff: Identification of conditionally essential genes in transposon sequencing studies. *BMC Bioinformatics* 18:326.
41. Gu Z, Gu L, Eils R, Schlesner M, Brors B. 2014. Circlize implements and enhances circular visualization in R. *Bioinformatics* 30:2811-2812.
42. Dorman MJ, Feltwell T, Goulding DA, Parkhill J, Short FL. 2018. The Capsule Regulatory Network of *Klebsiella pneumoniae* Defined by density-TraDISort. *mBio* 9:e01863-18.
43. Wickham H. 2016. *ggplot2: Elegant Graphics for Data Analysis*, 2 ed. Springer-Verlag, New York.
44. Carver T, Harris SR, Berriman M, Parkhill J, McQuillan JA. 2012. Artemis: An integrated platform for visualization and analysis of high-throughput sequence-based experimental data. *Bioinformatics* 28:464-469.

45. Turner KH, Wessel AK, Palmer GC, Murray JL, Whiteley M. 2015. Essential genome of *Pseudomonas aeruginosa* in cystic fibrosis sputum. *Proc Natl Acad Sci USA* 112:4110-4115.
46. Loper JE, Hassan KA, Mavrodi DV, Davis II EW, Lim CK, Shaffer BT, Elbourne LDH, Stockwell VO, Hartney SL, Breakwell K, Henkels MD, Tetu SG, Rangel LI, Kidarsa TA, Wilson NL, van de Mortel JE, Song C, Blumhagen R, Radune D, Hostetler JB, Brinkac LM, Durkin AS, Kluepfel DA, Wechter WP, Anderson AJ, Kim YC, Pierson III LS, Pierson EA, Lindow SE, Kobayashi DY, Raaijmakers JM, Weller DM, Thomashow LS, Allen AE, Paulsen IT. 2012. Comparative genomics of plant-associated *Pseudomonas* spp.: Insights into diversity and inheritance of traits involved in multitrophic interactions. *PLoS Genet* 8:e1002784.
47. Levy A, Gonzalez IS, Mittelviefhaus M, Clingenpeel S, Paredes SH, Miao J, Wang K, Devescovi G, Lebeis S, Jin Z, McDonald M, Klein AP, Meghan E. 2018. Genomic features of bacterial adaptation to plants. *Nat Genet* 50:138-150.
48. Suyama M, Bork P. 2001. Evolution of prokaryotic gene order: genome rearrangements in closely related species. *Trends Genet* 17:10-13.
49. Silby MW, Cerdeño-Tárraga AM, Vernikos GS, Giddens SR, Jackson RW, Preston GM, Zhang X-X, Moon CD, Gehrig SM, Godfrey SAC, Knight CG, Malone JG, Robinson Z, Spiers AJ, Harris S, Challis GL, Yaxley AM, Harris D, Seeger K, Murphy L, Rutter S, Squares R, Quail MA, Saunders E, Mavromatis K, Brettin TS, Bentley SD, Hothersall J, Stephens E, Thomas CM, Parkhill J, Levy SB, Rainey PB, Thomson NR. 2009. Genomic and genetic analyses of diversity and plant interactions of *Pseudomonas fluorescens*. *Genome Biol* 10:R51.
50. Winsor GL, Griffiths EJ, Lo R, Dhillon BK, Shay JA, Brinkman FSL. 2016. Enhanced annotations and features for comparing thousands of *Pseudomonas* genomes in the *Pseudomonas* genome database. *Nucleic Acids Res* 44:D646-D653.
51. Gao Y, Yurkovich JT, Seo SW, Kabimoldayev I, Dräger A, Chen K, Sastry AV, Fang X, Mih N, Yang L, Eichner J, Cho BK, Kim D, Palsson BO. 2018. Systematic discovery of uncharacterized transcription factors in *Escherichia coli* K-12 MG1655. *Nucleic Acids Res* 46:10682-10696.
52. Grazziotin AL, Vidal NM, Venancio TM. 2015. Uncovering major genomic features of essential genes in Bacteria and a methanogenic Archaea. *FEBS J* 282:3395-3411.

53. Royet K, Parisot N, Rodrigue A, Gueguen E, Condemine G. 2019. Identification by Tn-seq of *Dickeya dadantii* genes required for survival in chicory plants. Mol Plant Pathol 20:287-306.
54. Elbourne LDH, Tetu SG, Hassan KA, Paulsen IT. 2017. TransportDB 2.0: A database for exploring membrane transporters in sequenced genomes from all domains of life. Nucleic Acids Res 45:D320-D324.
55. Feissner RE, Richard-Fogal CL, Frawley ER, Kranz RG. 2006. ABC transporter-mediated release of a haem chaperone allows cytochrome c biogenesis. Mol Microbiol 61:219-231.
56. Davidson AL, Dassa E, Orelle C, Chen J. 2008. Structure, Function, and Evolution of Bacterial ATP-Binding Cassette Systems. Microbiol Mol Biol Rev 72:317-364.
57. Mavrodi DV, Loper JE, Paulsen IT, Thomashow LS. 2009. Mobile genetic elements in the genome of the beneficial rhizobacterium *Pseudomonas fluorescens* Pf-5. BMC Microbiol 9:8.
58. He J, Baldini RL, Déziel E, Saucier M, Zhang Q, Liberati NT, Lee D, Urbach J, Goodman HM, Rahme LG. 2004. The broad host range pathogen *Pseudomonas aeruginosa* strain PA14 carries two pathogenicity islands harboring plant and animal virulence genes. Proc Natl Acad Sci USA 101:2530-2535.
59. Carter MQ, Chen J, Lory S. 2010. The *Pseudomonas aeruginosa* pathogenicity island PAPI-1 is transferred via a novel type IV pilus. J Bacteriol 192:3249-3258.
60. Matsui H, Sano Y, Ishihara H, Shinomiya T. 1993. Regulation of pyocin genes in *Pseudomonas aeruginosa* by positive (*priN*) and negative (*priR*) regulatory genes. J Bacteriol 175:1257-1263.
61. Peng C, Gao F. 2014. Protein localization analysis of essential genes in prokaryotes. Sci Rep 4:6001.
62. Green ER, Meccas J. 2016. Bacterial Secretion Systems: An Overview. Microbiol Spectr 4:VMBF-0012-2015.
63. Kranz RG, Richard-Fogal C, Taylor J-S, Frawley ER. 2009. Cytochrome c Biogenesis: Mechanisms for Covalent Modifications and Trafficking of Heme and for Heme-Iron Redox Control. Microbiol Mol Biol Rev 73:510-528.

64. Mavridou DAI, Ferguson SJ, Stevens JM. 2012. The interplay between the disulfide bond formation pathway and cytochrome c maturation in *Escherichia coli*. FEBS Lett 586:1702-1707.
65. Ishiga Y, Ichinose Y. 2016. *Pseudomonas syringae* pv. *tomato* OxyR is required for virulence in tomato and *Arabidopsis*. Mol Plant-Microbe Interact 29:119-131.
66. Lim CK, Hassan KA, Penesyan A, Loper JE, Paulsen IT. 2013. The effect of zinc limitation on the transcriptome of *Pseudomonas protegens* Pf-5. Environ Microbiol 15:702-715.
67. Bough L, Gallagher LA, Patrapuvich R, Clifton MC, Gardberg AS, Edwards TE, Armour B, Begley DW, Dieterich SH, Dranow DM, Abendroth J, Fairman JW, Fox D, Staker BL, Phan I, Gillespie A, Choi R, Nakazawa-Hewitt S, Nguyen MT, Napuli A, Barrett L, Buchko GW, Stacy R, Myler PJ, Stewart LJ, Manoel C, Van Voorhis WC. 2013. Combining Functional and Structural Genomics to Sample the Essential *Burkholderia* Structome. PLoS ONE 8:e53851.
68. de Berardinis V, Vallenet D, Castelli V, Besnard M, Pinet A, Cruaud C, Samair S, Lechaplais C, Gyapay G, Richez C, Durot M, Kreimeyer A, Le Fèvre F, Schächter V, Pezo V, Döring V, Scarpelli C, Médigue C, Cohen GN, Marlière P, Salanoubat M, Weissenbach J. 2008. A complete collection of single-gene deletion mutants of *Acinetobacter baylyi* ADP1. Mol Syst Biol 4:174.
69. Grundy F, Henkin T. 2002. Synthesis of Serine, Glycine, Cysteine, and Methionine. In Sonenshein A, Losick R, Hoch J (ed), *Bacillus subtilis* and Its Closest Relatives. ASM Press, Washington DC. doi:10.1128/9781555817992.ch18.
70. Conway T. 1992. The Entner-Doudoroff pathway: history, physiology and molecular biology. FEMS Microbiol Lett 103:1-28.
71. Lerouge I, Vanderleyden J. 2002. O-antigen structural variation: Mechanisms and possible roles in animal/plant-microbe interactions. FEMS Microbiol Rev 26:17-47.
72. Seliger SS, Mey AR, Valle AM, Payne SM. 2001. The two TonB systems of *Vibrio cholerae*: Redundant and specific functions. Mol Microbiol 39:801-812.
73. Zhao Q, Poole K. 2000. A second *tonB* gene in *Pseudomonas aeruginosa* is linked to the *exbB* and *exbD* genes. FEMS Microbiol Lett 184:127-132.
74. Bentley R. 1990. The Shikimate Pathway — A Metabolic Tree with Many Branches. Crit Rev Biochem Mol Biol 25:307-384.

75. Tzin V, Galili G, Aharoni A. 2012. Shikimate Pathway and Aromatic Amino Acid Biosynthesis. eLS doi:10.1002/9780470015902.a0001315.pub2.
76. Lin S, Cronan JE. 2011. Closing in on complete pathways of biotin biosynthesis. Mol Biosyst 7:1811-1821.
77. Lutkenhaus J. 2007. Assembly Dynamics of the Bacterial MinCDE System and Spatial Regulation of the Z Ring. Annu Rev Biochem 76:539-562.
78. Bendezú FO, de Boer PAJ. 2008. Conditional Lethality, Division Defects, Membrane Involution, and Endocytosis in *mre* and *mrd* Shape Mutants of *Escherichia coli*. J Bacteriol 190:1792-1811.
79. Fujita Y, Matsuoka H, Hirooka K. 2007. Regulation of fatty acid metabolism in bacteria. Mol Microbiol 66:829-839.
80. Kang Y, Nguyen DT, Son MS, Hoang TT. 2008. The *Pseudomonas aeruginosa* PsrA responds to long-chain fatty acid signals to regulate the *fadBA5* β -oxidation operon. Microbiology 154:1584-1598.
81. Son MS, Nguyen DT, Kang Y, Hoang TT. 2008. Engineering of *FRT-lacZ* fusion constructs: Induction of the *Pseudomonas aeruginosa* *fadAB1* operon by medium and long chain-length fatty acids. Plasmid 59:111-118.
82. Deutscher MP. 2006. Degradation of RNA in bacteria: comparison of mRNA and stable RNA. Nucleic Acids Res 34:659-666.
83. Mackie GA. 2013. RNase E: At the interface of bacterial RNA processing and decay. Nat Rev Microbiol 11:45-57.
84. Scapin G, Blanchard JS. 1998. Enzymology of Bacterial Lysine Biosynthesis. In Purich DL (ed), Advances in Enzymology and Related Areas of Molecular Biology doi:10.1002/9780470123188.ch8. John Wiley & Sons, Inc., New York.
85. Parsons JB, Rock CO. 2013. Bacterial lipids: Metabolism and membrane homeostasis. Prog Lipid Res 52:249-276.
86. Akerley BJ, Rubin EJ, Novick VL, Amaya K, Judson N, Mekalanos JJ. 2002. A genome-scale analysis for identification of genes required for growth or survival of *Haemophilus influenzae*. Proc Natl Acad Sci USA 99:966-971.
87. Gerdes SY, Scholle MD, Campbell JW, Balázsi G, Ravasz E, Daugherty MD, Somera AL, Kyrpides NC, Anderson I, Gelfand MS, Bhattacharya A, Kapatral V, D'Souza M, Baev MV, Grechkin Y, Mseeh F, Fonstein MY, Overbeek R, Barabási A-L, Oltvai ZN,

- Osterman AL. 2003. Experimental Determination and System Level Analysis of Essential Genes in *Escherichia coli* MG1655. J Bacteriol 185:5673-5684.
88. Deutschbauer A, Price MN, Wetmore KM, Shao W, Baumohl JK, Xu Z, Nguyen M, Tamse R, Davis RW, Arkin AP. 2011. Evidence-Based Annotation of Gene Function in *Shewanella oneidensis* MR-1 Using Genome-Wide Fitness Profiling across 121 Conditions. PLoS Genet 7:e1002385.
89. Zawadzke LE, Bugg TDH, Walsh CT. 1991. Existence of two D-alanine:D-alanine ligases in *Escherichia coli*: cloning and sequencing of the *ddlA* gene and purification and characterization of the DdlA and DdlB enzymes. Biochemistry 30:1673-1682.
90. Barreteau H, Kovač A, Boniface A, Sova M, Gobec S, Blanot D. 2008. Cytoplasmic steps of peptidoglycan biosynthesis. FEMS Microbiol Rev 32:168-207.
91. Gonzales T, Robert-Baudouy J. 1996. Bacterial aminopeptidases: Properties and functions. FEMS Microbiol Rev 18:319-344.
92. Williams HD, Zlosnik JEA, Ryall B. 2006. Oxygen, Cyanide and Energy Generation in the Cystic Fibrosis Pathogen *Pseudomonas aeruginosa*, p 1-71. In Poole RK (ed), Adv Microb Physiol, vol 52. Academic Press.
93. Torres A, Kasturiarachi N, DuPont M, Cooper VS, Bomberger J, Zemke A. 2019. NADH dehydrogenases in *Pseudomonas aeruginosa* growth and virulence. Front Microbiol 10:75.
94. Rubin BE, Wetmore KM, Price MN, Diamond S, Shultzaberger RK, Lowe LC, Curtin G, Arkin AP, Deutschbauer A, Golden SS. 2015. The essential gene set of a photosynthetic organism. Proc Natl Acad Sci USA 112:E6634-E6643.
95. Gottesman S. 1996. Proteases and their targets in *Escherichia coli*. Annu Rev Genet 30:465-506.
96. Ito F, Tamiya T, Ohtsu I, Fujimura M, Fukumori F. 2014. Genetic and phenotypic characterization of the heat shock response in *Pseudomonas putida*. MicrobiologyOpen 3:922-936.
97. Ghazaei C. 2017. Role and mechanism of the Hsp70 molecular chaperone machines in bacterial pathogens. J Med Microbiol 66:259-265.
98. Patel PH, Suzuki M, Adman E, Shinkai A, Loeb LA. 2001. Prokaryotic DNA polymerase I: Evolution, structure, and "base flipping" mechanism for nucleotide selection. J Mol Biol 308:823-837.

- 918 99. Du S, Lutkenhaus J. 2017. Assembly and activation of the *Escherichia coli* divisome.
 919 Mol Microbiol 105:177-187.
 920