Sex differences in cortisol and memory following acute social stress **in amnesic mild cognitive impairment**

Kelly J. Murphy^{a,b,*}, Travis E. Hodges^c, Paul A.S. Sheppard^c, Angela K. Troyer^{a,b},
Elizabeth Hampson^d, & Liisa A.M. Galea^{c,*}

^a Neuropsychology and Cognitive Health Program, Baycrest, Toronto, ON, Canada.

^b Psychology Department, University of Toronto, Toronto, ON, Canada

^c Djavad Mowafaghian Centre for Brain Health, Department of Psychology, University of British Columbia, Vancouver, BC, Canada.

^d Psychology Department, Western University, London, ON, Canada.

* corresponding author; Email: liisa.galea@ubc.ca

ORCID IDs

KJM: 0000-0002-3827-5814

TEH: 0000-0002-5993-2633

PASS: 0000-0001-7432-1446

AKT: 0000-0001-5660-4127

EH: 0000-0003-4487-9013

LAMG: 0000-0003-2874-9972

Abstract: 300 words

Manuscript: 8746 words

Abstract

Objective: Older adults with amnesic mild cognitive impairment (aMCI) develop Alzheimer's type dementia approximately ten times faster annually than the normal population. Adrenal hormones are associated with aging and cognition. We investigated the relationship between acute stress, cortisol, and memory function in aMCI with an exploratory analysis of sex. **Method:** Salivary cortisol was sampled diurnally and during two test sessions, one session with the Trier Social Stress Test (TSST), to explore differences in the relationship between cortisol and memory function in age-normal cognition (NA) and aMCI. Participants with aMCI (n=6 women, 9 men; mean age=75) or similarly aged NA (n=9 women, 7 men, mean age=75) were given tests of episodic, associative, and spatial working memory with a psychosocial stressor (TSST) in the second session. **Results:** The aMCI group performed worse on the memory tests than NA as expected, and males with aMCI had elevated cortisol levels on test days. Immediate episodic memory was enhanced by social stress in NA but not in the aMCI group, indicating that stress-induced alterations in memory are different in individuals with aMCI. High cortisol was associated with impaired performance on episodic memory in aMCI males only. Cortisol in Session 1 moderated the relationship with spatial working memory, whereby higher cortisol was associated with worse performance in NA, but better spatial working memory in aMCI. In addition, effects of aMCI on perceived anxiety in response to stress exposure were moderated by stress-induced cortisol in a sex-specific manner. **Conclusions:** We show effects of aMCI on Test Session cortisol levels and effects on perceived anxiety, and stress-induced impairments in memory in males with aMCI in our exploratory sample. Future studies should explore sex as a biological variable as our findings suggests that effects at the confluence of aMCI and stress can be obfuscated without sex as a consideration.

KEYWORDS: diurnal, psychosocial stressor, men, women, aMCI, Trier Social Stress Test

Introduction

Normal aging results in declines in some cognitive domains, such as episodic memory, but not in others, such as experiential knowledge (Grady, 2012). Cognitive decline with aging is correlated with region-specific changes in prefrontal cortex and medial temporal lobe, a key change being hippocampal volume loss (Raz & Rodrigue, 2006) and these declines are accelerated in those with suspected Alzheimer's Disease (AD) (Shi et al., 2009). Older adults with mild cognitive impairment (MCI) develop clinical dementia of the AD type at a rate of 10-30% annually, depending on MCI subtype, whereas those without MCI develop dementia at a rate of only 1% to 2% annually (Busse et al., 2003; Lupien et al., 1998). Thus, it is critical to identify neurobiological factors that may distinguish MCI from normal aging, such as differences in cortisol levels and their response to stress.

Dysregulation of the hypothalamic-pituitary-adrenal (HPA) system, including the stress hormone cortisol, has been linked to memory performance, aging, and hippocampal volume (Lupien et al., 1998; Sindi et al., 2014; Justice, 2018). Indeed, participants with suspected AD have higher levels of plasma (morning, 24 h release) or morning cerebrospinal fluid (CSF) cortisol than those experiencing normal cognitive aging (Hartmann et al., 1997; Doecke et al., 2012; Laske et al., 2011). Morning (CSF) levels of cortisol are also higher in MCI due to suspected AD, referred to as amnesic MCI (aMCI), as compared to older adults experiencing normal aging (NA) or MCI of other types (Popp et al., 2015). Moreover, aMCI individuals with higher morning CSF cortisol levels experienced accelerated clinical worsening and cognitive decline than those with a lower levels of cortisol, with a higher proportion of males in the aMCI group (Popp et al., 2015). Intriguingly, despite higher levels of cortisol, MCI participants experience lower levels of perceived stress during cognitive performance compared to age-matched healthy controls (Guerdoux-Ninot & Trouillet, 2019). Previous studies have rarely examined multiple timepoints of cortisol, stress induced cortisol, diurnal cortisol, or have used biological sex as a discovery variable, all factors which may contribute to the findings (Hidalgo et al., 2019; Yan et al., 2018). Cortisol is well known to vary in a diurnal pattern (Adam et al., 2017) and diurnal patterns are flattened in dementia of the AD type (Rasmuson et al., 2011; Ferrari et al., 2001). Importantly, to our knowledge, stress-induced cortisol has not been studied in relation to aMCI status and memory performance. Acute stress may be a more salient variable of HPA dysregulation to examine whether it can perturbate memory. Indeed, normally aging older adults who show less cortisol reactivity to acute stress using the Trier Social Stress Test (TSST) are more at risk to develop cognitive decline characteristic of MCI within 5 years (de Souza-Talarico et al., 2020). Thus, it is important to examine not only diurnal fluctuations in cortisol but also response to acute stress to determine whether these

biomarkers modulate memory and how they may relate to progression to neurodegenerative disease. In addition, perceived stress, along with stress-induced cortisol, may be as important to investigate (Aggarwal et al., 2014). Furthermore, sex differences must be considered given previous studies have identified sex differences in cortisol levels in response to stress (Kudielka & Kirschbaum, 2005) which vary by age (for review see Hidalgo et al., 2019). Indeed, few studies to date have examined the relationships between cortisol, stress, memory, and potential differences between males and females in older populations (for review see Hidalgo et al., 2019), particularly with regards to MCI or AD.

Sex differences are seen in incidence of MCI (Gale et al., 2016; Koran et al., 2017; Mielke et al., 2014; Duarte-Guterman et al., 2020), with males more likely to develop MCI (both aMCI and non-amnestic subtypes) than females (Jack et al., 2019; Roberts et al., 2012), although there are conflicting reports that are likely due to methodological differences (e.g. Mielke et al., 2014). However, AD disproportionately affects females, with significant sex differences observed with regards to severity, neuropathological markers, and rates of cognitive decline (Irvine et al., 2012; Koran et al., 2017; Sohn et al., 2018). Sex differences in incidence of AD are not seen uniformly (Jack et al., 2019) and may depend on geographic location (Nebel et al., 2018) or greater longevity in women (Mielke et al., 2014). However, there are other sex differences in MCI to AD progression and symptom severity (Sundermann et al., 2017). For example, women tend to develop MCI at a later age, perhaps benefitting from their established superior verbal memory (Sundermann et al., 2017), but progress to AD more rapidly than men when adjusted for age (Irvine et al., 2012; Sohn et al., 2018). Sex differences in the trajectory of cognitive decline are also seen in MCI (Koran et al., 2017; Sohn et al., 2018). Indeed, females with more AD-associated neuropathology (total-tau and amyloid-beta [A β 42] in cerebrospinal fluid), show greater declines in hippocampal volume and cognition compared to males, particularly among MCI individuals using the ADNI database (Koran et al., 2017; Sohn et al., 2018). Furthermore cognitive differences in verbal learning, delayed recall, visual learning and memory between NA and MCI females were significantly greater than those between NA and MCI males (Gale et al., 2016). This pattern in sex differences persisted in those with AD (Gale et al., 2016). Thus, sex differences in severity and progression to AD are seen in individuals with MCI and identifying the biological causes of this phenomenon is critical to treatment and prevention.

Gonadal production of sex steroids is reduced, but not entirely eliminated, with age in women and, to a lesser extent, men; however, adrenal cortisol production increases with age (Laughlin & Barrett-Connor, 2000). Although both sexes show increased cortisol levels with increased age, this effect is 3 times more pronounced in females (Otte et al., 2005) and increased cortisol levels are linked

to poorer cognition and smaller hippocampal volume in older age (Lupien et al., 1998). Furthermore, as mentioned above females with MCI and AD present with greater declines in hippocampal atrophy and cognition than males (Irvine et al., 2012; Koran et al., 2017; Sohn et al., 2018), highlighting that the underlying pathophysiology of AD may be different in men and women and should be further explored. Although sex differences in AD have been identified, studies are scarce and even more so in MCI groups.

In this study we explored the relationship between diurnal fluctuations in cortisol, stress-induced cortisol, and memory performance in older adults experiencing aMCI and NA. A spatial working memory task known to be reliant on the integrity of the prefrontal cortex (Courtney et al., 1998) and an episodic and associative memory task known to be reliant on hippocampal integrity (Eichenbaum, 2017) were selected based on the number of glucocorticoid receptors and therefore sensitivity to cortisol in these brain regions (Dedovic et al., 2009) and the potential of fluctuations in cortisol to influence cognitive efficiencies. Because little is known as to whether there are sex differences in the relationship between aMCI and cortisol, we also used exploratory analyses of sex effects in the present study. We hypothesised that stress-induced cortisol, *via* the application of a psychosocial stressor (Trier Social Stress Test; Kirschbaum et al., 1993), would worsen memory scores and possibly alter perceived anxiety in individuals with aMCI compared to NA and that there would be sex differences in these effects.

Materials and methods

Participants

Older adults with age-normal memory (normal cognitive aging- NA) and with mild memory decline (aMCI) suggestive of neurodegenerative disease of the Alzheimer type (Albert et al., 2011) were recruited for this study and provided informed voluntary consent to participate. The following brief battery of neuropsychological measures were administered during Session 1 to confirm group membership: cognitive screening (Mini-Mental Status Exam (MMSE, Folstein & Folstein, 1974), expressive vocabulary (Vocabulary, Wechsler, 1997), attention tests involving auditory attention span (Digit Span, Wechsler, 1997) and speed and attention switching (Trail Making Tests A and B; -Spreen & Strauss, 1998), confrontation naming (Boston Naming Test, Kaplan et al., 1983), visuospatial construction, and immediate memory (Rey-Osterrieth Complex Figure-copy and immediate recall, Spreen & Strauss, 1998); and mood status (Hospital Anxiety and Depression Scale, HADS, Zigmond & Snaith, 1983). Criteria for establishing aMCI status were adherent to those described in Petersen (2004)

and Albert et al. (2011). Participants were classified with single domain amnesic MCI if memory performance was revealed to be the only cognitive domain among those tested (which included attention, psychomotor speed, memory, language, visual spatial ability, and executive function) for which age-scaled scores were lower than expected based on estimated verbal IQ (established on a test of expressive vocabulary) and based on demonstrated performance in the other cognitive domains examined (see Table 1).

Normal Cognitive Aging (NA) group

Fifteen older adults (age 61-86 years) experiencing NA were recruited *via* community talks, newspaper advertisements, and databases of research volunteers. Prior to invitation to participate, normal general cognitive status, using the Telephone Interview for Cognitive Status, and health status were confirmed in a telephone screening interview. At the first of two sessions, health history was further queried to verify that there was no history of a neurological, medical, or psychiatric disorder, substance abuse, or medications affecting cognition. As described, NA was confirmed during Session 1 by measuring performance on a brief battery of neuropsychological tests. Three participants initially recruited as NA were identified as meeting criteria for aMCI based on Session 1 interview and lower than age and education expected performance on immediate recall of a complex figure.

Amnesic Mild Cognitive Impairment (aMCI) group

Sixteen individuals (age 59-85 years), recruited from physician referrals, from databases of research volunteers, and from newspaper advertisement, were classified as meeting the National Institute on Aging-Alzheimer's Association classification criteria for aMCI (Albert et al., 2011). The single domain aMCI status of most of the aMCI participants had been previously established and the stability of this classification was confirmed by the interview and neuropsychological testing administered during Session 1. As previously stated, three of the participants initially recruited to the NA group were found to meet criteria for aMCI at Session 1 and were transferred to the aMCI group.

[INSERT TABLE 1 HERE]

Procedure

Participants completed alternate versions of episodic, associative, and spatial memory tasks across two test sessions (Figure 1) conducted 7-14 days apart. The TSST (Kirschbaum et al., 1993), a

psychosocial stressor, was applied during the second test session and is described below. Salivary cortisol was collected during both test sessions for all participants and diurnal samples were collected as described below. Saliva was chosen because it most closely represents bioavailable cortisol (i.e., the fraction of the circulating hormone that is biologically available to tissues).

Episodic memory

Two highly correlated versions (forms 5 and 6) of the Hopkins Verbal Learning Test-Revised (HVLT-R) were used (Session 1: form 6; Session 2: form 5; with the exception of one aMCI participant to whom they were presented in the opposite order). This task involves an oral presentation of a 12-item word list over three learning trials, followed by a 20-minute delayed recall trial and a forced choice yes/no recognition trial. The recognition trial consists of 24 items comprised of the 12 target words, six semantically related foils, and six un-related foils. Measures of interest included total recall across three learning trials, total delayed recall, and retention.

Associative recognition

A face-name associative recognition test, created by Troyer and colleagues (2011, 2012; modeled after Mayes et al., 2004), was used. Stimuli consisted of visually presented black-and-white images of faces (half male and half female) paired with aurally presented first names. Two versions of the task were used, each with 28 face-name pairs. During the task, 20 faces were individually presented on the computer screen for 6 seconds each with an inter-stimulus interval of 0.5 seconds; the examiner read the name associated with each face at the onset of each new face stimulus. Two study phases, differing only in stimulus presentation order, were administered in succession because our previous research indicated that item memory and association memory differences increase after repeated learning trials (Troyer et al., 2008). Only 16 of the 20 face-name pairs were considered test items as the first and last pairs in each study phase presentation were excluded to reduce primacy and recency effects on recognition accuracy. Following a 30 second delay, yes/no recognition testing was conducted with 24 face-name pairs presented, including eight intact pairs, eight recombined pairs, and eight new pairs presented in random order. During testing the examiner orally presented the name in the form of a question “Did I tell you this was [NAME]?” when the face appeared on the screen. Participants were instructed to say “yes” only to faces they had seen before that were paired with the correct name and “no” to faces they had not seen before, faces that were paired with the wrong name, or names they had not heard before. Immediately following testing, procedure verification was undertaken (i.e.

participants retold the yes/no rules to the examiner). Participants were presented with unique, but equivalent (Troyer et al., 2011), sets of face-name pairs during Sessions 1 and 2. Associative recognition calculated as the difference between the proportion of correctly identified intact pairs (same face-name pairs viewed at study) and proportion of false alarms to recombined pairs (different pairings of previously viewed faces and names at study) was the primary measure of interest. The decision to focus on associative recognition was based on our previous research demonstrating this measure was most sensitive to aMCI and hippocampal volume loss (Troyer et al., 2012).

Spatial working memory

This task was modeled after the stimuli and procedures of Duff and Hampson (2001). A 4x5 rectangular array, measuring approximately 27cm in length and 34cm in width, consisting of coloured squares (10 colours, each represented twice) that were hidden under removable covers, was presented on a tabletop at which participants were seated. The coloured squares were randomly arranged on a uniform white backing and completely concealed beneath uniform white covers that could be temporarily lifted by participants to reveal the coloured square beneath. Participants were instructed to find all 10 pairs of matching coloured squares in as few choices as possible by lifting the covers two at a time. Prior to beginning the task, participants were familiarised with the colours of the test stimuli by having them view and name a set of 10 individual coloured squares. Each time a matching pair was located on the stimulus array, the examiner placed an individual coloured square representing the colour of the pair discovered at the top of the rectangular array, so participants did not need to remember which colour pairs had been found. Measures of interest included: the number of choices (squares uncovered) made in discovering all 10 matching pairs (criterion) and the time taken to reach criterion. Participants were told they would be timed and that they should attempt to locate all 10 pairs in as few choices as possible. Once they reached criterion, a second trial was immediately administered, with a third trial administered following a 30-minute delay. Locations of the coloured squares was constant within session but changed between Sessions 1 and 2.

Psychosocial stressor (TSST)

The psychosocial stressor used in this study was modeled after the TSST (Kirschbaum et al., 1993). The stressor was introduced immediately following the first saliva collection at 10:10h. Participants were instructed to prepare a five-minute speech on the topic of ‘*The effect of tuna fishing on the dolphins and other ocean animals*’ to be presented to a panel of three evaluators, including the

examiner. They were given a pencil and paper and told to write down the points they would like to make in their speech for which they would have 10-minutes to prepare. The examiner then left the room and returned 10 minutes later, collected a saliva sample from the participant (anticipation period), and then led the participant to a conference room to give their speech. Participants were instructed to leave their written notes behind, to give their speech from memory, and to try and speak for five minutes, which was timed by the examiner. Immediately following the public speech, participants engaged in a five-minute serial subtraction task in which they were asked to count backwards aloud by 13 from the number 1022 as quickly and accurately as possible in front of the panel of evaluators. When an error was committed the participant was instructed to begin again from the number 1022. Because perceived stress/anxiety has been found to be lower in aMCI compared to NA individuals performing an attention-based task (Stroop-task; Guerdoux-Ninot & Trouillet, 2019), participants were asked if they found the test anxiety provoking and if so to rate their perceived anxiety during the public speech from 0 (low anxiety) to 100 (high anxiety). Lastly, participants were led back to the test room to undertake memory testing (episodic, associative, spatial) and to provide additional saliva samples as described below.

[INSERT FIGURE 1 HERE]

Saliva collection and analysis

Multiple salivary cortisol samples were taken using the Salivette™ method (Sarstedt Inc, Sarstedtstraße, Numbrecht, Germany) across both the test sessions (for all participants) and at home (to examine diurnal variations). Participants were instructed not to eat, drink, or smoke for at least 30 minutes prior to saliva collection and to rinse their mouths with water 5 minutes before collecting the saliva. For the test sessions, both sessions were conducted in the morning from 10:00h to 12:00h. Saliva was collected at 10:10h and 12:00h at the first test session and during the second session at 10:10h; immediately following the anticipation period to the application of a psychosocial stressor (~10:30h); 30 minutes following the application of the psychosocial stressor (~11:00h); and at about 12:00h at the conclusion of the second test session. In addition, basal salivary diurnal cortisol samples were requested from participants across three agreed upon days intervening between the test sessions. Five samples were collected per day on the following schedule: 30-minutes after awakening (ranged from 5:30 am to 8:30h), 09:00, 16:00, 19:00, and 21:00h. Phone call reminders and verifications were provided by the examiner for each of the 4 specified clock times on all three collection days.

Participants were given pre-labeled saliva collection tubes at the end of Session 1 and instructed to store their collected samples in the home refrigerator. Collection time of day was further verified by requiring participants to record the collection time on a label provided on the collection tube. The basal samples were gathered from participants when they returned for the second test session.

Saliva was centrifuged at 1500g and kept frozen at -20°C prior to analysis. Cotton-based collection is suitable for cortisol determinations (Büttler et al., 2018). Salivary cortisol was analyzed in duplicate by the Neuroendocrinology Assay Laboratory at the University of Western Ontario (EH). An established ¹²⁵I solid-phase radioimmunoassay was used (Norman et al., 2010), based on antibody and tracer obtained from Siemens Healthcare Diagnostics (Deerfield, IL). The Laboratory specialises in saliva determinations. Briefly, saliva was analyzed directly, without extraction, using a 200µL aliquot and an extended 3hr incubation at room temperature. The calibration curve was diluted 1:10 and ranged from 0-138 nmol/L. The intra-assay coefficient of variation calculated across low, medium, and high pools averaged 4.2% and the sensitivity of the assay was < 0.25 nmol across 3 assay runs. All samples from a given participant were analyzed in the same assay run and the average salivary cortisol concentration across the two duplicates (in nmol/L) was the value used for all statistical analyses. All cortisol data was log-transformed (log₁₀) due to non-normality.

We also examined the diurnal and TSST cortisol levels area under the curve (AUC) using two formulas for AUC one with respect to the ground (AUC_g) and one with respect to the increase (AUC_i) (Pruessner et al., 2003) using the following formulas:

$$AUC_g = \sum_{i=1}^{n-1} \frac{(m_{(i+1)} + m_i) * t_i}{2}$$

$$AUC_i = AUC_g - m_1 * \sum_{i=1}^{n-1} t_i$$

Where t_i - is the distance between time measurements, m₁ - is the first cortisol measurement

Data Analyses

Statistical analyses were performed using Statistica TIBCO Software (Palo Alto, CA, USA) and SPSS with the PROCESS package. Because we were examining sex differences as an exploratory factor, we initially conducted analyses on age and education level between the sexes (see Table 2). Due to differences in ages between groups we used age as a covariate in all analyses. General linear model repeated measures analyses of covariance (RMANCOVAs) were conducted on log-transformed cortisol measures (diurnal, session time) and cognitive measures (HVLt, SPWM, AR)

using group (NA, aMCI) and sex (male, female) as the between-subject factors and with either Session (1, 2) or time of day (30 min, 9 AM, 4PM, 7pm, 9PM) or time during test Session (10:10, 10:30, 11, 1130) as the within-subjects factor. On certain measures (represented in Table 1) general linear model ANCOVA were conducted using group (NA, aMCI) as the between-subject factor. *Post hoc* analyses used Newman-Keul's comparisons. Due to the small sample size, exploratory analyses on possible sex effects were run using a Bonferroni correction on *a priori* analyses, using a two-tailed significance criterion of $p=0.10$. Otherwise statistical significance was $p=0.05$ was used for all other statistical tests conducted. Chi-square analyses were completed for frequency of sexes between groups and for anxiety scores. For exploratory moderation analyses hierarchical linear regressions were performed to test the overarching hypothesis that the magnitude of Session 1 and Session 2 cortisol moderates the relationship between group (MCI, NA) and perceived anxiety during the TSST, episodic memory, associative memory, or working memory. Hierarchical linear regressions that included either Session 1 cortisol (AUCg) or Session 2 cortisol (AUCg) as a moderators were conducted for males and females separately. Dummy codes were created for participant group (MCI = 1, NA = 0), cortisol data were $\log_{10}$ transformed and standardized, and all other data were standardized. Regression models with significant interactions are reported.

Results

Males with aMCI had higher levels of cortisol than NA males during the test sessions.

Cortisol levels during the two test sessions were analyzed separately as the psychosocial stressor (TSST) was conducted during Session 2. For Session 1 (without the TSST), males with aMCI had significantly higher levels of cortisol than any other group at the first time point (10:10AM) (all $ps<0.001$; Cohens d: for NA males=0.79, NA females=1.99, and aMCI females=1.67); time by group by sex interaction: $F_{(1,27)}=3.65$, $p=0.066$, $\eta_p=0.12$; Figure 2A). There were also main effects of sex and time (main effect of sex: $F_{(1,27)}=16.24$, $p<0.001$, $\eta_p=0.37$; main effect of time: $F_{(1,27)}=19.515$, $p<0.0001$, $\eta_p=0.42$). For AUCg in Session 1, aMCI males had higher levels of cortisol than all other groups (all $ps<0.05$; Cohens d: for NA males=0.81, NA females=1.57, and aMCI females=1.95; interaction effect of sex by group: $F_{(1,27)}=4.013$, $p=0.055$, $\eta_p=0.129$; Figure 2B). There was also a main effect of sex ($F_{(1,27)}=10.25$, $p=0.003$, $\eta_p=0.28$) but not group ($p=0.37$; see Figure 2B). There were no significant differences using AUCi ($p>0.15$, group: $\eta_p=0.03$, sex $\eta_p=0.06$, group by sex $\eta_p=0.07$).

During Session 2, in which the TSST was performed, aMCI participants had higher cortisol levels than NA (main effect of group: $F_{(1,27)}=7.86$, $p<0.009$, $\eta_p=0.23$). Furthermore, males had higher

levels of cortisol than females (main effect of sex: $F_{(1, 27)}=13.58, p=0.001, \eta_p=0.33$). Furthermore, cortisol levels were highest 30min after the TSST was initiated compared to all other time points as expected (all $ps < 0.03$; main effect of time: $F_{(3, 81)}=3.95, p=0.01, \eta_p=0.13$). *A priori* analyses indicated that males with aMCI had significantly higher levels of cortisol during the first two time points in the second session prior to the TSST than all other groups (all $ps < 0.001$; Cohens d: timepoint 1: for NA males=1.45, NA females=2.33, and aMCI females=2.00, timepoint 2: for NA males=1.25, NA females=2.44, and aMCI females=1.82). Importantly, males with aMCI did not show any changes in cortisol levels across Session 2 (all $ps > 0.18$) whereas females with aMCI had significantly higher levels of cortisol 30min post-TSST than NA females ($p=0.003$ Cohens d=1.03) but not at any other time point (all $ps > 0.18$; Cohens d= 0.4-0.98; Figure 2C).

For AUCg, in Session 2 we found that males had higher levels of cortisol than females (main effect of sex: $F_{(1, 27)}=8.36, p<0.007, \eta_p=0.24$) and aMCI had higher levels than NA (main effect of group: $F_{(1, 27)}=4.202, p=0.05, \eta_p=0.135$). *A priori* we also found that aMCI males had higher AUCg than NA ($p=0.060$, one-tailed, 0.03; Cohens d: for NA males=0.77, NA females=2.39, and aMCI females=1.40) which was not evident in the females ($p=0.35$; Cohens d=0.73 between female groups) see Figure 2D). There were no significant differences using AUCi ($p>0.27$, group: $\eta_p=0.00$, sex $\eta_p=0.004$, group by sex $\eta_p=0.044$).

[INSERT FIGURE 2 HERE]

TSST was endorsed as anxiety provoking by females more than males with aMCI

Both sexes in the NA group endorsed the TSST as anxiety provoking, with 57% of participants indicating that the TSST was anxiety provoking. However, among aMCI participants 80% of females but only 11% of males indicated the TSST speech was anxiety provoking ($\chi^2=6.644, p<0.01$, Table 2). Of those participants who rated the TSST as anxiety provoking there was no significant difference in the rating of the anxiety level ($ps > 0.43$, main effect of group $\eta_p=0.001$, main effect of sex= $\eta_p=0.002$, interaction $\eta_p=0.07$; ratings (mean and standard deviation) and sample size of those that found the TSST anxiety provoking: NA females (n=4)=68.25±25.8, aMCI females (n=4)=50.25±26.10, NA males (n=4)=49.75±35.9; MCI males(n=1)= 64).

Social Stress enhanced immediate recall in the NA but not in participants with aMCI. aMCI participants performed worse on episodic, associative and spatial working memory than NA

Immediate recall of HVLT-R was enhanced after the TSST in Session 2 compared to Session 1 in the NA ($p=0.004$), but no such enhancement was seen in participants with aMCI ($p=0.245$; interaction: group by session: $F_{(1,27)}=7.6$, $p=0.010$, $\eta_p=0.22$, main effect of group: $F_{(1,27)}=41.4$, $p<0.001$, $\eta_p=0.61$). Breaking this down by sex, the NA groups, regardless of sex, showed enhanced recall in Session 2 compared to Session 1 (males ($p=0.02$, Cohens $d=1.14$) females ($p=0.04$, Cohens $d=0.61$)). However, aMCI males had impaired immediate recall ($p=0.04$, Cohen's $d=0.63$) on Session 2 following the stressor compared to Session 1, with no significant enhancement in the females with aMCI across sessions ($p=0.47$; Cohen's $d=0.25$, Figure 3A). This enhancement in recall following the TSST was also seen for delayed recall for the HVLT-R which was administered 40 minutes after the TSST with all groups scoring better on Session 2 than Session 1 (main effect of session $F_{(1,27)}=7.15$, $p=0.012$, $\eta_p=0.21$) and with aMCI scoring worse than NA ($F_{(1,27)}=63.22$, $p<0.001$, $\eta_p=0.70$; Figure 3B). For HVLT-R retention, aMCI participants performed worse compared to NA participants regardless of session (main effect of group $F_{(1,26)}=27.09$, $p<0.001$, $\eta_p=0.51$; Figure 3C).

[INSERT FIGURE 3 HERE]

For associative memory as expected aMCI participants remembered fewer face-name pairs than NA (main effect of group: $F_{(1,27)}=32.86$, $p<0.000$, $\eta_p=0.55$). There were no other main or interaction effects (all $ps>0.25$).

During the spatial working memory across sessions and trials, aMCI participants made more choices than NA across all trials (main effect of group: $F_{(1,27)}=13.6$, $p=0.001$, $\eta_p=0.34$). Furthermore, all participants required fewer choices by the delay trial (all $ps<0.02$; main effect of trial ($F_{(2,52)}=4.80$, $p=0.012$, $\eta_p=0.16$). There were no other significant main effects or interactions (all $ps>0.20$; all $\eta_p<0.09$; Figure 4B).

[INSERT FIGURE 4 HERE]

Diurnal CORT did not differ between male groups

Only 21 of 31 participants completed all five time points for cortisol collection across three consecutive days. We used multiple imputation to calculate missing values for all individuals with more than 75% of the data available (2 participants only had 2/15 and one had 9/15 samples available). Curiously these 3 participants were females with aMCI. As we did not want to use the imputed values for so many missing datapoints we performed the analyses only with males. Analysis of these 16 participants revealed only a main effect of time ($F_{(4,56)}=48.51$, $p<0.001$, $\eta_p=0.776$; Figure 5A), but no

main effect of group ($F_{(1,14)}=0.0349$, $p=0.85$, $\eta_p=0.002$ or interaction ($F_{(4,56)}=0.49$, $p=0.74$, $\eta_p=0.033$).
Post hoc analyses revealed that cortisol was higher at each early timepoint than all other timepoints except there was no difference between the two evening samples. There were no significant differences between groups in time of awakening.

We also calculated average cortisol AUCg and AUCi across all three days in males only. There were no significant effects for AUCg ($p=0.549$, $\eta_p=0.03$) or AUCi ($p=0.6$, $\eta_p=0.02$).

[INSERT FIGURE 5 HERE]

Session 1 Cortisol (AUCg) was associated with better spatial working memory in aMCI females, but the reverse in NA females

We next correlated Session 1 and Session 2 AUCg with episodic, associative and spatial working memory measures across group and sex, as we saw no significant differences in AUCi. Session 1 had the only significant correlations, with females showing positive associations with AUCg and better performance in spatial working memory in aMCI females (in trial 2 ($r=-0.839$, $p=0.037$) and the delay trial ($r=-0.952$, $p=0.003$)) but the opposite patterns in NA females (trial 2 ($r=0.796$, $p=0.01$)). These correlations were significantly different ($z=3.261$, $p=0.001$). In NA females, AUCg was positively associated with better HVLT-retention ($r=0.7017$, $p=0.035$), but negatively associated with associative memory ($r=-0.703$, $p=0.035$). In males, the only correlation was positive in Session 1 was associative memory and AUCg cortisol in aMCI males ($r=0.818$, $p=0.007$). There were no other significant correlations in males or for data in Session 2.

Session 1 cortisol (AUCg) moderates the relationship between aMCI and episodic memory retention in females, working memory in both females and males, and associative recognition in males.

We tested the hypothesis that the magnitude of cortisol (AUCg) moderates the relationship between participant group (aMCI, NA) and episodic (HVLT), associative (face-name), or working memory (spatial working memory) during Session 1 in males and females. Regressions revealed that higher cortisol predicted lower HVLT-R retention scores in aMCI females and higher HVLT-R retention scores in NA females (model: $F_{(3,11)}=11.64$, $p=0.001$, $R^2=0.760$, adjusted $R^2=0.695$; interaction: $b=-1.944$, $\beta=-1.124$, $t_{(14)}=-5.653$, $p<0.001$, $sr^2=0.696$). Further, high Session 1 cortisol predicted a lower number of choices in aMCI females and a higher number of choices in NA females during trial 1 (model: $F_{(3,11)}=10.2$, $p=0.002$, $R^2=0.735$, adjusted $R^2=0.663$; interaction: $b=-1.527$, $\beta=-$

1.562, $t_{(14)}=-4.413$, $p=0.001$, $sr^2=0.468$) and the delay trial (model: $F_{(3,11)}=11.21$, $p=0.001$, $R^2=0.754$, adjusted $R^2=0.686$; interaction: $b=-1.354$, $\beta=-1.228$, $t_{(14)}=-3.593$, $p=0.004$, $sr^2=0.289$) of the spatial working memory task. In males, high Session 1 cortisol predicted increased associative recognition in aMCI males versus a decline in NA males (model: $F_{(3,12)}=9.723$, $p=0.002$, $R^2=0.709$, adjusted $R^2=0.636$; interaction: $b=0.915$, $\beta=0.658$, $t_{(15)}=2.466$, $p=0.030$, $sr^2=0.148$). High Session 1 cortisol also marginally predicted a lower number of choices in aMCI males versus NA males for spatial working memory during trial 2 (model: $F_{(3,12)}=2.320$, $p=0.127$, $R^2=0.367$, adjusted $R^2=0.209$; interaction: $b=-1.389$, $\beta=-0.856$, $t_{(15)}=-2.177$, $p=0.050$, $sr^2=0.250$) (see Figure 6). Additionally, although no interaction was significant, group was a significant predictor for session 1 HVLT-R immediate recall and delayed recall in females and males and a predictor for face-name associative recognition in females (all $ps<0.05$) (see Table 3).

[INSERT FIGURE 6 HERE]

Session 2 cortisol (AUCg) moderates the relationship between aMCI and perceptions of anxiety in response to the TSST speech in males

We tested the hypothesis that the magnitude of Session 2 cortisol (AUCg) moderates the relationship between participant group (aMCI, NA) and episodic (HVLT), associative (face-name), working memory (spatial working memory), or perceptions of anxiety to the TSST speech during Session 2 in males and females. Regressions revealed that high Session 2 cortisol predicted low perceived TSST speech anxiety ratings in aMCI males versus high perceived TSST speech anxiety ratings in NA males (model: $F_{(3,12)}=7.106$, $p=0.005$, $R^2=0.640$, adjusted $R^2=0.550$; interaction: $b=-1.118$, $\beta=-0.832$, $t_{(15)}=-2.876$, $p=0.014$, $sr^2=0.248$; Figure 7). Similarly in females, the interaction of lower perceived TSST speech anxiety ratings in aMCI females versus NA females with high Session 2 cortisol was close to significant (model: $F_{(3,8)}=8.287$, $p=0.008$, $R^2=0.757$, adjusted $R^2=0.665$; interaction: $b=-1.266$, $\beta=-0.632$, $t_{(11)}=-2.217$, $p=0.057$, $sr^2=0.150$) (see Figure 7). The interaction also approached significance for HVLT-R delayed recall in males such that high Session 2 cortisol predicted increased Session 2 HVLT-R delayed recall in aMCI males versus no change in NA adult males (model: $F_{(3,12)}=22.995$, $p<0.001$, $R^2=0.852$, adjusted $R^2=0.815$; interaction: $b=0.597$, $\beta=0.368$, $t_{(15)}=1.984$, $p=0.071$, $sr^2=0.049$). Additionally, although no interaction was significant, group was a significant predictor for Session 2 HVLT-R immediate recall and delayed recall in females and

males and a predictor for Session 2 HVLT-R retention and face-name associative recognition in males (all $ps < 0.05$) (see Table 4).

[INSERT FIGURE 7 HERE]

Discussion

Cortisol levels were significantly higher in males with aMCI, an effect that was seen during the test sessions but not in diurnal cortisol, suggesting an effect of the test environment to elicit different cortisol responses among aMCI individuals (consistent with data in NA groups from Sindi et al., 2014). Psychosocial stress, as applied by the TSST, improved immediate verbal recall in NA, but not in participants with aMCI, in fact impairing recall in males with aMCI. Furthermore, our data revealed positive correlations between cortisol levels during Session 1 and spatial working memory in females only with opposing directions based on whether the females had aMCI or NA. Higher cortisol was related to better spatial working memory in aMCI females, but worse spatial working memory in NA females during Session 1. Exploratory moderation models revealed that cortisol moderated perceived anxiety during Session 2 and each of memory domains, dependent on sex and session. Cortisol during Session 1 moderated effects on spatial working memory in both sexes and associative recognition in males, with higher cortisol reducing performance in NA but improving performance in aMCI individuals. Cortisol during Session 1 also moderated effects of episodic memory retention in females such that high cortisol enhanced performance in aMCI but impaired performance in NA females. Stress-induced cortisol in Session 2 was associated with decreased perceived anxiety to the speech in males with aMCI but increased perceived anxiety in NA males, whereas in females cortisol moderated the effect on perceived anxiety positively in both groups but a lower anxiety rating with high cortisol in aMCI females than in NA females. While our sample size was small, our results are suggestive of sex differences in the relationship between cortisol in a testing environment and memory and perceived anxiety that depended on whether the participants were NA or had aMCI. These findings, while exploratory, suggest that sex must be considered when exploring relationships between stress biomarkers and memory.

Cortisol levels are higher in males with aMCI

Amnesic MCI males had higher salivary cortisol levels as shown in the first samples of both morning sessions conducted in the laboratory. This finding is partially consistent with past studies that

found increased morning serum cortisol levels in men but not women with AD (Rasmuson et al., 2011) or in salivary cortisol in individuals with MCI (Venero et al, 2013). In another study, salivary cortisol levels upon awakening were significantly higher in the non-amnesic and multidomain type but not in the aMCI compared to NA (Venero et al., 2013). Even though we found higher morning cortisol in aMCI males in the laboratory setting, cortisol levels did not differ between aMCI males and NA from diurnal samples taken at home, consistent with the Venero et al. (2013) study in which participants also took home samples. Rasmuson and colleagues (2011) found increased morning cortisol in males with AD compared to neurotypical participants they had a low number of participants. Nevertheless, it is compelling that in our findings increased morning levels of salivary cortisol are associated with aMCI in males, at least in the laboratory setting. In support of this, higher cortisol and greater variations in cortisol (and correlations with hippocampal volume) are seen in NA older adults when they were tested in an unfamiliar versus familiar environment (Sindi et al., 2014), suggesting that stressful environments influence correlates of memory. Combined with our findings concerning the influence of stress-induced cortisol on anxiety ratings in males and females with aMCI, our results suggest that further investigation into sex differences in cortisol levels is necessary in individuals with aMCI, and perhaps in the laboratory versus home setting.

What might the possible mechanisms be for the cortisol differences and moderation effects between sexes, and greater impairments in males with TSST on memory? Cortisol is only one output of the HPA axis and other outputs may be important to monitor such as alterations to alpha amylase. In addition, here we have captured acute and diurnal cortisol and it would be important to also examine the influence of allostatic load on these results as well as measures of chronic stress (Yan et al., 2018). In addition, HPA is a regulator of immune and metabolic function, which has been implicated as a driver of or in reaction to AD. Thus, other biomarkers such as cytokines (particularly IL-6) and C-reactive protein (CRP) may be fruitful areas of future testing. Indeed, plasma CRP was decreased in AD and MCI of both sexes and CSF IL-16 and IL-8 differed by sex dependent on APOEε4 genotype (Duarte-Guterman, Inkster, Albert, Barha, Robinson, Galea, 2020). There is a paucity of studies on aging, immune, and other biomarkers that have been implicated in AD with sex as a factor that has elicited calls for action (Mielke et al., 2018) and are findings provide further fodder for these calls.

Stress-induced increases in cortisol were associated with enhanced episodic memory in NA but not in aMCI

As expected, the TSST induced an increase in cortisol levels, which was associated with enhanced episodic memory (verbal recall) in the HVLT-R in NA but not in aMCI participants. These findings are consistent with a study by Wolf et al. (2002), which found that there were negative correlations between average cortisol and immediate recall of paragraphs in MCI participants but not in NA. Similarly, neurotypical older adults have been found to exhibit a positive correlation between high cortisol and memory performance, whereas aMCI subjects exhibit a negative correlation (de Souza-Talarico et al., 2010). Collectively, the present data and previous findings suggest that cortisol has opposing relationships with memory and recall in MCI versus normal aging that may differ in magnitude and direction by sex. Intriguingly we also found that aMCI males did not mount a stress-induced increase in cortisol with the TSST unlike the stress-induced increase in cortisol in the females with aMCI. These results are intriguing given that de Souza-Talarico et al. (2020) found that greater cortisol reactivity in the TSST was related to cognitive decline characteristic of future MCI after five years and their population was 80% female. This may explain why females with aMCI transition to AD at a greater rate than males with aMCI. The ability of the HPA axis to mount an appropriate stress response may be an important biomarker for AD transition with differences depending on sex.

Sex may influence the effects stress-induced cortisol on memory in aMCI

In a few of our findings, there were opposing effects of cortisol associations or effects of stress between aMCI and NA dependent on sex. This is intriguing and suggests that sex should be considered in future studies and research on age-related cognitive impairment. This is of particular relevance considering that a number of studies investigating memory and cortisol have had an imbalance of males or females in their test groups (e.g. de Souza-Talarico et al., 2010; de Souza-Talarico et al., 2020; Wolf et al., 2002). Furthermore, inconsistencies in the literature around the effects of acute social stress on memory likely depend on multiple factors including sex and age (for review see Hidalgo et al., 2019; Yan et al., 2018). However, due to a limited sample size in the current work, our sex-based analyses are exploratory and due caution should be paid when generalizing the results.

Our findings of sex differences are congruent with previous studies demonstrating epidemiological, symptomatic, and physiological differences between males and females with MCI. The prevalence of MCI has been found to be greater in males than females, with aMCI as the most common type (Petersen et al., 2010). Furthermore, the incidence of MCI is greater in males than in females (Roberts et al., 2012) and recent studies have uncovered sex-specific risk factors for MCI to AD progression (Kim et al., 2015). Although MCI is more prevalent in males, females exhibit faster

deterioration in cognitive and functional measures over time (Lin et al., 2015). Sex differences are also evident in cognitive and neurophysiological decline with AD, as females experience accelerated hippocampal atrophy and cognitive decline with AD (Ardekani et al., 2016; Irvine et al., 2012). These findings, in accord with our data, emphasise the need to account for sex differences in future research in memory and cognition. Intriguingly, decreases in hippocampal volume predict progression to probable AD (and MCI) in women, whereas increases in white matter hyperintensities in men predict progression to MCI (Burke et al., 2019). Optimistically, there is preliminary evidence that cognitive training in those with aMCI is more effective in women than men (Rahe et al., 2015).

The relationship between cortisol and memory may depend on brain health

Higher cortisol levels should not always be thought of as detrimental to brain function, particularly in the face of acute stress (McEwen, 2019). Indeed, in the present study we found enhanced delayed word-list retention on the HVLT-R in the female NA participants with high cortisol levels in Session 1 under the curve. However, the opposite relationship was found in females with aMCI, as higher cortisol under the curve was associated with reduced retention in the exploratory moderating analyses. These findings are consistent with at least two other studies (de Souza-Talarico et al., 2010; Wolf et al., 2002). De Souza-Talarico et al. (2010) showed a positive relationship between cortisol and delayed recall in NA but a negative relationship in people with MCI. Furthermore, Wolf et al. (2002) showed a negative correlation between average cortisol and immediate story recall in MCI participants but no relationship to average cortisol in NA. It is important to note that opposite patterns were seen in spatial working memory and associative memory, with high Session 1 cortisol associated with improved spatial working memory in aMCI males and females and improved associative recognition in aMCI males but impairments in both associative and spatial working memory in NA participants. During Session 2, high cortisol was associated with enhanced episodic memory (HVLT-R delayed recall) in aMCI males versus NA males, but this interaction had a low effect size. These findings in aMCI participants are in contrast to past findings of impaired cognitive performance in aMCI participants with high cortisol (Popp et al., 2015). However, it should be noted that those studies are based on high diurnal cortisol in aMCI participants and saliva samples were not collected on the same days that cognitive tasks were performed (Popp et al., 2015). Cortisol measures used in our moderation models were based on saliva samples taken in the laboratory throughout the days of Session 1 or Session 2. The impact of cortisol on memory task performance may change when aMCI participants are brought into the laboratory setting. Opposite relationships of cortisol to memory may also reflect

brain regions recruited during the task as spatial working memory heavily recruits the prefrontal cortex, episodic memory the medial temporal prefrontal cortex and hippocampus, and associative memory the entorhinal cortex (Courtney et al., 1998; Eichenbaum et al., 2017). Thus, our findings suggest that HPA function may be having opposing effects on memory performance in aMCI groups compared to normal cognitive aging groups.

Acute psychosocial stress improved immediate episodic recall in NA but not in aMCI participants. Although cortisol was not a moderating factor on episodic memory in Session 2, this may be due to a number of factors. Stress activates the HPA axis and it is possible had we measured more timepoints of salivary cortisol we may have seen a moderating effect. In addition, it is important to consider that other biomarkers of HPA activation such as corticotropin releasing hormone (CRH), adrenocorticotrophic hormone (ACTH), or sympathetic activation via the sympathetic-adrenal-medullary system (SAM: alpha amylase, epinephrine, heart rate variability) may have a moderating effect on memory with acute stress. Certainly, it is intriguing that the TSST had an adverse effect on episodic memory in aMCI but a positive effect in NA participants. Indeed, other research has found enhancing effects of the TSST on memory, depending on when TSST was administered relative to memory testing (encoding, retention, recall) that depends on age and sex (Hidalgo et al., 2019). Enhancing effects on episodic memory are seen in older (middle-aged) NA women with TSST (Almela et al., 2011). Others have found no effect of cortisol during TSST to moderate working memory in older NA individuals (Pulopulos et al., 2015), consistent in part to our findings that TSST did not influence spatial working memory in the present study. Furthermore, there are well known sex differences in the effects of stress in animal models (Goel et al., 2014). However, the effects of age and stress on learning are not as well studied. In light of these findings, we encourage the research community to make it a priority to examine sex as a factor in analyses of aging and cognition.

Stress-induced cortisol in Session 2 was associated with greater ratings of anxiety in response to the TSST speech in all groups except aMCI males which were associated with reduced anxiety

Our models revealed that Session 2 cortisol moderated the effect of aMCI on perception of anxiety for the TSST speech in both male and female participants. Greater Session 2 increases in cortisol, the session that involved stress exposure, was associated with greater perceptions of anxiety in NA males and females and to a lesser extent in aMCI females. This finding is similar to previous studies that have found higher anxiety scores in healthy male and female participants exposed to stressors (Ellenbogen et al., 2002). This relationship did not hold for males with aMCI for whom higher

levels of Session 2 cortisol were associated with reduced perceived anxiety in our study. Anxiety is found in a high percentage of patients with MCI, subjective cognitive decline, and AD (Banning et al., 2020), but a previous study by Guerdoux-Ninot and Trouillet (2019) found lower perceived stress in response to a Stroop test in male and female AD and aMCI participants compared to NA as the task became more effortful. In the present study, a large percentage of aMCI males did not characterize the TSST speech as anxiety provoking. A blunted perceived anxiety response to stress in aMCI males compared to NA males in the present study could be related to previous findings of increased apathy in male and female AD and aMCI participants compared to nonamnesic-MCI (naMCI) and NA participants (Ellison et al., 2008; Lanctôt et al., 2017). Indeed, past data has indicated that AD and MCI patients differ in the prevalence of symptoms of apathy with a greater prevalence in AD versus MCI groups (Siafarikas et al., 2018). However, these past studies did not analyse their males and females separately and could have missed effects of increased apathy driven by aMCI males. Moreover, these past studies did not examine the effects of cortisol levels on perceived stress or anxiety in aMCI groups. Nevertheless, our findings along with past findings suggest that, whereas increasing stress-induced cortisol is associated with increasing perceptions of anxiety in NA, male aMCI participants are less likely to perceive themselves as being anxious or stressed.

Limitations

This is an exploratory study given our low sample size and needs to be replicated with a larger population to examine sex-specific effects. Indeed, diurnal cortisol findings in aMCI females were missed because of a lack of samples. Missing diurnal data in aMCI females may reflect sex differences in partner support, whereby partners of females with aMCI may be less predisposed to assisting with remembering to engage in saliva sample collection than partners of males with aMCI might have been. In future, it may be important to measure perceived primary support and partner attitudes to ascertain why samples are missed. In the present study, we had only 2 aMCI females that completed all (or more than 75%) of the diurnal saliva sampling. However, when comparing these two females to the rest of the aMCI group with incomplete diurnal samples, they did not differ in age, education, TSST anxiety ratings, or on episodic or associative memory tasks. In addition to low sample size, and as previously mentioned, taking saliva samples at different timepoints and analysing samples for HPA activation biomarkers other than cortisol (ACTH, CRH, alpha amylase, epinephrine, heart rate variability) may have resulted in different moderating effects of aMCI on memory tasks. Nevertheless, even though cortisol is only one biomarker of stress our results did resemble the findings of other studies that

examined the effects of stress on memory function in aMCI versus NA participants. It should be noted that while linear trend lines had the best fit for our data we also attempted polynomial regressions (quadratic, cubic) for associations between cortisol and memory task performances specific to each session in aMCI versus NA and in male versus female participants. In addition to linear associations, some quadratic and cubic associations were also observed but the majority appeared to be driven by outliers. In a future study, a larger sample size in each group would help elucidate whether the association between cortisol and certain memory performance may be non-linear depending on aMCI and/or sex. All in all, a larger sample size would help account for some of the losses in home cortisol sampling and improve the power of our statistics to examine the influence of multiple other factors on our data.

A larger sample size would also permit examination of possible aMCI phenotypes. Other researchers have demonstrated heterogeneity within the aMCI subtype on memory performance measures that may be predictive of progression to AD type dementia (e.g., Sanborn et al., 2017). We chose to focus on the aMCI subtype in order to limit the possible influence of heterogeneous underlying neuropathologies to that of incipient AD. Widening the lens to include other MCI subtypes that are more likely to be of mixed etiology may produce different relationships between cortisol and memory than those demonstrated here. For example, as previously described, Guerdoux-Ninot and Trouillet (2019) found that effort increased perceived stress in NA adults and non-amnesic MCI patients but reduced it in aMCI and AD. Thus, it would be interesting to examine larger sample sizes that include MCI subtypes in a future study in order to further appreciate differences between subtypes and to further explore research demonstrating MCI subtypes may have different phenotypes (Edmonds et al., 2019; Sanborn et al., 2017). Certainly, our data demonstrate it would be important to consider the heterogeneity of aMCI and to determine whether biological sex may be a contributing factor to the heterogeneity.

Conclusions

The present study found relationships with cortisol (stress-induced and morning session cortisol) and aMCI and moderating effects of cortisol on some domains of memory. As expected, the aMCI participants performed more poorly across the memory measures as compared to NA; however within this we found response patterns influenced by biological sex. We found that males with aMCI had higher cortisol levels in the morning during the test sessions. We also found stress-induced impairment with episodic memory only in males with aMCI. Although our sex-based analyses are

exploratory due to the low sample size, sex differences were nonetheless observed. It is critical that future studies explore sex as a biological variable as we have presented evidence herein suggesting that effects at the confluence of aMCI and stress can be obfuscated or otherwise eliminated when males and females are combined instead of being considered separately. For real understanding and advancement to take place in this field, biological sex must be considered and statistically analyzed.

Estimates of the prevalence of MCI in the elderly show high variability, ranging from ~3-42% (Ward et al., 2012), due to differences in study methodology, especially with regards to the sample population (age, ethnicity, education-level, etc.) (Ward et al., 2012). Regardless, there is a health care burden associated with MCI (Ton et al., 2017) as those with MCI are more likely to develop AD (Busse et al., 2003; Lupien et al., 1998) and the health care burden of AD is more severe than that of MCI (Ton et al., 2017). The findings presented here indicate future studies should make examining sex differences (their nature, underlying mechanisms, outcomes, etc.) in aMCI a priority, as well as expand upon the influence of cortisol in aMCI and the interactions between these factors.

Acknowledgments

We gratefully acknowledge the participants for generously volunteering their time and effort. We thank Triti Namiranian, Angelina Polsinelli, Nicole D'Souza, Diana Smith, Preeyam Parikha, and Tallinn Splinter for assistance with data collection and management and Elizabeth Perez for editorial assistance.

Funding and Disclosure of Interest

This study was supported by grants from the Canadian Institutes of Health Research (CIHR-IA #131486 to LAMG, KJM, and AKT) and by the Morris Goldenberg Medical Research Endowment (to KJM). The funders had no role in the study design, collection, analyses or interpretation of the data. The authors have no further conflicts to disclose.

References

- Adam, E. K., Quinn, M. E., Tavernier, R., McQuillan, M. T., Dahlke, K. A., & Gilbert, K. E. (2017). Diurnal cortisol slopes and mental and physical health outcomes: A systematic review and meta-analysis. *Psychoneuroendocrinology*, 83, 25-41. doi:10.1016/j.psyneuen.2017.05.018
- Albert, M. S., DeKosky, S. T., Dickson, D., Dubois, B., Feldman, H. H., Fox, N. C., Gamst, A., Holtzman, D. M., Jagust, W. J., Petersen, R. C., Snyder, P. J., Carrillo, M. C., Thies, B., & Phelps, C. H. (2011). The diagnosis of mild cognitive impairment due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimer's & Dementia*, 7(3), 270–279. <https://doi.org/10.1016/j.jalz.2011.03.008>
- Aggarwal, N. T., Wilson, R. S., Beck, T. L., Rajan, K. B., Mendes de Leon, C. F., Evans, D. A., & Everson-Rose, S. A. (2014). Perceived stress and change in cognitive function among adults 65 years and older. *Psychosomatic Medicine*, 76(1), 80–85. doi: 10.1097/PSY.000000000000016
- Almela, M., Hidalgo, V., Villada, C., Espín, L., Gómez-Amor, J., & Salvador, A. (2011). The impact of cortisol reactivity to acute stress on memory: Sex differences in middle-aged people. *Stress (Amsterdam, Netherlands)*, 14(2), 117–127
- Ardekani, B. A., Convit, A., & Bachman, A. H. (2016). Analysis of the MIRIAD data shows Sex differences in hippocampal atrophy progression. *Journal of Alzheimer's Disease: JAD*, 50(3), 847–857. <https://doi.org/10.3233/JAD-150780>
- Banning, L. C., Ramakers, I. H., Köhler, S., Bron, E. E., Verhey, F. R., de Deyn, P. P., Claassen, J. A. H. R., Koek, H. L., Middelkoop, H. A. M., van der Flier, W. V., van der Lugt, A., Aalten, P., Alzheimer's Disease Neuroimaging Initiative, & Parelsnoer Institute Neurodegenerative Diseases study group. (2020). The association between biomarkers and neuropsychiatric symptoms across the Alzheimer's disease spectrum. *The American Journal of Geriatric Psychiatry*, 28(7), 735-744. <https://doi.org/10.1016/j.jagp.2020.01.012>
- Beluche, I., Carrière, I., Ritchie, K., & Ancelin, M. L. (2010). A prospective study of diurnal cortisol and cognitive function in community-dwelling elderly people. *Psychological Medicine*, 40(6), 1039–1049. <https://doi.org/10.1017/S0033291709991103>
- Burke, S. L., Hu, T., Fava, N. M., Li, T., Rodriguez, M. J., Schuldiner, K. L., Burgess, A., & Laird, A. (2019). Sex differences in the development of mild cognitive impairment and probable Alzheimer's disease as predicted by hippocampal volume or white matter hyperintensities. *Journal of Women & Aging*, 31(2), 140–164. <https://doi.org/10.1080/08952841.2018.1419476>

- 746 Busse, A., Bischkopf, J., Riedel-Heller, S. G., & Angermeyer, M. C. (2003). Mild cognitive
747 impairment: prevalence and incidence according to different diagnostic criteria. Results of the
748 Leipzig Longitudinal Study of the Aged (LEILA75+). *The British Journal of Psychiatry: The*
749 *Journal of Mental Science*, 182, 449–454.
- 750 Büttler, R. M., Bagci, E., Brand, H. S., Heijer, M. den, Blankenstein, M. A., & Heijboer, A. C. (2018).
751 Testosterone, androstenedione, cortisol and cortisone levels in human unstimulated, stimulated
752 and parotid saliva. *Steroids*, 138, 26–34. <https://doi.org/10.1016/j.steroids.2018.05.013>
- 753 Courtney, S. M., Petit, L., Maisog, J. M., Ungerleider, L. G., & Haxby, J. V. (1998). An area
754 specialized for spatial working memory in human frontal cortex. *Science*, 279(5355), 1347–
755 1351. <https://doi.org/10.1126/science.279.5355.1347>
- 756 Cullum, S., Huppert, F. A., McGee, M., Dening, T., Ahmed, A., Paykel, E. S., & Brayne, C. (2000).
757 Decline across different domains of cognitive function in normal ageing: Results of a
758 longitudinal population-based study using CAMCOG. *International Journal of Geriatric*
759 *Psychiatry*, 15(9), 853–862.
- 760 Dedovic, K., Duchesne, A., Andrews, J., Engert, V., & Pruessner, J. C. (2009). The brain and the stress
761 axis: The neural correlates of cortisol regulation in response to stress. *NeuroImage*, 47(3), 864–
762 871. <https://doi.org/10.1016/j.neuroimage.2009.05.074>
- 763 Delis D, Kaplan E, & Kramer JH. *Delis-Kaplan Executive Function System™*. San Antonio:
764 Psychological Corporation; 2001.
- 765 de Souza-Talarico, J. N., Alves, A. R., Brucki, S., Nitrini, R., Lupien, S. J., & Suchecki, D. (2020).
766 Cortisol reactivity to a psychosocial stressor significantly increases the risk of developing
767 Cognitive Impairment no Dementia five years later. *Psychoneuroendocrinology*, 115, 104601.
768 <https://doi.org/10.1016/j.psyneuen.2020.104601>
- 769 de Souza-Talarico, J. N., Chaves, E. C., Lupien, S. J., Nitrini, R., & Caramelli, P. (2010). Relationship
770 between cortisol levels and memory performance may be modulated by the presence or absence
771 of cognitive impairment: Evidence from healthy elderly, mild cognitive impairment and
772 Alzheimer’s disease subjects. *Journal of Alzheimer’s Disease: JAD*, 19(3), 839–848.
773 <https://doi.org/10.3233/JAD-2010-1282>
- 774 Dijckmans, B., Tortosa-Martínez, J., Caus, N., González-Caballero, G., Martínez-Pelegrin, B.,
775 Manchado-Lopez, C., Cortell-Tormo, J. M., Chulvi-Medrano, I., & Clow, A. (2017). Does the
776 diurnal cycle of cortisol explain the relationship between physical performance and cognitive

777 function in older adults?. *European review of aging and physical activity : official journal of the*
778 *European Group for Research into Elderly and Physical Activity*, 14, 6..

779 Doecke, J. D., Laws, S. M., Faux, N. G., Wilson, W., Burnham S. C., Lam, C-P., Mondal, A., Bedo, J.,
780 Bush, A. I., Brown, B., De Ruyck, K., Ellis, K. A., Fowler, C., Gupta, V. B., Head, R.,
781 Macaulay, S. L., Pertile, K., Rowe, C. C., Rembach, A., ... Australian Imaging Biomarker and
782 Lifestyle Research Group. (2012). Blood-based protein biomarkers for diagnosis of Alzheimer
783 disease. *Archives of Neurology*, 69(10), 1318-1325. doi:10.1001/archneurol.2012.1282

784 Donaldson, W. (1992). Measuring recognition memory. *Journal of Experimental Psychology: General*,
785 121(3), 275–277. <https://doi.org/10.1037/0096-3445.121.3.275>

786 Duarte-Guterman, P., Inkster A. M., Albert A. Y., Barha, C. K., Robinson, W. P., Galea L. A. M., &
787 Alzheimer's Disease Neuroimaging Initiative. (2020). Inflammation and epigenetic age in
788 Alzheimer's disease: Do sex and APOE matter? *bioRxiv*, 741777v4.
789 <https://doi.org/10.1101/741777>

790 Duff, S. J., & Hampson, E. (2001). A sex difference on a novel spatial working memory task in
791 humans. *Brain and Cognition*, 47(3), 470–493. <https://doi.org/10.1006/brcg.2001.1326>

792 Edmonds, E. C., McDonald, C. R., Marshall, A., Thomas, K. R., Eppig, J., Weigand, A. J., Delano-
793 Wood, L., Galasko, D. R., Salmon, D. P., Bondi, M. W., & Alzheimer's Disease Neuroimaging
794 Initiative. (2019). Early versus late MCI: Improved MCI staging using a neuropsychological
795 approach. *Alzheimer's & Dementia*, 15(5), 699-708. doi:10.1016/j.jalz.2018.12.009

796 Eichenbaum, H. (2017). Prefrontal-hippocampal interactions in episodic memory. *Nature Reviews*
797 *Neuroscience*, 18(9), 547–558. <https://doi.org/10.1038/nrn.2017.74>

798 Ellenbogen, M. A., Schwartzman, A. E., Stewart, J., & Walker, C. D. (2002). Stress and selective
799 attention: The interplay of mood, cortisol levels, and emotional information processing.
800 *Psychophysiology*, 39(6), 723-732.

801 Ellison, J. M., Harper, D. G., Berlow, Y., & Zeranski, L. (2008). Beyond the “C” in MCI: noncognitive
802 symptoms in amnesic and non-amnesic mild cognitive impairment. *CNS Spectrums*, 13(1),
803 66–72. <https://doi.org/10.1017/s1092852900016175>

804 Elosúa, M. R., & Contreras, M. J. (2017). Gender differences in verbal and visuospatial working
805 memory tasks in patients with mild cognitive impairment and Alzheimer disease. *Dementia and*
806 *Geriatric Cognitive Disorders Extra*, 7(1), 101-108.

807 Ferrari, E., Cravello, L., Muzzoni, B., Casarotti, D., Paltro, M., Solerte, S. B., Fioravanti, M., Cuzzoni,
808 G., Pontiggia, B., & Magri, F. (2001). Age-related changes of the hypothalamic-pituitary-

809 adrenal axis: Pathophysiological correlates. *European Journal of Endocrinology*, 144(4), 319-
810 329. doi:10.1530/eje.0.1440319.

811 Gale, S. D., Baxter, L., & Thompson, J. (2016). Greater memory impairment in dementing females
812 than males relative to sex-matched healthy controls. *Journal of Clinical and Experimental*
813 *Neuropsychology*, 38(5), 527–533. <https://doi.org/10.1080/13803395.2015.1132298>

814 Goel, N., Workman, J. L., Lee, T. T., Innala, L., & Viau, V. (2014). Sex differences in the HPA axis.
815 *Comprehensive Physiology*, 4(3), 1121–1155. <https://doi.org/10.1002/cphy.c130054>

816 Grady, C. (2012) Trends in Neurocognitive Aging. *Nat Rev Neurosci*, 13(7), 491–505.

817 Guerdoux-Ninot, E. & Trouillet, R. (2019). Impact of perceived stress on cognitive performance:
818 Moderating effect of mild cognitive impairment and Alzheimer’s disease. *Journal of Clinical*
819 *and Experimental Neuropsychology*, 41(4), 364-379. doi:10.1080/13803395.2018.1564250

820 Hartmann, A., Veldhuis, J. D., Deuschle, M., Standhardt, H., & Heuser, I. (1997). Twenty-four hour
821 cortisol release profiles in patients with Alzheimer’s and Parkinson’s disease compared to
822 normal controls: Ultradian secretory pulsatility and diurnal variation. *Neurobiology of Aging*,
823 18(3), 285–289.

824 Hidalgo, V., Pulopulos, M. M., & Salvador, A. (2019). Acute psychosocial stress effects on memory
825 performance: Relevance of age and sex. *Neurobiology of Learning and Memory*, 157, 48–60.
826 <https://doi.org/10.1016/j.nlm.2018.11.013>

827 Irvine, K., Laws, K. R., Gale, T. M., & Kondel, T. K. (2012). Greater cognitive deterioration in women
828 than men with Alzheimer’s disease: A meta analysis. *Journal of Clinical and Experimental*
829 *Neuropsychology*, 34(9), 989–998. <https://doi.org/10.1080/13803395.2012.712676>

830 Jack, C. R. Jr., Therneau, T. M., Weigand, S. D., Wiste, H. J., Knopman, D. S., Vemuri, P., Lowe, V.
831 J., Mielke, M. M., Roberts, R. O., Machulda, M. M., Graff-Radford, J., Jones, D. T., Schwarz,
832 C. G., Gunter, J. L., Senjem, M. L., Rocca, W. A., & Petersen, R. C. (2019). Prevalence of
833 biologically vs clinically defined Alzheimer spectrum entities using the National Institute on
834 Aging–Alzheimer’s Association research framework. *JAMA Neurology*, 76(10), 1174-1183.
835 <https://doi.org/10.1001/jamaneurol.2019.1971>

836 Justice N. J. (2018). The relationship between stress and Alzheimer's disease. *Neurobiology of stress*,
837 8, 127–133. <https://doi-org.ezproxy.library.ubc.ca/10.1016/j.ynstr.2018.04.002>

838 Kim, S., Kim, M. J., Kim, S., Kang, H. S., Lim, S. W., Myung, W., Lee, Y., Hong, C. H., Choi S. H.,
839 Na, D. L., Seo, S. W., Ku, B. D., Kim, S. Y., Kim, S. Y., Jeong, J. H., Park, S. A., Carroll, B. J.,
840 & Kim, D. K. (2015). Gender differences in risk factors for transition from mild cognitive

841 impairment to Alzheimer's disease: A CREDOS study. *Comprehensive Psychiatry*, 62, 114–
842 122. <https://doi.org/10.1016/j.comppsy.2015.07.002>

843 Kirschbaum, C., Pirke, K. M., & Hellhammer, D. H. (1993). The 'Trier Social Stress Test'--A tool for
844 investigating psychobiological stress responses in a laboratory setting. *Neuropsychobiology*,
845 28(1–2), 76–81. <https://doi.org/10.1159/000119004>

846 Koran, M. E. I., Wagener, M., Hohman, T. J., & Alzheimer's Neuroimaging Initiative. (2017). Sex
847 differences in the association between AD biomarkers and cognitive decline. *Brain Imaging*
848 *and Behavior*, 11(1), 205–213. <https://doi.org/10.1007/s11682-016-9523-8>

849 Kudielka, B. M., & Kirschbaum, C. (2005). Sex differences in HPA axis responses to stress: A review.
850 *Biological Psychology*, 69(1), 113–132. <https://doi.org/10.1016/j.biopsycho.2004.11.009>

851 Lanctôt, K. L., Agüera-Ortiz, L., Brodaty, H., Francis, P. T., Geda, Y. E., Ismail, Z., Marshall, G. A.,
852 Mortby, M. E., Onyike, C. U., Padala, P. R., Politis, A. M., Rosenberg, P. B., Siegel, E.,
853 Sultzer, D. L., & Abraham, E. H. (2017). Apathy associated with neurocognitive disorders:
854 recent progress and future directions. *Alzheimer's & Dementia*, 13(1), 84–100.

855 Laske C., Leyhe T., Stransky E., Hoffmann N., Fallgatter A. J., & Dietzsch J., (2011). Identification of
856 a blood-based biomarker panel for classification of Alzheimer's disease. *International Journal*
857 *of Neuropsychopharmacology*, 14(9), 1147–1155.

858 Laughlin, G. A., & Barrett-Connor, E. (2000). Sexual dimorphism in the influence of advanced aging
859 on adrenal hormone levels: The Rancho Bernardo Study. *The Journal of Clinical*
860 *Endocrinology and Metabolism*, 85(10), 3561–3568. <https://doi.org/10.1210/jcem.85.10.6861>

861 Lin, K. A., Choudhury, K. R., Rathakrishnan, B. G., Marks, D. M., Petrella, J. R., Doraiswamy, P. M.,
862 & Alzheimer's Disease Neuroimaging Initiative. (2015). Marked gender differences in
863 progression of mild cognitive impairment over 8 years. *Alzheimer's & Dementia (New York, N.*
864 *Y.)*, 1(2), 103–110. <https://doi.org/10.1016/j.trci.2015.07.001>

865 Lupien, S. J., de Leon, M., de Santi, S., Convit, A., Tarshish, C., Nair, N. P., Thakur, M., McEwen, B.
866 S., Hauger, R. L., & Meaney, M. J. (1998). Cortisol levels during human aging predict
867 hippocampal atrophy and memory deficits. *Nature Neuroscience*, 1(1), 69–73.
868 <https://doi.org/10.1038/271>

869 Mah, L., Binns, M. A., Steffens, D. C., & Alzheimer's Disease Neuroimaging Initiative. (2015).
870 Anxiety symptoms in amnesic mild cognitive impairment are associated with medial temporal
871 atrophy and predict conversion to Alzheimer disease. *The American Journal of Geriatric*
872 *Psychiatry*, 23(5), 466–476.

- 873 Mayes, A. R., Holdstock, J. S., Isaac, C. L., Montaldi, D., Grigor, J., Gummer, A., Cariga, P., Downes,
874 J. J., Tsivilis, D., Gaffan, D., Gong, Q., & Norman, K. A. (2004). Associative recognition in a
875 patient with selective hippocampal lesions and relatively normal item recognition.
876 *Hippocampus*, 14(6), 763–784. <https://doi.org/10.1002/hipo.10211>
- 877 McEwen B. S. (2019). The good side of "stress". *Stress (Amsterdam, Netherlands)*, 22(5), 524–525.
878 <https://doi.org/10.1080/10253890.2019.1631794>
- 879 Mielke, M. M., Ferretti, M. T., Iulita, M. F., Hayden, K., & Khachaturian, A. S. (2018). Sex and gender
880 in Alzheimer's disease - Does it matter? *Alzheimer's & Dementia*, 14(9), 1101–1103.
- 881 Mielke, M. M., Vemuri, P., & Rocca, W. A. (2014). Clinical epidemiology of Alzheimer's disease:
882 Assessing sex and gender differences. *Clinical Epidemiology*, 6, 37–48.
883 <https://doi.org/10.2147/CLEP.S37929>
- 884 Nebel, R. A., Aggarwal, N. T., Barnes, L. L., Gallagher, A., Goldstein, J. M., Kantarci, K.,
885 Mallampalli, M. P., Mormino, E. C., Scott, L., Yu, W. H., Maki, P. M., & Mielke, M. M.
886 (2018). Understanding the impact of sex and gender in Alzheimer's disease: A call to action.
887 *Alzheimer's & Dementia*, 14(9), 1171–1183. <https://doi.org/10.1016/j.jalz.2018.04.008>
- 888 Norman, R. M. G., Gawronski, B., Hampson, E., Sorrentino, R. M., Szeto, A., & Ye, Y. (2010).
889 Physical proximity in anticipation of meeting someone with schizophrenia: The role of explicit
890 evaluations, implicit evaluations and cortisol levels. *Schizophrenia Research*, 124(1–3), 74–80.
891 <https://doi.org/10.1016/j.schres.2010.07.021>
- 892 Otte, C., Hart, S., Neylan, T. C., Marmar, C. R., Yaffe, K., & Mohr, D. C. (2005). A meta-analysis of
893 cortisol response to challenge in human aging: importance of gender.
894 *Psychoneuroendocrinology*, 30(1), 80–91. <https://doi.org/10.1016/j.psychneuen.2004.06.002>
- 895 Petersen, R. C. (2004). Mild cognitive impairment as a diagnostic entity. *Journal of Internal Medicine*,
896 256, 183-194.
- 897 Petersen, R. C., Roberts, R. O., Knopman, D. S., Geda, Y. E., Cha, R. H., Pankratz, V. S., Boeve, B. F.,
898 Tangalos, E. G., Ivnik, R. J., & Rocca, W. A. (2010). Prevalence of mild cognitive impairment
899 is higher in men. The Mayo Clinic Study of Aging. *Neurology*, 75(10), 889–897.
900 <https://doi.org/10.1212/WNL.0b013e3181f11d85>
- 901 Popp, J., Wolfsgrubner, S., Heuser, I., Peters, O., Hüll, M., Schröder, J., Möller, H-J., Lewczuk, P.,
902 Schneider, A., Jahn, H., Luckhaus, C., Perneczky, R., Frölich, L., Wagner, M., Maier, W.,
903 Wiltfang, J., Kornhuber, J., & Jessen, F. (2015). Cerebrospinal fluid cortisol and clinical disease

- 904 progression in MCI and dementia of Alzheimer's type. *Neurobiology of Aging*, 36(2), 601–607.
- 905 <https://doi.org/10.1016/j.neurobiolaging.2014.10.031>
- 906 Pruessner, J. C., Kirschbaum, C., Meinlschmid, G., & Hellhammer, D. H. (2003). Two formulas for
- 907 computation of the area under the curve represent measures of total hormone concentration
- 908 versus time-dependent change. *Psychoneuroendocrinology*, 28(7), 916–931.
- 909 Pulpulos, M. M., Hidalgo, V., Almela, M., Puig-Perez, S., Villada, C., & Salvador, A. (2015). Acute
- 910 stress and working memory in older people. *Stress*, 18(2), 178-187.
- 911 Rahe, J., Liesk, J., Rosen, J. B., Petrelli, A., Kaesberg, S., Onur, O. A., Kessler, J., Fink, G. R., &
- 912 Kalbe, E. (2015). Sex differences in cognitive training effects of patients with amnesic mild
- 913 cognitive impairment. *Neuropsychology, Development, and Cognition. Section B, Aging,*
- 914 *Neuropsychology and Cognition*, 22(5), 620–638.
- 915 <https://doi.org/10.1080/13825585.2015.1028883>
- 916 Rasmuson, S., Näsman, B., & Olsson, T. (2011). Increased serum levels of dehydroepiandrosterone
- 917 (DHEA) and interleukin-6 (IL-6) in women with mild to moderate Alzheimer's disease.
- 918 *International Psychogeriatrics*, 23(9), 1386-1392. doi:10.1017/S1041610211000810
- 919 Raz, N. & Rodrigue, K. M. (2006). Differential aging of the brain: patterns, cognitive correlates and
- 920 modifiers. *Neuroscience and Biobehavioral Reviews*, 30(6), 730–748.
- 921 <https://doi.org/10.1016/j.neubiorev.2006.07.001>
- 922 Roberts, R. O., Geda, Y. E., Knopman, D. S., Cha, R. H., Pankratz, V. S., Boeve, B. F., Tangalos, E.
- 923 G., Ivnik, R. J., Rocca, W. A., & Petersen, R. C. (2012). The incidence of MCI differs by
- 924 subtype and is higher in men: The Mayo Clinic Study of Aging. *Neurology*, 78(5), 342–351.
- 925 <https://doi.org/10.1212/WNL.0b013e3182452862>
- 926 Sanborn V, Putcha D, & Tremont, G. (2017). Correlates of recognition memory performance in
- 927 amnesic mild cognitive impairment. *Journal of Clinical and Experimental Neuropsychology*,
- 928 40(2), 205-211. doi:10.1080/13803395.2017.1334043
- 929 Shi, F., Liu, B., Zhou, Y., Yu, C., & Jiang, T. (2009). Hippocampal volume and asymmetry in mild
- 930 cognitive impairment and Alzheimer's disease: Meta-analyses of MRI studies. *Hippocampus*,
- 931 19(11), 1055–1064. <https://doi.org/10.1002/hipo.20573>
- 932 Siafarikas, N., Selbaek, G., Fladby, T., Benth, J. Š., Auning, E., & Aarsland, D. (2018). Frequency and
- 933 subgroups of neuropsychiatric symptoms in mild cognitive impairment and different stages of
- 934 dementia in Alzheimer's disease. *International Psychogeriatrics*, 30(1), 103–113.

- 935 Sindi, S., Fiocco, A. J., Juster, R. P., Lord, C., Pruessner, J., & Lupien, S. J. (2014). Now you see it,
936 now you don't: Testing environments modulate the association between hippocampal volume
937 and cortisol levels in young and older adults. *Hippocampus*, 24(12), 1623–1632.
- 938 Spreen O, Strauss E. *A Compendium of Neuropsychological Tests: Administration, Norms, and*
939 *Commentary, 2nd Ed.* New York, NY, US: Oxford University Press; 1998.
- 940 Sohn, D., Shpanskaya, K., Lucas, J. E., Petrella, J. R., Saykin, A. J., Tanzi, R. E., Samatova, N. F., &
941 Doraiswamy, P. M. (2018). Sex differences in cognitive decline in subjects with high likelihood
942 of mild cognitive impairment due to Alzheimer's disease. *Scientific Reports*, 8(1), 7490.
943 <https://doi.org/10.1038/s41598-018-25377-w>
- 944 Spreen, O., & Strauss, E. (1998). *A compendium of neuropsychological tests: Administration, norms,*
945 *and commentary* (2nd ed.). New York: Oxford University Press.
- 946 Sundermann, E. E., Biegon, A., Rubin, L. H., Lipton, R. B., Landau, S., Maki, P. M., & Alzheimer's
947 Disease Neuroimaging Initiative. (2017). Does the female advantage in verbal memory
948 contribute to underestimating Alzheimer's Disease pathology in women versus men? *Journal of*
949 *Alzheimer's Disease: JAD*, 56(3), 947–957. <https://doi.org/10.3233/JAD-160716>
- 950 Ton, T. G. N., DeLeire, T., May, S. G., Hou, N., Tebeka, M. G., Chen, E., & Chodosh, J. (2017). The
951 financial burden and health care utilization patterns associated with amnesic mild cognitive
952 impairment. *Alzheimer's & Dementia*, 13(3), 217–224.
953 <https://doi.org/10.1016/j.jalz.2016.08.009>
- 954 Tortosa-Martínez, J., Manchado, C., Cortell-Tormo, J. M., & Chulvi-Medrano, I. (2018). Exercise, the
955 diurnal cycle of cortisol and cognitive impairment in older adults. *Neurobiology of stress*, 9,
956 40–47. <https://doi-org.ezproxy.library.ubc.ca/10.1016/j.ynstr.2018.08.004>
- 957 Troyer, A. K., D'Souza, N. A., Vandermorris, S., & Murphy, K. J. (2011). Age-related differences in
958 associative memory depend on the types of associations that are formed. *Neuropsychology,*
959 *Development, and Cognition. Section B, Aging, Neuropsychology and Cognition*, 18(3), 340–
960 352. <https://doi.org/10.1080/13825585.2011.553273>
- 961 Troyer, A. K., Murphy, K. J., Anderson, N. D., Craik, F. I. M., Moscovitch, M., Maione, A., & Gao, F.
962 (2012). Associative recognition in mild cognitive impairment: Relationship to hippocampal
963 volume and apolipoprotein E. *Neuropsychologia*, 50(14), 3721–3728.
964 <https://doi.org/10.1016/j.neuropsychologia.2012.10.018>
- 965 Troyer, A. K., Murphy, K. J., Anderson, N. D., Hayman-Abello, B. A., Craik, F. I. M., & Moscovitch,
966 M. (2008). Item and associative memory in amnesic mild cognitive impairment: Performance

967 on standardized memory tests. *Neuropsychology*, 22(1), 10–16. [https://doi.org/10.1037/0894-](https://doi.org/10.1037/0894-4105.22.1.10)
968 4105.22.1.10

969 Venero, C., Díaz-Mardomingo, C., Pereda-Pérez, I., García-Herranz, S., Utrera, L., Valencia, A., &
970 Peraíta, H. (2013). Increased morning salivary cortisol levels in older adults with nonamnesic
971 and multidomain mild cognitive impairment. *Psychoneuroendocrinology*, 38(4), 488–498.
972 <https://doi.org/10.1016/j.psyneuen.2012.07.007>

973 Wang, L. Y., Raskind, M. A., Wilkinson, C. W., Shofer, J. B., Sikkema, C., Szot, P., Quinn, J. F.,
974 Galasko, D. R., & Peskind, E. R. (2018). Associations between CSF cortisol and CSF
975 norepinephrine in cognitively normal controls and patients with amnesic MCI and AD
976 dementia. *International Journal of Geriatric Psychiatry*, 33(5), 763–768.
977 <https://doi.org/10.1002/gps.4856>

978 Ward, A., Arrighi, H. M., Michels, S., & Cedarbaum, J. M. (2012). Mild cognitive impairment:
979 Disparity of incidence and prevalence estimates. *Alzheimer's & Dementia*, 8(1), 14–21.
980 <https://doi.org/10.1016/j.jalz.2011.01.002>

981 Wolf, O. T., Convit, A., Thorn, E., & de Leon, M. J. (2002). Salivary cortisol day profiles in elderly
982 with mild cognitive impairment. *Psychoneuroendocrinology*, 27(7), 777–789.

983 Yan, Y., Dominguez, S., Fisher, D. W., & Dong, H. (2018). Sex differences in chronic stress responses
984 and Alzheimer's disease. *Neurobiology of stress*, 8, 120–126.
985

986 Table 1 *Descriptive Data for the Participant Groups*

	NA (<i>n</i> = 15)	aMCI (<i>n</i> = 16)	Cohen's <i>d</i>
Age (years)	75.3 (8.7)	74.6 (8.0)	0.08
Female:Male ratio	9:7	6:9	
Education (years)	14.4 (3.5)	14.9 (3.0)	0.15
MMSE	29.0 (1.0)	27.7 (1.9)	0.89
Vocabulary SS	13.6 (2.9)	13.8 (2.7)	0.07
Digit Span Forward SS	11.9 (3.4)	10.8 (2.9)	0.34
Digits Span Backward SS	13.3 (3.3)	12.7 (2.1)	0.01
TMT A SS	10.3(1.48)	9.47(1.92)	0.48
TMT B SS	12.69(1.89)	10.79(1.93)	0.99
Rey-Osterrieth Copy SS	8.94(2.14)	8.25(1.54)	0.37
Rey-Osterrieth Immediate Memory* SS	11.19(3.63)	7.25 (2.63)	1.24
Boston Naming* SS	12.68(2.75)	10.80(3.53)	0.59
HADS Depression Scale	2.6 (2.3)	2.6 (2.3)	0
HADS Anxiety Scale	5.1 (3.8)	5.6 (2.5)	0.16

987 *Note.* Mean scores with standard deviations in parentheses. NA= normal aging; aMCI = amnesic mild
988 cognitive impairment; MMSE = Mini-Mental Status Exam; Vocabulary = Expressive vocabulary;
989 Digits Span = Total attention span score for digits forward and backwards; TMT = Trail Making Test A
990 (number sequencing) and B (alternating number-letter sequencing); Rey-O Copy = visual construction
991 of the Rey-Osterreith Complex Figure; Rey-O Memory = immediate recall of the complex figure; SS =
992 age-corrected scaled score. HADS = Hospital Anxiety Depression Scale with scores < 7 considered
993 within normal limits. *indicates significantly different between NA and aMCI

994

995 **Table 2:** *Demographic Data for the Participant Groups*

	NA males (<i>n</i> = 7)	aMCI males (<i>n</i> = 9)	NA females (<i>n</i> = 9)	aMCI females (<i>n</i> = 6)	P values	Effect Size: η^2_p
Age (years)*	71.3 (8.8)	76.0 (7.7)	79.8 (5.9)	71.5 (8.0)	Sex ns Group ns Group by sex 0.025	Sex .02 Group .02 Group by sex .017
Education (years)	14.7 (3.3)	15.5 (2.2)	14.1 (3.6)	13.5 (2.7)	Sex ns Group ns Group by sex <i>ns</i>	Sex .05 Group .00 Group by sex .01

996 *Note.* Mean scores with standard deviations in parentheses. NA= normal aging; aMCI = amnesic mild
997 cognitive impairment * denotes significant difference.
998
999

Table 3. Session 1 cortisol moderation models: main effects and interactions.

	Model	Term	<i>B</i>	<i>SE</i>	β	<i>t</i>	<i>p</i>	<i>sr</i> ²
Males	HVLT-R Immediate Recall	GRP	-	0.396	-	-	0.002*	0.468
			1.566		0.921	3.960		
		S1BC	-					0.010
			0.134					
	HVLT-R Delayed Recall	GRP x S1BC	0.407					0.035
		GRP	-	0.410	-	-	0.017*	0.160
			1.138		0.538	2.773		
		S1BC	-					0.016
	HVLT-R Retention		0.210					
		GRP x S1BC	-					0.024
			0.415					
		GRP	-					0.084
	Associative Recognition		0.870					
		S1BC	-					0.002
			0.080					
		GRP x S1BC	-					0.057
	Spatial Working Memory – Trial 1	GRP	-	0.391	-	-	<0.001*	0.593
			1.933		1.036	4.940		
		S1BC	-					0.043
			0.307					
	Spatial Working Memory – Trial 2	GRP x S1BC	0.915	0.371	0.658	2.466	0.030*	0.148
		GRP	0.809					0.065
		S1BC	0.219					0.014
	Spatial Working Memory – Delay	GRP x S1BC	-					0.001
			0.104					
		GRP	1.484	0.673	0.682	2.206	0.048*	0.257
			0.549					
Females	HVLT-R Immediate Recall	GRP x S1BC	-	0.638	-	-	0.050*	0.250
			1.389		0.856	2.177		
		GRP	1.337					0.188
		S1BC	0.177					0.010
	HVLT-R Delayed Recall	GRP x S1BC	-					0.027
			0.479					
		GRP	-	0.691	-	-	0.025*	0.347
			1.785		0.794	2.582		
	HVLT-R Delayed Recall	S1BC	-					0.004
			0.187					
		GRP x S1BC	-					0.009
			0.334					
	HVLT-R Delayed Recall	GRP	-	0.389	-	-	0.001*	0.612
			1.944		1.054	5.002		
		S1BC	-					<0.001
			0.016					

		GRP x	-				0.023
		S1BC	0.438				
HVLT-R	GRP	-	0.338	-	-	<0.001*	0.696
Retention			1.913		1.124	5.653	
		S1BC	0.520				0.057
	GRP x	-	0.393	-	-	0.026*	0.145
	S1BC		1.016		0.870	2.582	
Associative	GRP	-	0.548	-	-	0.049*	0.194
Recognition			1.213		0.593	2.213	
		S1BC	-	0.522	-	-	0.062
			1.082		0.806	2.073	0.170
		GRP x	0.838				0.068
		S1BC					
Spatial	GRP	-					0.074
Working			0.522				
Memory –	S1BC	0.618	0.284	0.661	2.181	0.052	0.114
Trial 1	GRP x	-	0.346	-	-	0.001*	0.468
	S1BC		1.527		1.562	4.413	
Spatial Working	GRP	0.226					0.009
Memory – Trial	S1BC	0.074					0.001
2	GRP x	-					0.084
	S1BC	0.804					
Spatial	GRP	-					0.013
Working		0.245					
Memory –	S1BC	0.349					0.029
Delay	GRP x	-	0.377	-	-	0.004*	0.289
	S1BC		1.354		1.228	3.593	

Note. Male $n = 16$ per model. Female $n = 14-15$ per model. TSST = Trier social stress test. Group (GRP) = (1) amnesic mild cognitive impairment vs. (0) normal aging. SIC = stress-induced cortisol. DC = diurnal cortisol. * = $p < 0.05$. Effects and interactions with $p \sim 1.0$ have SE, β , and t values shown.

Table 4. Session 2 cortisol moderation models: main effects and interactions.

	Model	Term	<i>B</i>	<i>SE</i>	β	<i>t</i>	<i>P</i>	<i>sr</i> ²
Males	HVLT-R Immediate Recall	GRP	-	0.507	-	-	0.002*	0.387
			1.933		0.872	3.816		
		S2BC	0.154					0.010
		GRP x S2BC	-					<0.001
	HVLT-R Delayed Recall	GRP	-	0.327	-	-	<0.001*	0.653
			2.376		1.133	7.272		
		S2BC	-					<0.001
			0.030					
	HVLT-R Retention	GRP	-	0.579	-	-	0.017*	0.317
			1.597		0.789	2.757		
		S2BC	-					<0.001
			0.020					
	Associative Recognition	GRP	-	0.468	-	-	0.002*	0.431
			1.890		0.921	4.035		
		S2BC	-					0.001
			0.051					
	Spatial Working Memory – Trial 1	GRP	-	0.468	-	-	0.002*	0.431
			1.890		0.921	4.035		
		S2BC	-					0.001
			0.051					
	Spatial Working Memory – Trial 2	GRP	-	0.468	-	-	0.002*	0.431
			1.890		0.921	4.035		
		S2BC	-					0.001
			0.051					
	Spatial Working Memory – Delay	GRP	-	0.468	-	-	0.002*	0.431
			1.890		0.921	4.035		
		S2BC	-					0.001
			0.051					
	TSST Speech Task Anxiety	GRP	-	0.468	-	-	0.002*	0.431
			1.890		0.921	4.035		
		S2BC	-					0.001
			0.051					
	TSST Counting Task Anxiety	GRP	-	0.468	-	-	0.002*	0.431
			1.890		0.921	4.035		
		S2BC	-					0.001
			0.051					

		GRP x S2BC	0.673				0.073	
Females	HVLT-R Immediate Recall	GRP	-	0.373	-	-	0.025*	0.207
			0.966		0.611	2.589		
		S2BC	-					0.003
			0.101					
		GRP x S2BC	0.649					0.070
	HVLT-R Delayed Recall	GRP	-	0.594	-	-	0.041*	0.307
			1.390		0.745	2.341		
		S2BC	0.232					0.011
		GRP x S2BC	-					0.006
			0.232					
	HVLT-R Retention	GRP	-	0.645	-	-	0.067	0.251
			1.311		0.673	2.034		
		S2BC	0.285					0.015
		GRP x S2BC	-					0.049
			0.666					
	Associative Recognition	GRP	-					0.067
			0.672					
		S2BC	-					0.010
			0.237					
		GRP x S2BC	-					0.001
			0.082					
	Spatial Working Memory – Trial 1	GRP	0.393					0.017
		S2BC	0.173					0.004
		GRP x S2BC	0.267					0.006
	Spatial Working Memory – Trial 2	GRP	0.645					0.069
		S2BC	0.027					<0.001
		GRP x S2BC	-					0.024
			0.436					
	Spatial Working Memory – Delay	GRP	0.707					0.094
		S2BC	0.442					0.045
		GRP x S2BC	-					0.018
			0.359					
	TSST Speech Task Anxiety	GRP	-	0.495	-	-	0.034*	0.199
			1.265		0.599	2.557		
		S2BC	2.016	0.446	1.356	4.520	0.002*	0.622
		GRP x S2BC	-	0.571	-	-	0.057	0.150
			1.266		0.632	2.217		
	TSST Counting Task Anxiety	GRP	0.936					0.136
		S2BC	-					0.003
			0.119					
		GRP x S2BC	-					0.001
			0.081					

Note. Male n = 15-16 per model. Female n = 10-15 per model. TSST = Trier social stress test. Group (GRP) = (1) amnesic mild cognitive impairment vs. (0) normal aging. SIC = stress-induced cortisol.

1010 *DC = diurnal cortisol. * = $p < 0.05$. Effects and interactions with $p \sim 1.0$ have SE, β , and t values*
 1011 *shown.*

Figure Captions

Figure 1. Timeline of Sessions and Testing within Sessions. Exact times varied based on individual variability.

Figure 2. (A) Log₁₀ transformed salivary cortisol at two separate time points and (B) Log₁₀ transformed area under the curve with respect to the ground (AUCg) cortisol during Session 1 in normal aging (NA; $n = 9$ females, $n = 7$ males) and amnesic mild cognitive impaired (aMCI; $n = 6$ females, $n = 9$ males) participants. (C) Log transformed salivary cortisol at four separate time points and (D) Log₁₀ transformed AUCg cortisol during Session 2 in NA ($n = 9$ females, $n = 7$ males) and aMCI ($n = 6$ females, $n = 9$ males). # denotes a Main effect of sex, group, or time, $p < 0.05$. * denotes aMCI males differ from all other groups at a single time point or compared to all other groups for AUCg cortisol, $p < 0.05$. & denotes aMCI females or males differ from NA females or males respectively at a single time point, $p < 0.05$.

Figure 3. Hopkins Verbal Learning Test-Revised (HVLT-R) (A) immediate recall, (B) delayed recall, and (C) retention scores during session 1 and session 2 (after Trier Social Stress Test) in normal aging (NA; $n = 9$ females, $n = 7$ males) and amnesic mild cognitive impairment (aMCI; $n = 6$ females, $n = 9$ males). # denotes Main effect of group or session, $p < 0.05$. * denotes a significant increase from Session 1 and Session 2 in NA males and females, $p < 0.05$. & denotes a significant decrease from Session 1 and Session 2 in aMCI males, $p < 0.05$.

Figure 4. (A) Face-name associative recognition and (B) spatial working memory performance during Session 1 and Session 2 in normal aging (NA; $n = 9$ females, $n = 7$ males) and amnesic mild cognitive impaired (aMCI; $n = 6$ females, $n = 9$ males) participants. # denotes main effect of sex or group $p < 0.05$. * denotes fewer number of choices in delay trial compared to Trial 1 and Trial 2.

Figure 5. (A) Log₁₀ transformed diurnal salivary cortisol measures at five time points averaged across three consecutive days in normal aging (NA) males ($n = 7$) and amnesic mild cognitive impairment (aMCI; $n = 9$). (B) Log transformed area under the curve with respect to the ground (AUCg) diurnal cortisol in NA and aMCI males.

Figure 6. Standardized moderation effects of Session 1 cortisol (AUCg) on the relationship between participant group (normal aging (NA), amnesic mild cognitive impairment (aMCI)) and Session 1 (A)

associative recognition scores and (B) spatial working memory task Trial 2 choices in males ($n = 7$ NA, $n = 9$ aMCI) and (C) Hopkins Verbal Learning Test-Revised (HVLTR) retention scores, and (D) spatial working memory task Trial 1 and (E) delay choices in females ($n = 9$ NA, $n = 6$ aMCI).

Figure 7. Standardized moderation effects of Session 2 cortisol (AUCg) on the relationship between participant group (normal aging (NA), amnesic mild cognitive impairment (aMCI)) and Session 2 (A) perceived trier social stress test (TSST) speech anxiety ratings and (B) Hopkins Verbal Learning Test-Revised (HVLTR) delayed recall scores in males ($n = 7$ NA, $n = 9$ aMCI) and Session 2 (C) perceived anxiety rating for the Trier Social Stress Test in females ($n = 9$ NA, $n = 6$ aMCI).

Session 1

10:00	Consent, interview, and instructions
10:10	Saliva Sample #1
10:15	Face / Name Associative Memory Version A – Immediate
10:30	Hopkins Verbal Learning Test-Revised: Form 6 – Immediate
10:35	Spatial Working Memory Version A – Immediate
10:45	Trail Making Test – A & B
10:50	Face / Name Associative Memory Version A - Delay
10:58	Hopkins Verbal Learning Test-Revised: Form 6 - Delay
11:00	Mini-Mental Status Exam
11:05	Spatial Working Memory Version A – Delay (20-25 minutes)
11:10	WAIS-III Digit Span
11:20	Boston Naming Test (split half - odds)
11:30	Rey-Osterreith Figure Copy and Immediate Recall
11:40	WAIS-III Vocabulary (split half - odds)
11:50	Hospital Anxiety and Depression Scale
11:55	Saliva Sample #2, schedule at-home saliva collection, book session 2
12:05	Session ends

Session 2

10:00	Beck's Anxiety and Depression Inventories
10:10	Saliva sample #1 (~20 minutes before TSST)
10:13	Instructions and consent for Trier Social Stress Test
10:18	Trier Social Stress Test – preparation of speech
10:28	Saliva sample #2 (immediately before TSST Speech)
10:31	Trier Social Stress Test
10:43	Hopkins Verbal Learning Test-Revised: Form 5 – Immediate
10:48	Face / Name Associative Memory Version B – Immediate
11:03	Saliva Sample #3 (~30 minutes after beginning of TSST)
11:06	Spatial Working Memory Version B – Immediate
11:16	Hopkins Verbal Learning Test-Revised: Form 5 – Delay
11:18	Face / Name Associative Memory Version B – Delay
11:23	Spatial Working Memory Version B – Delay
11:28	Trail Making Test – A
11:33	Coping Strategies Scale, SF-36 Health Survey
11:43	Saliva Sample #4 (~60 minutes after beginning of TSST)
11:44	Session ends

1054

1055

1060
1061

1062

1063

SESSION 1: MODERATION MODELS

SESSION 2: MODERATION MODELS