Title: Inadequate level of knowledge, mixed outlook and poor adherence to COVID-19 prevention guideline among Ethiopians

Edessa Negera^{1, *}, Tesfaye Moti Demissie³, Ketema Tafess²

¹Hawassa University, Department of Biology, Ethiopia; ²College of Health Sciences, Arsi University, Ethiopia; ³ Medical Science Division, Centre for Vaccinology and Tropical Medicine, University of Oxford; * corresponding author

Abstract

COVID-19 has a potential to cause chaos in Ethiopia due to the country's already daunting economic and social challenges. Living and working conditions are highly conducive for transmission, as people live in crowded inter-generational households that often lack running water and other basic sanitary facilities. Thus, the aim of this study was to investigate the knowledge, attitudes and practices (KAP) of Ethiopians toward COVID-19 following the introduction of state of emergency by the Ethiopian government to curb the spread of the disease. A cross-sectional study design was conducted in nine regional states and two chartered cities. Data for demographic, Knowledge, attitude and practice toward COVID-19 were collected through telephone interview from 1570 participants. Descriptive and bivariate analyses using chi-square test, t-test or analysis of variance were performed as appropriate. Binary and multiple logistic regression analysis were used to measure the relationship between the categorical dependent variables and one or more socio-demographic independent variables with two-tailed at $\alpha=0.05$ significance level and 95% of confidence interval. The level of good knowledge, favourable attitude and good practice among the respondents were 42%, 53.8% and 24.3% respectively. Being rural resident, older than 50 years, having at least primary education, being resident of Amhara and Oromia regions were independent predictors of knowledge level. While being rural resident, married, employed, having at least basic education, being residents of Afar, Amhara, Gambela, Oromia and Somali regions were found

26 to be the best predictors of the attitude, being rural resident, government employee, having at
 27 least basic education, and living outside of the capital were the independent predictors of
 28 practice level of the respondents. The finding revealed that Ethiopians have inadequate level
 29 of knowledge and are generally have a mixed outlook on overcoming the pandemic with poor
 30 adherence to COVID-19 prevention practice. reinforcing preventive measures and intensifying
 31 sensitization campaigns to fill the knowledge gap and persuading people to follow the
 32 preventive measures set by the government with concurrent evaluation of the impacts of these
 33 measures on knowledge and practice is highly recommended to mitigate the disease.

1. Introduction

Coronavirus disease 2019 (COVID-19) is a pandemic disease caused by a novel human coronavirus (SARS-COV-2) first identified in china in December 2019 (1). The World Health Organization declared the outbreak of COVID-19 as a pandemic on the 11th of March 2020, after it has spread to 113 countries worldwide (2).

According to current evidence, the COVID-19 virus is primarily transmitted rapidly between people through respiratory droplets and contact routes (3). Airborne transmission has been suggested by some studies (4, 5). Recent experimental studies have examined the stability of SARS-CoV-2, showing that the virus remains infectious in aerosols for hours (5) and on surfaces up to days (5, 6). The mean incubation period of COVID-19 is about 3–9 days with a range between 0–24 days (3, 7). However, the mean time between successive cases in a chain of transmission is about 3-8 days suggesting that one becomes contagious before symptoms present about 2.5 days earlier from the onset of symptoms (8). Studies estimated that about 44% of transmission of COVID-19 occur before the onset of symptoms (9).

The first case of COVID-19 was confirmed in Ethiopia on 13th of March 2020 (10). The state of emergency was declared by the government on 8th of April 2020 to control the pandemic. The state of emergency includes closing schools, banning public gatherings and requiring employees to work from home (10). The introduction of the state of emergency has been welcomed by most citizens and institutions but was not without critiques from some political opposition parties. While the federal and regional governments announced measures such as suspending large gatherings and inter-city public transport, authorities have not introduced a comprehensive lockdown to try to contain the virus due to some real challenges. Firstly, most citizens live day-to-day and they may see a complete lockdown as counterproductive and unfair to those on the bottom rungs of society. Secondly, complete lockdown could worsen the life

of the vulnerable segments of society such as street children, internally displaced persons and refugees (11).

COVID-19 has the potential to cause chaos in Ethiopia due to the country's already daunting economic and social challenges. On one hand, the public health risks presented by COVID-19 are vast. Living and working conditions are highly conducive for transmission, as people live in crowded inter-generational households that often lack running water and other basic sanitary facilities. Allowing economic activity to continue unchecked could lead to huge infections within months, with serious cases quickly overwhelming the already weak health system (12).

Public health intervention measures are rapidly changing around the world to cope up with the rapid transmission of the disease and to minimize the risk for infection and spread of the disease. However, miscommunication regarding the threat of COVID-19 may lead to public confusion and inaction (13). Being older and having underlying health conditions, reduced ability to access and understand health information, inability to make well-informed decisions and failure to take optimal health-promoting are among the greatest risk factors for severe infection and death due to COVID-19 (13, 14). This situation is especially true when the health information itself is not timely, trusted, consistent, or actionable particularly in sub-Saharan African countries where health inequalities are worsened by lack of political commitment and good governance (15). As a result, a wide range of health disparities may exist by age, ethnicity, regions, political affiliations and socio-demographic status (15-17). Therefore, During COVID-19 pandemic when the understanding of critical preventive measures and ever-changing public health messages is most important, many vulnerable populations may be further marginalized by inadequate health communication, posing substantial risks to themselves and their communities.

81 On the other hand, the success of national preventive strategy of a pandemic is largely depends
 82 on the adherence of the public to the preventive measures established by the government. The
 83 public adherence is likely to be influenced by trust in government (18) and by public
 84 knowledge and attitude toward the pandemic (19). By evaluating the public knowledge, attitude
 85 and practice about COVID-19 pandemic, it is possible to explore attributes that influence the
 86 public in adopting practices and responsive behaviour toward COVID-19 prevention measures.

87 Thus, the aim of study was to investigate the knowledge, attitudes and practices (KAP) of
 88 Ethiopians toward COVID-19 following the introduction of state of emergency by the
 89 Ethiopian government to help curb the spread of COVID-19 following 55 cases and 2 fatalities
 90 on April 8, 2020 (20). we conducted a time sensitive KAP study in all regional states and
 91 chartered cities of Ethiopia from May 20, 2020 to June 20, 2020.

2. Materials and methods

2.1. Study design and site

A cross-sectional study was conducted from May 20 to June 20, 2020 in nine regional states and two chartered cities of Ethiopia. The regional states include Tigray, Afar, Amhara, Oromia, Harari, Somali, Gambela, Benishangul and Southern Nations, Nationalities and Peoples' Region. (SNNP). The two chartered cities are Addis Ababa (Finfinnee) and Dire Dawa.

2.2. Study population

The study population included all peoples living in the nine regional states and two chartered cities who are 18 years old and above during the study period.

2.3. Study procedure

This study was conducted after the government declared a state of emergency in the country to help curb the spread of COVID-19 following 55 cases and 2 fatalities on April 8, 2020. As face to face data collection was impossible, we opted to use telephone interview for enrolling potential participants. We ruled out the online enrolling methods for two reasons: Firstly, only about 17% of the population has access to the internet penetration but more than 45% of the population has access to either mobile network or fixed landline telephones in 2020 (21). Secondly, adult literacy is only 52% and social media penetration in Ethiopia is 5.5% by January 2020 (22). As a result, telephone interview was found to be the best alternative among the existing methods of data collection where face-to-face data collection is not possible.

2.4. Study tools

We adapted the survey tools from previous KAP studies towards COVID-19 (2, 13, 19). We included four core themes to assess the KAP of Ethiopians toward COVID-19. (i) demographic

data collection tools. (ii) knowledge about COVID-19, (iii) attitude toward COVID-19 and (iv) practices appropriate to curb covid-19 infection and transmission.

Demographic questions: Each respondent was asked for age, gender, residence, occupation, region, educational and marriage status.

COVID-19 Knowledge: Demonstrated knowledge of COVID-19 was assessed through asking the respondents 10 close-ended questions. The questions include the respondents' knowledge about the cause, transmission route, clinical symptoms, prevention and control measures of COVID-19. Respondents were given three response options: 'true', 'false' or 'I do not now'. A correct response was assigned 1 point, while an incorrect or I do not know response to the question was assigned 0 points. Each respondent achieves between 0 and 10 score points. Higher score indicates better knowledge of COVID-19 (supplementary (1)).

Attitude toward COVID-19: To assess the attitude of respondents toward COVID-19, we used 5 questions. Two questions include confidence of the respondent on COVID-19 prevention methods and confidence on the government to contain the spreading of coronavirus. They were also asked if they are in favour of the state of emergency introduced to curb the spread of COVID-19, their perception to get infected with coronavirus and if they believe traditional herbs and religious faith can cure COVID-19. For each question a 4-point Likert scale response options (strongly disagree, disagree, agree and straggly agree) were provided (supplementary (1)).

Practice towards COVID-19 prevention: To investigate how often respondents' practice COVID-19 prevention, 5-questions were used. The questions include how often the respondent wash its hand with soap, avoid non-essential travel, keep 2-meter social distancing, avoid social gatherings and avoid touching eyes, nose, and mouth with unwashed (unsensitized) hands. For

each question a 4-point Likert scale response options (never, sometimes, usually and always) were provided.

2.5. Data collection methods

Twenty data collectors were recruited and trained online. The data collectors were recruited from each region. All data collectors had previous experience of KAP data collection by phone. In most cases, phone numbers were randomly selected from local phone book. Following the verbal consent, each respondent was assured that the voice is not recorded, and any personal data is not required to participate in the study.

2.6. Operational definitions

Independent variables: all sociodemographic variables: age, sex, education, marital status, residence, employment and region are considered as independent variables.

Dependent Variables: knowledge, attitude and practice toward COVID-19 were considered as dependent variables for all statistical analysis.

Good level of knowledge: Respondents who scored above the mean score of the total respondents were considered as having good level of COVID-19 knowledge and those who scored below the mean as having poor level of COVID-19 Knowledge.

Favourable attitude: A response of agree or strongly agree to attitude questions were considered as favourable attitude for logistic regression analysis. However, for two questions: “I believe that traditional herbs and faith such as holy water can cure COVID-19; and I do not think I will get sick from COVID-19” a response of strongly disagree or disagree were considered as favourable attitude.

Good practice: a response of “usually” and “always” to each practice question was considered as having good practice towards COVID-19 prevention.

2.7. Statistical analysis

Descriptive statistics (means with standard deviations and percentage frequencies) were calculated for all survey responses. Associations between socio-demographic characteristics and responses to COVID-19 knowledge, attitude and practice were investigated in bivariate analyses using chi-square test, t-test or analysis of variance as appropriate. Binary and multiple logistic regression analyses were used to measure the relationship between the categorical dependent variables (knowledge, attitude and practice) and one or more socio-demographic independent variables. results of logistic regression have been reported as adjusted odds ratio (AOR). Data were analysed using Stata for Windows, version 16 (Stata Corp., College Station, TX, USA with two-tailed at $\alpha=0.05$ significance level and 95% of confidence interval

3. Result

3.1. Social and demographic characteristics

A total of 1570 participants completed the survey questionnaire of which 896 (57.1%) of the respondents were male. A third of the respondents were from rural residents. The age of the respondents ranges from 18 and 73 years with a mean $\pm$ standard deviation (SD) of 31.6 year $\pm$ 12.8. Among the respondents, 660 (42.0%) had college or university education while 260 (16.6%) are unable to read and write. Occupation-wise, about a third (34.9%) of them were government employees while a fifth (19.5%) of them were self-employed and one in 6 (15.9%) were farmers (Table 1).

179 Table 1. Sociodemographic characteristics of respondents (N=1570).

Demographic variable s		Number	Percentage (%)
Sex	male	896	57.1
	female	674	42.9
Residence	Urban	1094	69.7
	Rural	476	30.3
Ager group	18-29	860	54.8
	30-39	332	21.1
	40-49	174	11.1
	50-59	128	8.2
	60+	76	4.8
Marital status	single	662	42.2
	married	894	56.9
	widow	14	0.9
Education	illiterate	260	16.6
	basic	292	18.6
	primary	140	8.9
	secondary	218	13.9
	college/university	660	42.0
Occupation	farmer	250	15.9
	government employee	306	19.5
	self-employed	548	34.9
	student	466	29.7
Region/chartered city	Addis Ababa	326	20.8
	Afar	42	2.7
	Amhara	190	12.1
	Benishangual Gumuz	46	2.9
	Dire Dawa	52	3.3
	Harari	46	2.9
	Gambela	68	4.3
	Oromia	552	35.2
	Somali	60	3.8
	SNNPRS	90	5.7
	Tigray	98	6.2
	Total	1570	100.0

3.2. COVID-19 Knowledge

The overall correct response of the respondents to the knowledge questions about the COVID-19 varies from 14.0% to 73.6% (Table 1) with the mean knowledge score of 4.2 (SD=2.809, range 0-10) suggesting an overall of 42% ($4.2/10 \times 100$) correct rate on COVID-19 knowledge test. Knowledge scores significantly differ across each question. Nearly three-fourth (73.6%) of the respondents have heard about COVID-19 but only 42.4% and 37.8% knew COVID-19 transmission and clinical manifestations, respectively. Only 14% of the respondents knew that the asymptomatic transmission of COVID-19. While about 63.8% of the respondents had correctly answered COVID-19 prevention methods such as handwashing, social distancing, avoiding crowd gatherings and using face mask. only 18.6% of them knew that people who have COVID-19 symptoms should self-isolate at least for 7 days and their household contacts for 14 days. A third of the respondents consider that only elderly people or people with chronic health conditions develop severe condition if infected with COVID-19. Nearly 80% of the respondents did not know that currently there is no effective drug to cure COVID-19 nor vaccines to prevent the infection (Table 2).

Table 2. Summary of respondents' response to COVID-19 knowledge questions (N=1570).

no	knowledge questions	Responses n (%).		
		Yes	No	I do not know
1	I have heard about Coronavirus (Covid-19) virus	1156 (73.6)	414 (26.4)	
2	Coronavirus disease (COVID-19) is an infectious disease caused by a newly discovered coronavirus	414 (25.7)	136 (8.7)	1030 (65.6)
3	The main clinical symptoms of COVID-19 are fever, fatigue, continuous dry cough and shortness of breath	594 (37.8)	170 (10.8)	806 (51.3)
4	COVID-19 virus spreads primarily through droplets of saliva or discharge from the nose when an infected person coughs or sneezes	666 (42.4)	160 (10.2)	744 (47.4)
5	Only elderly people or people with chronic health condition develop sever condition if infected with COVID-19	454 (28.9)	458 (27.9)	678 (43.2)
6	Peoples infected with COVID-19 virus but not have any symptoms do not infect others	706 (45.0)	220 (14.0)	644 (41.0)
7	Children and young healthy adults do not necessarily take measures against COVID-19 as they are immune to the disease	450 (28.7)	436 (27.8)	684 (43.6)
8	Handwashing, social distancing, avoiding crowding and using facemask can reduce the risk of covid-19 infection	1002 (63.8)	154 (9.8)	414 (26.4)
9	People who have COVID-19 symptoms should self-isolate at least for 7 days and their household contacts for 14 days	292 (18.6)	222 (14.1)	1056 (67.3)
10	Currently there is no effective drug to cure COVID-19 nor vaccines to prevent infections	304 (19.4)	226 (14.4)	1040 (66.2)

198 The student t-test has shown that the level of COVID-19 knowledge was low among female
 199 respondents compared to male respondents ($P=0.05$). Older respondent (≥ 50 years) had low
 200 COVID-19 knowledge than the other age group ($F=8.38$, $P < 0.0001$). Similarly, there was a
 201 significant difference in the knowledge score of the respondents among age groups, residence,
 202 educational status, marital status and regions of residence (Table 3).

203 Table 3. Respondents knowledge score of COVID-19 by demographic variables (N=1750).

Demographic variables		Number of participants (%)	Knowledge score (mean $\pm$ SD), (maximum score 10)	Statistical test (t or F)	Significance (P)
Sex	male	896(57.1)	4.641 $\pm$ 0.09706	2.12	0.0343*
	Female	674 (42.9)	4.338 $\pm$ 0.1021		
	Total	1570 (100.00)	1570 (100.00)		
Age in ears	18-29	860 (54.8)	4.73023 $\pm$ 3.73023	8.38	< 0.0001*
	30-39	332 (21.1)	4.40964 $\pm$ 3.17861		
	40-49	174 (11.1)	4.77011 $\pm$ 3.1077		
	50-59	128 (8.2)	3.40625 $\pm$ 2.75349		
	60+	76 (4.8)	3.73684 $\pm$ 2.92743		
	Total	1570 (100.00)	1570 (100.00)		
Residence	urban	1094 (69.9)	4.724 $\pm$ 0.08727		< 0.0001*
	rural	476 (30.1)	4.000 $\pm$ 0.1163	4.73	
	Total	1570 (100.00)	1570 (100.00)		
Marital status	single	662 (42.2)	5.10876 $\pm$ 2.48206		
	married	894 (56.9)	4.0962 $\pm$ 2.95197	31.61	< 0.0001*
	widow	14 (0.9)	2.00 $\pm$ 0.784465		
	Total	1570 (100.00)	1570 (100.00)		
Occupation	government employed	548 (34.9)	7.03268 $\pm$ 2.85248		
	self-employed	306 (19.5)	3.37591 $\pm$ 2.32645	230.00	< 0.0001*
	farmer	250 (15.9)	2.576 $\pm$ 1.65868		
	student	466(29.7)	5.13008 $\pm$ 2.30906		
	Total	1570 (100.00)	1570 (100.00)		
Educational level	illiterate	260 (16.6)	2.53077 $\pm$ 1.53077	165.20	< 0.0001*
	basic	292(18.6)	2.93836 $\pm$ 2.25283		

	primary	140(8.9)	4.31429 ±2.73818		
	secondary	218(13.9)	3.92661±2.11079		
	college/university	660(42.0)	6.18485 ±2.63035		
	Total	1570 (100.00)	1570 (100.00)		
Region/chartered city	Addis Ababa	326 (6.2)	6.42331±3.13765		
	Afar	42 (2.7)	4.61905±2.368	12.40	< 0.0001*
	Amhara	190 (12.1)	4.32632±2.99362		
	Benishangul Gumuz	46 (2.9)	4.3913±2.6955		
	Dire Dawa	52 (3.3)	5.11538±2.55645		
	Harari	46 (2.9)	4.65217±2.54049		
	Gambela	68 (4.3)	4.64706±2.35483		
	Oromia	552 (35.2)	3.66667±2.47626		
	Somali	60 (3.8)	4.66667±2.28975		
	SNNPRS	90 (5.7)	4.24444±2.82966		
	Tigray	98 (20.8)	5.97959±2.54031		
	Total	1570 (100.00)	1570 (100.00)		

205 After grouping respondents into good (mean score ≥ 4.2) and poor (mean score < 4.2) level of
 206 knowledge using a bloom's cut of points, binary and multivariable logistic regression was
 207 performed. Age, occupation, residence and education have been significantly associated with
 208 COVID-19 knowledge of the respondents with correct prediction of 72.2% in multivariable
 209 regression analysis. Multivariable logistic regression analysis has shown that respondents of
 210 age group 50-59 (AOR= 0.562; (95% CI: 0.4285-0.708) and above 59 years old (AOR=0.387;
 211 (95%CI: 0.1924-0.4982) were less knowledgeable about COVID-19 compared to respondents
 212 in the age group of 18-29. Similarly, respondents those who had college or university education
 213 had more than 14 times higher knowledge (AOR=14.026; (95%CI: 9.56431-21.8764)
 214 compared to those who are unable to read and write. Respondents from the two most populous
 215 regions, Amhara (AOR=0.497 (95%CI: 0.2990-0.6542) and Oromia (AOR= 0.4512; 95%CI:
 216 0.3218-0.8026) regional states had poor level of knowledge compared with respondents from
 217 Addis Ababa (Finfinnee), the capital (Table 4).

218 Table 4. Logistic regression analysis of odds ratio for knowledge of the respondents in relation to COVID-19 potential risk factors.

4. Demographic variables		Univariable logistic regression			Multivariable logistic regression		
		Crude OR	95% CI	P-value	Adjusted OR	95% CI	P-value
Sex	Male	1.000			1		
	Female	0.921	0.747-1.135	0.440	0.864	0.7234-1.0923	0.621
Age in ears	18-29	1.000			1		
	30-39	0.829	0.637-1.080	0.165	0.923	0.6754-1.1073	0.243
	40-49	0.944	0.675-1.320	0.735	1.021	0.6891-1.4225	0.14
	50-59	0.473	0.307-0.728	0.001*	0.562	0.4285-0.708	0.004*
	60+	0.290	0.154-0.545	0.000*	0.387	0.1924-0.4982	0.001*
Residence	Urban	1.000					
	Rural	0.682	0.541-0.860	0.001*	0.772	0.5640-0.9245	0.004*
	Single	1.000					
Marital status	Married	0.524	0.425-0.646	0.000*	1.023	0.8776-1.2113	0.061
	Widow	0.000	0.621-1.135	0.998	0.8823	0.7645-0.9921	0.12
	Farmer	1.000					
Occupation	Government Employee	20.599	12.681-33.461	0.000*	16.4321	11.4564-27.9687	0.001*
	Self-employed	2.206	1.376-3.536	0.001*	1.76	1.0248-2.6843	0.008*
	Student	8.716	5.512-13.784	0.000*	5.0264	3.0245-10.2318	0.001*
Education	Illiterate	1.000					
	Basic	2.023	1.181-3.466	0.000*	1.276	0.9876-2.1346	0.086
	Primary	4.327	2.447-7.654	0.000*	3.0341	1.9842-5.6742	0.02*
	Secondary	3.740	2.197-6.366	0.000*	3.042	2.4420-6.2143	0.012*
	college/university	16.434	10.335-26.130	0.000*	14.026	9.56431-21.8764	0.001*
	Addis Ababa	1.000					

Region/chartered city	Afar	0.745	0.569-1.102	0.091	0.703	0.5423-1.097	0.110
	Amhara	0.513	0.349-0.756	0.001*	0.497	0.2990-0.6542	0.007*
	Benishangual Gumuz	0.596	0.302-1.259	0.127	0.586	0.2882-0.7549	0.340
	Dire Dawa	0.999	0.552-1.807	0.997	0.992	0.5286-1.577	0.481
	Harari	0.696	0.306-1.159	0.427	0.621	0.2764-1.112	0.67
	Gambela	0.743	0.431-1.280	0.284	0.709	0.476-1.1187	0.41
	Oromia	0.470	0.305-0.889	0.000*	0.4512	0.3218-0.8026	0.001*
	Somali	0.908	0.518-1.592	0.737	0.9	0.6234-1.4410	0.820
	SNNPRS	0.553	0.334-0.918	0.022*	0.54	0.3491-0.8823	0.230
	Tigray	1.672	1.061-2.634	0.027	1.02	0.7465-3.2438	0.160

219 OD= odds ratio; CI= confidence interval; * significant at $P \leq 0.05$

3.3. Attitude

Five questions were used to measure the attitude of the respondents towards COVID-19. More than 50% of the respondents either agree or strongly agree that traditional herbs and religious faith such as holy water can cure COVID-19. Half of the respondents think that it is unlikely to get sick from COVID-19. About 47% of the respondents do not agree with the principles of COVID-19 prevention methods such as hand washing, social distancing, avoiding non-essential travel and self-isolation (if have symptoms) and 57.1% of them disagree with the current introduction of state emergency to curb the spread of COVID-19. Only 40.4% of the respondents are confident that the government can prevent a nationwide outbreak of the disease (Table 5).

The association between the socio-demographic variables and COVID-19 attitude was assessed by chi-square test (Table 5). The overall favourable attitude response rate was 43.8%. Respondents in the age group of 40-49 years had more favourable attitude (50.6%) towards COVID-19 prevention methods and principles than the other age groups ($\chi^2=8.6$, $P \leq 0.05$). Interestingly rural residents had more favourable attitude (59.2%) than the corresponding urban residents (37.1%) ($\chi^2=65.996$, $P \leq 0.001$). There was strong association between occupation and attitude towards COVID-19 ($\chi^2 = 335.735$, $P \leq 0.0001$). While government employees had the highest favourable response (77.8%), self-employed respondents had the least favourable response rate (15.7%). The educational status of the respondents was also found to be strongly associated with the attitude of the respondents towards COVID-19 ($\chi^2=195.914$, $P \leq 0.001$). Respondents from Tigray (65.3%) and Amhara (63.2%) regions had relatively higher favourable attitude while respondents from Gambela (17.6%) and Afar (28.6%) had the lowest favourable attitude ($\chi^2 = 89.64$, $P \leq 0.001$) (Table 6).

243 Table 5. Summary of respondents' response to COVID-19 attitude questions (N=1570).

no	Attitude questions	Response n (%); N= 1570			
		Strongly disagree	Disagree	Agree	Strongly agree
1	I believe that traditional herbs and religious faith such as Holy water can cure COVID-19	588 (37.5)	186 (11.8)	290 (18.5)	506 (32.2)
2	I am in favour of state of emergency introduced to control the spread of the coronavirus	678 (43.2)	220 (14.0)	328 (20.9)	344 (21.9)
3	I don't think that I will get sick from the coronavirus	490 (31.2)	322(20.5)	286(18.2)	472 (30.1)
4	I am confident that the government can prevent a nationwide outbreak of the coronavirus	598 (38.1)	388 (21.5)	232 (14.8)	402 (25.6)
5	Hand washing, social distancing, avoiding non-essential travel and self-isolation (if have symptoms) greatly help to contain the spreading of coronavirus	386 (24.6)	356 (22.7)	262 (16.7)	566 (36.1)

244

245

246 Table 6. Respondents attitude score of COVID-19 by demographic variables (N=1570).

Demographic variables		Attitude response n (%)		Statistical test	p (two-sided)
		unfavourable	favourable	χ^2	
Sex	male	502 (56.0)	394 (44.0)	0.02	0.889
	Female	380 (56.4)	294 (43.6)		
Age in ears	18-29	478 (55.6)	382 (44.4)	8.60	0.072
	30-39	188 (56.6)	144 (43.4)		
	40-49	86 (49.4)	88 (50.60)		
	50-59	84 (65.6)	44 (34.4)		
	60+	46 (60.5)	30 (39.5)		
Residence	urban	688 (62.9)	406 (37.1)	65.996	0.000
	rural	194 (40.8)	282 (59.2)		
Marital status	single	332 (50.2)	330 (49.8)	20.54	0.000
	married	538 (60.2)	356 (39.8)		
	widow	12 (85.7)	2 (14.3)		
Occupation	government employee	68 (22.2)	238 (77.8)	335.735	0.000
	self-employed	462 (84.3)	86 (15.7)		
	farmer	114 (45.6)	136 (54.4)		
	student	238 (51.1)	228 (48.9)		
Education	illiterate	194 (74.6)	66 (25.4)	195.914	0.000
	basic	220 (75.3)	72 (24.7)		
	primary	72 (51.4)	68 (48.6)		
	secondary	152 (69.7)	66 (30.3)		
	college/university	244 (37.0)	416 (63.0)		
Region/chartered city	Addis Ababa	180 (55.2)	146 (44.8)	89.64	0.000
	Afar	30 (71.4)	12 (28.6)		

	Amhara	70 (36.8)	120 (63.2)		
	Benishangual Gumuz	24 (52.2)	22 (47.8)		
	Dire Dawa	24 (46.2)	28 (53.8)		
	Harari	24 (52.2)	22 (47.9)		
	Gambela	56 (82.4)	12 (17.6)		
	Oromia	346 (62.7)	206 (37.3)		
	Somali	44 (73.3)	16 (26.7)		
	SNNPRS	50 (55.6)	40 (44.4)		
	Tigray	34 (34.7)	64 (65.3)		
	overall	882 (56.2)	688 (43.8)		

248 Multivariable regression analysis has shown that residence, marital status, education,
 249 occupation and age are factors that most predict the attitude of the respondents with the correct
 250 predication of 73.2%. The odds of favourable attitude for being rural resident is twice as high
 251 as being urban resident (AOR=1.909; 95%CI: 1.436-2.538). On the other hand, being married
 252 (AOR=0.58; 95%CI: 0.398-0.854) and widow (AOR= 0.17; 95% CI: 0.034-0.854) were less
 253 likely to have favourable attitude compared to unmarried (single) respondents. Government
 254 employees had the highest odds of favourable attitude (OR=4.8) compared to farmers. As the
 255 educational level of the respondents increases, the odds of favourable attitude increase
 256 (OR=1.41, 1.68 and 3.01) for primary, secondary and college/university education
 257 respectively). While, Odds of favourable attitude for respondents from Amhara region is twice
 258 the odds of favourable attitude of respondents from Addis Ababa. The odds of favourable
 259 attitude of respondents from Afar, Gambela, Somali and Oromia are significantly low ($P \leq 0.05$)
 260 (Table 7).

261 Table 7. Logistic regression analysis of odds ratio for attitude of the respondents in relation to COVID-19 potential risk factors.

Demographic variables		Favourable attitude towards Covid-19 prevention measures					
		Univariable binary logistic regression			multivariable logistic regression		
		Crude OR	95% CI	P-value	adjusted OR	95% CI	P-value
Sex	male	1.00					
	Female	1.04	0.83-1.24	0.889	0.908	0.712-1.157	0.436
Age in ears	18-29	1.00					
	30-39	0.96	0.958-1.237	0.745	0.87	0.623-1.234	0.531
	40-49	1.28	0.924-1.774	0.138	1.03	0.7834-2.292	0.643
	50-59	0.66	0.444-0.967	0.033*	0.72	0.548-1.009	0.090
	60+	0.82	0.505-1.318	0.406	0.89	0.728-1.233	0.620
Residence	Urban	1.00					
	Rural	2.46	1.977-3.070	0.000*	1.909	1.436-2.538	0.000*
Marital status	Single	1.00					
	Married	0.61	0.435-0.855	0.00*	0.583	0.398-0.854	0.000*
	Widow	0.39	0.078-1.954	0.25	0.172	0.034-0.854	0.006*
Occupation	Farmer	1.00					
	Government employee	2.79	1.926-4.031	0.000*	4.758	2.144-7.321	0.000*
	Self employed	0.15	0.106-0.211	0.000*	0.559	0.423-0.992	0.042*
	Student	0.53	0.351-0.808	0.003*	0.852	0.582-1.113	0.110
Education	Illiterate	1.00					
	Basic	0.96	0.654-1.415	0.844	0.88	0.6783-1.194	0.740
	Primary	2.78	1.800-4.282	0.000*	1.41	1.09-3.247	0.023*
	Secondary	1.28	0.854-1.908	0.234	1.68	1.420-2.228	0.040*
	College/university	5.01	3.636-6.908	0.000*	3.01	2.078-5.193	0.000*

Region/chartered city	Addis Ababa	1.00			1		
	Afar	0.49	0.244-0.997	0.049*	0.52	0.242-0.8812	0.024*
	Amhara	2.11	1.465-3.050	0.000*	1.876	1.0429-3.378	0.036*
	Benishangual Gumuz	1.13	0.609-2.097	0.698	0.597	0.267-1.332	0.208
	Dire Dawa	1.44	0.799-2.588	0.225	0.841	0.499-1.416	0.514
	Harari	1.14	0.719-2.897	0.698	0.820	0.371-1.812	0.624
	Gambela	0.26	0.136-0.0511	0.000*	0.259	0.113-0.595	0.001*
	Oromia	0.73	0.556-0.969	0.029*	0.537	0.325-0.888	0.015*
	Somali	0.45	0.243-0.827	0.010*	0.245	0.112-0.535	0.000*
	SNNPRS	0.99	0.617-1.577	0.954	0.702	0.368-1.374	0.302
	Tigray	2.32	1.451-3.712	0.000*	1.016	0.473-2.181	0.968

3.4. Practice

The practice of the respondents to prevent COVI-19 was assessed by 5 questions. About 43% of the respondents never practice any of the COVID-19 prevention methods and only less than one fifth (19.1%) of the respondents follow COVID-19 prevention measures either usually or always. About a third (31.0%) of respondents never wash their hands with soap and 42.9% of the respondents never avoided touching their eyes, nose, and mouth with their unwashed or unsensitized hand. Four of five respondents (81.4%) do not keep 2-meter social distancing whenever they are out for essential reasons. More than half of the respondents never avoided social gatherings for religious, festivals, or other ceremonies since the introduction of the state of emergency (Table 8).

273 Table 8. Summary of respondents' response to COVID-19 practice questions (N=1570).

no	Practice questions	response n (%); N= 1570			
		never	sometimes	usually	always
1	I wash my hands with soap frequently	486 (31.0)	630 (40.1)	344 (21.9)	110 (7.0)
2	I avoided non-essential travel since the lockdown	682 (43.4)	598 (38.1)	214 (13.6)	76 (4.8)
3	I always keep the social distancing of at least 2-meters whenever I am out for essential reasons	710 (45.2)	568 (36.2)	228 (14.5)	64 (4.1)
4	I avoided social gatherings for religious, festivals, or other ceremonies since the state emergency is introduced	810 (51.6)	508(32.4)	190 (12.1)	62 (3.9)
5	I avoid touching my eyes, nose, and mouth with my unwashed hands.	678 (43.2)	686 (43.7)	160 (10.2)	46 (2,9)
	Overall	673 (42.9)	598 (38.1)	227 (14.5)	72 (4.6)

275 There was no statistically significant difference between gender, and among age and marital
 276 status in the implementation of COVID-19 preventive measures at individual level. On the
 277 other hand, there is statistically significant difference in practicing COVID-19 preventive
 278 measure among residence, education, occupation and region of the respondents ($P \leq 0.001$).
 279 While 27.6% of urban respondents practice COVID-19 prevention measures, only 16.8% of
 280 the corresponding rural residents practice these measures either usually or always. Only a
 281 small proportion of respondents practice COVID-19 prevention measures in SNNP, Gambela
 282 and Dire Dawa compared to respondents from other regions (Table 9).

283 Table 9. Respondents practice score of COVID-19 by demographic variables (N=1570).

Demographic variables		Response n (%)		Statistical test	P (two-sided)
		Never/sometimes	Usually/always	χ^2	
Sex	Male	663 (74.0)	233(26.0)	3.174	0.085
	Female	525 (77.9)	149(22.1)		
Age in ears	18-29	652 (75.8)	208 (24.2)	0.452	0.978
	30-39	249 (75.0)	83 (25.0)		
	40-49	132 (75.9)	42 (24.1)		
	50-59	99 (77.3)	29 (22.7)		
	60+	56 (73.7)	20 (26.3)		
Residence					
	Urban	792 (72.4)	302 (27.6)	21.007	0.000
	Rural	396 (83.2)	80 (16.8)		
Marital status	Single	486 (73.4)	176 (26.6)	3.429	0.180
	Married	692 (77.4)	202 (22.6)		
	Widow	10 (71.4)	4 (28.6)		
Occupation	Government employee	186 (60.8)	120 (39.2)	57.476	0. 000
	Self-employed	423 (77.2)	125 (22.8)		
	Farmer	219 (87.6)	31 (12.4)		
	Student	360 (77.3)	106 (22.7)		
Education	Illiterate	213 (81.9)	17 (18.1)	3.056	0.000
	Basic	240 (82.2)	52 (17.8)		
	Primary	114 (81.4)	26 (18.6)		
	Secondary	166 (76.1)	52 (23.9)		

	College/university	455 (68.9)	205 (31.1)		
Region/chartered city	Addis Ababa	185 (56.7)	141 (43.3)	95.236	0.000
	Afar	36 (85.7)	6 (14.3)		
	Amhara	155 (81.6)	35 (18.4)		
	Benishangual Gumuz	36 (78.3)	10 (21.7)		
	Dire Dawa	46 (88.5)	6 (11.5)		
	Harari	39 (84.8)	7 (15.2)		
	Gambela	60 (88.2)	8 (11.8)		
	Oromia	434 (78.6)	118 (21.4)		
	Somali	48 (80.0)	2 (20.0)		
	SNNPRS	80 (88.9)	10 (11.1)		
	Tigray	69 (70.4)	29 (29.6)		
	Overall	1188 (75.7)	382 (24.3)		

285 Multivariable logistic regression analysis has shown that residence, region, education and
 286 occupation are the most predicting factors (76.1%) of COVID-19 appropriate practical
 287 measures. Being rural resident has a probability of 43.0% not to practice COVID-19
 288 prevention measures than being urban resident. The odds of government employee are more
 289 than twice (AOD=2.268) compared with the odds of being farmer towards appropriate COVID-
 290 19 practical measures while the odds of being self-employed was not statistically significantly
 291 different compared with the odds of being a farmer. The odds of respondents from all other
 292 regions were significantly lower than the odds of Addis Ababa (Finfinnee) respondents (Table
 293 10).

Demographic variables		Good practice towards Covid-19 prevention					
		Univariable binary logistic regression			multivariable logistic regression		
		Crude OR	95% CI	P-value	adjusted OR	95% CI	P-value
Sex	male	1			1		
	Female	0.808	0.638-1.022	0.075	0.852	0.661-1.099	0.218
Age in ears	18-29	1			1		
	30-39	1.045	0.779-1.401	0.769	1.860	1.003-3.447	0.049
	40-49	0.997	0.682-1.459	0.989	1.587	0.861-2.995	0.139
	50-59	0.918	0.590-1.429	0.705	1.643	0.858-3.146	0.134
	60+	1.120	0.656-1.909	0.679	1.344	0.683-2.645	0.391
Residence	urban	1			1		
	rural	0.530	0.403-0.697	0.000*	0.570	0.413-0.786	0.001*
Marital status	single	1			1		
	married	.806	0.639-1.018	.070	1.224	0.342-4.383	0.757
	widow	1.105	0.342-3.567	.868	1.700	0.496-5.821	0.398
Occupation	farmer	1					
	government employee	4.558	2.934-7.081	0.000*	2.268	1.519-3.385	0.000*
	Self-employed	2.088	1.364-3.196	0.001*	1.102	0.584-2.081	0.765
	student	2.080	1.348-3.211	0.001*	1.372	0.875-2.151	0.168
Education	illiterate	1.000					
	basic	0.982	0.635-1.518	0.935	2.093	1.3923-3.146	0.000*
	primary	1.034	0.608-1.757	0.903	2.283	1.561-3.338	0.000*
	secondary	1.420	0.911-2.212	0.122	2.343	1.461-3.758	0.000*
	college/university	2.042	1,430-2.916	0.000*	1.494	1.036-2.153	0.031*
	Addis Ababa	1			1		

Region/chartered city	Afar	0.2187	0.090-0.533	0.001*	0.559	0.337-0.928	0.025*
	Amhara	0.2963	0.193-0.454	0.000*	0.236	0.095-0.585	0.002*
	Benishangul Gumuz	0.364	0.175-0.759	0.007*	0.367	0.234-0.573	0.000*
	Dire Dawa	0.1710	0.071-0.412	0.000*	0.519	0.371-0.727	0.000*
	Harari	0.235	0.102-0.452	0.001*	0.250	0.108-0.582	0.001*
	Gambela	0.175	0.081-0.378	0.000*	0.211	0.103-0.431	0.000*
	Oromia	0.357	0.265-0.481	0.000*	0.395	0.199-0.785	0.008*
	Somali	0.328	0.168-0.641	0.001*	0.234	0.107-0.515	0.000*
	SNNPRS	0.164	0.082-0.328	0.000*	0.406	0.191-0.861	0.019*
	Tigray	0.551	0.339-0.896	0.016*	0.154	0.064-0.373	0.000*

4. Discussions

COVID-19 is an infectious disease caused by severe acute respiratory coronavirus 2 (SARS-CoV-2) and its being pandemic poses a significant threat to public health (9). Given that the disease is pandemic threat with no vaccine or proven treatment drug, preventive measures are the most essential methods available to reduce the infection rates and control the spread of the disease. This implies that public adherence to preventive and control measures are essential to curb the disease. The extent of public adherence to preventive and control measures is affected by their knowledge, attitude and practice (KAP). Therefore, this study was envisioned to assess the KAP of the Ethiopian population for the control of COVID-19.

The finding has shown that the participants level of COVID-19 knowledge is unsatisfactory. Only 42% of the respondents achieved a satisfactory knowledge score and the result is lower than most previous reports from various countries (2, 13, 19, 23-25). Our study has shown that Ethiopians had lower level of COVID-19 knowledge than Chinese (80.5%) ((25), Saudi Arabia (81.6%)(24), Bangladesh (54-87%) (26), Malaysia (80.5%) (23) and Tanzania (84.4%) (27) but similar with a study reported from Cameroon (43,7%) (28).

Poor level of knowledge of Ethiopians towards COVID-19 could make the spread of the pandemic worst coupled with the country's limited capacity of testing, contact tracing and isolation of suspected cases. Until the end of June 2020, Ethiopia has tested only less than 300,000 peoples among the 114 million population indicating that Ethiopia is far from achieving the pandemic control.

Knowledge scores significantly differ across socio-demographic factors which underlines the need for targeted public awareness campaign. Although most of the respondents (73.6%) have heard about COVID-19, only 42.4% and 37.8% knew COVID-19 transmission and clinical manifestations, respectively. Furthermore, only less than 20% of the respondents knew that

currently there is no effective drug to cure COVID-19 nor vaccines to prevent the infection. This poor level of knowledge towards COVID-19 could be due to limited engagement of Ethiopian institutions, religious sectors, political parties and policy-decision makers to generate contextual knowledge platform to provide an easy and accessible ways of getting credible information. It could be also due to top-down community awareness campaigns, lack of engaging diverse groups by the government, communication overload as large and conflicting information are directed at federal, regional state, and local health bureau from multiple sources which challenges the credibility and acceptability of the information by the general public. However, these proposed scenarios need further investigation.

Multivariable logistic regression analysis has shown that respondents from Oromia and Amhara regions are 55% and 50% more likely to be less knowledgeable about COVID-19 respectively than other regions. The poor level of COVID-19 knowledge in these two regions could be due to the vastness of these regions to address every corner of the regions adequately using the limited resources and infrastructure the country has. The sporadic protests and instability in these two regions cannot also be ignored affecting the consolidation of the information delivered from the government. This finding is very important since it may indicate that more effort should be exerted to deliver the desired message in these two most populous regions of the country, which account for more than 65% of the population of the country, (29) to control the spread of the disease.

Majority of the respondents (59.6%) have no confidence on the government to control the nationwide outbreak of the disease and 57.1% of the respondents disagree with the current introduction of state of emergency to curb the spread of COVID-19 by the government. Unexpectedly, half of the respondents do not agree with the principles of COVID-19 prevention methods such as hand washing, social distancing, avoiding non-essential travel and self-isolation symptomatic patients. As a result, only one in five respondents practice COVID-

19 prevention measures satisfactorily. It is very difficult to explain why the attitude and practice of the public to fight COVID-19 is low unlike in other countries such as China (25), Egypt (30), Sudan (31) and Tanzania (27). One reason could be the top-down flow of government procedures to contain the virus may be perceived as authoritarian attempts by government to consolidate control, leading to a loss of faith in the government. This pinpoints that the government need to re-examine the mode of delivery of COVID-19 related messages to the public. Another reason could be related to poverty and food insecurity. The state of emergency weigh most heavily on the poor, who are often part of the informal economy and thus dependent on day-to-day making money to feed themselves and their families. For many of them, a day without work means a day without food. Hence, they may resist to follow government advice.

The attitude of the respondents toward COVID-19 significantly varies across some sociodemographic factors. Although rural residents had lower level of COVID-19 knowledge, a significantly higher proportion of (59.2%) them had favourable attitude than urban residents (37.1%) towards COVID-19. Multivariate logistic regression analysis has shown that the odds of favourable attitude of being a rural resident was twice than being an urban resident. This could be due to the fact that rural residents have limited access to various information sources particularly on social media (32). Hence, they largely depend on the information from the local and national governors. Rural residents are also less likely than the urban residents to translate service dissatisfaction into discontent with their government and hence have more trust in government and evaluate information from local and national officials more positively than urban peers (33).

Only one in 5 of the respondents adhered to COVID-19 prevention measures either usually or always. While a third of respondents never wash their hands with soap, more than 80% of the respondents had never applied a 2-meter social distancing rule at least once. Ethiopia is among one of the poorest sub-Sharan countries where a significant proportion of the population (70-

80%) have no access to adequate water supply, sanitation and hygiene facilities (34) which challenges the handwashing practice to limit the spreading of COVID-19. Maintaining social distance is unattainable practice in Ethiopia since majority of the citizens live in slums. About 80% of Addis Ababa, the capital of the country, is considered slum areas, characterized by widespread sanitation challenges where families live in crowded rooms and are exposed to health and safety risks (35).

The ability of the government to persuade people to internalize the externality they would impose on the community by not reducing their mobility is one of the most important elements of compliance (36). This is particularly important where the ability to enforce lockdowns on a large scale is unattainable. Evidence from different countries indicates that political beliefs, coupled with differences in media consumption, have important implications for risk perceptions and compliance with social distancing and the Ethiopian government should revisits its COVID-19 prevention strategy (36).

Strengthening of government measures, community-based sensitization and health education programs will go a long way to prepare the community to overcome the pandemic. Providing this education to the population will help fill the knowledge gaps reflected and correct the misperception regarding the attitude of the population towards the disease which ultimately improves the practice.

5. Conclusions and recommendations

In summary, the present study was able to provide a comprehensive investigation of the knowledge, attitude and practice of Ethiopians toward COVID-19. The finding suggest that Ethiopians have inadequate level of knowledge on COVID-19 and are generally have a mixed outlook on overcoming the pandemic. The poor adherence to COVID-19 prevention practice by the public underlines the need for urgent reinforcing preventive measures and intensifying

394 sensitization campaigns to fill the knowledge gap of the population with regards to COVID-19
 395 and to persuade people to follow the preventive measures set by the government. At the same
 396 time, there is a need to evaluate the impact of these measures on the knowledge and practices
 397 of the population with the progression of the pandemic in Ethiopia. Finally, we believe that the
 398 study will inspire the healthcare authorities, and media to spread more COVID-19 related
 399 accurate knowledge which ultimately results in better preventive practices toward COVID-19.
 400 Since no proven medicine or vaccine is developed yet, the best way to curb the spreading of
 401 the disease is maximizing knowledge and preventive practices toward COVID-19 at all levels.

402 6. References

- 403 1. Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, et al. A Novel Coronavirus from Patients with
404 Pneumonia in China, 2019. *N Engl J Med*. 2020;382(8):727-33.
- 405 2. Olum R, Chekwech G, Wekha G, Nassozi DR, Bongomin F. Coronavirus Disease-2019:
406 Knowledge, Attitude, and Practices of Health Care Workers at Makerere University Teaching
407 Hospitals, Uganda. *Frontiers in Public Health*. 2020;8(181).
- 408 3. Siordia JA. Epidemiology and clinical features of COVID-19: A review of current literature.
409 *Journal of Clinical Virology*. 2020;127:104357.
- 410 4. Zhang R, Li Y, Zhang AL, Wang Y, Molina MJ. Identifying airborne transmission as the
411 dominant route for the spread of COVID-19. *Proceedings of the National Academy of Sciences*.
412 2020:202009637.
- 413 5. van Doremalen N, Bushmaker T, Morris DH, Holbrook MG, Gamble A, Williamson BN, et al.
414 Aerosol and Surface Stability of SARS-CoV-2 as Compared with SARS-CoV-1. *New England Journal of*
415 *Medicine*. 2020;382(16):1564-7.
- 416 6. Chin AWH, Chu JTS, Perera MRA, Hui KPY, Yen H-L, Chan MCW, et al. Stability of SARS-CoV-2
417 in different environmental conditions. *The Lancet Microbe*. 2020;1(1):e10.
- 418 7. Chan JF-W, Yuan S, Kok K-H, To KK-W, Chu H, Yang J, et al. A familial cluster of pneumonia
419 associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a
420 family cluster. *The Lancet*. 2020;395(10223):514-23.
- 421 8. He X, Lau EHY, Wu P, Deng X, Wang J, Hao X, et al. Temporal dynamics in viral shedding and
422 transmissibility of COVID-19. *Nat Med*. 2020;26(5):672-5.
- 423 9. Guan W-j, Ni Z-y, Hu Y, Liang W-h, Ou C-q, He J-x, et al. Clinical characteristics of 2019 novel
424 coronavirus infection in China. *medRxiv*. 2020:2020.02.06.20020974.
- 425 10. WHO. FIRST CASE OF COVID-19 CONFIRMED IN ETHIOPIA". WHO | Regional Office for Africa,
426 . Retrieved June 15, 2020. 2020.
- 427 11. Yesuf N. Ethiopia needs a State of Emergency to combat COVID-19 Addis Ababa, Ethiopia
428 Ethiopia Insight 2020 [
- 429 12. Manyazewal T, Matlakala MC. Implementing health care reform: implications for
430 performance of public hospitals in central Ethiopia. *J Glob Health*. 2018;8(1):010403-.
- 431 13. Wolf MS, Serper M, Opsasnick L, O'Connor RM, Curtis LM, Benavente JY, et al. Awareness,
432 Attitudes, and Actions Related to COVID-19 Among Adults With Chronic Conditions at the Onset of
433 the U.S. Outbreak: A Cross-sectional Survey. *Ann Intern Med*. 2020.
- 434 14. Paasche-Orlow MK, Parker RM, Gazmararian JA, Nielsen-Bohman LT, Rudd RR. The
435 prevalence of limited health literacy. *J Gen Intern Med*. 2005;20(2):175-84.
- 436 15. Heaton TB, Crookston B, Pierce H, Amoateng AY. Social inequality and children's health in
437 Africa: a cross sectional study. *International Journal for Equity in Health*. 2016;15(1):92.
- 438 16. Deaton AS, Tortora R. People in sub-Saharan Africa rate their health and health care among
439 the lowest in the world. *Health Aff (Millwood)*. 2015;34(3):519-27.
- 440 17. Coates MM, Kamanda M, Kintu A, Arikpo I, Chauque A, Mengesha MM, et al. A comparison
441 of all-cause and cause-specific mortality by household socioeconomic status across seven INDEPTH
442 network health and demographic surveillance systems in sub-Saharan Africa. *Global Health Action*.
443 2019;12(1):1608013.
- 444 18. Birkhäuser J, Gaab J, Kossowsky J, Hasler S, Krummenacher P, Werner C, et al. Trust in the
445 health care professional and health outcome: A meta-analysis. *PloS one*. 2017;12(2):e0170988.
- 446 19. Yap J, Lee VJ, Yau TY, Ng TP, Tor P-C. Knowledge, attitudes and practices towards pandemic
447 influenza among cases, close contacts, and healthcare workers in tropical Singapore: a cross-
448 sectional survey. *BMC Public Health*. 2010;10(1):442.
- 449 20. Reuters T. Ethiopia and Liberia declare states of emergency to curb coronavirus. Health
450 NEWS. Addis Ababa2020.

21. Lancaster H. Ethiopia - Telecoms, Mobile and Broadband - Statistics and Analyses. Sydney, AUSTRALIA: Paul Budde Communication Pty Ltd; 2020.
22. Max R, Hannah R, Esteban O-O. How many internet users does each country have? accessible at <https://ourworldindata.org/internet>. 2020.
23. Azlan AA, Hamzah MR, Sern TJ, Ayub SH, Mohamad E. Public knowledge, attitudes and practices towards COVID-19: A cross-sectional study in Malaysia. *PloS one*. 2020;15(5):e0233668.
24. Al-Hanawi MK, Angawi K, Alshareef N, Qattan AMN, Helmy HZ, Abudawood Y, et al. Knowledge, Attitude and Practice Toward COVID-19 Among the Public in the Kingdom of Saudi Arabia: A Cross-Sectional Study. *Frontiers in Public Health*. 2020;8(217).
25. Zhong B-L, Luo W, Li H-M, Zhang Q-Q, Liu X-G, Li W-T, et al. Knowledge, attitudes, and practices towards COVID-19 among Chinese residents during the rapid rise period of the COVID-19 outbreak: a quick online cross-sectional survey. *International Journal of Biological Sciences*. 2020;16(10):1745-52.
26. Karim A, Akter M, Mazid AHMT, Pullock OS, Aziz TT, Hayee S, et al. Knowledge and attitude towards COVID-19 in Bangladesh: Population-level estimation and a comparison of data obtained by phone and online survey methods. *medRxiv*; 2020.
27. rugarabamu s, Byanaku A, Ibrahim M. Knowledge, attitudes, and practices (KAP) towards COVID-19: A quick online cross-sectional survey among Tanzanian residents. *medRxiv*. 2020:2020.04.26.20080820.
28. Nicholas T, Mandaah VF, Esemu K, S, Vanessa ABT, Gilchrist KTD, Vanessa LF, et al. CVOVID-19 knowledge, attitudes and practices in a conflict affected area of the South West Region of Cameroon. . *Pan African Medical Journal*. 2020;35(34).
29. Baker J. State, Governance and the Creation of Small Towns in Ethiopia. *The European Journal of Development Research*. 2019;31(1):34-52.
30. Abdelhafiz AS, Mohammed Z, Ibrahim ME, Ziady HH, Alorabi M, Ayyad M, et al. Knowledge, Perceptions, and Attitude of Egyptians Towards the Novel Coronavirus Disease (COVID-19). *J Community Health*. 2020:1-10.
31. Khawla Nasr Aldeen Altayb M, Malaz Malik Yousif S, Mohammed Tayseer Bashir A. Knowledge, attitudes, and practices surrounding COVID-19 among Sudan citizens during the pandemic: an online cross-sectional study. *Sudan Journal of Medical Sciences (SJMS)*. 2020;15(5).
32. Amoah PA, Edusei J, Amuzu D. Social Networks and Health: Understanding the Nuances of Healthcare Access between Urban and Rural Populations. *Int J Environ Res Public Health*. 2018;15(5):973.
33. Brinkerhoff DW, Wetterberg A, Wibbels E. Distance, services, and citizen perceptions of the state in rural Africa. *Governance*. 2018;31(1):103-24.
34. Admasie A, Debebe A. Estimating Access to Drinking Water Supply, Sanitation, and Hygiene Facilities in Wolaita Sodo Town, Southern Ethiopia, in Reference to National Coverage. *Journal of Environmental and Public Health*. 2016;2016:8141658.
35. Habitat. Slums in Addis Ababa: the road ahead. <https://www.habitatforhumanity.org.uk/blog/2017/11/slum-housing-ethiopia-modernise/> 2017 [
36. Bavel JJV, Baicker K, Boggio PS, Capraro V, Cichocka A, Cikara M, et al. Using social and behavioural science to support COVID-19 pandemic response. *Nature Human Behaviour*. 2020;4(5):460-71.