1 Title: Development of a free radical scavenging probiotic to mitigate coral bleaching

2 Running title: Making a probiotic to mitigate coral bleaching

3

4 Ashley M. Dungan^{a#}, Dieter Bulach^b, Heyu Lin^c, Madeleine J. H. van Oppen^{a,d}, Linda L. Blackall^a

5

6 ^aSchool of Biosciences, The University of Melbourne, Melbourne, VIC, Australia

7 ^bMelbourne Bioinformatics, The University of Melbourne, Melbourne, VIC, Australia

8 ^cSchool of Earth Sciences, The University of Melbourne, Melbourne, VIC, Australia

9 ^dAustralian Institute of Marine Science, Townsville, QLD, Australia

10

11

12 [#]Address correspondence to Ashley M. Dungan, adungan31@gmail.com

13

14 Abstract word count: 211 words

15 Text word count: 4838 words

16

17 Keywords: symbiosis, *Exaiptasia diaphana*, *Exaiptasia pallida*, probiotic, antioxidant, ROS,
18 *Symbiodiniaceae*, bacteria

ABSTRACT

Corals are colonized by symbiotic microorganisms that exert a profound influence on the animal's health. One noted symbiont is a single-celled alga (from the family *Symbiodiniaceae*), which provides the coral with most of its fixed carbon. During thermal stress, hyperactivity of photosynthesis results in a toxic accumulation of reactive oxygen species (ROS). If not scavenged by the antioxidant network, ROS may trigger a signaling cascade ending with the coral host and algal symbiont disassociating; this process is known as bleaching. Our goal was to construct a probiotic comprised of host-associated bacteria able to neutralize free radicals such as ROS. Using the coral model, the anemone *Exaiptasia diaphana*, and pure bacterial cultures isolated from the model animal, we identified six strains with high free radical scavenging ability belonging to the families *Alteromonadaceae*, *Rhodobacteraceae*, *Flavobacteriaceae*, and *Micrococcaceae*. In parallel, we established a "negative" probiotic consisting of genetically related strains with poor free radical scavenging capacities. From their whole genome sequences, we explored genes of interest that may contribute to their potential beneficial roles, which may help facilitate the therapeutic application of a bacterial probiotic. In particular, the occurrence of key pathways that are known to influence ROS in each of the strains has been inferred from the genomes sequences and are reported here.

IMPORTANCE

Coral bleaching is tightly linked to the production of reactive oxygen species (ROS), which accumulates to a toxic level in algae-harboring host cells leading to coral-algal dissociation. Interventions targeting ROS accumulation, such as the application of exogenous antioxidants, have shown promise for maintaining the coral-algal partnership. With the feasibility of administering antioxidants directly to corals being low, we aim to develop a probiotic to neutralize toxic ROS during a thermal stress event. This probiotic can be tested with corals or a coral model to assess its efficacy in improving coral resistance to environmental stress.

INTRODUCTION

Coral reefs are among the most biologically and economically valuable ecosystems on Earth (1-3). While they cover less than 0.1% of the ocean floor (4), coral reefs support fisheries, tourism, pharmaceuticals and coastal development with a global value of \$8.9 trillion “international \$”/year (5). Corals and other reef organisms have been dying, largely due to anthropogenic influences such as climate change (6, 7), which has led to an increased frequency, intensity and duration of summer heat waves that cause coral bleaching (8, 9).

The coral holobiont, which is the sum of the coral animal and its associated microbiota, including algae, fungi, protozoans, bacteria, archaea and viruses (10), is an ecosystem engineer. By secreting a calcium carbonate skeleton, the reef structure rises from the ocean floor, forming the literal foundation of the coral reef ecosystem. The success of corals to survive and build up reefs over thousands of years (11) is tightly linked to their obligate yet fragile symbiosis with endosymbiotic dinoflagellates of the family *Symbiodiniaceae* (12).

Intracellular *Symbiodiniaceae* translocate photosynthetically fixed carbon to the coral host (13, 14) in exchange for inorganic nutrients and location in a high light environment with protection from herbivory (15, 16). During periods of thermal stress, the relationship between the coral host and their *Symbiodiniaceae* can break down, resulting in a separation of the partners and significantly, fixed carbon shortage for the host. This phenomenon, ‘coral bleaching’, is devastating to the host and detrimental to the reef system. The ecosystem-wide effects of bleaching on the coral include reduced skeletal growth and reproductive activity, a lowered capacity to shed sediments, and an inability to resist invasion of competing species and diseases. Severe and prolonged bleaching can cause partial to total colony death, resulting in diminished reef growth, the transformation of reef-building communities to alternate, non-reef building community types, bioerosion and ultimately the disappearance of reef structures (12).

There are several hypotheses detailing the mechanisms driving bleaching (see 17, 18-20), with a common theme being the overproduction and toxic accumulation of reactive oxygen species (ROS) from the algal symbiont. Excess ROS are generated by a number of pathways including

heat damage to both chloroplast and mitochondrial membranes (21, 22), and are shown to play a central role in inter-partner communication of a stress response (17). Once generated, ROS causes damage to many cell components including photosystem II reaction centers in the *Symbiodiniaceae*, specifically at the D1 and D2 proteins (23, 24), thus interfering with the supply of fixed carbon to the holobiont. Once damaged, *Symbiodiniaceae* are no longer able to maintain their role in the symbiotic relationship with corals and separate from the host tissue via *in situ* degradation, exocytosis, host cell detachment, host cell apoptosis or host cell necrosis (17).

Probiotics are preparations of viable microorganisms that are introduced to a host to alter their microbial community in a way that is beneficial to the system. Microbiome engineering through the addition of probiotics has been postulated as a key strategy to facilitate adaptation to changing environmental conditions by enhancing corals with the metabolic capabilities of the introduced probiotic bacterial strains (25-30). The differences in the bacterial species composition of healthy and thermally stressed corals (31-36) and the coral model *Exaiptasia diaphana* (37-39) suggest a role for microbiome engineering in cnidarian health. A disruption to the bacterial community of *Pocillopora damicornis* with antibiotic treatment diminished the resilience of the holobiont during thermal stress, whereas intact microbial communities conferred resilience to thermal stress and increased the rate of holobiont recovery after bleaching events (40). The relative stability of coral-associated bacterial communities has also been linked to coral heat tolerance; for instance, the bacterial community of heat sensitive *Acropora hyacinthus* corals shifted when transplanted to thermal stress conditions, whereas heat-tolerant *A. hyacinthus* corals harbored a stable bacterial community (41).

In recent years, researchers have begun to explore the use of probiotics in corals and the model organism for corals, *E. diaphana*. To inhibit the progression of white pox disease, caused by pathogenic *Serratia marcescens*, an *Alphaproteobacteria* cocktail containing several *Marinobacter* spp. isolates was applied to *E. diaphana* (42). These introduced strains were able to inhibit both biofilm formation and swarming of *S. marcescens*, which halted disease progression. The *Marinobacter*-based probiotic was deemed effective as anemones exposed to

both the cocktail and pathogen survived after seven days, while anemones in the *S. marcescens* control treatment died. A bacterial consortium native to the coral *Mussismilia harttii* was selected to degrade water-soluble oil fractions(43). This bioremediation strategy reduced the negative impacts of oil on *M. harttii* health and accelerated the degradation of petroleum hydrocarbons (43). Coral microbiomes have also been manipulated through addition of a consortium of native or seawater derived bacteria to the surface of *P. damicornis* to mitigate the effects of thermal stress (44). The results from this study suggest the consortium was able to partially mitigate coral bleaching.

Our goal was to identify bacterial strains suitable for use in a probiotic to mitigate the effects of thermal stress in *E. diaphana*. Given the potential role of ROS in the bleaching process, our focus was to select diverse *E. diaphana*-sourced bacterial strains with antioxidant properties while avoiding potential pathogens. Antioxidant properties were measured using the stable free radical 2,2-diphenyl-1-picrylhydrazyl (DPPH), which is reduced in the presence of an antioxidant molecule, undergoing a color change from a violet to a colorless solution.

RESULTS

Diversity of culturable bacteria associated with *E. diaphana*. A total of 842 isolates were obtained from four genotypes of Great Barrier Reef (GBR)-sourced *E. diaphana*. There were no significant differences in bacterial colony forming units (CFUs) between the four genotypes, regardless of growth medium, with 5.9-10.3 x 10³ cells per anemone on Reasoner's 2A agar (R2A) and 6.3-10.4 x10³ cells per anemone on marine agar (MA) (p>0.05). Partial 16S rRNA gene sequences (~1000 bp) were used to identify the closest matches from the NCBI database using the Basic Local Alignment Search Tool (BLASTn). In total there were 109 species in 64 genera, 27 families and six phyla (Fig. 1). The most abundant genera were *Alteromonas*, *Labrenzia*, and *Ruegeria* (Table 1). Gram-positive bacteria comprised 23 species, including *Microbacterium* (31 isolates) and *Micrococcus* (28 isolates). Eight genera were found to be associated with all four genotypes (Table 1); these eight genera made up 59.4% of all *E. diaphana*-associated bacterial isolates.

Table 1: Bacterial genera associated with all four genotypes of GBR-sourced *E. diaphana* (AIMS1-4).

Genus	Class	No. of species	AIMS1	AIMS2	AIMS3	AIMS4	*Total Isolates
<i>Alteromonas</i>	<i>Gamma-proteobacteria</i>	6	10	52	24	30	116
<i>Labrenzia</i>	<i>Alpha-proteobacteria</i>	4	11	10	26	38	85
<i>Marinobacter</i>	<i>Gamma-proteobacteria</i>	4	7	25	10	13	55
<i>Muricauda</i>	<i>Flavobacteriia</i>	1	16	11	10	5	42
<i>Roseovarius</i>	<i>Alpha-proteobacteria</i>	3	9	12	5	6	32
<i>Ruegeria</i>	<i>Alpha-proteobacteria</i>	3	29	8	5	39	81
<i>Shimia</i>	<i>Alpha-proteobacteria</i>	2	2	3	9	40	54
<i>Vibrio</i>	<i>Gamma-proteobacteria</i>	3	3	7	31	15	56

*521 out of the 842 isolates obtained from the four *E. diaphana* genotypes sourced from the Great Barrier Reef.

FIG 1: Neighbor-Joining tree showing an overview of the phylogenetic relationship between the 842 *E. diaphana*-associated bacterial isolates inferred using partial 16S rRNA sequences. These isolates covered six phyla indicated by shading over the tree with *Proteobacteria* split into the classes *Gammaproteobacteria* and *Alphaproteobacteria*. The positions of selected probiotic strains are highlighted by arrows with blue arrows indicating the high FRS strains and orange arrows indicating low FRS strains.

Bacterial probiotic selection. Of the original 842 isolates, 709 were screened for their ability to scavenge free radicals. Those isolates were divided into three categories, positive (144), weakly positive (121), and negative (444). Ninety-eight strains representing eight families and 18 genera were then quantitatively assessed for FRS. There was no clear pattern of free radical scavenging (FRS) capacity at the family level (Fig. 2) with strain-specific responses evident. From these isolates, probiotic members were selected by choosing *E. diaphana*-associated bacteria, where conspecific or congeneric pairs of strains displayed a high (“positive”) and low (“negative”) FRS ability (Fig. 2; Table 2). Of the 12 selected probiotic members, seven were catalase positive and five were catalase negative (Table 2). In each probiotic set (i.e., high or low FRS strains), none of the selected isolates showed antagonistic activity against one other as evidenced by the absence of any zone of inhibition and growth from each combination of isolates on a plate.

149

150 **FIG 2:** Quantitative FRS ability of *E. diaphana*-associated bacteria isolates, separated by Family. Families with high relative
 151 abundance among all cultured bacteria (*Rhodobacteraceae* – A, *Alteromonadaceae* – B, *Pseudoalteromonadaceae* – C,
 152 *Flavobacteriaceae* – D, and *Micrococcaceae* – E) were separately analyzed to identify strains with a high FRS ability (blue arrows) and
 153 a corresponding conspecific or congeneric strain with a low FRS ability (orange arrows). In each panel, the light dashed vertical line
 154 on the left represents the mean FRS of a 0.025% (w/v) ascorbic acid standard, the middle dark dashed vertical line is the mean FRS
 155 for 0.05% (w/v) ascorbic acid standard, and the far right dashed line is the mean FRS of the 0.075% (w/v) ascorbic acid standard.

156 **Table 2:** Overview of probiotic high FRS bacterial strains and conspecific/congeneric low FRS control strains. All sequence data can
157 be found under BioProject# PRJNA574193. References to each probiotic candidate at the Genus level are identified in the last three
158 columns.

Strain	Bacteria species (Phylum or Class)	#FRS (% $\pm$ SE)	*Positive vs Negative	*Strain vs Growth Medium	Catalase	<i>E. diaphana</i> Literature	Coral Literature	Probiotic Literature
MMSF01163 (n=3)	<i>Alteromonas oceani</i> (Gamma- proteobacteria)	61.7 $\pm$ 5.2	0.065	0.004	Negative	(45-47)	(27, 48- 50)	(51-54)
MMSF00404 (n=4)		35.5 $\pm$ 13.7		0.571	Negative			
MMSF00958 (n=3)	<i>Alteromonas macleodii</i> (Gamma- proteobacteria)	62.0 $\pm$ 4.2	0.175	0.017	Negative			
MMSF00257 (n=3)		30.3 $\pm$ 0.9		0.431	Positive			
MMSF00132 (n=8)	<i>Labrenzia aggregata</i> (55) (Alpha- proteobacteria)	53.6 $\pm$ 8.7	0.016	0.101	Negative	(46)	(35, 50)	(56)
MMSF00249 (n=3)		14.0 $\pm$ 3.6		0.103	Positive			
MMSF01190 (n=3)	<i>Marinobacter salsuginis</i> (Gamma- proteobacteria)	62.0 $\pm$ 4.5	0.041	<0.0001	Negative	(45, 46, 57)	(49, 58, 59)	(42)
MMSF00964 (n=3)		43.7 $\pm$ 5.8		0.020	Positive			
MMSF00068 (n=6)	<i>Micrococcus luteus</i> (Actinobacteria)	56.3 $\pm$ 7.3	0.210	0.002	Positive	(46)	(60)	(51, 61, 62)
MMSF00107 (n=3)	<i>Micrococcus yunnanensis</i> (Actinobacteria)	38.0 $\pm$ 8.0		0.306	Positive			

MMSF00046 (n=3)	<i>Winogradskyella</i> <i>poriferorum</i> (63)	73.3 ± 1.9	0.284	0.010	Negative	^	(43, 64)	NA
MMSF00910 (n=3)	(<i>Bacteroidetes</i>)	36.0 ± 3.5		0.174	Negative			

159 # Mean R2A FRS (% ± SE) was 27.2 ± 2.3% (n=12) and it was catalase negative.

160 ^While there are no instances of *Winogradskyella* specifically identified in current *E. diaphana* literature, there are several
161 *Flavobacteriaceae* that are not resolved to the genus level from metabarcoding data (38, 39, 45, 47).

162 *Indicates p values for pairwise comparisons from respective one-way analysis of variance (Tukey HSD) or Kruskal-Wallis rank sum
163 test (Dunn test) with **bold** values representing significant differences.

Comparative genomics. As part of the characterization of the 12 isolates (six positive and six conspecific or congeneric negative FRS isolates), a draft genome sequence was determined for each isolate. A summary of the data and metrics for the draft genome sequences is presented in Table S1. The diversity of the six pairs of isolates is indicated by the %G+C range (35% to 72%) and genome size (2.4 Mb to 6.8 Mb). Each isolate pair was classified as the same species except the *Micrococcus* stains. Isolates MMSF00068 (high FRS strain) and MMSF00107 (low FRS strain) are classified as *Micrococcus luteus* by both 16S rRNA gene and genome-based methods, but MMSF00107 is classified as *Micrococcus yunnanensis* by NCBI. These isolates have the smallest genomes among the isolates (2.43 Mb and 2.48 Mb, respectively) and the highest G+C content of 72.8% and 72.4%, respectively. The classification of isolate MMSF00107 is uncertain, particularly given that the genome sequence for the type strain for *M. yunnanensis* is not yet available.

Core genome comparison of the draft genome sequences provides an overview of the genetic relationship between the conspecific or congeneric isolate pairs. A wide range of genome variation between pairs was observed, this ranged from ~190,000 core genome single nucleotide polymorphisms (SNP) differences between the *Alteromonas oceani* strains, to fewer than five core genome SNPs between the *Labrenzia aggregata* isolates (Table S2). Importantly, this is not a comprehensive estimate of isolate identity, e.g. the genome sequences of *Winogradskyella poriferorum* isolates MMSF00910 and MMSF00046 are nearly identical with fewer than ten pairwise SNP differences in the core genome; but there are accessory genome differences.

Genes of interest. The annotated genome sequences of each selected probiotic member were searched for key genes of interest (Table S3-S4). Dimethylsulfoniopropionate (DMSP) cleavage to dimethylsulfide (DMS) was identified by presence of one or more of the DMSP lyase genes; *dddP*, *dddD*, *dddL*, *dddW*, and *dddQ*. Only *L. aggregata* strains contained DMSP lyase genes (*dddP* and *dddL*) in their whole genome sequences. DMSP biosynthesis was identified by the presence of *dsyB*, which is the only described gene for an enzyme in the DMSP biosynthesis pathway. The *cobP* gene was used as an indicator for the presence of the dynamic vitamin B₁₂

pathway, which contains 27 genes (Table S4). Again, only the *L. aggregata* isolates contained *cobP*. Catalase positive strains were identified by the presence of *katG*; all positive and negative FRS strains contained *katG* except the *Micrococcus* spp. strains, in which *kata* and *katE* were detected.

16S rRNA gene copy number.

The 16S rRNA gene copy numbers of the 12 draft genomes were estimated using a read depth approach (Table S1). The copy numbers were similar within pairs of isolates, in which the pair of isolates MMSF00257 and MMSF00958 contained the most copies (5.15 and 4.79, respectively), corresponding with copy numbers in a known closed genome of *A. macleodii* (65). *Marinobacter* isolates MMSF00046 and MMSF00910 contained the fewest copies (1.03 and 0.77, respectively).

DISCUSSION

The 842 *E. diaphana* bacterial isolates reported here comprise 109 species from 64 genera and six phyla. Using metabarcoding, studies of microbiomes associated with *E. diaphana* have revealed a similar diversity at the phylum level for *E. diaphana* sourced from the GBR (37), Hawaii (strain H2; 45), Pacific and Caribbean (57), Atlantic (strain CC7; 47) and Red Sea (38), as well as stony corals (66). Thus, our culture collection of *E. diaphana* bacterial isolates captures the diversity of the *E. diaphana*-associated microbiome. The *E. diaphana* probiotics generated in this study were assembled from 12 isolated bacterial strains selected based on their FRS ability since free radical production, specifically ROS, is relevant in coral bleaching. The broader culture collection contains bacteria with a wide range of FRS capacity, and our probiotic comprises greater bacterial diversity compared to others (42, 44).

The consistent and frequent reporting of our probiotic bacteria genera in *E. diaphana* and coral studies (Table 2) suggests these bacteria likely have key functions in cnidarian holobionts. Among these potential functions are the production of the antioxidant DMSP and the breakdown of DMSP to other antioxidants (67). *L. aggregata* has been reported to produce

DMSP in the absence of any methylated sulfur compounds with *dsyB* identified as the first DMSP biosynthesis gene in any organism (68). The *dsyB* gene was found in the genome sequences of both high and low FRS *L. aggregata* strains (Table S3). Many *E. diaphana*-sourced bacterial species, specifically relatives of our selected probiotic members, are implicated in the degradation of DMSP to DMS (*Alteromonas* spp., (69); *Labrenzia* spp., (70)). *dddP* codes for the enzyme responsible for cleaving DMSP to DMS and acrylate and was used (from the Prokka annotation) as an indicator of a DMSP degradation genotype. Only the *L. aggregata* isolates contained genes responsible for DMSP degradation (Table S3).

Carotenoids are among the strongest antioxidants and are highly reactive against both ROS and free radicals (71-75). Carotenoids are lipid-soluble pigments, and in bacteria they give an orange-yellow hue to colonies. Two of the five selected probiotic genera produce orange/yellow colonies (*Winogradskyella*, *Micrococcus*), and there is evidence of carotenoid production by marine *Flavobacteriaceae* (75, 76) and *Micrococcus* (77). A marine *Flavobacteriaceae* (strain GF1) was found to produce the potent antioxidant carotenoid zeaxanthin that protected *Symbiodiniaceae* from thermal and light stress (78).

Vitamin B₁₂ is a cofactor involved in the production of the amino acid methionine, which is needed to synthesize every protein and in diverse metabolic pathways including generation of the antioxidants glutathione and DMSP (79). Vitamin B₁₂ is synthesized by many heterotrophic bacteria (80). Genomic evidence suggests that *Symbiodiniaceae* have lost the capacity to synthesize vitamin B₁₂ (81), which is in agreement with other work showing that free-living *Symbiodiniaceae* rely on bacterial symbionts for this important cofactor (82). The genes involved in the biosynthesis of vitamin B₁₂ have been found in coral-associated bacteria, specifically *L. aggregata* cultured from the Caribbean coral, *Orbicella faveolata* (83). Eighteen genes, including the *cobP* gene (80), are annotated in each of *L. aggregata* isolates (MMSF00132 & MMSF00249), suggesting both are capable of vitamin B₁₂ biosynthesis. None of the other sequenced isolates had genes related to vitamin B₁₂ biosynthesis, except in the *Marinobacter salsuginis* isolates, where a *cobO* gene was detected, strongly indicating that none of the remaining ten isolates were capable of vitamin B₁₂ biosynthesis (Table S4).

Bacteria have developed highly specific mechanisms to protect themselves against oxidative stress, in particular using enzymatic antioxidants such as catalase/peroxidase and superoxide dismutase (84). It has been suggested that increasing the *in hospite* concentration of catalase in the coral holobiont by the application of a probiotic with catalase-positive organisms, could possibly minimize the impact of thermal stress by neutralizing hydrogen peroxide (H₂O₂) (29). Here we tested all probiotic candidates for catalase production using a standard H₂O₂ assay (85). Catalase participates in cellular antioxidant defense by enzymatically breaking down H₂O₂ to H₂O and O₂. Hydroperoxidase I (HPI, *katG*) is present during aerobic growth and is transcriptionally controlled at different levels, and hydroperoxidase II (HPII, *katE*) is induced during stationary phase (86, 87). The phenotypically determined catalase positive strains (MMSF00249, *L. aggregata*; MMSF00257, *A. macleodii*; MMSF00964, *M. salsuginis*) had a catalase-peroxidase gene (HPI, *katG*); but all strains had at least one *katG*, *katA* (catalase), or *katE* (catalase HPII) gene (Table S3). The lack of correlation between genotype and phenotype in relation to catalase activity may be associated with the culture conditions used or the bacterial growth phase during the catalase assay. The inconsistency between the catalase and DPPH phenotype and FRS ability (in many cases high FRS isolates were catalase negative while the low FRS strain was catalase positive), justifies our approach of using FRS analysis, instead of using the presence of catalase genes as a proxy for FRS.

A critical characteristic in the selection of probiotic members is the maintenance and proliferation of the inoculated bacteria in the host over time (51) and their potential transmission to the next generation. There is evidence that corals release bacteria with their offspring such as *Alteromonas* (48), *Flavobacteriaceae* (48), *Rhodobacteraceae* (88), and *Marinobacter* (58). While broadcast spawning corals do not appear to transfer their bacterial symbionts with their gametes (vertical transfer) (89), the brooding coral *Porites astreoides* transmits some bacteria vertically to planulae with two bacterial taxa (*Roseobacter* clade-associated bacteria and *Marinobacter* spp.) consistently and stably associating with juvenile *P. astreoides* (58). *Marinobacter* potentially has antioxidant properties (90), while *Roseobacter* spp. might be beneficial in facilitating larval settlement. If adult corals stably associate with

inoculated probiotic candidates like *Marinobacter*, *Alteromonas*, and *Winogradskyella*, they may be passed on to offspring and thus have a long-term positive impact on these individuals.

The probiotic members were chosen from a highly diverse pool of *E. diaphana*-sourced bacterial isolates. While the selected probiotic members are phylogenetically diverse, potentially promising probiotic bacteria in the culture collection were omitted based on our selection criteria. For example, *Ruegeria* spp. were reported to breakdown DMSP and participate in denitrification (83). *Muricauda* isolates had high FRS abilities, but they did not grow consistently in the selected medium and therefore were excluded from the consortium. *Muricauda* isolates have genes for denitrification (83), can breakdown DMSP (70), and produce potent carotenoids (91) that can mitigate thermal and light stresses in *Symbiodiniaceae* cultures (78). *Muricauda* will be included in future probiotic evaluations.

Interactions among members of the microbiota associated with marine animals are undoubtedly complex. Results presented in this manuscript show that pure cultured bacteria from *E. diaphana* can scavenge free radicals, albeit at a strain-specific rate. Inoculation with high FRS strains could be beneficial to the host under high oxidative stress conditions, such as those that contribute to coral bleaching. Outlined here is the start of a complex process to identify, evaluate, and select durable and useful probiotics that can buffer the coral host against climactic variation. Conspecific or congeneric pairs of bacteria provide an opportunity to determine the genetic basis for measured phenotypic differences between the pairs. An essential element of future work will be to investigate the stability of the phenotypic differences observed and this stability may be reflected in the nature of the genetic differences between the pairs of strains.

MATERIALS AND METHODS

Isolation of bacterial isolates. GBR origin *E. diaphana* were maintained in the laboratory at 26 °C (92) and used to isolate probiotic candidates. Sixteen individuals from each of four *E. diaphana* genotypes (AIMS1-4) were collected using sterile disposable pipets and gently transferred to filter-sterilized (0.2 µm) reverse osmosis (RO) water reconstituted Red Sea Salt™

(Red Sea; RSS) at ~34 parts per thousand (ppt) salinity (fRSS). This was done to remove some of the influence of the external seawater on the bacterial community. After 30 min, each anemone was transferred to a sterile glass homogenizer with 1 mL of fRSS. Each homogenate was used to prepare serial dilutions from 10^{-1} to 10^{-4} . From each dilution, 50 μ L was spread plated onto three replicate plates each of MA (Difco™ Marine Agar 2216) and R2A (CM0906, Oxoid) supplemented with 40 g L⁻¹ RSS and incubated at 26 °C. After one week, CFU counts were completed. Individual bacterial isolates were sub-cultured to purification from plates with <100 CFUs onto the initial isolation medium. All purified bacterial isolates were resuspended in 40% glycerol, aliquoted into 1.2 mL cryotubes and stored at -80 °C.

Identification of *E. diaphana*-sourced isolates. Colony PCR with the universal bacterial primers 27f (5' – AGA GTT TGA TCM TGG CTC AG – 3') and 1492r (5' – TAC GGY TAC CTT GTT ACG ACT T – 3') (93) was used to generate 16S rRNA gene amplicons from each isolate. Briefly, cells from each pure culture were suspended in 20 μ L Milli-Q water and denatured at 95 °C for 10 min. The suspension was then centrifuged at 2,000 x *g* at 4 °C for two minutes and the supernatant was used as the DNA template for PCR amplification. The PCR was performed with 20 μ L Mango Mix™ (Bioline), 0.25 μ M of each primer and 2 μ L of DNA template in a final volume of 40 μ L. The thermal cycling protocol was as follows: 95 °C for 5 min; 35 cycles of 95 °C for 1 min, 50 °C for 1 min, and 72 °C for 1 min; and a final extension of 10 min at 72 °C. Amplicons were purified and Sanger sequenced on an ABI sequencing instrument by Macrogen Inc. (Seoul, South Korea) or by the Australian Genome Research Facility (AGRF) using the 1492r primer. Trimmed high quality read data from each isolate was used for presumptive identification by querying the 16S rRNA gene sequences via BLASTn. For some isolates the near-complete 16S rRNA gene sequence was determined by sequencing with additional primers (27f, 357f (5'-CCT ACG GGA GGC AGC AG-3', (94)), 926f (5'-CCG TCA ATT CMT TTR AGT TT-3', (95)), 519r (5'-GWA TTA CCG CGG CKG CTG-3', (94)), 926r (5'-AAA CTR AAA MGA ATT GAC GG-3', (95)), and 1492r). The six reads for each isolate were aligned using Geneious Prime 2019.1.2 (<https://www.geneious.com>) via the Geneious global alignment default settings with automatic determination of read direction. From this alignment, a consensus sequence for the 16S rRNA gene was constructed based on the frequency of a base and its quality (from chromatogram

data) in each alignment column. The consensus sequence length for each of the six isolate pairs varied from 1352 to 1495 nucleotides. GenBank accession numbers for sequences are shown in Table 1.

Qualitative free radical scavenging assay. DPPH is a stable free radical that is purple in its oxidized state but becomes white-yellow when reduced by antioxidants, and has been used to identify FRS marine bacteria (96, 97). To qualitatively assess *E. diaphana*-associated bacteria isolates for FRS ability, a sterile Whatman #1 filter paper was gently pressed against fresh (2-4 days old) colonies from a streak plate. Plates (with filter paper) were then incubated overnight at 26 °C. The following day, filter papers were removed with forceps, allowed to dry in a fume hood for 30 min, and 500 µL of a 0.2 mM DPPH (Cat# D9132, Sigma-Aldrich) solution in methanol was applied with a pipette over individual colonies. As a positive control, a few drops of 0.1% (w/v) L-ascorbic acid (Cat# A7631, Sigma-Aldrich) were placed on a separate filter. The response of each isolate to DPPH was recorded within 3 min of DPPH application; a positive response was recorded if a white-yellow halo appeared around individual colonies within 1 min, a weak positive response was assigned to strains that had a halo form between 1 and 3 min after DPPH application, and a negative response was listed for strains that failed to form a halo (Fig. 3). Approximately 700 isolates were screened using the qualitative DPPH assay.

FIG 3: DPPH is a stable free radical that is purple in its oxidized state. When reduced by an antioxidant, a white-yellow halo will appear around individual bacteria colonies (A), this was qualitatively deemed a positive response. Isolates that did not have a halo around colonies within 3 min of DPPH application were deemed negative (B).

Quantitative free radical scavenging assay. To quantitatively assess the FRS ability, select isolates were grown in R2A broth (Table S5). A volume of 50 mL of autoclaved medium was added to sterile 250 mL Erlenmeyer flasks and each flask was inoculated with an isolate colony grown from a R2A or MA plate culture. Cultures were grown with shaking (150 rpm; Ratek orbital incubator) at 37°C for 48 h. Sterile uninoculated R2A was used as a control. A minimum of three replicate cultures were grown per isolate. After 48 h, the optical density of each culture was measured at 600 nm (OD₆₀₀, CLARIOstar PLUS, BMG Labtech), and the cultures (including negative medium controls) were centrifuged at 3000 x *g* at 4°C for 30 min (Allegra X-12R) to pellet the bacterial cells. The cell-free supernatants (CFSs) were collected, frozen at -80 °C, freeze dried (Alpha 1-4 LDplus, Martin Christ), and stored under inert gas in a dark, dry environment until analysis. Antioxidants were extracted from the CFSs by resuspending at 50 mg mL⁻¹ in 100% methanol, sonicating (Branson 2510) for 5 min, then centrifuging at 3000 x *g* at 4 °C for 5 min. Quantitative DPPH assays were run by creating a 1:1 solution of 0.2 mM DPPH in methanol and CFS extract to a final volume of 1 mL, vortexing, and incubation in the dark at room temperature for 30 min. Samples were then vortexed briefly, and three 300 µL replicates of each sample were transferred into a well of a 96 well plate. Decolourization of DPPH was determined by measuring the decrease in absorbance at 517 nm (Enspire 2300 plate reader, Perkin Elmer), and the FRS activity was calculated according to the formula, % DPPH scavenging activity = (Control – Sample) / Control × 100, where, Control is the absorbance of the DPPH control (1:1 0.2 mM DPPH:methanol), and Sample is the absorbance of CFS extract in DPPH. All samples were measured against a 100% methanol blank. Positive controls consisting of 0.01 - 0.001% (w/v) L-ascorbic acid were run on each 96-well plate. FRS activity ranged from 0-90%.

Catalase assay. The pelleted cells from above were resuspended in 2 mL fRSS and 500 µL H₂O₂ giving a final concentration of 16 mM. If bubbles appeared, the organism was considered catalase positive. If there were no bubbles, the organism was classified as catalase negative.

Inhibition testing. Each paired set of high and low FRS strains were inoculated crosswise along the middle of MA plates to test for antagonism. Plates were kept at 26 °C and monitored daily

for up to 7 days for antagonistic activity by documenting the presence or absence of both inoculated isolates and if there was a zone of inhibition between them.

Phylogenetic analysis. All partial 16S rRNA gene sequences (842) were aligned with reference sequences (72) of closely related organisms using Geneious Prime 2019.1.2 (<https://www.geneious.com>). This alignment was used to construct a neighbor-joining phylogenetic tree using the Jukes-Cantor method. Maximum-likelihood dendrograms were generated with bootstrap values of 1000.

Whole genome sequence analysis. Positive FRS strains along with conspecific or congeneric negative FRS strains were selected for genome sequencing; in total, six pairs of isolates were sequenced. Genomic DNA was isolated from a single colony using a JANUS Chemagic Workstation and Chemagic Viral DNA/RNA kit (PerkinElmer). Libraries were prepared with the Nextera XT DNA sample preparation kit (Illumina). Readsets were produced using the Illumina sequencing platform (Instrument: Illumina NextSeq 500, 150 base, paired-end) and the whole genome shotgun (WGS) method. Read depth coverage was approximately 100 times assuming a genome size of 4 M bases.

Illumina readsets for each isolate were assembled using Skesa (98) and the draft genome sequence annotated using Prokka (99). A genome sequence based taxonomic classification for each isolate was determined using Kraken2 (100) with the Genome Taxonomy Database (GTDB; 101) as the curated genomic data source. Classification was primarily based on the genome sequence of related isolates (within the relevant species where possible), which were obtained from GenBank. In situations where genomes of taxonomically relevant individuals were available, a species level classification was possible. Where available, closed genome sequences from GenBank were used for comparative genomics analysis. For each of the six pairs of isolates, core genome (i.e. genes shared between the isolate pair) comparisons were performed, as implemented in Nullarbor (<https://github.com/tseemann/nullarbor>), with phylogenies inferred using SNP differences. Genes of interest for DMSP synthesis and degradation, vitamin B₁₂ synthesis, and catalase were identified from the annotated genome

sequence (GFF format) produced by Prokka; specific genes were identified by both name and Refseq accession number.

16S rRNA gene copy number estimation.

The 16S rRNA gene copy number of the 12 draft genomes was predicted by the 16Stimator pipeline (102). Briefly, all the 12 genomic assemblies were submitted to the rapid annotations using subsystems technology (RAST) server (103), and the positions of 16S rRNA and a set of single-copied housekeeping genes (Table S6) were extracted from the RAST annotations. The clean read sets were mapped back to the corresponding genomic assemblies by Bowtie 2 (104) to determine the read depth of each position. Finally, the 16S copy number of each isolate was calculated by dividing the median depth of 16S gene by the median depth of the single-copied housekeeping genes after the read depths were calibrated by the model parameters provided by 16Stimator.

Statistical analysis. CFU counts were analyzed in R (v3.6.2, 105) by first checking the assumptions of equal variance and homogeneity. An analysis of variance test was used to detect differences in the mean number of bacterial colonies from each anemone genotype by solid growth media (R2A or MA). A one-way analysis of variance (one-way ANOVA; 106) was used to determine if there were significant differences between FRS abilities of selected positive (high FRS), negative (low FRS), and media controls, and pairwise comparisons were performed using Tukey's HSD (107, 108). Each probiotic pair and media control was tested to determine if data met the assumptions of normality and homoscedasticity. If either assumption was violated, the non-parametric Kruskal-Wallis rank sum test (109) was used with a Dunn test (110) for multiple comparisons (p-values adjusted with the Benjamini-Hochberg method (111)) with the R package "FSA" (112).

Data availability. WGS raw reads are freely available in the Sequence Read Archive under BioProject PRJNA574193; the complete data set is listed in Table S1.

431 **ACKNOWLEDGEMENTS**

432 This research was supported by the Australian Research Council Discovery Project grant
 433 DP160101468 (to MJHvO and LLB). MJHvO acknowledges Australian Research Council Laureate
 434 Fellowship FL180100036. We are grateful to Leon Hartman, Giulia Holland, and Shona Elliot-
 435 Kerr for their contributions in the preliminary culturing and screening of anemone-associated
 436 bacteria. Dr. Gayle Philip contributed with bacterial whole genome sequence analysis and Leon
 437 Hartman assisted with figure designs and reviewed the manuscript. Xavier Smith assisted with
 438 bacterial inhibition tests. Whole genome sequencing was organized by Dr. Glen Carter at the
 439 Peter Doherty Institute, Melbourne, Australia.

440 AMD, MvO and LLB conceived and designed the study. AMD performed the sampling and
 441 sample processing. AMD, DB, and HL completed bioinformatic analyses. AMD wrote the first
 442 draft. All authors edited and approved the final manuscript.

- 443 1. Cesar H, Burke L, Pet-Soede L. 2003. The economics of worldwide coral reef degradation. (CEEC)
- 444 CEEC,
- 445 2. Alder J, Artherton R, Ash N, Diop S, Heileman S, Huber M, Arias-Izasa F, Koranteng K, Lacambra
- 446 C, McLeod K, Payet E, Perera N, Tang LD, Spalding M, Zahedi K. 2006. Marine and coastal
- 447 ecosystems and human well-being: A synthesis report based on the findings of the Millennium
- 448 Ecosystem Assessment. Programme UNE,
- 449 3. Fisher R, O'Leary RA, Low-Choy S, Mengersen K, Knowlton N, Brainard RE, Caley MJ. 2015.
- 450 Species richness on coral reefs and the pursuit of convergent global estimates. *Current Biology*
- 451 25:500-505.
- 452 4. Spalding M, M. Grenfell A. 1997. New estimates of global and regional coral reef areas, vol 16.
- 453 5. de Groot R, Brander L, van der Ploeg S, Costanza R, Bernard F, Braat L, Christie M, Crossman N,
- 454 Ghermandi A, Hein L, Hussain S, Kumar P, McVittie A, Portela R, Rodriguez LC, ten Brink P, van
- 455 Beukering P. 2012. Global estimates of the value of ecosystems and their services in monetary
- 456 units. *Ecosystem Services* 1:50-61.
- 457 6. Stuart-Smith RD, Brown CJ, Ceccarelli DM, Edgar GJ. 2018. Ecosystem restructuring along the
- 458 Great Barrier Reef following mass coral bleaching. *Nature* 560:92-96.
- 459 7. Hughes TP, Kerry JT, Alvarez-Noriega M, Alvarez-Romero JG, Anderson KD, Baird AH, Babcock
- 460 RC, Beger M, Bellwood DR, Berkemans R, Bridge TC, Butler IR, Byrne M, Cantin NE, Comeau S,
- 461 Connolly SR, Cumming GS, Dalton SJ, Diaz-Pulido G, Eakin CM, Figueira WF, Gilmour JP, Harrison
- 462 HB, Heron SF, Hoey AS, Hobbs JA, Hoogenboom MO, Kennedy EV, Kuo CY, Lough JM, Lowe RJ,
- 463 Liu G, McCulloch MT, Malcolm HA, McWilliam MJ, Pandolfi JM, Pears RJ, Pratchett MS, Schoepf
- 464 V, Simpson T, Skirving WJ, Sommer B, Torda G, Wachenfeld DR, Willis BL, Wilson SK. 2017.
- 465 Global warming and recurrent mass bleaching of corals. *Nature* 543:373-377.
- 466 8. Hoegh-Guldberg O, Jacob D, Taylor M, Bolaños TG, Bindi M, Brown S, Camilloni I, Diedhiou A,
- 467 Djalante R, Ebi K. 2019. The human imperative of stabilizing global climate change at 1.5° C.
- 468 *Science* 365:eaaw6974.
- 469 9. Hughes TP, Anderson KD, Connolly SR, Heron SF, Kerry JT, Lough JM, Baird AH, Baum SF,
- 470 Berumen ML, Bridge T, Claar DC, Eakin CM, Gilmour JP, Graham NAI, Harrison HB, Hobbs JA,
- 471 Hoey AS, Hoogenboom M, Lowe RJ, McCulloch MT, Pandolfi JM, Pratchett MS, Schoepf V, Torda
- 472 G, Wilson SK. 2018. Spatial and temporal patterns of mass bleaching of corals in the
- 473 Anthropocene. *Science* 359:80-83.
- 474 10. Rohwer F, Seguritan V, Azam F, Knowlton N. 2002. Diversity and distribution of coral-associated
- 475 bacteria. *Mar Ecol Prog Ser* 243:10.
- 476 11. Devlin-Durante MK, Miller MW, Precht WF, Baums IB. 2016. How old are you? Genet age
- 477 estimates in a clonal animal. *Molecular Ecology* 25:5628-5646.
- 478 12. Glynn PW. 1996. Coral reef bleaching: facts, hypotheses and implications. *Global Change Biology*
- 479 2:495-509.
- 480 13. Muscatine L, Porter JW. 1977. Reef corals: mutualistic symbioses adapted to nutrient-poor
- 481 environments. *Bioscience* 27:454-460.
- 482 14. Tremblay P, Grover R, Maguer J-F, Hoogenboom M, Ferrier-Pagès C. 2014. Carbon translocation
- 483 from symbiont to host depends on irradiance and food availability in the tropical coral
- 484 *Stylophora pistillata*. *Coral Reefs* 33:1-13.
- 485 15. Venn AA, Loram JE, Douglas AE. 2008. Photosynthetic symbioses in animals. *J Exp Bot* 59:1069-
- 486 80.
- 487 16. Yellowlees D, Rees TAV, Leggat W. 2008. Metabolic interactions between algal symbionts and
- 488 invertebrate hosts. *Plant, Cell & Environment* 31:679-694.
- 489 17. Weis VM. 2008. Cellular mechanisms of Cnidarian bleaching: stress causes the collapse of
- 490 symbiosis. *Journal of Experimental Biology* 211:3059-3066.

- 491 18. Cuning R, Baker AC. 2012. Excess algal symbionts increase the susceptibility of reef corals to
492 bleaching. *Nature Climate Change* 3:259-262.
- 493 19. Wiedenmann J, D'Angelo C, Smith EG, Hunt AN, Legiret F-E, Postle AD, Achterberg EP. 2012.
494 Nutrient enrichment can increase the susceptibility of reef corals to bleaching. *Nature Climate*
495 *Change* 3:160.
- 496 20. Wooldridge SA. 2013. Breakdown of the coral-algae symbiosis: towards formalising a linkage
497 between warm-water bleaching thresholds and the growth rate of the intracellular
498 zooxanthellae. *Biogeosciences* 10:1647-1658.
- 499 21. Dykens JA, Shick JM, Benoit C, Buettner GR, WINSTON GW. 1992. Oxygen radical production in
500 the sea anemone *Anthopleura elegantissima* and its endosymbiotic algae. *Journal of*
501 *Experimental Biology* 168:219-241.
- 502 22. Nii CM, Muscatine L. 1997. Oxidative Stress in the Symbiotic Sea Anemone *Aiptasia pulchella*
503 (Carlgren, 1943): Contribution of the Animal to Superoxide Ion Production at Elevated
504 Temperature. *Biol Bull* 192:444-456.
- 505 23. Lesser MP. 2004. Experimental biology of coral reef ecosystems. *Journal of Experimental Marine*
506 *Biology and Ecology* 300:217-252.
- 507 24. Warner ME, Fitt WK, Schmidt GW. 1999. Damage to photosystem II in symbiotic dinoflagellates:
508 a determinant of coral bleaching. *Proceedings of the National Academy of Sciences* 96:8007-
509 8012.
- 510 25. Damjanovic K, Blackall LL, Webster NS, van Oppen MJH. 2017. The contribution of microbial
511 biotechnology to mitigating coral reef degradation. *Microb Biotechnol* 10:1236-1243.
- 512 26. van Oppen MJH, Gates RD, Blackall LL, Cantin N, Chakravarti LJ, Chan WY, Cormick C, Crean A,
513 Damjanovic K, Epstein H, Harrison PL, Jones TA, Miller M, Pears RJ, Peplow LM, Raftos DA,
514 Schaffelke B, Stewart K, Torda G, Wachenfeld D, Weeks AR, Putnam HM. 2017. Shifting
515 paradigms in restoration of the world's coral reefs. *Glob Chang Biol* 23:3437-3448.
- 516 27. Damjanovic K, van Oppen MJH, Menéndez P, Blackall LL. 2019. Experimental Inoculation of Coral
517 Recruits With Marine Bacteria Indicates Scope for Microbiome Manipulation in *Acropora tenuis*
518 and *Platygyra daedalea*. *Frontiers in Microbiology* 10.
- 519 28. van Oppen MJ, Oliver JK, Putnam HM, Gates RD. 2015. Building coral reef resilience through
520 assisted evolution. *Proc Natl Acad Sci U S A* 112:2307-13.
- 521 29. Peixoto RS, Rosado PM, Leite DC, Rosado AS, Bourne DG. 2017. Beneficial Microorganisms for
522 Corals (BMC): Proposed Mechanisms for Coral Health and Resilience. *Front Microbiol* 8:341.
- 523 30. Epstein HE, Smith HA, Torda G, van Oppen MJ. 2019. Microbiome engineering: enhancing
524 climate resilience in corals. *Frontiers in Ecology and the Environment* 17:100-108.
- 525 31. Vega Thurber R, Willner-Hall D, Rodriguez-Mueller B, Desnues C, Edwards RA, Angly F, Dinsdale
526 E, Kelly L, Rohwer F. 2009. Metagenomic analysis of stressed coral holobionts. *Environ Microbiol*
527 11:2148-63.
- 528 32. Epstein HE, Torda G, van Oppen MJH. 2019. Relative stability of the *Pocillopora acuta*
529 microbiome throughout a thermal stress event. *Coral Reefs* 38:373-386.
- 530 33. Sunagawa S, Woodley CM, Medina M. 2010. Threatened corals provide underexplored microbial
531 habitats. *PLoS One* 5:e9554.
- 532 34. Mouchka ME, Hewson I, Harvell CD. 2010. Coral-associated bacterial assemblages: current
533 knowledge and the potential for climate-driven impacts. *Integr Comp Biol* 50:662-74.
- 534 35. Littman R, Willis BL, Bourne DG. 2011. Metagenomic analysis of the coral holobiont during a
535 natural bleaching event on the Great Barrier Reef. *Environ Microbiol Rep* 3:651-60.
- 536 36. Pootakham W, Mhuanong W, Yoocha T, Putchim L, Jomchai N, Sonthirod C, Naktang C,
537 Kongkachana W, Tangphatsornruang S. 2019. Heat-induced shift in coral microbiome reveals

- several members of the Rhodobacteraceae family as indicator species for thermal stress in *Porites lutea*. *MicrobiologyOpen* 8:e935.
37. Hartman LM, van Oppen MJ, Blackall LL. 2020. The Effect of Thermal Stress on the Bacterial Microbiome of *Exaiptasia diaphana*. *Microorganisms* 8:20.
38. Ahmed HI, Herrera Sarrias M, Liew YJ, Aranda M. 2019. Long-term temperature stress in the coral model *Aiptasia* supports the “Anna Karenina principle” for bacterial microbiomes. *Frontiers in microbiology* 10:975.
39. Plovie A. 2010. Comparison of bacterial communities associated with healthy and bleached *Aiptasia pallida*, a novel model organism for coral studies: Implications and variation during bleaching. *Biochemistry and Biotechnology* Liège University.
40. Gilbert JA, Hill R, Doblin MA, Ralph PJ. 2012. Microbial consortia increase thermal tolerance of corals. *Marine Biology* 159:1763-1771.
41. Ziegler M, Seneca FO, Yum LK, Palumbi SR, Voolstra CR. 2017. Bacterial community dynamics are linked to patterns of coral heat tolerance. *Nat Commun* 8:14213.
42. Alagely A, Krediet CJ, Ritchie KB, Teplitski M. 2011. Signaling-mediated cross-talk modulates swarming and biofilm formation in a coral pathogen *Serratia marcescens*. *ISME J* 5:1609-20.
43. dos Santos FAH, Duarte GA, Rachid CT, Chaloub RM, Calderon EN, Marangoni LF, Bianchini A, Nudi AH, do Carmo FL, van Elsas JD, Rosado AS, Castro CB, Peixoto RS. 2015. Impact of oil spills on coral reefs can be reduced by bioremediation using probiotic microbiota. *Sci Rep* 5:18268.
44. Rosado PM, Leite DCA, Duarte GAS, Chaloub RM, Jospin G, Nunes da Rocha U, P. Saraiva J, Dini-Andreote F, Eisen JA, Bourne DG, Peixoto RS. 2018. Marine probiotics: increasing coral resistance to bleaching through microbiome manipulation. *The ISME Journal* doi:10.1038/s41396-018-0323-6.
45. Herrera M, Ziegler M, Voolstra CR, Aranda M. 2017. Laboratory-Cultured Strains of the Sea Anemone *Exaiptasia* Reveal Distinct Bacterial Communities. *Frontiers in Marine Science* 4.
46. Binsarhan M. 2016. Comparative analysis and culturing of the microbial community of *Aiptasia pallida*, A Sea Anemone Model for Coral Biology.
47. Röthig T, Costa RM, Simona F, Baumgarten S, Torres AF, Radhakrishnan A, Aranda M, Voolstra CR. 2016. Distinct Bacterial Communities Associated with the Coral Model *Aiptasia* in Aposymbiotic and Symbiotic States with *Symbiodinium*. *Frontiers in Marine Science* 3.
48. Ceh J, van Keulen M, Bourne DG. 2013. Intergenerational transfer of specific bacteria in corals and possible implications for offspring fitness. *Microb Ecol* 65:227-31.
49. Rothig T, Yum LK, Kremb SG, Roik A, Voolstra CR. 2017. Microbial community composition of deep-sea corals from the Red Sea provides insight into functional adaption to a unique environment. *Sci Rep* 7:44714.
50. Chiu JM, Li S, Li A, Po B, Zhang R, Shin PK, Qiu JW. 2012. Bacteria associated with skeletal tissue growth anomalies in the coral *Platygyra carnosus*. *FEMS Microbiol Ecol* 79:380-91.
51. Hai NV. 2015. The use of probiotics in aquaculture. *Journal of Applied Microbiology* 119:917-935.
52. Kesarcodi-Watson A, Kaspar H, Lategan MJ, Gibson L. 2010. *Alteromonas macleodii* 0444 and *Neptunomonas* sp. 0536, two novel probiotics for hatchery-reared Greenshell™ mussel larvae, *Perna canaliculus*. *Aquaculture* 309:49-55.
53. Riquelme C, Araya R, Vergara N, Rojas A, Guaita M, Candia M. 1997. Potential probiotic strains in the culture of the Chilean scallop *Argopecten purpuratus* (Lamarck, 1819). *Aquaculture* 154:17-26.
54. Haryanti H, Sugama K, Tsumura S, Nishijima R. 2017. ENHANCE PRODUCTION OF BLACK TIGER SHRIMP *Penaeus monodon* POSTLARVAE BY PROBIOTIC BACTERIUM *Alteromonas* sp. *Indonesian Fisheries Research Journal* 7:1-6.

55. Biebl H, Pukall R, Lünsdorf H, Schulz S, Allgaier M, Tindall BJ, Wagner-Döbler I. 2007. Description of *Labrenzia alexandrii* gen. nov., sp. nov., a novel alphaproteobacterium containing bacteriochlorophyll a, and a proposal for reclassification of *Stappia aggregata* as *Labrenzia aggregata* comb. nov., of *Stappia marina* as *Labrenzia marina* comb. nov. and of *Stappia alba* as *Labrenzia alba* comb. nov., and emended descriptions of the genera *Pannonibacter*, *Stappia* and *Roseibium*, and of the species *Roseibium denhamense* and *Roseibium hamelinense*.
56. Mancuso M, Rappazzo A, Genovese M, El Hady M, Ghonimy A, Ismail M, Reda R, Cappello S, Genovese L, Maricchiolo G. 2015. In vitro selection of bacteria and isolation of Probiotics from farmed *Sparus aurata* with potential for use as probiotics. *International Journal of Animal Biology* 1:93-98.
57. Brown T, Otero C, Grajales A, Rodriguez E, Rodriguez-Lanetty M. 2017. Worldwide exploration of the microbiome harbored by the cnidarian model, *Exaiptasia pallida* (Agassiz in Verrill, 1864) indicates a lack of bacterial association specificity at a lower taxonomic rank. *PeerJ* 5:e3235.
58. Sharp KH, Distel D, Paul VJ. 2012. Diversity and dynamics of bacterial communities in early life stages of the Caribbean coral *Porites astreoides*. *ISME J* 6:790-801.
59. Littman RA, Willis BL, Pfeiffer C, Bourne DG. 2009. Diversities of coral-associated bacteria differ with location, but not species, for three acroporid corals on the Great Barrier Reef. *FEMS Microbiol Ecol* 68:152-63.
60. Kellogg CA, Piceno YM, Tom LM, DeSantis TZ, Gray MA, Zawada DG, Andersen GL. 2013. Comparing bacterial community composition between healthy and white plague-like disease states in *Orbicella annularis* using PhyloChip G3 microarrays. *PLoS One* 8:e79801.
61. El-Rhman AMA, Khat tab YA, Shalaby AM. 2009. *Micrococcus luteus* and *Pseudomonas* species as probiotics for promoting the growth performance and health of Nile tilapia, *Oreochromis niloticus*. *Fish & Shellfish Immunology* 27:175-180.
62. Osman H, Ibrahim TB, Soliman W, Aboud O. 2010. Improvement growth and immune status using a potential probiotic bacteria *Micrococcus* species among cultured *Oreochromis niloticus*. *New York Science Journal* 3:5-11.
63. Lau SCK, Tsoi MMY, Li X, Plakhotnikova I, Dobretsov S, Lau KWK, Wu M, Wong P-K, Pawlik JR, Qian P-Y. 2005. *Winogradskyella poriferorum* sp. nov., a novel member of the family Flavobacteriaceae isolated from a sponge in the Bahamas. *International Journal of Systematic and Evolutionary Microbiology* 55:1589-1592.
64. Franco A, Busse H-J, Schubert P, Wilke T, Kämpfer P, Glaeser S. 2018. *Winogradskyella pocilloporae* sp. Nov. Isolated from healthy tissue of the coral *Pocillopora damicornis*, vol 68.
65. Gonzaga A, López-Pérez M, Martín-Cuadrado A-B, Ghai R, Rodríguez-Valera F. 2012. Complete Genome Sequence of the Copiotrophic Marine Bacterium *Alteromonas macleodii* Strain ATCC 27126^T. *Journal of Bacteriology* 194:6998-6998.
66. Blackall LL, Wilson B, van Oppen MJ. 2015. Coral-the world's most diverse symbiotic ecosystem. *Mol Ecol* 24:5330-47.
67. Sunda W, Kieber DJ, Kiene RP, Huntsman S. 2002. An antioxidant function for DMSP and DMS in marine algae. *Nature* 418:317-20.
68. Curson AR, Liu J, Bermejo Martinez A, Green RT, Chan Y, Carrion O, Williams BT, Zhang SH, Yang GP, Bulman Page PC, Zhang XH, Todd JD. 2017. Dimethylsulfoniopropionate biosynthesis in marine bacteria and identification of the key gene in this process. *Nat Microbiol* 2:17009.
69. Raina J-B, Tapiolas D, Willis BL, Bourne DG. 2009. Coral-Associated Bacteria and Their Role in the Biogeochemical Cycling of Sulfur. *Applied and Environmental Microbiology* 75:3492-3501.

632 70. Hatton AD, Shenoy DM, Hart MC, Mogg A, Green DH. 2012. Metabolism of DMSP, DMS and
633 DMSO by the cultivable bacterial community associated with the DMSP-producing dinoflagellate
634 *Scrippsiella trochoidea*. Biogeochemistry 110:131-146.

635 71. Fiedor J, Burda K. 2014. Potential role of carotenoids as antioxidants in human health and
636 disease. Nutrients 6:466-488.

637 72. Fiedor J, Fiedor L, Haeßner R, Scheer H. 2005. Cyclic endoperoxides of β -carotene, potential pro-
638 oxidants, as products of chemical quenching of singlet oxygen. Biochimica et Biophysica Acta
639 (BBA) - Bioenergetics 1709:1-4.

640 73. Flórez LV, Biedermann PHW, Engl T, Kaltenpoth M. 2015. Defensive symbioses of animals with
641 prokaryotic and eukaryotic microorganisms. Natural Product Reports 32:904-936.

642 74. Asker D, Beppu T, Ueda K. 2007. Mesoflavibacter zeaxanthinifaciens gen. nov., sp. nov., a novel
643 zeaxanthin-producing marine bacterium of the family Flavobacteriaceae. Systematic and applied
644 microbiology 30:291-296.

645 75. Shindo K, Kikuta K, Suzuki A, Katsuta A, Kasai H, Yasumoto-Hirose M, Matsuo Y, Misawa N,
646 Takaichi S. 2007. Rare carotenoids, (3R)-saproxanthin and (3R,2'S)-myxol, isolated from novel
647 marine bacteria (Flavobacteriaceae) and their antioxidative activities. Applied Microbiology and
648 Biotechnology 74:1350.

649 76. Gammone M, Riccioni G, #039, Orazio N. 2015. Marine Carotenoids against Oxidative Stress:
650 Effects on Human Health. Marine Drugs 13:6226.

651 77. Mohana D, Thippeswamy S, Abhishe R. 2013. Antioxidant, antibacterial, and ultraviolet-
652 protective properties of carotenoids isolated from Micrococcus spp. Radiation Protection and
653 Environment 36:168-174.

654 78. Motone K, Takagi T, Aburaya S, Miura N, Aoki W, Ueda M. 2020. A Zeaxanthin-Producing
655 Bacterium Isolated from the Algal Phycosphere Protects Coral Endosymbionts from
656 Environmental Stress. mBio 11:e01019-19.

657 79. Croft MT, Lawrence AD, Raux-Deery E, Warren MJ, Smith AG. 2005. Algae acquire vitamin B 12
658 through a symbiotic relationship with bacteria. Nature 438:90.

659 80. Raux E, Schubert H, Warren M. 2000. Biosynthesis of cobalamin (vitamin B12): a bacterial
660 conundrum. Cellular and Molecular Life Sciences CMLS 57:1880-1893.

661 81. Matthews JL, Raina JB, Kahlke T, Seymour JR, van Oppen MJ, Suggett DJ. Symbiodiniaceae-
662 bacteria interactions: rethinking metabolite exchange in reef-building corals as multi-partner
663 metabolic networks. Environmental Microbiology.

664 82. Agostini S, Suzuki Y, Casareto BE, Nakano Y, Hidaka M, Badrun N. 2009. Coral symbiotic complex:
665 Hypothesis through vitamin B12 for a new evaluation. Galaxea, Journal of Coral Reef Studies
666 11:1-11.

667 83. Smith SM. 2018. Complementarity in the Coral Holobiont: A Genomic Analysis of Bacterial
668 Isolates of *Orbicella faveolata* and *Symbiodinium* spp.

669 84. Imlay JA. 2018. Where in the world do bacteria experience oxidative stress? Environ Microbiol
670 doi:10.1111/1462-2920.14445.

671 85. Taylor WI, Achanzar D. 1972. Catalase Test as an Aid to the Identification of
672 *Enterobacteriaceae*. Applied Microbiology 24:58-61.

673 86. Cabiscol E, Tamarit J, Ros J. 2000. Oxidative stress in bacteria and protein damage by reactive
674 oxygen species. Int Microbiol 3:3-8.

675 87. von Ossowski I, Mulvey MR, Leco PA, Borys A, Loewen PC. 1991. Nucleotide sequence of
676 *Escherichia coli* katE, which encodes catalase HPII. Journal of bacteriology 173:514-520.

- 677 88. Damjanovic K, Menéndez P, Blackall LL, van Oppen MJH. 2020. Mixed-mode bacterial
678 transmission in the common brooding coral *Pocillopora acuta*. *Environmental Microbiology*
679 22:397-412.
- 680 89. Sharp KH, Ritchie KB, Schupp PJ, Ritson-Williams R, Paul VJ. 2010. Bacterial acquisition in
681 juveniles of several broadcast spawning coral species. *PLoS One* 5:e10898.
- 682 90. Sharp KH, Sneed JM, Ritchie KB, McDaniel L, Paul VJ. 2015. Induction of Larval Settlement in the
683 Reef Coral *Porites astreoides* by a Cultivated Marine *Roseobacter* Strain. *Biol Bull* 228:98-107.
- 684 91. Prabhu S, Rekha P, Arun A. 2014. Zeaxanthin biosynthesis by members of the genus *Muricauda*.
685 *Pol J Microbiol* 63:115-119.
- 686 92. Dungan AM, Hartman LM, Tortorelli G, Belderok R, Lamb AM, Pisan L, McFadden GI, Blackall LL,
687 van Oppen MJ. 2020. *Exaiptasia diaphana* from the great barrier reef: a valuable resource for
688 coral symbiosis research. *Symbiosis*:1-12.
- 689 93. Lane D. 1991. 16S/23S rRNA sequencing. *Nucleic acid techniques in bacterial systematics*:115-
690 175.
- 691 94. Muyzer G, De Waal EC, Uitterlinden AG. 1993. Profiling of complex microbial populations by
692 denaturing gradient gel electrophoresis analysis of polymerase chain reaction-amplified genes
693 coding for 16S rRNA. *Appl Environ Microbiol* 59:695-700.
- 694 95. Lane DJ, Pace B, Olsen GJ, Stahl DA, Sogin ML, Pace NR. 1985. Rapid determination of 16S
695 ribosomal RNA sequences for phylogenetic analyses. *Proceedings of the National Academy of*
696 *Sciences* 82:6955-6959.
- 697 96. Takao T, Kitatani F, Watanabe N, Yagi A, Sakata K. 1994. A simple screening method for
698 antioxidants and isolation of several antioxidants produced by marine bacteria from fish and
699 shellfish. *Bioscience, Biotechnology, and Biochemistry* 58:1780-1783.
- 700 97. Velho-Pereira S, Parvatkar P, Furtado IJ. 2015. Evaluation of antioxidant producing potential of
701 halophilic bacterial bionts from marine invertebrates. *Indian Journal of Pharmaceutical Sciences*
702 77:183-189.
- 703 98. Souvorov A, Agarwala R, Lipman DJ. 2018. SKESA: strategic k-mer extension for scrupulous
704 assemblies. *Genome Biol* 19:153.
- 705 99. Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068-9.
- 706 100. Wood DE, Salzberg SL. 2014. Kraken: ultrafast metagenomic sequence classification using exact
707 alignments. *Genome Biol* 15:R46.
- 708 101. Parks DH, Chuvochina M, Chaumeil P-A, Rinke C, Mussig AJ, Hugenholtz P. 2020. A complete
709 domain-to-species taxonomy for Bacteria and Archaea. *Nature Biotechnology*
710 doi:10.1038/s41587-020-0501-8.
- 711 102. Perisin M, Vetter M, Gilbert JA, Bergelson J. 2016. 16Stimator: statistical estimation of ribosomal
712 gene copy numbers from draft genome assemblies. *The ISME journal* 10:1020-1024.
- 713 103. Brettin T, Davis JJ, Disz T, Edwards RA, Gerdes S, Olsen GJ, Olson R, Overbeek R, Parrello B, Pusch
714 GD. 2015. RASTtk: a modular and extensible implementation of the RAST algorithm for building
715 custom annotation pipelines and annotating batches of genomes. *Scientific reports* 5:8365.
- 716 104. Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nature methods*
717 9:357.
- 718 105. R Core Team. 2019. A language and environment for statistical computing. Vienna, Austria: R
719 Foundation for Statistical Computing; 2012. URL <https://www.R-project.org>.
- 720 106. Chambers JM, Hastie TJ. 1992. Statistical models in S, vol 251. Wadsworth & Brooks/Cole
721 Advanced Books & Software Pacific Grove, CA.
- 722 107. Miller RG. 1981. Normal univariate techniques, p 37-108, *Simultaneous statistical inference*.
723 Springer.
- 724 108. Yandell B. 2017. *Practical data analysis for designed experiments*. Routledge.

- 725 109. Holland M, Wolfe D. 1973. Nonparametric statistical methods. John Wiley & Sons, New York.
726 110. Dunn OJ. 1964. Multiple comparisons using rank sums. *Technometrics* 6:241-252.
727 111. Ferreira J, Zwinderman A. 2006. On the Benjamini–Hochberg method. *The Annals of Statistics*
728 34:1827-1849.
729 112. Ogle DH. 2017. FSA: fisheries stock analysis. R package version 08 17:636.

730

731 **Table S1:** Isolate Genome Sequence Data Summary. Strains are presented as high FRS (grey) followed by low FRS (white). 16S rRNA
 732 gene presumptive identity is derived from the NCBI classification of near-complete 16S rRNA gene sequences. *We were unable to
 733 determine the 16S rRNA copy number of isolate MMSF00068 due to the presence of contaminant sequences in the WGS.

Isolate	Genus Level Classification	Genus % Confidence	Family	Family % Confidence	SRA Accession	Sample Accession
MMSF00257	<i>Alteromonas</i>	96.65	<i>Alteromonadaceae</i>	96.74	SRR10186803	SAMN12851724
MMSF00958	<i>Alteromonas</i>	92.93	<i>Alteromonadaceae</i>	92.98	SRR10186806	SAMN12851731
MMSF01163	<i>Alteromonas</i>	9.28	<i>Alteromonadaceae</i>	12.01	SRR10186808	SAMN12851721
MMSF00404	<i>Alteromonas</i>	8.15	<i>Alteromonadaceae</i>	10.49	SRR10186805	SAMN12851732
MMSF00132	<i>Labrenzia</i>	90.9	<i>Rhodobacteraceae</i>	90.91	SRR10186800	SAMN12851727
MMSF00249	<i>Labrenzia</i>	90.83	<i>Rhodobacteraceae</i>	90.83	SRR10186799	SAMN12851728
MMSF00964	<i>Marinobacter</i>	59.99	<i>Alteromonadaceae</i>	60.1	SRR10186802	SAMN12851725
MMSF01190	<i>Marinobacter</i>	65.77	<i>Alteromonadaceae</i>	65.85	SRR10186797	SAMN12851730
MMSF00068	<i>Micrococcus</i>	71.81	<i>Micrococcaceae</i>	72.88	SRR10186807	SAMN12851722
MMSF00107	<i>Micrococcus</i>	83.18	<i>Micrococcaceae</i>	84.87	SRR10186798	SAMN12851729
MMSF00046	<i>Winogradskyella</i>	49.63	<i>Flavobacteriaceae</i>	49.72	SRR10186801	SAMN12851726
MMSF00910	<i>Winogradskyella</i>	54.44	<i>Flavobacteriaceae</i>	54.53	SRR10186804	SAMN12851723

734

Isolate	16S rRNA gene presumptive identity	Confidence	Reads	Total Bases	G+C%	Avg. Read Length	Max Read Length	Avg Quality
MMSF00257	<i>Alteromonas macleodii</i>	1.00	2233080	334079592	44.7	149	151	30.9
MMSF00958	<i>Alteromonas macleodii</i>	1.00	2971718	443870758	44.6	149	151	31.9
MMSF01163	<i>Alteromonas oceani</i>	0.83	2757340	412741366	48.7	149	151	33.7
MMSF00404	<i>Alteromonas oceani</i>	0.83	2067800	306005114	48.7	147	151	33.3
MMSF00132	<i>Labrenzia aggregata</i>	0.98	3758944	557529936	59.2	148	151	31.1
MMSF00249	<i>Labrenzia aggregata</i>	0.98	3333098	497032699	59.3	149	151	30.7
MMSF00964	<i>Marinobacter salsuginis</i>	1.00	1713198	256635737	57.1	149	151	31
MMSF01190	<i>Marinobacter salsuginis</i>	1.00	3239568	483874048	57.1	149	151	33.1
MMSF00068	<i>Micrococcus luteus</i>	0.99	2658898	398496031	72.4	149	151	30.3

Supplemental Tables

MMSF00107	<i>Micrococcus yunnanensis</i>	0.99	2323654	348617190	72.8	150	151	31.9
MMSF00046	<i>Winogradskyella poriferorum</i>	0.59	1930584	285978916	35	148	151	31.8
MMSF00910	<i>Winogradskyella poriferorum</i>	0.59	3939880	579510587	35.6	147	151	33.9

735

Isolate	Est. Read Coverage	Contigs in Draft Genome	Bases in Draft Genome	Min Contig	Avg Contig	Max Contig	N50	16S rRNA gene copy number
MMSF00257	69	136	4831263	515	35523	210048	72973	5.15
MMSF00958	94	42	4732026	620	112667	580903	303003	4.79
MMSF01163	75	100	5507488	526	55074	394030	155957	4.18
MMSF00404	51	111	6014142	502	54181	256548	93725	3.97
MMSF00132	82	34	6792087	855	199767	1019681	302139	3.57
MMSF00249	73	44	6791310	851	154347	1286212	294458	3.02
MMSF00964	56	41	4588310	512	111910	932206	291012	3.67
MMSF01190	110	40	4404819	529	110120	769441	404351	3.21
MMSF00068	160	316	2484978	509	7863	78799	12670	*
MMSF00107	143	501	2435904	515	4862	42365	7717	1.75
MMSF00046	83	104	3446386	553	33138	299826	78549	1.03
MMSF00910	168	50	3456468	553	69129	355406	162623	0.77

736

737 **Table S2:** Pairwise comparison of the genome sequences between the pairs of isolates

NCBI Classification	Type strain	Core Genome	SNP difference
<i>Alteromonas macleodii</i>	ATCC 27126	85%	~60000
<i>Alteromonas oceani</i>	S35	60%	~190000
<i>Labrenzia aggregata</i>	IAM12614	55%	5
<i>Marinobacter salsuginis</i>	SD14B	15%	~120000
<i>Micrococcus</i> spp.	NCTC2665	70%	~35000
<i>Winogradskyella poriferorum</i>	NA	NA	10

738

739 **Table S3:** Search outcomes for genes of interest.

Isolate	NCBI Classification	Gene	Gene Classification	Contig	Start	End	Strand
MMSF00132	<i>Labrenzia aggregata</i>	<i>dddL</i>	positive	Contig_16_56.34	104067	104756	-
MMSF00249	<i>Labrenzia aggregata</i>	<i>dddL</i>	positive	Contig_18_45.1167	104045	104734	-
MMSF00132	<i>Labrenzia aggregata</i>	<i>dddP</i>	positive x2 (Copy 1)	Contig_6_57.2063	50787	52046	+
MMSF00132	<i>Labrenzia aggregata</i>	<i>dddP</i>	positive x2 (Copy 2)	Contig_6_57.2063	650583	651881	-
MMSF00249	<i>Labrenzia aggregata</i>	<i>dddP</i>	positive x2 (Copy 1)	Contig_17_46.7174	317316	318575	+
MMSF00249	<i>Labrenzia aggregata</i>	<i>dddP</i>	positive x2 (Copy 2)	Contig_17_46.7174	917112	918410	-
MMSF00132	<i>Labrenzia aggregata</i>	<i>dsyB</i>	positive	Contig_32_56.2994	210397	211419	+
MMSF00249	<i>Labrenzia aggregata</i>	<i>dsyB</i>	positive	Contig_4_46.0396	460215	461237	+
MMSF00046	<i>Winogradskyella poriferorum</i>	<i>katG</i>	positive	Contig_15_55.0851	16430	18661	+
MMSF00068	<i>Micrococcus luteus</i>	<i>katG</i>	ND; katA & katE detected	NA	NA	NA	NA
MMSF00107	<i>Micrococcus yunnanensis</i>	<i>katG</i>	ND; katA & katE detected	NA	NA	NA	NA
MMSF00132	<i>Labrenzia aggregata</i>	<i>katG</i>	positive	Contig_6_57.2063	425873	428044	-
MMSF00249	<i>Labrenzia aggregata</i>	<i>katG</i>	positive	Contig_17_46.7174	692402	694573	-
MMSF00257	<i>Alteromonas macleodii</i>	<i>katG</i>	positive	Contig_101_46.6864	22130	24502	+
MMSF00404	<i>Alteromonas oceani</i>	<i>katG</i>	positive x2 (Copy 1)	Contig_26_39.8992	70873	73035	-
MMSF00404	<i>Alteromonas oceani</i>	<i>katG</i>	positive x2 (Copy 2)	Contig_52_41.1626	55994	58225	-
MMSF00910	<i>Winogradskyella poriferorum</i>	<i>katG</i>	positive	Contig_39_101.483	25621	27852	+
MMSF00958	<i>Alteromonas macleodii</i>	<i>katG</i>	positive	Contig_21_66.072	22086	24458	+
MMSF00964	<i>Marinobacter salsuginis</i>	<i>katG</i>	positive	Contig_25_36.8412	50391	52565	-
MMSF01163	<i>Alteromonas oceani</i>	<i>katG</i>	positive x2 (Copy 1)	Contig_46_63.2119	54248	56479	-
MMSF01163	<i>Alteromonas oceani</i>	<i>katG</i>	positive x2 (Copy 2)	Contig_86_59.9301	369720	371882	-
MMSF01190	<i>Marinobacter salsuginis</i>	<i>katG</i>	positive	Contig_22_84.8909	7780	9954	+
MMSF00132	<i>Labrenzia aggregata</i>	<i>cobA</i>	positive	Contig_6_57.2063	552920	553741	-

Supplemental Tables

MMSF00249	<i>Labrenzia aggregata</i>	<i>cobA</i>	positive	Contig_17_46.7174	819449	820270	-
MMSF00132	<i>Labrenzia aggregata</i>	<i>cobB</i>	positive	Contig_6_57.2063	551611	552930	-
MMSF00249	<i>Labrenzia aggregata</i>	<i>cobB</i>	positive	Contig_17_46.7174	818140	819459	-
MMSF00132	<i>Labrenzia aggregata</i>	<i>cobC</i>	positive	Contig_6_57.2063	842748	843392	-
MMSF00249	<i>Labrenzia aggregata</i>	<i>cobC</i>	positive	Contig_17_46.7174	1109277	1E+06	-
MMSF00132	<i>Labrenzia aggregata</i>	<i>cobD</i>	positive x2	Contig_26_56.6932	208207	209184	-
MMSF00132	<i>Labrenzia aggregata</i>	<i>cobD</i>	positive x2	Contig_26_56.6932	209194	210195	-
MMSF00249	<i>Labrenzia aggregata</i>	<i>cobD</i>	positive x2	Contig_15_47.504	84229	85230	+
MMSF00249	<i>Labrenzia aggregata</i>	<i>cobD</i>	positive x2	Contig_15_47.504	85240	86217	+
MMSF00132	<i>Labrenzia aggregata</i>	<i>cobH</i>	positive	Contig_6_57.2063	545294	545950	+
MMSF00249	<i>Labrenzia aggregata</i>	<i>cobH</i>	positive	Contig_17_46.7174	811823	812479	+
MMSF00132	<i>Labrenzia aggregata</i>	<i>cobI</i>	positive	Contig_6_57.2063	547369	548139	+
MMSF00249	<i>Labrenzia aggregata</i>	<i>cobI</i>	positive	Contig_17_46.7174	813898	814668	+
MMSF00132	<i>Labrenzia aggregata</i>	<i>cobK</i>	positive	Contig_6_57.2063	555025	555786	+
MMSF00249	<i>Labrenzia aggregata</i>	<i>cobK</i>	positive	Contig_17_46.7174	821554	822315	+
MMSF00132	<i>Labrenzia aggregata</i>	<i>cobL</i>	positive	Contig_6_57.2063	546203	547417	+
MMSF00249	<i>Labrenzia aggregata</i>	<i>cobL</i>	positive	Contig_17_46.7174	812732	813946	+
MMSF00132	<i>Labrenzia aggregata</i>	<i>cobM</i>	positive	Contig_6_57.2063	550028	550801	+
MMSF00249	<i>Labrenzia aggregata</i>	<i>cobM</i>	positive	Contig_17_46.7174	816557	817330	+
MMSF00132	<i>Labrenzia aggregata</i>	<i>cobN</i>	positive	Contig_26_56.6932	201965	205723	+
MMSF00249	<i>Labrenzia aggregata</i>	<i>cobN</i>	positive	Contig_15_47.504	88701	92459	-
MMSF00132	<i>Labrenzia aggregata</i>	<i>cobO</i>	positive	Contig_26_56.6932	205720	206358	+
MMSF00249	<i>Labrenzia aggregata</i>	<i>cobO</i>	positive	Contig_15_47.504	88066	88704	-
MMSF00964	<i>Marinobacter salsuginis</i>	<i>cobO</i>	positive	Contig_25_36.8412	168373	168930	+
MMSF01190	<i>Marinobacter salsuginis</i>	<i>cobO</i>	positive	Contig_21_84.3861	85977	86534	+
MMSF00132	<i>Labrenzia aggregata</i>	<i>cobP</i>	positive	Contig_26_56.6932	200255	200806	+
MMSF00249	<i>Labrenzia aggregata</i>	<i>cobP</i>	positive	Contig_15_47.504	93618	94169	-
MMSF00132	<i>Labrenzia aggregata</i>	<i>cobQ</i>	positive	Contig_26_56.6932	206715	208205	+
MMSF00249	<i>Labrenzia aggregata</i>	<i>cobQ</i>	positive	Contig_15_47.504	86219	87709	-

Supplemental Tables

MMSF00132	<i>Labrenzia aggregata</i>	<i>cobU</i>	positive	Contig_24_56.1446	78969	79994	+
MMSF00249	<i>Labrenzia aggregata</i>	<i>cobU</i>	positive	Contig_13_45.8339	40345	41370	+
MMSF00132	<i>Labrenzia aggregata</i>	<i>cobV</i>	positive	Contig_24_56.1446	77990	78862	-
MMSF00249	<i>Labrenzia aggregata</i>	<i>cobV</i>	positive	Contig_13_45.8339	39366	40238	-
MMSF00132	<i>Labrenzia aggregata</i>	<i>cbiD</i>	positive	Contig_6_57.2063	553941	555035	+
MMSF00249	<i>Labrenzia aggregata</i>	<i>cbiD</i>	positive	Contig_17_46.7174	820470	821564	+
MMSF00964	<i>Marinobacter salsuginis</i>	<i>cbiO</i>	positive	Contig_36_38.8121	420124	420720	+

740

741

742 **Table S4:** Summary of vitamin B₁₂ biosynthesis pathway genes. A “+” indicates the presence of the gene in the respective isolate,
 743 whereas a “-” represents the absence of that gene. Genes in red were not found in any isolate.

Isolate	Genus Level Classification	CobP as indicator	Gene Count	<i>CobA</i>	<i>CobB</i>	<i>CobC</i>	<i>CobD</i>	<i>CobE</i>	<i>CobF</i>	<i>CobG</i>
MMSF01163	<i>Alteromonas</i>	-	0	-	-	-	-	-	-	-
MMSF00257	<i>Alteromonas</i>	-	0	-	-	-	-	-	-	-
MMSF00958	<i>Alteromonas</i>	-	0	-	-	-	-	-	-	-
MMSF00404	<i>Alteromonas</i>	-	0	-	-	-	-	-	-	-
MMSF00132	<i>Labrenzia</i>	+	18	+	+	+	+ x2	-	-	-
MMSF00249	<i>Labrenzia</i>	+	18	+	+	+	+ x2	-	-	-
MMSF00964	<i>Marinobacter</i>	-	2	-	-	-	-	-	-	-
MMSF01190	<i>Marinobacter</i>	-	1	-	-	-	-	-	-	-
MMSF00068	<i>Micrococcus</i>	-	0	-	-	-	-	-	-	-
MMSF00107	<i>Micrococcus</i>	-	0	-	-	-	-	-	-	-
MMSF00910	<i>Winogradskyella</i>	-	0	-	-	-	-	-	-	-
MMSF00046	<i>Winogradskyella</i>	-	0	-	-	-	-	-	-	-

744

Isolate	Genus Level Classification	<i>CobH</i>	<i>CobI</i>	<i>CobJ</i>	<i>CobK</i>	<i>CobL</i>	<i>CobM</i>	<i>CobN</i>	<i>CobO</i>	<i>CobP</i>	<i>CobQ</i>
MMSF01163	<i>Alteromonas</i>	-	-	-	-	-	-	-	-	-	-
MMSF00257	<i>Alteromonas</i>	-	-	-	-	-	-	-	-	-	-
MMSF00958	<i>Alteromonas</i>	-	-	-	-	-	-	-	-	-	-
MMSF00404	<i>Alteromonas</i>	-	-	-	-	-	-	-	-	-	-
MMSF00132	<i>Labrenzia</i>	+	+	-	+	+	+	+	+	+	+
MMSF00249	<i>Labrenzia</i>	+	+	-	+	+	+	+	+	+	+
MMSF00964	<i>Marinobacter</i>	-	-	-	-	-	-	-	+	-	-
MMSF01190	<i>Marinobacter</i>	-	-	-	-	-	-	-	+	-	-
MMSF00068	<i>Micrococcus</i>	-	-	-	-	-	-	-	-	-	-

Supplemental Tables

MMSF00107	<i>Micrococcus</i>	-	-	-	-	-	-	-	-	-	-	-
MMSF00910	<i>Winogradskyella</i>	-	-	-	-	-	-	-	-	-	-	-
MMSF00046	<i>Winogradskyella</i>	-	-	-	-	-	-	-	-	-	-	-

745

Isolate	Genus Level Classification	<i>CobP</i>	<i>CobQ</i>	<i>CobR</i>	<i>CobS</i>	<i>CobT</i>	<i>CobU</i>	<i>CobV</i>	<i>CobW</i>	<i>CobX</i>	<i>CobY</i>	<i>CobZ</i>	<i>Cbi</i>
MMSF01163	<i>Alteromonas</i>	-	-	-	-	-	-	-	-	-	-	-	-
MMSF00257	<i>Alteromonas</i>	-	-	-	-	-	-	-	-	-	-	-	-
MMSF00958	<i>Alteromonas</i>	-	-	-	-	-	-	-	-	-	-	-	-
MMSF00404	<i>Alteromonas</i>	-	-	-	-	-	-	-	-	-	-	-	-
MMSF00132	<i>Labrenzia</i>	+	+	-	-	-	+	+	-	-	-	-	cbiD
MMSF00249	<i>Labrenzia</i>	+	+	-	-	-	+	+	-	-	-	-	cbiD
MMSF00964	<i>Marinobacter</i>	-	-	-	-	-	-	-	-	-	-	-	cbiO
MMSF01190	<i>Marinobacter</i>	-	-	-	-	-	-	-	-	-	-	-	-
MMSF00068	<i>Micrococcus</i>	-	-	-	-	-	-	-	-	-	-	-	-
MMSF00107	<i>Micrococcus</i>	-	-	-	-	-	-	-	-	-	-	-	-
MMSF00910	<i>Winogradskyella</i>	-	-	-	-	-	-	-	-	-	-	-	-
MMSF00046	<i>Winogradskyella</i>	-	-	-	-	-	-	-	-	-	-	-	-

746

Supplemental Tables

Table S5: Composition of R2A broth adjusted to suit marine bacteria. Final pH = 7.2 +/- 0.2 at 26 °C. R2A broth was made by suspending 43.12 g of combined reagents in 1 L of MilliQ water, dissolving the medium completely, and sterilization by autoclaving at 121°C for 15 min.

Component	grams L ⁻¹	Supplier
Casein acid hydrolysate	0.500	Cat#C0501, Sigma-Aldrich
Yeast extract	0.500	Cat#LP0021, Oxoid
Proteose peptone	0.500	Cat#211684, ThermoFisher
Dextrose	0.500	Cat#G360, PhytoTech Laboratories
Starch, soluble	0.500	Cat#AJA526, Univar
Dipotassium phosphate	0.300	Cat#P3786, Sigma-Aldrich
Magnesium sulfate	0.024	Cat#M2643, Sigma-Aldrich
Sodium pyruvate	0.300	Cat#P2256, Sigma-Aldrich
Red Sea Salt™	40.00	Cat#R11065, Red Sea

Supplemental Tables

751 **Table S6:** Single-copy housekeeping genes extracted from the RAST annotations

Gene	Function	NCBI
16S	SSU rRNA ## 16S rRNA, small subunit ribosomal RNA	16S ribosomal RNA
23S	LSU rRNA ## 23S rRNA, large subunit ribosomal RNA	23S ribosomal RNA
fusA	Translation elongation factor G	elongation factor G
gyrB	DNA gyrase subunit B (EC 5.99.1.3)	DNA gyrase subunit B
gyrA	DNA gyrase subunit A (EC 5.99.1.3)	DNA gyrase subunit A
pyrG	CTP synthase (EC 6.3.4.2)	CTP synthetase
lepA	Translation elongation factor LepA	GTP-binding protein LepA
recA	RecA protein	recA protein
recG	ATP-dependent DNA helicase RecG (EC 3.6.4.12)	ATP-dependent DNA helicase RecG
rpoB	DNA-directed RNA polymerase beta subunit (EC 2.7.7.6)	DNA-directed RNA polymerase subunit beta
rpoD	RNA polymerase sigma factor RpoD	RNA polymerase sigma factor RpoD
atpD	ATP synthase beta chain (EC 3.6.3.14)	FOF1 ATP synthase subunit beta
ppk	Polyphosphate kinase (EC 2.7.4.1)	polyphosphate kinase
polA	DNA polymerase I (EC 2.7.7.7)	DNA polymerase I
pheSa	Phenylalanyl-tRNA synthetase alpha chain (EC 6.1.1.20)	phenylalanyl-tRNA synthetase subunit alpha
pheSb	Phenylalanyl-tRNA synthetase beta chain (EC 6.1.1.20)	phenylalanyl-tRNA synthetase subunit beta
ileS	Isoleucyl-tRNA synthetase (EC 6.1.1.5)	isoleucyl-tRNA synthetase
leuS	Leucyl-tRNA synthetase (EC 6.1.1.4)	leucyl-tRNA synthetase
aspS	Aspartyl-tRNA synthetase (EC 6.1.1.12)	aspartyl-tRNA synthetase
alaS	Alanyl-tRNA synthetase (EC 6.1.1.7)	alanyl-tRNA synthetase
argS	Arginyl-tRNA synthetase (EC 6.1.1.19)	arginyl-tRNA synthetase
hisS	Histidyl-tRNA synthetase (EC 6.1.1.21)	histidyl-tRNA synthetase
valS	Valyl-tRNA synthetase (EC 6.1.1.9)	valyl-tRNA synthetase
srp1	SSU ribosomal protein S1p	30S ribosomal protein S1
srp2	SSU ribosomal protein S2p (SAe)	30S ribosomal protein S2
srp3	SSU ribosomal protein S3p (S3e)	30S ribosomal protein S3
srp4	SSU ribosomal protein S4p (S9e)	30S ribosomal protein S4
srp5	SSU ribosomal protein S5p (S2e)	30S ribosomal protein S5
srp6	SSU ribosomal protein S6p	30S ribosomal protein S6
srp7	SSU ribosomal protein S7p (S5e)	30S ribosomal protein S7
srp8	SSU ribosomal protein S8p (S15Ae)	30S ribosomal protein S8
srp9	SSU ribosomal protein S9p (S16e)	30S ribosomal protein S9
srp10	SSU ribosomal protein S10p (S20e)	30S ribosomal protein S10

Supplemental Tables

srp11	SSU ribosomal protein S11p (S14e)	30S ribosomal protein S11
srp12	SSU ribosomal protein S12p (S23e)	30S ribosomal protein S12
srp13	SSU ribosomal protein S13p (S18e)	30S ribosomal protein S13
srp14	SSU ribosomal protein S14p (S29e) @ SSU ribosomal protein S14p (S29e), zinc-independent	30S ribosomal protein S14
srp15	SSU ribosomal protein S15p (S13e)	30S ribosomal protein S15
srp16	SSU ribosomal protein S16p	30S ribosomal protein S16
srp17	SSU ribosomal protein S17p (S11e)	30S ribosomal protein S17
srp18	SSU ribosomal protein S18p @ SSU ribosomal protein S18p, zinc-independent	30S ribosomal protein S18
srp19	SSU ribosomal protein S19p (S15e)	30S ribosomal protein S19
srp20	SSU ribosomal protein S20p	30S ribosomal protein S20
srp21	SSU ribosomal protein S21p	30S ribosomal protein S21
lrp1	LSU ribosomal protein L1p (L10Ae)	50S ribosomal protein L1
lrp2	LSU ribosomal protein L2p (L8e)	50S ribosomal protein L2
lrp3	LSU ribosomal protein L3p (L3e)	50S ribosomal protein L3
lrp4	LSU ribosomal protein L4p (L1e)	50S ribosomal protein L4
lrp5	LSU ribosomal protein L5p (L11e)	50S ribosomal protein L5
lrp6	LSU ribosomal protein L6p (L9e)	50S ribosomal protein L6
lrp7	LSU ribosomal protein L7/L12 (P1/P2)	50S ribosomal protein L7
lrp8	LSU ribosomal protein L8p	50S ribosomal protein L8
lrp9	LSU ribosomal protein L9p	50S ribosomal protein L9
lrp10	LSU ribosomal protein L10p (P0)	50S ribosomal protein L10
lrp11	LSU ribosomal protein L11p (L12e)	50S ribosomal protein L11
lrp12	LSU ribosomal protein L12p	50S ribosomal protein L12
lrp13	LSU ribosomal protein L13p (L13Ae)	50S ribosomal protein L13
lrp14	LSU ribosomal protein L14p (L23e)	50S ribosomal protein L14
lrp15	LSU ribosomal protein L15p (L27Ae)	50S ribosomal protein L15
lrp16	LSU ribosomal protein L16p (L10e)	50S ribosomal protein L16
lrp17	LSU ribosomal protein L17p	50S ribosomal protein L17
lrp18	LSU ribosomal protein L18p (L5e)	50S ribosomal protein L18
lrp19	LSU ribosomal protein L19p	50S ribosomal protein L19
lrp20	LSU ribosomal protein L20p	50S ribosomal protein L20
lrp21	LSU ribosomal protein L21p	50S ribosomal protein L21
lrp22	LSU ribosomal protein L22p (L17e)	50S ribosomal protein L22
lrp23	LSU ribosomal protein L23p (L23Ae)	50S ribosomal protein L23
lrp24	LSU ribosomal protein L24p (L26e)	50S ribosomal protein L24
lrp25	LSU ribosomal protein L25p	50S ribosomal protein L25
lrp26	LSU ribosomal protein L26p	50S ribosomal protein L26
lrp27	LSU ribosomal protein L27p	50S ribosomal protein L27
lrp28	LSU ribosomal protein L28p	50S ribosomal protein L28

Supplemental Tables

lrp29	LSU ribosomal protein L29p (L35e)	50S ribosomal protein L29
lrp30	LSU ribosomal protein L30p (L7e)	50S ribosomal protein L30
lrp31	LSU ribosomal protein L31p @ LSU ribosomal protein L31p, zinc-independent	50S ribosomal protein L31

752