North American deer mice are susceptible to SARS-CoV-2

Bryan D. Griffin¹, Mable Chan¹, Nikesh Tailor¹, Emelissa J. Mendoza¹, Anders Leung¹, Bryce M. Warner^{1,2}, Ana T. Duggan³, Estella Moffat⁴, Shihua He¹, Lauren Garnett^{1,2}, Kaylie N. Tran¹, Logan Banadyga¹, Alixandra Albietz¹, Kevin Tierney¹, Jonathan Audet¹, Alexander Bello¹, Robert Vendramelli¹, Amrit S. Boese¹, Lisa Fernando¹, L. Robbin Lindsay^{1,5}, Claire M. Jardine⁶, Heidi Wood¹, Guillaume Poliquin^{2,7,8}, James E. Strong^{1,2,7}, Michael Drebot^{1,2}, David Safronetz^{1,2}, Carissa Embury-Hyatt⁴, Darwyn Kobasa^{1,2,*}

Affiliations:

¹Zoonotic Diseases and Special Pathogens Program, National Microbiology Laboratory, Public Health Agency of Canada, 1015 Arlington Street, Winnipeg, Manitoba, Canada R3E 3R2

²Department of Medical Microbiology and Infectious Diseases, College of Medicine, Faculty of Health Sciences, University of Manitoba, 745 Bannatyne Avenue, Winnipeg, Manitoba, Canada R3E 0J9

³Science Technology Cores and Services, National Microbiology Laboratory, Public Health Agency of Canada, 1015 Arlington Street, Winnipeg, Manitoba, Canada R3E 3R2.

⁴National Centre for Foreign Animal Disease, Canadian Food Inspection Agency, 1015 Arlington Street, Winnipeg, Manitoba, Canada R3E 3M4

⁵Department of Entomology, University of Manitoba, Winnipeg, Manitoba, Canada R3T 2N2,

⁶Department of Pathobiology, Canadian Wildlife Health Cooperative, Department of Pathobiology, University of Guelph, Guelph, Ontario, Canada N1G 2W1

⁷Pediatrics & Child Health, College of Medicine, Faculty of Health Sciences, University of Manitoba, 745 Bannatyne Avenue, Winnipeg, Manitoba, Canada R3E 0J9

⁸Office of the Scientific Director, National Microbiology Laboratories, Public Health Agency of Canada, Winnipeg, MB, Canada R3E 3R2

*Corresponding author: darwyn.kobasa@canada.ca

Keywords: Severe Acute Respiratory Syndrome-coronavirus 2, SARS-CoV-2, coronavirus disease 2019, COVID-19, pandemic, angiotensin-converting enzyme 2 (Ace2), virology, infection, host specificity, species specificity, zoonoses, zoonanthroponosis, reverse zoonosis, animal reservoir, re-emergence, virus shedding, virus transmission, real-time polymerase chain reaction, *Cricetidae*, *Mesocricetus auratus*, deer mice, *Peromyscus maniculatus*, white footed mouse, *Peromyscus leucopus*

Abstract

The zoonotic spillover of the pandemic SARS-coronavirus 2 (SARS-CoV-2) from an animal reservoir, currently presumed to be the Chinese horseshoe bat, into a naïve human population has rapidly resulted in a significant global public health emergency. Worldwide circulation of SARS-CoV-2 in humans raises the theoretical risk of reverse zoonosis events with wildlife, reintroductions of SARS-CoV-2 into permissive non-domesticated animals, potentially seeding new host reservoir species and geographic regions in which bat SARS-like coronaviruses have not historically been endemic. Here we report that North American deer mice (*Peromyscus maniculatus*) and some closely related members of the *Cricetidae* family of rodents possess key amino acid residues within the angiotensin-converting enzyme 2 (ACE2) receptor known to confer SARS-CoV-2 spike protein binding. *Peromyscus* rodent species are widely distributed across North America and are the primary host reservoirs of several emerging pathogens that repeatedly spill over into humans including *Borrelia burgdorferi*, the causative agent of Lyme disease, deer tick virus, and Sin Nombre orthohantavirus, the causative agent of hantavirus pulmonary syndrome (HPS). We demonstrate that adult deer mice are susceptible to SARS-CoV-2 infection following intranasal exposure to a human isolate, resulting in viral replication in the upper and lower respiratory tract with little or no signs of disease. Further, shed infectious virus is detectable in nasal washes, oropharyngeal and rectal swabs, and viral RNA is detectable in feces and occasionally urine. We further show that deer mice are capable of transmitting SARS-CoV-2 to naïve deer mice through direct contact. The extent to which these observations may translate to wild deer mouse populations remains unclear, and the risk of reverse zoonosis and/or the potential for the establishment of *Peromyscus* rodents as a North American reservoir for SARS-CoV-2 is unknown. Nevertheless, efforts to monitor wild, peri-domestic *Peromyscus* rodent populations are likely warranted as the SARS-CoV-2 pandemic progresses.

Main text

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is the causative agent of coronavirus disease 2019 (COVID-19), an acute respiratory illness that emerged in Wuhan, China in late 2019¹, initiating a global pandemic associated with millions of infections, hundreds of thousands of deaths, and severe social and economic disruption. SARS-CoV-2, a positive-sense RNA virus of the betacoronavirus genus, is closely related to the high-consequence respiratory viruses, SARS-CoV and Middle East respiratory syndrome coronavirus (MERS-CoV). While the majority of SARS-CoV-2 infections result in asymptomatic^{2,3} or mild respiratory disease^{4,5} some individuals progress to more severe disease that can result in admittance to an intensive care unit (ICU) and/or the need for oxygen therapy or invasive mechanical ventilation and can result in serious multi-organ sequelae and death. The severity of disease and likelihood of mortality often correlates with advanced age, compromised immune status, and the pre-existence of certain cardiovascular, pulmonary, and metabolic comorbidities^{4,6}. Human-to-human transmission of SARS-CoV-2 is believed to occur primarily through short-range respiratory droplets and aerosols emitted by infected individuals during coughing, sneezing, speaking, breathing, or as a result of aerosol-generating medical procedures⁷. Individuals ranging from pre-symptomatic, asymptomatic, mildly symptomatic to fulminant symptomatic are thought to be capable of transmitting SARS-CoV-2^{8,9}.

Like the other non-seasonal human coronaviruses, SARS-CoV^{10,11} and MERS-CoV^{12,13}, SARS-CoV-2 is believed to be of bat origin^{11,14,15} although related viruses have also been identified in wild pangolins¹⁶. Examinations of amino acid sequences of the SARS-CoV-2 receptor, angiotensin-converting enzyme 2 (Ace2), from various vertebrate species predicts that additional wild animal species may be susceptible^{17,18}. Several experimental animal models of SARS-CoV-2 infection have been reported including transgenic mice that express human Ace2 (hAce2)^{19,20} and mouse-adapted SARS-CoV-2^{21,22} as well as tree shrews²³, hamsters^{24,25}, ferrets^{26,27}, fruit bats²⁸, rhesus macaques²⁹, cynomolgus macaques³⁰, marmosets³⁰, and African green monkeys³¹ that are innately susceptible to infection with wildtype SARS-CoV-2. Experimental transmission by both direct and indirect contact has been reported for hamsters^{24,25}, ferrets²⁷, and domestic cats^{26,32}. Lastly, there have been documented instances of unintentional human-to-animal transmission (zoonothroponosis) with several species of captive or domesticated animals including tigers³³, mink³⁴, cats^{26,35}, and dogs³⁶.

After Syrian hamsters (*Mesocricetus auratus*) were identified to be a SARS-CoV-2-susceptible host species^{24,25}, we sought to determine whether or not deer mice (*Peromyscus maniculatus*) were also susceptible hosts since they, like hamsters, are members of the *Cricetidae* family of rodents (Fig. 1a). *Peromyscus* species rodents are susceptible to persistent infections with and are the natural host reservoirs of several important zoonotic pathogens including: *Borrelia burgdorferi*, the causative agent of Lyme disease^{37,38}, deer tick virus (DTV)³⁹, and Sin Nombre orthohantavirus (SNV)^{40,41}. Deer mice and closely related white-footed mice (*Peromyscus leucopus*) are widely distributed across North America (Fig. 1b). We examined deer mouse Ace2 for congruence with human Ace2 (hAce2) at key amino acid residues known to confer efficient binding to the SARS-CoV-2 Spike (S) (Fig. 1c) and came to a conclusion consistent with recent reports that deer mice were likely to be susceptible hosts^{17,42}. We found that deer mouse Ace2 differed from hAce2 S-contacting residues at four locations and that the amino acid differences were unlikely to have a detrimental effect on S binding efficiency⁴³. We further found that Ace2 belonging to the white-footed mouse had in common all but three hAce2 S-contacting residues, and white-footed mouse Ace2 has since been shown to bind S and confer entry to SARS-CoV-2 into Hela cells¹⁸.

Here, eight to thirty two-week old male or female deer mice (*P. maniculatus rufinus*; in-house colony) were inoculated with 10⁵ TCID₅₀ of SARS-CoV-2 by an intranasal route (i.n.) and monitored daily for clinical signs and weight loss for 21 days or necropsied at 2 and 4 day post-infection (dpi). SARS-CoV-2-exposed deer mice seldom displayed clinical signs (occasionally presenting with ruffled fur), and no animals succumbed to infection (Fig. 2a) or lost weight (Fig. 2b) at any point post-infection with no differences noted related to age. At 2 dpi high levels of viral RNA (vRNA; ~10⁸-10¹⁰ genome equivalents/g) were detected in the nasal turbinates and lungs, and infectious virus (up to 3.3x10⁷ TCID₅₀/g and 1.5x10⁷ TCID₅₀/g) was detected in the nasal turbinates and lungs (Fig. 2 c-d). At 4 dpi infectious virus detected in the nasal turbinates had declined by several logs while lungs titers had declined to a lesser extent (ranging from 3x10³ to 1.4x10⁵ TCID₅₀/g and 6.8x10⁴ to 3.4x10⁶ TCID₅₀/g, respectively). Age did not make a significant contribution to the viral burden detected in the tissues (Extended Data Fig. 1). At 2 and 4 dpi high levels of vRNA (~10⁵ – 10⁸ genome copies/g) were detected in the small intestine and colon tissues (Fig. 2 e-f), and infectious virus was present at a low level. To monitor viral shedding, oropharyngeal and rectal swabs were collected every 2 days from 2-8 dpi (Fig. 2g-h). High levels of vRNA (~10⁵ – 10⁷ genome copies/ml) were detected on 2 and 4 dpi in oropharyngeal swabs, and both vRNA and infectious virus peaked at 2 dpi and declined

thereafter (Fig. 2g). Moderate levels of vRNA ($\sim 10^3 - 10^5$ genome copies/ml) were detected on 2 and 4 dpi in rectal swabs, and vRNA peaked at 2 dpi and declined thereafter (Fig. 2h). Infectious virus in rectal swabs was detected on 2, 4, 6, and 8 dpi at or near the limit of detection (Fig. 2h). Viral RNA was detected in nasal washes at 6 dpi in the four deer mice for which samples were obtained ($\sim 10^5 - 10^6$ genome copies/ml) (Fig. 2i). An additional experimental group of deer mice ($n = 5$) were inoculated i.n. with 10^6 TCID₅₀ of SARS-CoV-2. In this group vRNA was detected in urine samples at 6 dpi in one deer mouse ($\sim 10^4$ genome copies/ml) (Fig. 2j) and feces at 4 dpi in all deer mice ($\sim 10^4 - 10^6$ genome copies/g) (Fig. 2k). Viral RNA in blood was detected on 1 dpi, indicating transient viremia that declined in most animals by 3 dpi (Fig. 2l). Studies to examine the vectorial capacity of selected tick species such as *Ixodes scapularis* for SARS-CoV2 may be warranted. An additional experimental group of deer mice ($n = 4$) was inoculated i.n. with 10^5 TCID₅₀ of SARS-CoV-2, and tissues were harvested at 21 dpi (Fig. 2m). In this group vRNA was detected in the nasal turbinates ($\sim 10^5 - 10^7$ genome copies/g) and lungs ($\sim 10^4 - 10^6$ genome copies/g) of all deer mice and in the small intestine of two deer mice ($\sim 10^4$ genome copies/g), indicating persistence of vRNA in organs that to our knowledge has not been reported in other animal models studied thus far.

Lesions in animals infected at 10^5 TCID₅₀ were largely absent from nasal turbinates except for mild neutrophilic infiltration in the submucosa at 4 dpi (Fig. 3a, left panel). Abundant vRNA was primarily detected in epithelial cells at 2 and 4 dpi (Fig. 3a, middle panels) and anti-genomic RNA could be detected occasionally in epithelial cells at 2 dpi only (Fig. 3a, right panels). Histopathologic examination of lung tissue revealed lesions that included perivascular and peribronchiolar infiltrations of histiocytes and neutrophils with occasional multinucleated syncytial cells (Fig. 3b, left panel). Occasional discrete foci of interstitial pneumonia were observed (Fig. 3b, left panel). Abundant vRNA was primarily detected in bronchiolar epithelial cells and occasionally in the interstitium (Fig. 3b, middle panels). Anti-genomic RNA could be detected in individual scattered bronchiolar epithelial cells (Fig. 3b, right panels). Lung lesions were largely absent at 21 dpi although some foci of inflammation with syncytia were identified, and vRNA was not detected in the lungs by ISH despite qRT-PCR positivity (Extended Data Fig. 2).

Blood biochemistry and hematological parameters were compared for uninfected ($n = 5$) and SARS-CoV-2-infected deer mice ($n = 3$; 10^6 TCID₅₀) (Fig. 4a-e). Both groups showed similar levels of white blood cells; however, infected deer mice had reduced lymphocyte counts and elevated neutrophil counts (Fig. 4a). These data appear to recapitulate the trend in hematologic values observed in COVID-19 patients, where lymphocytopenia with elevated neutrophil counts can occur⁴⁴. One out of three infected deer mice had a dramatically elevated neutrophil-to-lymphocyte rate (NLR), a clinical metric that has been found to be elevated in human patients during severe COVID-19⁴⁵, and the NLR was significantly elevated in all infected deer mice ($P < 0.05$, Mann-Whitney test). The lack of a difference in alanine aminotransferase (ALT) (Fig. 4c), and the reduced values for blood albumin (ALB) (Fig. 4d) and blood urea nitrogen (BUN) (Fig. 4e), when uninfected and infected deer mice were compared suggests that infection did not cause liver and kidney impairment. Additional serum biochemical values were unremarkable (Extended Data Fig. 3). Transcriptional profiling of host cytokine expression in the lungs by real-time PCR using a previously reported panel that includes TNF α , IL-6, IL10, and IFN α ⁴⁶ was carried out 2 dpi (Fig. 4f) and compared to uninfected controls ($n = 6$). TNF α and IL-6 mRNA expression was elevated in all infected mice (mean fold changes of $2^{2.7}$ and $2^{3.5}$, respectively). IL-

10 mRNA expression was elevated in five out of six infected animals and reduced in one (mean fold change of $2^{1.5}$). In contrast, IFN α was elevated in two and reduced in four infected deer mice (mean fold change of $2^{-0.8}$). Overall, the mRNA levels of inflammatory cytokines (TNF α , IL-6, and IL10) were slightly elevated compared to what is seen in the lungs of deer mice ($\sim 2 - 5$ fold increase) following experimental infection with SNV⁴⁷. All five deer mice infected at 10^5 TCID₅₀ had detectable serum IgG titers against mixed spike/nucleoprotein (S/N) antigen as assessed by ELISA at 14 dpi (OD 2.7 – 2.8 at 1:100) (Fig. 4g), and neutralizing antibodies by 28 dpi (plaque reduction neutralization test (PRNT₉₀) 1:40 to 1:320) (Fig. 4h).

Transmission by direct contact was examined between SARS-CoV-2-infected and naïve deer mice (experimental design schematic, Extended Data Fig. 4). Five adult male and female deer mice, infected i.n. with 10^6 TCID₅₀ SARS-CoV-2 were each transferred to a new cage at 1 dpi and co-housed with a single naïve contact deer mouse (1:1 ratio; zero days post-direct contact (dpc)). Oropharyngeal and rectal swab samples taken every other day revealed that initially one contact animal (DM7, Fig. 5 a-b) rapidly became infected and shed moderate levels of vRNA as early as 2 dpc. The level of vRNA in both swab samples from DM7 continued to increase for the next 4 (oropharyngeal swabs) or 2 (rectal swabs) days then declined. The remaining contact deer mice had detectable vRNA in oropharyngeal and rectal swab samples at low levels from 2 - 4 dpc which declined until 10 dpc when shed vRNA was again detected in both oropharyngeal and rectal swabs from all contact deer mice (including DM7). Interestingly, despite the high challenge dose administered to donor deer mice only low amounts of infectious virus were detected in swabs (Fig. 5c-d). Infectious virus was detected in oropharyngeal and rectal swabs of contact deer mouse DM7 at 4 dpc and an oropharyngeal swab sample from contact deer mouse DM6 at 10 dpc. An additional transmission study was carried out in the same manner and nasal wash, nasal turbinates, and lung samples were collected from contact animals at 2 and 4 dpc (Fig. 5e-f). Viral RNA was detected in nasal wash, nasal turbinates, and the lungs of contact deer mice DM19 (2 dpc) and DM30 (4 dpc), as well as in the nasal wash of deer mouse DM17 (2 dpc), DM26 (4 dpc) and DM28 (4 dpc). IgG against mixed SARS-CoV-2 spike/nucleoprotein antigens were detected in deer mouse DM7 at 21 dpc, indicating seroconversion and confirming that direct contact transmission had occurred (Fig. 5g). A similar, but more stringent transmission study was carried out using a lower dose of virus (10^5 TCID₅₀), and donor and naïve deer mice were mixed at 2 dpi (Fig. 5h-i). Oropharyngeal and rectal swab samples taken every other day revealed that contact deer mice had detectable vRNA in both oropharyngeal and rectal swab samples from 2 - 4 dpc at a low levels, which declined until 6-10 dpc after which shed vRNA was again detected in oropharyngeal swabs from four out of five and rectal swabs from three out of five contacts. At 21 dpc vRNA was detected in the oropharyngeal swabs of two of five contact and was not detected in any rectal swab samples.

The described experiments demonstrate that adult deer mice are susceptible to experimental infection with a human isolate of SARS-CoV-2, resulting in asymptomatic infection or mild disease with lesions limited to mild lung pathology, despite a high viral burden and elevated levels of inflammatory cytokines in the lungs and seroconversion. The detected RNA viremia in SARS-CoV-2-infected deer mice appears to be more transient than what is observed in deer mice experimentally infected with SNV⁴⁸; however, the viral shedding and the propensity for direct contact transmission appears to be greater^{41,48}. The relatively low amount of shed infectious virus from nasal, oral, and rectal routes consistently result in direct contact transmission from infected

to co-housed naive deer mice. SNV is also detectable in deer mouse feces and urine, and zoonotic transmission of SNV into humans is thought to result from unintentional ingestion or inhalation of contaminated deer mouse excreta⁴⁹. Additional studies should be carried out to determine if vRNA detected in the feces and occasionally urine is indicative of the presence infectious virus.

Concerns have justifiably been raised about the potential for reverse zoonosis, human-to-animal transmission of SARS-CoV-2 into susceptible wild animal host species⁵⁰. The introduction of novel pathogens into susceptible wildlife hosts can have devastating effects on wildlife populations⁵¹. The findings reported here suggest that the impacts of SARS-CoV-2 on infected *Peromyscus* species rodents are likely to be minimal; however, wild deer mouse populations may be more or less susceptible to infection than the experimentally housed animals we have described. An additional concern is the potential for zoonotic maintenance of SARS-CoV-2 in an animal reservoir and/or geographic region where SARS-like coronaviruses had not previously been endemic. The findings reported here are concerning in light of the fact that *Peromyscus* species rodents tolerate persistent infection with and serve as the primary reservoirs for several emerging zoonotic pathogens that spillover into humans, including *Borrelia burgdorferi*^{37,38}, DTV³⁹, and SNV^{40,41}. It should be acknowledged that although deer mice are widely distributed in North America and can live in close proximity to humans, the actual risk of human-to-deer mouse transmission remains unknown. Further, it remains unclear how well the transmission studies we describe translate to the risk of sustained transmission within wild deer mouse populations. The theoretical risk of zoonotic transmission of SARS-CoV-2 back to humans from *Peromyscus* species rodents would depend on additional unknowns such as whether infectious virus is present in excreta at high enough levels to initiate human infection and whether other wild or domestic animal species might serve as an intermediary host. Efforts to monitor wild, especially peri-domestic *Peromyscus* rodent populations are likely warranted as the SARS-CoV-2 pandemic continues to progress.

Lastly, we suggest that the deer mouse model of SARS-CoV-2 infection, as a largely outbred small animal model, may prove useful for studying viral pathogenesis and for evaluating the protective efficacy of experimental vaccines and therapeutics.

References

1. Zhu, N. *et al.* A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N. Engl. J. Med.*, doi:10.1056/NEJMoa2001017 (2020).
2. Poletti, P. *et al.* Probability of symptoms and critical disease after SARS- CoV-2 infection, Preprint at: <https://arxiv.org/pdf/2006.08471.pdf> (2020).
3. O, B. *et al.* Estimating the Extent of True Asymptomatic COVID-19 and Its Potential for Community Transmission: Systematic Review and Meta-Analysis Preprint at: <https://www.medrxiv.org/content/10.1101/2020.05.10.20097543v2> (2020).
4. Huang, C. *et al.* Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* **395**, 497-506, doi:10.1016/S0140-6736(20)30183-5 (2020).
5. Guan, W. J. *et al.* Clinical Characteristics of Coronavirus Disease 2019 in China. *N. Engl. J. Med.* **382**, 1708-1720, doi:10.1056/NEJMoa2002032 (2020).
6. Zhou, F. *et al.* Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* **395**, 1054-1062, doi:10.1016/S0140-6736(20)30566-3 (2020).
7. Prather, K. A., Wang, C. C. & Schooley, R. T. Reducing transmission of SARS-CoV-2. *Science* **368**, 1422-1424, doi:10.1126/science.abc6197 (2020).

273 8 Li, R. *et al.* Substantial undocumented infection facilitates the rapid dissemination of novel coronavirus
274 (SARS-CoV-2). *Science* **368**, 489-493, doi:10.1126/science.abb3221 (2020).
275 9 Moghadas, S. M. *et al.* The implications of silent transmission for the control of COVID-19 outbreaks.
276 *Proc. Natl. Acad. Sci. U. S. A.*, doi:10.1073/pnas.2008373117 (2020).
277 10 Guan, Y. *et al.* Isolation and characterization of viruses related to the SARS coronavirus from animals in
278 southern China. *Science* **302**, 276-278, doi:10.1126/science.1087139 (2003).
279 11 Hu, B. *et al.* Discovery of a rich gene pool of bat SARS-related coronaviruses provides new insights into
280 the origin of SARS coronavirus. *PLoS Pathog.* **13**, e1006698, doi:10.1371/journal.ppat.1006698 (2017).
281 12 Memish, Z. A. *et al.* Middle East respiratory syndrome coronavirus in bats, Saudi Arabia. *Emerg. Infect.*
282 *Dis.* **19**, 1819-1823, doi:10.3201/eid1911.131172 (2013).
283 13 Lau, S. K. P. *et al.* Receptor Usage of a Novel Bat Lineage C Betacoronavirus Reveals Evolution of Middle
284 East Respiratory Syndrome-Related Coronavirus Spike Proteins for Human Dipeptidyl Peptidase 4
285 Binding. *J. Infect. Dis.* **218**, 197-207, doi:10.1093/infdis/jiy018 (2018).
286 14 Zhou, P. *et al.* A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature*,
287 doi:10.1038/s41586-020-2012-7 (2020).
288 15 Ge, X. Y. *et al.* Isolation and characterization of a bat SARS-like coronavirus that uses the ACE2 receptor.
289 *Nature* **503**, 535-538, doi:10.1038/nature12711 (2013).
290 16 Lam, T. T. *et al.* Identifying SARS-CoV-2-related coronaviruses in Malayan pangolins. *Nature*,
291 doi:10.1038/s41586-020-2169-0 (2020).
292 17 Damas, J. *et al.* Broad Host Range of SARS-CoV-2 Predicted by Comparative and Structural Analysis of
293 ACE2 in Vertebrates, Preprint at: <https://www.ncbi.nlm.nih.gov/pubmed/32511356>
294 (2020).
295 18 Liu, Y. *et al.* Functional and Genetic Analysis of Viral Receptor ACE2 Orthologs Reveals Broad Potential
296 Host Range of SARS-CoV-2, Preprint at: <https://www.biorxiv.org/content/10.1101/2020.04.22.046565v2>
297 (2020).
298 19 Bao, L. *et al.* The pathogenicity of SARS-CoV-2 in hACE2 transgenic mice. *Nature*, doi:10.1038/s41586-
299 020-2312-y (2020).
300 20 Hassan, A. O. *et al.* A SARS-CoV-2 Infection Model in Mice Demonstrates Protection by Neutralizing
301 Antibodies. *Cell*, doi:10.1016/j.cell.2020.06.011 (2020).
302 21 Dinno, K. H. *et al.* A mouse-adapted SARS-CoV-2 model for the evaluation of COVID-19 medical
303 countermeasures, Preprint at: <https://www.ncbi.nlm.nih.gov/pubmed/32511406> (2020).
304 22 Gu, H. *et al.* Rapid adaptation of SARS-CoV-2 in BALB/c mice: Novel mouse model for vaccine efficacy,
305 Preprint at: <https://www.biorxiv.org/content/10.1101/2020.05.02.073411v1> (2020).
306 23 Zhao, Y. *et al.* Susceptibility of tree shrew to SARS-CoV-2 infection, Preprint at:
307 <https://www.biorxiv.org/content/10.1101/2020.04.30.029736v1> (2020).
308 24 Chan, J. F. *et al.* Simulation of the clinical and pathological manifestations of Coronavirus Disease 2019
309 (COVID-19) in golden Syrian hamster model: implications for disease pathogenesis and transmissibility.
310 *Clin. Infect. Dis.*, doi:10.1093/cid/ciaa325 (2020).
311 25 Sia, S. F. *et al.* Pathogenesis and transmission of SARS-CoV-2 in golden hamsters. *Nature*,
312 doi:10.1038/s41586-020-2342-5 (2020).
313 26 Shi, J. *et al.* Susceptibility of ferrets, cats, dogs, and other domesticated animals to SARS-coronavirus 2.
314 *Science* **368**, 1016-1020, doi:10.1126/science.abb7015 (2020).
315 27 Kim, Y. I. *et al.* Infection and Rapid Transmission of SARS-CoV-2 in Ferrets. *Cell Host Microbe* **27**, 704-
316 709 e702, doi:10.1016/j.chom.2020.03.023 (2020).
317 28 Schlottau, K. *et al.* SARS-CoV-2 in fruit bats, ferrets, pigs, and chickens: an experimental transmission
318 study. *The Lancet Microbe*, doi:10.1016/S2666-5247(20)30089-6.
319 29 Munster, V. J. *et al.* Respiratory disease in rhesus macaques inoculated with SARS-CoV-2. *Nature*,
320 doi:10.1038/s41586-020-2324-7 (2020).
321 30 Lu, S. *et al.* Comparison of SARS-CoV-2 infections among 3 species of non-human primates, Preprint at:
322 <https://www.biorxiv.org/content/10.1101/2020.04.08.031807v1> (2020).
323 31 Woolsey, C. *et al.* Establishment of an African green monkey model for COVID-19, Preprint at:
324 <https://www.ncbi.nlm.nih.gov/pubmed/32511377> (2020).
325 32 Halfmann, P. J. *et al.* Transmission of SARS-CoV-2 in Domestic Cats. *N. Engl. J. Med.*,
326 doi:10.1056/NEJMc2013400 (2020).
327 33 United States Department of Agriculture. USDA Statement on the Confirmation of COVID-19 in a Tiger in
328 New York. (2020).

- 34 Oreshkova, N., Molenaar, R.-J., Vreman, S., Harders, F. & Munnink, B. B. O. SARS-CoV2 infection in farmed mink, Netherlands, April 2020, Preprint at: <https://www.biorxiv.org/content/10.1101/2020.05.18.101493v1> (2020).
- 35 Zhang, Q. *et al.* SARS-CoV-2 neutralizing serum antibodies in cats: a serological investigation, Preprint at: <https://www.biorxiv.org/content/10.1101/2020.04.01.021196v1> (2020).
- 36 Sit, T. H. C. *et al.* Infection of dogs with SARS-CoV-2. *Nature*, doi:10.1038/s41586-020-2334-5 (2020).
- 37 Bosler, E. M., Ormiston, B. G., Coleman, J. L., Hanrahan, J. P. & Benach, J. L. Prevalence of the Lyme disease spirochete in populations of white-tailed deer and white-footed mice. *Yale J. Biol. Med.* **57**, 651-659 (1984).
- 38 Brown, R. N. & Lane, R. S. Natural and experimental *Borrelia burgdorferi* infections in woodrats and deer mice from California. *J. Wildl. Dis.* **30**, 389-398, doi:10.7589/0090-3558-30.3.389 (1994).
- 39 El Khoury, M. Y. *et al.* Potential role of deer tick virus in Powassan encephalitis cases in Lyme disease-endemic areas of New York, U.S.A. *Emerg. Infect. Dis.* **19**, 1926-1933, doi:10.3201/eid1912.130903 (2013).
- 40 Knust, B. & Rollin, P. E. Twenty-year summary of surveillance for human hantavirus infections, United States. *Emerg. Infect. Dis.* **19**, 1934-1937, doi:10.3201/eid1912.131217 (2013).
- 41 Botten, J. *et al.* Persistent Sin Nombre virus infection in the deer mouse (*Peromyscus maniculatus*) model: sites of replication and strand-specific expression. *J. Virol.* **77**, 1540-1550, doi:10.1128/JVI.77.2.1540-1550.2003 (2003).
- 42 Luan, J., Lu, Y., Jin, X. & Zhang, L. Spike protein recognition of mammalian ACE2 predicts the host range and an optimized ACE2 for SARS-CoV-2 infection. *Biochem. Biophys. Res. Commun.* **526**, 165-169, doi:10.1016/j.bbrc.2020.03.047 (2020).
- 43 Procko, E. The sequence of human ACE2 is suboptimal for binding the S spike protein of SARS coronavirus 2, Preprint at: <https://www.biorxiv.org/content/10.1101/2020.03.16.994236v3> (2020).
- 44 Gong, J. *et al.* A Tool to Early Predict Severe 2019-Novel Coronavirus Pneumonia (COVID-19) : A Multicenter Study using the Risk Nomogram in Wuhan and Guangdong, China, Preprint at: <https://www.medrxiv.org/content/10.1101/2020.03.17.20037515v1.full.pdf> (2020).
- 45 Liu, J. *et al.* Neutrophil-to-Lymphocyte Ratio Predicts Severe Illness Patients with 2019 Novel Coronavirus in the Early Stage, Preprint at: <https://www.medrxiv.org/content/10.1101/2020.02.10.20021584v1> (2020).
- 46 Oko, L. *et al.* Profiling helper T cell subset gene expression in deer mice. *BMC Immunol* **7**, 18, doi:10.1186/1471-2172-7-18 (2006).
- 47 Schountz, T. *et al.* Kinetics of immune responses in deer mice experimentally infected with Sin Nombre virus. *J. Virol.* **86**, 10015-10027, doi:10.1128/JVI.06875-11 (2012).
- 48 Warner, B. M. *et al.* Development and Characterization of a Sin Nombre Virus Transmission Model in *Peromyscus maniculatus*. *Viruses* **11**, doi:10.3390/v11020183 (2019).
- 49 Brocato, R. L. & Hooper, J. W. Progress on the Prevention and Treatment of Hantavirus Disease. *Viruses* **11**, doi:10.3390/v11070610 (2019).
- 50 Leroy, E. M., Ar Gouilh, M. & Brugere-Picoux, J. The risk of SARS-CoV-2 transmission to pets and other wild and domestic animals strongly mandates a one-health strategy to control the COVID-19 pandemic. *One Health*, 100133, doi:10.1016/j.onehlt.2020.100133 (2020).
- 51 Blehert, D. S. *et al.* Bat white-nose syndrome: an emerging fungal pathogen? *Science* **323**, 227, doi:10.1126/science.1163874 (2009).

Methods References

- 52 Hall, E. *The mammals of North America*. (John Wiley and Sons, 1981).
- 53 Griffin, B. D. *et al.* Establishment of an RNA polymerase II-driven reverse genetics system for Nipah virus strains from Malaysia and Bangladesh. *Sci Rep* **9**, 11171, doi:10.1038/s41598-019-47549-y (2019).
- 54 Corman, V. M. *et al.* Detection of 2019 novel coronavirus (2019-nCoV) by real-time RT-PCR. *Euro Surveill* **25**, doi:10.2807/1560-7917.ES.2020.25.3.2000045 (2020).
- 55 Schountz, T., Shaw, T. I., Glenn, T. C., Feldmann, H. & Prescott, J. Expression profiling of lymph node cells from deer mice infected with Andes virus. *BMC Immunol* **14**, 18, doi:10.1186/1471-2172-14-18 (2013).
- 56 Livak, K. J. & Schmittgen, T. D. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* **25**, 402-408, doi:10.1006/meth.2001.1262 (2001).

- 57 Mendoza, E. J., Manguiat, K., Wood, H. & Drebot, M. Two Detailed Plaque Assay Protocols for the
Quantification of Infectious SARS-CoV-2. *Curr Protoc Microbiol* **57**, ecpmc105, doi:10.1002/cpmc.105
(2020).
- 58 Madeira, F. *et al.* The EMBL-EBI search and sequence analysis tools APIs in 2019. *Nucleic Acids Res.* **47**,
W636-W641, doi:10.1093/nar/gkz268 (2019).
- 59 Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7: improvements in
performance and usability. *Mol. Biol. Evol.* **30**, 772-780, doi:10.1093/molbev/mst010 (2013).
- 60 Castresana, J. Selection of conserved blocks from multiple alignments for their use in phylogenetic
analysis. *Mol. Biol. Evol.* **17**, 540-552, doi:10.1093/oxfordjournals.molbev.a026334 (2000).
- 61 Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies.
Bioinformatics **30**, 1312-1313, doi:10.1093/bioinformatics/btu033 (2014).
- 62 Darriba, D. *et al.* ModelTest-NG: A New and Scalable Tool for the Selection of DNA and Protein
Evolutionary Models. *Mol. Biol. Evol.* **37**, 291-294, doi:10.1093/molbev/msz189 (2020).

Figure Legends

Figure 1: Deer mouse range and predicted susceptibility to SARS-CoV-2.

a, Phylogeny showing the relationships among selected members of the *Cricetidae* family with mice and humans. **b**, Geographical distribution of deer mice (based on data from Hall, 1981)⁵². **c**, Alignment of human Ace2 (hAce2) amino acid residues known to confer efficient binding of the RBD of SARS-CoV-2 spike with the corresponding Ace2 amino acid residues from selected members of the *Cricetidae* family and other naturally or experimentally susceptible host species. The physiochemical properties of the amino acids residues are indicated: non-polar (yellow), polar (green), acidic (red), and basic residues (blue).

Figure 2: SARS-CoV-2 infection of adult deer mice.

a - m Eight to thirty two-week old female or male deer mice (*P. maniculatus*) were inoculated with 10⁵ TCID₅₀ or 10⁶ TCID₅₀ of SARS-CoV-2 by an intranasal route (i.n.) of administration and compared to age-matched uninfected controls. Bars indicate means. Dashed lines and dotted lines indicate the limit of detection for the TCID₅₀ assay and qRT-PCR assay, respectively. **a-b**, Kaplan-Meier curve depicting survival data (**a**) and weight data (**b**) over the course of 21 days following SARS-CoV-2 exposure (challenge dose, 10⁵ TCID₅₀). **c-f**, Infectious viral load (filled in circles, left axis) and vRNA levels (empty circles, right axis) in the (**c**) nasal turbinates, (**d**) lung, (**e**) small intestine, and (**f**) colon (challenge dose, 10⁵ TCID₅₀). **g-h**, Infectious viral load (filled in circles with bars, left axis) and viral RNA levels (empty circles, right axis) in (**g**) oropharyngeal swabs and (**h**) rectal swab samples (challenge dose, 10⁵ TCID₅₀). **i**, Viral RNA levels in nasal washes at 6 dpi (infectious dose, 10⁵ TCID₅₀). **j-l**, Viral RNA levels in the (**j**) urine, (**k**) feces, and (**l**) blood at the indicated times post-infection (infectious dose, 10⁶ TCID₅₀). **m**, Viral RNA levels in the nasal turbinates, lung, and small intestines at 21 days post-infection (infectious dose, 10⁵ TCID₅₀). Data were collected from two independent experiments. (a-b, n = 6; c-h, n = 3/6 for uninfected/infected; i-l, n = 5; m, n = 4).

Figure 3: Histopathology and virus distribution.

Hematoxylin/eosin (H&E) staining (left) and in situ hybridization (ISH) using antisense probes that detect the SARS-CoV-2 genome/mRNA (middle) and sense probes that detects anti-genomic RNA (right) were carried out on **a**) nasal turbinates and **b**) lung tissue of uninfected and SARS-CoV-2-infected deer mice (10⁵ TCID₅₀ i.n. route) at 2 and 4 dpi. Positive detection of viral genomic RNA/mRNA or anti-genomic RNA is indicated by magenta staining (middle/right panels and insets). Arrows indicate perivascular infiltrations of histiocytes and neutrophils

(black), peribronchiolar infiltrations of histiocytes and neutrophils (orange), mild neutrophilic infiltration in the submucosa (blue), occasionally observed discrete foci of interstitial pneumonia (purple), occasionally observed multinucleated syncytial cells (magenta), and anti-genomic RNA occasionally in individual scattered bronchiolar epithelial cells (green and right inset). The magnification is 20x for H&E and 20x for ISH unless otherwise indicated. Scale bars = 100 μ m, 50 μ m, and 20 μ m for 10x, 20x, and 40x, respectively.

Figure 4: Deer mouse host response to SARS-CoV-2 infection.

a-e, Hematological levels and serum biochemistry were measured in uninfected and SARS-CoV-2-infected deer mice (infectious dose 10^6 TCID₅₀, i.n. route) at 3 dpi, including **(a)** white blood cell, lymphocyte, and neutrophil counts, **(b)** the neutrophil-to-lymphocyte ratio, **(c)** alanine aminotransferase (ALT), **(d)** blood albumin (ALB), and **(e)** blood urea nitrogen (BUN). **f**, Cytokine gene expression was measured for TNF α , IL-6, IL-10, and IFN α in the lungs of SARS-CoV-2-infected deer mice and displayed relative to age-matched mock-infected animals. Gene expression was normalized using GAPDH as a control. **g**, IgG antibody response against SARS-CoV-2 mixed spike and nucleoprotein (S/N) antigens were assessed by ELISA using serum collected on the indicated days post-infection (infectious dose, 10^5 TCID₅₀). **h**, Neutralizing antibody against SARS-CoV-2 was measured by PRNT₉₀ using serum collected at 28 days after SARS-CoV-2 infection (infectious dose, 10^5 TCID₅₀). Dotted lines indicate the limit of detection. Bars indicate mean, error bars indicate SEM (c-e) or 95% confidence interval (f). (a-b, n = 5/3 for uninfected/infected; c-e, n = 5/4 for uninfected/infected, f, n = 6, g-h, n = 5 or 6) * = $P < 0.05$, ns = $P > 0.05$, Mann-Whitney test (b), unpaired t test (c-e).

Figure 5: Transmission of SARS-CoV-2 between deer mice through direct contact.

a-g, Adult male and female deer mice were exposed to 10^6 TCID₅₀ SARS-CoV-2 by an i.n. route of infection. At 1 dpi individual inoculated donor deer mice were transferred to a new cage and co-housed with a single naïve deer mouse (1:1 ratio) to assess SARS-CoV-2 transmission by direct contact. Deer mice were either maintained in direct contact throughout the 10 day study for serial swabbing or humanely euthanized on 2 or 4 days post-contact (dpc) for tissue collection. **a-d**, Viral RNA levels (**a**, **b**) and infectious viral loads (**c**, **d**) were measured in oropharyngeal swabs and rectal swab samples every other day from -1 to 10 days after direct contact was initiated. Oropharyngeal or rectal swab samples obtained from the same animal are demarcated with a unique colour, and samples derived from co-housed pairs share the same symbol. **e-f**, An additional transmission study was carried out in the same manner and nasal wash, nasal turbinates, and lung samples were collected from contact animals at **(e)** 2 dpc and **(f)** 4 dpc. **g**, IgG antibody response (reciprocal serum dilution) against SARS-CoV-2 S and N in donor and direct-contact exposed animals were assessed by ELISA using serum collected at 21 dpc. **h-i**, In an additional study adult male and female deer mice were exposed to 10^5 TCID₅₀ SARS-CoV-2 by an intranasal route of infection. At 2 dpi individual inoculated donor deer mice were transferred to a new cage and co-housed with a single naïve deer mouse (1:1 ratio) to assess SARS-CoV-2 transmission by direct contact (n = 5). Viral RNA levels were measured in **(h)** oropharyngeal swabs and **(i)** rectal swab samples every other day from 0 to 14 days after direct contact was initiated, and final swab samples were collected at 21 dpi and assessed. Dotted or dashed lines indicate the limit of detection. Bars indicate means. Data were collected from two independent experiments.

Methods

Ethics statement

The experiments described in this study were carried out at the National Microbiology Laboratory (NML) at the Public Health Agency of Canada as described in the Animal use document AUD# H-20-006. Experiment were approved by the Animal Care Committee located at the Canadian Science Center for Human and Animal Health in accordance with the guidelines provided by the Canadian Council on Animal Care. All procedures were performed under anesthesia, and all efforts were made to minimize animal suffering and to reduce the number of animals used. All infectious work was performed under biosafety level 3 (BSL-3) conditions or higher.

Viruses

The SARS-CoV-2 strain used in these studies (SARS-CoV-2; hCoV- 19/Canada/ON-VIDO-01/2020, GISAID accession# EPI_ISL_425177) was isolated from a clinical specimen obtained at the Sunnybrook Research Institute (SRI)/ University of Toronto on VeroE6 cells and provided by the Vaccine and Infectious Disease Organization (VIDO) with permission. The P1 virus was subsequently passaged at a 1:1000 dilution on mycoplasma-free VeroE6 cells (ATCC) in Dulbecco's Modified Eagle's Medium (Hyclone) containing 1% L-glutamine and 0.5 µg/ml of TPCK-trypsin and harvested when 80% cytopathic effect (CPE) became evident. The P2 virus stock was clarified by centrifugation at 6000 x g for 5 minutes and stored at -80°C until thawed for deer mouse infections. The virus stock was titrated on Vero cells by conventional TCID₅₀ assay, as described previously⁵³.

Deer mouse challenge experiments

Deer mice (*Peromyscus maniculatus rufinus*) used in these studies were supplied by a breeding colony housed at the University of Manitoba in a pathogen-free facility. All the deer mice were acclimated for a minimum of one week prior to the initiation of experimental procedures. Deer mice were randomly assigned to their respective groups and were housed in a temperature-controlled, light-cycled facility. Deer mice were supplied with food and water ad libitum and were monitored daily throughout the course of the experiments. Thirteen to thirty two-week old male or female deer mice were infected with 10⁵ or 10⁶ TCID₅₀ of SARS-CoV-2 by an intranasal route (i.n.) of administration in a 50 µl volume. Blinding of the animal experiments was not performed. Sample sizes were calculated a priori, and no animals were excluded from the data analysis although at times sample volumes were insufficient to allow downstream analyses.

Blood, fluids, feces, swab, and tissue collection

All deer mice were exsanguinated via cardiac puncture under deep isoflurane anesthesia prior to euthanization. Whole blood and serum was collected in BD microtainer tubes (K₂-EDTA or serum, respectively) (Becton, Dickinson and Company) as per the manufacturer's instructions. Serum samples were collected from the retro-orbital sinus under deep isoflurane sedation at 14, 21, and 28 dpi for challenge studies and at 21 dpc for transmission studies. Oropharyngeal and rectal swabs were taken using fine tip Dryswab Fine Tip rayon swabs (MWE cat #MW113). Harvested tissue samples for infectious assays were flash frozen and stored at -80 °C until later use, and swab samples were placed in 1 ml MEM supplemented with 1% heat-inactivated FBS, 1x L-glutamine,

and 2x penicillin–streptomycin, flash frozen, and stored at -80°C until later use. Tissue samples harvested for vRNA detection were immersed in RNeasy lysis buffer (Qiagen) at 4°C for 1 day, then stored at -80°C until later use. For urine collection deer mice were gently raised from their cages using soft-tip mouse handling forceps and placed immediately onto a 35-mm petri dish, a process that consistently allowed a minimum of 25 μl of urine to be collected. Feces samples were collected fresh from the isoflurane chamber following sedation, homogenized in 0.5 ml of 0.89% NaCl per 80 mg of feces using a Bead Ruptor Elite Bead Mill Homogenizer (Omni International) with a stainless steel bead at 4 m/s for 30 seconds. The sample was then clarified for 20 minutes at 4000 x g and the supernatant was filtered through a 0.2 μm filter. Viral RNA was then isolated using the QIAamp Viral RNA Mini kit (Qiagen), as per the manufacturer's instructions.

Infectious virus in tissues and swab samples

For infectious virus assays, thawed tissue samples were weighed and placed in 1 ml of MEM supplemented with 1% heat-inactivated FBS, 1x L-glutamine before being homogenized in a Bead Ruptor Elite Bead Mill Homogenizer (Omni International) at 4 m/s for 30 seconds and clarified by centrifugation at 1500 x g for 6 minutes. Swab samples were vortexed for 10 seconds and clarified at 2400 rpm for 5 minutes. Tissue homogenates or swab samples were serially diluted 10-fold in MEM supplemented with 1% heat-inactivated FBS, 1x L-glutamine, and 2x penicillin–streptomycin. One hundred microliter volumes of neat samples (swabs only) and sample dilutions were added to 96-well plates of 95% confluent Vero cells containing 50 μl of the same medium in replicates of three and incubated for 5 days at 37°C with 5% CO_2 . Plates were scored for the presence of cytopathic effect on day 5 after infection.

Viral RNA copies in tissues and swab samples

For vRNA copy number analysis, tissue samples were thawed and weighed and homogenized in 600 μl RLT buffer using a Bead Ruptor Elite Bead Mill Homogenizer (Omni International) with a stainless steel bead at 4 m/s for 30 seconds. Viral RNA from 30 mg tissue samples was extracted with the RNeasy Plus Mini kit (Qiagen), and vRNA from swab samples was extracted with the QIAamp Viral RNA Mini kit (Qiagen). A SARS-CoV-2 E-specific real-time RT–PCR assay was used for the detection of vRNA⁵⁴. RNA was reverse transcribed and amplified using the primers E_Sarbeco_F1 (5'- ACAGGTACGTTAATAGTTAATAGCGT-3') and E_Sarbeco_R2 (5'- ATATTGCAGCAGTACGCACACA-3') and probe E_Sarbeco_P1 (5'-FAM-ACACTAGCCATCCTTACTGCGCTTCG-BBQ-3') using the TaqPath 1-Step Multiplex Master Mix kit (Applied Biosystems) on a QuantStudio 5 real-time PCR system (Applied Biosystems), as per manufacturer's instructions. A standard curve was generated in parallel for each plate using synthesized DNA and used for the quantification of viral genome copy numbers. The Design and Analysis Software version 1.5.1 (ThermoFisher Scientific) was used to calculate the cycle threshold values, and a cycle threshold value ≤ 36 for both replicates was considered positive.

Blood counts and biochemistry

Complete blood counts were carried out using a VetScan HM5 hematology system (Abaxis Veterinary Diagnostics), as per manufacturer instructions. Analysis of serum biochemistry was performed with a VetScan VS2 analyzer (Abaxis Veterinary Diagnostics), as per manufacturer instructions.

Transcriptional profiling of host responses

RNA was extracted from tissues using the RNeasy plus mini kit (Qiagen) following the manufacturer's instructions. A two-step qRT-PCR reaction in triplicate was performed on a Quantstudio 5 (Applied Biosystems). Reverse transcription was carried out with the Superscript III RT first strand synthesis kit (Invitrogen) following DNase elimination of genomic DNA. The RT reaction was performed using 100 ng of template RNA mixed with random hexamers as primers in a 20 μ L reaction volume for 5 min at 65 °C followed by 10 min at 25 °C and 50 min at 50 °C. The qPCR reaction was carried out using the PowerUp SYBR Green Master Mix (Applied Biosystems) with 2 μ L of cDNA. The cycle parameters for qPCR were 2 min at 50 °C and 2 min at 95°C followed by 40 cycles of 3 s at 95 °C and 30 s at 60 °C. The qPCR reactions were performed in a 20 μ L volume with oligonucleotide pairs specific for deer mouse TNF α , IL6, IL10, IFN α ⁵⁵ at a concentration of 2 μ M. The fold change in gene expression of TNF α , IL6, IL10, IFN α in SARS-CoV-2-infected versus uninfected deer mice were calculated using the $\Delta\Delta$ Ct method with GAPDH as a reference gene⁵⁶. GraphPad Prism's multiple t test was used to perform the second subtraction so as not to lose the variation of the mock animals.

SARS-CoV-2-S-specific enzyme-linked immunosorbent assay (ELISA)

SARS-CoV-2 spike/nucleoprotein (S/N)-specific IgG antibody responses were assessed using an in-house assay. Briefly, a 1:100 dilution (or a serial dilution) of deer mouse serum was carried out in duplicate and added to plates pre-coated with both the spike and nucleoprotein (S/N) antigens in the same assay wells. Deer mouse IgG was detected with a KPL peroxidase-labeled polyclonal goat antibodies against *Peromyscus leucopus* IgG (H+L) (Sera Care).

Virus neutralization assay

Deer mouse serum samples were collected and stored at -80°C. SARS-CoV-2 stocks were titrated and used in the PRNT⁵⁷. Briefly, serum was heat-inactivated at 56°C for 30 minutes and diluted 2-fold from 1:40 to 1:1280 in DMEM supplemented with 2% FBS. Diluted sera was incubated with 50 plaque forming units of SARS-CoV-2 at 37°C and 5% CO₂ for 1 hour. The sera-virus mixtures were added to 24-well plates containing Vero E6 cells at 100% confluence, followed by incubation at 37 °C and 5% CO₂ for 1 hour. After adsorption, 1.5% carboxymethylcellulose diluted in MEM supplemented with 4% FBS, L-glutamine, non-essential amino acids, and sodium bicarbonate was added to each well and plates were incubated at 37°C and 5% CO₂ for 72 hours. The liquid overlay was removed and cells were fixed with 10% neutral-buffered formalin for 1 hour at room temperature. The monolayers were stained with 0.5% crystal violet for 10 minutes and washed with 20% ethanol. Plaques were enumerated and compared to a 90% neutralization control. The PRNT-90 endpoint titre was defined as the highest serum dilution resulting in 90% reduction in plaques. PRNT-90 titers $\geq$ 1:40 were considered positive for neutralizing antibodies.

Histopathology and vRNA in situ hybridization

Tissues were fixed in 10% neutral phosphate buffered formalin for two to four weeks. Subsequently, routine processing was carried out and tissue samples were sectioned at 5 μ m. A set of slides was stained with hematoxylin and eosin for histopathologic examination. RNA in situ hybridization (ISH) was carried out using RNAscope 2.5 HD Detection Reagent-Red (Advanced Cell Diagnostics), according to the manufacturer's instructions. Briefly, formalin-fixed, paraffin-embedded tissue samples of the various tissues were mounted on slides, baked in

a dry oven for 1 hour at 60 °C, and deparaffinized. Tissue sections were then pre-treated with RNAscope H₂O₂ to block endogenous peroxidases for 10 minutes at room temperature. Target retrieval was carried out using the RNAscope Target Retrieval Reagent for 15 minutes. RNAscope Protease Plus Reagent was then applied for 15 minutes at 40 °C. The probes targeting SARS-CoV-2 RNA (V-nCoV2019-S probe, ref#848561) or anti-genomic RNA (V-nCoV2019-S-sense ref#845701) were designed and manufactured by Advanced Cell Diagnostics. The negative probe was also obtained from Advanced Cell Diagnostics (Reference # 310034). The stained tissues were counterstained with Gills I Hematoxylin, and the final images were captured using a light microscope equipped with a digital camera.

SARS-CoV-2 transmission studies

Eight to thirty two-week old female or male deer mice were challenged with 10⁵ or 10⁶ TCID₅₀ of SARS-CoV-2 by an intranasal route (i.n.) of administration in a 50 µl volume. At 24 or 48 hpi (for 10⁶ TCID₅₀ and 10⁵ TCID₅₀, respectively) experimentally infected deer mice were placed into a fresh cage with a naïve cage mate to assess direct contact transmission (donor:contacts at 1:1 ratio). Deer mice were monitored daily for clinical signs and/or weight loss. Oropharyngeal and rectal swab samples were taken on alternate days up to 10 DPI or 14 DPI (for 10⁶ TCID₅₀ and 10⁵ TCID₅₀, respectively), and oropharyngeal and rectal swabs as well as serum samples were collected at 21 dpc.

Bioinformatic analyses

Ace2 sequences were aligned with Clustal Omega⁵⁸. For the phylogenetic analysis, mitochondrial genome sequences (see Accession codes) were aligned with MAFFT v7.467⁵⁹, with regions of poor alignment trimmed with Gblocks v0.91b⁶⁰ resulting in a final alignment of 15,393bp in 115 blocks of minimum 5bp lengths. A maximum likelihood phylogeny was constructed with RAXML v8.2.12⁶¹ using the GTR+I+G4 substitution model as selected by modeltest-ng⁶².

Data analysis

Results were analyzed and graphed using Prism 8 software (Graphpad Software). As appropriate, statistical analyses were performed using ANOVA with multiple comparison correction, the multiple t test, or the unpaired t test with Welch's correction or Mann-Whitney test.

Accession codes

Ace2 sequences: Deer Mouse (*Peromyscus maniculatus bairdii*; XP_006973269), White-footed Mouse (*Peromyscus leucopus*; XP_028743609), Syrian Hamster (*Mesocricetus auratus*; XP_005074266), Chinese Hamster (*Cricetulus griseus*; XP_003503283), Mouse (*Mus Musculus*; ABN80105), Human (*Homo sapiens*; ACT66268), Masked Palm Civet (AAX63775), Fruit Bat (*Rousettus aegyptiacus*; XP_015974412), Chinese rufous horseshoe bat (*Rhinolophus sinicus*; ADN93472.1), Domestic cat (*Felis catus*; AAX59005), Ferret (*Mustela putorius furo*; BAE53380), and Prairie Vole (*Microtus ochrogaster*; XP_005358818). Mitochondrial genomes: *Peromyscus maniculatus* (NC_039921.1), *Peromyscus leucopus* (NC_037180.1), *Mesocricetus auratus* (NC_013276.1), *Cricetulus griseus* (NC_007936.1), *Mus musculus* (NC_005089.1).

Data availability

All relevant data are available from the authors upon request.

661

662 **Acknowledgements**

663 We thank Samira Mubareka of Sunnybrook Research Institute and University of Toronto for
 664 providing the clinical sample and Darryl Falzarano of Vaccine and Infectious Disease
 665 Organization - International Vaccine Centre (VIDO-InterVac) for isolation of the SARS-CoV-2
 666 strain used in these studies. The authors also thank Michelle French, Kimberly Azaransky,
 667 Stephanie Kucas and Christine DeGraff of the Veterinary Technical Services at the National
 668 Microbiology Laboratory (NML) for their technical assistance during the course of this work.
 669 The work described in this manuscript was financially supported by the Public Health Agency of
 670 Canada and the Canadian Food Inspection Agency.

671

672 **Author Contributions**

673 The experiments were conceived of and designed by BDG, DS, and DK. The phylogenetic
 674 analysis was performed by ATD. The Ace2 alignment analysis was carried out by BDG and EM.
 675 The virus was propagated and titered by AL, and DK. Animal husbandry, infection and sample
 676 collection was carried out by BDG, MC, NT, BMW, AA, KT, and DK. Infectious virus assays
 677 were performed by BDG and MC. Sample inactivation and RNA extractions were carried out by
 678 BDG, NT, AL, BMW, LG, KNT, RV, and ABo. Real-time RT-PCR was performed by BDG
 679 and NT. Transcriptional profiling of host responses was carried out by BMW and NT. Blood
 680 counts, biochemistry, and pathology analysis was performed and interpreted by BDG, AB, GP,
 681 JS, CEH, and DK. ELISA was developed and carried out by AL and SH. PRNT assays were
 682 developed and carried out by EM, HW, and MD. Histopathology and ISH was carried out by
 683 EMO and pathology was assessed by CEH. Statistical analysis was performed by JA. Figures
 684 were prepared by BDG, EM, and LB. Interpretation of potential impacts on wild deer mice and
 685 deer mouse ecology was done by LRL and CMJ. The initial draft of the manuscript was written
 686 by BDG and DK, with all other authors contributing to editing into the manuscript into its final
 687 form. The work was managed and supervised by BDG, MC, LF, CEH, and DK.

688

689 **Conflict of Interest**

690 The authors declare no conflicts of interest.

691 **Extended Data**

692 **Extended Data Figure 1.** Viral burden in the tissues by age.

693 **Extended Data Figure 2.** Histological analysis of lungs at 21 days following SARS-CoV-2
 694 exposure.

695 **Extended Data Figure 3.** Serum biochemical values.

696 **Extended Data Figure 4.** Schematic of deer mouse SARS-CoV-2 transmission studies 1 and 2
 697 design.

698

a

b

c

Species	ACE2, SARS-CoV-2 spike contacting residues																			Congruence with hAce2, spike contacting residues	
																				AA identity	AA polarity
Human (hAce2)	Q24	T27	K31	H34	E35	E37	D38	Y41	Q42	L45	L79	M82	Y83	N90	Q325	E329	N330	K393	G394	-	-
Syrian hamster	Q	T	K	Q	E	E	D	Y	Q	L	L	N	Y	N	Q	E	N	K	G	89.5% (17/19)	89.5% (17/19)
Chinese hamster	Q	T	K	Q	E	E	D	Y	Q	L	L	N	Y	N	Q	G	N	K	G	84.2% (16/19)	84.2% (16/19)
Cat	L	T	K	H	E	E	E	Y	Q	L	L	T	Y	N	Q	E	N	K	G	84.2% (16/19)	89.5% (17/19)
White-footed mouse	Q	I	K	Q	E	E	D	Y	Q	L	L	N	Y	N	Q	E	N	K	G	84.2% (16/19)	84.2% (16/19)
Deer mouse	Q	I	K	Q	E	E	D	Y	Q	L	L	N	Y	N	Q	R	N	K	G	78.9% (15/19)	78.9% (15/19)
Prairie Vole	D	A	K	Q	E	E	D	Y	Q	L	L	S	Y	N	Q	E	N	K	D	73.7% (14/19)	73.7% (14/19)
Fruit bat	L	T	K	T	E	E	D	Y	Q	L	L	T	Y	D	E	E	K	K	G	68.4% (13/19)	68.4% (13/19)
Chinese rufous horseshoe bat	E	I	K	T	K	E	D	H	Q	L	L	N	Y	N	E	N	N	K	G	57.9% (11/19)	63.2% (12/19)
Masked Palm Civet	L	T	T	Y	E	Q	E	Y	Q	V	L	T	Y	D	Q	E	N	K	G	57.9% (11/19)	68.4% (13/19)
Ferret	L	T	K	Y	E	E	E	Y	Q	L	H	T	Y	D	E	Q	N	K	R	52.6% (10/19)	57.9% (11/19)
Mouse	N	T	N	Q	E	E	D	Y	Q	L	T	S	F	T	Q	A	N	H	G	52.6% (10/19)	68.4% (13/19)

Figure 1: Deer mouse range and predicted susceptibility to SARS-CoV-2. a, Phylogeny showing the relationships among selected members of the *Cricetidae* family with mice and humans. b, Geographical distribution of deer mice (based on data from Hall, 1981)³². c, Alignment of human Ace2 (hAce2) amino acid residues known to confer efficient binding of the RBD of SARS-CoV-2 spike with the corresponding Ace2 amino acid residues from selected members of the *Cricetidae* family and other naturally or experimentally susceptible host species. The physiochemical properties of the amino acids residues are indicated: non-polar (yellow), polar (green), acidic (red), and basic residues (blue).

Figure 2: SARS-CoV-2 infection of adult deer mice. a-m Eight to thirty-two week old female or male deer mice (*P. maniculatus*) were inoculated with 10^5 TCID_{50} or 10^6 TCID_{50} of SARS-CoV-2 by an intranasal route (i.n.) of administration and compared to age-matched uninfected controls. Bars indicate means. Dashed lines and dotted lines indicate the limit of detection for the TCID_{50} assay and qRT-PCR assay, respectively. a-b, Kaplan-Meier curve depicting survival data (a) and weight data (b) over the course of 21 days following SARS-CoV-2 exposure (challenge dose, 10^5 TCID_{50}). c-f, Infectious viral load (filled in circles, left axis) and vRNA levels (empty circles, right axis) in the (c) nasal turbinates, (d) lung, (e) small intestine, and (f) colon (challenge dose, 10^5 TCID_{50}). g-h, Infectious viral load (filled in circles with bars, left axis) and viral RNA levels (empty circles, right axis) in (g) oropharyngeal swabs and (h) rectal swab samples (challenge dose, 10^5 TCID_{50}). i, Viral RNA levels in nasal washes at 6 dpi (infectious dose, 10^5 TCID_{50}). j-l, Viral RNA levels in the (j) urine, (k) feces, and (l) blood at the indicated times post-infection (infectious dose, 10^5 TCID_{50}). m, Viral RNA levels in the nasal turbinates, lung, and small intestines at 21 days post-infection (infectious dose, 10^5 TCID_{50}). Data were collected from two independent experiments. (a-b, n = 6; c-h, n = 3/6 for uninfected/infectious; i-l, n = 5; m, n = 4).

Figure 3: Histopathology and virus distribution. Hematoxylin/eosin (H&E) staining (left) and in situ hybridization (ISH) using anti-sense probes that detect the SARS-CoV-2 genome/mRNA (middle) and sense probes that detect anti-genomic RNA (right) were carried out on **a**) nasal turbinates and **b**) lung tissue of uninfected and SARS-CoV-2-infected deer mice (1×10^5 TCID₅₀ i.n. route) at 2 and 4 dpi. Positive detection of viral genomic RNA/mRNA or anti-genomic RNA is indicated by magenta staining (middle/right panels and insets). Arrows indicate perivascular infiltrations of histiocytes and neutrophils (black), peribroncholar infiltrations of histiocytes and neutrophils (orange), mild neutrophilic infiltration in the submucosa (blue), occasionally observed discrete foci of interstitial pneumonia (purple), occasionally observed multinucleated syncytial cells (magenta), and anti-genomic RNA occasionally in individual scattered bronchiolar epithelial cells (green and right inset). The magnification is 20x for H&E and 20x for ISH unless otherwise indicated. Scale bars = 100 μ m, 50 μ m, and 20 μ m for 10x, 20x, and 40x, respectively.

Figure 4: Deer mouse host response to SARS-CoV-2 infection. **a-e**, Hematological levels and serum biochemistry were measured in uninfected and SARS-CoV-2-infected deer mice (infectious dose 10^6 TCID₅₀, i.n. route) at 3 dpi, including **(a)** white blood cell, lymphocyte, and neutrophil counts, **(b)** the neutrophil-to-lymphocyte ratio, **(c)** alanine aminotransferase (ALT), **(d)** blood albumin (ALB), and **(e)** blood urea nitrogen (BUN). **f**, Cytokine gene expression was measured for TNF α , IL-6, IL-10, and IFN α in the lungs of SARS-CoV-2-infected deer mice and displayed relative to age-matched mock-infected animals. Gene expression was normalized using GAPDH as a control. **g**, IgG antibody response against SARS-CoV-2 mixed spike and nucleoprotein (S/N) antigens were assessed by ELISA using serum collected on the indicated days post-infection (infectious dose, 10^5 TCID₅₀). **h**, Neutralizing antibody against SARS-CoV-2 was measured by PRNT₉₀ using serum collected at 28 days after SARS-CoV-2 infection (infectious dose, 10^5 TCID₅₀). Dotted lines indicate the limit of detection. Bars indicate mean, error bars indicate SEM (c-e) or 95% confidence interval (f). (a-b, n = 5/3 for uninfected/infected; c-e, n = 5/4 for uninfected/infected, f, n = 6, g-h, n = 5 or 6) * = $P < 0.05$, ns = $P > 0.05$, Mann-Whitney test (b), unpaired t test (c-e).

Figure 5: Transmission of SARS-CoV-2 between deer mice through direct contact. a-g. Adult male and female deer mice were exposed to 10^6 TCID₅₀ SARS-CoV-2 by an i.n. route of infection. At 1 dpi individual inoculated donor deer mice were transferred to a new cage and co-housed with a single naïve deer mouse (1:1 ratio) to assess SARS-CoV-2 transmission by direct contact. Deer mice were either maintained in direct contact throughout the 10 day study for serial swabbing or humanely euthanized on 2 or 4 days post-contact (dpc) for tissue collection. a-d. Viral RNA levels (a, b) and infectious viral loads (c, d) were measured in oropharyngeal swabs and rectal swab samples every other day from -1 to 10 days after direct contact was initiated. Oropharyngeal or rectal swab samples obtained from the same animal are demarcated with a unique colour, and samples derived from co-housed pairs share the same symbol. e-f. An additional transmission study was carried out in the same manner and nasal wash, nasal turbinates, and lung samples were collected from contact animals at (e) 2 dpc and (f) 4 dpc. g. IgG antibody response (reciprocal serum dilution) against SARS-CoV-2 S and N in donor and direct-contact exposed animals were assessed by ELISA using serum collected at 21 dpc. h-i. In an additional study adult male and female deer mice were exposed to 10^6 TCID₅₀ SARS-CoV-2 by an intranasal route of infection. At 2 dpi individual inoculated donor deer mice were transferred to a new cage and co-housed with a single naïve deer mouse (1:1 ratio) to assess SARS-CoV-2 transmission by direct contact (n = 5). Viral RNA levels were measured in (h) oropharyngeal swabs and (i) rectal swab samples every other day from 0 to 14 days after direct contact was initiated, and final swab samples were collected at 21 dpi and assessed. Dotted or dashed lines indicate the limit of detection. Bars indicate means. Data were collected from two independent experiments.