Light might suppress both types of sound-evoked anti-predator flight in moths

Theresa Hügel*, Holger R. Goerlitz*

Acoustic and Functional Ecology Group, Max Planck Institute for Ornithology, Seewiesen,
Germany

***corresponding authors:**

Acoustic and Functional Ecology Group

Max Planck Institute for Ornithology

Eberhard-Gwinner-Str. 11

82319 Seewiesen, Germany

Email: thuegel@orn.mpg.de, hgoerlitz@orn.mpg.de

ORCID:

Theresa Hügel: 0000-0003-0500-1988

Holger Goerlitz: 0000-0002-9677-8073

ABSTRACT

Urbanization exposes wild animals to increased levels of light, affecting particularly nocturnal animals. Artificial light at night might shift the balance of predator-prey interactions, for example of nocturnal echolocating bats and eared moths. Moths exposed to light show less last-ditch manoeuvres in response to attacking close-by bats. In contrast, the extent to which negative phonotaxis, moths' first line of defence against distant bats, is affected by light is unclear. Here, we aimed to quantify the overall effect of light on both types of sound-evoked anti-predator flight, last-ditch manoeuvres and negative phonotaxis. We caught moths at two light traps, which were alternately equipped with speakers that presented ultrasonic playbacks to simulate hunting bats. While the light field was omnidirectional, attracting moths equally from all directions, the sound field was directional, thus eliciting only negative phonotaxis, or negative phonotaxis and last-ditch manoeuvres, depending on approach direction. We could not detect an effect of sound playback on number of caught moths, suggesting that light might suppress both types anti-predator flight. In contrast, our playback elicited evasive flight in moths released in a dark flight room. Showing no effect of a treatment, however, is difficult. We discuss potential alternative explanations for our results, and call for further studies to investigate how light interferes with animal behaviour.

KEY WORDS

ALAN, Chiroptera, Noctuidae, Lepidoptera, playback experiment

INTRODUCTION

Light pollution by artificial light at night has increased substantially over the last decades (Falchi et al., 2016) (Fouquet, 2006; Hölker et al., 2010), adversely affecting plants and animals (Davies and Smyth, 2018; Knop et al., 2017; Longcore and Rich, 2004). The effects of light range from single individual's orientation, reproduction and communication (Longcore and Rich, 2004) to whole communities, for example, by shifting the balance of predator-prey interactions (Bailey et al., 2019; Davies et al., 2013, 2014; Miller Colleen R. et al., 2017; Russo et al., 2019; Yurk and Trites, 2000). Echolocating bats and eared moths constitute a globally occurring predator-prey system of high ecological relevance (Boyles et al., 2011; Kasso and Balakrishnan, 2013; Kunz et al., 2011; Van Toor et al., 2019). Their interactions take place in the darkness of the night and are exclusively mediated by sound. Echolocating bats hunt by emitting ultrasonic calls (Fenton, 2003; Schnitzler et al., 2003), which eared moths can hear and react to with evasive flight (Roeder, 1962; Ter Hofstede and Ratcliffe, 2016). Evasive flight likely consists of two stages: negative phonotaxis to fly away from distant bats (stage 1), and last-ditch evasive manoeuvres such as erratic flight or (power) dives to escape nearby attacking bats. Corresponding to the differences in bat distance, negative phonotaxis is elicited at received sound pressure levels that are about 20 dB fainter than those that elicit last-ditch manoeuvres (Agee, 1969; Roeder, 1962, 1964, 1967).

Artificial night at light is of increasing concern for both bats and moths. While some bats may profit from exploiting prey accumulated at lights (Cravens et al., 2018; Rydell, 1992), other species are negatively affected while foraging, commuting and roosting (Mathews et al., 2015; Stone et al., 2009, 2015; Straka et al., 2016, 2020). Moths are strongly attracted to light sources, causing reduced time spent foraging (Macgregor et al., 2019; van Langevelde Frank et al., 2017), disrupted navigation (Owens and Lewis, 2018), and in turn

reduced pollination (Macgregor et al., 2017) and population decline (van Langevelde Frank et al., 2017; Wilson et al., 2018). Furthermore, light increases the predation risk of moths, for two reasons. The accumulations of moths around light sources attract bats, thereby increasing the predation pressure on moths (Cravens et al., 2018; Rydell, 1992). In addition, light interferes with the moths' sound-evoked anti-predator evasive flight response. In one set of studies, the sound-evoked evasive flight of moths was compared under lit and unlit conditions, showing that light reduces the evasive flight. Wakefield et al. (Wakefield et al., 2015) showed that only 24% of moths performed last-ditch power-dives under LED illumination compared to 60% of moths in darkness; that is, the light inhibited last-ditch manoeuvres in 60% of the moths that would react in darkness. Similarly, Svensson & Rydell (Svensson and Rydell, 1998) reported that moths within a radius of 1 m around a light source showed ~60% less last-ditch manoeuvres than moths in darkness (where 100% of moths reacted). Minnaar et al. (Minnaar et al., 2015) reported the most extreme effect: in darkness, bat diet (as proxy for moth capture rate) was best explained by a model that included evasive flight of moths. In contrast, with light, bat diet was best explained by a model that included a 100% reduction in moth evasive flight, suggesting that the light inhibited both stages of evasive flight (negative phonotaxis and last-ditch manoeuvres).

In another set of studies, light exposure was kept constant while the sound received by the moth was manipulated. Those results showed that moths still exhibited some degree of evasive flight under illumination. Acharya & Fenton (Acharya and Fenton, 1999) compared last-ditch manoeuvres in eared and deafened moths under illumination, showing that 48% of eared moths exhibited last-ditch manoeuvres when preyed on by bats, whereas deafened moths did not. Agee & Webb (Agee and Webb, 1969) and Treat (Treat, 1962) compared the number of caught moths at light traps with and without ultrasonic stimuli. Depending on

sound stimulus and moth species, ultrasound playback reduced captures by 51-86 % (in *Heliothis zea*, Noctuidae, (Agee and Webb, 1969) and 8-49 % (nine tympanate moth families with a at least 39 caught individuals (Treat, 1962)) of the captures at the silent trap.

In summary, the first set of studies shows that light suppresses the sound-triggered evasive flight in 60-100% of the moths that would react in darkness. Contrasting this, the second set of studies shows that even in the light, sound can still trigger evasive flight in 8-86% of the moths. Noteworthy, these studies either only reported effects of light on last-ditch manoeuvres (Svensson and Rydell, 1998; Wakefield et al., 2015), or the results can be sufficiently explained by effects of light on last-ditch manoeuvres, as all moths had to fly through fields of high sound pressure level before entering the light trap (Treat, 1962) (Agee and Webb, 1969). Only the modelling results of Minnaar et al. (Minnaar et al., 2015) suggest that light completely suppresses both evasive flight responses under natural conditions. Therefore, while several lines of evidence suggest that last-ditch manoeuvres are suppressed by light pollution (with variable effect sizes), we lack a similar understanding of the effect of light on negative phonotaxis. If the light-induced suppression of negative phonotaxis is as strong as for last-ditch manoeuvres, this will strongly affect the predator-prey interactions between bats and moths, because negative phonotaxis is elicited over larger distances and larger spatial volumes than last-ditch manoeuvres. Here, we advanced the light-trap approach of Agee & Webb and Treat (Agee and Webb, 1969; Treat, 1962) to investigate the effects of light on both stages of evasive flight, negative phonotaxis and last-ditch manoeuvres. We combined the omnidirectional light field of light traps (attracting moths equally from all directions) with a directional ultrasonic playback that should elicit different stages of evasive flight depending on approach direction. The received sound pressure level will increase as moths approach the light trap, independent of approach direction. However,

depending on their approach direction, received sound pressure levels will reach different maximum levels, sufficient to elicit either only negative phonotaxis (off-axis approach), or negative phonotaxis and last-ditch manoeuvres (on-axis approach). We compared moth captures at the light traps with and without acoustic playback, to measure the effect of light on the sound-evoked evasive flight. Hypothesizing in line with Minaar et al. (Minaar et al., 2015) that light suppresses both stages of evasive flight, we predicted equal moth counts at both light traps. In contrast, if negative phonotaxis was not suppressed or anti-predator flight was not completely suppressed, we predicted lower moth counts at the ultrasonic than the silent trap.

METHODS

Setup, study site, moth capture and measurement

We compared the number of moths caught at two light traps, one of which was equipped with a loudspeaker. We set up two equal light traps (Sylvania, blacklight, F15W/350BL-T8, USA, Fig. 1A); one next to a path in a forest (trap A) and the other one at 30 m distance in the forest (trap B), close to the Max Planck Institute for Ornithology, Seewiesen, Germany. Both traps hung freely at ~1.7 m above ground, radiating light at 360° in the horizontal plane (Fig. 1A). Between July 19 and August 16 2018, we collected data over 15 rainless nights, based on the number of nights sampled in similar previous studies (6 nights (Agee and Webb, 1969) and 2-12 nights (Treat, 1962), respectively). Each test night, we alternatingly equipped one of the two traps with two loudspeakers (Ultrasonic Dynamic Speaker ScanSpeak, Avisoft Bioacoustics, Glienicke, Germany); both broadcasting an ultrasonic

Figure 1: (A) Experimental setup. The photo shows one of the light traps with two loudspeakers attached above the trap, both of which broadcast the acoustic stimulus in opposite directions. (B) Side view of the biologically relevant sound field around the light trap with attached loudspeakers. Coloured areas indicate areas with minimum sound pressure levels of 60 and 80 dB SPL re. 20 μ Pa, which are biologically relevant acoustic thresholds for eliciting negative phonotaxis and last-ditch manoeuvres, respectively, in eared moths. Schematic indicates position and dimension of the light trap. (C) Top view of the biologically relevant sound field (coloured areas) and light field (dashed lines). Dashed lines indicate the distance over which 5% of released noctuid (10 m) and geometrid (23 m) moths are recaptured at the light trap, respectively (Merckx and Slade, 2014), as an indicator of the maximum range of light attraction.

stimulus (see below) to simulate echolocating bats. The two loudspeakers were fixed back to back facing in opposite directions and were mounted above the respective light trap at ~2 m above ground. Thus, each trap was associated for 7-8 nights with ultrasound simulating an echolocating bat. Each test night, lights and acoustic playback were turned on in the evening (between 20:30h and 23:40h) and turned off the next morning (between 7:00h and 9:40h). We emptied the traps each morning and counted all moths of the three ear-possessing families Noctuidae, Geometridae and Erebidae. We measured each individuals' body length along the middorsal line (from the head to the end of the abdomen), to correct for the fact that larger moths have more sensitive hearing than smaller moths (Surlykke et al., 1999; Ter Hofstede et al., 2013). For those individuals whose body length could not be measured (e.g., due to missing abdomen, N = 137, 15.5%), we either used the mean value of the species or, if this was not possible (N = 1), the mean value of the family. For statistical analysis, we binned individuals into six categories of body length (1.0 to 3.0 cm, bin width 0.5 cm).

Ultrasonic playback stimulus design and evaluation

We simulated predation pressure by echolocating bats by repeatedly playing a short bat-like ultrasonic pure tone pulse. Pulse frequency was 35 kHz, matching most moths' best hearing threshold around 20-50 kHz (Noctuidae: (Nakano et al., 2015; Ter Hofstede et al., 2013); Erebidae: (ter Hofstede et al., 2008); Pyralidae: (Skals and Surlykke, 2000); Geometridae: (Rydell et al., 1997; Surlykke et al., 1997); Sphingidae/Drepanidae: (Nakano et al., 2015)). Pulse duration was 10 ms including 2 ms linear rise and fall times, corresponding to the calls of European open space bats (Obrist et al., 2004; Skiba, 2014) and optimizing information transfer to the moths (Gordon and Ter Hofstede, 2018). The pulse interval was 100 ms, matching the call interval of searching bats (60 - 200 ms, (Holderied and von Helversen,

2003; Skiba, 2014)). On-axis sound pressure level (SPL) was 98 dB SPL re. 20 μ Pa RMS at 1 m distance (see below for a detailed description of the sound field). This stimulus was presented continuously in a loop throughout the night via the loudspeakers using Avisoft-RECORDER software (Avisoft Bioacoustics), a sound card with amplifier (Avisoft UltraSoundGate 116, Avisoft Bioacoustics) and a laptop computer.

To test the effect of our acoustic stimulus in darkness, without the potentially suppressing influence of light, we exposed free-flying moths in a dark flight room (5.3 m x 3.5 x 3 m³) to the same acoustic stimulus. We caught moths at trap A over the course of four nights and tested them within 30 hours after capture. We placed moths on the ground of the flight room and recorded the flight paths of upward-flying moths with an IR-sensitive camera (Sony HDR-CX560, Sony, Tokio, Japan) under IR illumination (850 nm, Mini IR Illuminator TV6700, ABUS Security-Center, Wetter, Germany). Using the same audio equipment as described above, we manually started the sound presentation when a moth flew in front of the speaker. We subsequently categorized the video-recorded flight paths as “reaction” when the flight direction, level of erraticness or both changed with stimulus onset (for examples, see supplementary video); as “no reaction” when we did not observe those changes; or as “ambiguous” when we could not clearly assign the flight path to one of the previous categories.

Overlap of sound and light field

The range and geometry of the presented light and sound fields differed. While the light was emitted omnidirectionally in the horizontal plane, the sound field was directional (Fig. 1). We estimated the biologically relevant range for attracting moths by light and the biologically

relevant sound fields for triggering moths' evasive flight based on literature values and our own measurements.

Light traps can attract released moths over up to 80 m distance, yet recaptures dramatically decrease beyond 15 m and depend on family (Merckx and Slade, 2014; Truxa and Fiedler, 2012). Family-specific models estimated the 5%-recapture rate at a distance of 10 ± 6 m (mean $\pm$ SEM) for Noctuidae and 23 ± 12 m for Geometridae (Merckx and Slade, 2014). Note, however, that these distances were obtained with a different light source than ours (6W actinic vs. 15W blacklight in our case), and that the distance over which light attracts moths must not be equivalent to the distance over which light interferes with evasive flight. We still used these family-specific distances as first approximation for the biologically relevant light fields, which attract moths to our light traps (Fig. 1B, dashed lines).

Figure 2: Number of caught moths per night and per playback treatment. (A) Number of caught moths at both light traps, with and without playback. Boxplots show median, quartiles and whiskers (up to 1.5 x inter-quartile range) of the daily counts (dots). (B) Daily counts of caught moths per night and per trap for all 15 nights. White letters at the base of the bars indicate the trap.

To estimate the effect of the playback, we measured the sound pressure level (SPL) of the pulse playback in front of the loudspeaker (on-axis) in steps of 5° up to 90° off-axis (for details, see SI). The on-axis source level at 1 ms distance was 97 dB SPL re. 20 µPa RMS, i.e., within the lower range of call levels emitted by free-flying bats (100-120 dB peSPL @ 1 m (Holderied and von Helversen, 2003), with the corresponding RMS-SPL-levels being ~3-7 dB lower than the peSPL-levels (Lewanzik and Goerlitz, 2017; Seibert et al., 2015)). With increasing off-axis angle, the source level became fainter by up to ~30 dB at 45°, resulting in a minimum playback level of 70 dB SPL RMS @ 1m. We then calculated the angle-dependent distances around the loudspeaker where the playback would reach biologically relevant levels of 60 and 80 dB SPL RMS. We chose 60 and 80 dB SPL as approximate thresholds for eliciting negative phonotaxis and last-ditch manoeuvres, respectively, based on several lines of evidence. Negative phonotaxis and last-ditch manoeuvres are likely triggered at levels just above the thresholds of the moths' auditory receptor neurons A1 and A2, respectively (Gordon and Ter Hofstede, 2018; Madsen and Miller, 1987; Roeder, 1974). The best thresholds of the A1 neuron are at about 35-55 dB SPL, and of the A2 neuron at about ~52-72 dB SPL (Gordon and Ter Hofstede, 2018; Madsen and Miller, 1987; Surlykke, 2003; Ter Hofstede and Ratcliffe, 2016; Ter Hofstede et al., 2013; Waters and Jones, 1996). Behavioural thresholds in moths are largely unknown, but those that are known tend to be about 10 dB higher than neuronal thresholds (reviewed in (Lewanzik and Goerlitz, 2017)). We thus defined 60 and 80 dB SPL RMS as thresholds that will likely elicit negative phonotaxis and last-ditch manoeuvres in most moth species, respectively, and calculated their isolines of constant sound pressure levels. SPL-Isolines varied with the angle around the loudspeaker, ranging from 3.6 to 14.6 m for 60 dB SPL RMS, and from 0 to 5.6 m for 80 dB SPL RMS (Fig. 1B). In summary, we thus presented an highly directional sound field in an

omnidirectional light field. Thus, moths that were attracted to the light trap in the on-axis direction of the loudspeaker experienced gradually increasing SPLs sufficiently high to first elicit negative phonotaxis and later last-ditch manoeuvres. In contrast, moths that were attracted from the side (off-axis to the loudspeaker) experienced gradually increasing SPLs that were only sufficiently high to elicit negative phonotaxis, but not last-ditch manoeuvres.

Statistical analysis

To test for an effect of light on the moths' evasive flight, we fitted linear models to the logarithmized moth count data as a function of the fixed effects *playback*, *trap*, *moth family*, and *moth body length*, and the interactions of *playback* and *moth family*, and of *playback* and *moth body length*.

We preferred a linear model with logarithmized count data over a negative binomial model as the linear model enabled us to do detailed power analysis, even though both model types could be fitted similarly well. Although we have repeated measures over 15 nights, we did not include date as a random factor as it only explained a minor proportion of the variance in the data.

We defined the moth family Noctuidae as intercept, as it had the largest sample size and thus was the most reliable reference. To test which factors significantly contributed to the model fit, we conducted backwards model reduction (Lewis et al., 2011). Hence, the full model was successively reduced, by stepwise removing factors, starting with the factor having the highest p-value of the t-statistics provided by the model summary. We compared models with likelihood ratio tests using a F-statistic and AICs. None of the interaction terms nor *trap* contributed significantly to model fit and were thus excluded. *Moth family* and *moth body length* contributed significantly to model fit (see results). We also kept *playback*

as factor in the final model, as we aimed to analyse the effect of playback on the number of caught moths. Our final model thus included *playback*, *moth family* and *moth body length* as fixed factors, without any interactions.

We evaluated the power of our model for the effect sizes found in the field and in the flight room by a randomization approach of our real dataset. We first randomized the factor *playback* and then added an effect size as determined in the flight room or in the field to those logarithmized moth counts where the *playback* was “on”. We ran this simulation 10000 times for both effect sizes, and each time compared models (final model vs. final model without *playback*) with likelihood ratio tests using a F-statistic to test for a significant effect of *playback* on the model fit. The proportion of significant effects of *playback* per 10000 simulations equals our power to detect an effect of the tested effect size.

All statistical analyses were conducted in R version 3.3.2 (R Foundation for Statistical Computing, Vienna, Austria) using the packages “lme4” (Bates et al., 2015), “MASS” (Venables and Ripley, 2002), “blme4” (Korner-Nievergelt et al., 2015), “DHARMa” (Hartig, 2019), “car” (Fox, 2019), “effectsize” (Kassambara, 2019), and RNOmni” (McCaw, 2018) for statistics and the packages “ggplot2” (Wickham, 2020), “dplyr” (Wickham, 2018), “cowplot” (Wilke, 2019), and “ggthemes” (Arnold, 2018) for data sorting and plotting. For further details, see R-script included in SI.

RESULTS

Of 33 moths tested in free-flight in the dark flight room, ten moths (30.3%) showed an evasive reaction in response to the acoustic stimulus. Twelve moths (36.4%) did not react, and eleven moths (33.3%) were categorized as “ambiguous”. The moths’ distance to the

speaker at stimulus onset was about 1-2 m. Our acoustic stimulus thus was audible and elicited an evasive reaction in about one third of the tested moths. When considering the ambiguous reactions, the stimulus might even be audible to a larger proportion of up to two thirds of the moth population.

In the field experiment, we caught a total of 878 moths over 15 nights, with a median of ~23 moths caught per night and light trap (Fig. 2A), yet with large fluctuations between nights and smaller fluctuations between traps (7-116 moths per night and trap; Fig. 2B). We mostly caught moths of the family Noctuida (80.9 %), followed by Geometridae (11.1%) and Erebidae (7.8%), and more smaller than larger moths. Accordingly, both *moth family* and *moth body length* contributed significantly to our final model after stepwise model reduction (likelihood ratio tests: *moth family*, $F(153,2) = 31.92$, $p < 0.001$; *moth body length*, $F_{(153,3)} = 34.43$, $p < 0.001$).

Although the total number of moths caught at trap A (506 moths) was 41% higher than the total number caught at trap B (365 moths), this was largely driven by one night (night 10: 115 vs 29 moths; Fig. 2B). Across all 15 nights, the nightly capture rate did not significantly differ between traps (likelihood ratio test for factor *trap*: $F_{(151,1)} = 0.53$, $p = 0.468$; Fig.2A).

We could not detect an effect of our acoustic playback on the nightly capture rates (Fig. 2A; modelled effect size of factor *playback* on logarithmized moth at the playback trap relative to the silent trap: -0.06 (95% CI: -0.16 – + 0.04), corresponding to 86% (95% CI: 69 – 109%) capture rate at the playback trap relative to the silent trap; likelihood ratio test of factor *playback*, $F_{(152,1)} = 1.53$, $p = 0.218$).

We analysed the power of our experiment, both for the effect size observed in the field (-0.06) and the flight room. In the flight room, at least 30% of the moths reacted to our

acoustic playback. Assuming that those reacting moths would not be caught in the light trap, the capture rate at the ultrasonic trap would be 70% relative to the silent trap, resulting in an expected effect size of the playback of -0.16 (in logarithmized moth counts). The power is the probability of rejecting the null hypothesis (i.e., obtaining a statistically significant result at a chosen significance level), given that the null hypothesis is false. Based on our dataset and sample size, our field experiment had a very low power of only 21% to detect an effect as small as the one observed in the field. In contrast, we had a sufficiently high power of 87% to detect an effect as large as the one observed in the flight room. This suggests that the lack of a significant effect of *playback* in the field might have been caused by a light-induced decrease of the effect size compared to the dark flight room – i.e., a light-induced suppression of sound-evoked anti-predator flight.

DISCUSSION

Echolocating bats and eared insects are a textbook example of sound-based predator-prey interactions (Ter Hofstede and Ratcliffe, 2016). Increasing light pollution (Fouquet, 2006; Hölker et al., 2010), however, severely impacts both bats and moths (e.g. (Cravens et al., 2018; Macgregor et al., 2019; Stone et al., 2009), with potential cascading effects on their predator-prey-interactions, population dynamics and ecosystems (Minnaar et al., 2015; Russo et al., 2019). While good evidence exists that light reduces the sound-evoked last-ditch manoeuvres of eared moths, making them more vulnerable to nearby bats, the effect of light on the moths' first line of defence, negative phonotaxis in response to distant bats, is unclear. Here, we compared moth captures at two light traps. One trap was silent, while the other trap broadcast bat-like ultrasonic stimuli to trigger last-ditch manoeuvres and negative

phonotaxis under dark conditions. We did not find a significant reduction in the number of caught moths at the ultrasonic light trap. There is however a high bar for showing no effect of a treatment (such as the ultrasonic playback in our case). The power of our experiment was too low to detect significant changes in moth count as small as we observed in the field. In contrast, our field experiment had sufficient power to detect an effect of the playback as large as the one observed in the dark flight room, if it had also occurred under lit field conditions. One conclusion thus is that the light suppressed both types of the moths' sound-evoked anti-predator flight, negative phonotaxis and last-ditch manoeuvres.

There are, however, alternative explanations in addition to a light-induced suppression of anti-predator flight. The field and flight room experiments differed not only in the light level, but also in temporal (full night vs. short-term sound exposure) and spatial (variable distances between moth and loudspeaker vs close-range to the loudspeaker) parameters. In addition, physiological and behavioural states of the moths will likely differ between free-flying moths in the field and captured and released moths in the flight room.

The continuous ultrasonic playback over a full night might cause the moths to habituate to the playback. Habituation was previously suggested as an explanation for playback-independent capture rates of male *Helicoverpa zea* moths at pheromone traps (Gillam et al 2011). However, although the playback was on throughout the night, the exposure of individual moths to the playback will likely be shorter and depends on their spatio-temporal movement patterns. Indeed, the studies of Treat (Treat, 1962) and Agee & Webb (Agee and Webb, 1969), which we based our study on, also presented ultrasonic playbacks throughout the night, and were able to detect differences between the silent and ultrasonic trap, arguing against habituation. We also believe that differences in stimulus design cannot explain the differences between experiments. Treat (Treat, 1962) and

Agee & Webb (Agee and Webb, 1969) broadcast multiple stimuli varying in pulse rate, frequency, duration, and sound pressure level (ranges: 0.7-155 pulses/s; 12.5-200 kHz; 2-10 ms; SPL: ~60-100 dB SPL @ 1 m distance), all of which elicited varying degrees of evasive flight in eared moths. The acoustic properties of our stimulus was within this range and did elicit evasive flight under dark control conditions, but seems not to elicit evasive flight under lit conditions.

When assuming that light indeed suppressed the moths' sound-evoked antipredator-flight in our experiment, the question arises why this was not the case in the similar light-trap experiments by Treat (Treat, 1962) and Agee & Webb (Agee and Webb, 1969). We propose that the differences in the geometry and overlap of light- and sound-fields might explain these contrasting results. In the previous setups ((Treat, 1962), (Agee and Webb, 1969)), sound and light fields almost overlapped and were emitted within a relatively narrow angle. Before entering the light trap, approaching moths thus passed through high sound pressure levels that would likely elicit last-ditch manoeuvres. As both studies caught fewer moths in the ultrasonic trap than the silent traps, this suggests that the playback still elicited some anti-predator flight (likely last-ditch manoeuvres) despite the light. Specifically, for stimuli similar to ours, the relative capture rates between the ultrasonic and silent traps were 35% vs. 65% for noctuid moths and a 37.5 kHz tone pulses (Treat, 1962), and 15% vs. 85% for two species of the families Noctuidae and Pyralidae and a 30 kHz tone pulses (Agee and Webb, 1969). In contrast, our setup combined an omnidirectional light field with a directional sound field, thus exposing the moths to different sound pressure levels (SPL) depending on approach direction. When approaching the trap on-axis of the loudspeaker's main axis, received SPLs ranged from low to high, which should first elicit negative phonotaxis and later last-ditch manoeuvres. In contrast, when approaching the traps off-

axis, SPLs remained so low to only elicit negative phonotaxis. As our capture rates did not differ among the light traps – indicating that the playback evoke anti-predator flight – we suggest that the light strongly reduced the negative phonotaxis of eared moths. Even if the moths still exhibited some degree of last-ditch manoeuvre close to the trap (as shown by Treat and Agee & Webb, (Agee and Webb, 1969; Treat, 1962), these manoeuvres might have brought the moth into a position off-axis to the loudspeaker (either to the side or below the main axis) into a region of low SPL. From there, no further last-ditch manoeuvres would have been elicited due to the low SPL, while the light kept attracting the moth into the trap and suppressed negative phonotaxis. Our results are in line with those of Minnaar et al. (Minnaar et al., 2015), who found that a model of escape behaviour in moths assuming 0% efficiency best explained bats’ diet under lit conditions. We therefore suggest that light suppresses not only last-ditch manoeuvres, as previously shown, but also negative phonotaxis.

In summary, our results underline the strong effect of light on eared moths, and suggest that both types of anti-predator flight are suppressed by light. It is important to note, though, that our study design tested the effect of light only indirectly, by testing the effect of sound on light-mediated moth captures, not the effect of light on sound-mediated evasive flight. In addition, showing no effect is difficult, and is complicated by natural variation in capture rates between nights, and the variability of moth behaviour (Hügel & Goerlitz 2019), potentially complicating simple answers. If increasing artificial light at night suppresses both negative phonotaxis and last-ditch manoeuvres, moths are not only unable to escape nearby predators, but also unable to avoid distant predators by flying away. Similarly, fish are attracted to lit areas, where they are “trapped” and preyed upon by seals (Yurk and Trites, 2000) and other fish (Becker et al., 2013). The increasing levels of light

408 pollution demand for further studies to understand the mechanism(s) of how light attracts
409 animals and interferes with their behaviour.
410

Ethics

Permission to carry out fieldwork was provided by the relevant German authority (Landratsamt Starnberg, permit # 55.1-8646.NAT_02-8-41).

Data accessibility

All data, R-scripts and a flight video are provided as electronic supplementary material.

Author contributions

TH and HRG conceived of and designed the study. TH collected and analysed the data. TH and HRG wrote the manuscript. All authors gave final approval for publication and agree to be held accountable for the work performed therein.

Competing interests

We have no competing interests.

Funding

This study was supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation, Emmy Noether grant 241711556 to HRG) and the International Max Planck Research School for Organismal Biology (student grant to T.H.)

Acknowledgements

We thank the IMPRS for support, the participants of the Scientific Writing Course 2019 for comments on previous versions of the manuscript, and the Max Planck Institute for Ornithology Seewiesen for excellent infrastructure and support.

REFERENCES

- Acharya, L., and Fenton, M.B. (1999). Bat attacks and moth defensive behaviour around street lights. *Can J Zool* 77, 27–33.
- Agee, H.R. (1969). Response of flying bollworm moths and other tympanate moths to pulsed ultrasound. *Ann Entomol Soc Am* 62, 801–807.
- Agee, H.R., and Webb, J.C. (1969). Effects of ultrasound on capture of *Heliothis zea* and *Ostrinia nubilalis* moths in traps equipped with ultraviolet lamps. *Ann Entomol Soc Am* 62, 1248–1252.
- Arnold, J.B. (2018). ggthemes package | R Documentation.
- Bailey, L.A., Bringham, R.M., Bohn, S.J., Boyles, J.G., and Smit, B. (2019). An experimental test of the allotonic frequency hypothesis to isolate the effects of light pollution on bat prey selection. *Oecologia* 190, 367–374.
- Bates, D., Maechler, M., Bolker, B., and Walker, S. (2015). Fitting linear mixed-effects models using lme4. *J Stat Softw* 67, 1–48.
- Becker, A., Whitfield, A.K., Cowley, P.D., Järnegren, J., and Næsje, T.F. (2013). Potential effects of artificial light associated with anthropogenic infrastructure on the abundance and foraging behaviour of estuary-associated fishes. *J Appl Ecol* 50, 43–50.
- Boyles, J.G., Cryan, P.M., McCracken, G.F., and Kunz, T.H. (2011). Economic importance of bats in agriculture. *Science* 332, 41–42.
- Cravens, Z.M., Brown, V.A., Divoll, T.J., and Boyles, J.G. (2018). Illuminating prey selection in an insectivorous bat community exposed to artificial light at night. *J Appl Ecol* 55, 705–713.
- Davies, T.W., and Smyth, T. (2018). Why artificial light at night should be a focus for global change research in the 21st century. *Glob Chang Biol* 24, 872–882.
- Davies, T.W., Bennie, J., Inger, R., Ibarra, N.H., and Gaston, K.J. (2013). Artificial light pollution: are shifting spectral signatures changing the balance of species interactions? *Glob Chang Biol* 19, 1417–1423.
- Davies, T.W., Duffy, J.P., Bennie, J., and Gaston, K.J. (2014). The nature, extent, and ecological implications of marine light pollution. *Front Ecol Environ* 12, 347–355.
- Falchi, F., Cinzano, P., Duriscoe, D., Kyba, C.C.M., Elvidge, C.D., Baugh, K., Portnov, B.A., Rybnikova, N.A., and Furgoni, R. (2016). The new world atlas of artificial night sky brightness. *Sci Adv* 2, e1600377.
- Fenton, M.B. (2003). Eavesdropping on the echolocation and social calls of bats. *Mammal Rev* 33, 193–204.
- Fouquet, R. (2006). Seven centuries of energy services: the price and use of light in the United Kingdom (1300–2000). *Energy Journal* 27, 139–177.

470 Fox, J. (2019). car package | R Documentation.

471 Gordon, and Ter Hofstede, H.M. (2018). The influence of bat echolocation call duration and
472 timing on auditory encoding of predator distance in noctuid moths. *J Exp Biol* 221,
473 jeb171561.

474 Hartig, F. (2019). DHARMA: Residual Diagnostics for Hierarchical (Multi-Level / Mixed)
475 Regression Models.

476 ter Hofstede, H.M., Ratcliffe, J.M., and Fullard, J.H. (2008). Nocturnal activity positively
477 correlated with auditory sensitivity in noctuid moths. *Biol Lett* 4, 262–265.

478 Holderied, M.W., and von Helversen, O. (2003). Echolocation range and wingbeat period
479 match in aerial-hawking bats. *Proc Royal Soc B* 270, 2293–2299.

480 Hölker, F., Moss, T., Griefahn, B., Kloas, W., Voigt, C., Henckel, D., Hänel, A., Kappeler, P.,
481 Völker, S., Schwoppe, A., et al. (2010). The dark side of light: a transdisciplinary research
482 agenda for light pollution policy. *Ecol Soc* 15.

483 Hügel, T., and Goerlitz, H.R. (2019): Species-specific strategies increase unpredictability of
484 escape flight in eared moths. *Functional Ecology* 33, 1674-1683.

485 Gillam, E.H., Westbrook, J.K., Schleider, P.G., and McCracken, G.F. (2001). Virtual bats and
486 real insects: effects of echolocation on pheromone-tracking behavior of male corn earworm
487 moths, *Helicoverpa zea*. *The Southwestern Naturalist* 56(1), 103-107.

488 Kassambara, A. (2019). rstatix package | R Documentation.

489 Kasso, M., and Balakrishnan, M. (2013). Ecological and economic importance of bats (order
490 Chiroptera). *ISRN Ecol Article ID 187415*, 9 pages.

491 Knop, E., Zoller, L., Ryser, R., Gerpe, C., Hörler, M., and Fontaine, C. (2017). Artificial light at
492 night as a new threat to pollination. *Nature* 548, 206–209.

493 Korner-Nievergelt, F., Rpth, T., von Felten, S., Guelat, J., Almasi, B., and Korner-Nievergelt, P.
494 (2015). *blmeco*.

495 Kunz, T.H., Braun de Torrez, E., Bauer, D., Lobova, T., and Flemming, T.H. (2011). Ecosystem
496 services provided by bats. *Ann NY Acad Sci* 1223, 1–38.

497 Lewanzik, D., and Goerlitz, H.R. (2017). Continued source level reduction during attack in the
498 low-amplitude bat *Barbastella barbastellus* prevents moth evasive flight. *Funct Ecol* 32,
499 1251–1261.

500 Lewis, F., Butler, A., and Gilbert, L. (2011). A unified approach to model selection using the
501 likelihood ratio test. *Methods Ecol Evol* 2, 155–162.

502 Longcore, T., and Rich, C. (2004). Ecological light pollution. *Front Ecol Environ* 2, 191–198.

- 503 Macgregor, C.J., Evans, D.M., Fox, R., and Pocock, M.J.O. (2017). The dark side of street
504 lighting: impacts on moths and evidence for the disruption of nocturnal pollen transport.
505 Glob Chang Biol 23, 697–707.
- 506 Macgregor, C.J., Pocock, M.J.O., Fox, R., and Evans, D.M. (2019). Effects of street lighting
507 technologies on the success and quality of pollination in a nocturnally pollinated plant.
508 Ecosphere 10, e02550.
- 509 Madsen, B.M., and Miller, L.A. (1987). Auditory input to motor neurons of the dorsal
510 longitudinal flight muscles in a noctuid moth (*Barathra brassicae* L.). J. Comp. Physiol. 160,
511 23–31.
- 512 Mathews, F., Roche, N., Aughney, T., Jones, N., Day, J., Baker, J., and Langton, S. (2015).
513 Barriers and benefits: implications of artificial night-lighting for the distribution of common
514 bats in Britain and Ireland. Philos Trans R Soc Lond B Biol Sci 370, 20140124.
- 515 McCaw, Z. (2018). RNOmni package | R Documentation.
- 516 Merckx, T., and Slade, E.M. (2014). Macro-moth families differ in their attraction to light:
517 implications for light-trap monitoring programmes. Insect Conserv Divers 7, 453–461.
- 518 Miller Colleen R., Barton Brandon T., Zhu Likai, Radeloff Volker C., Oliver Kerry M., Harmon
519 Jason P., and Ives Anthony R. (2017). Combined effects of night warming and light pollution
520 on predator–prey interactions. P Roy Soc P B Biol Sci 284, 20171195.
- 521 Minnaar, C., Boyles, J.G., Minnaar, I.A., Sole, C.L., and McKechnie, A.E. (2015). Stacking the
522 odds: light pollution may shift the balance in an ancient predator-prey arms race. J Appl Ecol
523 52, 522–531.
- 524 Nakano, R., Takanashi, T., and Surlykke, A. (2015). Moth hearing and sound communication.
525 J Comp Physiol A 201, 111–121.
- 526 Obrist, M.K., Boesch, R., and Fluckiger, P.F. (2004). Variability in echolocation call design of
527 26 Swiss bat species: consequences, limits and options for automated field identification
528 with a synergetic pattern recognition approach. Mammalia 68, 307–322.
- 529 Owens, A.C.S., and Lewis, S.M. (2018). The impact of artificial light at night on nocturnal
530 insects: a review and synthesis. Ecol Evol 1–22.
- 531 Roeder, K.D. (1962). The behaviour of free flying moths in the presence of artificial ultrasonic
532 pulses. Anim Behav 10, 300–304.
- 533 Roeder, K.D. (1964). Aspects of the noctuid tympanic nerve response having significance in
534 the avoidance of bats. J Insect Physiol 10, 529–546.
- 535 Roeder, K.D. (1967). Turning tendency of moths exposed to ultrasound while in stationary
536 flight. J Insect Physiol 13, 873–888.
- 537 Roeder, K.D. (1974). Responses of the less sensitive acoustic sense cells in the tympanic
538 organs of some noctuid and geometrid moths. J Insect Physiol 20, 55–66.

- 539 Russo, D., Cosentino, F., Festa, F., De Benedetta, F., Pejic, B., Cerretti, P., and Ancillotto, L.
540 (2019). Artificial illumination near rivers may alter bat-insect trophic interactions. *Environ*
541 *Pollut* 252, 1671–1677.
- 542 Rydell, J. (1992). Exploitation of insects around streetlamps by bats in Sweden. *Funct Ecol* 6,
543 744–750.
- 544 Rydell, J., Skals, N., Surlykke, A., and Svensson, A.M. (1997). Hearing and bat defence in
545 geometrid winter moths. *Proc Biol Sci* 264, 83–88.
- 546 Schnitzler, H.-U., Moss, C.F., and Denzinger, A. (2003). From spatial orientation to food
547 acquisition in echolocating bats. *Trends Ecol Evol* 18, 386–394.
- 548 Seibert, A.-M., Koblitz, J.C., Denzinger, A., and Schnitzler, H.-U. (2015). Bidirectional
549 echolocation in the bat *Barbastella barbastellus*: different signals of low source level are
550 emitted upward through the nose and downward through the mouth. *PLOS ONE* 10,
551 e0135590.
- 552 Skals, N., and Surlykke, A. (2000). Hearing and evasive behaviour in the greater wax moth,
553 *Galleria mellonella* (Pyralidae). *Physiol Entomol* 25, 354–362.
- 554 Skiba, R. (2014). Europäische Fledermäuse (Magdeburg: VerlagsKG Wolf).
- 555 Stone, E.L., Jones, G., and Harris, S. (2009). Street lighting disturbs commuting bats. *Curr Biol*
556 19, 1123–1127.
- 557 Stone, E.L., Harris, S., and Jones, G. (2015). Impacts of artificial lighting on bats: a review of
558 challenges and solutions. *Mammalian Biology* 80, 213–219.
- 559 Straka, T.M., Lentini, P.E., Lumsden, L.F., Wintle, B.A., and van der Ree, R. (2016). Urban bat
560 communities are affected by wetland size, quality, and pollution levels. *Ecol Evol* 6, 4761–
561 4774.
- 562 Straka, T.M., Greif, S., Schultz, S., Goerlitz, H.R., and Voigt, C.C. (2020). The effect of cave
563 illumination on bats. *Global Ecology and Conservation* 21, e00808.
- 564 Surlykke, A. (2003). Hearing in hooktip moths (Drepanidae: Lepidoptera). *J Exp Biol* 206,
565 2653–2663.
- 566 Surlykke, A., Filskov, M., Sanderson, C.G., Lal, D., Peters, B., Handb, Rosner, G., and Winkler
567 (1997). Hearing in geometrid moths. *Naturwissenschaften* 84, 356–359.
- 568 Surlykke, A., Filskov, M., Fullard, J.H., and Forrest, E. (1999). Auditory relationships to size in
569 noctuid moths: bigger is better. *Naturwissenschaften* 86, 238–241.
- 570 Svensson, A.M.A., and Rydell, J. (1998). Mercury vapour lamps interfere with the bat defence
571 of tympanate moths (*Operophtera spp.*; Geometridae). *Animal Behav* 55, 223–226.
- 572 Ter Hofstede, H.M., and Ratcliffe, J.M. (2016). Evolutionary escalation: the bat-moth arms
573 race. *J Exp Biol* 219, 1589–1602.

- 574 Ter Hofstede, H.M., Goerlitz, H.R., Ratcliffe, J.M., Holderied, M.W., and Surlykke, A. (2013).
 575 The simple ears of noctuid moths are tuned to the calls of their sympatric bat community. *J*
 576 *Exp Biol* 216, 3954–3962.
- 577 Treat, A.E. (1962). Comparative moth catches by an ultrasonic and a silent light trap. *Ann*
 578 *Entomol Soc Am* 55, 716–720.
- 579 Truxa, C., and Fiedler, K. (2012). Attraction to light - from how far do moths (Lepidoptera)
 580 return to weak artificial sources of light? *Eur J Entomol* 109, 77–84.
- 581 van Langevelde Frank, van Grunsven Roy H. A., Veenendaal Elmar M., and Fijen Thijs P. M.
 582 (2017). Artificial night lighting inhibits feeding in moths. *Biol Lett* 13, 20160874.
- 583 Van Toor, M.L., O'Mara, M.T., Abedi-Lartey, M., Wikelski, M., Fahr, J., and Dechmann, D.K.N.
 584 (2019). Linking colony size with quantitative estimates of ecosystem services of African fruit
 585 bats. *Curr Biol* 29, R237–R238.
- 586 Venables, W.N., and Ripley, B.D. (2002). *Modern Applied Statistics with S* (New York:
 587 Springer).
- 588 Wakefield, A., Jones, G., Stone, E.L., Harris, S., Jones, G., and Harris, S. (2015). . *R Soc*
 589 *Open Sci* 2, 150291.
- 590 Waters, D.A., and Jones, G. (1996). The peripheral auditory characteristics of noctuid moths:
 591 responses to the search-phase echolocation calls of bats. *J Exp Biol* 199, 847–856.
- 592 Wickham, H. (2018). *dplyr* package | R Documentation.
- 593 Wickham, H. (2020). *ggplot2* | R Documentation.
- 594 Wilke, C. (2019). *cowplot* package | R Documentation.
- 595 Wilson, J.F., Baker, D., Cheney, J., Cook, M., Ellis, M., Freestone, R., Gardner, D., Geen, G.,
 596 Hemming, R., Hodgers, D., et al. (2018). A role for artificial night-time lighting in long-term
 597 changes in populations of 100 widespread macro-moths in UK and Ireland: a citizen-science
 598 study. *J Insect Conserv* 22, 189–196.
- 599 Yurk, H., and Trites, A.W. (2000). Experimental attempts to reduce predation by harbor seals
 600 on out-migrating juvenile salmonids. *Trans Am Fish Soc* 129, 1360–1366.

601