ER β regulated ovarian kisspeptin plays an important role in oocyte maturation

¹V. Praveen Chakravarthi, ¹Subhra Ghosh, ^{2,4}Katherine F. Roby, ^{3,4}Michael W. Wolfe, and ^{1,4}M. A. Karim Rumi*

¹Department of Pathology and Laboratory Medicine, ²Department of Anatomy and Cell Biology, ³Department of Molecular and Integrative Physiology, ⁴Institute for Reproduction and Perinatal Research, University of Kansas Medical Center, Kansas City, Kansas, USA

Running head: Ovarian kisspeptin induces oocyte maturation

***Corresponding author:** M. A. Karim Rumi, Department of Pathology and Laboratory Medicine, University of Kansas Medical Center, Kansas City, KS 66160, USA. Email: mrumi@kumc.edu

Keywords: Kisspeptin; kisspeptin receptor; estrogen receptor β ; gonadotropin; granulosa cells; and oocyte maturation.

Grants support: This study was supported in part by the grant supports from KUMC SOM, COBRE and K-INBRE.

Disclosure: The authors do not have any conflict of interest.

Abstract

Kisspeptin (KISS1) signaling in the hypothalamic-pituitary (H-P) axis plays essential role in regulating gonadotropin secretion. KISS1 and KISS1 receptor (KISS1R) are also expressed in the ovary; however, the role of intraovarian KISS1 signaling remains largely unclear. Granulosa cell (GC)-specific expression of KISS1, and oocyte-specific expression of KISS1R indicate that GC-derived KISS1 may act on oocytes. Expression of KISS1 in GCs is induced by gonadotropins but it is absent in estrogen receptor β knockout ($Er\beta^{null}$) rats. We also observed that gonadotropin stimulation failed to induce maturation of $Er\beta^{null}$ oocytes. Interestingly, KISS1 treatment of cumulus oocyte complexes (COCs) isolated from antral follicles promotes *in vitro* maturation of oocytes. Treatment of oocytes with KISS1 induced intracellular Ca^{2+} release, and increased activation of MAP kinase ERK1/2. KISS1 treatment also induced the expression of oocyte genes that are crucial for differentiation of GCs, and maturation of oocytes. Our findings suggest that ovarian KISS1-signaling plays an important role in gonadotropin induced follicle development and oocyte maturation.

1. Introduction

Estrogen receptor β (ER β) is the predominant estrogen receptor in the ovary (Byers et al., 1997; Couse et al., 2004; Pelletier and El-Alfy, 2000; Slomczynska et al., 2001) and disruption of ER β signaling leads ovulatory defects (Asadi et al., 2013; Lang-Muritano et al., 2018). Failure of ovulation in *Erb β ^{null}* mice and rats is associated with defective follicle maturation and an attenuated preovulatory gonadotropin surge (Couse et al., 2005; Rumi et al., 2017). However, treatment with exogenous gonadotropins fails to induce ovulation, suggesting a primary ovarian defect due to the loss of ER β (Rumi et al., 2017). To identify the ER β -regulated genes that may affect ovarian follicle development, oocyte maturation, and ovulation, we performed RNA-sequencing analyses of gonadotropin stimulated granulosa cells (GCs) and oocytes (Chakravarthi et al., 2020; Chakravarthi et al., 2019; Khristi et al., 2018). We identified *Kiss1* to be one of the most-differentially expressed gene in *Erb β ^{null}* GCs (Chakravarthi et al., 2020; Khristi et al., 2018). GCs showed a marked induction in *Kiss1* expression in response to gonadotropin stimulation, which was absent in *Erb β ^{null}* rats (Chakravarthi et al., 2020; Chakravarthi et al., 2018; Khristi et al., 2018).

Hypothalamic KISS1 regulates gonadotropin secretion, which is essential for gonadal development, onset of puberty, and induction of ovulation (Javed et al., 2015; Moore et al., 2018). In recent studies, expression of KISS1 and KISS1R in the ovaries has been detected across species (Castellano et al., 2006; Gaytan et al., 2009; Shahed and Young, 2009; Terao et al., 2004). However, the physiological role of ovarian KISS1 signaling remains largely unknown.

While the expression of *Kiss1* is markedly higher in GCs, *Kiss1r* is detected predominantly in the oocytes, suggesting that GC-derived KISS1 may act on oocytes (Chakravarthi et al., 2018).

Bidirectional interactions between GCs and oocytes play vital roles in follicle development and ovulation (Craig et al., 2007; Emori and Sugiura, 2014; McNatty et al., 2003; Monniaux, 2016; Russell and Robker, 2007; Wigglesworth et al., 2013). While oocyte maturation is regulated by GC-derived factors, maturing oocytes contribute to GC differentiation and cumulus cell expansion (Craig et al., 2007; Emori and Sugiura, 2014; McNatty et al., 2003; Monniaux, 2016; Russell and Robker, 2007; Wigglesworth et al., 2013). KISS1 expressed in GCs represent the former group of signaling molecules. KISS1 expression in the ovary coincides with the preovulatory gonadotropin surge (Castellano et al., 2006) and intrafollicular KISS1 levels positively correlate with the stages of oocyte maturation (Taniguchi et al., 2017). Taken together, we hypothesize that GC-derived KISS1 may act on oocytes to effect oocyte maturation, and an attenuated expression of KISS1 in *Erβ^{null}* GCs may cause failure of follicle development and oocyte maturation despite gonadotropin stimulation.

2. Materials and methods

2.1 Animal models

Wildtype and *Erβ^{null}* Holtzman Sprague-Dawley (HSD) rats were studied for intraovarian kisspeptin signaling. *Erβ^{null}* rat model was generated by targeted deletion of the exon 3 in the *Erβ* gene (Rumi et al., 2017). Exon 3 deletion caused a frameshift and null mutation (Rumi et al.,

2017). Rats were screened for the presence of mutations by PCR using tail-tip DNA samples (RED extract-N-Amp Tissue PCR Kit, Sigma-Aldrich) as described previously (Rumi et al., 2014; Rumi et al., 2017). All procedures were performed in accordance with the protocols approved by the University of Kansas Medical Center Animal Care and Use Committee.

2.2 Gonadotropin treatment

4-wk-old $Er\beta^{null}$ and age-matched wildtype female rats were used to evaluate gonadotropin-induced follicular development. Synchronized follicular growth was performed by intraperitoneal injection of 30 IU pregnant mare serum gonadotropin (PMSG; BioVendor LLC, Asheville, NC). Forty-eight hours after the PMSG injection, 30 IU of human chorionic gonadotropin (hCG; BioVendor) was injected intraperitoneally. Rats were sacrificed prior to gonadotropin treatment (Basal), 48h after PMSG treatment (PMSG), 4h and 24h after hCG injection to PMSG treated rats and their ovaries were collected (**Fig. 1B**) as described previously (Chakravarthi et al., 2020; Chakravarthi et al., 2019; Chakravarthi et al., 2018).

2.2 Isolation of oocytes and GCs

Cumulus oocyte complexes (COCs) were isolated from the large antral follicles by needle puncture under microscopic observation as described previously (Chakravarthi et al., 2020; Chakravarthi et al., 2019; Chakravarthi et al., 2018). All cumulus cells were removed from the oocytes by mechanical pipetting followed by repeated washings into fresh media using capillary suction. GCs were collected by filtration of the remaining cells through 40 μ M cell strainers (Fisher Scientific).

110

111 **2.3 RNA-sequencing of GCs and oocytes**

112 Oocytes and GCs were collected from 4-wk-old wildtype and *Erβ^{null}* rats with or without
 113 gonadotropin treatment. Total RNA was extracted using TRI Reagent (Millipore-Sigma)
 114 following the manufactures instructions. Approximately 500ng of total RNA was used for the
 115 RNA-seq library preparation. Libraries were prepared by using a TruSeq standard mRNA kit
 116 (Illumina) following manufacturer's instructions. The cDNA libraries were sequenced at the
 117 Molecular Biology Core Laboratory of Mayo Clinic (Rochester, MN). RNA-Seq data were
 118 analyzed using CLC Genomics Workbench (Qiagen Bioinformatics). Gene expression values
 119 were reported as TPM (Transcripts per million base pairs). All the RNA seq data are available at
 120 SRA (SRX6376730, SRX6376729, SRX6376735, SRX6376752, SRX6376751, SRX6376757,
 121 SX6761695, SRX6761696, SRX6761697, SRX6761698, SRX6761699 and SRX67617) and the
 122 data articles are published in Data in Brief journal (Chakravarthi et al., 2020; Chakravarthi et al.,
 123 2019) .

124

125 **2.4 In vitro maturation of rat oocytes**

126 COCs were collected from the antral follicles of 4-wk-old wildtype and *Erβ^{null}* rats 48h
 127 after PMSG treatment. After washing three times in *in vitro* maturation media (BO-IVM, IVF
 128 Biosciences, UK) the COCs were placed in 20 µl drops of IVM medium in 35-mm culture
 129 dishes, covered with light weight mineral oil, and incubated at 37°C in a humidified CO₂
 130 incubator for an additional 24h (Kona et al., 2016; Praveen Chakravarthi et al., 2015). At the end
 131 of incubation, oocytes were denuded of the cumulus cells by repeated pipetting through a fine

bore glass pipette and examined microscopically for extrusion of the polar body as indicative of oocyte maturation to MII stage (Kona et al., 2016; Praveen Chakravarthi et al., 2015).

2.5 Calcium release assay

4-wk-old wildtype rats were administered with PMSG, and 48h later COCs were isolated. COCs were incubated with 5 μ M Fluo-3AM (ThermoFisher Scientific) solution containing 20% Pluronic F-127 (ThermoFisher Scientific) for 30 min in a 37°C incubator. All cumulus cells were removed from the oocytes by pipetting followed by repeated washings into fresh media using capillary suction and oocytes were placed into HBSS drops containing 1nM of rat kisspeptin-10 (KP-10) (Tocris Bioscience, Bristol, UK). Oocytes were monitored under a fluorescent microscope for 3 min and images were recorded every 30 sec.

2.6 Protein extraction and western blotting

COCs were collected from 4-wk-old wildtype rats 48h after PMSG administration. Oocytes and GCs were purified as described in Section 2.2 and proteins were extracted in 1X SDS lysis buffer (62.5mM Tris-HCl pH 6.8, 2% SDS, 42 mM dithiothreitol, 10% glycerol, and 0.01% bromophenol blue), containing protease and phosphatase inhibitors (Cell Signaling Technologies, Danvers, MA). In addition, oocytes were cultured with or without 1nM or 10nM KP-10. After treatment, oocytes were denuded mechanically, proteins were extracted in 1XSDS buffer. Oocyte or GC cell-lysates in 1XSDS buffer were sonicated to shear DNA and reduce viscosity, heat denatured, and separated on a 4-20% SDS-PAGE. Electrophoresed proteins were transferred from the gel to PVDF membranes, blocked with 5% skim milk in TBST (1XTBS

buffer containing 0.1% Tween-20), and incubated overnight at 4°C with specific primary antibodies at the appropriate dilution in blocking solution (**Supplementary Table 1**). After removing the unbound primary antibody solution, membranes were washed with TBST, blocked, and incubated with peroxidase conjugated anti-mouse, or anti-rabbit secondary antibodies (Jackson ImmunoResearch, West Grove, PA) at a dilution of 1:10,000 to 50,000, and the immunoreactive signals were visualized with Luminata Crescendo HRP substrate (Millipore Sigma, Burlington, MA).

2.7 Preparation cDNA and RT-qPCR analyses

Total RNA was extracted from the GCs and oocytes using TRI Reagents (Sigma-Aldrich). 1000 ng of total RNA from each sample was used for the preparation of cDNAs using High-Capacity cDNA Reverse Transcription Kits (Applied Biosystems, Foster City, CA). RT-qPCR amplification of cDNAs was carried out in a 10µl reaction mixture containing Applied Biosystems Power SYBR Green PCR Master Mix (Thermo Fisher Scientific). Amplification and fluorescence detection of RT-qPCR were carried out on Applied Biosystems Quant Studio Flex 7 Real Time PCR System (Thermo Fisher Scientific). The $\Delta\Delta CT$ method was used for relative quantification of target mRNA expression level normalized to Rn18s (18S rRNA). A list of qPCR primers is shown in **Supplementary Table 2**.

2.8 Statistical analyses

Ovulation counting was performed on 6 individual rats of the same genotype at each time point. Each RNA-sequencing library was prepared using pooled RNA samples from three

individual wildtype or $Er\beta^{null}$ rats and each group of RNA-sequencing consisted of three libraries. Gene expression analyses were performed on at least 6 individual rats. The experimental results are expressed as mean $\pm$ SE. Statistical comparisons between two means were determined with Student's t-test while comparisons among multiple means were evaluated with ANOVA and the significance of mean differences was determined by Duncan post hoc test, with $P \leq 0.05$ considered a significant level of difference. All statistical calculations were done with SPSS 22 (IBM, Armonk, NY).

3. Results

3.1. $ER\beta$ is essential for oocyte maturation

8-12 wk old adult $Er\beta^{null}$ rats failed to ovulate during a natural estrous cycle (**Fig. 1A**). Administration of exogenous gonadotropins also failed to induce ovulation in 4-wk-old $Er\beta^{null}$ rats (**Fig. 1C**). We observed that $Er\beta^{null}$ oocytes isolated from the antral follicles prior to gonadotropin stimulation appeared similar to those isolated from wildtype follicles (**Fig. 1D** and **E**). However, gonadotropin treatment resulted in an abnormal maturation of the $Er\beta^{null}$ oocytes (**Fig. 1F-K**). Development of abnormal and fragmented nuclei in $Er\beta^{null}$ oocytes became more evident after hCG treatment (**Fig. 1H-K**).

3.2. $Kiss1$ is expressed in GCs and $Kiss1r$ in oocytes

We observed a prominent expression of $Kiss1$ in gonadotropin treated GCs (**Fig. 2A**) while $Kiss1r$ in oocytes (**Fig. 2B**). RNA-sequencing revealed a high level of $Kiss1$ transcripts in

wildtype GCs, but the expression was severely reduced or absent in $Er\beta^{null}$ GCs (**Fig. 2C**). These findings were validated by qRT-PCR analyses (**Fig. 2D**). In contrast to *Kiss1*, *Kiss1r* expression did not show any significant difference between the wildtype or $Er\beta^{null}$ oocytes (**Fig. 2E and F**).

3.3. *KISS1* treatment induces *in vitro* maturation of oocytes

We observed that $Er\beta^{null}$ oocytes exhibit defects in *in vitro* maturation. While approximately 55% of wildtype oocytes mature to MII stage (**Fig. 3A**), less than 35% of $Er\beta^{null}$ oocytes undergo *in vitro* maturation (**Fig. 3D**). Remarkably, treatment with rat kisspeptin 10 (KP-10) increased the rate of oocyte maturation in both wildtype (**Fig. 3B and C**) and $Er\beta^{null}$ oocytes (**Fig. 3E and F**).

3.4. *KISS1* signaling in rat oocytes

Western blot analyses detected the expression of KISS1R in oocytes but not in GCs (**Fig. 4A**). Treatment of oocytes with rat KP-10 induced a prominent intracellular Ca^{2+} release (**Fig. 4B-E**). KP-10 treatment also activated the ERK1/2 in oocytes (**Fig. 4F**), but not the AKT (**Fig. 4G**). While 1nM of KP-10 induced marked activation of ERK1/2, an increased concentration failed to show such an effect.

3.5. *KP-10* treatment induces genes involved in oocyte maturation

Expression of genes known to be involved in regulating oocyte maturation were assessed by RT-qPCR analyses (**Fig. 5**). KP-10 treatment upregulated the expression of *Gdf9* and *Bmp15*, (**Fig. 5 A and B**), whereas it downregulated the expression of *Bmp7* (**Fig. 5 C**). Expression of

genes involved in protein kinase activation required for oocyte maturation including *cMos*, *Kit* and *Aurka* (**Fig. 5 D-F**) were also increased by KP-10 stimulation of oocytes. Similarly, KP-10 treatment increased the expression of genes involved in meiosis including *Wee2*, *Cdc25b* and *Rec8* (**Fig. 5 G-I**). RNA binding proteins like *Dazl*, *Btg4* and *Papbc1l*, which are known to induce mRNA stability and selective activation of genes were also upregulated with KP-10 treatment (**Fig. 5 J-L**).

4. Discussion

We have studied the role of ovarian derived KISS1 using an *Erb^{null}* mutant rat model that fails to induce *Kiss1* gene expression in GCs in response to gonadotropins (Chakravarthi et al., 2020; Chakravarthi et al., 2018; Khristi et al., 2018) and suffer from defective oocyte maturation. Intraovarian *Kiss1* gene expression is induced by the preovulatory gonadotropin surge that initiates the final steps of follicle development and oocyte maturation (Castellano et al., 2006). We also detected a dramatic upregulation of *Kiss1* gene expression in GCs following the administration of hCG to PMSG primed rats (Chakravarthi et al., 2020; Chakravarthi et al., 2018; Khristi et al., 2018). The expression pattern of *Kiss1* and *Kiss1r* (Chakravarthi et al., 2018) suggest that GC-derived KISS1 may act on the KISS1R in oocytes to regulate oocyte maturation (**Fig. 6**). An absence of *Kiss1* gene induction in *Erb^{null}* GCs may be linked to the defective oocyte maturation in *Erb^{null}* rats in response to gonadotropin treatment.

We observed that treatment of oocytes with a low dose (1nM) of KP-10 improved *in vitro* maturation of rat oocytes. Previous studies on pig (Saadeldin et al., 2012) and sheep (Byri et al.,

2017) oocytes also support our findings. Although *Erβ^{null}* oocytes showed a reduced rate of *in vitro* maturation, both of wildtype and *Erβ^{null}* oocytes responded to rat KP-10 treatment showing a similar improvement. This can be explained by similar levels of *Kiss1r* expression in wildtype and *Erβ^{null}* oocytes (Fig. 2E, F) and we would assert normal intracellular signaling induced by KP-10.

KISS1R protein is expressed in rat oocytes and responds to KISS1 treatment. Intracellular Ca^{2+} release immediately after KP-10 treatment further verifies the presence of KISS1R and indicates specific KISS1/KISS1R signaling. KISS1R is a G-protein coupled receptor that signals through intracellular Ca^{2+} release. Activation of MAPK kinase ERK1/2 following kisspeptin treatment has been reported with KISS1/KISS1R signaling in hypothalamic neurons (d'Anglemont de Tassigny and Colledge, 2010) and we have confirmed that this signaling pathway is maintained in oocytes. Conversely, we did not observe activation of AKT by KP-10 in oocytes, but this is a typical response to KISS1/KISS1R signaling in the hypothalamic neurons (d'Anglemont de Tassigny and Colledge, 2010). In oocytes, Ca^{2+} stored in the endoplasmic reticulum and generates a Ca^{2+} efflux that is important for signaling and inducing oocyte maturation (Kline, 2000; Stricker et al., 1998; Tosti, 2006; Wang and Machaty, 2013). Ca^{2+} efflux has also been reported to induce RAF/MEK/ERK pathway through activation of RAS (Atherfold et al., 1999; Li et al., 2005). Furthermore, ERK1/2 activation has also been found essential for oocyte maturation (Choi et al., 1996). We observed that KP-10 induced Ca^{2+} efflux in oocytes, which resulted in an upregulation of ERK1/2 activation. Thus KP-10 induction of both Ca^{2+} efflux and ERK1/2 activation may be responsible for the enhanced oocyte maturation that was observed in the current study.

265

266 To elucidate the mechanisms downstream of Ca^{2+} signaling and ERK1/2 activation, we
 267 investigated transcript levels of genes that play important roles in oocyte maturation. We
 268 observed that KP-10 treatment upregulated the expression of *Gdf9*, *Bmp15*, *cMos*, *Kit*, *Aurka*,
 269 *Wee2*, *Cdc25b*, *Dazl*, *Btg4*, and *Papbc1l*. BMP15 and GDF9 are known to promote cumulus
 270 expansion and regulate oocyte maturation (Chang et al., 2016; McNatty et al., 2005a; McNatty et
 271 al., 2005b; Persani et al., 2014). ERK/MAPK pathway has been shown to induce the expression
 272 of *Bmp15* and *Gdf9* which are essential for growth and differentiation of GCs during oocyte
 273 maturation (Mottershead et al., 2012; Reader et al., 2011). KP-10 treatment reduced the
 274 expression of BMP7 which has been shown to inhibit oocyte maturation (Yang et al., 2018).
 275 cMOS has been reported to be involved in the spindle formation, activation of maturation
 276 promoting factor, and oocyte maturation (Araki et al., 1996; Chang et al., 2016; Choi et al.,
 277 1996; Hashiba et al., 2001). KIT/KITL signaling also enhances the oocyte maturation in *in vitro*
 278 by overcoming the inhibitory effect of NPPC signaling (Medvedev et al., 2011; Ye et al., 2009).
 279 AURKA is a serine /threonine kinase that accumulates on microtubule organizing centers and is
 280 involved in the resumption of meiosis (Saskova et al., 2008). WEE2 is involved in the inhibitory
 281 phosphorylation of CDC2 that drives the exit from metaphase II (MII), promoting oocyte
 282 maturation (Oh et al., 2011). CDC25 is a dual specificity phosphatase that activates cyclin
 283 dependent kinases and regulates meiotic maturation of oocytes (Lincoln et al., 2002). *Dazl*
 284 encodes an RNA binding protein that plays an important role in spindle assembly, MI-MII
 285 transition and oocyte maturation (Chen et al., 2011). PABPC, a poly A binding protein that
 286 increases stability of selective mRNAs, promotes mRNA translation and improves oocyte
 287 maturation (Lowther and Mehlmann, 2015). In addition, BTG4 is involved in selective

degradation of oocyte mRNA that is required for maternal-zygotic transition (Yu et al., 2016). We speculate that these KP-10 induced changes in gene expression play important role in promoting oocyte maturation.

It is well established that KISS1 regulation of oocyte maturation and induction of ovulation is mediated indirectly through gonadotropin secretion from the H-P axis. Our findings indicate that the role for KISS1 in regulating female fertility can be explained beyond the HP-axis to intrafollicular expression of KISS1 and signaling, which plays an important role in regulating oocyte maturation.

Acknowledgements: This study was supported in part by the grant supports from KUMC SOM, COBRE and K-INBRE.

5. Figure legends

Fig. 1. Abnormal oocyte maturation in $Er\beta^{null}$ rats. 8-12 wk-old $Er\beta^{null}$ adult rats are infertile and they failed to ovulate either during natural estrus cycle (**A**) or induced by exogenous gonadotropins to 4 wk-old rats (**C**). Oocytes isolated from 4-wk-old wildtype (WT) and $Er\beta^{null}$ rat ovaries at different stages of gonadotropin treatment were examined after removal of cumulus cells. Although unstimulated (Basal **D** and **E**) oocytes of both genotypes appeared similar, abnormal division and fragmentation was observed in $Er\beta^{null}$ oocytes after PMSG 48h (**F** and **G**), 4h after hCG (**H** and **I**), and 24h after hCG (**J** and **K**) treatment.

Fig. 2. *Kiss1* and *Kiss1r* expression in rat ovary. Oocytes and granulosa cells (GCs) were isolated from the ovaries collected 4h after hCG administration to PMSG treated 4-wk-old rats. RT-qPCR analyses showed that expression of *Kiss1* was remarkably higher in (GCs) (A), whereas *Kiss1r* in the oocytes (B). RNA sequencing analyses showed a significantly higher expression of *Kiss1* in wildtype (WT) GCs compared to *Erβ^{null}* GCs (C), but a similar level of *Kiss1r* expression in both WT and *Erβ^{null}* oocytes (E). RT-qPCR data from GCs and oocytes further confirmed the results of RNA-sequencing (D and F). RNA-sequencing data are presented in triplicates. RT-qPCR data are represented as mean ± SE. n = 6. *P ≤ 0.05.

Fig. 3. Role of KISS1 in oocyte maturation. 48h after PMSG treatment, cumulus oocyte complex (COCs) were isolated from 4-wk-old wildtype and *Erβ^{null}* rats and cultured in BO-IVM in presence or absence of 1nM of rat kisspeptin 10 (KP-10) for 20h. Approximately 55% of wildtype (A and C) and only 35% of *Erβ^{null}* oocytes achieved meiotic maturation (MII) in BO-IVM culture (control) (D and F). Remarkably, addition of KP-10 to BO-IVM increased the *in vitro* maturation to 78% of wildtype (B and C) and 56% of *Erβ^{null}* oocytes (E and F). Each treatment group contained at least 25 oocytes at each time point and repeated three times. Cont, Control.

Fig. 4. Role of KISS1 in inducing Calcium efflux and ERK signaling. The expression of KISS1R was significantly higher in oocytes compared to granulosa cells (GCs) as shown by western blot analysis (A). Oocytes from PMSG treated rats were examined for calcium efflux using a Fluo-3 assay kit. Calcium efflux increased after the treatment with rat kisspeptin 10 (KP-10) (E) compared to control (C). Treatment with 1nM of KP-10 also markedly elevated the

phosphorylation of ERK1/2 (F) but not AKT (G) in oocytes. Ph, Phase contrast; Fluo, Fluorescence microscopy; Cont, Control.

Fig. 5. KP-10 treatment regulated genes involved in inducing oocyte maturation. 48h after PMSG treatment, cumulus oocyte complex (COCs) were isolated from 4-wk-old wildtype female rats and treated with 1nM of rat kisspeptin 10 (KP-10) for 30 min. KP-10 treatment upregulated the mRNA levels of *Gdf9*, *Bmp15*, but downregulated *Bmp7* (A-C). It also upregulated the mRNA levels of protein kinases *cMos*, *Kit*, and *Aurka* (D-F), as well as *Wee2* and *Cdc25b* (G and H) important for meiotic maturation. Moreover, KP-10 treatment upregulated the mRNA levels of RNA binding proteins *Dazl*, *Btg4*, and *Papbc1l* (J-L) required for selective mRNA stability during oocyte maturation. RT-qPCR data represent the mean $\pm$ SE. *P < 0.05, n = 6.

Fig. 6. Intraovarian role of ovarian derived KISS1. Based on our findings, we conclude that the preovulatory LH surge induces the expression of KISS1 (KP) by granulosa cells (GCs), which is ER β -dependent. GC-derived KP acts on the KISS1R (KR) expressed in oocytes to increase Ca²⁺ release, ERK phosphorylation, and modulation of genes, which are important for inducing oocyte maturation.

6. References

Araki, K., K. Naito, S. Haraguchi, R. Suzuki, M. Yokoyama, M. Inoue, S. Aizawa, Y. Toyoda, and E. Sato. 1996. Meiotic abnormalities of c-mos knockout mouse oocytes: activation

after first meiosis or entrance into third meiotic metaphase. *Biology of reproduction*.
55:1315-1324.

Asadi, M., S. Ghafouri-Fard, D. Zare-Abdollahi, A. Ebrahim-Habibi, and N. Matin. 2013.
Estrogen receptor mutation in a girl with primary amenorrhea. *Clinical genetics*. 83:497-
498.

Atherfold, P.A., M.S. Norris, P.J. Robinson, E.W. Gelfand, and R.A. Franklin. 1999. Calcium-
induced ERK activation in human T lymphocytes. *Molecular immunology*. 36:543-549.

Byers, M., G.G. Kuiper, J.A. Gustafsson, and O.K. Park-Sarge. 1997. Estrogen receptor-beta
mRNA expression in rat ovary: down-regulation by gonadotropins. *Molecular
endocrinology (Baltimore, Md.)*. 11:172-182.

Byri, P., A. Gangineni, K.R. Reddy, and K.B.P. Raghavender. 2017. Effect of kisspeptin on in
vitro maturation of sheep oocytes. *Veterinary world*. 10:276-280.

Castellano, J.M., M. Gaytan, J. Roa, E. Vigo, V.M. Navarro, C. Bellido, C. Dieguez, E. Aguilar,
J.E. Sanchez-Criado, A. Pellicer, L. Pinilla, F. Gaytan, and M. Tena-Sempere. 2006.
Expression of KiSS-1 in rat ovary: putative local regulator of ovulation? *Endocrinology*.
147:4852-4862.

Chakravarthi, V.P., S. Ghosh, E. Dai, D. Pathak, and M.A.K. Rumi. 2020. Transcriptome
datasets of ESR2-regulated genes in rat granulosa cells during gonadotropin-induced
follicle maturation. *Data in brief*. 30:105405.

Chakravarthi, V.P., S. Ghosh, R. Roy, E. Dai, D. Pathak, and M.A.K. Rumi. 2019.
Transcriptome datasets of gonadotropin-induced ESR2-regulated genes in rat oocytes.
Data in brief. 27:104786.

- Chakravarthi, V.P., V. Khristi, S. Ghosh, S. Yerrathota, E. Dai, K.F. Roby, M.W. Wolfe, and M.A.K. Rumi. 2018. ESR2 Is Essential for Gonadotropin-Induced Kiss1 Expression in Granulosa Cells. *Endocrinology*. 159:3860-3873.
- Chang, H.M., J. Qiao, and P.C. Leung. 2016. Oocyte-somatic cell interactions in the human ovary-novel role of bone morphogenetic proteins and growth differentiation factors. *Human reproduction update*. 23:1-18.
- Chen, J., C. Melton, N. Suh, J.S. Oh, K. Horner, F. Xie, C. Sette, R. Blelloch, and M. Conti. 2011. Genome-wide analysis of translation reveals a critical role for deleted in azoospermia-like (Dazl) at the oocyte-to-zygote transition. *Genes & development*. 25:755-766.
- Choi, T., K. Fukasawa, R. Zhou, L. Tessarollo, K. Borror, J. Resau, and G.F. Vande Woude. 1996. The Mos/mitogen-activated protein kinase (MAPK) pathway regulates the size and degradation of the first polar body in maturing mouse oocytes. *Proceedings of the National Academy of Sciences of the United States of America*. 93:7032-7035.
- Couse, J.F., M.M. Yates, B.J. Deroo, and K.S. Korach. 2005. Estrogen receptor-beta is critical to granulosa cell differentiation and the ovulatory response to gonadotropins. *Endocrinology*. 146:3247-3262.
- Couse, J.F., M.M. Yates, R. Sanford, A. Nyska, J.H. Nilson, and K.S. Korach. 2004. Formation of cystic ovarian follicles associated with elevated luteinizing hormone requires estrogen receptor-beta. *Endocrinology*. 145:4693-4702.
- Craig, J., M. Orisaka, H. Wang, S. Orisaka, W. Thompson, C. Zhu, F. Kotsuji, and B.K. Tsang. 2007. Gonadotropin and intra-ovarian signals regulating follicle development and atresia:

the delicate balance between life and death. *Frontiers in bioscience : a journal and virtual library*. 12:3628-3639.

d'Anglemont de Tassigny, X., and W.H. Colledge. 2010. The role of kisspeptin signaling in reproduction. *Physiology (Bethesda, Md.)*. 25:207-217.

Emori, C., and K. Sugiura. 2014. Role of oocyte-derived paracrine factors in follicular development. *Animal science journal = Nihon chikusan Gakkaiho*. 85:627-633.

Gaytan, F., M. Gaytan, J.M. Castellano, M. Romero, J. Roa, B. Aparicio, N. Garrido, J.E. Sanchez-Criado, R.P. Millar, A. Pellicer, H.M. Fraser, and M. Tena-Sempere. 2009. KiSS-1 in the mammalian ovary: distribution of kisspeptin in human and marmoset and alterations in KiSS-1 mRNA levels in a rat model of ovulatory dysfunction. *American journal of physiology. Endocrinology and metabolism*. 296:E520-531.

Hashiba, Y., Y. Asada, O. Heikinheimo, S.E. Lanzendorf, and S. Mizutani. 2001. Microinjection of antisense c-mos oligonucleotides prevents the progression of meiosis in human and hamster oocytes. *Fertility and sterility*. 76:143-147.

Javed, Z., U. Qamar, and T. Sathyapalan. 2015. The role of kisspeptin signalling in the hypothalamic-pituitary-gonadal axis--current perspective. *Endokrynologia Polska*. 66:534-547.

Khristi, V., V.P. Chakravarthi, P. Singh, S. Ghosh, A. Pramanik, A. Ratri, S. Borosha, K.F. Roby, M.W. Wolfe, and M.A.K. Rumi. 2018. ESR2 regulates granulosa cell genes essential for follicle maturation and ovulation. *Molecular and cellular endocrinology*. 474:214-226.

Kline, D. 2000. Attributes and dynamics of the endoplasmic reticulum in mammalian eggs. *Current topics in developmental biology*. 50:125-154.

- Kona, S.S., V. Praveen Chakravarthi, A.V. Siva Kumar, D. Srividya, K. Padmaja, and V.H. Rao. 2016. Quantitative expression patterns of GDF9 and BMP15 genes in sheep ovarian follicles grown in vivo or cultured in vitro. *Theriogenology*. 85:315-322.
- Lang-Muritano, M., P. Sproll, S. Wyss, A. Kolly, R. Hurlimann, D. Konrad, and A. Biason-Laubert. 2018. Early-Onset Complete Ovarian Failure and Lack of Puberty in a Woman With Mutated Estrogen Receptor beta (ESR2). *The Journal of clinical endocrinology and metabolism*. 103:3748-3756.
- Li, D.W., J.P. Liu, Y.W. Mao, H. Xiang, J. Wang, W.Y. Ma, Z. Dong, H.M. Pike, R.E. Brown, and J.C. Reed. 2005. Calcium-activated RAF/MEK/ERK signaling pathway mediates p53-dependent apoptosis and is abrogated by alpha B-crystallin through inhibition of RAS activation. *Molecular biology of the cell*. 16:4437-4453.
- Lincoln, A.J., D. Wickramasinghe, P. Stein, R.M. Schultz, M.E. Palko, M.P. De Miguel, L. Tessarollo, and P.J. Donovan. 2002. Cdc25b phosphatase is required for resumption of meiosis during oocyte maturation. *Nature genetics*. 30:446-449.
- Lowther, K.M., and L.M. Mehlmann. 2015. Embryonic Poly(A)-Binding Protein Is Required During Early Stages of Mouse Oocyte Development for Chromatin Organization, Transcriptional Silencing, and Meiotic Competence. *Biology of reproduction*. 93:43.
- McNatty, K.P., J.L. Juengel, K.L. Reader, S. Lun, S. Myllymaa, S.B. Lawrence, A. Western, M.F. Meerasahib, D.G. Mottershead, N.P. Groome, O. Ritvos, and M.P. Laitinen. 2005a. Bone morphogenetic protein 15 and growth differentiation factor 9 co-operate to regulate granulosa cell function. *Reproduction (Cambridge, England)*. 129:473-480.
- McNatty, K.P., J.L. Juengel, K.L. Reader, S. Lun, S. Myllymaa, S.B. Lawrence, A. Western, M.F. Meerasahib, D.G. Mottershead, N.P. Groome, O. Ritvos, and M.P. Laitinen. 2005b.

Bone morphogenetic protein 15 and growth differentiation factor 9 co-operate to regulate granulosa cell function in ruminants. *Reproduction (Cambridge, England)*. 129:481-487.

McNatty, K.P., J.L. Juengel, T. Wilson, S.M. Galloway, G.H. Davis, N.L. Hudson, C.L. Moeller, M. Cranfield, K.L. Reader, M.P. Laitinen, N.P. Groome, H.R. Sawyer, and O. Ritvos. 2003. Oocyte-derived growth factors and ovulation rate in sheep. *Reproduction (Cambridge, England) Supplement*. 61:339-351.

Medvedev, S., H. Pan, and R.M. Schultz. 2011. Absence of MSY2 in mouse oocytes perturbs oocyte growth and maturation, RNA stability, and the transcriptome. *Biology of reproduction*. 85:575-583.

Monniaux, D. 2016. Driving folliculogenesis by the oocyte-somatic cell dialog: Lessons from genetic models. *Theriogenology*. 86:41-53.

Moore, A.M., L.M. Coolen, D.T. Porter, R.L. Goodman, and M.N. Lehman. 2018. KNDy Cells Revisited. *Endocrinology*. 159:3219-3234.

Mottershead, D.G., L.J. Ritter, and R.B. Gilchrist. 2012. Signalling pathways mediating specific synergistic interactions between GDF9 and BMP15. *Molecular human reproduction*. 18:121-128.

Oh, J.S., A. Susor, and M. Conti. 2011. Protein tyrosine kinase Wee1B is essential for metaphase II exit in mouse oocytes. *Science (New York, N.Y.)*. 332:462-465.

Pelletier, G., and M. El-Alfy. 2000. Immunocytochemical localization of estrogen receptors alpha and beta in the human reproductive organs. *The Journal of clinical endocrinology and metabolism*. 85:4835-4840.

Persani, L., R. Rossetti, E. Di Pasquale, C. Cacciatore, and S. Fabre. 2014. The fundamental role of bone morphogenetic protein 15 in ovarian function and its involvement in female fertility disorders. *Human reproduction update*. 20:869-883.

Praveen Chakravarthi, V., S.S. Kona, A.V. Siva Kumar, M. Bhaskar, and V.H. Rao. 2015. Quantitative expression of antiapoptotic and proapoptotic genes in sheep ovarian follicles grown in vivo or cultured in vitro. *Theriogenology*. 83:590-595.

Reader, K.L., D.A. Heath, S. Lun, C.J. McIntosh, A.H. Western, R.P. Littlejohn, K.P. McNatty, and J.L. Juengel. 2011. Signalling pathways involved in the cooperative effects of ovine and murine GDF9+BMP15-stimulated thymidine uptake by rat granulosa cells. *Reproduction (Cambridge, England)*. 142:123-131.

Rumi, M.A., P. Dhakal, K. Kubota, D. Chakraborty, T. Lei, M.A. Larson, M.W. Wolfe, K.F. Roby, J.L. Vivian, and M.J. Soares. 2014. Generation of Esr1-knockout rats using zinc finger nuclease-mediated genome editing. *Endocrinology*. 155:1991-1999.

Rumi, M.A.K., P. Singh, K.F. Roby, X. Zhao, K. Iqbal, A. Ratri, T. Lei, W. Cui, S. Borosha, P. Dhakal, K. Kubota, D. Chakraborty, J.L. Vivian, M.W. Wolfe, and M.J. Soares. 2017. Defining the Role of Estrogen Receptor beta in the Regulation of Female Fertility. *Endocrinology*. 158:2330-2343.

Russell, D.L., and R.L. Robker. 2007. Molecular mechanisms of ovulation: co-ordination through the cumulus complex. *Human reproduction update*. 13:289-312.

Saadeldin, I.M., O.J. Koo, J.T. Kang, D.K. Kwon, S.J. Park, S.J. Kim, J.H. Moon, H.J. Oh, G. Jang, and B.C. Lee. 2012. Paradoxical effects of kisspeptin: it enhances oocyte in vitro maturation but has an adverse impact on hatched blastocysts during in vitro culture. *Reproduction, fertility, and development*. 24:656-668.

- Saskova, A., P. Solc, V. Baran, M. Kubelka, R.M. Schultz, and J. Motlik. 2008. Aurora kinase A controls meiosis I progression in mouse oocytes. *Cell cycle (Georgetown, Tex.)*. 7:2368-2376.
- Shahed, A., and K.A. Young. 2009. Differential ovarian expression of KiSS-1 and GPR-54 during the estrous cycle and photoperiod induced recrudescence in Siberian hamsters (*Phodopus sungorus*). *Molecular reproduction and development*. 76:444-452.
- Slomczynska, M., M. Duda, and J. Galas. 2001. Estrogen receptor alpha and beta expression in the porcine ovary. *Folia histochemica et cytobiologica*. 39:137-138.
- Stricker, S.A., R. Silva, and T. Smythe. 1998. Calcium and endoplasmic reticulum dynamics during oocyte maturation and fertilization in the marine worm *Cerebratulus lacteus*. *Developmental biology*. 203:305-322.
- Taniguchi, Y., A. Kuwahara, A. Tachibana, Y. Yano, K. Yano, Y. Yamamoto, M. Yamasaki, T. Iwasa, K. Hinokio, T. Matsuzaki, and M. Irahara. 2017. Intra-follicular kisspeptin levels are related to oocyte maturation and gonadal hormones in patients who are undergoing assisted reproductive technology. *Reproductive medicine and biology*. 16:380-385.
- Terao, Y., S. Kumano, Y. Takatsu, M. Hattori, A. Nishimura, T. Ohtaki, and Y. Shintani. 2004. Expression of KiSS-1, a metastasis suppressor gene, in trophoblast giant cells of the rat placenta. *Biochimica et biophysica acta*. 1678:102-110.
- Tosti, E. 2006. Calcium ion currents mediating oocyte maturation events. *Reproductive biology and endocrinology : RB&E*. 4:26.
- Wang, C., and Z. Machaty. 2013. Calcium influx in mammalian eggs. *Reproduction (Cambridge, England)*. 145:R97-r105.

512 Wigglesworth, K., K.B. Lee, M.J. O'Brien, J. Peng, M.M. Matzuk, and J.J. Eppig. 2013.
513 Bidirectional communication between oocytes and ovarian follicular somatic cells is
514 required for meiotic arrest of mammalian oocytes. *Proceedings of the National Academy*
515 *of Sciences of the United States of America*. 110:E3723-3729.

516 Yang, Y., L. Shu, Q. Jiang, H. Huang, and H. Ye. 2018. Does the bone morphogenetic protein 7
517 inhibit oocyte maturation by autocrine/paracrine in mud crab? *General and comparative*
518 *endocrinology*. 266:119-125.

519 Ye, Y., K. Kawamura, M. Sasaki, N. Kawamura, P. Groenen, M.D. Gelpke, R. Rauch, A.J.
520 Hsueh, and T. Tanaka. 2009. Kit ligand promotes first polar body extrusion of mouse
521 preovulatory oocytes. *Reproductive biology and endocrinology : RB&E*. 7:26.

522 Yu, C., S.Y. Ji, Q.Q. Sha, Y. Dang, J.J. Zhou, Y.L. Zhang, Y. Liu, Z.W. Wang, B. Hu, Q.Y. Sun,
523 S.C. Sun, F. Tang, and H.Y. Fan. 2016. BTG4 is a meiotic cell cycle-coupled maternal-
524 zygotie-transition licensing factor in oocytes. *Nature structural & molecular biology*.
525 23:387-394.

526

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6